

# Isotopic exchange on the diurnal scale between near-surface snow and lower atmospheric water vapor at Kohnen station, East Antarctica

François Ritter<sup>1,2</sup>, Hans Christian Steen-Larsen<sup>2,3</sup>, Martin Werner<sup>1</sup>,  
Valérie Masson-Delmotte<sup>2</sup>, Anais Orsi<sup>2</sup>, Melanie Behrens<sup>1</sup>, Gerit Birnbaum<sup>1</sup>,  
Johannes Freitag<sup>1</sup>, Camille Risi<sup>4</sup>, and Sepp Kipfstuhl<sup>1</sup>

<sup>1</sup>Alfred-Wegener-Institut (AWI) Helmholtz-Zentrum für Polar und Meeresforschung, Bremerhaven, Germany.

<sup>2</sup>Laboratoire des Sciences du Climat et de l'Environnement (LSCE), IPSL/CEA-CNRS-UVSQ, Saclay, Gif-sur-Yvette, France.

<sup>3</sup>Center for Ice and Climate, Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark.

<sup>4</sup>Laboratoire de Météorologie Dynamique (LMD), IPSL/CNRS-UPMC, Paris, France.

*Correspondence to:* François Ritter (fritte2@uic.edu)

**Abstract.** Quantifying the magnitude of post-depositional processes affecting the isotopic composition of surface snow is essential for a more accurate interpretation of ice core data. To achieve this, high temporal resolution measurements of both lower atmospheric water vapor and surface snow isotopic composition are required. This study presents the first continuous measurements of water vapor isotopes performed in East Antarctica (Kohnen station) from December 2013 to January 2014 using a laser spectrometer. Observations have been compared with the outputs of two atmospheric general circulation models (AGCMs) equipped with water vapor isotopes: ECHAM5-wiso and LMDZ5Aiso. During our monitoring period, the signals in the 2 m air temperature  $T$ , humidity mixing ratio  $q$  and both water vapor isotopes  $\delta D$  and  $\delta^{18}O$  are dominated by the presence of diurnal cycles. Both AGCMs simulate similar diurnal cycles with a mean amplitude 30 % to 70 % lower than observed, possibly due to an incorrect simulation of the surface energy balance and the boundary layer dynamics. In parallel, snow surface samples were collected each hour during 35 h, with a sampling depth of 2-5 mm. A diurnal cycle in the isotopic composition of the snow surface is observed in phase with the water vapor, reaching a peak-to-peak amplitude of 3 ‰ for  $\delta D$  over 24 h (compared to 36 ‰ for  $\delta D$  in the water vapor). A simple box model treated as a closed system has been developed to study the exchange of water molecules between an air and a snow reservoir. In the vapor, the box model simulations show too much isotopic depletion compared to the observations. Mixing with other sources (advection, free troposphere) has to be included in order to fit the observations. At the snow surface, the simulated isotopic values are close to the observations with a snow reservoir of  $\sim 5$  mm depth (range of the snow sample depth). Our analysis suggests that fractionation

occurs during sublimation and that vapor-snow exchanges can no longer be considered insignificant for the isotopic composition of near surface snow in polar regions.

## 1 Introduction

Thanks to the design of mass spectrometers and their application to stable water isotopes since the  
25 1950s, precipitation has long been sampled for laboratory stable isotope analyses in order to trace  
atmospheric processes related to the hydrological cycle (e.g., Dansgaard, 1964). Past changes in  
precipitation isotopic composition have also been investigated using a variety of natural archives.  
Among these, ice cores form one of the most direct records of the isotopic composition of past  
precipitation. In Antarctica, stable water isotope measurements are central for past climate recon-  
30 structions from these ice cores, through atmospheric distillation processes connecting temperature,  
condensation and isotopic composition (e.g., Masson-Delmotte et al., 2008). However, the relation-  
ship between precipitation isotopic composition and climate is complex, as it is affected by frac-  
tionation taking place at most phase transition, at evaporation, during atmospheric transport, and  
condensation.

35 While spatial relationships have been documented using surface snow data (Masson-Delmotte  
et al., 2008), only few studies have examined the drivers of Antarctic precipitation isotopic com-  
position variability at timescales associated with synoptic or diurnal events (Fujita and Abe, 2006;  
Schlosser et al., 2004). In this study we focus on these timescales. The climatic controls on precipita-  
tion isotopic composition have been investigated under different climatic contexts thanks to the im-  
40 plementation of stable water isotopes into Atmospheric General Circulation Models (AGCMs) (e.g.,  
Joussaume et al., 1984). Classically, the mean isotopic composition of precipitation simulated by at-  
mospheric models is directly compared to ice core data, thereby ignoring potential post-depositional  
processes which may transform the initial precipitation signal in the upper part of the firn. In this  
study we aim at investigating one such post-depositional process: the isotopic exchange between  
45 atmospheric water vapor and snow.

The possibility to monitor isotopic exchanges between surface snow and low level atmospheric  
water vapor has emerged thanks to recent technological development. New laser spectrometers have  
been released by two companies (Los Gatos Research Inc. and Picarro Inc.) in the 2000s, creating a  
substantial advance within the field of water isotope research. These analyzers are able to perform  
50 continuous and in-situ measurements of the humidity mixing ratio (defined as the mass of water  
vapor divided by the mass of dry air) and the stable water vapor isotope concentrations. Prior to  
this advance, water vapor could only be collected using tedious cold trapping methods (e.g., Jacob  
and Sonntag, 1991; Steen-Larsen et al., 2011; Angert et al., 2008), deployed until sufficient amounts  
were collected to allow subsequent transfer to vials and later mass spectrometer analyses. Such a  
55 task was almost impossible to perform routinely (e.g., Schwarz et al., 1998).

The first implementation of continuous in-situ isotopic monitoring of surface water vapor isotopic  
composition above an ice sheet was achieved at NEEM, NW Greenland, during summer field sea-  
sons 2010-2012 (Steen-Larsen et al., 2013, 2014). In parallel, repeated sampling of surface snow  
was also implemented for laboratory isotopic analyses. The combined snow and vapor datasets re-

60 vealed changes in the isotopic composition of surface snow in response to synoptic changes in the atmospheric water vapor isotope values, and exhibited a strong diurnal variability in near-surface water vapor isotopic composition. Over several days and weeks, parallel variations between the surface snow isotopic composition and the near-surface vapor isotopic composition were identified in between snowfall events. This was surprising, as the snow isotopic composition was expected to be  
65 controlled by the isotopic composition of precipitation. The variations were interpreted to reflect interactions between the snow surface and the lower atmosphere occurring during surface snow metamorphism. This new finding has potential implications for the interpretation of ice core records, and for the comparison of ice core data with atmospheric model results. It is unclear however, whether surface air-snow exchanges could lead to significant changes in the snow isotopic composition at  
70 colder sites, such as those of the Antarctic Plateau.

Here, we report the first surface vapor isotope measurements performed above the Antarctic ice sheet. These measurements were performed at the German Kohnen station, a deep ice coring site with intermediate temperature and a humidity mixing ratio high enough in the summer for making accurate measurements of the water vapor isotopic composition. In parallel, snow surface samples  
75 were collected for comparison with the vapor on a diurnal scale. The surface of the ice sheet around Kohnen is characterized by the presence of large sastrugi, created by wind redistribution and sublimation of snow, hence producing considerable variability in the snow surface age, origin and density. In particular, very hard dunes sticking up above the mean surface level may be half a year old (Birnbauer et al., 2010). A previous study performed at Vostok station (Antarctica) reported a large  
80 variability in the isotopic composition of the snow surface (10 cm depth) over an 1 km transect, with a maximum variation of 30 ‰ in  $\delta D$  over 100 m horizontally (Ekaykin et al., 2002). The isotopic composition of the snow surface at Kohnen station is expected to show a similar spatial variability, depending on the age and the origin of the snow patch. The importance of the sublimation and condensation processes in the isotopic composition of different snow patches will be investigated. The  
85 term “condensation” (rather than “deposition”) is preferred in this paper to describe the water phase change from vapor to solid in order to avoid a possible confusion with “post-depositional processes”.

After a brief overview of the Kohnen station environment (Sect. 2), this article details the successful implementation of continuous measurements of water vapor isotopic composition during the months of December 2013 and January 2014. Section 3 (material and methods) describes our  
90 protocol for water vapor data processing, and reports the accuracy of the data. We also report the parallel surface snow sampling over 35 hours and introduce simulations performed with two AGCMs equipped with stable water isotopes, LMDZ5Aiso and ECHAM5-wiso. In Sect. 4, we present observed and simulated values, first for the day to day variability and then we focus on the diurnal cycles. We compare the diurnal variability of the isotopic composition of the water vapor and of the  
95 very first layer of surface snow through a box model. The last section summarizes our key conclusions and perspectives.

## 2 Kohnen station environment

The German Kohnen station [75°00'S, 0°04'E] is located on the Antarctic Plateau in Dronning Maud Land, 550 km from the South Atlantic coast line and 2892 m above sea level. Near the station, the surface elevation has a gentle slope of  $\sim 1.3 \pm 0.3 \text{ m.km}^{-1}$  with a direction of  $\sim 61^\circ$ . This place is characterized by katabatic winds ( $\sim 8 \text{ m.s}^{-1}$ ) on diurnal time scale, which form around 3 h and vanish around 15 h (UTC time) (Van As et al., 2005). The katabatic regime can be interrupted by the influence of synoptic systems, responsible for 20 % of snowfall events at Kohnen station (Schlosser et al., 2010). From December 2013 to January 2014, five snowfall events were observed and no significant snow accumulation was detected from daily measurements of snow surface height within a precision of  $\sim 1 \text{ mm}$  (protocol described in Sect. 3.5).

The mean temperature for the month of January 2014 was  $-25^\circ\text{C}$ , similar to the climatological average ( $-25 \pm 2^\circ\text{C}$ , 1998 to 2013). Both temperature values are based on hourly data from the permanent Automatic Weather Station at Kohnen (hereafter AWS9, described in Sect. 3.1). During clear sky conditions, the net radiation at the surface is predominantly positive during the day (i.e., a dominant energy gain by shortwave radiation from the sun despite a high albedo of the snow surface) and negative during the night (i.e., a net loss by longwave emissions from the surface), imprinting a strong diurnal cycle to the surface temperature, with a peak-to-peak amplitude of  $14^\circ\text{C}$  (Van As et al., 2005).

## 3 Material and methods

### 3.1 Weather observations

Weather observations are reported every hour and precipitation events are labeled "snowfall", "light snowfall" or "diamond dust". A snowfall event leaves a visible accumulation on flat surfaces (for example transport boxes) whereas during a light snowfall event or a diamond dust event no visible accumulation is observed. During the time of observations, snowfall and light snowfall events were almost always associated with 6/8 to 8/8 cover of low-level clouds whereas the cloud cover of low-level clouds was almost always 0/8 to 2/8 in case of diamond dust.

Two Automatic Weather Stations (AWS) were also installed at Kohnen (Fig. 1). AWS9 is permanent and performs measurements since  $\sim 1998$ . AWS 13/14 was only temporarily installed at Kohnen for the summer season, from December 2013 to January 2014. Hourly averages of the following parameters measured at 2 m height are available: air pressure, temperature, relative humidity, wind speed and wind direction.

### 3.2 Water vapor sampling system

Measurements of water vapor isotopes in the near surface atmosphere were performed from 2013/12/17  
130 to 2014/01/21 using a Los Gatos Research Inc. analyzer (hereafter simply analyzer), type DTL-  
100. It continuously measured the humidity mixing ratio and the ratio of two stable water isotopes:  
 $R_{18O} = [^1H_2^{18}O]/[^1H_2^{16}O]$  and  $R_D = [^1H^2H^{16}O]/[^1H_2^{16}O]$  (Baer et al., 2002). We will use through-  
out this paper the standard  $\delta$  notation in ‰:

$$\delta^* = \left( \frac{R_*}{R_{VSMOW}} - 1 \right) \times 1000 \quad (1)$$

135 where  $\delta^*$  stands for  $\delta D$  or  $\delta^{18}O$ , and  $R_{VSMOW}$  is the ratio of the Vienna Standard Mean Ocean Water.  
The instrumental temporal resolution is 2 Hz, but we report measurements averaged over 11 minutes  
to increase the signal to noise ratio. The precision decreases with humidity, leading us to exclude all  
measurements performed below 500 ppmv.

The analyzer was calibrated using a stream of water vapor with known constant isotopic compo-  
140 sition generated by the Water Vapor Isotope Standard Source (WVISS, Los Gatos Research Inc.).  
The WVISS allowed control over the amount of dilution of the vapor stream resulting in a vapor  
stream of adjustable humidity level. A working standard was created at the beginning of the cam-  
paign by melting surface snow and subsequently stored in a sealed glass container. Samples were  
taken from the working standard every two weeks for later laboratory isotopic analysis to check for  
145 stability. No significant drift was observed. The working standard was calibrated against VSMOW,  
SLAP (Standard Light Antarctic Precipitation) and GISP (Greenland Ice Sheet Precipitation) stan-  
dards at the Alfred Wegener Institute, Bremerhaven (hereafter AWI). Its isotopic composition was:  
 $\delta^{18}O = -44.44 \pm 0.03 \text{ ‰}$ , and  $\delta D = -345.5 \pm 0.1 \text{ ‰}$ .

Figure 1 shows the location of the instrument compared to the main wind direction ( $\sim 61^\circ$  true  
150 north) and its distance to the base. The large clean area prevents any perturbations from local human  
activity. Measurements were performed at three different heights above the snow surface: 0.2, 0.9 and  
3 m. Insulated and heated copper tubes were used to suck in air to the analyzer following the setup of  
Steen-Larsen et al. (2013). An air-filter and a snowfall protection were placed on each inlet to prevent  
sucking in snow crystals. The three copper tubes and the WVISS were connected to a device called  
155 the Multiport Input Unit (Los Gatos Research Inc.). This device was controlled by the analyzer to  
switch the valves and alternate between the inlet measurements and WVISS calibrations. Dry air was  
flushed through the system to check for leaks. The order of measurements during the campaign was  
the following: 12 minutes calibration followed by three cycles of 33 minutes, measuring 11 minutes  
at each inlet.

### 160 3.3 Calibrations

The calibration protocol follows Steen-Larsen et al. (2013). In short, one calibration is applied to the  
humidity mixing ratio and then three types of calibrations of the measured water vapor isotopes are

performed: instrumental humidity-isotope response calibration, VSMOW-SLAP calibration and drift correction. More details about the calibration procedures are given in the supplementary material.

165 The humidity mixing ratio is calibrated against the relative humidity measured by the AWS9. This relative humidity has been previously calibrated following the protocol of Anderson (1994), setting its maximum values equal to 100 % of humidity. Then the relative humidity is converted into the humidity mixing ratio  $q$  using the surface pressure (Goff-Gratch equation with respect to ice) and the air temperature at 2 m height. We finally calculate a fit from AWS9 to analyzer  $q$ , with  
170 a 2nd order polynomial to get the function correcting the humidity mixing ratio measured during the campaign. This function is  $f(q) = a + b \times q + c \times q^2$  with  $a = 80 \pm 80$ ,  $b = 0.59 \pm 0.14$  and  $c = (0.23 \pm 0.06) \times 10^{-3}$ . We have checked the linear relationship between  $\delta D$  and  $q$  (shown in Table 5) with/without this calibration and conclude that the slope is not sensitive.

The instrumental humidity-isotope response calibration was obtained by varying the humidity  
175 level of the vapor stream produced by the WVISS while measuring the same standard. As the water vapor isotopic composition generated by the WVISS is assumed to be constant, the observed variation of the water vapor isotopes can be attributed to the instrumental humidity response. Five humidity calibrations were performed during the campaign (2013/12/06, 12/14, 12/28 and 2014/01/13, 01/28). For each of them, we continuously generated a stream of water vapor from the “Working Standard water”, using the WVISS during  $\sim 20$  hours. We forced the humidity to vary from  
180  $\sim 4000$  ppmv to 300 ppmv by changing the amount of dilution. The humidity isotope response in-between calibration periods is assumed to vary linearly.

The VSMOW-SLAP calibration is carried out by measuring vapor from the WVISS generated when evaporating standards of known isotopic composition referenced against the VSMOW-SLAP  
185 scale. Four different water standards (NZE, NEEM, TALOS, OC3) with a known isotopic composition were brought to Kohnen station in 40 glass bottles of 10 cl. During a VSMOW-SLAP calibration, each standard was vaporized and measured during 15+15 minutes at two different humidity levels, which were used to check the accuracy of the humidity correction. This calibration lasted  $\sim 6$  hours and has been reproduced four times on 2013/12/06, 2013/12/15 and 2014/01/08, 2014/01/22. Very  
190 small variations in the VSMOW-SLAP calibration slope were observed (see Table 1) and we have therefore simply calculated the mean value to obtain our conversion slopes :  $\alpha_{\delta D} = 1.43 \pm 0.02$  and  $\alpha_{\delta^{18}O} = 1.00 \pm 0.03$  (uncertainties are the standard error of the mean value).

Finally the measured isotopic value is corrected for the drift by measuring vapor generated by the WVISS when evaporating the prepared working standard. This measurement is performed every  
195 111 minutes during 12 minutes and linear interpolation is assumed between each ‘drift’-correction measurement.

### 3.4 Precision and accuracy of measurements

We use the following notation to describe the error propagation, with  $\delta^*$  standing either for  $\delta D$  or  $\delta^{18}O$ .

- 200 1. The raw isotopic composition (direct output from the analyzer without any corrections) averaged over 11 minutes is  $\delta_{\text{raw}}^* \pm d\delta_{\text{raw}}^*$  with  $d\delta_{\text{raw}}^*$  the standard error associated with the mean value.
2. The humidity isotope response associated to  $\delta_{\text{raw}}^*$  is  $\Gamma_* \pm d\Gamma_*$  with  $d\Gamma_*$  the uncertainty associated with the humidity correction.  $d\Gamma_*$  is for a given  $q$  and a given time the absolute difference  
205 between two humidity isotope responses from two consecutive humidity calibrations.  $d\Gamma_*$  is therefore taken as the maximum possible error on the humidity correction.
3. The slope conversion to the VSMOW-SLAP international scale is  $\alpha_* \pm d\alpha_*$  with  $d\alpha_*$  the uncertainty associated with the slope.  $d\alpha_*$  is the standard error of the mean value of the slopes from the 4 different VSMOW calibrations.
- 210 4. The drift correction is  $\mu_* \pm d\mu_*$  with  $d\mu_*$  the uncertainty associated with the drift correction.  $d\mu_*$  has been estimated at the end of the campaign by performing an extra 24 hours calibration with a stable humidity.
5. The corrected isotopic composition is  $\delta_{\text{corr}}^* \pm d\delta_{\text{corr}}^*$  with  $d\delta_{\text{corr}}^*$  the final uncertainty containing both the precision and the accuracy on the corrected measurements.

215 We apply the three corrections (humidity isotope response, conversion to the VSMOW-SLAP scale, drift removal) to calculate the corrected isotopic composition at a given time:

$$\delta_{\text{corr}}^* = \alpha_* \times (\delta_{\text{raw}}^* - \Gamma_*) - \mu_*$$

We obtain a final uncertainty on the corrected isotopic composition by applying an error propagation calculation, assuming no correlation between the three corrections:

$$(d\delta_{\text{corr}}^*)^2 = \overbrace{\alpha_*^2 \times ((d\delta_{\text{raw}}^*)^2 + (d\Gamma_*)^2)}^{P_*^2} + (d\mu_*)^2 + \underbrace{(d\alpha_*)^2 \times (\delta_{\text{raw}}^* - \Gamma_*)^2}_{A_*^2}$$

220

With  $A^*$  standing for the accuracy of the measurements and  $P^*$  standing for the precision of the measurements. We have attributed the part  $A^*$  to the accuracy because the uncertainty on the VSMOW correction will affect the mean value of the data over the campaign. Each correction depends on the time of the measurement (the drift varies through time, the humidity isotope response as well) and Table 2 summarizes the different orders of magnitude of the parameters with the estimated  
225 precision/accuracy.



We estimate a precision of the measurements of 3.0 ‰ for  $\delta D$  and of 0.9 ‰ for  $\delta^{18}O$ . We estimate an accuracy of the measurements of 11 ‰ for  $\delta D$  and of 2.5 ‰ for  $\delta^{18}O$ . When focusing on the mean diurnal cycle, we will get a higher precision by calculating hourly averages over 18 days (Sect. 4.2).

### 3.5 Surface snow sampling and analysis

In order to detect any snow accumulation or erosion (due to snowfall events or wind drift), 100 thin wood sticks were distributed every meter along a 100 m transect in a clean area and daily measured with a folding ruler. No accumulation or erosion was detected within a precision of 1 mm.

The Snow Surface Diurnal Cycle experiment (SSDC) was devoted to the detection of a diurnal cycle in the isotopic composition of surface snow. Keeping in mind the possible variability in the isotopic composition of the snow surface, three different areas with consistent surface snow texture were selected, based on visual observation (the border of the snow patch was visible) and subjective assessment of the hardness. The snow sampling protocol is based on the assumption that the isotopic composition of a snow patch at a given time is homogeneous. Patch 1 was made of hard ice, patch 2 of compact snow and patch 3 was composed of soft snow. Five adjacent samples for each patch were sampled every hour (15 samples per hour) during a 35-hour period, from 2014/01/08 to 2014/01/10 (as it is shown in Fig. 2 indicated by the SSDC label). The sample depth is estimated between 2 and 5 millimeters, the tool used was a cake shovel. Samples were scraped into a plastic bag, which was sealed and shipped for subsequent isotope measurements at the AWI, using two water isotope analyzers Picarro, type L1102-i and L2120-i. The protocol followed Geldern and Barth (2012).

### 3.6 Atmospheric simulations

ECHAM5-wiso (Werner et al., 2011) is the isotopic version of the atmospheric general circulation model ECHAM5 (Roeckner et al., 2003). Simulations from ECHAM5-wiso are implicitly nudged to the European Center for Medium-Range Weather Forecasts (ECMWF) ERA-interim reanalyses data (Berrisford et al., 2011) using 6-hourly pressure, temperature, divergence and vorticity fields (Rast et al., 2013). Sea surface temperatures and sea ice coverage are derived from the ERA-interim data set too. For our purpose this model has been run with a high vertical and horizontal T106L31 resolution (31 levels, 1.1 ° in longitude × 1.1 ° in latitude). The lowest model level (about 60 m above the surface) has been selected followed by a bi-linear interpolation of nearby model grid points to the location of Kohnen base [75°00'S, 0°04'E]. The simulation has been started in 1979 and any potential model spin-up bias, e.g. caused by the initialization of the atmosphere in terms of humidity and its isotopic composition, can be safely neglected for our study period.

LMDZ5Aiso (hereafter LMDZiso) is the isotopic version of LMDZ5A, part of the atmospheric general circulation model IPSL-CM5A used in the Coupled Model Intercomparison Project (CMIP5) (Risi et al., 2010). For our purpose this model has been run with a vertical resolution of 39 levels and

with a stretched grid designed to refine the horizontal resolution in Antarctica: the resolution is of 0.30 ° latitude x 2.5 ° longitude in Antarctica, yielding a resolution of about 33 km in latitude x 72 km in longitude around Kohlen. Simulations are constrained by sea surface temperature (SST) data from the National Centers for Environmental Prediction (NCEP) and nudged to the 6-hourly ECMWF analyses using only the wind fields. There could theoretically be some inconsistencies between the winds and the SSTs from different reanalyses datasets, but the impact should be very small due to the overall consistency between the two reanalyses datasets and due to the strong nudging of the winds, preventing any drift. Because such high-resolution simulation is costly, the simulation has been started in January 2013 but inspection of simulated time series show that the spinup is sufficient.

Three selected outputs from both models are calculated at a specific height, 10 m for the wind speed and wind direction, and at 2 m for temperature. However, the reader should notice that simulated parameters such as humidity or water vapor isotopes are extracted from the first vertical model level (which represents a height of 60 m above ground) whereas the in-situ observations are measured close to the surface. Furthermore, at Kohlen Station, the 2 m temperature in ECHAM5-wiso is calculated from the surface energy balance equation, assuming a constant surface albedo of 0.8. This might also lead to further differences between simulation results and observations.

## 4 Results and discussion

Section 4.1 will be devoted to the study of the day-to-day variability over the study period of both air observations at 3 m and simulations from ECHAM5-wiso and LMDZiso. Section 4.2 will focus on the diurnal scale, showing subtle differences between the 0.2 and 3 m inlets in the mean value of 18 selected days (labeled as horizontal orange bars in Fig. 2) and comparing it with AGCM outputs. Finally, Section 4.3 will study the diurnal cycle in the snow surface, by comparing the results from the snow surface samples collected during the SSDC experiment (labeled as the horizontal purple bar in Fig. 2) with isotopic simulations from a snow surface-air model running as a closed system.

### 4.1 Observed and simulated day-to-day variability

In order to estimate the magnitude of the day-to-day variability, days with data gaps larger than 8 h have been removed from the data-set (on 12/16, 12/28, 12/29, 01/13, 01/14, 01/17, 01/18 and 01/21) and then 29 daily mean values of  $q$ ,  $T$ , d-excess and  $\delta D$  have been calculated for the observations and the model outputs. Average and standard deviation of these parameters over the 29 mean values are detailed in Table 3.

The signal from the temperature, humidity mixing ratio and water vapor isotopes measurements is stable, with a much lower variability on the day-to-day scale compared to the diurnal scale (Fig. 2). The mean humidity mixing ratio measured at 3 m over the study period is 1100 ppmv (Table 3),

with a maximum of 2200 ppmv on 12/22 at 15 h UTC. It coincides with the highest temperature, reaching  $-15.5\text{ }^{\circ}\text{C}$  at 17 h UTC on the same day. By contrast, the driest and coldest conditions are encountered on 01/19, at 2 h UTC, and is estimated at 150 ppmv for the humidity mixing ratio (therefore below the limit of confidence of our instrument, 500 ppmv) and measured at  $-35.8\text{ }^{\circ}\text{C}$  for the air temperature. Deuterium mean value at 3 m is  $-410\text{ }_{\text{‰}}$ , with a range of variation from  $-360\text{ }_{\text{‰}}$  (on 12/18 at 17 h UTC) to  $-470\text{ }_{\text{‰}}$  (on 01/21 at 3 h UTC).

We now compare the model performances for these daily mean values. ECHAM5-wiso correctly simulates these mean conditions for the temperature, humidity mixing ratio and deuterium while LMDZiso produces a mean humidity mixing ratio 40 % lower than observed and shows a positive offset of  $56\text{ }_{\text{‰}}$  in deuterium (Table 3). We notice that the mean radiative input (in longwave and shortwave) measured at the surface by the AWS 9 is  $583\text{ W}\cdot\text{m}^{-2}$ , compared to only  $552\text{ W}\cdot\text{m}^{-2}$  for LMDZiso. An incorrect simulation of the cloud cover (and subsequently the precipitation) is likely related to this offset in LMDZiso. The surface energy balance determines the mean surface temperature ( $-27\text{ }^{\circ}\text{C}$  for LMDZiso compared to  $-24\text{ }^{\circ}\text{C}$  for ECHAM5-wiso), which itself impacts the sublimation rate and the 2 m air temperature via sensible and latent heat exchanges with the lower atmosphere. The radiative offset present in LMDZiso could explain the low simulated values of temperature and humidity mixing ratio.

Observations depict a day-to-day variability of 200 ppmv for the humidity mixing ratio,  $9\text{ }_{\text{‰}}$  for deuterium and  $4\text{ }^{\circ}\text{C}$  for temperature (Table 3). Both models underestimate the variability of temperature and humidity mixing ratio. One explanation concerning temperature could be that both models fail to capture the very cold events observed on 12/30 and 01/19. They also underestimate the variations of temperature and humidity mixing ratio observed over several days of relatively clear sky from 12/20 to 12/24, with an amplitude in  $T$  ( $q$ ) of  $0.9\text{ }^{\circ}\text{C}$  (250 ppmv) for ECHAM5-wiso against  $4.2\text{ }^{\circ}\text{C}$  (490 ppmv) for the observations. The deuterium has a higher day-to-day variability in ECHAM5-wiso because of the simulation of high depletion events which strongly deviate from the observations (Fig. 2). For example on 01/05 at 6 h the simulated value is as low as  $-520\text{ }_{\text{‰}}$  against  $-430\text{ }_{\text{‰}}$  for the top inlet. These depletion events do not correspond to any parallel signal in the simulated meteorological data (cloud cover, wind speed, temperature or humidity mixing ratio) and further analyses will be necessary to understand these artifacts.

We finally focus on deuterium excess. The observed mean value at 3 m is  $\sim 30\text{ }_{\text{‰}}$ , with a strong relative variability of  $9\text{ }_{\text{‰}}$  (Table 3). ECHAM5-wiso simulates deuterium excess values close to the observations within these uncertainties, while LMDZiso shows a low signal of  $15\text{ }_{\text{‰}}$  with almost no variability. Some of the air masses simulated by ECHAM5-wiso have even a correct deuterium excess variability despite the fact that the model is not able to simulate correctly the magnitude of the  $\delta D$  depletion during these days, as reported above (e.g., on 01/05 and 01/13). This might be explained by a stronger dependency of deuterium excess variability to climate conditions during

evaporation processes in the vapor source regions, while the  $\delta D$  signal is understood to be more directly controlled by climate conditions near or at Kohnen station.

Table 5 presents a synthesis of the linear relationships between the deuterium and different parameters (temperature, humidity mixing ratio,  $\delta^{18}O$  and d-excess) for the observed and simulated values on two different time scales (day-to-day and diurnal). The left column corresponds to the 29 daily mean values we have calculated before. Only  $\delta D$  vs  $\delta^{18}O$  presents a strong linear relationship on this time scale, because the day to day variability of other parameters (mixing ratio, temperature) is much weaker than their diurnal variations (Sect. 4.2). The strong linear relationship on the diurnal scale highlights the importance of the local processes, which will be investigated through a box model in Sect. 4.3.

## 4.2 Observed and simulated diurnal cycles

We have chosen 18 days showing a strong diurnal variation in the humidity mixing ratio and the isotopes for the purpose of stacking them: for each of these days and for each parameter, the daily mean value has been subtracted to obtain the anomalies. Then from these anomalies we have calculated for each hour (from 1 to 24, hours UTC) the average of the 18 values and its associated standard deviation. For visualization purposes, we have calculated the average of the 24 standard deviations, called hereafter MSD for 'Mean Standard Deviation' (each error bar in Fig. 3 represents  $\pm 1$  MSD). The MSD associated with the wind direction has been obtained differently, because this parameter is calculated from the U and V wind components. Knowing the standard deviation  $\sigma_U$  and  $\sigma_V$  associated with the U and V stacks, a Monte-Carlo simulation has been performed to calculate for each run a different wind direction from random values of  $U \pm 2\sigma_U$  and  $V \pm 2\sigma_V$ . The MSD associated with the wind direction is the average of the maximum range of values obtained from the Monte-Carlo simulation (n=10000). Also, the saturated mixing ratio has been shifted to preserve the difference ( $q_{sat} - q$ ) and allow a meaningful comparison with the humidity mixing ratio anomalies. The average temperature (average humidity mixing ratio at 3 m) over the 18 selected days is  $-23.6 \pm 1.9$  °C (1130  $\pm$  170 ppmv), which is close to the mean values over the campaign ( $-23$  °C, 1100 ppmv) and indicates that these days constitute a representative sample of the campaign. We will also frequently refer to the measurements performed by Van As et al. (2005) at Kohnen from 2002/01/08 to 2002/02/09 in temperature and specific humidity at two different heights: surface and 1 m.

The mean wind speed is  $4.4 \pm 0.1$  m.s<sup>-1</sup> over the 18 selected days, with a diurnal variation from 5.8 m.s<sup>-1</sup> (12 h UTC) to 2.4 m.s<sup>-1</sup> (20 h UTC). The mean wind direction over the 18 selected days is  $46 \pm 50$  °. These measurements are therefore consistent with the presence of katabatic winds toward the slope direction of the terrain (61 °). Both models fail to simulate the pattern of katabatic winds. The mean wind direction is  $32 \pm 27$  ° for ECHAM5-wiso and  $21 \pm 24$  ° for LMDZiso, and their mean wind velocity at 10 m is only 3.1 m.s<sup>-1</sup> for ECHAM5-wiso and 1.9 m.s<sup>-1</sup> for LMDZiso. They also

show a low diurnal variability, whereas Van As et al. (2005) observed at the same height variations higher than  $2 \text{ m}\cdot\text{s}^{-1}$  over 24 h. The underestimation might be due to the horizontal resolution, which  
370 is too coarse to represent properly the katabatic winds, especially in LMDZiso.

We focus on the peak-to-peak amplitude of the diurnal cycles, summarized in Table 4. The diurnal amplitude of observed surface air temperature at 2 m is  $10.0 \text{ }^\circ\text{C}$  (within a MSD of  $1.0 \text{ }^\circ\text{C}$ ). Van As et al. (2005) found an amplitude of  $\sim 11 \text{ }^\circ\text{C}$  at 1 m, which is consistent with our observations. For the measurements from the 3 m inlet, we find a diurnal amplitude for  $q$  of 930 ppmv and for  $\delta D$  of 36 ‰.  
375 The magnitude of mean daily  $\delta D$  and temperature variations is similar to that observed in Greenland at NEEM ( $\sim 35 \text{ }^\circ\text{C}$  for  $\delta D$  and  $\sim 10 \text{ }^\circ\text{C}$ ), despite much larger humidity variations in Greenland with  $\sim 2300 \text{ ppmv}$  (Steen-Larsen et al., 2013). Both models underestimate the amplitude of the diurnal cycles in the air temperature at 2 m by more than 50 %. The surface temperature simulated by both AGCMs has a peak-to-peak amplitude of  $7 \text{ }^\circ\text{C}$ , compared to  $\sim 14 \text{ }^\circ\text{C}$  for the measurements of  
380 Van As et al. (2005). Variations of the surface temperature at Kohnen are supposed to be driven on the first order by the radiative budget. We have therefore compared the radiative budget of the AWS 9, ECHAM5-wiso and LMDZiso. Both models show good agreement with the observations for the net shortwave budget at the surface. However, the longwave radiative components are more difficult to simulate. Downward longwave emissions are related to the cloud cover (greenhouse effect) and  
385 snowfalls, whereas upward longwave emissions are related to the surface temperature and emissivity of the surface. Both models show difficulties simulating a proper cloud cover and snowfall events, and the variation in their surface temperature is 50 % lower than observed. That explains the disagreement between the observations and the AGCMs with respect to the longwave radiative budget, leading to a wrong simulation of the surface temperature.

The diurnal amplitude observed at 3 m is 930 ppmv for  $q$  and 36 ‰ for  $\delta D$ . The diurnal variation in the water vapor isotopes and temperature is similar at NEEM in Greenland ( $\sim 35 \text{ }^\circ\text{C}$  for  $\delta D$  and  $\sim 10 \text{ }^\circ\text{C}$  for  $T$ ), despite much larger diurnal variations in the humidity mixing-ratio with a peak-to-peak amplitude of  $\sim 2300 \text{ ppmv}$  (Steen-Larsen et al., 2013). Both AGCMs simulate a lower amplitude for the humidity mixing-ratio likely due to the low variation in their surface temperature,  
395 but interestingly ECHAM5-wiso manages to simulate the right magnitude of the diurnal deuterium variability (32 ‰), associated with a large MSD of 15 ‰ (Table 4). The model-data comparison is hampered by the fact that the simulated humidity mixing-ratio is only available from the first grid level of the AGCMs, and is therefore an average value over the first  $\sim 60 \text{ m}$ . The height and stability of the boundary layer is particularly difficult to simulate over ice, and have a certain impact  
400 on the presence or absence of diurnal cycles (Holtslag et al., 2013). A proper understanding of the simulation of the boundary layer by the AGCMs would require relevant output parameters such as the boundary layer depth or stability classes, which have not been implemented yet. Further analyses will therefore be necessary to understand the different behavior of LMDZiso and ECHAM5-wiso.

We now compare the amplitudes at 0.2 and 3 m for the humidity mixing ratio and the  $\delta D$ . The absolute difference between the amplitudes measured from the bottom and the top inlets is 80 ppmv for  $q$  (20 % of  $\Delta q$ ) and 4 ‰ for  $\delta D$ . Within a MSD of  $\sim 125$  ppmv for  $q$  and  $\sim 6$  ‰ for  $\delta D$ , we cannot conclude that the amplitude is decreasing significantly with height.

In the observations, diurnal air temperature variations occur in phase with  $\delta D$  and humidity mixing ratio measured at 0.2 m, with minima at 2 h and maxima at 16 h. While  $\delta D$  observations from 0.2 and 3 m seem synchronous within uncertainties, the top inlet presents a clear lag of 1 hour behind the bottom inlet with respect to the measured humidity mixing ratio. The same delay was observed by Van As et al. (2005) from the surface to 1 m in both temperature and humidity mixing ratio. Their observations showed minima and maxima for  $T$  and  $q$  at  $\sim 3$  h (time UTC) and  $\sim 16$  h at 1 m against  $\sim 2$  h and  $\sim 15$  h at the surface. The diurnal observations from NEEM (Steen-Larsen et al., 2013) were also in phase with  $T$ ,  $q$  and  $\delta D$  at  $\sim 1$  m, however humidity mixing ratio and  $\delta D$  measured on a much higher tower ( $\sim 13$  m) showed a delay of only  $\sim 1$  h with the observations at 1 m. In the simulations from ECHAM5-wiso and LMDZiso, temperature diurnal variations occur approximately synchronous with the observations at Kohlen. However, simulated humidity mixing ratio and deuterium variations are delayed by  $\sim 3$  hours compared to the observations from the 0.2 and 3 m inlets. This could be explained by the fact that the temperature at 2 m is driven by the surface radiative budget while the timing of changes in humidity/isotopes may reflect boundary layer dynamics which are less accurately simulated.

The d-excess values also depict a diurnal cycle anti-correlated to  $\delta D$ , with an amplitude for the 0.2 m inlet (3 m inlet) of 21 ‰ (15 ‰) but associated with a large MSD of 7 ‰ (6 ‰). This anti-correlation is expected from the d-excess linear definition: at very low temperature, d-excess is influenced by distillation, and increases as  $\delta D$  decreases. ECHAM5-wiso underestimates the diurnal amplitude variability and LMDZiso fails to simulate any diurnal variability in the d-excess.

As previously reported for daily mean values (Table 5, Sect. 4.1), close linear relationships are observed between  $q$ ,  $\delta D$  and d-excess for hourly mean values, highlighting the importance of local fluxes. These relationships are better simulated for the diurnal cycle, but ECHAM5-wiso tends to overestimate the associated slopes compared to the observations. LMDZiso strongly overestimates the slope of  $\delta D$ - $\delta^{18}O$ . ECHAM5-wiso is able to capture the diurnal anti-correlation of d-excess and  $\delta D$ . The slope calculated on the hourly scale for  $\delta D$  vs  $\delta^{18}O$  has a value of  $5.99 \pm 0.12$ , whereas Steen-Larsen et al. (2013) calculated at NEEM a slope of  $6.47 \pm 0.07$ , in a warmer and more humid air ( $\sim 3000$  ppmv against  $\sim 1200$  ppmv at Kohlen).

### 4.3 Air-snow exchanges

In order to document the isotopic exchange between the surface snow and the overlying water vapor, snow surface samples were collected from three snow patches, with five juxtaposed replicas per patch and per hour (protocol described in Section 3.5). In the isotopic composition of the snow

440 surface presented in Figure 4, each hourly data point is the average of the five snow samples, with  
a standard deviation of 0.03 ‰ for  $\delta^{18}O$  and 0.2 ‰ for  $\delta D$ . We identify a clear diurnal cycle in the  
snow with a significant peak-to-peak amplitude of  $\sim 3$  ‰ for  $\delta D$  and  $\sim 0.4$  ‰ for  $\delta^{18}O$ , in phase  
with the diurnal cycle in the air (Fig. 4). The mean deuterium value of the snow patches varies from  
-296 ‰ to -316 ‰, showing that the texture of the snow patch and its isotopic composition could  
445 be related (Table 6). This observation confirms the spatial variability previously observed at Vostok  
in the isotopic composition of the snow surface (10 cm depth), with variations up to 30 ‰ in  $\delta D$   
over 100 m horizontally (Ekaykin et al., 2002). Both AGCMs manage to simulate a similar isotopic  
composition for the snow surface, with on average a deuterium value of -330 ‰ for ECHAM5-wiso  
and -299 ‰ for LMDZiso. As expected, the isotopic composition of the snow surface simulated by  
450 the AGCMs depends on the snowfall events only, with a variation in  $\delta D$  of 6 ‰ for ECHAM5-wiso  
over the study period (no variation is simulated by LMDZiso). Unfortunately, the cooling phase  
during the night of 2014/01/09 is restricted compared to the usual strong decrease in temperature or  
humidity shown in Fig. 3 because the presence of a cloud cover. This meteorological event is likely  
to have impacted  $T$ ,  $q$  and  $\delta D$  during the cooling phase.

455 We notice that the vapor is close to or at saturation by looking at the saturated mixing-ratio cal-  
culated in Fig. 4. We can therefore expect condensation to occur during the night (and sublimation  
during the day as the diurnal cycle is observed to be approximately symmetrical) and an isotopic  
exchange between the lower atmospheric water vapor and the surface snow. This raises the question,  
if and by what magnitude condensation and sublimation processes might affect the surface snow and  
460 lower the water vapor isotopic composition.

In order to address this question, we set up a simple box model as a closed system containing  
two interacting and homogeneous reservoirs. Figure 5 depicts the schematics of this model, and  
introduces our notations. As this system is closed, the variation of moisture in the air (and its isotopic  
composition) is only due to condensation/sublimation during the cooling/warming phase and we  
465 have mass conservation of the water molecules:  $\forall t, m_t^v + m_t^s = m_0^v + m_0^s$  with  $t = 0$  the start of the  
cooling phase and  $v$  and  $s$  indices representing vapor and snow, respectively. Our simulation is based  
on the values from the mean stack of the 18 diurnal cycles instead of the specific day corresponding  
to the snow sampling because of the unusually high temperature and humidity during the night of  
2014/01/09. We split our analysis into two parts: the cooling phase (from  $\sim 17$  h to  $\sim 2$  h UTC) and  
470 the warming phase (from  $\sim 3$  h to  $\sim 15$  h UTC). More details concerning the equations present in the  
box model are given in the supplementary material.

### 4.3.1 Cooling phase

The deposition of the condensate on the snow surface during the cooling phase is expected to reach a  
maximum height of  $\zeta^{\max} \sim 0.1$  mm (calculation detailed in Fig. 5). As the depth of our surface snow  
475 samples is  $\sim 2 - 5$  mm, we mix the condensate with the snow reservoir. From  $t$  to  $t + 1$ , an amount

( $m_t^v - m_{t+1}^v$ ) condensates and the isotopic ratio of the condensate in equilibrium with the vapor is  $\alpha_t R_t^v$  with  $R$  the isotopic ratio of the heavy isotope and  $\alpha_t$  the associated fractionation coefficient with respect to ice calculated from the air temperature measured at 2 m at time  $t$ . Assuming an immediate removal of the condensate from the air reservoir and an immediate mixing with the snow reservoir, we obtain for the isotopic composition of the vapor  $\delta_t^v$  and the isotopic composition of the snow  $\delta_t^s$ :

$$\delta_{t+1}^v + 1000 = A_t^v (\delta_t^v + 1000) \quad (2)$$

$$\delta_{t+1}^s + 1000 = B_t^s (\delta_t^s + 1000) + B_t^v (\delta_t^v + 1000) \quad (3)$$

with

$$A_t^v = \frac{q_t}{q_{t+1}} - \alpha_t \left( \frac{q_t}{q_{t+1}} - 1 \right)$$

$$B_t^s = \frac{\rho_s h_0 + \rho_d H_0 (q_0 - q_t)}{\rho_s h_0 + \rho_d H_0 (q_0 - q_{t+1})}$$

$$B_t^v = \alpha_t \frac{\rho_d H_0 (q_t - q_{t+1})}{\rho_s h_0 + \rho_d H_0 (q_0 - q_{t+1})}$$

We will consider the equilibrium case (relative humidity RH set to 1) but also the supersaturated case (RH = 1.1) by replacing  $\alpha_t$  with the equivalent fractionation coefficient (Jouzel and Merlivat, 1984), which takes into account the kinetic effects in a supersaturated environment. The required input parameters for simulating  $\delta_t^v$  are  $q_t$ ,  $\delta_0^v$ ,  $\alpha_t$  and RH (set to 1 or 1.1). The required input parameters for simulating  $\delta_t^s$  are  $q_t$ ,  $\delta_t^v$  (measured, not simulated),  $\alpha_t$ ,  $h_0$ ,  $H_0$ , RH,  $\rho_s$ ,  $\rho_d$  and  $\delta_0^s$ .

Figure 6 presents the simulation of  $\delta D_t^v$  during the cooling phase, based on equations (2) and the isotopic variation of the condensate (Eq. (3) with  $h_0 = 0$  mm), which does not depend on  $H_0$ ,  $\rho_s$  or  $\rho_d$ . We have used fractionation coefficients with respect to ice given by Merlivat and Nief (1967) and Ellehoj et al. (2013).

The amplitude of the simulated isotopic composition of the vapor is for each case three times larger than observed. The box model is closed, so any change in the vapor is forced to condensate in order to keep the mass conservation equation. In reality, there is advection of air masses with different moisture or temperature into and out of the box. Exchanges with the free troposphere are also possible. This could partly contribute to the decrease of humidity during the cooling phase in an open system instead of a pure condensation process in a closed system. Nevertheless, our simplistic approach leads to the conclusion that about 40% of the diurnal vapor mixing ratio variation is sufficient to simulate the right order of magnitude of isotopic variations, based on equilibrium



fractionation. This is consistent with the results of ECHAM5-wiso: while this atmospheric model underestimates the diurnal variability of humidity (by 40%), it does correctly capture the diurnal  
510 variability of deuterium.

The isotopic variation of the condensate is  $\sim 6\text{‰}$  and decreases in phase with the vapor. We define  $(\delta D_0^s)_{\text{eq}}$  as the deuterium value of the condensate at equilibrium with the initial vapor. The value of  $(\delta D_0^s)_{\text{eq}}$  is  $-295\text{‰}$  using the fractionation coefficient from Merlivat and Nief (1967). This result is consistent with the mean value of the isotopic composition of the three snow patches (Table 6).

515 Figure 7 presents the simulation of  $\delta D_t^s$  during the cooling phase, based on equations (3). We have used  $\rho_d = 0.95 \text{ kg.m}^{-3}$ , calculated from the surface pressure, temperature and relative humidity measured at Kohnen, and  $\rho_s = 340 \text{ kg.m}^{-3}$ , calculated from 100 daily snow samples collected at Kohnen during the period of air measurement. The initial depth of the snow reservoir  $h_0$  correspond the depth of the SSDC snow samples, i.e., between 2 mm and 5 mm. The air reservoir is considered  
520 as the part of the lower atmosphere affected by convection and turbulence within a time scale of about one hour. The parameter  $H_t$  is therefore the mixing-layer height. Sodar measurements performed at Dome C (Antarctica) showed magnitudes between 10 m and 300 m (Pietroni et al., 2012; Casasanta et al., 2014). In this study, we set the mixing-layer height as a constant and consider two cases:  $H_0 = 50 \text{ m}$  or  $H_0 = 100 \text{ m}$ .

525 This simple parametrization allows us to test the sensitivity of the box model to the variations of  $H_t$ . We also have chosen two initial isotopic compositions of the snow surface as two distinct cases: (i)  $\delta D_0^s = -310\text{‰}$ , below  $(\delta D_0^s)_{\text{eq}}$  and (ii)  $\delta D_0^s = -290\text{‰}$ , above  $(\delta D_0^s)_{\text{eq}}$ .

In the first case, the mixing between the condensate and the snow surface will tend toward the equilibrium through a positive trend; in the second case, a negative trend is predicted. This is due to  
530 the difference between the isotopic composition of the condensate and the snow surface, positive or negative at a given time  $t$ . The response of the model shown in Fig. 7 depends strongly on parameters that are not well constrained. These parameters are the box sizes (a snow reservoir with a depth above 1 cm will keep a constant isotopic composition), the fractionation coefficients (disagreement between Merlivat and Nief (1967) and Ellehoj et al. (2013)) or the value of  $\delta D_0^v$ , which could be  
535 measured with an accuracy of  $11\text{‰}$  only (see Table 2). However, we are able to conclude that the condensation of water vapor has an effect on the isotopic composition of the top 2 mm of the snow surface.

#### 4.3.2 Warming phase

It is generally assumed that no fractionation occurs during sublimation. Using Greenland data, Steen-  
540 Larsen et al. (2011) and Landais et al. (2012) showed that on average the snow surface isotopes and the water vapor isotopes are in equilibrium, and estimated that the value of the equilibrium factor lies between the fractionation coefficient  $\alpha^{\text{ice}}$  with respect to ice (Merlivat and Nief, 1967; Ellehoj et al., 2013) and the fractionation coefficient  $\alpha^{\text{water}}$  with respect to water (Majoube, 1971). In this study,

we test different hypotheses to obtain a range of prediction of the isotopic variation in the vapor and  
 545 the snow surface. From  $t$  to  $t + 1$ , an amount  $(m_{t+1}^v - m_t^v)$  sublimates and the isotopic ratio  $R_t^{\text{sub}}$  of  
 the sublimate will be tested under three different assumptions: (i) no fractionation occurs and  $R_t^{\text{sub}} = R_t^s$ , (ii) the sublimate is formed in equilibrium with the snow and  $R_t^{\text{sub}} = R_t^s/\alpha_t$ , (iii) the kinetic  
 effect due to subsaturation is taken into account and a thin layer of liquid water above the snow with  
 the same isotopic composition is considered. Following Merlivat and Jouzel (1979) we have in case  
 550 (iii):

$$R_t^{\text{sub}} = \frac{1-k}{1-\text{RH}} \left( \frac{R_t^s}{\alpha_t} - \text{RH} \times R_t^v \right)$$

With the relative humidity RH set equal to 0.9, and the kinetic fractionation factor k given by  
 $k_{\delta^{18}\text{O}} = 6.2\%$  and  $k_{\delta\text{D}} = 5.5\%$ . We present the equations for the vapor and the snow surface for  
 the case (iii) only, noticing that cases (i) and (ii) are mathematically obtained from case (iii).

555 Assuming an immediate removal of the sublimate from the snow reservoir and an immediate  
 mixing with the molecules contained in the air reservoir, we have for the isotopic composition of the  
 vapor:

$$\delta_{t+1}^v + 1000 = E_t^v (\delta_t^v + 1000) + E_t^s (\delta_t^s + 1000) \quad (4)$$

with

$$560 E_t^v = \frac{q_t}{q_{t+1}} - \text{RH} \times \frac{1-k}{1-\text{RH}} \left( 1 - \frac{q_t}{q_{t+1}} \right)$$

$$E_t^s = \frac{1}{\alpha_t} \frac{1-k}{1-\text{RH}} \left( 1 - \frac{q_t}{q_{t+1}} \right)$$

The equations for the isotopic composition of the snow are:

$$\delta_{t+1}^s + 1000 = F_t^s (\delta_t^s + 1000) + F_t^v (\delta_t^v + 1000) \quad (5)$$

565 with

$$F_t^s = \frac{\rho_s h_0 + \rho_d H_0 \left( q_0 - q_t - \frac{1}{\alpha_t} \frac{1-k}{1-\text{RH}} (q_{t+1} - q_t) \right)}{\rho_s h_0 + \rho_d H_0 (q_0 - q_{t+1})}$$

$$F_t^v = \text{RH} \times \frac{1-k}{1-\text{RH}} \times \frac{\rho_d H_0 (q_{t+1} - q_t)}{\rho_s h_0 + \rho_d H_0 (q_0 - q_{t+1})}$$

Figure 8 displays the measurements and simulations performed for the isotopic composition of  
 570 the deuterium and d-excess of the vapor during the warming phase. When no fractionation occurs  
 during sublimation, the simulated variation of the deuterium is two times higher than observed. If we  
 were sublimating a block of solid ice, it would be conceivable that only the very surface atoms would  
 be able to sublimate, and the system would not fractionate. However, in the presence of very porous  
 snow, there are a very large numbers of water molecules participating in the snow-air interface,

575 and it is possible that snow would behave more like a liquid than like a solid in this respect, and  
fractionate. We tested for the presence of fractionation by running the box model with a variety of  
available fractionation factors for air over ice and water (Merlivat and Nief, 1967; Ellehoj et al.,  
2013; Majoube, 1971), and find that using  $\alpha^{\text{ice}}$  the model underestimates the variations in  $\delta D$  (Fig.  
8). The true fractionation factor at sublimation is probably lower than  $\alpha^{\text{ice}}$ , but the crude nature of  
580 our model prevents us from quantifying it precisely.

Figure 9 presents the simulated  $\delta D$  of the surface snow during the warming phase. We notice  
that  $\delta_{t+1}^s = \delta_t^s$  when no fractionation occurs as it is assumed in ECHAM5-wiso and LMDZ-iso. A  
difference between equilibrium and subsaturation has to be noticed in Eq. (5) due to the coefficient  
 $F_t^v$ . If  $F_t^v = 0$  (equilibrium), there is no influence of the vapor on the isotopic composition of the  
585 surface snow and  $\delta_0^s$  will not have a significant impact on  $\delta_t^s$ : any patch of snow will share the same  
isotopic variation whatever its initial isotopic composition is. If  $F_t^v \neq 0$  (subsaturation), the isotopic  
composition of the snow is affected by the isotopic composition of the vapor, and in that case the  
variation of the isotopic composition of a snow patch during the warming phase will depend on its  
initial isotopic composition  $\delta_0^s$  (Fig. 7).

590 We focus here only on  $\delta D_0^s = -320 \text{‰}$ , which is the average of the 100 daily snow samples collected  
at Kohnen over the measurement period. Our data from the three snow patches consistently depict  
a positive trend during the warming phase, with an amplitude between 3 and 7 ‰ for  $\Delta\delta D$  (Fig.  
9). Simulations with different values of  $H_0$ ,  $h_0$  and  $\alpha_t$  share the same positive trend with a peak-to-  
peak amplitude between 1 and 8 ‰, which is of the same order of magnitude as the observations.  
595 We notice that the uncertainties related to the heights of the reservoirs have greater impacts on the  
simulated snow surface isotopic values than the different fractionation coefficients. As a result, we  
are not able to constrain the fractionation factor at sublimation, but we observe that since the surface  
snow isotopic composition is changing, the sublimation process must be associated with an isotopic  
fractionation.

## 600 5 Conclusions

Continuous measurements of temperature, humidity mixing ratio and water vapor isotopes were  
performed during summer 2013/2014 at Kohnen station in East-Antarctica. These data highlight  
a strong diurnal cycle, in contrast with rather stable day-to-day mean levels over one month of  
observations. During our monitoring period, the surface vapor isotopic composition was therefore  
605 more driven by local processes than by synoptic changes. Outputs from the two AGCMs (ECHAM5-  
wiso and LMDZiso) show in general good agreements with the observations. However, the surface  
temperature variations simulated by the models have an amplitude 50 % lower than observed by  
Van As et al. (2005), likely due to the difficulty to simulate the longwave radiative budget (related to  
the cloud cover and snowfall events). Moreover, the strong katabatic winds observed at Kohnen are

610 not properly simulated by the AGCMs. The simulation of processes in the polar boundary layer and associated inversion is also known to be a challenge for AGCMs (Holtslag et al., 2013). This could explain why the amplitude of the diurnal cycles is lower in the models compared to the observations.

We have investigated the diurnal isotopic response of the upper thin layer of snow surface to the atmospheric variations. A continuous hourly sampling over 35 hours of the first  $\sim 2\text{-}5$  mm of the snow surface of three different snow patches reveals a significant variability in both  $\delta D$  and  $\delta^{18}O$  615 during a period without snowfall events. As these variations in the surface snow isotopic composition follow the diurnal trend in the air, this result confirms the observations of Steen-Larsen et al. (2013) at NEEM who also observed parallel variations between the snow surface isotopic composition and lower atmosphere isotopic composition. These observations were however with a higher variation 620 on a day-to-day scale. In their case, they reported larger variations in the parallel variations in the isotopic composition of both surface snow (5 mm) and vapor, reaching 10 ‰ over 5 days.

Two important consequences can be inferred from the snow sample diurnal observations: (1) post-depositional processes have a significant impact on the isotopic composition of the snow surface and (2) the sublimation process is fractionating. These two points are not included in classical isotopic 625 theory and therefore not implemented in atmospheric models.

In order to determine the contributions of condensation and sublimation to the isotopic variations of the vapor and surface snow, we developed a simple model describing the isotopic exchange between two reservoirs contained in a closed system: a water vapor column and a thin snow surface layer. We find that the observed isotopic variations in the water vapor phase ( $\delta D^v$ ) are about half of 630 what simple condensation/evaporation equilibrium would dictate. This is likely due to advection, exchanges with the free troposphere and variations in the boundary layer height. Additionally, for our observed isotopic composition of surface (2-5mm) snow changes during both the warming and the cooling phase (peak-to-peak variation of  $\sim 3$  ‰ for  $\delta D$ ), our crude model is able to reproduce these observations, although the model results depend strongly on the size of the reservoir chosen. According our box model, no diurnal cycle in the isotopic composition of the snow surface is expected 635 from a depth of 1 cm or above. We do observe an increase in  $\delta D^s$  during the sublimation process, which indicates that water isotopes undergo fractionation during sublimation. The only doubt we could emit on this result is related to the wind drift. Effectively, the isotopic variability observed on the diurnal scale in a snow patch could also be attributed to the renewal of the snow surface by the

640 wind, which mixes the surface of the snow with ice crystals coming from other snow patches. Assuming that fractionation occurs during sublimation, the uncertainties in the model geometry and in air advection prevent us from being able to determine the fractionation coefficient. Our water vapor isotope data suggest however that it is smaller than  $\alpha^{ice}$ . Further analyses are required to quantify the impact of post-depositional processes on the isotopic signal from ice core data.

645 The day-to-day variations in water vapor isotopic composition have a much smaller amplitude than the diurnal cycle, because no large synoptic event was recorded during our monitoring period.

Expanding the temporal framework of such monitoring is a pre-requisite in order to better understand the importance of horizontal advection, and to evaluate the processes at play during the winter season. Our observations show the possible importance of surface snow - surface vapor exchanges  
650 for the isotopic composition recorded in ice cores. This stresses the potential of isotopic monitoring of snow-air interactions for the study of fractionation processes during water phase change, but also underscore the importance of improvements in analytical accuracy under low humidity conditions. This constitutes an experimental challenge for future works.

*Acknowledgements.* This first paper is dedicated to Dominique Hironde, former teacher of history of sciences,  
655 for being a role model for scientific curiosity and fascination toward reality. The scientific campaign would not have been possible without the efficient work of the AWI logistic staff. Special thanks also to Carleen Reijmer for providing the data of AWS9, to Michael Schaefer, Gert Koenig-Langlo and Bernd Loose for their contribution to the meteorological observations and to Max Berkelhammer for his valuable contribution to the final version of the manuscript. LMDZiso simulations were performed on the Ada supercomputer at IDRIS un-  
660 der GENCI project 0292. ECHAM5-wiso simulations were performed on AWI's high-performance computing systems with support by the Scientific Computing research group.

## References

- Anderson, P.: A method for rescaling humidity sensors at temperatures well below freezing, *Journal of Atmospheric and Oceanic Technology*, 11, 1388–1391, 1994.
- 665 Angert, A., LEE, J.-E., and Yakir, D.: Seasonal variations in the isotopic composition of near-surface water vapour in the eastern Mediterranean, *Tellus B*, 60, 674–684, 2008.
- Baer, D. S., Paul, J. B., Gupta, M., and O’Keefe, A.: Sensitive absorption measurements in the near-infrared region using off-axis integrated cavity output spectroscopy, in: *International Symposium on Optical Science and Technology*, pp. 167–176, International Society for Optics and Photonics, 2002.
- 670 Berrisford, P., Dee, D., Poli, P., Brugge, R., Fielding, K., Fuentes, M., Kållberg, P., Kobayashi, S., Uppala, S., and Simmons, A.: *The ERA-Interim archive Version 2.0*, Shinfield Park, Reading, 2011.
- Birbaum, G., Freitag, J., Brauner, R., König-Langlo, G., Schulz, E., Kipfstuhl, S., Oerter, H., Reijmer, C. H., Schlosser, E., Faria, S. H., et al.: Strong-wind events and their influence on the formation of snow dunes: observations from Kohlen station, Dronning Maud Land, Antarctica, *Journal of glaciology*, 56, 891–902, 2010.
- 675 Casasanta, G., Pietroni, I., Petenko, I., and Argentini, S.: Observed and modelled convective mixing-layer height at Dome C, Antarctica, *Boundary-Layer Meteorology*, 151, 597–608, 2014.
- Dansgaard, W.: Stable isotopes in precipitation, *Tellus A*, 16, 1964.
- Ekaykin, A. A., Lipenkov, V. Y., Barkov, N. I., Petit, J. R., and Masson-Delmotte, V.: Spatial and temporal variability in isotope composition of recent snow in the vicinity of Vostok station, Antarctica: implications for ice-core record interpretation, *Annals of Glaciology*, 35, 181–186, 2002.
- 680 Ellehoj, M., Steen-Larsen, H. C., Johnsen, S. J., and Madsen, M. B.: Ice-vapor equilibrium fractionation factor of hydrogen and oxygen isotopes: Experimental investigations and implications for stable water isotope studies, *Rapid Communications in Mass Spectrometry*, 27, 2149–2158, 2013.
- 685 Fujita, K. and Abe, O.: Stable isotopes in daily precipitation at Dome Fuji, East Antarctica, *Geophysical Research Letters*, 33, n/a–n/a, doi:10.1029/2006GL026936, <http://dx.doi.org/10.1029/2006GL026936>, 118503, 2006.
- Geldern, R. and Barth, J. A.: Optimization of instrument setup and post-run corrections for oxygen and hydrogen stable isotope measurements of water by isotope ratio infrared spectroscopy (IRIS), *Limnology and Oceanography: Methods*, 10, 1024–1036, 2012.
- 690 Holtslag, A., Svensson, G., Baas, P., Basu, S., Beare, B., Beljaars, A., Bosveld, F., Cuxart, J., Lindvall, J., Steeneveld, G., et al.: Stable atmospheric boundary layers and diurnal cycles: challenges for weather and climate models, *Bulletin of the American Meteorological Society*, 94, 1691–1706, 2013.
- Jacob, H. and Sonntag, C.: An 8-year record of the seasonal variation of  $2\text{H}$  and  $18\text{O}$  in atmospheric water vapour and precipitation at Heidelberg, Germany, *Tellus B*, 43, 291–300, doi:10.1034/j.1600-0889.1991.t01-2-00003.x, <http://dx.doi.org/10.1034/j.1600-0889.1991.t01-2-00003.x>, 1991.
- 695 Joussau, S., Sadourny, R., and Jouzel, J.: A general circulation model of water isotope cycles in the atmosphere, *Nature*, 311, 24–29, doi:10.1038/311024a0, <http://dx.doi.org/10.1038/311024a0>, 1984.
- Jouzel, J. and Merlivat, L.: Deuterium and oxygen 18 in precipitation: Modeling of the isotopic effects during snow formation, *Journal of Geophysical Research: Atmospheres (1984–2012)*, 89, 11 749–11 757, 1984.
- 700


- Landais, A., Steen-Larsen, H. C., Guillevic, M., Masson-Delmotte, V., Vinther, B., and Winkler, R.: Triple isotopic composition of oxygen in surface snow and water vapor at NEEM (greenland), *Geochimica et Cosmochimica Acta*, 77, 304–316, 2012.
- 705 Majoube, M.: Fractionnement en oxygene-18 et en deuterium entre l'eau et sa vapeur, *J. Chim. phys*, 68, 1423–1436, 1971.
- Masson-Delmotte, V., Hou, S., Ekaykin, A., Jouzel, J., Aristarain, A., Bernardo, R., Bromwich, D., Cattani, O., Delmotte, M., Falourd, S., et al.: A review of Antarctic surface snow isotopic composition: Observations, atmospheric circulation, and isotopic modeling, *Journal of Climate*, 21, 3359–3387, 2008.
- Merlivat, L. and Jouzel, J.: Global climatic interpretation of the deuterium-oxygen 18 relationship for precipitation, *Journal of Geophysical Research*, 84, 5029–5033, 1979.
- 710 Merlivat, L. and Nief, G.: Fractionnement isotopique lors des changements d'état solide-vapeur et liquide-vapeur de l'eau à des températures inférieures à 0° C, *Tellus A*, 19, 1967.
- Pietroni, I., Argentini, S., Petenko, I., and Sozzi, R.: Measurements and parametrizations of the atmospheric boundary-layer height at Dome C, Antarctica, *Boundary-layer meteorology*, 143, 189–206, 2012.
- 715 Rast, S., Brokopf, R., Cheedela, S.-K., Esch, M., Gayler, V., Kirchner, I., Kornbluh, L., Rhodin, A., Schmidt, H., Schulzweida, U., and Wieners, K.-H.: User manual for ECHAM6, *Reports on Earth System Science*, 13, 2013.
- Risi, C., Bony, S., Vimeux, F., and Jouzel, J.: Water-stable isotopes in the LMDZ4 general circulation model: Model evaluation for present-day and past climates and applications to climatic interpretations of tropical isotopic records, *Journal of Geophysical Research: Atmospheres* (1984–2012), 115, 2010.
- 720 Roeckner, E., Bäuml, G., Bonaventura, L., Brokopf, R., Esch, M., Giorgetta, M., Hagemann, S., Kirchner, I., Kornblueh, L., Manzini, E., et al.: The atmospheric general circulation model ECHAM 5. PART I: Model description, 2003.
- Schlosser, E., Reijmer, C., Oerter, H., and Graf, W.: The influence of precipitation origin on the  $\delta^{18}\text{O}$ -T relationship at Neumayer Station, Ekströmisen, Antarctica, *Annals of Glaciology*, 39, 41–48, 2004.
- 725 Schlosser, E., Manning, K., Powers, J., Duda, M., Birnbaum, G., and Fujita, K.: Characteristics of high-precipitation events in Dronning Maud Land, Antarctica, *Journal of Geophysical Research: Atmospheres*, 115, 2010.
- Schwarz, G., Kowski, P., and Gernandt, H.: Saisonale Variation von Deuterium und Sauerstoff-18 in Luftfeuchte und Niederschlag über Antarktika, *Isotopes in Environmental and Health Studies*, 34, 159–168, doi:10.1080/10256019708036343, <http://dx.doi.org/10.1080/10256019708036343>, PMID: 22087493, 1998.
- 730 Steen-Larsen, H., Masson-Delmotte, V., Hirabayashi, M., Winkler, R., Satow, K., Prié, F., Bayou, N., Brun, E., Cuffey, K., Dahl-Jensen, D., et al.: What controls the isotopic composition of Greenland surface snow?, *Climate of the Past*, 10, 377–392, 2014.
- 735 Steen-Larsen, H. C., Masson-Delmotte, V., Sjolte, J., Johnsen, S. J., Vinther, B. M., Bréon, F.-M., Clausen, H., Dahl-Jensen, D., Falourd, S., Fettweis, X., et al.: Understanding the climatic signal in the water stable isotope records from the NEEM shallow firn/ice cores in northwest Greenland, *Journal of Geophysical Research: Atmospheres* (1984–2012), 116, 2011.
- 740 Steen-Larsen, H. C., Johnsen, S. J., Masson-Delmotte, V., Stenni, B., Risi, C., Sodemann, H., Balslev-Clausen, D., Blunier, T., Dahl-Jensen, D., Ellehøj, M., et al.: Continuous monitoring of summer surface water vapor

isotopic composition above the Greenland Ice Sheet, *Atmospheric Chemistry and Physics*, 13, 4815–4828, 2013.


Van As, D., Van Den Broeke, M., and Van De Wal, R.: Daily cycle of the surface layer and energy balance on the high Antarctic Plateau, *Antarctic Science*, 17, 121–133, 2005.

745 Werner, M., Langebroek, P. M., Carlsen, T., Herold, M., and Lohmann, G.: Stable water isotopes in the ECHAM5 general circulation model: Toward high-resolution isotope modeling on a global scale, *Journal of Geophysical Research: Atmospheres*, 116, n/a–n/a, doi:10.1029/2011JD015681, <http://dx.doi.org/10.1029/2011JD015681>, d15109, 2011.







**Figure 1.** Location of Kohnen station in Antarctica (bottom right panel). Location of the two Automatic Weather Stations (AWS, left panel) and location of the measurement tent connected to the three inlets (central panel). The right schematic presents the set-up with the three inlets, the multiport (MIU), the Water Vapor Isotope Standard Source (WVISS) and the analyzer measuring the humidity mixing ratio  $q$  and isotopes  $\delta D$  and  $\delta^{18}O$  in the vapor.




**Figure 2.** Hourly observed (black and green) and simulated (red and blue) humidity mixing ratio ( $q$ ), deuterium ( $\delta D$ ), air temperature at 2 m ( $T_{2m}$ , measured with the AWS from the 2013/2014 season) and d-excess (d-exc) at Kohonen station 2013/2014. Hourly observed precipitation events are labelled light snowfall and snowfall (light and dark cyan). The Snow Surface Diurnal Cycle (SSDC) experiment period is indicated in purple, and the 18 selected days for the diurnal cycles study in orange.



**Figure 3.** Stack of 18 diurnal cycles of the wind direction and wind speed (simulated at 10 m for the models, measured at 2 m for the observations), air temperature at 2 m (measured with the AWS from the 2013/2014 season), humidity mixing ratio  $q$ , saturated mixing ratio range calculated with the surface pressure and temperature at 2 m  $\pm 1$  °C, deuterium  $\delta D$  and d-excess. Error bars represent  $\pm 1$  mean standard deviation (MSD): average of the 24 standard deviations associated with the hourly mean values.



**Figure 4.** Observations performed during the Snow Surface Diurnal Cycle (SSDC) experiment from the 2014/01/09 to the 2014/01/10. The three different snow patches are labelled 'Hard', 'Medium' and 'Soft' from their texture. Each hourly data point is the average of the measurements made on 5 snow surface samples per patch.



$\rho_d, \rho_s$  : dry air density and snow density  
 $m_t^v, m_t^s$  : water vapor mass and snow mass

$$\begin{array}{l|l} h_t = h_0 + \zeta_t & m_t^s = \rho_s h_t S \\ H_t = H_0 - \zeta_t & m_t^v = q_t \rho_d H_t S \end{array}$$


**Snow accumulation during condensation:**

$$\zeta_{t+1} \approx \zeta_t + \frac{\rho_d}{\rho_s} H_0 (q_t - q_{t+1})$$


**Order of magnitude of the parameters:**

$$\begin{array}{l|l} \rho_d = 0.95 \text{ kg.m}^{-3} & h_0 \approx 2.5 \times 10^{-3} \text{ m} \\ \rho_s = 300\text{-}380 \text{ kg.m}^{-3} & q_t \in 5\text{-}9 \times 10^{-4} \text{ kg/kg} \\ H_0 \approx 50\text{-}100 \text{ m} & \zeta^{\max} \approx 1.1\text{-}1.4 \times 10^{-4} \text{ m} \end{array}$$


**Figure 5.** Schematic of the box model and description of the input parameters (isotopes excluded).  $\zeta^{\max}$  is the maximum value of the surface height anomaly at the end of the cooling phase, calculated with two extrem values of ( $H_0, \rho_s$ ). The height of the SSSC snow samples gives an estimation of  $h_0$ .  $\rho_d$  and  $\rho_s$  have been calculated from 100 snow samples per day, surface pressure, temperature and relative humidity observed at Kohnen from 2013/12 to 2014/01.



**Figure 6.** Measured and simulated (box model) deuterium anomalies of the vapor and simulated deuterium anomaly of the condensate ( $h_0=0$  mm) during the cooling phase, from 17 h to 2 h UTC. Observations come from the stack of the 18 diurnal cycles. Equivalent fractionation coefficients have been calculated from Jouzel and Merlivat (1984) for RH=1.1.



**Figure 7.** Simulated (box model) deuterium anomalies of the snow surface during the cooling phase, with  $h_0$ ,  $H_0$ , RH,  $\alpha_t$  and  $\delta D_0^s$  as varying inputs parameters. Fractionation coefficients at equilibrium come from Merlivat and Nief (1967) and Ellehoj et al. (2013), calculated with  $T$  at 2 m. Equivalent fractionation coefficients have been calculated from Jouzel and Merlivat (1984) for RH = 1.1.



**Observations**


— 3.0 m inlet measurements

**Simulations**

- $\alpha_t = 1$  (no fractionation)
- $\alpha_t^{\text{ice}}$  from (Merlivat and Nief, 1967)
- $\alpha_t^{\text{ice}}$  from (Ellehoj et al., 2013)
- $\alpha_t^{\text{water}}$  from (Majoube, 1971)
- $\alpha_t^{\text{water}}$  from (Majoube, 1971) taking into account the kinetic effect at subsaturation (RH = 0.9)

**Figure 8.** Measured and simulated (box model) deuterium and d-excess anomalies of the vapor during the warming phase. Kinetic effects occurring at subsaturation (RH = 0.9) have been calculated following (Merlivat et al., 1979) in a smooth regime.





**Figure 9.** Simulated (box model) deuterium anomalies of the snow surface during the warming phase, with  $h_0$ ,  $H_0$  and the fractionation coefficients as varying inputs parameters. The initial isotopic composition of the snow surface has been taken as the average of the 100 daily snow samples collected at Kohnen over the period of measurements,  $\delta D_0^s = -320 \text{ ‰}$ .

day of calibration	$\delta D$ (standards) vs $\delta D$ (measured)	$\delta^{18}O$ (standards) vs $\delta^{18}O$ (measured)	standards used
2013/12/06	$1.45 \pm 0.02$	$1.03 \pm 0.07$	NZE, JASE, TD1
2013/12/15	$1.40 \pm 0.03$	$0.93 \pm 0.05$	DML, TD1, NZE, JASE
2014/01/08	$1.38 \pm 0.01$	$1.01 \pm 0.03$	NZE, OC3, TALOS, NEEM
2014/01/22	$1.47 \pm 0.01$	$1.01 \pm 0.01$	JASE, TD1, NZE, DML

**Table 1.** Conversion slopes calculated from four VSMOW-SLAP calibrations with different standards. Data have been corrected with respect to humidity before calculating the slopes. Uncertainties represent 1 standard error on the slopes.

	related to $\delta D$			related to $\delta^{18}O$		
	avg	min	max	avg	min	max
$\alpha_*$	1.43	/	/	1.00	/	/
$d\alpha_*$	0.02	/	/	0.03	/	/
$\delta_{\text{raw}}^*$	-552	-586	-519	-84.3	-94.6	-77.7
$d\delta_{\text{raw}}^*$	0.3	0.1	0.9	0.1	0.05	0.4
$ \Gamma_* $	4	0	9	1.7	0	4
$d\Gamma_*$	2	0.1	5	0.4	0	2
$d\mu_*$	1.4	/	/	0.7	/	/
$A_*$	11	10	12	2.5	2.2	2.9
$P_*$	3.0	1.4	7.3	0.9	0.7	2.0

**Table 2.** Order of magnitude of the parameters involved in the error propagation calculation for  $\delta D$  and  $\delta^{18}O$ . 'avg', 'min' and 'max' are the mean, the minimum and the maximum value of the parameter over the campaign. 'A' stands for accuracy and 'P' for precision. Every parameter is in ‰, except  $\alpha_*$  and  $d\alpha_*$  which are dimensionless.

	q (ppmv)	$\delta D$ (‰)	d-exc (‰)	T 2m (°C)
AWS 13/14	/	/	/	$-23 \pm 4$
0.2 m inlet	$1000 \pm 200$	$-413 \pm 9$	$33 \pm 11$	/
3.0 m inlet	$1100 \pm 200$	$-409 \pm 9$	$30 \pm 9$	/
ECHAM	$1120 \pm 110$	$-411 \pm 15$	$26 \pm 5$	$-23.1 \pm 1.7$
LMDZiso	$700 \pm 100$	$-355 \pm 10$	$15 \pm 1$	$-24.0 \pm 1.4$

**Table 3.** Mean values over 29 daily averages from the measurement period (days with more than 8 hours of data gap have been removed). Uncertainties represent  $\pm 1$  standard deviation of the mean value. ECHAM stands for ECHAM5-wiso.

	$\Delta q(\text{ppmv})$	$\Delta \delta D(\text{‰})$	$\Delta d\text{-exc}(\text{‰})$	$\Delta T_{2m}(\text{°C})$
AWS 13/14	/	/	/	$10.1 \pm 1.0$
0.2 m inlet	$1010 \pm 130$	$40 \pm 7$	$21 \pm 7$	/
3.0 m inlet	$930 \pm 120$	$36 \pm 6$	$15 \pm 6$	/
ECHAM	$280 \pm 60$	$32 \pm 15$	$7 \pm 4$	$4.9 \pm 0.4$
LMDZiso	$160 \pm 100$	$7 \pm 4$	$0.6 \pm 0.4$	$3.0 \pm 1.3$

**Table 4.** Peak-to-peak amplitude of the 18 selected diurnal cycles. Uncertainties represent  $\pm 1$  mean standard deviation (average of the 24 standard deviations of the hourly mean values). ECHAM stands for ECHAM5-wiso.

	(1) daily mean values (n = 29)	(2) hourly mean values (n = 24)
$\delta D$ (‰) vs q (ppmv)		
3.0 m inlet	$\alpha = 0.04 \pm 0.01$ $r^2 = 0.31$	$\alpha = 0.037 \pm 0.001$ $r^2 = 0.98$
ECHAM5-wiso	$\alpha = 0.03 \pm 0.02$ $r^2 = 0.05$	$\alpha = 0.071 \pm 0.008$ $r^2 = 0.76$
LMDZiso	$\alpha = 0.06 \pm 0.01$ $r^2 = 0.37$	$\alpha = 0.034 \pm 0.005$ $r^2 = 0.69$
$\delta D$ (‰) vs T 2m (°C)		
3.0 m inlet	$\alpha = 3.1 \pm 1.1$ $r^2 = 0.24$	$\alpha = 3.37 \pm 0.11$ $r^2 = 0.98$
ECHAM5-wiso	$\alpha = 6 \pm 4$ $r^2 = 0.10$	$\alpha = 4.7 \pm 0.9$ $r^2 = 0.54$
LMDZiso	$\alpha = 3.9 \pm 1.1$ $r^2 = 0.33$	$\alpha = 1.5 \pm 0.3$ $r^2 = 0.49$
$\delta D$ (‰) vs $\delta^{18} O$ (‰)		
3.0 m inlet	$\alpha = 6.2 \pm 0.3$ $r^2 = 0.94$	$\alpha = 5.99 \pm 0.12$ $r^2 = 0.99$
ECHAM5-wiso	$\alpha = 6.64 \pm 0.19$ $r^2 = 0.98$	$\alpha = 6.57 \pm 0.04$ $r^2 = 0.99$
LMDZiso	$\alpha = 7.5 \pm 0.1$ $r^2 = 0.99$	$\alpha = 7.41 \pm 0.02$ $r^2 = 0.99$
d-excess (‰) vs $\delta D$ (‰)		
3.0 m inlet	$\alpha = -0.21 \pm 0.06$ $r^2 = 0.32$	$\alpha = -0.32 \pm 0.03$ $r^2 = 0.86$
ECHAM5-wiso	$\alpha = -0.18 \pm 0.03$ $r^2 = 0.50$	$\alpha = -0.22 \pm 0.01$ $r^2 = 0.98$
LMDZiso	$\alpha = -0.06 \pm 0.02$ $r^2 = 0.34$	$\alpha = -0.08 \pm 0.01$ $r^2 = 0.97$

**Table 5.** Slopes and determination coefficients calculated from two data sets : (1) the 29 daily mean values from the campaign, and (2) the 24 hourly mean values from the stack composed of 18 selected diurnal cycles. Uncertainties represent 1 standard error on the slopes.

	$\overline{\delta D}$ (‰)	$\overline{\delta^{18}O}$ (‰)
air observations	-407	-54.4
patch soft	-296	-37.4
patch medium	-301	-37.0
patch hard	-316	-38.8

**Table 6.** Isotopic mean values over the SSDC period. The standard error on the mean value is below 1 ‰ for  $\delta D$  and below 0.1 ‰ for  $\delta^{18}O$ .