

Supplementary material to the paper submitted to TCD:

Using SAR satellite data time series for regional glacier mapping

Solveig H. Winsvold¹, Andreas Kääb¹, Christopher Nuth¹, Liss M. Andreassen², Ward van Pelt³, and Thomas Schellenberger¹

¹ Department of Geosciences, University of Oslo, P.O. Box 1047 Blindern, 0316 Oslo, Norway

² Section for Glaciers, Ice and Snow, Hydrology Department, Norwegian Water Resources and Energy Directorate, P.O. Box 5091 Majorstua, 0301 Oslo, Norway

³ Department of Earth Sciences, Uppsala University, Villav. 16, 752 36 Uppsala, Sweden

The supplements consist of two animations of Sentinel-1 backscatter time-series over glaciers:

- 1) TCD_Supl_Animation_KNG_Winsvold_tc-2017-136.avi
 - a. Sentinel-1 backscatter time-series images from 2015 to 2016 over Kongsvegen, Svalbard. The heatplot shows backscatter values (sigma 0) plotted along a centerline with a spacing of 300 m (y-axis) up-glacier in time. White regions in the heatplot represent missing Sentinel-1 acquisitions.
- 2) TCD_Supl_Animation_HELL_Winsvold_tc-2017-136.avi
 - a. Sentinel-1 backscatter time-series images from 2014 to 2016 over Hellstugubreen, Norway. The heatplot shows backscatter values (sigma 0) plotted along a centerline with a spacing of 100 m (y-axis) up-glacier in time. White regions in the heatplot represent missing Sentinel-1 acquisitions.