

1. Comments from Editor

Comments:

1. Line 21: change “especially” to “particularly”.
2. Line 22: change “distributed physically based” to “distributed, physically-based”.
3. Line 23: change “the long-term (from 1971 to 2013)” to “long-term (1971-2013)”.
4. Line 24-25: Delete “which is located on the”.
5. Line 25: Delete “carefully”.
6. Line 26: Delete “the”.
7. Line 27: change “the” to “spatio-temporal”.
8. Line 27: delete “the” before “hydrology”.
9. Line 27: change “the results” to “our results”.
10. Line 28: change “showed” to “show”.
11. Line 28: change “permafrost area” to “area with permafrost”.
12. Line 28-29: change “especially” to “predominantly”. Delete “frozen”.
13. Line 33: change “the” to “an”.
14. Line 58: Add “(QTP)”.
15. Line 58: change “Intensive” to “Those studies that include intensive”.
16. Line 60: “(e.g., Cite an example or two).
17. Line 60-61: change “have not been” to “are not typically”. change “available” to “available in many areas”.
18. Line 66: “As an alternat strategy, hydrological”.
19. Line 69: change “which could” to “ “but these cann not”.
20. Line 77: change “for” to “to”.
21. Line 82: change “Qinghai-Tibetan Plateau (QTP)” to “(QTP)”.
22. Line 93: Delete “the”.
23. Line 96: Delete “the”.
24. Line 105: Delete “solid”.
25. Line 107: Change “basin located on the” to “basin,”.
26. Line 108: change “On the basis of previous studies” to “Specifically”. Also, append to the end of the previous paragraph.
27. Line 115: change “of ” to “ranging from”.
28. Line 121: delete “,”.
29. Line 122: delete “,”.
30. Line 122-123: change “,” to “;” change “i.e.” to “one at”.
31. Line 124-125: change “and” to “and the other at”. Delete “see”.
32. Line 132: change “by” to “from”.
33. Line 137: Add “(LAI)” after “index”.
34. Line 140: change “leaf area index (LAI)” to “LAI”.
35. Line 155: Add “a” before “lack”. Change “measurement” to “measurements”.
36. Line 169-170: change “and the study catchment” to “which”. change “a” to “each”.
37. Line 173-174: change to “Additional hillslope properties include soil and vegetation types”.
38. Line 260: change “Eq” to “Equation”.

39. Line 276: change “simulate the permafrost we consider an underground depth of 50 m. We” to “simulate permafrost we”.
40. Line 282: change “the first layer” to “the first soil layer at the surface”. Delete “soil” before “layer”.
41. Line 283-284: change “thickness increases” to “thicknesses increase”. change “m. and then decreases linearly with depth to” “m. Then thickness decreases linearly with depth reaching”.
42. Line 284-285: change “There” to “From 1.7 m to 3 m depth, there”. change “from 1.7 m to 3.0 m to try to replicate” to “to try to capture”.
43. Line 309: change “to” to “in order to”.
44. Line 355: add “,” after “depth”.
45. Line 368-371: Line is needed in methods section to describe how simulated versus observed discharge are compared or assessed.
46. Line 371: delete “see”.
47. Line 410-415: This text should be moved to Methods.
48. Line 431-432: Delete “the”. Change “the development of taliks” to “that talik development”.
49. Line 436: change “as shown in Figure 10(a),” to “(Figure 10(a))”.
50. Line 437: change “decreased, and the ground temperature in the deep layer (with” to “decreased and the ground temperature in deep layers (depths...”
51. Line 441: add “surface”. change “frozen depths” to “depth to the base of permafrost”
52. Line 459: delete “the”
53. Line 466: change “increasing trend of the” to “increase in”
54. Line 468: change “annual water storage in the top 0-3 m layer” to “annual soil liquid water storage (0-3 m)”.
55. Line 469-450: delete “the”. Add “soil” before “water”. Delete “of the top 0-3 m”.
56. Line 476: change “the frozen soil” to “permafrost”
57. Line 483: change “as shown in Figure 10(a),” to “(Figure 10(a))”.
58. Line 485: change “as shown in Figure 10(b),” to “(Figure 10(b))”.
59. Line 488: delete “the”
60. Line 490-491: change “increased” to “increase in”. delete “the”.
61. Line 497: add “area of” before “seasonally”.
62. Line 498-499: change “runoff in the permafrost area was lower than in the seasonally frozen ground” to “the inverse was true”
63. Line 501: change “1980 and that changed” to “1980, but degraded”
64. Line 502: delete “the”.
65. Line 503: delete “the” before “runoff” and add “hydrological” before “recession”.
66. Line 510: change “larger” to “large”.
67. Line 583-584: change to “In areas with high groundwater flow rates, laterally advected heat flux may increase the thawing of permafrost”.
68. Line 592: change “challenging. This work will be done in a future study.” to “challenging, but is part of our ongoing research.”.
69. Line 602: change “grounds” to “ground”.
70. Line 606-607: change “that the increasing trend of” to “A trend of increasing”. change “is” to “of”.

71. Line 633: delete “in the future”.
72. Line 635: At your discretion, I recommend acknowledging that suggestions from reviewers have substantially improved the paper.
73. Line 638: change “for this paper are properly cited and referred in the reference list” to “cited in this paper are available from the references”.
74. Figure 1. Change color of boreholes sites to black so the they are immediately visible. Figure looks great!
75. Figure 6. Not sure why the figure looks like this after I exported to Microsoft Word, but anyway, please change “(m³/s to “(m³ · s⁻¹)” in axis titles.
76. Figure 7. Please change caption to “... “October). Results from linear regressions are indicated.”
77. Figure 8. Please change caption to “... “permafrost. Results from linear regressions are indicated.”
78. Figure 11. In order to better separate the two pairs of panels, I suggest moving the “(a) Freezing season” and “(b Thawing season” titles outside of the panels. This will provide a bit of further separation between the pairs. Change caption to “...Data and regression results are shown. The upper...”
79. Figure 12. You use “groundwater” in some places, and “ground water” in others. Please be consistent and make appropriate change. Please change caption to “... “water. “Resu from linear regressions are indicated.”
80. Figure 13. Caption: Change “changes” to “showing changes”.
81. Table1. Add “.” After “1”.
82. Table2. Add “.” After “2”.component (mm·y⁻¹)”. If you make this insertion, you can delete “(mm/yr)” from each of the column headers.

2. Author's responses

Thanks for handling our manuscript and the detail suggestions to revise the manuscript. Following comments from the editor, we have revised our manuscript. The details are given bellow.

Comments:

1. Line 21: change “especially” to “particularly”.

Reply: We have revised as suggested (Please see line 21 in the revised clean version manuscript).

2. Line 22: change “distributed physically based” to “distributed, physically-based”.

Reply: We have revised as suggested (Please see line 22 in the revised clean version manuscript).

3. Line 23: change “the long-term (from 1971 to 2013)” to “long-term (1971-2013)”.

Reply: We have revised as suggested (Please see line 23 in the revised clean version manuscript).

4. Line 24-25: Delete “which is located on the”.

Reply: We have revised as suggested (Please see line 24 in the revised clean version manuscript).

5. Line 25: Delete “carefully”.

Reply: We have revised as suggested (Please see line 25 in the revised clean version manuscript).

6. Line 26: Delete “the”.

Reply: We have revised as suggested (Please see line 26 in the revised clean version manuscript).

7. Line 27: change “the” to “spatio-temporal”.

Reply: We have revised as suggested (Please see line 26 in the revised clean version manuscript).

8. Line 27: delete “the” before “hydrology”.

Reply: We have revised as suggested (Please see line 27 in the revised clean version manuscript).

9. Line 27: change “the results” to “our results”.

Reply: We have revised as suggested (Please see line 27 in the revised clean version manuscript).

10. Line 28: change “showed” to “show”.

Reply: We have revised as suggested (Please see line 27 in the revised clean version manuscript).

11. Line 28: change “permafrost area” to “area with permafrost”.

Reply: We have revised as suggested (Please see line 27 in the revised clean version manuscript).

12. Line 28-29: change “especially” to “predominantly”. Delete “frozen”.

Reply: We have revised as suggested (Please see line 28-29 in the revised clean version manuscript).

13. Line 33: change “the” to “an”.

Reply: We have revised as suggested (Please see line 32 in the revised clean version manuscript).

14. Line 58: Add “(QTP)”.

Reply: We have revised as suggested (Please see line 57 in the revised clean version manuscript).

15. Line 58: change “Intensive” to “Those studies that include intensive”.

Reply: We have revised as suggested (Please see line 58 in the revised clean version manuscript).

16. Line 60: “(e.g., Cite an example or two).

Reply: We have revised as “(e.g., Cheng and Wu, 2007; Wu et al., 2010)” (Please see line 59-60 in the revised clean version manuscript).

17. Line 60-61: change “have not been” to “are not typically”. change “available” to “available in many areas”.

Reply: We have revised as suggested (Please see line 60-61 in the revised clean version manuscript).

18. Line 66: “As an alternat strategy, hydrological”.

Reply: We have revised as suggested (Please see line 66 in the revised clean version manuscript).

19. Line 69: change “which could” to “ “but these cann not”.

Reply: We have revised as suggested (Please see line 69 in the revised clean version manuscript).

20. Line 77: change “for” to “to”.

Reply: We have revised as suggested (Please see line 77 in the revised clean version manuscript).

21. Line 82: change “Qinghai-Tibetan Plateau (QTP)” to “(QTP)”.

Reply: We have revised as suggested (Please see line 82 in the revised clean version manuscript).

22. Line 93: Delete “the”.

Reply: We have revised as suggested (Please see line 92 in the revised clean version manuscript).

23. Line 96: Delete “the”.

Reply: We have revised as suggested (Please see line 95 in the revised clean version manuscript).

24. Line 105: Delete “solid”.

Reply: We have revised as suggested (Please see line 104 in the revised clean version manuscript).

25. Line 107: Change “basin located on the” to “basin,”.

Reply: We have revised as suggested (Please see line 106 in the revised clean version manuscript).

26. Line 108: change “On the basis of previous studies” to “Specifically”. Also, append to the end of the previous paragraph.

Reply: We have revised as suggested (Please see line 106 in the revised clean version

manuscript).

27. Line 115: change “of” to “ranging from”.

Reply: We have revised as suggested (Please see line 113 in the revised clean version manuscript).

28. Line 121: delete “,”.

Reply: We have revised as suggested (Please see line 119 in the revised clean version manuscript).

29. Line 122: delete “,”.

Reply: We have revised as suggested (Please see line 120 in the revised clean version manuscript).

30. Line 122-123: change “,” to “;” change “i.e.,” to “one at”.

Reply: We have revised as suggested (Please see line 120-121 in the revised clean version manuscript).

31. Line 124-125: change “and” to “and the other at”. Delete “see”.

Reply: We have revised as suggested (Please see line 121-123 in the revised clean version manuscript).

32. Line 132: change “by” to “from”.

Reply: We have revised as suggested (Please see line 130 in the revised clean version manuscript).

33. Line 137: Add “(LAI)” after “index”.

Reply: We have revised as suggested (Please see line 135 in the revised clean version manuscript).

34. Line 140: change “leaf area index (LAI)” to “LAI”.

Reply: We have revised as suggested (Please see line 138 in the revised clean version manuscript).

35. Line 155: Add “a” before “lack”. Change “measurement” to “measurements”

Reply: We have revised as suggested (Please see line 152 in the revised clean version manuscript).

36. Line 169-170: change “and the study catchment” to “which”. change “a” to “each”.

Reply: We have revised as suggested (Please see line 166 in the revised clean version manuscript).

37. Line 173-174: change to “Additional hillslope properties include soil and vegetation types”.

Reply: We have revised as suggested (Please see line 170 in the revised clean version manuscript).

38. Line 260: change “Eq” to “Equation”.

Reply: We have revised as suggested (Please see line 256 in the revised clean version manuscript).

39. Line 276: change “simulate the permafrost we consider an underground depth of 50 m. We” to “simulate permafrost we”.

Reply: We have revised as suggested (Please see line 272 in the revised clean version manuscript).

40. Line 282: change “the first layer” to “the first soil layer at the surface”. Delete “soil” before “layer”.

Reply: We have revised as suggested (Please see line 278 in the revised clean version

manuscript).

41. Line 283-284: change “thickness increases” to “thicknesses increase”. change “m. and then decreases linearly with depth to” “m. Then thickness decreases linearly with depth reaching”.

Reply: We have revised as suggested (Please see line 278-280 in the revised clean version manuscript).

42. Line 284-285: change “There” to “From 1.7 m to 3 m depth, there”. change “from 1.7 m to 3.0 m to try to replicate” to “to try to capture”.

Reply: We have revised as suggested (Please see line 280-281 in the revised clean version manuscript).

43. Line 309: change “to” to “in order to”.

Reply: We have revised as suggested (Please see line 312 in the revised clean version manuscript).

44. Line 355: add “,” after “depth”.

Reply: We have revised as suggested (Please see line 358 in the revised clean version manuscript).

45. Line 368-371: Line is needed in methods section to describe how simulated versus observed discharge are compared or assessed.

Reply: We have added a sentence in the section 3.3 as “The Nash-Sutcliffe efficiency and relative error are calculated using observed and simulated discharge to evaluate the model performance.” (Please see line 306-307 in the revised clean version manuscript).

46. Line 371: delete “see”.

Reply: We have revised as suggested (Please see line 374 in the revised clean version manuscript).

47. Line 410-415: This text should be moved to Methods.

Reply: We have moved it to section 3.2 (Please see line 286-291 in the revised clean version manuscript).

48. Line 431-432: Delete “the”. Change “the development of taliks” to “that talik development”.

Reply: We have revised as suggested (Please see line 429-430 in the revised clean version manuscript).

49. Line 436: change “as shown in Figure 10(a),” to “(Figure 10(a))”.

Reply: We have revised as suggested (Please see line 434 in the revised clean version manuscript).

50. Line 437: change “decreased, and the ground temperature in the deep layer (with” to “decreased and the ground temperature in deep layers (depths...”.

Reply: We have revised as suggested (Please see line 435 in the revised clean version manuscript).

51. Line 441: add “surface”. change “frozen depths” to “depth to the base of permafrost”.

Reply: We have revised as suggested (Please see line 439-440 in the revised clean version manuscript).

52. Line 459: delete “the”.

Reply: We have revised as suggested (Please see line 457 in the revised clean version manuscript).

53. Line 466: change “increasing trend of the” to ““increase in”.

Reply: We have revised as suggested (Please see line 464 in the revised clean version manuscript).

54. Line 468: change “annual water storage in the top 0-3 m layer” to “annual soil liquid water storage (0-3 m)”.

Reply: We have revised as suggested (Please see line 466-467 in the revised clean version manuscript).

55. Line 469-450: delete “the”. Add “soil” before “water”. Delete “of the top 0-3 m”.

Reply: We have revised as suggested (Please see line 467 in the revised clean version manuscript).

56. Line 476: change “the frozen soil” to “permafrost”.

Reply: We have revised as suggested (Please see line 474 in the revised clean version manuscript).

57. Line 483: change “as shown in Figure 10(a).” to “(Figure 10(a))”.

Reply: We have revised as suggested (Please see line 481 in the revised clean version manuscript).

58. Line 485: change “as shown in Figure 10(b).” to “(Figure 10(b))”.

Reply: We have revised as suggested (Please see line 483 in the revised clean version manuscript).

59. Line 488: delete “the”.

Reply: We have revised as suggested (Please see line 485 in the revised clean version manuscript).

60. Line 490-491: change “increased” to “increase in”. delete “the”.

Reply: We have revised as suggested (Please see line 487-488 in the revised clean version manuscript).

61. Line 497: add “area of” before “seasonally”.

Reply: We have revised as suggested (Please see line 494 in the revised clean version manuscript).

62. Line 498-499: change “runoff in the permafrost area was lower than in the seasonally frozen ground” to “the inverse was true”.

Reply: We have revised as suggested (Please see line 495 in the revised clean version manuscript).

63. Line 501: change “1980 and that changed” to “1980, but degraded”.

Reply: We have revised as suggested (Please see line 497 in the revised clean version manuscript).

64. Line 502: delete “the”.

Reply: We have revised as suggested (Please see line 499 in the revised clean version manuscript).

65. Line 503: delete “the” before “runoff” and add “hydrological” before “recession”.

Reply: We have revised as suggested (Please see line 499 in the revised clean version manuscript).

66. Line 510: change “larger” to “large”.

Reply: We have revised as suggested (Please see line 507 in the revised clean version manuscript).

67. Line 583-584: change to “In areas with high groundwater flow rates, laterally advected heat flux may increase the thawing of permafrost”.

Reply: We have revised as suggested (Please see line 580-581 in the revised clean version manuscript).

68. Line 592: change “challenging. This work will be done in a future study.” to “challenging, but is part of our ongoing research.”.

Reply: We have revised as suggested (Please see line 589 in the revised clean version manuscript).

69. Line 602: change “grounds” to “ground”.

Reply: We have revised as suggested (Please see line 599 in the revised clean version manuscript).

70. Line 606-607: change “that the increasing trend of” to “A trend of increasing”. change “is” to “of”.

Reply: We have revised as suggested (Please see line 603-604 in the revised clean version manuscript).

71. Line 633: delete “in the future”.

Reply: We have revised as suggested (Please see line 630 in the revised clean version manuscript).

72. Line 635: At your discretion, I recommend acknowledging that suggestions from reviewers have substantially improved the paper.

Reply: We have added a sentence “The authors would like to thank the editor and reviewers for their constructive suggestions, which have substantially improved the paper.” In the manuscript (Please see line 635-636 in the revised clean version manuscript).

73. Line 638: change “for this paper are properly cited and referred in the reference list” to “cited in this paper are available from the references”.

Reply: We have revised as suggested (Please see line 637 in the revised clean version manuscript).

74. Figure 1. Change color of boreholes sites to black so the they are immediately visible. Figure looks great!

Reply: We have revised this figure as suggested (Please see Figure 1 in the revised clean version manuscript).

75. Figure 6. Not sure why the figure looks like this after I exported to Microsoft Word, but anyway, please change “(m³/s to “(m³ · s⁻¹)” in axis titles.

Reply: We have revised this figure as suggested (Please see Figure 6 in the revised clean version manuscript).

76. Figure 7. Please change caption to “... “October). Results from linear regressions are indicated.”

Reply: We have revised this figure as suggested (Please see Figure 7 in the revised clean version manuscript).

77. Figure 8. Please change caption to “... “permafrost. Results from linear regressions are indicated.”

Reply: We have revised this figure as suggested (Please see Figure 8 in the revised

clean version manuscript).

78. Figure 11. In order to better separate the two pairs of panels, I suggest moving the “(a) Freezing season” and “(b Thawing season” titles outside of the panels. This will provide a bit of further separation between the pairs. Change caption to “...Data and regression results are shown. The upper...”.

Reply: We have revised this figure as suggested (Please see Figure 11 in the revised clean version manuscript).

79. Figure 12. You use “groundwater” in some places, and “ground water” in others. Please be consistent and make appropriate change. Please change caption to “... “water. “Resu from linear regressions are indicated.”

Reply: We have revised this figure as suggested (Please see Figure 12 in the revised clean version manuscript). And we use “groundwater” for the whole manuscript to keep consistency.

80. Figure 13. Caption: Change “changes” to “showing changes”.

Reply: We have revised this figure as suggested (Please see Figure 13 in the revised clean version manuscript).

81. Table1. Add “.” After “1”.

Reply: We have revised as suggested (Please see Table 1 in the revised clean version manuscript).

82. Table2. Add “.” After “2”. “component (mm·y-1)”. If you make this insertion, you can delete “(mm/yr)” from each of the column headers.

Reply: We have revised as suggested (Please see Table 2 in the revised clean version manuscript).

3. Author's changes in manuscript

1. Modified Figure 1, Figure 6 and Figure 11 as suggested.
2. Modified figure captions of Figure 7, Figure 8, Figure 11, Figure 12 and Figure 13 as suggested.
3. Modified Table 2 as suggested.
4. Modified all the texts as suggested.

**Change in Frozen Soils and the Effects on Regional Hydrology, Upper
Heihe Basin, Northeastern Qinghai-Tibetan Plateau**

Bing Gao¹, Dawen Yang^{2*}, Yue Qin², Yuhan Wang², Hongyi Li³, Yanlin Zhang³, and
Tingjun Zhang⁴

¹ School of Water Resources and Environment, China University of Geosciences,
Beijing 100083, China

² State Key Laboratory of Hydrosience and Engineering, Department of Hydraulic
Engineering, Tsinghua University, Beijing 100084, China

³ Cold and Arid Regions Environmental and Engineering Research Institute, Chinese
Academy of Sciences, Lanzhou, Gansu 730000, China

⁴ Key Laboratory of West China's Environmental Systems (MOE), College of Earth
and Environmental Sciences, Lanzhou University, Lanzhou, 730000, China

** Corresponding author:* Dawen Yang (yangdw@tsinghua.edu.cn)

To be submitted to: The Cryosphere, August 2017

ABSTRACT:

Frozen ground has an important role in regional hydrological cycles and ecosystems, ~~particularly~~especially on the Qinghai-Tibetan Plateau (QTP), which is characterized by high elevations and a dry climate. This study modified a distributed, physically-based hydrological model and applied it to simulate ~~the~~ long-term (~~from~~ 1971 ~~to~~ 2013) changes in frozen ground and the effects on hydrology in the upper Heihe basin, ~~which is located on the~~ northeastern QTP. The model was ~~carefully~~ validated against data obtained from multiple ground-based observations. Based on ~~the~~ model simulations, we analyzed spatio-temporal~~the~~ changes in frozen soils and their effects on ~~the~~ hydrology. ~~The~~Our results showed that the area with permafrost ~~area~~ shrank by 8.8% (approximately 500 km²), predominantly~~especially~~ in areas with elevations between 3500 m and 3900 m. The maximum ~~frozen~~ depth of seasonally frozen ground decreased at a rate of approximately 0.032 m·decade⁻¹, and the active layer thickness over the permafrost increased by approximately 0.043 m·decade⁻¹. Runoff increased significantly during the cold season (November-March) due to ~~the~~an increase in liquid soil moisture caused by rising soil temperatures. Areas in which permafrost changed into seasonally frozen ground at high elevations showed especially large increases in runoff. Annual runoff increased due to increased precipitation, the base flow increased due to changes in frozen soils, and the actual evapotranspiration increased significantly due to increased precipitation and soil warming. The groundwater storage showed an increasing trend, indicating that a reduction in permafrost extent enhanced the groundwater recharge.

41 **KEYWORDS:** permafrost; seasonally frozen ground; soil moisture; ground
42 temperature; runoff

1. Introduction

Global warming has led to significant changes in frozen soils, including both permafrost and seasonally frozen ground at high latitudes and high elevations (Hinzman et al., 2013; Cheng and Wu, 2007). Changes in frozen soils can greatly affect land-atmosphere interactions and the energy and water balances of the land surface (Subin et al., 2013; Schuur et al., 2015), altering soil moisture, water flow pathways and stream flow regimes (Walvoord and Kurylyk, 2016). Understanding the changes in frozen soils and their impacts on regional hydrology is important for water resources management and ecosystem protection in cold regions.

Previous studies based on either experimental observations or long-term meteorological or hydrological observations have examined changes in frozen soils and their impacts on hydrology. Several studies reported that permafrost thawing might enhance base flow in the Arctic and the Subarctic (Walvoord and Striegl, 2007; Jacques and Sauchyn, 2009; Ye et al., 2009), as well as in northeastern China (Liu et al., 2003; Duan et al., 2017). A few studies reported that permafrost thawing might reduce river runoff (here, runoff is defined as all liquid water flowing out of the study area), especially on the Qinghai-Tibetan Plateau (QTP) (e.g., Qiu, 2012; Jin et al., 2009). ~~Those studies that include intensive~~Intensive field observations of frozen soils have typically been performed at small spatial scales over short periods (e.g., Cheng and Wu, 2007; Wu et al., 2010). Consequently, regional patterns and long-term trends are not typically~~have not been~~ captured. Long-term meteorological and hydrological observations are available in many areas~~available~~, but they do not provide information

on soil freezing and thawing processes (McClelland et al., 2004; Liu et al., 2003; Niu et al., 2011). Therefore, previous observation-based studies have not provided a sufficient understanding of the long-term changes in frozen soils and their impact on regional hydrology (Woo et al., 2008).

As an alternate strategy, Hydrological models have been coupled with soil freezing-thawing schemes to simulate impacts of the changes in frozen soils on catchment hydrology. Several hydrological models (Rawlins et al., 2003; Chen et al., 2008) used simple freezing-thawing schemes, but these cannot~~which could not~~ simulate the vertical soil temperature profiles. The modified VIC model (Cherkauer and Lettenmaier, 1999) and the CLM model (Oleson et al., 2010) simulate vertical soil freezing-thawing processes, but they simplify the flow routing using linear schemes. Subin et al. (2013) and Lawrence et al. (2015) used the CLM model to simulate global changes in permafrost. Cuo et al. (2015) used the VIC to simulate frozen soil changes and their hydrological impacts at the plot scale in the headwaters of the Yellow River. The GEOTop model (Endrizzi et al., 2014) simulates three-dimensional water flux and vertical heat transfer in soil, but it is difficult to apply ~~for~~to regional investigations. Wang et al. (2010) and Zhang et al. (2013) incorporated frozen soil schemes in a distributed hydrological model and showed improved performance in a small mountainous catchment. More regional studies are necessary to better understand the frozen soil changes and their impacts on regional hydrologic processes and water resources.

The ~~Qinghai-Tibetan Plateau~~ (QTP) is known as Asia's water tower, and runoff

changes on the plateau have significant impacts on water security in downstream regions (Walter et al., 2010); hence, such changes have attracted considerable attention in recent years (Cuo et al., 2014). The QTP is characterized by high elevations and a cold climate. Consequently, cryospheric processes have great impacts on its hydrological processes (Cheng and Jin, 2013; Cuo et al., 2014). The thickness of permafrost on the QTP varies from 1 to 130 m, and the temperature ranges between -0.5 and -3.5 °C (Yang et al., 2010). Compared with Arctic and Subarctic soils, the frozen soils on the QTP are more sensitive to increased air temperature (Yang et al., 2010), and changes in the frozen soils may have more significant impacts on the regional hydrology.

Clear increases in ~~the~~ annual and seasonal air temperatures have been observed on the QTP (Li et al., 2005; Liu and Chen, 2000; Zhao et al., 2004). Several studies have shown changes in frozen soils based on long-term observations. For example, Cheng and Wu (2007) analyzed ~~the~~ soil temperature profiles from boreholes on the QTP and found that the active layer thickness of frozen soils increased by 0.15-0.50 m during the period of 1996-2001. Zhao et al. (2004) observed a decreasing trend of freezing depth in the seasonally frozen soils at 50 stations. Several studies have analyzed the relationship between changes in frozen soils and river discharge using observational data (Zhang et al., 2003; Jin et al., 2009; Niu et al., 2011). However, the spatio-temporal characteristics of the long-term changes in frozen soils are not sufficiently clear. Based on comprehensive field experiments (Cheng et al., 2014), a hydrological model coupling cryospheric processes and hydrological processes has been developed (Yang

et al., 2015; Gao et al., 2016). This model provides a solid basis upon which to analyze the spatio-temporal changes in frozen soils and their impacts on the regional hydrology in the upper Heihe basin, located on the northeastern QTP.

~~Specifically~~On the basis of previous studies, this study aims to: (1) explore the spatial and temporal changes in frozen soils using a distributed hydrological model with comprehensive validation and (2) analyze the hydrological responses to the changes in frozen soils during the past 40 years in the upper Heihe basin.

2. Study Area and Data

The Heihe River is one of the major inland basins in northwestern China. As shown in Figure 1, the upper reaches of the Heihe River, representing a drainage area of 10,009 km², are located on the northeastern QTP at elevations ranging from 2200 to 5000 m. The upper reaches of this river provide the majority of the water supplied to the middle and lower reaches (Cheng et al., 2014). The annual precipitation in the upper Heihe basin ranges from 200 to 700 mm, and the mean annual air temperature ranges from -9 to 5 °C. Permafrost dominates the high elevation region above 3700 m (Wang et al., 2013), and seasonal frozen ground covers the remaining portion of the study area. Glaciers are found at elevations above 4000 m, and cover approximately 0.8% of the upper Heihe basin. The upper Heihe basin contains two tributaries, each with a hydrological station; one at Qilian (on the eastern tributary) and the other at Zhamashike (on the western tributary). The outlet of the upper Heihe basin also features a hydrological station, namely, Yingluoxia (see Figure 1).

The spatial data used in this study includes atmospheric forcing data, land surface

data and actual evapotranspiration data based on remote sensing. The atmospheric forcing data include a 1-km gridded dataset of daily precipitation, air temperature, sunshine hours, wind speed and relative humidity. The gridded daily precipitation was interpolated from observations at meteorological stations (~~see~~ Figure 1) provided by the China Meteorological Administration (CMA) using the method developed by Wang et al. (2017). The other atmospheric forcing data were interpolated ~~by~~ from observations at meteorological stations using the inverse distance weighted method. The interpolation of air temperature considers the elevation-dependent temperature gradient provided by the HiWATER experiment (Li et al., 2013).

The land surface data used to run the model include land use, topography, leaf area index (LAI), and soil parameters. The topography data were obtained from the Shuttle Radar Topography Mission (SRTM) dataset (Jarvis et al., 2008) with a spatial resolution of 90 m. The land use/cover data were provided by the Institute of Botany, Chinese Academy of Sciences (Zhou and Zheng, 2014). The ~~leaf area index (LAI)~~ data with 1-km resolution were developed by Fan (2014). The soil parameters were developed by Song et al. (2016) and include the saturated hydraulic conductivity, residual soil moisture content, saturated soil moisture content, soil sand content, soil clay content and soil organic content. Monthly actual evapotranspiration data with 1-km resolution during the period of 2002-2012 were estimated based on remote sensing data (Wu et al., 2012; Wu, 2013).

The field observation data used in this study include river discharge, soil temperature, frozen depth, soil moisture and borehole observations. Daily river

discharge data were obtained from the Hydrology and Water Resources Bureau of Gansu Province. The CMA provided daily soil temperature data collected at the Qilian station from January 1, 2004 to December 31, 2013, and daily frozen depth data collected at the Qilian and Yeniugou stations from January 1, 2002 to December 31, 2013. We obtained ground temperature observations from six boreholes, whose location are shown in Figure 1, from Wang et al. (2013). We used the observations at specific dates instead of annual averages due to a lack of continuous measurements. The borehole depths are 100 m for T1, 69 m for T2, 50 m for T3, 90 m for T4, and 20 m for T5 and T7. The HiWATER experiment (Li et al., 2013; Liu et al., 2011) provided the soil moisture data from January 1 to December 31, 2014 at the A'rou Sunny Slope station (100.52 E, 38.09 N).

3. Methodology

3.1 Brief introduction of the hydrological model

This study used the distributed eco-hydrological model GBEHM (geomorphology-based ecohydrological model), which was developed by Yang et al. (2015) and Gao et al. (2016). The GBEHM is a spatial distributed model for large-scale river basins. It employs geomorphologic properties to reduce the lateral two-dimensions into one-dimension for flow routing within a sub-catchment, which greatly improves the computational efficiency while retaining the spatial heterogeneity in water flow paths at the basin scale. As shown in Figure 2, the GBEHM used a 1-km grid system to discretize the study catchment, and the study catchment which was divided into 251 sub-catchments. Each sub-catchment was further divided into flow-intervals along its

main stream. To capture the sub-grid topography, each 1-km grid was represented by a number of hillslopes with an average length and gradient, but different aspect, which were estimated from the 90-m DEM. Additional hillslope properties include soil and vegetation types~~The terrain properties of a hillslope include the slope length, slope gradient, slope aspect, soil type and vegetation type~~ (Yang et al., 2015).

The hillslope is the basic unit in the hydrological simulation of the water and heat transfers (both conduction and convection) in the vegetation canopy, snow/glacier, and soil layers. The canopy interception, radiation transfer in the canopy and the energy balance of the land surface are described using the methods of SIB2 (Sellers et al., 1985, 1996). The surface runoff on the hillslope is solved using the kinematic wave equation. The groundwater aquifer is considered as individual storage unit corresponding to each grid. Exchange between the groundwater and the river water is calculated using Darcy's law (Yang et al., 1998, 2002; Cong et al., 2009).

The model runs with a time step of 1 hour. Runoff generated from the grid is the lateral inflow into the river over the same flow interval in the corresponding sub-catchment. Flow routing in the river network is calculated using the kinematic wave equation following the sequence determined by the Horton-Strahler scheme (Strahler, 1957). The model is driven by the atmosphere forcing data and land surface data introduced in section 2.

3.2 Simulation of cryospheric processes

The simulation of cryospheric processes in the GBEHM includes glacier ablation, snow melting, and soil freezing and thawing.

(1) Glacier ablation

Glacier ablation is simulated using the following energy balance model (Oerlemans, 2001):

$$Q_M = SW(1-\alpha) + LW_{in} - LW_{out} - Q_H - Q_L - Q_G + Q_R \quad (1)$$

where Q_M is the net energy absorbed by the surface of the glacier ($\text{W}\cdot\text{m}^{-2}$); SW is the incoming shortwave radiation ($\text{W}\cdot\text{m}^{-2}$); α is the surface albedo; LW_{in} is the incoming longwave radiation ($\text{W}\cdot\text{m}^{-2}$); LW_{out} is the outgoing longwave radiation ($\text{W}\cdot\text{m}^{-2}$); Q_H is the sensible heat flux ($\text{W}\cdot\text{m}^{-2}$); Q_L is the latent heat flux ($\text{W}\cdot\text{m}^{-2}$); Q_R is the energy from rainfall ($\text{W}\cdot\text{m}^{-2}$); and Q_G is the penetrating shortwave radiation ($\text{W}\cdot\text{m}^{-2}$). The surface albedo is calculated as follows (Oerlemans and Knap, 1998):

$$\alpha = \alpha_{snow} + (\alpha_{ice} - \alpha_{snow})e^{-h/d^*} \quad (2)$$

where α_{snow} is the albedo of snow on the glacier surface; α_{ice} is the albedo of the ice surface; h is the snow depth on the glacier surface (m); d^* is a parameter describing the snow depth effect on the albedo (m).

The amount of melt water is calculated as (Oerlemans, 2001):

$$M = \frac{Q_M}{L_f} dt \quad (3)$$

where dt is the time step used in the model (s) and L_f is the latent heat of fusion ($\text{J}\cdot\text{kg}^{-1}$).

(2) Snow melt

A multi-layer snow cover model is used to describe the mass and energy balance of snow cover. The snow parametrization is based on Jordan (1991), and two constituents, namely, ice and liquid water, are used to describe each snow layer. For each snow layer,

219 temperature is solved using an energy balance approach (Bartelt and Lehnin, 2002):

$$220 \quad C_s \frac{\partial T_s}{\partial t} - L_f \frac{\partial \rho_i \theta_i}{\partial t} = \frac{\partial}{\partial z} \left(K_s \frac{\partial T}{\partial z} \right) + \frac{\partial I_R}{\partial z} + Q_R \quad (4)$$

221 where C_s is the heat capacity of snow ($\text{J} \cdot \text{m}^{-3} \cdot \text{K}^{-1}$); T_s is the temperature of the snow
 222 layer (K); ρ_i is the density of ice ($\text{kg} \cdot \text{m}^{-3}$); θ_i is the volumetric ice content; K_s
 223 is the thermal conductivity of snow ($\text{W} \cdot \text{m}^{-1} \cdot \text{K}^{-1}$); L_f is the latent heat of ice fusion ($\text{J} \cdot \text{kg}^{-1}$);
 224 I_R is the radiation transferred into the snow layer ($\text{W} \cdot \text{m}^{-2}$); and Q_R is the energy
 225 delivered by rainfall ($\text{W} \cdot \text{m}^{-2}$), which is only considered for the top snow layer. The solar
 226 radiation transfer in the snow layers and the snow albedo are simulated using the
 227 SNICAR model, which is solved using the method developed by Toon et al. (1989). Eq.
 228 (4) is solved using an implicit centered finite difference method, and a Crank-Nicholson
 229 scheme is employed.

230 The mass balance of the snow layer is described as follows (Bartelt and Lehnin, 2002):

$$231 \quad \frac{\partial \rho_i \theta_i}{\partial t} + M_{iv} + M_{il} = 0 \quad (5)$$

$$232 \quad \frac{\partial \rho_l \theta_l}{\partial t} + \frac{\partial U_l}{\partial z} + M_{lv} - M_{il} = 0 \quad (6)$$

233 where ρ_l is the density of the liquid water ($\text{kg} \cdot \text{m}^{-3}$); θ_l is the volumetric liquid water
 234 content; U_l is the liquid water flux ($\text{kg} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$); M_{iv} is the mass of ice that changes into
 235 vapour within a time step ($\text{kg} \cdot \text{m}^{-3} \cdot \text{s}^{-1}$); M_{il} is the mass of ice that changes into liquid
 236 water within a time step ($\text{kg} \cdot \text{m}^{-3} \cdot \text{s}^{-1}$); and M_{lv} is the mass of liquid water that changes
 237 into vapour within a time step ($\text{kg} \cdot \text{m}^{-3} \cdot \text{s}^{-1}$). The liquid water flux of the snow layer is
 238 calculated as follows (Jordan, 1991):

$$239 \quad U_l = -\frac{k}{\mu_l} \rho_l^2 g \quad (7)$$

240 where k is the hydraulic permeability (m^2), μ_l is dynamic viscosity of water at 0 °C

($1.787 \cdot 10^{-3} \text{ N} \cdot \text{s} \cdot \text{m}^{-2}$), ρ_l is the density of liquid water ($\text{kg} \cdot \text{m}^{-3}$) and g is gravitational acceleration ($\text{m} \cdot \text{s}^{-2}$). The water flux of the bottom snow layer is considered snowmelt runoff.

(3) Soil freezing and thawing

The energy balance of the soil layer is solved as follows (Flerchinger and Saxton, 1989):

$$C_s \frac{\partial T}{\partial t} - \rho_i L_f \frac{\partial \theta_i}{\partial t} - \frac{\partial}{\partial z} (\lambda_s \frac{\partial T}{\partial z}) + \rho_l c_l \frac{\partial q_l T}{\partial z} = 0 \quad (8)$$

where C_s is the volumetric soil heat capacity ($\text{J} \cdot \text{m}^{-3} \cdot \text{K}^{-1}$); T is the temperature (K) of the soil layers; z is the vertical depth of the soil (m); θ_i is the volumetric ice content; ρ_i is the density of ice ($\text{kg} \cdot \text{m}^{-3}$); λ_s is the thermal conductivity ($\text{W} \cdot \text{m}^{-1} \cdot \text{K}^{-1}$); ρ_l is the density of liquid water ($\text{kg} \cdot \text{m}^{-3}$); and c_l is the specific heat of liquid water ($\text{J} \cdot \text{kg}^{-1} \cdot \text{K}^{-1}$). In addition, q_l is the water flux between different soil layers ($\text{m} \cdot \text{s}^{-1}$) and is solved using the 1-D vertical Richards equation. The unsaturated soil hydraulic conductivity is calculated using the modified van Genuchten's equation (Wang et al., 2010), as follows:

$$K = f_{ice} K_{sat} \left(\frac{\theta_l - \theta_r}{\theta_s - \theta_r} \right)^{1/2} \left[1 - \left(1 - \left(\frac{\theta_l - \theta_r}{\theta_s - \theta_r} \right)^{-1/m} \right)^m \right]^2 \quad (9)$$

where K is the unsaturated soil hydraulic conductivity ($\text{m} \cdot \text{s}^{-1}$); K_{sat} is the saturated soil hydraulic conductivity ($\text{m} \cdot \text{s}^{-1}$); θ_l is the volumetric liquid water content; θ_s is the saturated water content; θ_r is the residual water content; m is an empirical parameter in van Genuchten's equation and f_{ice} is an empirical hydraulic conductivity reduction factor that is calculated using soil temperature as follows (Wang et al., 2010):

$$f_{ice} = \exp[-10(T_f - T_{soil})], \quad 0.05 \leq f_{ice} \leq 1 \quad (10)$$

where T_f is 273.15 K and T_{soil} is the soil temperature.

Equation (8) solves the soil temperature with the upper boundary condition as the heat flux into the uppermost soil layer. When the ground is not covered by snow, the heat flux from the atmosphere into the uppermost soil layer is expressed as follows (Oleson et al., 2010):

$$h = S_g + L_g - H_g - \lambda E_g + Q_R \quad (11)$$

where h is the upper boundary heat flux into the soil layer ($\text{W}\cdot\text{m}^{-2}$); S_g is the solar radiation absorbed by the uppermost soil layer ($\text{W}\cdot\text{m}^{-2}$); L_g is the net long wave radiation absorbed by the ground ($\text{W}\cdot\text{m}^{-2}$), H_g is the sensible heat flux from the ground ($\text{W}\cdot\text{m}^{-2}$); λE_g is the latent heat flux from the ground ($\text{W}\cdot\text{m}^{-2}$); and Q_R is the energy delivered by rainfall ($\text{W}\cdot\text{m}^{-2}$). When the ground is covered by snow, the heat flux into the uppermost soil layer is calculated as follows:

$$h = I_p + G \quad (12)$$

where I_p is the radiation that penetrates the snow cover, and G is the heat conduction from the bottom snow layer to the uppermost soil layer. Eq (8) is solved using a finite difference scheme with an hourly time step, similar to the solution of Eq (4).

There are no available observations of the geothermal heat flux for the northeastern QTP. To simulate the permafrost we consider an underground depth of 50 m. We assume an upward thermal heat flux at the bottom boundary and estimate its value to be $0.14 \text{ W}\cdot\text{m}^{-2}$ at a depth of 50 m using the average geothermal gradient from the 4 boreholes (T1-T4) shown in Figure 3, which is reasonable based on a comparison with the observations ($0.02 \text{ W}\cdot\text{m}^{-2}$ to $0.16 \text{ W}\cdot\text{m}^{-2}$) from the interior of the QTP (Wu et al., 2010).

The vertical soil column is divided into 39 layers in the model (~~see~~ Figure 2). The 1.7 m topsoil layer is subdivided into 9 layers. The first soil layer at the surface is 0.05 m, and the ~~soil~~ layer thicknesses increases with depth linearly from 0.05 m to 0.3 m at a depth of 0.8 m. ~~Then~~ thicknesses ~~decreases~~ linearly with depth ~~to reaching~~ 0.1 m at a depth of 1.7 m. From 1.7 m to 3.0 m, There ~~there~~ are 12 soil layers with a constant thickness of 0.1 m ~~from 1.7 m to 3.0 m~~ to try to ~~replicate~~ capture the maximum freezing depths according to field observations. From the depth of 3 m to 50 m, there are 18 layers with thicknesses increasing exponentially from 0.1 m to 12 m. The liquid soil moisture, ice content, and soil temperature of each layer is calculated at each time step. The soil heat capacity and soil thermal conductivity are estimated using the method developed by Farouki (1981). Permafrost is defined as ground with a temperature at or below 0 °C for at least two consecutive years (Woo, 2012). This study differentiated permafrost from seasonally frozen ground based on the simulated vertical ground temperature profile in each grid. For each year in each grid, the frozen ground condition was determined by searching the ground temperature profile within a four-year window from the previous three years to the current year.

3.3 Model calibration

To initialize the model, we first estimated the soil temperature profiles based on the assumption that there is a linear relationship between the ground temperature at a given depth below the surface and elevation. This temperature-elevation relationship is estimated from the observed ground temperatures in 6 boreholes (see Figure 1). Next, the model had a 500-year spin up run to specify the initial values of the hydrological

variables (e.g., soil moisture, soil ice content, ground temperature, and groundwater table) by repeating the atmospheric forcing data from 1961 to 1970.

This study used the period of 2002 to 2006 for model calibration and the period of 2008 to 2012 for model validation. The daily ground temperature at the Qilian station and the frozen depths at the Qilian and Yeniugou stations were used to calibrate the ground surface reflectance according to vegetation type. The other parameters, such as groundwater hydraulic conductivity, were calibrated according to the observed baseflow discharge in the winter season at the Qilian, Zhamashike and Yingluoxia stations. The Nash-Sutcliffe efficiency and relative error are calculated using observed and simulated discharge to evaluate the model performance. We calibrated the surface retention capacity and surface roughness to match the observed flood peaks, and calibrated the leaf reflectance, leaf transmittance and maximum Rubisco capacity of the top leaf based on the remote sensing evapotranspiration data. Table 1 shows the major parameters used in the model.

We also simulated the hydrological processes without the frozen soil scheme in order to investigate the impact of frozen soils. In this case, the phase transition of soil water between the solid and the liquid is not considered, although the ground temperature is still simulated. Other processes are simulated in the same manner as in the normal run.

4. Results

4.1 Validation of the hydrological model

We conducted a comprehensive validation of the GBEHM using the ground temperature profiles observed from six boreholes, the long-term observations of ground

temperature and frozen depths from the Qilian and Zhamashike stations, the soil moisture observations from the A'rou Sunny Slope station, the long-term observations of streamflow from the three hydrological stations shown in Figure 1 and the monthly actual evapotranspiration estimated from remote sensing data.

Figure 3 shows the comparison of the model-simulated and observed ground temperature profiles at the six boreholes. The model generally captured the vertical distribution of the ground temperature at T1, T2, T3 and T4 in the permafrost area, but the temperatures were overestimated above 20 m depth for T1 and T3. Good agreement between the simulated and observed ground temperature profiles below the depth of 20 m is probably due to fitting of initial values. Therefore, the deep ground temperatures are stable, which is confirmed by the comparison of temperature profiles in different years, as shown in Figure S1 in the supplementary material. Figure S1 also illustrates that the temperatures above 20 m have shown significant increasing trends over the past 40 years. The errors in simulating the vertical temperature profile near the surface might be caused by simplification of the 3-D topography. At T5, which is located in seasonally frozen ground, the simulated ground temperature profile did not agree well with the observed profile at depths of 4-20 m. This error might also be related to heterogeneity in the ground properties, especially the thermal conductivity and heat capacity, since no such information was available. The model simulation agrees well with the borehole observations at T7, which is located in the transition zone from permafrost to seasonally frozen ground. Therefore, the model can identify the boundary between the permafrost and seasonally frozen ground.

We also validated the model simulation of the freezing/thawing cycles based on long-term observations of ground temperature and frozen depth. Figure 4 compares the simulated ground temperature with the observed temperature at the Qilian station, which is located in seasonally frozen ground (observed daily ground temperature data are available from 2004). Generally, the model simulations accurately captured the seasonal changes in the ground temperature profile. Validation of the ground temperature at different depths (0.05 m, 0.1 m, 0.2 m, 0.4 m, 0.8 m, 0.16 m, and 0.32 m) showed that the root mean square error (RMSE) decreases with increasing depth. The RMSE was approximately 2.5 °C for the uppermost three depths (0.05 m, 0.10 m and 0.2 m). The RMSE for depths of 0.4 m and 0.8 m were 1.7 °C and 1.5 °C, respectively, and the RMSE for a depth of 3.2 m was 0.9 °C. Uncertainties in the simulations may be related to the ground heat capacity and thermal conductivity estimated according to Farouki (1981), and the results are similar to the findings by Ou et al. (2016) using the Northern Ecosystem Soil Temperature (NEST) model. We compared the model-simulated daily frozen depth with in situ observations at the Qilian and Yeniugou stations from 2002 to 2014, as shown in Figure 5. The model reproduced well the daily variations in frozen depth, although the depth was underestimated by approximately 0.5 m at the Yeniugou station. In general, the validation of ground temperature and frozen depth indicates that the model effectively captured the freezing and thawing processes in the upper Heihe basin.

Furthermore, we used the the observed hourly liquid soil moisture at the A'rou Sunny Slope station for an additional independent validation. Figure S2 in the supplementary

material shows the comparison between the simulated and observed liquid soil moisture at different depths from January 1 to December 31, 2014. This comparison demonstrates that the model simulation of liquid soil moisture is reasonable.

Figure 6 compares the model simulated and the observed daily streamflow discharge at the Yingluoxia, Qilian and Zhamashike stations. The model simulations agreed well with the observations. The model simulations captured the flood peaks and the magnitude of base flow in both the calibration and validation periods. For the Yingluoxia, Qilian and Zhamashike stations, the Nash-Sutcliffe efficiency (NSE) coefficients were 0.64, 0.63 and 0.72, respectively, in the calibration period and 0.64, 0.60, and 0.73, respectively, in the validation period. The relative error (RE) was within 10% for both the calibration and validation periods (see Figure 6). Figure S3 in the supplementary material shows the comparison of the model-simulated monthly actual evaporation data and the remote sensing-based evaporation data for the entire calibration and validation periods. The GBEHM simulation showed similar temporal variations in actual evapotranspiration compared with the remote sensing based estimation, and the RMSE of the simulated monthly evapotranspiration was 9.1 mm in the calibration period and 7.1 mm in the validation period.

We also compared the model-simulated river discharges with and without the frozen soil scheme. Table S1 in the supplementary material shows that the model with the frozen soil scheme achieves a better simulation of the daily hydrograph than the model without the frozen soil scheme. Figure S4 in the supplementary material shows that the model without the frozen soil scheme overestimates the river discharge in the freezing

season and underestimates flood peaks in the warming season.

4.2 Long-term changes in frozen soils

In the upper Heihe basin, the ground surface starts to freeze in November and begins to thaw in April (Wang et al., 2015a). From November to March, the ground surface temperature is below 0°C in both the permafrost and seasonally frozen ground regions, and precipitation mainly falls in the period from April to October. Therefore, to investigate the changes in frozen soils and their hydrological impact, a year is subdivided into two seasons, i.e., the freezing season (November to March) and the thawing season (April to October). Increasing precipitation and air temperature in the study area in both seasons over the past 50 years were reported in a previous study (Wang et al., 2015b). Compared to the decadal mean for 1971 to 1980, the annual mean air temperature for the 2001 to 2010 period was approximately 1.2 °C higher, with a larger increase in the freezing season (1.4 °C) than in the thawing season (1.1 °C) (Table S2).

Figure 7 shows the changes in the basin-averaged ground temperature in the freezing and thawing seasons. The ground temperature increased in all seasons, especially over the past 30 years. The increasing trend of ground temperature was larger in the freezing season than in the thawing season. In the freezing season (Figure 7(a)), the top layer ground temperature was lower than the deep layer temperature. The linear trend of the top layer (0-0.5 m) ground temperature was 0.49 °C·decade⁻¹ and the trend of the deep layer (2.5-3 m) temperature was 0.32 °C·decade⁻¹. The ground temperature in the deep layer (2.5-3 m) changed from -0.7 °C in the 1970s to approximately 0 °C in the most

recent decade. In the thawing season (Figure 7(b)), the increasing trend of the top layer (0-0.5 m) ground temperature ($0.29\text{ }^{\circ}\text{C}\cdot\text{decade}^{-1}$) was greater than that of the deep layer (2.5-3 m) temperature ($0.22\text{ }^{\circ}\text{C}\cdot\text{decade}^{-1}$). The warming trend was larger in shallow ground layers; this is because the surface heat flux is impeded by the thermal inertia as it penetrates to greater depths.

~~Permafrost is defined as ground with a temperature at or below $0\text{ }^{\circ}\text{C}$ for at least two consecutive years (Woo, 2012). This study differentiated permafrost from seasonally frozen ground based on the simulated vertical ground temperature profile in each grid. For each year in each grid, the frozen ground condition was determined by searching the ground temperature profile within a four-year window from the previous three years to the current year.~~ Figure 8 shows the change in permafrost area during 1971-2013. As shown in Figure 8(a), the permafrost areas decreased by approximately 8.8% (from 5700 km^2 in the 1970s to 5200 km^2 in the 2000s), indicating an evident decrease in the permafrost extent in the upper Heihe basin in the past 40 years.

Figure 8 (b) shows the changes in the basin-averaged maximum frozen depth in the seasonally frozen ground areas and active layer thickness in the permafrost areas. The basin-averaged annual maximum frozen depth showed a significant decreasing trend ($0.032\text{ m}\cdot\text{decade}^{-1}$). In addition, the maximum frozen depth had a significantly negative correlation with the annual mean air temperature ($r = -0.71$). Simulated active layer thickness in the permafrost regions increased ($0.043\text{ m}\cdot\text{decade}^{-1}$), and correlated positively with the annual mean air temperature ($p = 0.005$).

Figure 9 shows the frozen soil distributions in the periods of 1971 to 1980 and 2001

to 2010. Comparing the frozen soil distributions of the two periods, we observed major changes in the frozen soils on sunny slopes at elevations between 3500 and 3900 m, especially in the west tributary, where large areas of permafrost changed into seasonally frozen ground. Figure S5, illustrating the taliks simulated in the period of 2001-2010, shows that ~~the~~ taliks were mainly located on the edge of the permafrost area and ~~the~~ that talik development ~~of taliks~~ was not significant.

Figure 10 shows the monthly mean ground temperatures for areas with elevations between 3300 and 3500 m and over areas with elevations between 3500 and 3700 m in the upper Heihe basin. In the areas with elevations between 3300 and 3500 m located in the seasonally frozen ground region, ~~as shown in~~ (Figure 10(a)), the frozen depth decreased, and the ground temperature in ~~the~~ deep layers (~~with~~ depths greater than 2 m) increased. Figure 10(b) shows that the increase in ground temperature was larger in the area with higher elevation (3500-3700 m). This figure shows that the thickness of the permafrost layer decreased as the ground temperature increased, and the permafrost changed into seasonally frozen ground after 2000. The surface thaw depths changed slowly compared with the ~~depth to the base of permafrost~~ frozen depths as shown in Figure 10, which may be primarily due to the geothermal heat flux. Additionally, the faster increase in the air temperature in the freezing season ($0.41\text{ }^{\circ}\text{C decade}^{-1}$) than in the thawing season ($0.26\text{ }^{\circ}\text{C decade}^{-1}$) may be another reason.

4.3 Changes in the water balance and runoff

Table 2 shows the decadal changes in the annual water balance from 1971 to 2010 based on the model simulation. The annual precipitation, annual runoff and annual

runoff ratio exhibited the same decadal variation; however the annual evapotranspiration maintained an increasing trend starting in the 1970s that was consistent with the rising air temperature and soil warming. Although the actual evapotranspiration increased, the runoff ratio remained stable during the past 4 decades because of the increased precipitation.

Figure 11 and Table 2 show the changes in runoff (both simulated and observed) in different seasons. The model-simulated and observed runoff both exhibited significant increasing trends in the freezing season and in the thawing season. Therefore, the model simulation effectively reproduced the observed long-term changes. In the freezing season, since there was no glacier or snow melting (see Table 2), the runoff was mainly ~~the~~ subsurface flow (groundwater flow and lateral flow from the unsaturated zone). In the thawing season, as shown in Table 2, snowmelt runoff contributed approximately 14% of the total runoff, whereas glacier runoff contributed only a small fraction of the total runoff (approximately 2.2%). Rainfall runoff was the major component of the total runoff in the thawing season, and the runoff increase in the thawing season was mainly due to increased precipitation and snowmelt. As shown in Figure 11, the actual evapotranspiration increased significantly in both seasons due to increased precipitation and ground warming. The ~~increasing-increase trend of in the~~ actual evapotranspiration was greater in the thawing season than in the freezing season.

Figure 12 shows the changes in the basin-averaged annual liquid soil water storage ~~in the top (0-3 m layer)~~ and ~~the~~ groundwater storage. The annual liquid soil water storage ~~of the top 0-3 m~~ showed a significant increasing trend, especially in the most

recent 3 decades. This long-term change in liquid water storage was similar to the runoff change in the freezing season, as shown in Figure 11 (a), exhibiting a correlation coefficient of 0.79. The annual ice water storage in the top 0-3 m soil layers showed a significant decreasing trend due to frozen soil changes. Annual groundwater storage showed a significantly increasing trend especially in the most recent 3 decades, which indicates that the groundwater recharge has increased with ~~the frozen soil~~permafrost degradation.

5. Discussion

5.1 Impact of frozen soil changes on the soil moisture and runoff

We have plotted the long-term changes in the spatially averaged liquid soil moistures in the areas with elevations between 3300 and 3500 m and in the areas with elevations between 3500 and 3700 m in Figure S6 in the supplementary material. In the seasonally frozen ground at elevations of 3300-3500 m, the liquid soil moisture increased slightly due to the decrease in the frozen depth, ~~as shown in~~ (Figure 10(a)). At elevations of 3500-3700 m, the liquid soil moisture in the deep layer increased significantly since the 1990s, due to the change of the permafrost into seasonally frozen ground, ~~as shown in~~ (Figure 10 (b)).

In the freezing season, since the surface ground is frozen, runoff is mainly subsurface flow coming from ~~the~~ seasonally frozen ground. Runoff has the highest correlation ($r = 0.82$) with the liquid soil moisture in the freezing season, which indicates that the frozen soil changes were the primary cause of the increased in liquid soil moisture, resulting in increased runoff in the freezing season. During the past 40 years, parts of ~~the~~

permafrost changed into seasonally frozen ground and the frozen depth of the seasonally frozen ground decreased, leading to increases in the liquid soil moisture in the deep layers during the freezing season. The increase in liquid soil moisture also increased the hydraulic conductivity, which enhanced the subsurface flow. Figure 13(c) shows the seasonal pattern of runoff from the entire basin. From April to October (the thawing season), runoff in the permafrost area was much larger than in the area of seasonally frozen ground; however, in the freezing season the inverse was true~~runoff in the permafrost area was lower than in the seasonally frozen ground~~. Figure S7 in the supplementary material shows runoff changes from a typical area (with elevations of 3500-3700 m) that featured permafrost during the period of 1971 to 1980, ~~and but~~ degraded~~that changed~~ to seasonally frozen ground during the period of 2001 to 2010. This illustrates that ~~the~~-thawing of ~~the~~-permafrost increased ~~the~~-runoff in the freezing season and slowed hydrological recession processes in autumn. Figure S4 illustrates the increase in freezing season runoff and the shift in the seasonal flow patterns simulated by the model without the frozen soil scheme.

Figure 13 shows the large difference in runoff variation with elevation between the freezing and thawing seasons. In the freezing season, the runoff change from the 1970s to the 2000s in the areas of seasonally frozen ground (mainly located below 3500 m, see Figure 9) was relatively small. The areas with elevations of 3500-3900 m showed large changes in runoff. This pattern is due to the shift from permafrost to seasonally frozen ground in some areas in the elevation range of 3500 to 3900 m, as simulated by the model, particularly for sunny hillslopes (see Figure 9). This finding illustrates that

a change from permafrost to seasonally frozen ground has a larger impact on the runoff than a change in frozen depth in areas of seasonally frozen ground. In the thawing season, runoff increased with elevation due to the increase in precipitation with increasing elevation, and the magnitude of the runoff increase was mainly determined by magnitude of the precipitation increase (Gao et al., 2016). Precipitation in the region with elevations below 3100 m was low, and the air temperature was high. Hence, runoff in this region was lower during 2001-2010 than during 1971-1980 because of greater evapotranspiration.

5.2 Comparison with the previous similar studies

In this study, the model simulation showed that the thawing of frozen soils led to increased freezing season runoff and base flow in the upper Heihe basin. This result is consistent with previous findings based on observations in high latitude regions (Walvoord and Striegl, 2007; Jacques and Sauchyn, 2009; Ye et al., 2009) and in northeast China (Liu et al., 2003). However, those studies did not consider spatial variability. This study found that the impact of the frozen soil thawing on runoff varied regionally. In the upper Heihe basin (see Figure 13), the change in the freezing season runoff was strongly affected by the change from permafrost to seasonally frozen ground in the higher-elevation region and by the evaporation increase in the lower-elevation region due to rising air temperature. However, runoff at the basin scale mainly came from the higher-elevation regions.

This study also showed that the thawing of frozen soils increased the liquid soil moisture in the upper Heihe basin, which is consistent with the finding of Subin et al.

(2013) using the CLM model to simulate northern high-latitude permafrost regions, and the findings of Cuo et al. (2015) using the VIC model to simulate 13 sites on the QTP. In contrast, Lawrence et al. (2015) found that permafrost thawing reduced soil moisture based on CLM model simulations of the global permafrost region. This finding might be related to the uncertainties in the soil water parameters and the high spatial heterogeneity of soil properties, which are difficult to consider in a global-scale model. Subin et al. (2013) and Lawrence et al. (2015) simulated the soil moisture changes in the active layer of permafrost over large areas with coarse spatial resolution. Unlike those studies, this study investigated the spatio-temporal variability in soil moisture using a high spatial resolution and analyzed the impacts of frozen soil changes.

Jin et al. (2009) found decreased soil moisture and runoff due to permafrost degradation based on observations at the plot scale in the source area of the Yellow River basin. These results are different from those in the present study, possibly due to the difference in the hydrogeological structure and soil hydraulic parameters between the source area of the Yellow River and the upper Heihe basin. Wang et al. (2015a) estimated the increasing trend of the maximum frozen depth in the seasonally frozen ground to be $0.04 \text{ m} \cdot \text{decade}^{-1}$ during 1972-2006 in the Heihe River basin based on plot observations, which is consistent with the results in this study. The increase in groundwater storage illustrated in this study is also consistent with the findings of Cao et al. (2012) based on GRACE data, which showed that groundwater storage increased during the period of 2003~2008 in the upper Heihe basin.

5.3 Uncertainty in simulation of the frozen soils

Estimation of the change in permafrost area is a great challenge due to such complex factors as climatology, vegetation, and geology. Guo et al. (2013) reported that the permafrost area for the whole QTP decreased from approximately $175.0 \times 10^4 \text{ km}^2$ in 1981 to $151.5 \times 10^4 \text{ km}^2$ in 2010, with a relative change of 13.4%. Wu et al. (2005) reported that the permafrost area decreased by 12% from 1975 to 2002 in the Xidatan basin of the QTP based on a ground penetration radar survey. Jin et al. (2006) found an area reduction of 35.6% in island permafrost in Liangdaohe, which is located along the southern portion of the Qinghai-Tibet Highway, from 1975 to 1996. Compared with the borehole observations by Wang et al. (2013) shown in Figure 2, our model slightly overestimated the soil temperature in permafrost areas, possibly leading to an overestimation of the rate of permafrost area reduction.

There were two major uncertainties in the frozen soils simulation: uncertainty in the simulation of the land surface energy balance and uncertainty in the simulation of the soil heat-water transfer processes (Wu et al., 2016). Uncertainty in the land surface energy balance simulation might result from uncertainty in the radiation and surface albedo estimates due to the complex topography, vegetation cover and soil moisture distribution, thereby introducing uncertainties into the estimated ground temperature and soil heat flux. The uncertainty in the simulation of soil heat-water transfer processes might result from the soil water and heat parameters and the bottom boundary conditions of heat flux. For example, the soil depth and the fraction of rock in soil can greatly affect the ground temperature simulation. Permafrost degradation is closely related to the thermal properties of rocks and soils, the geothermal flow and the initial

ground temperature and soil ice conditions. Sub-grid topography may also affect the frozen soil simulation. For example, active layer thickness is different between the low-elevation valleys and higher-elevation slopes due to the temperature inversion caused by the accumulation of cold air in valleys (Bonnaventure et al., 2012; Zhang et al., 2013; O'Neill et al., 2015). ~~In areas with high groundwater flow rates, laterally advected heat flux may increase the thawing of permafrost~~ ~~the laterally advected heat flux may increase the thawing of permafrost, especially in areas with high groundwater flow rates~~ (Kurylyk et al., 2016; Sjöberg et al., 2016). Not considering the lateral heat flux may lead to an underestimation of talik development and thawing rates of permafrost. In addition, uncertainties in the input data, particularly solar radiation (which is estimated using interpolated sunshine hour data from a limited number of observational stations) and precipitation (which is also interpolated based on observations at these stations), may also influence the results of the model simulation. Due to the complexity of the distributed model and the large number of model parameters, quantifying the overall simulation uncertainty is challenging. ~~This but is part of our ongoing research work will be done in a future study.~~

6. Conclusions

This work carefully validated a distributed hydrological model coupled with cryospheric processes in the upper Heihe River basin using available observations of soil moisture, soil temperature, frozen depth, actual evaporation and streamflow discharge. Based on the model simulations from 1971 to 2013 in the upper Heihe River, the long-term changes in frozen soils were investigated, and the effects of the frozen

soil changes on the hydrological processes were explored. Based on these analyses, we have reached the following conclusions:

(1) The model simulation suggests that 8.8% of the permafrost areas degraded into seasonally frozen grounds in the upper Heihe River basin during 1971-2013, predominantly between elevations of 3500 m and 3900 m. The results indicate that the decreasing trend of the annual maximum frozen depth of the seasonally frozen ground is $0.032 \text{ m} \cdot \text{decade}^{-1}$, which is consistent with previous observation-based studies at the plot scale. Additionally, our work indicates ~~a trend of increasing that the increasing trend of~~ active layer thickness in the permafrost regions ~~is of~~ $0.043 \text{ m} \cdot \text{decade}^{-1}$.

(2) The model-simulated runoff trends agree with the observed trends. In the freezing season (November-March), based on the model simulation, runoff was mainly sourced from subsurface flow, which increased significantly in the higher elevation regions where significant frozen soil changes occurred. This finding implies that the runoff increase in the freezing season is primarily caused by frozen soil changes (permafrost degradation and reduced seasonally frozen depth). In the thawing season (April-October), the model simulation indicates that runoff was mainly sourced from rainfall and showed an increasing trend at higher elevations, which can be explained by the increase in precipitation. In both the freezing and thawing seasons, the model-simulated runoff decreased in the lower-elevation regions, which can be explained by increased evaporation due to rising air temperatures.

(3) The model-simulated changes in soil moisture and ground temperature indicate that the annual storage of liquid water increased, especially in the most recent three

decades, due to frozen soil changes. The annual ice water storage in the top 0-3 m of soil showed a significant decreasing trend due to soil warming. The model simulated annual groundwater storage had an increasing trend, which is consistent with the changes observed by the GRACE satellite. Therefore, groundwater recharge in the upper Heihe basin has increased in recent decades.

(4) The model simulation indicated that regions where permafrost changed into seasonally frozen ground had larger changes in runoff and soil moisture than the areas covered by seasonally frozen ground throughout the study period.

For a better understanding of the changes in frozen soils and their impact on ecohydrology, the interactions among soil freezing-thawing processes, vegetation dynamics and hydrological processes need to be investigated in future studies. There are uncertainties in simulations of frozen soils and hydrological processes that also warrant further investigation ~~in the future~~.

Acknowledgements: This research was supported by the major plan of “Integrated Research on the Ecohydrological Processes of the Heihe Basin” (Project Nos. 91225302 and 91425303) funded by the National Natural Science Foundation of China (NSFC). The authors would like to thank the editor and reviewers for their constructive suggestions, which have substantially improved the paper. All data cited in this paper are available from the references~~for this paper are properly cited and referred in the reference list~~. The model code with a working example is freely available from our website (<https://github.com/gb03/GBEHM>) or upon request from the corresponding

659 author (yangdw@tsinghua.edu.cn).

660

661 **References**

662 Bartelt P. and M. Lehning: A physical snowpack model for the swiss avalanche warning: Part I :

663 numerical model, Cold Regions Sci. and Tech., 35(3), 123-145, doi: 10.1016/S0165-

664 232X(02)00074-5, 2002.

665 Bonnaventure PP, Lewkowicz AG, Kremer M, Sawada MC: A Permafrost Probability Model for the

666 Southern Yukon and Northern British Columbia, Canada, Permafrost and Periglac. Process., 23,52-

667 68, doi: 10.1002/ppp.1733, 2012.

668 Cao Y., Nan Z. and Hu X.: Estimating groundwater storage changes in the Heihe river basin using

669 GRACE, in: IEEE International Geoscience and Remote Sensing Symposium (IGARSS), Munich,

670 Germany, 22–27 July 2012, 798-801, 2012.

671 Chen, R., Lu, S., Kang, E., Ji, X., Zhang, Z., Yang, Y., Qing, W.: A distributed water-heat coupled model

672 for mountainous watershed of an inland river basin of Northwest China (I) model structure and

673 equations, Environ. Geol., 53, 1299-1309, doi: 10.1007/s00254-007-0738-2,2008.

674 Cheng, G. and Jin, H.: Permafrost and groundwater on the Qinghai-Tibet Plateau and in northeast China,

675 Hydrogeol. J., 21, 5-23, doi: 10.1007/s10040-012-0927-2, 2013.

676 Cheng, G., Li, X., Zhao, W., Xu, Z., Feng, Q., Xiao, S., Xiao, H.: Integrated study of the water-

677 ecosystem-economy in the Heihe River Basin. Nat. Sci. Rev., 1(3), 413-428, doi:

678 10.1093/nsr/nwu017, 2014.

679 Cheng, G., and Wu T.: Responses of permafrost to climate change and their environmental significance,

680 Qinghai-Tibet Plateau, J. Geophys. Res., 112, F02S03, doi:10.1029/2006JF000631, 2007.

681 Cherkauer, K. A., and D. P. Lettenmaier: Hydrologic effects of frozen soils in the upper Mississippi River
682 basin, *J. Geophys. Res.*, 104, 19,599-19,610, doi: 10.1029/1999JD900337, 1999.

683 Cong Z T, Yang D W, Gao B, et al.: Hydrological trend analysis in the Yellow River basin using a
684 distributed hydrological model, *Water Resour Res*, 45: W00A13, doi: 10.1029/2008WR006852,
685 2009.

686 Cuo, L., Y. Zhang, F. Zhu, and L. Liang.: Characteristics and changes of streamflow on the Tibetan
687 Plateau: A review, *J. Hydrol.: Reg. Stud.*, 2, 49-68, doi: 10.1016/j.ejrh.2014.08.004, 2014.

688 Cuo, L., Y. Zhang, T. J. Bohn, L. Zhao, J. Li, Q. Liu, and B. Zhou: Frozen soil degradation and its effects
689 on surface hydrology in the northern Tibetan Plateau, *J. Geophys. Res. Atmos.*, 120,
690 doi:10.1002/2015JD023193, 2015.

691 Duan L., Man X., Kurylyk B. L. and Cai T.: Increasing winter baseflow in response to permafrost thaw
692 and precipitation regime shifts in northeastern China, *Water*, 9(1), 25, doi:10.3390/w9010025, 2017.

693 Endrizzi S, Gruber S, Dall'Amico M, and R. Rigon: GEOTop 2.0: simulating the combined energy and
694 water balance at and below the land surface accounting for soil freezing, snow cover and terrain
695 effects, *Geosci. Model Dev.*, 7: 2831-2857, doi:10.5194/gmd-7-2831-2014, 2014.

696 Fan, W.: Heihe 1km LAI production, Heihe Plan Science Data Center at Lanzhou,
697 doi:10.3972/heihe.090.2014.db, 2014.

698 Farouki, O.T.: The thermal properties of soils in cold regions, *Cold Regions Sci.and Tech.*, 5, 67-75, doi:
699 10.1016/0165-232X(81)90041-0, 1981.

700 Flerchinger G., Saxton K.: Simultaneous heat and water model of a freezing snow-residue-soil system I.
701 Theory and development, *Trans. ASAE*, 32, 565-571, doi: 10.13031/2013.31040, 1989.

702 Gao B., Qin Y., Wang YH, Yang DW, and Zheng YR: Modeling Ecohydrological Processes and Spatial

703 Patterns in the Upper Heihe Basin in China, *Forests*, 7(1),10, doi:10.3390/f7010010, 2016.

704 Guo, D., and H. Wang: Simulation of permafrost and seasonally frozen ground conditions on the Tibetan

705 Plateau, 1981–2010, *J. Geophys. Res. Atmos.*, 118, 5216-5230, doi:10.1002/jgrd.50457, 2013.

706 Hinzman, L.D., C.J. Deal, A.D. McGuire, S.H. Mernild, I.V. Polyakov, and J.E. Walsh: Trajectory of

707 the Arctic as an integrated system, *Ecol. Appl.*, 23(8),1837-1868, doi:10.1890/11-1498.1, 2013.

708 Jacques St., J.-M., and Sauchyn D. J.: Increasing winter baseflow and mean annual streamflow from

709 possible permafrost thawing in the Northwest Territories, Canada, *Geophys. Res. Lett.*, 36, L01401,

710 doi:10.1029/2008GL035822, 2009.

711 Jarvis, A., Reuter, H.I., Nelson, A., Guevara, E.: Hole-filled seamless SRTM data, Version 4,

712 International Centre for Tropical Agriculture (CIAT), 2008.

713 Jin, H., He, R., Cheng, G., Wu, Q., Wang, S., Lu, L. and Chang X.: Changes in frozen ground in the

714 Source Area of the Yellow River on the Qinghai–Tibet Plateau, China, and their eco-environmental

715 impacts, *Environ. Res. Lett.*, 4(4), 045206, doi:10.1088/1748-9326/4/4/045206, 2009.

716 Jin, H.J., Zhao, L., Wang, S.L., Jin, R.: Thermal regimes and degradation modes of permafrost along

717 the Qinghai–Tibet Highway, *Science in China D: Earth Sciences*, 49 (11), 1170-1183, 2006.

718 Jordan R.: A one-dimensional temperature model for a snow cover, Technical Documentation for

719 SNTHERM.89, Cold Regions Research and Engineering Lab, Hanover NH, 49 pp., 1991.

720 Kurylyk, B. L., M. Hayashi, W. L. Quinton, J. M. McKenzie, and C. I. Voss: Influence of vertical and

721 lateral heat transfer on permafrost thaw, peatland landscape transition, and groundwater flow, *Water*

722 *Resour. Res.*, 52, 1286-1305, doi:10.1002/2015WR018057, 2016.

723 Lawrence, D.M., C.D. Koven, S.C. Swenson, W.J. Riley, and A.G. Slater: Permafrost thaw and resulting

724 soil moisture changes regulate projected high-latitude CO₂ and CH₄ emissions, *Environ. Res. Lett.*,

10, doi:10.1088/1748-9326/10/9/094011, 2015.

Li, D.L., Zhong, H.L., Wu, Q.B., Zhang, Y.J., Hou, Y.L., Tang, M.C.: Analyses on changes of surface temperature over Qinghai–Xizang Plateau, *Plateau Meteorology*, 24, 291-298, 2005 (in Chinese).

Li, X., Cheng, G.D., Liu, S.M., Xiao, Q., Ma, M.G., Jin, R., Che, T., Liu, Q.H., Wang, W.Z., Qi, Y., Wen, J.G., Li, H.Y., Zhu, G.F., Guo, J.W., Ran, Y.H., Wang, S.G., Zhu, Z.L., Zhou, J., Hu, X.L., Xu, Z.W.: Heihe Watershed Allied Telemetry Experimental Research (HiWATER): Scientific Objectives and Experimental Design, *B. Am. Meteorol. Soc.*, 94(8), 1145-1160, doi: 10.1175/BAMS-D-12-00154.1, 2013.

Liu, X. and Chen, B.: Climate warming in the Tibetan Plateau during recent decades, *Int. J. Climatol.*, 20(1), 1729-1742, doi: 10.1002/1097-0088(20001130)20:14<1729::AID-JOC556>3.0.CO;2-, 2000.

Liu, S., Xu Z., Wang W., Bai J., Jia Z., Zhu M., and Wang J.: A comparison of eddy-covariance and large aperture scintillometer measurements with respect to the energy balance closure problem, *Hydrol. Earth Syst. Sc.*, 15(4), 1291-1306, doi:10.5194/hess-15-1291-2011, 2011.

Liu J., N. Hayakawab, Lu M., Dong S., and Yuan J.: Hydrological and geocryological response of winter streamflow to climate warming in Northeast China, *Cold Regions Sci.and Tech.*, 37,15-24, doi: 10.1016/S0165-232X(03)00012-0, 2003.

Wu T., Li S., Cheng G. and N Z.: Using ground-penetrating radar to detect permafrost degradation in the northern limit of permafrost on the Tibetan Plateau, *Cold Regions Sci.and Tech.*, 41, 211-219, 2005, doi:10.1016/j.coldregions.2004.10.006.

Niu L., Ye B., Li J., and Sheng Y.: Effect of permafrost degradation on hydrological processes in typical basins with various permafrost coverage in Western China, *China Earth Sci.*, 54(4), 615-624, doi: 10.1007/s11430-010-4073-1, 2011.

Oerlemans, J. and Knap, W.H.: A 1 year record of global radiation and albedo in the ablation zone of Morteratschgletscher, Switzerland, *J. Glaciol.*, 44, 231-238, doi: 10.3198/1998JoG44-147-231-238,

1998.

Oerlemans, J.: *Glaciers and Climate Change*, Lisse: Swets & Zeitlinger, 2001.

Oleson, K.W., Lawrence, D.M., Bonan, G.B., Flanner, M.G., Kluzek, E., Lawrence, P.J., Levis, S., Swenson, S.C., Thornton, P.E., Dai, A., Decker, M., Dickinson, R., Feddema, J., Heald, C.L., Hoffman, F., Lamarque, J., Mahowald, N., Niu, G., Qian, T., Randerson, J., Running, S., Sakaguchi, K., Slater, A., Stöckli, R., Wang, A., Yang, Z., Zeng, X., Zeng, X.: Technical Description of version 4.0 of the Community Land Model (CLM), NCAR Technical Note NCAR/TN-47+STR, National Center for Atmospheric Research, Boulder, CO, 257 pp., 2010.

Qiu J.: Thawing permafrost reduces river runoff, *Nature*, doi:10.1038/nature.2012.9749, 2012.

Ou, C., B. Leblon, Y. Zhang, A. LaRocque, K. Webster, and J. McLaughlin: Modelling and mapping permafrost at high spatial resolution using Landsat and RADARSAT images in northern Ontario, Canada: Part 1 - Model calibration, *International Journal of Remote Sensing*, doi: 10.1080/01431161.2016.1157642, 2016.

O'Neill, H. B., Burn, C. R., Kokelj, S. V. & Lantz, T. C.: 'Warm' tundra: atmospheric and near-surface ground temperature inversions across an alpine treeline in continuous permafrost, western arctic, Canada. *Permafrost and Periglac. Process.* 26, 103–118, doi: 10.1002/ppp.1838., 2015.

Rawlins M., Lammers R., Frolking S., Fekete B. and Vorosmarty C.: Simulating pan-Arctic runoff with a macro-scale terrestrial water balance model, *Hydrol. Process.*, 17, 2521-2539, doi: 10.1002/hyp.1271, 2003.

Rawlins, M.A., D.J. Nicolsky, K.C. McDonald, and V.E. Romanovsky: Simulating soil freeze/thaw dynamics with an improved pan-Arctic water balance model, *J. Adv. Model. Earth Syst.*, 5:659-675, doi:10.1002/jame.20045, 2013.

Rigon R., Bertoldi G., and Over TM: *GEOtop: A distributed hydrological model with coupled water*

772 and energy budgets, *J. Hydrometeorol.*, 7, 371–388, doi: 10.1175/JHM497.1, 2006.

773 Schuur, E.A.G., A.D. McGuire, C. Schädel, G. Grosse, J.W. Harden, D.J. Hayes, G. Hugelius, C.D.

774 Koven, P. Kuhry, D.M. Lawrence, S.M. Natali, D. Olefeldt, V.E. Romanovsky, K. Schaefer, M.R.

775 Turetsky, C.C. Treat, and J.E. Vonk: Climate change and the permafrost carbon feedback, *Nature*

776 520,171-179 doi:10.1038/nature14338, 2015.

777 Sellers, P.J.: Canopy reflectance, photosynthesis, and transpiration, *Int. J. Remote Sens.*, 8, 1335-1372,

778 doi: 10.1080/01431168508948283, 1985.

779 Sellers, P.J.; Randall, D.A.; Collatz, G.J.; Berry, J.A.; Field, C.B.; Dazlich, D.A.; Zhang, C.; Collelo,

780 G.D.; Bounoua, L.: A Revised Land Surface Parameterization (SiB2) for Atmospheric GCMS.

781 Part I: Model Formulation, *J. Clim.*, 9, 676-705, doi: 10.1175/1520-

782 0442(1996)009<0676:ARLSPF>2.0.CO;2, 1996.

783 Sjöberg, Y., E. Coon, A. B. K. Sannel, R. Pannetier, D. Harp, A. Frampton, S. L. Painter, and S. W. Lyon:

784 Thermal effects of groundwater flow through subarctic fens: A case study based on field

785 observations and numerical modeling, *Water Resour. Res.*, 52, 1591-1606,

786 doi:10.1002/2015WR017571, 2016.

787 Song X., Brus DJ, Liu F., Li D., Zhao Y., Yang J. and Zhang G.: Mapping soil organic carbon content by

788 geographically weighted regression: A case study in the Heihe River Basin, China, *Geoderma*,

789 261,11–22, doi: 10.1016/j.geoderma.2015.06.024, 2016.

790 Strahler A N: Quantitative analysis of watershed geomorphology, *Eos, Transactions American*

791 *Geophysical Union*, 38(6), 913-920, doi: 10.1029/TR038i006p00913, 1957.

792 Subin Z.M., Koven C.D., Riley W.J., Torn M.S., Lawrence D.M. and Swenson S.C.: Effects of Soil

793 Moisture on the Responses of Soil Temperatures to Climate Change in Cold Regions, *J. Clim.*,

794 26,3139-3158, doi: 10.1175/JCLI-D-12-00305.1, 2013.

795 Toon, O.B., McKay, C.P., Ackerman, T.P., and Santhanam, K.: Rapid calculation of radiative heating
796 rates and photodissociation rates in inhomogeneous multiple scattering atmospheres, *J. Geophys.*
797 *Res.* 94(D13), 16,287-16,301, doi: 10.1029/JD094iD13p16287, 1989.

798 Walter W. Immerzeel, Ludovicus P. H. van Beek, and Marc F. P. Bierkens.: Climate Change Will Affect
799 the Asian Water Towers, *Science*, 328, 1382-1385, doi: 10.1126/science.1183188, 2010.

800 Walvoord, M. A. and B. L. Kurylyk: Hydrologic Impacts of Thawing Permafrost—A Review, *Vadose*
801 *Zone J.*, doi:10.2136/vzj2016.01.0010, 2016.

802 Walvoord, M. A., and R. G. Striegl: Increased groundwater to stream discharge from permafrost thawing
803 in the Yukon River basin: Potential impacts on lateral export of carbon and nitrogen, *Geophys. Res.*
804 *Lett.*, 34, L12402, doi:10.1029/2007GL030216, 2007.

805 Wang, L., Koike, T., Yang, K., Jin, R., Li, H.: Frozen soil parameterization in a distributed biosphere
806 hydrological model, *Hydrol. Earth Syst. Sc.*, 14(3), 557-571, doi: 10.5194/hess-14-557-2010, 2010.

807 Wang Q., Zhang T., Wu J., et al.: Investigation of permafrost distribution over the upper reaches of the
808 Heihe River in the Qilian Mountains, *Journal of Glaciology and Geocryology*, 35(1), 19-29, 2013
809 (in Chinese).

810 Wang Q., Zhang T., Peng X., Cao B., and Wu Q.: Changes of soil thermal regimes in the Heihe River
811 Basin over Western China, *Arct., Antarct., and Alpine Res.*, 47(2), 231-241, doi:
812 10.1657/AAAR00C-14-012, 2015a.

813 Wang, Y., Yang, D., Lei, H. and Yang, H.: Impact of cryosphere hydrological processes on the river runoff
814 in the upper reaches of Heihe River, *J. Hydraul. Eng.*, 46, 1064-1071, 2015b (In Chinese).

815 Wang, Y., Yang, H., Yang, D., Qin Y., Gao B. and Cong ZT.: Spatial interpolation of daily precipitation

816 in a high mountainous watershed based on gauge observations and a regional climate model
817 simulation, *J. Hydrometeorol.*, 18, 845-862, 2017, doi: 10.1175/JHM-D-16-0089.1.

818 Woo, M.-K., Kane, D. L., Carey, S. K. and Yang, D.: Progress in permafrost hydrology in the new
819 millennium, *Permafrost Periglac. Process.*, 19, 237-254, doi:10.1002/ppp.613, 2008.

820 Woo M K.: *Permafrost Hydrology*, Springer-Verlag, Berlin Heidelberg, 2012.

821 Wu, B.F., Yan, N.N., Xiong, J., Bastiaanssen, W., Zhu, W.W., Stein, A.: Validation of ETWatch using
822 field measurements at diverse landscapes: A case study in Hai Basin of China. *J. Hydrol.*, 436, 67-
823 80, doi: 10.1016/j.jhydrol.2012.02.043, 2012.

824 Wu, B.F.: *Monthly Evapotranspiration Datasets (2000–2012) with 1 km Spatial Resolution over the*
825 *Heihe River Basin, Heihe Plan Science Data Center at Lanzhou, China*, doi:
826 10.3972/heihe.115.2013.db, 2013.

827 Wu, M., Jansson P. E., Tan X., Wu J., and Huang, J.: Constraining parameter uncertainty in simulations
828 of water and heat dynamics in seasonally frozen soil using limited observed data, *Water*, 8(2), 64,
829 doi:10.3390/w8020064, 2016.

830 Wu, Q., Zhang T. and Liu Y.: Permafrost temperatures and thickness on the Qinghai-Tibet Plateau, *Glob.*
831 *Planet. Change*, 72, 32-38, doi: 10.1016/j.gloplacha.2010.03.001, 2010.

832 Yang, D.W., Gao, B., Jiao, Y., Lei, H.M., Zhang, Y.L., Yang, H.B., Cong, Z.T.: A distributed scheme
833 developed for eco-hydrological modeling in the upper Heihe River, *China Earth Sci.*, 58(1), 36-45,
834 doi: 10.1007/s11430-014-5029-7, 2015.

835 Yang, D.W., Herath, S., and Musiak, K.: Development of a geomorphology-based hydrological model
836 for large catchments, *Annu. J. Hydraul. Eng.*, 42, 169-174, doi: 10.2208/prohe.42.169, 1998.

837 Yang, D.W., Herath, S., and Musiak, K.: A hillslope-based hydrological model using catchment area

838 and width functions, Hydrol. Sci. J., 47, 49-65, doi: 10.1080/02626660209492907, 2002.

839 Yang, M., F. E. Nelson, N. I. Shiklomanov, D. Guo, and G. Wan, Permafrost degradation and its
840 environmental effects on the Tibetan Plateau: A review of recent research, Earth Sci. Rev., 103, 31-
841 44, doi: 10.1016/j.earscirev.2010.07.002, 2010.

842 Ye, B., D. Yang, Z. Zhang, and D. L. Kane: Variation of hydrological regime with permafrost coverage
843 over Lena Basin in Siberia, J. Geophys. Res., 114, D07102, doi:10.1029/2008JD010537, 2009.

844 Zhao, L., C. L. Ping, D. Q. Yang, G. D. Cheng, Y. J. Ding, and S. Y. Liu: Changes of climate and
845 seasonally frozen ground over the past 30 years in Qinghai-Xizang (Tibetan) Plateau, China, [Glob.](#)
846 [Planet. Change](#)~~Global Planet. Change~~, 43, 19-31, doi: 10.1016/j.gloplacha.2004.02.003, 2004.

847 Zhang, Y.L., Cheng, G.D., Li, X., Han, X.J., Wang, L., Li, H.Y., Chang, X.L., Flerchinger, G.N.:
848 Coupling of a simultaneous heat and water model with a distributed hydrological model and
849 evaluation of the combined model in a cold region watershed, Hydrol. Process., 27(25), 3762-3776,
850 doi: 10.1002/hyp.9514, 2013.

851 Zhang, Y., Ohata, T., and Kadota, T.: Land-surface hydrological processes in the permafrost region of the
852 eastern Tibetan Plateau, J. Hydrol., 283, 41-56, doi: 10.1016/S0022-1694(03)00240-3, 2003.

853 Zhang, Y., X. Wang, R. Fraser, I. Olthof, W. Chen, D. McLennan, S. Ponomarenko, and W. Wu: Modelling
854 and mapping climate change impacts on permafrost at high spatial resolution for an Arctic region
855 with complex terrain, The Cryosphere, 7, 1121-1137, doi:10.5194/tc-7-1121-2013, 2013.

856 Zhou, J.H. and Zheng, Y.R.: Vegetation Map of the upper Heihe basin, Version 2.0, Heihe Plan Science
857 Data Center at Lanzhou, China, doi:10.3972/heihe.426.2014.db, 2014.

858

859 **Figure caption:**

860 Figure 1. The Study area, hydrological stations, borehole observation and flux tower stations.

861 Figure 2. Model structure and vertical discretization of soil column.

862 Figure 3. Comparison of the simulated and the observed soil temperature at borehole observation
863 sites, and the observed data is provided by Wang et al. (2013).

864 Figure 4. Daily soil temperature at the Qilian station: (a) observation; (b) simulation; (c) difference
865 (simulation - observation).

866 Figure 5. Comparison of the simulated and observed daily frozen depths during the period of 2002-
867 2014 at: (a) the Qilian station, (b) the Yeniugou station.

868 Figure 6. Comparison of the simulated and the observed daily river discharge at: (a) the Yingluoxia
869 Gauge, (b) the Qilian Gauge, and (c) the Zhamashike Gauge. For each gauge, the upper and lower
870 panels show the calibration and validation periods, respectively. Nash-Sutcliffe efficiency and
871 relative error coefficients are indicated.

872 Figure 7. Simulated ground temperature changes in: (a) the freezing season (from November to
873 March) (b) the thawing season (from April to October). Results from linear regressions are
874 indicated.~~Simulated ground temperature changes in: (a) the freezing season (from November to~~
875 ~~March) (b) the thawing season (from April to October).~~

876 Figure 8. Change of the frozen soils in the upper Heihe basin: (a) areas of permafrost and basin
877 averaged annual air temperature; (b) the basin averaged annual maximum depths of seasonally
878 frozen ground and thaw above permafrost. Results from linear regressions are indicated.~~Change of~~
879 ~~the frozen soils in the upper Heihe basin: (a) areas of permafrost and basin averaged annual air~~
880 ~~temperature; (b) the basin averaged annual maximum depths of seasonally frozen ground and thaw~~
881 ~~above permafrost.~~

Figure 9. Distribution of permafrost and seasonally frozen ground for two periods: (a) 1971-1980 and (b) 2001-2010; (c) Area where permafrost degraded to seasonally frozen ground from (a) to (b); Percentage of permafrost area with respect to elevation on the (d) sunny and (e) the shaded slopes for the two periods. Note that (d) and (e) share a legend.

Figure 10. Spatially averaged monthly ground temperatures simulated from 1971 to 2013 for two elevation intervals: (a) seasonally frozen ground between 3300 and 3500 m; (b) permafrost that degraded to seasonally frozen ground between 3500 and 3700 m.

Figure 11. Runoff and simulated evapotranspiration in (a) the freezing season and (b) the thawing season. Trend lines are for simulated data and regression results are shown. The upper two panels are for freezing season and the lower two panels are for thawing season.
~~Runoff and simulated evapotranspiration in (a) the freezing season and (b) the thawing season. Trend lines are for simulated data. The up two panels are for freezing season and the lower two panels are for thawing season.~~


Figure 12. Basin averaged annual water storage (equivalent water depth) changes simulated over the period of 1971 to 2013 for: (a) liquid water in the top layer of the ground (0-3 m); (b) ice in the top layer of the ground (0-3 m); (c) and groundwater. Results from linear regressions are indicated.
~~Basin averaged annual water storage (equivalent water depth) changes simulated over the period of 1971 to 2013 for: (a) liquid water in the top layer of the ground (0-3 m); (b) ice in the top layer of the ground (0-3 m); (c) and groundwater.~~

Figure 13. Model simulated runoff showing changes from the 1971-1980 period to the 2001-2010 period with elevation for (a) the freezing season and (b) the thawing season, and (c) monthly averaged seasonal runoff in permafrost and seasonally frozen ground for the period of 2001-2010.

905

906

907



908




Figure 1. The Study area, hydrological stations, borehole observation and flux tower stations.




Figure 2. Model structure and vertical discretization of soil column.




Figure 3. Comparison of the simulated and the observed soil temperature at borehole observation sites, and the observed data is provided by Wang et al. (2013).




Figure 4. Daily soil temperature at the Qilian station: (a) observation; (b) simulation; (c) difference (simulation - observation).






Figure 5. Comparison of the simulated and observed daily frozen depths during the period of 2002-2014 at: (a) the Qilian station, (b) the Yeniugou station.

--- Simulation — Observation — Precipitation


(a) Yingluoxia



(b) Qilian



(c) Zhamashike






Figure 6. Comparison of the simulated and the observed daily river discharge at: (a) the Yingluoxia Gauge, (b) the Qilian Gauge, and (c) the Zhamashike Gauge. For each gauge, the upper and lower panels show the calibration and validation periods, respectively. Nash-Sutcliffe efficiency and relative error coefficients are indicated.




Figure 7. Simulated ground temperature changes in: (a) the freezing season (from November to March) (b) the thawing season (from April to October). Results from linear regressions are indicated.




Figure 8. Change of the frozen soils in the upper Heihe basin: (a) areas of permafrost and basin averaged annual air temperature; (b) the basin averaged annual maximum depths of seasonally frozen ground and thaw above permafrost. Results from linear regressions are indicated.




Figure 9. Distribution of permafrost and seasonally frozen ground for two periods: (a) 1971-1980 and (b) 2001-2010; (c) Area where permafrost degraded to seasonally frozen ground from (a) to (b); Percentage of permafrost area with respect to elevation on the (d) sunny and (e) the shaded slopes for the two periods. Note that (d) and (e) share a legend.




Figure 10. Spatially averaged monthly ground temperatures simulated from 1971 to 2013 for two elevation intervals: (a) seasonally frozen ground between 3300 and 3500 m; (b) permafrost that degraded to seasonally frozen ground between 3500 and 3700 m.





Figure 11. Runoff and simulated evapotranspiration in (a) the freezing season and (b) the thawing season. Trend lines are for simulated data and regression results are shown. The upper two panels are for freezing season and the lower two panels are for thawing season.






Figure 12. Basin averaged annual water storage (equivalent water depth) changes simulated over the period of 1971 to 2013 for: (a) liquid water in the top layer of the ground (0-3 m); (b) ice in the top layer of the ground (0-3 m); (c) and ground-water.

Results from linear regressions are indicated.




Figure 13. Model simulated runoff showing changes from the 1971-1980 period to the 2001-2010 period with elevation for (a) the freezing season and (b) the thawing season, and (c) monthly averaged seasonal runoff in permafrost and seasonally frozen ground for the period of 2001-2010.

987 **Table list:**

988 Table 1. Major parameters of the GBEHM model.

989 Table 2. Changes in annual basin water balance and runoff components in different seasons.

990

991 Table 1. Major parameters of the GBEHM model

Parameters	Coniferous Forest	Shrub	Steppe	Alpine Meadow	Alpine Sparse Vegetation	Desert
Surface retention capacity (mm)	30.0	25.0	10.0	15.0	15.0	5.0
Surface roughness (Manning coefficient)	0.5	0.3	0.1	0.1	0.1	1.0
Soil reflectance to visible light	0.20	0.20	0.20	0.28	0.14	0.11
Soil reflectance to near-infrared radiation	0.225	0.225	0.225	0.28	0.225	0.225
Leaf reflectance to visible light	0.105	0.105	0.105	0.105	0.105	—
Leaf reflectance to near-infrared radiation	0.35	0.58	0.58	0.58	0.58	—
Leaf transmittance to visible light	0.05	0.07	0.07	0.07	0.07	—
Leaf transmittance to near-infrared radiation	0.10	0.25	0.25	0.25	0.25	—
Maximum Rubisco capacity of top leaf ($10^{-5} \text{ mol} \cdot \text{m}^{-2} \cdot \text{s}^{-1}$)	6.0	6.0	3.3	3.3	3.0	—
Plant root depth (m)	2.0	1.0	0.40	0.40	0.1	0.0
Intrinsic quantum efficiency ($\text{mol} \cdot \text{mol}^{-1}$)	0.08	0.08	0.05	0.05	0.05	—
Canopy top height (m)	9.0	1.9	0.3	0.3	0.2	—
Leaf length (m)	0.055	0.055	0.3	0.3	0.04	—
Leaf width (m)	0.001	0.001	0.005	0.005	0.001	—
Stem area index	0.08	0.08	0.05	0.05	0.08	—

992 Table 2. Changes in annual basin water balance and runoff components ($\text{mm} \cdot \text{y}^{-1}$) in different seasons

Decade	P ($\text{mm} \cdot \text{yr}^{-1}$)	E ($\text{mm} \cdot \text{yr}^{-1}$)	Sim R ($\text{mm} \cdot \text{yr}^{-1}$)	Obs R ($\text{mm} \cdot \text{yr}^{-1}$)	Runoff ratio (Obs)	Runoff ratio (Sim)	Runoff components ($\text{mm} \cdot \text{yr}^{-1}$)					
							Freezing season (from November to March)			Thawing season (from April to October)		
							T	G	S	T	G	S
1971-1980	439.1	282.1	154.1	143.8	0.33	0.35	18.5	0.0	0.0	135.6	3.5	13.8
1981-1990	492.8	300.8	188.5	174.1	0.35	0.38	20.5	0.0	0.0	168.0	3.1	27.8
1991-2000	471.0	307.6	161.9	157.4	0.33	0.34	20.5	0.0	0.0	141.4	3.8	18.4
2001-2010	504.3	319.0	180.6	174.3	0.35	0.36	26.2	0.0	0.0	154.3	3.7	24.1

993 Note: P means precipitation, E means actual evaporation, R means runoff, T means total runoff, G
994 means glacier runoff and S means snowmelt runoff, Sim means simulation and Obs means
995 observation.