

Near-surface temperature inversion during summer at Summit, Greenland, and its relation to MODIS-derived surface temperatures

Alden C. Adolph^{1,2}, Mary R. Albert¹, Dorothy K. Hall^{3,4}

¹Thayer School of Engineering, Dartmouth College, Hanover, NH, USA

5 ²Department of Physics, St. Olaf College, Northfield, MN, USA

³Earth System Science Interdisciplinary Center / University of Maryland, College Park, MD, USA

⁴Cryospheric Sciences Laboratory, NASA Goddard Space Flight Center, Greenbelt, MD, USA

Correspondence to: Alden C. Adolph (Adolph1@stolaf.edu)

Abstract

10

As rapid warming of the Arctic occurs, it is imperative that climate indicators such as temperature be monitored over large areas to understand and predict the effects of climate changes. Temperatures are traditionally tracked using in situ 2 m air temperatures and can also be assessed using remote sensing techniques. Remote sensing is especially valuable over the Greenland Ice Sheet where few ground based air temperature measurements exist. Because of the presence of surface-based temperature inversions in ice-covered areas, differences between 2 m air temperature and the temperature of the actual snow surface (referred to as “skin” temperature) can be significant and are particularly relevant when considering validation and application of remote sensing temperature data. We present results from a field campaign extending from 8 June through 18 July 2015, near Summit Station in Greenland to study surface temperature using the following measurements: skin temperature measured by an infrared (IR) sensor, 2 m air temperature measured by a National Oceanic and Atmospheric Administration (NOAA) meteorological station; and a MODerate-resolution Imaging Spectroradiometer (MODIS) surface temperature product. Our data indicate that 2 m air temperature is often significantly higher than snow skin temperature measured in situ, and this finding may account for apparent biases in previous studies of MODIS products that used 2 m air temperature for validation. This inversion is present during summer months when incoming solar radiation and wind speed are both low. As compared to our in situ IR skin temperature measurements, after additional cloud masking, the MOD/MYD11 Collection 6 surface-temperature standard product has an RMSE of 1.0°C and a mean bias of 0.4°C, spanning a range of temperatures from -35°C to -5°C. For our study area and time series, MODIS surface temperature products agree with skin surface temperatures better than previous studies indicated, especially at temperatures below -20°C where other studies found a significant cold bias. We show that the apparent “cold bias” present in other comparisons of 2 m air temperature and MODIS surface temperature may be a result of the near-surface temperature inversion. Further investigation of how in situ IR skin temperatures compare to MODIS surface temperature at lower temperatures (below -35°C) is warranted to determine whether a cold bias exists for those temperatures.

25

30

35 1. Introduction

The Arctic is experiencing warming at a more rapid rate than the rest of the world (Stocker, 2014), but the impacts of this increased temperature extend beyond the polar region. Declining sea ice extent and retreat of glaciers contribute to a powerful ice-albedo feedback that results in further warming on a large scale. This increased warming leads to declining mass balance of the Greenland Ice Sheet, contributing to global sea level rise. Quantifying current and future ice sheet mass balance remains an active area of research (e.g. Rignot et al., 2011; Rae et al., 2012; Vernon et al., 2013) and is critical to improving projections of sea level rise. Declining Greenland Ice Sheet mass balance is driven in part by changes in surface energy balance, which drives surface temperature and surface melt (Box, 2013; van den Broeke et al., 2016). Furthermore, the energy balance at the snow surface controls the interactions between the snow surface and the atmospheric surface layer. The net surface energy balance is defined by the net shortwave and longwave radiation, as well as sensible and latent heat fluxes, and heat flux from the underlying snow and ice. The net radiation at the surface affects the stability of the near-surface atmosphere and the extent to which turbulent heat exchange occurs between the snow surface and the lower atmosphere, impacting both local and regional circulation and climate.

Surface temperature is a critical component for understanding ice sheet mass balance and for tracking changes in surface energy balance, however making accurate measurements of surface temperature across the vast expanse of the Greenland Ice Sheet over a long period of time is challenging (Reeves Eyre and Zeng, 2017). The installation of automatic weather stations (AWS) across the ice sheet has begun to provide point meteorological data at many locations through programs such as Greenland Climate Network (GC-Net) (e.g. Steffen et al., 1996; Steffen and Box, 2001; Shuman et al., 2001) and the Programme for Monitoring of the Greenland Ice Sheet (PROMICE), which monitors both skin and air temperatures (e.g. Ahlstrøm, et al, 2008; van As et al., 2011; Fausto et al., 2012). In addition, thermal infrared satellite remote sensing provides the opportunity to collect surface temperature with large spatial coverage and sub-daily to weekly temporal resolution, depending on cloud conditions. In this study, we will focus on the MODerate-resolution Imaging Spectroradiometer (MODIS) thermal infrared land surface temperature (LST) product.

“Surface” temperatures in climatological studies often refer to 2 m air temperature (Hudson and Brandt, 2005) as it is a standard measurement at meteorological stations around the globe; however, remotely-sensed surface temperatures from satellite-borne sensors in the cryosphere measure the radiometric surface temperature, which is the actual “skin” temperature of the surface at the snow/air interface (Warren and Brandt, 2008). In the polar regions, the high albedo of snow in the visible part of the spectrum means relatively little solar radiation is absorbed, and high emissivity of snow at longer wavelengths as compared to the emissivity of the atmosphere often leads to temperatures at the surface that are lower than the air above, a phenomenon called an inversion. The presence of surface-based inversions in the hundreds of meters of the lower atmosphere in the polar regions has long been established (Sverdrup, 1926) and is defined by measuring temperature at two different heights to determine the magnitude of the temperature difference over the relevant height difference. Inversions have been

characterized in Greenland and the wider Arctic (Reeh, 1989; Kahl, 1990, Overland and Guest, 1991) as well as in Antarctica (Philpot and Zillman, 1970). Conditions that cause inversions are most frequently met in winter when incoming radiation is low. Surface-based inversions have typically been studied with 2m air temperature as the “base” of the inversion and the height of the inversion extending hundreds of meters or more into the atmosphere. However, work by Hudson and Brandt (2005) demonstrated the presence of a surface-based temperature inversion below 2 m in the winter of 2001 at South Pole in Antarctica, showing that the largest temperature gradient was in the 20 cm nearest to the snow surface. Good (2016) presents measurements of skin temperature and 2 m air temperature, and finds that at polar sites, during snow-covered seasons in fall, winter, and spring, these two temperatures generally agree well, with the caveat that there is a reduced amplitude of diurnal cycle temperatures at 2 m.

Hall et al. (2008) analysed 2 m air temperature data and skin temperature data from across Greenland and discussed conditions that lead to near-surface thermal stratification over snow-covered areas. Incoming solar irradiance and wind speed are two major controls on thermal stratification. Temperature inversions occur when the incoming solar irradiance is small (i.e. during night) and the snow surface emits longwave radiation; the net radiation at the surface is negative, causing heat transport from the air to the snow surface. The opposite phenomenon of temperature lapse can occur when there is significant incoming solar irradiance resulting in net positive radiation at the surface, with higher temperatures closer to the ground surface and upward heat transport from the snow surface to the air. Winds can serve to neutralize these temperature gradients by mixing air masses.

In recent years, studies have been conducted on surface energy balance and near-surface processes in Greenland (e.g. Miller et al., 2013; 2015; 2017; Berkelhammer et al., 2016) and Antarctica (e.g. van As et al., 2005; van den Broeke et al., 2006; Kuipers Munneke et al., 2012). At our study site at Summit, Greenland, Miller et al. (2013) studied the inversions over two years but consider the 2 m air temperature to be the base of these inversions, and they did not investigate the surface processes beneath 2 m height. They find that inversions are prevalent in winter months and are less intense during summer months and that the presence of clouds results in weaker inversions. In Miller et al. (2015) the impact of clouds on the surface energy budget at Summit is further investigated, and the warming effect of clouds on 2 m air temperatures is shown in all seasons. Details of the Summit, Greenland surface energy balance are extensively documented in Miller et al. (2017). Berkelhammer et al. (2016) discuss the impacts of the surface-based temperature inversions on boundary-layer dynamics, showing that the stability of the atmosphere prevents mixing and ultimately limits accumulation at Summit. These recent studies have investigated near-surface processes at Summit because of the importance of surface energy balance and turbulent snow/atmosphere exchange in climate monitoring and ultimately prediction of larger scale circulation and future change in ice mass balance. Though some surface temperature measurements at Summit have been made (Berkelhammer et al. 2016), controls on surface temperature gradients in the lowest 2 meters of the atmosphere, which are most relevant for the remote sensing community and also have important implications for changing ice sheet dynamics, have not been explicitly studied at Summit, Greenland.

100 In remote sensing validation studies or use of remotely sensed temperatures, this distinction between 2 m air
temperature and skin temperature is important and has been demonstrated in polar regions. In work using satellite data to study
warming trends in the Arctic, Comiso (2003) presents a dataset from an Arctic sea ice study showing correlation between 2 m
air temperature and skin temperature that had been averaged monthly. Over sea ice, there was an average offset of 0.34°C
105 negative offset, perhaps due to inversions that are not well understood. Indeed, best practices for thermal remote sensing
validation indicate that ground-based radiance measurements that yield a skin temperature provide the best validation of remote
sensing land surface temperature products (Guillevic et al., 2017). Because these data have not always been available, previous
studies have used a variety of measurement types for remote sensing surface temperature validation.

A number of validation studies present results acquired over various time scales and in different locations to determine
110 the accuracy of the MODIS surface temperature products in the cryosphere (Hall et al., 2004, 2008; Koenig and Hall, 2010;
Westermann et al., 2012; Hachem et al., 2012; Shuman et al., 2014; Østby et al., 2014; Shamir and Georgakakos, 2014; Hall
et al., 2015; Williamson et al., 2017). Table 1 provides summary statistics related to the results of many of these validation
studies and is discussed in further detail in the discussion section. Overall, a negative bias is present in nearly all validation
studies, where the MODIS surface temperature is lower than the measured skin or 2 m air temperatures, and this bias is
115 particularly prevalent at temperatures below -20°C. Some studies (e.g., Hall et al., 2004, 2008; Shuman et al., 2014) use 2 m
air temperature to validate the MODIS surface temperature products, which may be part of the reason for the biases that are
consistently present. Shuman et al. 2014 acknowledge that differences between 2 m air temperature and skin temperature
caused by inversions could cause bias in their comparison to MODIS, but at the time there was insufficient data to suggest
whether inversions would persist in central Greenland and in the very near-surface. Other studies use thermochrons, either
120 shielded (e.g., Hall et al., 2015) or during darkness (Koenig and Hall, 2010). However, Westermann et al. (2012) and Østby
et al. (2014) both use pyrometers to measure thermal longwave radiation and estimate surface (skin) temperature, and these
studies also find a cold bias in the MODIS surface temperatures. Østby et al. (2014) indicate that this bias is present at lower
temperatures during the winter (and that there is a slight warm bias in the MODIS temperatures during summer), whereas
Westermann et al. (2012) show a cold bias at higher temperatures. Identifying if and when this bias is indeed present is critical
125 to the use of the MODIS surface temperature products over the ice sheet. We hypothesize that a cold bias between 2 m air
temperature and skin surface temperature could be indicative of physical processes of temperature inversion and not any issue
of MODIS instrument calibration, and coupled datasets can be used to further develop our understanding of temperature
processes in polar regions.

In the summer of 2015, we conducted a field campaign near Summit Station, Greenland to measure skin and near-
130 surface air temperature to study near-surface thermal stratification and determine its impact in validation of the MODIS land
surface temperature product. We use our original dataset to determine how summertime meteorological conditions impact
near-surface inversions (beneath 2 m height) on the ice sheet at Summit. Furthermore, we provide a validation of MODIS land
surface temperatures, and show that the use of 2 m air temperature for MODIS validation is not recommended due to the

presence of near-surface inversions. Lastly, we use in situ cloud data to show that the accuracy of the MODIS surface
135 temperature product could be improved through stricter cloud masking.

2. Methods

2.1. In situ Measurements

To characterize snow skin temperature, an autonomous measurement station was installed approximately 10 km
140 NNW of Summit, Greenland (indicated on a map in Figure 1) at an undisturbed site for 40 days between June 8 and July 18,
2015. A Campbell Scientific/Apogee Precision Infrared (IR) Radiometer [Model: SI-111] was used to measure skin
temperature of the snow. The instrument covers the wavelength range from 8 to 14 μm . It has a stated absolute accuracy of
 $\pm 0.5^\circ\text{C}$ from -40°C to -10°C , and $\pm 0.2^\circ\text{C}$ from -10°C to 65°C . The sensor was factory calibrated within several months of its
deployment. The sensor was mounted on a horizontal rod extending approximately 60 cm out from the supporting tripod, and
145 the sensor was approximately 60 cm from the surface, pointed directly downward. The field of view of the sensor is 22° half
angle, so the legs of the tripod did not affect the measurements. Figure 2 shows an image of the sensor setup.

Summit Station was the location of the Greenland Ice Sheet Program 2 (GISP2) deep core site and has operated
continuously as a year-round station for nearly a decade. The National Atmospheric and Oceanic Administration (NOAA) has
operated a meteorological station at Summit, measuring the 2 m air temperature using a shielded Logan PT139 sensor.
150 Additionally, wind speed and incoming solar radiation data were also measured as part of the NOAA station data (NOAA
ESRL Global Monitoring Division, 2017). The data provided by NOAA and used in this paper have a one minute temporal
frequency, and we take a 30 minute average of the data so that the 2 m air temperature is comparable to the IR skin temperature
measurements. Further details of the 2 m air measurements are outlined in Shuman et al. (2014). Additionally, through the
Integrated Characterization of Energy, Clouds, Atmospheric state, and Precipitation at Summit (ICECAPS) project, a number
155 of instruments to monitor cloud, atmosphere, and precipitation were installed at Summit in 2010. One of these instruments is
the millimetre wavelength cloud radar (MMCR), a custom-built Doppler 35 GHz radar that measures reflectivity, mean
Doppler velocity, Doppler spectra, and Doppler spectrum width (data available at <http://www.archive.arm.gov>). More
information about the MMCR can be found in Moran et al. (1998). We use MMCR data in this study to detect the presence of
clouds and determine the accuracy of the MODIS cloud mask, again employing the higher temporal frequency measurements
160 and calculating 30-minute averages so that the data are comparable to our in situ skin temperature measurements.

2.2 Remote Sensing of Surface Temperature with MODIS

There are many different remote sensing instruments that measure radiance in the thermal infrared part of the
electromagnetic spectrum to determine skin temperature, including the Advanced Very High Resolution Radiometer
(AVHRR), the Advanced Thermal Emission and Reflection Radiometer (ASTER), the Enhanced Thematic Mapper Plus
165 (ETM+), and the MODIS. The theoretical basis for determining temperature of a snow surface based on measured thermal
infrared radiance is described by Hook et al. (2007) and Hall et al. (2008) as follows:

$$LS_\lambda = [\epsilon_\lambda L_{bb,\lambda}(T) + (1 - \epsilon_\lambda)L_{sky,\lambda}] \tau_\lambda + L_{atm,\lambda}$$

170 where $L_{s\lambda}$ is the radiance measured by the sensor on a given satellite, ϵ_{λ} is the surface emissivity at a given wavelength, $L_{bb,\lambda}(T)$ is the spectral radiance from a black body as a function of temperature, $L_{sky,\lambda}$ is the spectral downwelling radiance from the atmosphere on the surface, τ_{λ} is the spectral transmittance through the atmosphere, and $L_{atm,\lambda}$ is the spectral radiance upwelling from atmospheric emission and scattering. If emissivity, sky radiance, transmittance, and path radiance are known, surface temperature can be determined through measurements of the radiance at the sensor. In the measurements of snow, the resulting temperature is representative of the top several microns of the surface at the snow/air interface because of the penetration depth of radiation at the wavelengths used, so it is indeed a skin temperature (Warren and Brandt, 2008).

175 The MODIS instrument produces widely-used land surface temperature (LST), which we use as the remote sensing product in this work. This instrument, aboard the Terra and Aqua satellites, has been collecting radiance data from 24 February 2000 to present. The surface temperature products of the Greenland Ice Sheet are used as a baseline to investigate surface temperature trends (e.g. Hall et al. 2012), to monitor melt events on the ice sheet (Hall et al., 2013), and as input for surface mass balance or snowpack modeling (Fréville et al., 2014; Shamir and Georgakakos, 2014; Navari et al., 2016). In this study, we use the MOD/MYD11 Collection 6 product, where MOD refers to the Terra MODIS product and MYD refers to the Aqua MODIS product. This product has a pixel size of 1km x 1 km.

The MOD/MYD11 algorithm was developed to map land surface temperature (Wan and Dozier, 1996; Wan, 2008, 2014) using radiance in MODIS bands 31 and 32, which correspond to a center wavelength of 11 μ m and 12 μ m, respectively. The algorithm used to estimate temperature is referred to as a “split window” technique because the differences between the 11 μ m and 12 μ m bands are used to account for atmospheric effects on the measured radiance. MOD/MYD11 estimates an emissivity value based on land cover (assessed from bands 3-7, 13, and 16-19), presence of water vapor and estimated air temperature atmospheric profiles using MODIS sounding channels (Wan and Dozier, 1996). Emissivity can vary widely because MOD/MYD11 is a global product that estimates land surface temperature on all types of land cover types. Because this study focuses on consistently snow-covered land, there was not significant variability in the emissivity; in band 32 the emissivity is 0.990 for each data point, and in band 31, the emissivity fluctuates between either 0.992 or 0.994 as determined from MOD/MYD11. For cloud masking, MOD/MYD11 uses MOD/MYD35, the standard MODIS cloud mask product which uses data from multiple MODIS bands for cloud detection. This product gives a probability that a pixel is clear. MOD/MYD11 masks out anything below 95% probability of a clear pixel. The accuracy of the MOD/MYD11 product is limited by the uncertainties of radiative modelling, the uncertainty of absorption and scattering coefficients of aerosols and water vapor, and the atmospheric profiles of temperature and water vapor (Wan and Dozier, 1996). For surfaces with a known emissivity, the accuracy of the MOD/MYD11 is within 1°C (Wan, 1999). For further information on the MOD/MYD11 algorithm and associated uncertainties, consult Wan and Dozier (1996) and Wan (1999, 2008, 2014).

200 Previous MODIS surface temperature validation studies have used Collection 5 (C5) products; Collection 6 (C6) products started to become available in 2014. Improvements were made in the C6 MODIS algorithms, most notably to rectify degradation of some sensors on the Terra satellite. However the sensor degradation was largely affecting bands in the visible part of the spectrum and not in the thermal infrared part of the spectrum used to calculate surface temperature (Lyapustin et

al., 2014; Polashenski et al., 2015; Casey et al., 2017). MOD/MYD11 C6 benefits from improved stability of emissivity values and improved algorithms to account for viewing angle over its C5 counterpart (Wan, 2014). Additionally, in C6, the calibration of bands 31 and 32 (used in surface temperature calculation) is improved. Supplemental Figure 1 shows comparisons of C5 and C6 data at our study site over the time period of interest. On average, C6 results in temperatures 0.2°C higher than C5. The temperatures differences are larger at higher temperatures. Finally, the cloud mask algorithms are improved in C6 (Riggs et al., 2017), resulting in a less strict cloud mask over Greenland.

The high latitude location of Summit, Greenland puts it within the field of view of the MODIS instruments on Terra and Aqua multiple times each day. To compare in situ measurements to the temporally coincident MODIS collections, we use swath-level products whose file names contain the UTC time of collection within ± 5 minutes. Within each swath, we find the 1 km x 1 km square pixel in which our measurement site is located by minimizing distance between pixel nadir point and our in situ measurement site. Comparisons between temperatures from the MODIS products and the in situ measurements that are within 30 minutes of one another are used in the analysis. As skin and near-surface air temperatures can fluctuate within a span of 30 minutes, the non-synchronicity may introduce some error in the comparisons, but errors should be random and non-systematic, as 30-minute windows of both increasing and decreasing temperature are included in the analysis.

3. Results and Discussion

3.1. Near-Surface Temperature Measurements

The IR skin temperature measurements operated continuously during the 40-day campaign. The station was visited several times between June 8 and June 25, though no maintenance was required, and then left unmaintained for the remainder of the measurement period. A time series of the IR skin temperature is presented in Figure 3. The snow skin temperature varied between approximately -34°C and -2°C during the measurement period. Gray vertical bars in the figure indicate the presence of clouds as detected by the MMCR radar, and while the diurnal cycles are clear throughout the time series, there is more high frequency fluctuation in temperature during cloudy periods.

Our IR skin temperature measurements are compared in a subset time series to the 2 m air temperature measurements at Summit Station in Figure 4. This time window shows a clear sky period when diurnal cycles are clear and conditions for inversion are most favorable. Thermal stratification in the lowest several meters of the atmosphere is prominently seen in the difference between 2 m air temperature and IR skin temperature (Figure 4). 2 m air temperature and IR skin temperature are similar during peak solar irradiance, with the mean difference in temperature equal to -0.32°C when incoming solar radiation is greater than 600 W m⁻². There is a larger difference between the two during the night-time, with 2 m air temperature higher than skin temperature by an average of 2.4°C when incoming radiation is less than 200 W m⁻². This near-surface inversion is due to low incoming solar radiation and emission of longwave radiation from the snow surface during the night. This stable condition prevents turbulent heat exchange and allows the inversion to persist. Figure 5a shows a direct comparison between the 2 m air temperature measured at the NOAA weather station at Summit and the in situ IR skin temperature measured 10km NNW of Summit. As the inversions appear diurnal in nature, the measurements are quite similar at higher temperatures (above -10°C, mean difference is -0.16°C), but at lower temperatures, there is increased discrepancy between 2 m temperature and

snow skin temperature (below -20°C , mean difference is 3.5°C). Figure 5b shows a histogram of the differences between the same 2 m air temperature and IR skin temperature. There is a clear skew in the histogram, indicating that 2 m air temperature is most frequently higher than skin temperature (in 68% of measurements). This is true in both clear and cloudy sky conditions, where the percentage of measurements for which air temperature exceeds skin temperature is 70% in clear sky conditions and 240 65% in cloudy sky conditions.

Figure 6 shows the magnitude of the temperature difference between 2 m and snow skin temperature as a function of concurrent wind speed, with the color of the marker indicating the concurrent incoming solar radiation. It is clear that increasing wind speed serves to reduce any temperature gradient in the lower meters of the atmosphere, and that at peak solar radiation, there are no inversions present. These differences are much higher at lower wind speeds; a stronger wind shear allows the 245 system to overcome the stability in temperature and promotes heat flux from the air to the snow surface. Weaker winds cannot overcome the temperature stability so the temperature differences persist. Specifically, for the data presented here, at incoming solar radiation above 600 W m^{-2} or wind speeds greater than approximately 7 m s^{-1} , there were no inversions greater than 2°C in the 2 m above the snow surface.

The presence of this near-surface thermal inversion is of particular interest in the context of previous MODIS surface 250 temperature comparison studies. Several studies have used 2 m air temperature to compare to MODIS surface temperature products (Hall et al., 2004, 2008; Shuman et al., 2014). These studies consistently report a "cold bias" in the MODIS surface temperatures (see Table 1), where MODIS surface temperature is lower than concurrently measured 2 m air temperature. In Shuman et al. (2014), a comparison of MOD29 to 2 m air temperature results in a cold bias of approximately 3°C , and the authors note that the disagreement was larger for lower temperatures. Previous studies acknowledge that near-surface 255 stratification may be part of the cause of the discrepancy, but also highlight other potential causes such as issues of calibration of the MODIS instruments at very low (less than approximately -20°C) temperatures (Wenny et al., 2012; Xiong et al., 2015), errors in cloud masking, and potential atmospheric interference. The data presented in Figure 5 show that near-surface thermal stratification may play quite a large role in the discrepancies found between MOD29 and 2 m air temperatures (see Figure 1 of Shuman et al. (2014)). Inversions, which are present during periods of lower incoming solar radiation, and thus frequently 260 lower temperature, result in offsets between skin and 2 m air temperature. Because the MODIS products provide skin temperature (Warren and Brandt, 2008), the difference seen in Shuman et al. (2014) between 2 m air temperature and MODIS temperature at these lower temperatures could in fact be a signature of inversions, which the authors indeed acknowledge but did not have the data to explore. Comparisons of 2 m air temperature to MODIS surface temperature allow us to see how potentially pervasive these inversions could be, though further measurements are needed to determine their presence in non- 265 summer seasons.

Hall et al. (2008) present a figure (their Figure 2) similar to our Figure 5a, in which measured IR skin temperature is plotted vs. 2 m air temperature measured at Summit Station in Greenland from 2000 to 2001. However, they found a consistent offset between 2 m air temperature and skin temperature (of approximately 1°C), a trend that does not vary with temperature. In contrast, our measurements show that the offset is larger at lower temperatures than at higher temperatures and has a much

270 larger magnitude than 1°C; inversions up to 12°C were measured in our data (Figure 5c). The mean differences are reported
above as $-0.16 \pm 0.88^\circ\text{C}$ when temperatures are above -10°C and as $3.5 \pm 2.4^\circ\text{C}$ when temperatures are below -20°C . A paired t-
test shows that these means are not equal to one another with a p-value of less than 0.001. In the summer, inversions are present
only when solar radiation is low, and therefore temperatures are typically low, so discrepancies between 2 m air temperature
and skin temperature only occur during periods of high solar zenith angle. During day time in summer, when there is more
275 incoming radiation and temperatures are typically higher, there is good agreement between measured 2 m air temperature and
skin temperature. Because the Hall et al (2008) data span a longer time scale over all seasons, it is possible that the seasonality
effects of studying only summer are the root of the differences in our results. However, because inversions are known to be
more persistent in the winter than in the summer, we might expect that the trend of larger offsets at lower temperatures would
be even more pronounced when all seasons are included. Future studies, beyond our analysis here, that incorporate all seasons
280 are needed to investigate this discrepancy and determine conditions under which 2 m air temperature is, or is not, a good proxy
for snow skin temperature.

3.2. In situ Temperature Comparisons to MODIS Surface Temperature Products

3.2.1. IR Skin Temperature Comparison

285 Figure 7 shows a time series of a subset of the measurement period with the 30-minute IR skin temperature
measurements overlain with the MOD/MYD11 LSTs. MOD/MYD11 does not provide a surface temperature when the cloud
mask indicates that there are clouds present, which is why there are some gaps in the data (i.e. at day 186/187). Most of the
time series shown in Figure 7 is during a consistently cloudless period. Terra (MOD) passes over Summit several times in the
latter half of the day as temperatures are dropping. Aqua (MYD) passes over Summit as temperatures are typically increasing
290 within the diurnal cycle. The algorithm to calculate temperature from measured radiance is the same in the two different
satellites. Figure 7 shows that there is generally good agreement between IR skin temperature and both MOD11 and MYD11
products. This is also evident in Figure 8a, where MOD/MYD11 products combine to yield an RMSE of 1.6°C ($n=374$) when
compared with IR skin temperature, and there is a mean bias of $-0.7 \pm 1.4^\circ\text{C}$. Separate results for Terra and Aqua are not
significantly different (see Table 1). In contrast to the results from Shuman et al. (2014), there does not seem to be an increase
295 in the difference between MODIS surface temperature and in situ temperature as temperatures decrease.

While we do not believe that 2 m air temperature is a good proxy for skin temperature, for demonstration purposes,
we have compared the 2 m air temperature measurements to the MOD/MYD11 product in Figure 8b. In doing so, we find an
RMSE of 3.1°C and a mean bias of $1.9 \pm 2.5^\circ\text{C}$ ($n=374$). This comparison results in a similar RMSE to Shuman et al. (2014) of
 3.5°C , though the mean bias of our comparison is slightly less than their bias was at 3°C . This comparison further illustrates
300 the importance of using skin temperatures in MODIS validation studies. Shuman et al. (2014) were unable to conclusively say
that any of their bias was a result of using 2 m air temperature instead of skin temperature, and in fact they did not think it was
likely that any inversion effects would cause the gradually increasing bias with decreasing temperature because there was

insufficient research on the presence of near-surface inversions in the dry snow zone in Greenland. The comparison of Figure 8a and 8b shows that at least in the summer, inversions were likely to have played a large role in their 2014 results.

305 As compared to other MODIS validation studies, these results indicate a closer match between in situ measurements and MODIS temperature products as indicated by smaller RMSE and mean bias (see Table 1). While the length of our study is short in comparison to many of the other works referenced, the use of a different in situ sensor is likely a key factor, and there is still a significant range of temperatures captured within our study. In comparing our results to other studies, it is also important to consider that we are using a Collection 6 product, which has seen improvements from previous versions. The
310 Collection 5 cloud mask was more conservative over the Greenland Ice Sheet than is the Collection 6 cloud mask. If we consider only swaths that are considered cloud free by both C5 and C6 (n=341) and compare the MODIS surface temperature to our IR skin temperature, we find that the C6 performs slightly better than C5 with a lower RMSE (C6: 1.44°C, C5: 1.57°C) and lower mean bias (C6: -0.70°C, C5: -0.93°C). The comparisons are shown graphically in Figure S2 of the supplement.

However, there are still some differences between IR skin temperature and MODIS surface temperature in our
315 validation study. To investigate the root of these discrepancies, we consider the sensitivity of the difference between MOD/MYD11 surface temperature and in situ skin temperature as a function of the following parameters: IR skin temperature, solar zenith angle, and sensor viewing angle. These results are presented in Figure 9. The only significant relationship is between temperature difference and MODIS sensor view angle ($p = 0.0029$). The viewing angle varies between 0° and 66°, and the slope of the trend (-0.01°C/°) indicates that at larger viewing angles, there is a larger difference between the MODIS
320 surface temperature and our measured IR skin temperature, but it does not explain much of the variance, as the R^2 value is only 0.02. There is not a significant trend with temperature or with solar zenith angle. As these variables do not explain much of the difference, we believe that the differences may be due to insufficient cloud masking (discussed in the following section) and perhaps to imperfect synchronicity of measurements, where in situ skin measurements represent an average of 30 minutes but the MODIS measurement represents a shorter time window. Yu et al. (1995) suggest that ice crystal precipitation present
325 during inversions may also cause differences between in situ and satellite skin temperatures, though they caused a warm bias rather than a cold bias.

3.2.2. Using In situ Cloud Data to Improve MODIS Surface Temperature

Using the MMCR data from Summit, we identify periods when there were clouds present above Summit Station.
330 While our IR skin temperature measurements were 10km away, we believe that this is still a relatively good proxy for cloudiness, as we resample the data to cover a 30 minute window, so we feel it is more reflective of a larger area. Figure 10 shows a comparison of IR skin temperature to the MOD/MYD11 reduced data, when cloud-affected pixels are removed. There is an improvement in the RMSE of the data comparison when the cloud-affected data are removed (from 1.6°C to 1.0°C) and the mean bias is also reduced from -0.7°C to -0.4°C. In determining the strictness of the cloud mask used, there is a trade-off
335 due to the need to mask out all cloud contaminated pixels but not overflag data, which results in the generation of false positives and removes pixels that were in fact clear. In comparing the MMCR data to the MOD/MYD11, we find that of the 1059 times

that the site was within the field of view of the satellites in June and July of 2015, there were 585 instances when both MMCR and MODIS detected cloud cover, 288 instances when both MMCR and MODIS indicated clear sky. This indicates 82% agreement. There were 86 false negatives (where MMCR indicates clouds and MODIS does not) and 100 false positives (where MMCR indicates clear sky, and MODIS indicates clouds). Østby et al. (2014) also use in situ cloud data to filter out MODIS surface temperatures that are impacted by the presence of clouds in their study in Svalbard. They found an overall false negative rate of 17%, whereas our false negative rate was 8%. Their work shows that the MOD35 cloud mask performs more poorly in the winter than in the summer, so the difference in false negatives is likely due to more favourable conditions for effective cloud masking due to constant sunlight during our measurement period.

345

4. Conclusions

Data collected during a 40-day field campaign at Summit, Greenland in June and July of 2015 are used to improve understanding of near-surface temperature on an ice sheet, particularly with respect to MODIS LST retrieval products. We find that at Summit, 2 m air temperature is often significantly higher than skin temperature during the summer months, particularly at periods of low incoming solar radiation and low wind speed. This result is important because previous studies that have used 2 m air temperature to validate MODIS surface temperature products have concluded that there was a cold bias in the MODIS data, but our results indicate that the MODIS data has only a very slight cold bias (-0.7°C), and the 2 m air temperature is not necessarily representative of skin temperature. Indeed, it is because of the differences between 2 m air temperature and MODIS temperature that we began to see the pervasiveness of the inversion. We do find that there is a slight cold bias in the MOD/MYD11 surface temperature products as compared to in situ IR skin temperature, but it is not as large as previous studies have reported, and the RMSE is 1.6°C. The lower RMSE and mean bias are likely a result of measuring the skin temperature using an IR instrument directly (instead of using 2 m air temperature, which resulted in an RMSE of 3.1°C and a mean bias of 1.9°C). During our study period, we measured temperatures down to approximately -30°C. In the future, we plan to extend studies of this type to longer spans of time to determine if these results also are representative of lower temperatures. Furthermore, the validation presented in this study of the strong correlation between MODIS surface temperature and snow skin temperature in the summer lays a groundwork for inversions to be studied more extensively in locations where 2 m air temperature is currently measured. Finally, by using in situ cloud radar data, we confirm, as has been noted in previous studies, that the MODIS cloud mask did not remove all cloud-obscured data from the dataset. When we remove data that were cloud-obscured using the MMCR, the RMSE of MOD/MYD11 improves to 1.0°C. This indicates that stricter cloud-masking in the MODIS surface temperature products could improve the accuracy of the data collected.

365

Acknowledgements

We thank the National Science Foundation and Vasilii Petrenko for the opportunity to conduct this field work, and Polar Field Services and staff at Summit Station for logistic support. Data including 2 m air temperature, wind speed, and irradiance were provided by NOAA's Earth System Research Laboratory Global Monitoring Division. Cloud radar data were provided by the

370

ICECAPS program. This work was funded by NSF-GRFP 2014186404 and NSF-1506155. Dorothy Hall was funded by NASA-NNX16AP80A.

References

375

Ahlstrøm, A.P. and PROMICE project team: A new programme for monitoring the mass loss of the Greenland ice sheet, In Bennike O. and Higgins A.K. eds. Review of survey activities 2007, GEUS, Copenhagen (Geological Survey of Denmark and Greenland Bulletin 15), 2008.

380 Berkelhammer, M., Noone, D. C., Steen-Larsen, H. C., Bailey, A., Cox, C. J., O'Neill, M. S., Schneider, D., Steffen, K., and White, J. W. C.: Surface-atmosphere decoupling limits accumulation at Summit, Greenland, *Science Advances*, 2, doi:10.1126/sciadv.1501704, 2016.

Box, J. E.: Greenland ice sheet surface mass balance reconstruction. Part II: surface mass balance (1840-2010), *Journal of Climate*, 26, 6974-6989, doi: 10.1175/JCLI-D-12-00518.1, 2013.

385 Casey, K. A., Polashenski, C. M., Chen, J., & Tedesco, M.: Impact of MODIS sensor calibration updates on Greenland Ice Sheet surface reflectance and albedo trends. *The Cryosphere*, 11(4), 1781, 2017. <https://doi.org/10.5194/tc-11-1781-2017>.

Comiso, J. C.: Warming trends in the Arctic from clear sky satellite observations. *Journal of Climate*, 16(21), 3498-3510, 2003.

Fausto, R., van As, D. and Ahlstrøm, A.: Assessing the accuracy of Greenland ice sheet ice ablation measurements by pressure transducer, *Journal of Glaciology*, 58, 1144-1150, doi: 10.3189/2012JoG12J075, 2012.

390 Fréville, H., Brun, E., Picard, G., Tatarinova, N., Arnaud, L., Lanconelli, C., Reijmer, C., and van den Broeke, M.: Using MODIS land surface temperatures and the Crocus snow model to understand the warm bias of ERA-Interim reanalyses at the surface in Antarctica, *The Cryosphere*, 8, 1361–1373, doi:10.5194/tc-8-1361-2014, 2014.

Good, E. J.: An in situ-based analysis of the relationship between land surface “skin” and screen-level air temperatures, *Journal of Geophysical Research: Atmospheres*, 121, 8801–8819, doi:10.1002/2016JD025318, 2016.

395 Guillevic, P., Göttsche, F., Nickeson, J., Hulley, G., Ghent, D., Yu, Y., Trigo, I., Hook, S., Sobrino, J.A., Remedios, J., Román, M. and Camacho, F.: Land Surface Temperature Product Validation Best Practice Protocol. Version 1.0. In P. Guillevic, F. Göttsche, J. Nickeson and M. Román (Eds.), *Best Practice for Satellite-Derived Land Product Validation* (p. 60): Land Product Validation Subgroup (WGCV/CEOS), doi:10.5067/doc/ceoswgcv/lpv/lst.001. https://lpvs.gsfc.nasa.gov/PDF/CEOS_LST_PROTOCOL_Oct2017_v1.0.0.pdf

400 Hachem, S., Duguay, C. R., and Allard, M.: Comparison of MODIS-derived land surface temperatures with ground surface and air temperature measurements in continuous permafrost terrain, *The Cryosphere*, 6, 51–69, doi:10.5194/tc-6-51-2012, 2012.

Hall, D. K., Key, J. R., Casey, K. A., Riggs, G. A., and Cavalieri, D. J.: Sea ice surface temperature product from MODIS, *IEEE transactions on geoscience and remote sensing*, 42, 1076–1087, doi:10.1109/TGRS.2004.825587, 2004.

- 405 Hall, D. K., Box, J. E., Casey, K. A., Hook, S. J., Shuman, C. A., and Steffen, K.: Comparison of satellite-derived and in situ observations of ice and snow surface temperatures over Greenland, *Remote Sensing of Environment*, 112, 3739–3749, doi: 10.1016/j.rse.2008.05.007, 2008.
- Hall, D. K., Comiso, J. C., DiGirolamo, N. E., Shuman, C. A., Key, J. R., and Koenig, L. S.: A satellite-derived climate-quality data record of the clear-sky surface temperature of the Greenland ice sheet, *Journal of Climate*, 25, 4785–4798, 410 doi:10.1175/JCLI-D-11-00365.1, 2012.
- Hall, D. K., Comiso, J. C., DiGirolamo, N. E., Shuman, C. A., Box, J. E., and Koenig, L. S.: Variability in the surface temperature and melt extent of the Greenland ice sheet from MODIS, *Geophysical Research Letters*, 40, 2114–2120, doi:10.1002/grl.50240, 2013.
- Hall, D. K., Nghiem, S. V., Rigor, I. G., and Miller, J. A.: Uncertainties of temperature measurements on snow-covered land 415 and sea ice from in situ and MODIS data during BROMEX, *Journal of Applied Meteorology and Climatology*, 54, 966–978, doi:10.1175/JAMC-D-14-0175.1, 2015.
- Hook, S. J., Vaughan, R. G., Tonooka, H., and Schladow, S. G.: Absolute Radiometric In-Flight Validation of Mid Infrared and Thermal Infrared Data From ASTER and MODIS on the Terra Spacecraft Using the Lake Tahoe, CA/NV, USA, Automated Validation Site, *IEEE Transactions on Geoscience and Remote Sensing*, 45, 1798–1807, 420 doi:10.1109/TGRS.2007.894564, 2007.
- Hudson, S. R. and Brandt, R. E.: A look at the surface-based temperature inversion on the Antarctic Plateau, *Journal of Climate*, 18, 1673–1696, doi:10.1175/JCLI3360.1, 2005.
- Koenig, L. S. and Hall, D. K.: Comparison of satellite, thermochron and air temperatures at Summit, Greenland, during the winter of 2008/09, *Journal of Glaciology*, 56, 735–741, 2010.
- 425 Kuipers Munneke, P., van den Broeke, M. R., King, J. C., Gray, T., and Reijmer, C. H.: Near-surface climate and surface energy budget of Larsen C ice shelf, Antarctic Peninsula, *The Cryosphere*, 6, 353–363, doi:10.5194/tc-6-353-2012, 2012.
- Lyapustin, A., Wang, Y., Xiong, X., Meister, G., Platnick, S., Levy, R., Franz, B., Korkin, S., Hilker, T., Tucker, J., et al.: Scientific impact of MODIS C5 calibration degradation and C6+ improvements, *Atmospheric Measurement Techniques*, 7, 4353–4365, doi:10.5194/amt-7-4353-2014, 2014.
- 430 Miller, N. B., Turner, D. D., Bennartz, R., Shupe, M. D., Kulie, M. S., Cadeddu, M. P., and Walden, V. P.: Surface-based inversions above central Greenland, *Journal of Geophysical Research: Atmospheres*, 118, 495–506, doi:10.1029/2012JD018867, 2013.
- Miller, N. B., Shupe, M. D., Cox, C. J., Walden, V. P., Turner, D. D., and Steffen, K.: Cloud Radiative Forcing at Summit, Greenland, *Journal of Climate*, 28, 6267–6280, doi:10.1175/JCLI-D-15-0076.1, 2015.
- 435 Miller, N. B., Shupe, M. D., Cox, C. J., Noone, D., Persson, P. O. G., and Steffen, K.: Surface energy budget responses to radiative forcing at Summit, Greenland, *The Cryosphere*, 11, 497–516, doi:10.5194/tc-11-497-2017, 2017.

- Moran, K. P., Martner, B. E., Post, M., Kropfli, R. A., Welsh, D. C., and Widener, K. B.: An unattended cloud-profiling radar for use in climate research, *Bulletin of the American Meteorological Society*, 79, 443–455, doi:0.1175/1520-0477(1998)079<0443:AUCPRF>2.0.CO;2, 1998.
- 440 Navari, M., Margulis, S., Bateni, S., Tedesco, M., Alexander, P., and Fettweis, X.: Feasibility of improving a priori regional climate model estimates of Greenland ice sheet surface mass loss through assimilation of measured ice surface temperatures, *The Cryosphere*, 10, 103–120, doi:10.5194/tc-10-103-2016, 2016.
- NOAA ESRL Global Monitoring Division. Meteorology measurements at Summit Station, Greenland. Compiled by the Observatory Operations Group. National Oceanic and Atmospheric Administration (NOAA), Earth System Research
- 445 Laboratory (ESRL), Global Monitoring Division (GMD): Boulder, Colorado, USA, 2017.
- Østby, T. I., Schuler, T. V., and Westermann, S.: Severe cloud contamination of MODIS Land Surface Temperatures over an Arctic ice cap, Svalbard, *Remote Sensing of Environment*, 142, 95–102, doi:10.1016/j.rse.2013.11.005, 2014.
- Overland, J. E., and Guest, P. S.: The Arctic snow and air temperature budget over sea ice during winter. *Journal of Geophysical Research: Oceans* (1978–2012), 96(C3), 4651–4662. 1991.
- 450 Polashenski, C. M., Dibb, J. E., Flanner, M. G., Chen, J. Y., Courville, Z. R., Lai, A. M., Schauer, J. J., Shafer, M. M., and Bergin, M.: Neither dust nor black carbon causing apparent albedo decline in Greenland’s dry snow zone: Implications for MODIS C5 surface reflectance, *Geophysical Research Letters*, 42, 9319–9327, doi:10.1002/2015GL065912, 2015.
- Rae, J., Aðalgeirsdóttir, G., Edwards, T., Fettweis, X., Gregory, J., Hewitt, H., Lowe, J., Lucas-Picher, P., Mottram, R., Payne, A., et al.: Greenland ice sheet surface mass balance: evaluating simulations and making projections with regional climate
- 455 models, *The Cryosphere*, 6, 1275–1294, doi:10.5194/tc-6-1275-2012, 2012.
- Reeh, N.: Parameterization of melt rate and surface temperature on the Greenland ice sheet, *Polarforschung*, 59, 113–128, 1989.
- Reeves Eyre, J. E. J. and Zeng, X.: Evaluation of Greenland near surface air temperature datasets, *The Cryosphere*, 11, 1591–1605, doi:10.5194/tc-11-1591-2017, 2017.
- 460 Riggs, G. A., Hall, D. K., & Román, M. O. (2017). Overview of NASA's MODIS and Visible Infrared Imaging Radiometer Suite (VIIRS) snow-cover Earth System Data Records. *Earth System Science Data*, 9(2), 765.
- Rignot, E., Velicogna, I., van den Broeke, M. R., Monaghan, A., and Lenaerts, J. T.: Acceleration of the contribution of the Greenland and Antarctic ice sheets to sea level rise, *Geophysical Research Letters*, 38, doi:10.1029/2011GL046583, 2011.
- Shamir, E. and Georgakakos, K. P.: MODIS Land Surface Temperature as an index of surface air temperature for operational
- 465 snowpack estimation, *Remote Sensing of Environment*, 152, 83–98, doi:10.1016/j.rse.2014.06.001, 2014.
- Shuman, C. A., Steffen, K., Box, J. E., and Stearns, C. R.: A dozen years of temperature observations at the Summit: Central Greenland automatic weather stations 1987–99, *Journal of Applied Meteorology*, 40, 741–752, doi:10.1175/1520-0450(2001)040<0741:ADYOTO>2.0.CO;2, 2001.

- Shuman, C. A., Hall, D. K., DiGirolamo, N. E., Mefford, T. K., and Schnaubelt, M. J.: Comparison of near-surface air
470 temperatures and MODIS ice-surface temperatures at Summit, Greenland (2008–13), *Journal of Applied Meteorology and
Climatology*, 53, 2171–2180, doi:10.1175/JAMC-D-14-0023.1, 2014.
- Steffen, K. and Box, J.: Surface climatology of the Greenland ice sheet: Greenland Climate Network 1995–1999, *Journal of
Geophysical Research: Atmospheres*, 106, 33 951–964, doi:10.1029/2001JD900161, 2001.
- Steffen, K., Box, J., and Abdalati, W.: Greenland climate network: GC-Net, US Army Cold Regions Research and Engineering
475 (CRREL), CRREL Special Report, pp. 98–103, 1996.
- Stocker, T. F.: *Climate change 2013: the physical science basis: Working Group I contribution to the Fifth assessment report
of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 2014.
- Sverdrup, H. U.: Scientific work of the Maud Expedition, 1922–1925, *The Scientific Monthly*, 22, 400–410, 1926.
- Van As, D., Van Den Broeke, M., and Van De Wal, R., Daily cycle of the surface layer and energy balance on the high
480 Antarctic Plateau, *Antarctic Science*, 17(1), 121–133, doi: 10.1017/S095410200500252X, 2005
- Van As D., Fausto, R.S. and PROMICE Project Team: Programme for monitoring of the Greenland Ice Sheet (PROMICE):
first temperature and ablation records, In Bennike O., Garde A.A. and Watt W.S. eds., *Review of survey activities 2010.*
GEUS, Copenhagen, 73–76 (*Geological Survey of Denmark and Greenland Bulletin 23*), 2011.
- Van Den Broeke, M., Reijmer, C., Van As, D., and Boot, W., Daily cycle of the surface energy balance in Antarctica and the
485 influence of clouds, *International Journal of Climatology*, 26(12), 1587–1605, doi: 10.1002/joc.1323, 2006.
- van den Broeke, M. R., Enderlin, E. M., Howat, I. M., Kuipers Munneke, P., Noël, B. P. Y., van de Berg, W. J., van Meijgaard,
E., and Wouters, B.: On the recent contribution of the Greenland ice sheet to sea level change, *The Cryosphere*, 10, 1933–
1946, doi:10.5194/tc-10-1933-2016, 2016.
- Van Tricht, K., Lhermitte, S., Lenaerts, J. T., Gorodetskaya, I. V., L’Ecuyer, T. S., Noël, B., van den Broeke, M. R., Turner,
490 D. D., and Van Lipzig, N. P.: Clouds enhance Greenland ice sheet meltwater runoff, *Nature communications*, 7,
doi:10.1038/ncomms10266, 2016.
- Vernon, C. L., Bamber, J., Box, J., Van den Broeke, M., Fettweis, X., Hanna, E., and Huybrechts, P.: Surface mass balance
model intercomparison for the Greenland ice sheet, *Cryosphere (The)*, 7, 599–614, doi:10.5194/tc-7-599-2013, 2013.
- Wan, Z.: MODIS Land-Surface Temperature Algorithm Theoretical Basis Document (LST ATBD) Version 3.3,
495 https://modis.gsfc.nasa.gov/data/atbd/atbd_mod11.pdf, 1999.
- Wan, Z.: New refinements and validation of the MODIS land-surface temperature/emissivity products, *Remote sensing of
Environment*, 112, 59–74, doi:10.1016/j.rse.2006.06.026, 2008.
- Wan, Z.: New refinements and validation of the collection-6 MODIS land-surface temperature/emissivity product, *Remote
sensing of Environment*, 140, 36–45, doi:10.1016/j.rse.2013.08.027, 2014.
- 500 Wan, Z. and Dozier, J.: A generalized split-window algorithm for retrieving land-surface temperature from space, *IEEE
Transactions on geoscience and remote sensing*, 34, 892–905, doi:10.1109/36.508406, 1996.

- Warren, S. G. and Brandt, R. E.: Optical constants of ice from the ultraviolet to the microwave: A revised compilation, *Journal of Geophysical Research*, 113, doi:10.1029/2007jd009744, 2008.
- Wenny, B., Xiong, X., and Madhavan, S.: Evaluation of Terra and Aqua MODIS thermal emissive band calibration consistency, in: *SPIE Remote Sensing*, pp. 853 317–853 317, International Society for Optics and Photonics, doi:10.1117/12.974230, 2012.
- Westermann, S., Langer, M., and Boike, J.: Systematic bias of average winter-time land surface temperatures inferred from MODIS at a site on Svalbard, Norway, *Remote Sensing of Environment*, 118, 162 – 167, doi:http://dx.doi.org/10.1016/j.rse.2011.10.025, 2012.
- 510 Williamson, S. N., Hik, D. S., Gamon, J. A., Jarosch, A. H., Anslow, F. S., Clarke, G. K., and Rupp, T. S.: Spring and summer monthly MODIS LST is inherently biased compared to air temperature in snow covered sub-Arctic mountains, *Remote Sensing of Environment*, 189, 14–24, doi:10.1016/j.rse.2016.11.009, 2017.
- Yu, Y., Rothrock, D. A., and Lindsay, R. W.: Accuracy of sea ice temperature derived from the advanced very high resolution radiometer, *Journal of Geophysical Research: Oceans (1978–2012)*, 100(C3), 4525-4532, 1995.
- 515 Xiong, X., Wu, A., Wenny, B. N., Madhavan, S., Wang, Z., Li, Y., Chen, N., Barnes, W. L., and Salomonson, V. V.: Terra and Aqua MODIS Thermal Emissive Bands On-Orbit Calibration and Performance, *IEEE Transactions on Geoscience and Remote Sensing*, 53, 5709–5721, doi:10.1109/TGRS.2015.2428198, 2015.




Figure 1: Map indicating the location of Summit, Greenland, the study site for remote sensing and in situ temperature comparisons. Contour lines represent elevation change of 500 m. Latitude and longitude coordinates for the measurement site are 72.65923°N, 38.57067°W.



530 Figure 2: Image of the IR skin temperature sensor and tripod set up.




Figure 3: Time series of skin temperature at Summit, Greenland measured with SI-111 IR thermometer (blue). Grey bars indicate presence of clouds as detected by a millimeter cloud radar at Summit Station.






Figure 4: Time series of IR skin temperature, 2 m air temperature, during a clear sky period near Summit, Greenland.


540



545 Figure 5: a) Comparison of 2 m air temperature to IR skin temperature at Summit, Greenland during June and July 2015. The difference between air and skin temperature is largest at lower temperatures. b) Histogram of the difference between 2 m air temperature and IR skin temperature during the study period in June and July of 2015 at Summit, Greenland during all sky conditions and c) clear sky and cloudy sky conditions (as detected by MMCR data) separated. The difference is skewed to positive temperature differences indicating higher air temperatures than skin temperatures.




550 Figure 6: Difference between 2m air temperature and IR skin temperature showing the presence of strong surface-based inversions at low wind speeds and low values of incoming solar radiation (indicated by the marker colour).



560 Figure 7: Time series as shown in Figure 3 with only a temporal subset of data presented to clearly show the diurnal cycle of temperature during fairly clear conditions. Note that the MOD/MYD11 product shows good agreement with IR skin temperature throughout the diurnal cycle.

565



570






Figure 8: a) Direct comparison of in situ IR skin temperature data with MOD/MYD11 C6 surface temperatures. Agreement between satellite and ground-based measurements is quite good (RMSE = 1.6°C, n=374), and there is not a noticeable difference between the performance of the MOD11 and MYD11 temperature products, on the Terra and Aqua satellites, respectively. b) Direct comparison of 2 m air temperature with MOD/MYD C6 surface temperatures. This is illustrative of bias that may be inferred if 2 m air temperature is used in validation studies when inversions are present. RMSE = 3.1°C.

575



580 Figure 9: Difference in temperature measured from MOD/MYD11 and in situ IR skin temperature measurements as a function of a) IR skin temperature, b) solar zenith angle, and c) MODIS viewing angle. The only significant relationship is that the temperature difference is sensitive to the MODIS viewing angle. While the relationship is statistically significant, it is not a strong control on the temperature difference.



585 Figure 10: Comparison of MOD/MYD11 to in situ IR skin temperature after cloud-affected data are removed. The RMSE is 1.0°C and the mean bias is -0.4°C.

590 Table 1: Summary statistics from recent literature comparing MODIS surface temperature products to in situ surface temperature measurements in snow-covered regions.

Study	Location	Temperature Range	Temperature Measurement	MODIS Product	MODIS Collection	Sample Number	RMSE	Bias
Hall et al. 2004	South Pole	-70°C to -20°C	2m Air	29	4	255	1.7°C	-1.2°C
Hall et al. 2008	Greenland AWS	-40°C to 0°C	2m Air	11	4	48	2.1°C	-0.3°C
Koenig and Hall 2008	Summit, Greenland	-41°C to -20°C	Thermochron Skin	11	5	62	3.1°C	-3.4°C
		-60°C to -20°C	2m Air	11	5	259	4.1°C	-5.5°C
Westermann et al. 2012	Svalbard	-40°C to 0°C	IR Skin	11	5			~-3°C
Shuman et al. 2014	Summit, Greenland	-60°C to 0°C	2m Air	29	5	2536 2270	All: 5.3°C Filtered: 3.5°C	~-3°C
Østby et al. 2014	Svalbard	-45°C to 0°C	IR Skin	11	5	3941 1065	All: 5.3°C Filtered: 3.0°C	All: -3.3°C Filtered: -0.3°C
Hall et al. 2014	Barrow, Alaska	-42°C to -20°C	Thermochron Skin	11	5	69		-2.3±3.9°C
				11	5	84		0.6±2.0°C
This Study	Summit, Greenland	-30°C to 0°C	IR Skin	11	6	374	All: 1.6°C	All: -0.7±1.4°C
						288	Cloud Filter: 1.0°C	Cloud Filter: -0.4±0.9°C
					MOD11	6	207	1.8°C
				MYD11	6	167	1.4°C	-0.6±1.3°C