

Response to reviewers:

The role of subtemperate slip in thermally-driven ice stream margin migration

M. Haseloff, C. Schoof, O. Gagliardini

We thank both reviewers for their time and effort to review our manuscript. We have responded to the two reviews point-by-point below. We are listing reviewer comments in black, and respond to these comments in blue. Page, line, and equation numbers refer to the original manuscript unless indicated otherwise.

A marked-up version of our manuscript with tracked changes is attached to the end of this document.

Anonymous Referee #1

General Comments:

The manuscript by Haseloff and coworkers addresses the enlargement of ice streams by the migration of margins that are controlled by a transition between temperate and subtemperate basal temperatures. Using a boundary layer approach, the analysis leads to predictions of outward migration rate that are formulated in terms of parameters that can be obtained from the behavior of the adjacent ice stream and ice ridge, with the ultimate goal of enabling the primary results of this study to be incorporated within large-scale ice sheet models. Several special cases are shown to admit simplified expressions that agree with detailed numerical results in particular limits that may be of relevance to different portions of the Siple Coast ice streams. The writing is clear and succinct, and the treatment and results are substantial and worthy of publication without major revisions.

Specific Comments:

The analysis is very involved and appears to be technically sound. To further improve the utility of this work and its reception by the broader community, I strongly encourage minor revisions that are aimed at providing further context and physical insight. A dense mathematical treatment is no doubt warranted, but some brevity could be sacrificed to improve the communication of this substantial effort. For example, consider enlightening the audience as to why the focus is on ice stream enlargement rather than the more general case – (e.g., is it considered that a narrowing of ice stream width is less relevant or is it a harder problem for some reason?) the title does not reflect this one-sided focus. How were the particular parameter values listed at the bottom of table 1 chosen and where might they be considered relevant (or what ranges of values might be considered typical)? Figure 10 illustrates three different regimes of behavior, which is helpful, but some brief discussion of the likely importance of these different cases could be beneficial. The mathematics is interesting, but the physical insight it provides is what makes the effort worthwhile and this aspect of the paper could be improved.

We thank the reviewer for these suggestions. To address the question of narrowing versus widening, note that the physics is different. For widening, we have changed the following passage immediately before equation 8:

“On the temperate (stream-ward) side of the ice–bed interface, we assume that the basal shear stress is negligible compared with the shear stresses in the ice, leading to a free slip boundary condition: “

and to elaborate on this, we have introduced the following additional paragraphs immediately after equation 8:

“In posing this boundary condition for an ice stream that is actively widening, we are assuming that an infinitesimal amount of melting of the bed suffices to allow for slip: once the thermal barrier at the bed is breached, we only need a very thin ice-free layer in order for slip to occur. This is consistent at least with the idea of a plastic bed, where slip can happen on a plane, or with a hard bed.

To the extent that additional degrees of freedom (other than temperature) are involved in sliding, the main concern would be presumably water pressure at the bed or within the till, rather than the thickness of the unfrozen till layer. Our assumption of a free slip once the melting point is reached is best justified (see Haseloff et al, 2015) if we suppose that the unfrozen bed is hydraulically well-connected, so that the water pressure in the parts of the bed that have just become unfrozen quickly equilibrates with water pressure elsewhere under the ice stream (and hence basal friction is comparable to the rest of the active ice stream). Shear stresses experienced by the margins of the ice stream are large compared with basal drag throughout the ice stream (Haseloff et al, 2015), and this implies that basal friction is small at leading order everywhere where the melting point is reached. There are undoubtedly other, more elaborate models for basal shear stress at the unfrozen bed; ours is the simplest possible case to analyse.”

The case of freezing is quite different, because here, the depth of the unfrozen till layer (if present) does matter: while sliding can potentially be activated by only melting out a small portion of the bed, shutting sliding down requires freezing all the way through any deformable till layer. We acknowledge this now in the following updated passage in the introduction

“Under these conditions, the inwards migration (or narrowing) of an ice stream requires freezing of the entire sediment column (Schoof, 2012, appendix B). As melt water can be supplied to sections of the bed with active freezing from other regions of the ice stream via subglacial drainage, this necessitates taking into account the ice-stream wide energy balance and the thermal response of bed. Consequently, the inwards migration of ice streams is the result of insufficient heat dissipation over the width of the entire ice stream (Haseloff, 2015). However, as shown in Haseloff (2015) this process can at least in principle be modelled with large-scale ice sheet models without recourse to a boundary layer.”

In the paragraph that follows, we also expand on the difference between widening and narrowing:

“In this scenario the outwards migration of ice stream margins requires melting of the frozen sediment under the ice ridge. By contrast with a narrowing ice stream, it is however not

necessary for the entire thickness of the sediment column to melt out: only part of it needs to be unfrozen to permit sliding, and we will later idealize this by assuming that sliding is possible as soon as the melting point is reached at the bed. This however also underlines the asymmetry between widening and narrowing of an ice stream, which motivates us to focus on the harder problem of widening, which requires heat to be transferred into the bed. Several studies show that a strong gradient in basal resistance created by a thermal transition at the bed leads to significant englacial heat production in the ice stream margins. “

The values of the ice stream thickness h_s , marginal inflow q_r and lateral shear stress τ_s are chosen to be representative of the upper margin of Whillans ice stream studied in Harrison et al (1998) and Echelmeyer & Harrison (1999) which the authors observed to migrate at a rate between 7 – 30 m/year. We have amended the caption of table 1 to clarify this.

To clarify which of the different parameterizations might be relevant where, we have added the following paragraph to the discussion section:

“Since the parameter ranges where equations (54)-(56) are applicable for a given ice stream depends on the yield strength of the subtemperate region, knowledge of the basal properties is necessary. There are only very few observations which would allow an estimate of the subtemperate basal yield strength (Holdsworth, 1974; Echelmeyer and Zhongxiang (1987); Cuffey et al, 1999). Using the values reported in Cuffey et al. (1999) for Meserve Glacier, Antarctica, gives an approximate basal yield stress of 380 kPa (neglecting the dependence on the thickness of the interfacial water layers). As the typical lateral shear stress of an ice stream margin can be estimated from $\tau_s = \rho g \sin \alpha W_s/2$ to be in the range of 180 kPa to 310 kPa (Joughin et al, 2002, table 3), this suggests that equation (55) will be appropriate in most circumstances.”

In the heat balance treatment, it appears that the bed is ice free and so can warm from subtemperate to temperate conditions without latent heat effects. If the physical situation were different, I would have anticipated that this would strongly retard the rate of margin migration. Some further explanation should be provided for why only sensible heat need be considered beneath the ridge.

This is hopefully largely answered above.

To reiterate, we imagine the frozen part of bed to be either bed rock or an ice-sediment matrix, and therefore not necessarily ice free. The point is however that only a very small amount of the ice in the ice-sediment matrix (if present) needs to melt to create a layer that can support motion. We idealize this by saying that sliding becomes possible as soon as the melting point is reached, without requiring the full thickness of sediment to thaw out (similar physics is effectively assumed in e.g. Robel et al. (2014). Of course, latent heat effects will continue to enter into the energy balance of the bed: for our travelling wave formulation, this happens on the *warm* side, where the bed generally has a positive energy balance. That positive energy balance naturally goes into melting – and will melt out any remaining ice in the pore space of till (or equally, “dirty basal ice”, which may be the same thing) before melting clean ice. This is an effect we could diagnose after the fact, but it does not enter into the dynamical part of our model.

Similarly, in the description surrounding Figure 3, I found the discussion surrounding the enthalpy gradient model unsatisfying. To be clear, are the contours in panels c, g, and h

showing ice at +3C? Perhaps a brief discussion in the supplementary information might be used to outline why it's okay once again to neglect latent heat.

This paper is probably not the place to explain the basics of enthalpy formulations, but we have expanded the text in the description of figure 3 (see below). The basic point is that latent heat is **not** neglected. Rather than interpreting T as temperature, it becomes necessary in regions where $T > 0$ to interpret ρ^*c^*T as the enthalpy content of the ice (i.e., the latent heat content), rather than as something proportional to actual temperature. This is a common approach in temperate ice modelling (see Aschwanden et al, 2012; Schoof and Hewitt, 2016; Hewitt and Schoof, 2017). The reason why this works is that ρ^*c^*T in the derivation of the heat equation starts out as internal energy (or enthalpy) content per unit volume, and the conservation law can be extended to cover latent heat as well.

By solving the heat equation in regions where the internal energy content comes in the form of latent rather than specific heat (the temperate regions, where our model computes $T > 0$), we make a very specific assumption about the flux of that energy: by retaining the heat equation, we are assuming that internal energy flows down its own spatial gradient, with a diffusivity that remains the same as when internal energy takes the form of specific heat (the “cold” part of the domain). The assumption that internal energy flows down its own gradient is the cornerstone of the widely used enthalpy gradient model as described at length in Aschwanden et al; our model goes a step further and assumes equal diffusivities in the cold and temperate regions. Neither assumption is really likely to be a good one (Schoof and Hewitt, 2016, Hewitt and Schoof, 2017).

In practice, because we mostly focus on high-Peclet-number situations with advection from the cold side, the effect of incorrectly modelling temperate ice is likely to be confined near the regions where our model predicts $T > 0$, and those regions are pushed to the right of the origin when we have lateral advection and subtemperate sliding. V_m is ultimately controlled by the behaviour of the temperature field around the origin (where the two inequalities in (17) must both be satisfied within the a given neighbourhood of $Y = 0$). It is therefore likely that our treatment of temperate ice has little effect on the predicted V_m in that situation (large Pe and moderate or small τ); for little or no subtemperate slip, a more careful look at temperate ice modelling is warranted, as we point out in section 5.

The text alteration we have made is the following, in section 3.2:

*“Note that all the temperature fields shown in figure 3 have $T > 0$ in some parts of the ice. The boundaries of these regions are marked by a bold red line. In these regions of the ice stream, we solve the same heat equation as in the remainder of the domain (see also Schoof, 2012, Haseloff et al, 2015). Obviously, ice cannot have a temperature in excess of its melting point, and T cannot be interpreted as temperature where $T > 0$. Effectively, we assume a very special case of an enthalpy gradient model (Aschwanden et al, 2012, see also Schoof and Hewitt, 2016, Hewitt and Schoof, 2017): where $T > 0$, the product $\rho^*c_p^*T$ (which is generally the specific heat content per unit volume of ice) must instead be interpreted as the latent heat content per unit volume of the ice. That is, ρ^*c^*T should be interpreted as $\rho_w^*L^*\phi$, where ρ_w is the density of water, L is latent heat per unit mass, and ϕ is the volumetric moisture content of the ice. This allows us to identify $\phi = \rho^*c_p^*T/(\rho_w^*L)$, so T is nothing more than a proxy for moisture content when $T > 0$.*”

By solving the heat equation where $T > 0$, we make two main assumptions. First, qualitatively, we assume that moisture flows down gradients of moisture, which is the assumption common to enthalpy gradient models and permits the same diffusive model to be applied regardless of whether the melting point has been reached or not. The second, quantitative assumption we make is that the corresponding diffusivity remains the same for cold and temperate regions. This is consistent with prior work, but also an obvious area for future model improvement. We will return to a discussion of the limitations imposed by this assumption in Section 5.”

At the end of section 5, we have added:

“Finally, we have assumed that the dynamics of temperate ice can be represented by a particular version of an enthalpy gradient model (Aschwanden et al., 2012). We expect future iterations of our model to incorporate ice dynamics which can account for gravity-driven moisture transport in temperate ice (e.g., Schoof and Hewitt, 2016). A more sophisticated treatment of temperate ice is likely to be particularly relevant when temperate ice forms near the transition from a cold to a temperate bed: as we have seen, the Peclet number in a shear margin is large, and temperate ice formation down-flow from the cold-temperate transition is consequently unlikely to affect the temperature field close to the transition, as T is dominated by advection. This makes our results with moderate to small τ_c (or, more accurately, with moderate to small τ) likely to be the most robust to changes in the temperate ice model, since temperate ice forms some distance inside the ice stream in that case (see for instance Fig. 3). We leave a deeper investigation to future work.”

What is it about $V_m < 0$ that necessitates a different treatment strategy? I understand that there has been a deliberate decision to focus on cases with $V_m > 0$, but it would be helpful to give the reader some insight into why.

This is hopefully covered by the above; to reiterate, we have included the following passages in the introduction

“Under these conditions, the inwards migration (or narrowing) of an ice stream requires freezing of the entire sediment column (Schoof, 2012, appendix B). As melt water can be supplied to sections of the bed with active freezing from other regions of the ice stream via subglacial drainage, this necessitates taking into account the ice-stream-wide energy balance and the thermal response of bed. Consequently, the inwards migration of ice streams is the result of insufficient heat dissipation over the width of the entire ice stream (Haseloff, 2015). However, as shown in Haseloff (2015) this process can at least in principle be modelled with large-scale ice sheet models.

In this scenario the outwards migration of ice stream margins requires melting of the frozen sediment under the ice ridge. By contrast with a narrowing ice stream, it is however not necessary for the entire thickness of the sediment column to melt out: only part of it needs to be unfrozen to permit sliding, and we will later idealize this by assuming that sliding is possible as soon as the melting point is reached at the bed. This however also underlines the asymmetry between widening and narrowing of an ice stream, which motivates us to focus on the harder problem of widening, which requires heat to be transferred into the bed. Several studies show that a strong gradient in basal resistance created by a thermal transition at the bed leads to significant englacial heat production in the ice stream margins (...).”

Technical Corrections:

Equation 10 doesn't need the :
Removed.

The density of the bed in Table 1 must be wrong.
Density, heat capacity, and thermal conductivity of the bed are set to the same values as in the ice, as we envision the bed as an ice/sediment mixture.

Line 22 on page 13 should read “constrained” not constraint
Corrected.

In many of the figures the symbols are so large as to obstruct the underlying lines (e.g. fig. 4). These could be shrunk or even eliminated in favor of combinations of dashes and dots.
The symbols are shrunk in figures 4, 6, 7, and 9 of the revised version of the manuscript.

The caption for figure 6c is confusing and the symbols for Θ_1 - Θ_3 are difficult to discern. This could be improved. As it stands, the description at the end of 4.3 describes results that are very difficult to see.
The figure and caption have been updated.

On line 32 of page 21 we are told that the stress scale can be found from (37), but further steps are required - maybe show this in the supplement.
We have added extra information to the text to that should clarify how the stress scale can be found.

On line 16 of page 28, give the physical explanation for the upper limit on χ - the number by itself is not particularly helpful.
We have added the following statement to the text:

“The upper limit on χ corresponds to the lower limit on heat dissipation which is required for outwards migration.”

Anonymous Referee #2

This paper derives a parameterization of the rate at which thermally-controlled shear margins migrate outwards based on several physical parameters that can be constrained observationally or theoretically. The model is based on a boundary-layer approximation published previously by the authors and the resulting governing equations are solved through the standard software package Elmer/Ice. The paper is well structured and clearly written.

To the best of my knowledge and understanding, the models and results presented here are novel, relevant, and mathematically and physically sound. The conclusions are thoroughly backed up by the results presented in the manuscript. There are no results that appear to be unsupported. I recommend this manuscript for publication after minor modifications have been made, as it would constitute a valuable contribution to the community and readers of The Cryosphere.

That being said, I think there are a couple of opportunities for improving this manuscript that the authors should consider.

(1) A key goal of this paper is to propose a parameterization of the rate of margin migration that can be incorporated into large-scale ice-sheet models to capture this process. I like the ambition, but I think it would be valuable for the authors to state more explicitly that this parametrization is probably not generally applicable. There is the obvious issue of topographic controls, which are certainly relevant in some cases. Then, there is the assumption that the shear margin is thermally controlled and I think it would be useful for readers to learn whether the authors argue that they believe all margins are thermally controlled or whether they simply focus on this subset for the purposes of this study. Finally, the boundary-layer approach might not be equally applicable everywhere? For example, it is not clear to me that a margin that is wide as compared to the ice thickness scale (e.g., the Whillans ice plain with margins of 30km+) would be well described by this particular approximation.

We thank the reviewer for these suggestions, which we have incorporated by making the following changes:

- To emphasize that we are only interested in ice streams that are not topographically controlled, we have changed the first sentence in the discussion to

“In this study, we have investigated how different physical processes determine the widening of ice streams that are not topographically confined.”

- We have also added the following clarifying sentence to the model description after we describe the model geometry, which assumes a flat bed:

“Note however that this assumption does not exclude the application of our results to ice streams with a weak topographic control, as found in many regions of the Siple Coast: this assumption merely requires the elevation differences of the bed to be sufficiently small that it does not vary significantly over the lateral distance of a few ice thicknesses.”

- A discussion of thermal vs hydrological controls is given in the second paragraph of the discussion/conclusion section, where we underline that we are interested in one

particular version of shear margins (and by implication admit that, for a particular margin, different physics might apply).

We do not think that most readers would identify an ice plain as an ice stream margin in the classical sense, especially if they look at the geometry in Figure 1, or if they see the paper in the context of prior literature (Raymond, 1996; Jacobson and Raymond, 1998; Schoof, 2006; Suckale et al., 2014; Perol and Rice, 2015). While the question of how such regions work mechanically is perfectly valid, we felt that it would actually be confusing to discuss them in the paper.

(2) The boundary-layer model is very similar to the approach previously published by the same authors in JFM. One difference that stands out is the inclusion of subtemperate sliding leading to a different stress boundary condition. Are there other differences that I've missed? Given that the model approach as such has been published already, I'm not sure that it is necessary to introduce the model quite in this level of detail. Instead, I would suggest emphasizing the differences between the models more clearly. Given that the audience of TC is less theoretically inclined than the readers of JFM, I suggest a concise summary of the governing equations and boundary conditions together with a summary of the key assumptions behind the boundary-layer approach and the limitations entailed for applying this idea to field data. For example, the surface correction s' does not actually couple back to the mechanical model, so would this approach be suited to think about the Thwaites shear margin given the rapid thinning rates and differences in ice thickness in ridge/stream? In the discussion section, the authors argue that they make two simplifying assumption, (1) assuming subtemperate slip and (2) ignoring temperature-dependence of stress and viscosity. I would argue that there are a lot more assumptions entailed in the model setup. I think it would be helpful to spell these out explicitly in the discussion.

The reviewer raises two main points, to which we respond separately:

1. The model should be written more concisely
2. The differences to the published paper in JFM should be made clearer, and the limitations of the model should be made clearer

Point 1:

Since reviewer #1 states above: "A dense mathematical treatment is no doubt warranted, but some brevity could be sacrificed to improve the communication of this substantial effort.", we have opted to not further shorten the model description, as this could make the paper even harder to read. In our collective experience publishing other modelling papers, truncated model descriptions usually lead to requests for a complete model (even if this has been stated in large part elsewhere).

Point 2:

We have added the following to the introduction:

"The purpose of this paper is therefore twofold: (i) to use the margin boundary layer model of Haseloff et al. (2015) to investigate how subtemperate slip changes the heat production and temperature field in the ice stream margin, and thereby the rate at which ice streams can migrate outwards and (ii) to derive parameterizations of margin migration rate which can be used in large scale ice sheet models. Both of these points go beyond the work in Haseloff et al (2015); the parameterizations we derive in particular show how the limit of rapid advection of heat across the shear margin can be used to simplify the boundary layer

model and arrive at tractable forms of the migration rate that could be implemented in computational models either in the form of semi-analytical formulae or lookup tables.”

We also have added further information about the derivation in section 2, to explicitly highlight that most simplifications of the model arise from the derivation of the boundary layer model given in Haseloff et al (2015), rather than from ad-hoc simplifications. For instance, we have added on page 3, line 23:

“The asymptotic analysis in Haseloff et al (2015) shows that the boundary layer evolves rapidly in comparison to the ice stream and ice ridge, and is consequently quasi-static with the only time-dependence arising from the moving transition between a frozen and a temperate bed at $\pm y_m(x,t)$. Moreover, Haseloff et al (2015) show that the surface of the ice stream margin is flat at leading order and located at $z=h_s$, where h_s is the ice thickness of the ice stream.”

To answer the question about the surface correction s' , the fact that this does not couple back into the model but is simply computed diagnostically comes directly out of the asymptotic expansions in Haseloff et al (2015); this is once more not an ad hoc approximation but the natural leading order form of the problem. In order for this approximation (which appears to be common to previous work as well, such as Raymond 1996, Jacobson and Raymond 1998, Schoof 2004, Suckale et al 2014, all of whom use a flat upper surface for their computational domains) not to hold, an $O(1)$ surface slope would have to exist in the shear margin – meaning, a surface angle comparable with 45 degrees. To our knowledge, such steep slopes do not usually occur at the edges of ice streams. (As with all asymptotics, our theory is never exact except in the limit, so we would expect to see deviations for moderate surface slopes, but the point of our work is clearly to get a tractable leading order result.)

With regard to the last point raised (“In the discussion section, the authors argue that they make two simplifying assumption, (1) assuming subtemperate slip and (2) ignoring temperature-dependence of stress and viscosity. I would argue that there are a lot more assumptions entailed in the model setup. I think it would be helpful to spell these out explicitly in the discussion.”), it would be helpful to know exactly which other simplifying assumption the referee has in mind. We chose to focus in the discussion on the physics that is not prescribed by the geometrical assumptions made in Haseloff et al (2015; long, narrow ice stream, ice stream width much greater than ice thickness) which gives most of the boundary conditions and the parallel-sided strip geometry of our boundary layer model. Two additional caveats we discuss are (3) the treatment of temperate ice (which we now discuss in greater detail at the very end of the discussion/conclusions section), and (4) the free slip boundary condition for $Y > 0$. For (3), we have added the following at the end of the paper:

“Finally, we have assumed that the dynamics of temperate ice can be represented by a particular version of an enthalpy gradient model (Aschwanden et al., 2012). We expect future iterations of our model to incorporate ice dynamics which can account for gravity-driven moisture transport in temperate ice (e.g., Schoof and Hewitt, 2016). A more sophisticated treatment of temperate ice is likely to be particularly relevant when temperate ice forms near the transition from a cold to a temperate bed: as we have seen, the Peclet number in a shear margin is large, and temperate ice formation down-flow from the cold-temperate transition is consequently unlikely to affect the temperature field close to the transition, as T is dominated by advection. This makes our results with moderate to small τ_c (or, more accurately, with moderate to small τ) likely to be the most robust to changes in the temperate ice model, since

temperate ice forms some distance inside the ice stream in that case (see for instance Fig. 3). We leave a deeper investigation to future work.”

In the modelling section, we have also expanded on our current formulation, adding

*“Note that all the temperature fields shown in figure 3 have $T > 0$ in some parts of the ice. The boundaries of these regions are marked by a bold red line. In these regions of the ice stream, we solve the same heat equation as in the remainder of the domain (see also Schoof, 2012, Haseloff et al, 2015). Obviously, ice cannot have a temperature in excess of its melting point, and T cannot be interpreted as temperature where $T > 0$. Effectively, we assume a very special case of an enthalpy gradient model (Aschwanden et al, 2012, see also Schoof and Hewitt, 2016, Hewitt and Schoof, 2017): where $T > 0$, the product $\rho * c_p * T$ (which is generally the specific heat content per unit volume of ice) must instead be interpreted as the latent heat content per unit volume of the ice,. That is, $\rho * c * T$ should be interpreted as $\rho_w * L * \phi$, where ρ_w is the density of water, L is latent heat per unit mass, and ϕ is the volumetric moisture content of the ice. This allows us to identify $\phi = \rho * c_p * T / (\rho_w * L)$, so T is nothing more than a proxy for moisture content when $T > 0$.*

By solving the heat equation where $T > 0$, we make two main assumptions. First, qualitatively, we assume that moisture flows down gradients of moisture, which is the assumption common to enthalpy gradient models and permits the same diffusive model to be applied regardless of whether the melting point has been reached or not. The second, quantitative assumption we make is that the corresponding diffusivity remains the same for cold and temperate regions. This is consistent with prior work, but also an obvious area for future model improvement. We will return to a discussion of the limitations imposed by this assumption in Section 5.”

With regard to point (4) (friction on the warm side $Y > 0$), we have added the following text in section 2:

“In posing this boundary condition for an ice stream that is actively widening, we are assuming that an infinitesimal amount of melting of the bed suffices to allow for slip: once the thermal barrier at the bed is breached, we only need a very thin ice-free layer in order for slip to occur. This is consistent at least with the idea of a plastic bed, where slip can happen on a plane, or with a hard bed.

To the extent that additional degrees of freedom (other than temperature) are involved in sliding, the main concern would be presumably water pressure at the bed or within the till, rather than the thickness of the unfrozen till layer. Our assumption of a free slip once the melting point is reached is best justified (see Haseloff et al, 2015) if we suppose that the unfrozen bed is hydraulically well-connected, so that the water pressure in the parts of the bed that have just become unfrozen quickly equilibrates with water pressure elsewhere under the ice stream (and hence basal friction is comparable to the rest of the active ice stream). Shear stresses experienced by the margins of the ice stream are large compared with basal drag throughout the ice stream (Haseloff et al, 2015), and this implies that basal friction is small at leading order everywhere where the melting point is reached. There are undoubtedly other, more elaborate models for basal shear stress at the unfrozen bed; ours is the simplest possible case to analyse.”

(3) I was very surprised to read that the authors are unable to incorporate the temperature-dependence of the viscosity and the basal yield stress because of the large computational times apparently required to capture this. After all, this is “just” a 2D model? Ignoring the temperature-dependence of these two parameters strikes me as a very significant drawback of this analysis, both because both parameters are sensitively dependent on temperature (e.g., likely multiple orders of magnitude in the viscosity) and because both of these parameters are very consequential for margin migration. I am convinced that a more complete solution to this problem within reasonable computational time is well within the reach of modern numerics. That being said, I realize that switching the numerical approach would require an unreasonable amount of time by the authors, so I think the current route is acceptable. That being said, I suggest that the authors qualify the generality of their solutions very clearly and point out that ignoring temperature-dependence is likely not a minor detail.

We agree with the reviewer that in particular ignoring the temperature-dependence of the subtemperate yield stress is not a minor detail. However, in Haseloff et al (2015), figure 4a we show explicitly that a grid refinement to approximately 10^{-6} ice thicknesses is necessary to correctly model thermally-driven margin migration for $O(1)$ parameters of α and Pe . Even further grid refinement is necessary to resolve the very small nested thermal boundary layers for the case of $\alpha \gg 1$ and $Pe \gg 1$, which required us to refine the grid down to 10^{-9} ice thicknesses. The necessity of this grid refinement has been confirmed with the analytic solutions provided in different asymptotic limits, which match the numerical solutions well. In the presence of subtemperate slip, this grid refinement has to be extended over the length of the subtemperate slip region, leading to ice-thickness wide regions with extremely high grid refinement. Additionally, each calculation of V_m requires 20 or more individual calculations within the bisection method, as the migration velocity has to be found iteratively. All these requirements make the computational effort involved here significantly greater than for more conventional two-dimensional computational problems, in which the decoupling that we use to our advantage here would not be of great consequence.

Concerning the temperature-dependence of the viscosity, we have added the following text to the discussion:

“However, our boundary layer model is not intended to reproduce velocity and temperature profiles over the entirety of an ice stream, but to focus on processes in the ice stream margin that to leading order control margin migration. The asymptotic analysis of sections 4.2-4.4 would remain structurally the same if we accounted for a temperature-dependent viscosity, though the form of the englacial dissipation term would change, as would, in most cases, the velocity. The most robust result in this regard is likely to be the $\tau \sim O(1)$ result of section 4.4, in particular the result (50) for small τ and a wide subtemperate slip region. For $\tau \sim O(1)$ or small, englacial dissipation affected by ice temperatures does not enter into the leading order basal energy balance that determines the margin migration rate, and the advection velocity that appears in that energy balance problem is set by flow at the ice thickness scale. In other words, the thermal boundary layer described by equation (47) is not changed by having a temperature-dependent viscosity. For a wide subtemperate slip region, where lateral flow takes the form of a plug flow, it then turns out that the derivation of equation (50) in the present paper (see section S6 of the supplementary material) remains the same for a temperature-dependent viscosity. Advection above the basal thermal boundary layer then simply preserves the vertical temperature profile imposed by far field conditions in the ridge, and the same depth-integrated calculation as in the supplementary material can be applied.”

References:

Aschwanden, A., Bueler, E., Khroulev, C., and Blatter, H.: An enthalpy formulation for glaciers and ice sheets, *Journal of Glaciology*, 58, 441–457, doi:10.3189/2012JoG11J088, 2012.

Hewitt, I. J. and Schoof, C.: Models for polythermal ice sheets and glaciers, *The Cryosphere*, 11, 541–551, doi:10.5194/tc-11-541-2017, <https://www.the-cryosphere.net/11/541/2017/>, 2017.

Jacobson, H. P. and Raymond, C. F.: Thermal effects on the location of ice stream margins, *Journal of Geophysical Research*, 103, 12 111– 12 122, doi:10.1029/98JB00574, 1998.

Raymond, C.: Shear margins in glaciers and ice sheets, *Journal of Glaciology*, 42, 90–102, 1996.

Robel, A., Schoof, C., and Tziperman, E.: Rapid grounding line migration induced by internal ice stream variability, *Journal of Geophysical Research*, 119, 2430–2447, doi:10.1002/2014JF003 251, 2014.

Schoof, C.: On the mechanics of ice-stream shear margins, *Journal of Glaciology*, 50, 208–218, doi:10.3189/172756504781830024, 2004.

Schoof, C. and Hewitt, I.: A model for polythermal ice incorporating gravity-driven moisture transport, *Journal of Fluid Mechanics*, 797, 504–535, doi:10.1017/jfm.2016.251, 2016.

Suckale, J., Platt, J. D., Perol, T., and Rice, J. R.: Deformation-induced melting in the margins of the West Antarctic ice streams, *Journal of Geophysical Research*, 119, 1004–1025, doi:10.1002/2013JF003008, 2014.

The role of subtemperate slip in thermally-driven ice stream margin migration

Marianne Haseloff^{1,2}, Christian Schoof², and Olivier Gagliardini³

¹Atmospheric and Oceanic Sciences Program, Princeton University, Princeton, NJ 08540, USA

²Department of Earth, Ocean and Atmospheric Sciences, The University of British Columbia, Vancouver, BC V6T 1Z4, Canada

³Univ. Grenoble Alpes, CNRS, IRD, Grenoble INP, IGE, 38000 Grenoble, France

Correspondence to: Marianne Haseloff (marianne.haseloff@earth.ox.ac.uk)

Abstract. The amount of ice discharged by an ice stream depends on its width, and the widths of unconfined ice streams such as the Siple Coast ice streams in West Antarctica have been observed to evolve on decadal to centennial timescales. Thermally-driven widening of ice streams provides a mechanism for this observed variability through melting of the frozen beds of adjacent ice ridges. This widening is driven by the heat dissipation in the ice stream margin, where strain rates are high, and at the bed of the ice ridge, where subtemperate sliding is possible. The inflow of cold ice from the neighboring ice ridges impedes ice stream widening. Determining the migration rate of the margin requires resolving conductive and advective heat transfer processes on very small scales in the ice stream margin, and these processes cannot be resolved by large scale ice sheet models. Here, we exploit the thermal boundary layer structure in the ice stream margin to investigate how the migration rate depends on these different processes. We derive a parameterization of the migration rate in terms of parameters that can be estimated from observations or large scale model outputs, including the lateral shear stress in the ice stream margin, the ice thickness of the stream, the influx of ice from the ridge, and the bed temperature of the ice ridge. This parameterization will allow the incorporation of ice stream margin migration into large-scale ice sheet models.

1 Introduction

The Siple Coast ice streams are fast moving regions within the West Antarctic ice sheet. They exhibit temporal changes on decadal to centennial time scales in their spatial configuration, for example slow down and reactivation cycles and changes in ice stream width (Stephenson and Bindschadler, 1988; Retzlaff and Bentley, 1993; Harrison et al., 1998; Hamilton et al., 1998; Echelmeyer and Harrison, 1999; Fahnestock et al., 2000; Conway et al., 2002; Catania et al., 2006; Catania et al., 2012; Bindschadler et al., 2000; Stearns et al., 2005; Hulbe and Fahnestock, 2007). The widening and narrowing of ice streams can strongly affect mass discharge from an ice stream: simplified ice stream models show that the ice stream velocity and discharge strongly increase with stream width (Raymond, 2000). Correctly modeling the evolution of ice streams, including the migration of their margins, is therefore essential for reliable predictions of the evolution of the Antarctic ice sheet (Bamber et al., 2000)

Ice streams are bordered by slowly moving ice, called ice ridges, and the close proximity of fast to slowly moving ice is reflected in a sharp gradient in basal resistance between ridge and stream (Bentley et al., 1998). The question of margin

migration is tightly linked to the question of how this gradient is sustained. In the absence of freezing in the bed, subglacial drainage can in principle widen ice streams, if water transport is in the direction of effective pressure gradients (Haseloff, 2015). In this scenario, infinite widening can be suppressed by the formation of ice ridges (e.g. in Kyrke-Smith et al., 2014). This leads to a gradient in ice overburden pressure that counteracts gradients in effective pressure, so that there is no net water pressure gradient driving flow of water towards the ridge. Alternatively, Perol et al. (2015) propose the existence of a channel co-located with the ice stream margin, which theoretically locks the position of the margin into place: margin migration now requires a reorganization of the subglacial drainage system.

However, if freezing in the bed is possible, a thermal barrier can form in the bed which suppresses widening through subglacial drainage (Haseloff, 2015). The existence of such a thermal barrier is supported by radar observations under some ice streams and ridges, where strong contrasts in basal reflectivity from stream to ridge have been interpreted as transitions from a temperate to a frozen bed (Bentley et al., 1998; Catania et al., 2003). ~~Under these conditions, the inwards migration of ice streams can now be the result of insufficient heat dissipation over the width of the entire ice stream, which leads to freezing of the ice stream bed.~~ Under these conditions, the inwards migration (or narrowing) of an ice stream requires freezing of the entire sediment column (Schoof, 2012, appendix B). As melt water can be supplied to sections of the bed with active freezing from other regions of the ice stream via subglacial drainage, this necessitates taking into account the ice-stream-wide energy balance. Consequently, the inwards migration of ice streams is the result of insufficient heat dissipation over the width of the entire ice stream (Haseloff, 2015). However, as shown in Haseloff (2015) this process can at least in principle be modelled with large-scale ice sheet models without recourse to a boundary layer.

In this scenario the outwards migration of ice stream margins requires melting of the frozen sediment under the ice ridge. By contrast with a narrowing ice stream, it is however not necessary for the entire thickness of the sediment column to melt out: only part of it needs to be unfrozen to permit sliding, and we will later idealize this by assuming that sliding is possible as soon as the melting point is reached at the bed. This however also underlines the asymmetry between widening and narrowing of an ice stream, which motivates us to focus on the harder problem of widening, which requires heat to be transferred into the bed. Several studies show that a strong gradient in basal resistance created by such a thermal transition leads to significant englacial heat production in the ice stream margins (Raymond, 1996; Jacobson and Raymond, 1998; Schoof, 2006; Suckale et al., 2014; Perol and Rice, 2015). Combined with conductive heat transfer, this heat dissipation can lead to the outwards migration of the margins by warming the bed outside the active ice stream (Schoof, 2012; Haseloff et al., 2015). This migration is counteracted by advective cooling through the inflow of ice from the ice ridge, driven by an elevation difference between ice ridge and ice stream. The rate of migration is highly sensitive to the relative strength of these two processes (Jacobson and Raymond, 1998; Haseloff et al., 2015).

Existing studies that derive a migration rate from this competition between dissipation, conduction, and advection (Schoof, 2012; Haseloff et al., 2015) assume that the transition from a temperate to a frozen bed is co-located with an abrupt transition from free slip to no slip. However, it is unlikely that such a transition occurs in reality: the basal shear stress goes to infinity at a no-slip to free-slip transition. Therefore a no slip boundary condition on the cold side requires slip to be suppressed there for any amount of basal shear stress (Fowler, 2013). Instead, we expect sliding to occur, either due to mechanical failure or due to

a residual premelted water film at the ice–bed contact (Fowler, 1986; Echelmeyer and Zhongxiang, 1987; Cuffey et al., 1999; Schoof, 2004; Platt et al., 2016; Elsworth and Suckale, 2016). Both of these processes would lead to subtemperate sliding, that is, sliding at temperatures below the melting point. Additionally, the high stress concentrations may be alleviated by mechanical failure or damage production in the ice itself (Pralong and Funk, 2005).

5 In the presence of subtemperate slip, we expect significant changes to the velocity field, which is responsible for advection of heat, and to the spatial distribution of heat dissipation. In particular, heat is then dissipated at the frozen ice–bed interface. This is the very location where warming has to occur in order for the ice stream margin to migrate outwards. We therefore expect subtemperate slip to have a significant influence on the rate at which ice stream margins can migrate.

To determine the rate of margin migration, we have to consider the thermal and mechanical transitions from ice ridge to
10 ice stream flow, which take place over a distance of just a few ice thicknesses. This is narrow in comparison to the width of the ice ridge and the ice stream, and can be captured by a boundary layer model (Haseloff et al., 2015). The physics captured by the boundary layer model is not necessarily included in large scale ice sheet models, and requires very high resolution of the computational grid (Haseloff et al., 2015). The purpose of this paper is therefore twofold: (i) to use the margin boundary layer model of Haseloff et al. (2015) to investigate how subtemperate slip changes the heat production and temperature field
15 in the ice stream margin, and thereby the rate at which ice streams can migrate outwards and (ii) to derive a parameterizations of the margin migration rate which can be used in large scale ice sheet models. Both of these points go beyond the work in Haseloff et al. (2015): the parameterizations we derive in particular show how the limit of rapid advection of heat across the shear margin can be used to simplify the boundary layer model and arrive at tractable forms of the migration rate that could be implemented in computational models either in the form of semi-analytical formulae or lookup tables.

20 This paper is laid out as follows: we state the model in section 2. Typical solutions of the model are presented in section 3, where we also explain how we determine the migration velocity. The dependence of the migration velocity on forcing parameters is determined in section 4, and in sections 4.3–4.4 we derive a parameterization of the migration velocity as function of these forcing parameters. We discuss our results in section 5.

2 The model

25 The model for ice stream margins we use here is derived in Haseloff et al. (2015). Let (x, y', z) be a fixed coordinate system. The model assumes a well-developed ice stream, whose principal flow direction is aligned with the positive x -direction, as shown in figure 1a. The y' -axis is transverse to the ice stream, and the z -axis is vertical. The ice stream is bordered by slowly moving ice ridges. The model for the ice stream margin is located at the transition from ice stream flow regime to ice ridge flow regime, providing the coupling between the two.

30 In contrast to typical ‘shallow’ ice stream and ice sheet models (Fowler and Larson, 1978; Morland and Johnson, 1980; Hutter, 1983; Muszynski and Birchfield, 1987; MacAyeal, 1989; Blatter, 1995; Pattyn, 2003), which assume a small aspect ratio between vertical and lateral extent, the width of the margin region captured by the boundary layer model is of the order of the ice stream thickness. Consequently, the far fields of the ice ridge and ice stream are attained at $y \pm \infty$ (see figure 1b).




Figure 1. Panel a: Large scale model geometry: we assume an ice stream flowing in positive x -direction. At its sides, the ice stream is bordered by slowly moving ice ridges. Panel b: boundary layer geometry, which moves at the rate $v_m = \partial y_m / \partial t$ through the ice stream/ridge geometry shown in panel a.

The asymptotic analysis in Haseloff et al. (2015) shows that the boundary layer evolves rapidly in comparison to the ice stream and ice ridge, and is consequently quasi-static with the only time-dependence arising from the moving transition between a frozen and a temperate bed at $\pm y_m(x, t)$. Moreover, Haseloff et al. (2015) show that the surface of the ice stream margin is flat at leading order and located at $z = h_s$, where h_s is the ice thickness of the ice stream.

- 5 The ice-bed interface is assumed to be flat and located at $z = 0$. Note however that this assumption does not preclude the application of our results to ice streams with a weak topographic control, as found in many regions of the Siple Coast: this assumption merely requires the elevation gradient of the bed to be sufficiently small that the bed elevation does not vary significantly over lateral distance of a few ice thicknesses.

We define the margin location $y' = -y_m(x, t)$ as the point where the bed goes from being at the melting point to below the melting point. To facilitate our analysis, we immediately change to a moving coordinate frame in which the transition from a temperate to a frozen bed is stationary at $y = 0$, see figure 1b. In other words, if y' is the stationary coordinate with $y' = 0$ in the ice stream center and the margin at $y' = -y_m(x, t)$, then the lateral coordinate y of the margin model is linked to y' through

$$y = y' + y_m(x, t), \quad (1)$$

- 15 with $y = 0$ at the transition from frozen to temperate bed (see figure 1b). y increases towards the ice stream, with negative y corresponding to the ice ridge side of the domain. The boundary layer is moving at velocity $-\partial y_m / \partial t$ with respect to the fixed y' -axis.

The boundary layer is effectively two-dimensional: the ice stream is much longer than a single ice thickness, and therefore along-flow variations in mechanical and thermal conditions in the x -direction are much smaller than corresponding variations

in the transverse and vertical directions. In other words, the x -coordinate is passive and (y, z) are the independent variables in the model.

We assume that the thermal state of the ice-bed interface controls the basal boundary conditions for the ice. This requires us to model the thermal response not only of the ice but also of the bed. We therefore specifically include the bed in the domain, and apply a geothermal heat flux at $z \rightarrow -\infty$. At the lateral domain boundaries at $y \pm \infty$, far field boundary conditions are determined by coupling with stream and ridge.

Force balance can be separated into a downstream component, with u denoting the velocity component in the x -direction, and a transverse component in the (y, z) -plane, with (v, w) denoting the corresponding transverse velocity plane. In the downstream direction, the velocity u is determined by

$$10 \quad \frac{\partial}{\partial y} \left(\eta \frac{\partial u}{\partial y} \right) + \frac{\partial}{\partial z} \left(\eta \frac{\partial u}{\partial z} \right) = 0 \quad (2)$$

with η the viscosity. The transverse velocity field is determined by the two-dimensional Stokes and mass balance equations

$$\frac{\partial}{\partial y} \left(2\eta \frac{\partial v}{\partial y} \right) + \frac{\partial}{\partial z} \left[\eta \left(\frac{\partial v}{\partial z} + \frac{\partial w}{\partial y} \right) \right] - \frac{\partial p}{\partial y} = 0, \quad (3a)$$

$$\frac{\partial}{\partial y} \left[\eta \left(\frac{\partial v}{\partial z} + \frac{\partial w}{\partial y} \right) \right] + \frac{\partial}{\partial z} \left(2\eta \frac{\partial w}{\partial z} \right) - \frac{\partial p}{\partial z} = 0, \quad (3b)$$

$$\frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = 0. \quad (3c)$$

15 The viscosity η depends on all three velocity components through Glen's law (Paterson, 1994)

$$\eta = \frac{A^{-1/n}}{2^{1/n}} \left[\left| \frac{\partial u}{\partial y} \right|^2 + \left| \frac{\partial u}{\partial z} \right|^2 + \left| \frac{\partial v}{\partial z} + \frac{\partial w}{\partial y} \right|^2 + 2 \left| \frac{\partial v}{\partial y} \right|^2 + 2 \left| \frac{\partial w}{\partial z} \right|^2 \right]^{\frac{1-n}{2n}} \quad (4)$$

with A the usual viscosity parameter and n the rheology exponent. For simplicity, we neglect the effect of temperature on viscosity here.

The ice stream imposes a lateral shear stress τ_s as a far-field boundary condition. Additionally, the plug flow in the ice stream requires a vertically-uniform across-stream velocity in this far-field, so

$$20 \quad \eta \frac{\partial u}{\partial y} \rightarrow \tau_s, \quad \frac{\partial v}{\partial z} \rightarrow 0, \quad w \rightarrow 0 \quad \text{for } y \rightarrow \infty. \quad (5)$$

Towards the ice ridge, we expect a shearing flow in the transverse direction and negligible flow in the downstream direction

$$\left. \begin{array}{l} u \rightarrow 0 \\ v \rightarrow \frac{(n+2) q_r}{(n+1) h_s} \left[1 - \left(1 - \frac{z}{h_s} \right)^{n+1} \right] \\ w \rightarrow 0 \end{array} \right\} \text{for } y \rightarrow -\infty \quad (6)$$

with q_r the ice flux from the ice ridge towards the ice stream and h_s the ice thickness of the ice stream; the form of v corresponds to a 'shallow ice'-type shearing flow with a temperature-independent rate factor A .

We assume that basal melting has a negligible effect on ice velocities, so

$$w = 0 \quad \text{at } z = 0. \quad (7)$$

On the temperate (stream-ward) side of the ice–bed interface, we assume that the basal shear stress is negligible compared with the shear stresses in the ice, leading to a free slip boundary condition:

$$5 \quad \eta \frac{\partial u}{\partial z} = \eta \frac{\partial v}{\partial z} = 0 \quad \text{for } y > 0, z = 0. \quad (8)$$

In posing this boundary condition for an ice stream that is actively widening, we are assuming that an infinitesimal amount of melting of the bed suffices to allow for slip: once the thermal barrier at the bed is breached, we only need a very thin ice-free layer in order for slip to occur. This is consistent at least with the idea of a plastic bed, where slip can happen on a plane, or with a hard bed.

- 10 To the extent that additional degrees of freedom (other than temperature) are involved in sliding, the main concern would presumably be water pressure at the bed or within the till, rather than the thickness of the unfrozen till layer. Our assumption of a free slip once the melting point is reached is best justified (see Haseloff et al, 2015) if we suppose that the unfrozen bed is hydraulically well-connected, so that the water pressure in the parts of the bed that have just become unfrozen quickly equilibrates with water pressure elsewhere under the ice stream (and hence basal friction is comparable to the rest of the active
- 15 ice stream). Shear stresses experienced by the margins of the ice stream are large compared with basal drag throughout the ice stream (Haseloff et al, 2015), and this implies that basal friction is small at leading order everywhere where the melting point is reached. There are undoubtedly other, more elaborate models for basal shear stress of the unfrozen bed; ours is the simplest possible case to analyse.

- Where the bed is frozen, we consider two different possibilities. The first assumes that no slip is possible, so that the basal
- 20 boundary condition for $y < 0$ is

$$u = v = 0 \quad \text{for } y < 0, z = 0. \quad (9a)$$

We also investigate the possibility of slip at significant basal friction on the frozen bed. For simplicity we use the simplest possible version of this problem and assume that the frozen ice-bed contact fails at a fixed yield stress τ_c (Schoof, 2004, 2010)

$$\left. \begin{array}{l} \text{either} \quad \eta \frac{\partial u}{\partial z} = \tau_c \frac{u}{\sqrt{u^2+v^2}}, \quad \eta \frac{\partial v}{\partial z} = \tau_c \frac{v}{\sqrt{u^2+v^2}}, \quad \sqrt{u^2+v^2} > 0 \\ \text{or} \quad \quad \quad \sqrt{(\eta \frac{\partial u}{\partial z})^2 + (\eta \frac{\partial v}{\partial z})^2} \leq \tau_c, \quad \sqrt{u^2+v^2} = 0 \end{array} \right\} \text{for } y < 0, z = 0. \quad (9b)$$

- 25 The no-slip case (9a) can be obtained formally by putting $\tau_c = \infty$ in (9b).

The upper surface is traction-free, and flat at leading order. In practice, this implies vanishing shear stress and normal velocity, with vanishing normal stress accounted for by a first order correction to the constant leading order surface elevation: **if** the actual upper surface is located at $h_s + s'$ with $s' \ll h_s$, then

$$\eta \frac{\partial u}{\partial z} = \eta \frac{\partial v}{\partial z} = w = 0, \quad 2\eta \frac{\partial w}{\partial z} - p + \rho g s' = 0 \quad \text{at } z = h_s. \quad (10)$$

Thus, even though our model geometry is a parallel-sided strip, it takes into account the first order surface slope towards the ice ridge.

Note that we have formulated the flow problem in such a way that it can be solved without reference to the temperature field. Physically, we however require that the temperature T is below the melting point T_m for $y < 0$, $z = 0$ and that the temperature is at the melting point for $y \geq 0$, $z = 0$. To ensure that these conditions are met, we have to solve the heat equation in the ice ($0 < z < h_s$) and in the bed ($z < 0$). The englacial heat production in the ice is balanced by conductive and advective heat transport, as well as a pseudo-advective term which is the result of the ice stream margin migrating at the rate

$$v_m := \frac{\partial y_m}{\partial t} \quad (11)$$

into the ice ridge. (Physically, this term represents the effect of having to warm the initially cold ice in the ice ridge as the margin migrates into the ridge.) In the bed, no heat is dissipated and the bed is assumed to be static, so that we have a balance between the same pseudo-advective term and diffusion of heat:

$$\rho c_p \left(v_m \frac{\partial T}{\partial y} + v \frac{\partial T}{\partial y} + w \frac{\partial T}{\partial z} \right) - k \left(\frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2} \right) = a \quad \text{for } 0 < z < h_s, \quad (12a)$$

$$\rho_{\text{bed}} c_{p,\text{bed}} v_m \frac{\partial T}{\partial y} - k_{\text{bed}} \left(\frac{\partial^2 T}{\partial y^2} + \frac{\partial^2 T}{\partial z^2} \right) = 0 \quad \text{for } z < 0 \quad (12b)$$

where we used the margin migration velocity v_m as defined by (11). ρ and ρ_{bed} are the densities of ice and bed, respectively, c_p and $c_{p,\text{bed}}$ are specific heat capacities, and k and k_{bed} are thermal conductivities (see table 1). The heat production term a provides the thermo-mechanical coupling:

$$a = \frac{A^{-1/n}}{2^{1/n}} \left(\left| \frac{\partial u}{\partial y} \right|^2 + \left| \frac{\partial u}{\partial z} \right|^2 + \left| \frac{\partial v}{\partial z} + \frac{\partial w}{\partial y} \right|^2 + 2 \left| \frac{\partial v}{\partial y} \right|^2 + 2 \left| \frac{\partial w}{\partial z} \right|^2 \right)^{\frac{1+n}{2n}}. \quad (13)$$

Advection from the ice ridge prescribes a far-field temperature profile $T_r(z)$, while there are no significant lateral temperature gradients towards the ice stream far field

$$T = T_r(z) \quad \text{for } y \rightarrow -\infty, \quad \frac{\partial T}{\partial y} \rightarrow 0 \quad \text{for } y \rightarrow \infty. \quad (14)$$

To be consistent with a conduction-dominated temperature field, we assume for the far field ridge temperature

$$T_r = \begin{cases} T_s + \frac{q_{\text{geo}}}{k} (h_s - z) & \text{if } z \geq 0 \\ T_s + \frac{q_{\text{geo}}}{k} \left(h_s - \frac{k}{k_{\text{bed}}} z \right) & \text{if } z < 0. \end{cases} \quad (15)$$

We will show in section 4 that the migration velocity is sensitive only to the far field temperature at the bed, so the specific form of T_r is immaterial. Here we have assumed a linear profile for simplicity.

At the surface at $z = h_s$, we assume a constant surface temperature T_s , and towards $z \rightarrow -\infty$ we assume a constant geothermal heat flux q_{geo}

$$T = T_s \quad \text{at } z = h_s, \quad -k_{\text{bed}} \frac{\partial T}{\partial z} \rightarrow q_{\text{geo}} \quad \text{as } z \rightarrow -\infty. \quad (16)$$

Finally, at the bed, we impose the following boundary conditions and inequality constraints:

$$T < T_m \quad \text{and} \quad -k \left. \frac{\partial T}{\partial z} \right|^+ + k_{\text{bed}} \left. \frac{\partial T}{\partial z} \right|^- = \begin{cases} 0 & \text{if } \tau_c = \infty \\ \tau_c \sqrt{u^2 + v^2} & \text{if } \tau_c < \infty \end{cases} \quad \text{for } y < 0, z = 0, \quad (17a)$$

$$T = T_m \quad \text{and} \quad -k \left. \frac{\partial T}{\partial z} \right|^+ + k_{\text{bed}} \left. \frac{\partial T}{\partial z} \right|^- < \begin{cases} 0 & \text{if } \tau_c = \infty \\ \infty & \text{if } \tau_c < \infty \end{cases} \quad \text{for } y > 0, z = 0. \quad (17b)$$

The two cases in τ_c correspond to whether subtemperate slip is or is not possible (τ_c finite or infinite, respectively).

- 5 The equalities in (17) arise from the construction of our model: we have chosen the location $y = 0$ to separate a region with a temperate bed ($y > 0$) from one where the bed temperature must be below the melting point, but where it is not otherwise prescribed. In the latter case, a flux condition is necessary to ensure conservation of energy at the bed. Consequently, the temperature inequality in (17a) is an intrinsic part of how we have defined our domain, with the transition from a subtemperate to temperate bed occurring at $y = 0$ in our traveling coordinate system.
- 10 The flux constraint in (17b) by contrast is really a constraint on freezing rates on the temperate side of the thermal transition at the bed, and in imposing it we are assuming that the margin is migrating into the ice ridge at a rate $v_m > 0$. A local analysis of the temperature field near $y = 0$ (see appendix A for a summary and the supplementary material, section S2 and Schoof (2012) for details) demonstrates that, if the temperature constraint in (17a)₁ is satisfied, then the net heat flux out of the bed for small $y > 0$ either approaches $+\infty$ or equals the basal dissipation rate attained at small negative $y < 0$. When the margin
- 15 migrates towards the ice ridge, we assume an infinite basal freezing rate cannot occur on the temperate side of the thermal transition at $y = 0, z = 0$, leaving only the possibility of a finite heat flux (the second version of the inequality in (17b)₂). For the case of no subtemperate slip, this corresponds neatly to having no freezing at the bed at all on the temperate side of the transition (see also Haseloff et al., 2015; Schoof, 2012). For the case of subtemperate slip, our formulation of having an abrupt transition from a finite value of τ_c to free slip implies a discontinuity in basal heat production, and consequently
- 20 requires a finite but non-zero basal freezing rate near the origin. In order to maintain the bed at the melting point, that finite freezing rate has to be compensated for by subglacial drainage (that is, a finite supply of latent heat into the very tip of the temperate bed region). We anticipate that future versions of this model will consider smooth transitions from subtemperate to temperate sliding, for instance by allowing the yield stress to approach zero continuously as function of T . This however is computationally extremely onerous, and we persist with our simpler version of the basal physics here.
- 25 The inequality constraints serve the role of determining a unique migration rate v_m . If we were to dispense with them, we could solve (12a)–(16) with an arbitrary choice of v_m . However, for fixed model parameters, an arbitrary choice of v_m will see one of the inequality constraints in (17) violated, and their role is therefore to specify the migration rate (see supplementary material, section S1, and Schoof (2012)). That migration rate is then a function of geometrical and forcing parameters such as h_s, τ_s, q_r , and q_{geo} . τ_s and q_r in particular represent the far field forcing due to coupling with ice stream and ice ridge.
- 30 For instance, from the perspective of the ice stream, τ_s is the lateral shear stress it imposes on the boundary, while from the perspective of the ice ridge, q_r is the rate at which it supplies mass to the ice stream through the margin.

Table 1. Parameter values used in the sample calculations presented here. Ice stream thickness h_s , lateral inflow of ice from the ridge q_r and marginal lateral shear stress τ_s are highlighted as they represent coupling with ice ridge and ice stream dynamics. h_s and τ_s correspond to the values observed at the upper margin of Whillans ice stream (Harrison et al., 1998), which migrates at a rate of 7 to 30 m year⁻¹ (Hamilton et al., 1998; Harrison et al., 1998; Echelmeyer and Harrison, 1999). The q_r estimate is based on an inflow velocity of 10 m year⁻¹.

Description	Symbol	Value	Units
viscosity parameter	A	$\pm 1.6 \times 10^{-15}$	kPa ⁻³ s ⁻¹
rheological exponent	n	3	
specific heat capacity	$c_p, c_{p,\text{bed}}$	2	kJ kg ⁻¹ K ⁻¹
acceleration due to gravity	g	9.81	m s ⁻²
thermal conductivity	k, k_{bed}	2.3	W m ⁻¹ K ⁻¹
geothermal heat flux	q_{geo}	56×10^{-2}	W m ⁻²
density of ice	ρ	920	kg m ⁻³
density of bed	ρ_{bed}	920	kg m ⁻³
surface temperature	T_s	-20 -25	°C
melting point	T_m	0	°C
basal yield stress of ice ridge	τ_c		kPa
ice stream thickness	h_s	800 900	m
marginal inflow of ice	q_r	10 ⁴	m ² year ⁻¹
marginal lateral shear stress	τ_s	200	kPa

3 Solution of the model

We solve the coupled mechanical and thermal system (2)–(17) with the finite element solver Elmer/Ice (Gagliardini et al., 2013). The computational domain is a relatively large, elongated rectangle which represents the margin cross-section in the (y, z) -plane. It consists of an ice and a sediment sub-domain. We apply the boundary conditions (5)–(6) and (14)–(16) at the relevant sides of the domain, rather than at $\pm\infty$.

The solutions to the problem are uniquely determined by the lateral shear stress τ_s , the ice thickness h_s , marginal inflow of ice from the ridge q_r , the geothermal heat flux q_{geo} , and the surface temperature T_s , in addition to material properties such as thermal conductivity, heat capacity, density, rheological parameters for the ice, and the basal yield stress τ_c . We will treat the majority of these material properties as fixed (see table 1), but consider carefully the effect of changing the basal yield stress.

3.1 Ice flow and heat production

We begin with solutions to the ice flow problem (2)–(10). In our model, we are treating viscosity and basal yield stress as independent from temperature. At present, this is necessary to allow the computation of more than a handful of solutions in a reasonable amount of time, mostly due to the difficulty involved in computing the migration rate from the inequality constraints




Figure 2. Influence of the subtemperate yield stress τ_c on the mechanical fields in the ice stream margin. τ_s is the lateral shear stress in the margin. Rows of panels are labeled *a-d*, with suffixes 1-3 indicating columns. Panels *a*₁–*c*₁: contours of downstream velocity component u at contour intervals of 10 m/year; bold line indicates $u = 10$ m/year. Panels *a*₂–*c*₂: transverse velocity field (v, w) shown as arrows, shading indicates magnitude of the transverse velocity. Panels *a*₃–*c*₃: contours of heat production a at contour intervals 10^{-4} W m^{-3} , $a = 8 \times 10^{-3} \text{ W m}^{-3}$ shown as a bold line. Panels *d*₁–*d*₃ show the velocities and heat dissipation at the bed. Note that no heat is dissipated at the bed where the velocities are zero. Calculations were done with the values listed in table 1 and τ_c as indicated.

(17). The latter requires very fine grids and a costly iterative scheme (Haseloff et al., 2015). We anticipate that future versions of the model will consider two-way coupling between the mechanical and thermal processes, but in our simplified version, we are able to compute solutions to the mechanical problem in isolation: given h_s , τ_s , q_r , τ_c and the rheological properties A and n , we are able to compute velocity and pressure in the ice.

- 5 The downstream velocity u is vertically uniform in the ice stream far field and increases with a prescribed lateral gradient of $2A\tau_s^n$, see (5). Panels *a*₁–*c*₁ of figure 2 show contours of u for different τ_c plotted in the (y, z) plane, computed using the parameters in table 1. Panels *a*₂–*c*₂ show the corresponding across-flow field (v, w) , whose magnitude is significantly smaller than the downstream velocity. Hence gradients of the downstream velocity u dominate the heat production rate, while the velocity (v, w) in the transverse plane accounts for the advection.

In case of no slip on the cold side of the margin ($y < 0$), stress is concentrated around the transition from no slip to free slip at the origin. This translates to very high dissipation rates (a) as shown in figure 2, panel a_3 . It can be shown that there is in fact a singularity with $a \sim 1/r$ in shear stress and consequently the heat production rate at the origin in this case (Rice, 1967, and supplementary material, section S3).

5 For decreasing τ_c (rows b and c of figure 2), the slip transition around the origin is smoothed, as a finite region with $y < 0$ forms where the yield stress τ_c is attained and sliding occurs. Panels d_1 and d_2 show the velocities u and v at the bed ($z = 0$). Allowing slip for $y < 0$ leads to two major changes in the heat production. First, the englacial heat production is reduced and the singularity at $y = 0$ is at least partially alleviated (see figure $2b_3-c_3$; the local analysis with $n = 1$ presented in the supplementary material, section S2 indicates that a may still have a logarithmic singularity). Secondly, heat dissipation is
 10 introduced along the ice–bed interface (see panel d_3 of figure 2). We will discuss below how this shift in the location of heat production affects the temperature field in the ice.

Panels a_2-c_2 of figure 2 show the transition of the transverse velocity component v from a shearing flow to a plug flow. At the boundaries of the domain the vertical velocity component w is zero. However, near the origin, we can observe a downwards motion of ice towards the bed, offsetting the accelerating transverse flow. As for u , allowing for subtemperate slip in a small
 15 region at $y < 0$ leads to a more gradual increase in velocities around the origin (see panels d_1 and d_2 of figure 2).

3.2 Temperature field

To solve the heat equation (12), we need to know the migration rate v_m , which enters as a parameter. Without the inequality constraints (17a)₁–(17b)₂, any value of v_m could be used to solve the heat equation. However, with an arbitrary choice of v_m , one or the other of these two inequalities is generally violated, which allows us to determine v_m with an adapted bisection
 20 method: the upper limit of the search interval is a migration velocity that is too big and therefore leads to a singular freezing rate on the ice stream side for $y > 0$, in violation of (17b)₂. The lower limit of the search interval leads to temperatures exceeding the melting point at the frozen side for $y < 0$, violating (17a)₁. At every iteration, we halve the search interval and continue the search in the upper half if (17a)₁ is violated at the midpoint and in the lower half otherwise (see supplementary section S1 for details).

25 In figure 3 we show temperature fields calculated with migration velocities that satisfy the inequality constraints (17a)–(17b). Each panel shows solutions obtained with a different combination of lateral shear stress τ_s , lateral inflow of cold ice q_r and basal yield stress τ_c . Increasing τ_s while holding all the other parameters constant (top row) leads to more heating of the ice around the slip transition point and inside the ice stream. Consequently, the migration velocity increases. This gives the bed and the ice to the left of the transition point less time to heat up, and temperatures decrease there. The opposite effect can be
 30 observed for an increase of the lateral inflow of ice (second row), which reduces the migration velocity and leads to a warmer bed. However, due to greater advection velocities, temperatures in the ice are lower (figure 3d–f).

Introducing subtemperate slip (decreasing τ_c) leads to additional heat dissipation along the ice–bed interface on the ridge side ($y < 0$). This heat can thaw the frozen bed, thereby increasing the migration velocity. However, as in the case of increasing τ_s this gives the ice less time to heat up. Consequently, temperatures in the ice decrease as τ_c does.




Figure 3. The effect of shear heating (represented by τ_s), advection (represented by q_r), and subtemperate basal sliding (represented by τ_c) on the temperature field in an ice stream margin, same plotting scheme as in figure 2. Solid black lines show contours in 3°C 5°C intervals with the contour of $T = 0^\circ\text{C}$ in red. Thin dashed lines indicate the ice–bed interface and the location of the cold-temperate transition at the ice–bed interface. Red shading indicates the formation of temperate ice. Note that T should be interpreted as a proxy for moisture content when $T > 0$. In this case we can identify $\phi = \rho c_p T / (\rho_w L)$ with ϕ the volumetric moisture content of the ice, ρ_w the density of water, and L latent heat per unit mass. Top row has $q_r = 0$, $\tau_c = \infty$ (i.e., no subtemperate slip), and values of τ_s as indicated above the panel. Middle row has $\tau_s = 200$ kPa and $\tau_c = \infty$. Bottom row has $\tau_s = 200$ kPa and $q_r = 10^4$ m² year⁻¹. The yellow lines in panels g to i mark the extent of the subtemperate slip region. ~~The far-field ice-ridge bed temperature was set to $T_b = -10^\circ\text{C}$ to facilitate visualization, leading to relatively small migration velocities.~~ All other values as listed in table 1.

Note that all the temperature fields shown in figure 3 have $T > 0$ in some parts of the ice. The boundaries of these regions are marked by a bold red line. In these regions of the ice stream, we solve the same heat equation as in the remainder of the domain (see also Schoof, 2012; Haseloff et al., 2015). Obviously, ice cannot have a temperature in excess of its melting point,

and T cannot be interpreted as temperature where $T > 0$. Effectively, we assume a very special case of an enthalpy gradient model (Aschwanden et al., 2012; Schoof and Hewitt, 2016; Hewitt and Schoof, 2017): where $T > 0$, the product $\rho c_p T$ (which is generally the specific heat content per unit volume of ice) must instead be interpreted as the latent heat content per unit volume of the ice. That is, $\rho c_p T$ should be interpreted as $\rho_w L \phi$, where ρ_w is the density of water, L is latent heat per unit mass, and ϕ is the volumetric moisture content of the ice. This allows us to identify $\phi = \rho c_p T / (\rho_w L)$, so T is nothing more than a proxy for moisture content when $T > 0$.

By solving the heat equation where $T > 0$, we make two main assumptions. First, qualitatively, we assume that moisture flows down gradients of moisture, which is the assumption common to enthalpy gradient models and permits the same diffusive model to be applied regardless of whether the melting point has been reached or not. The second, quantitative assumption we make is that the corresponding diffusivity remains the same for cold and temperate regions. This is consistent with prior work, but also an obvious area for future model improvement. We will return to a discussion of the limitations imposed by this assumption in section 5.

~~In these regions of the ice stream margin temperate ice forms, that is, ice at the melting point. T now has to represent melt water fraction, rather than temperature. At this point, we assume that we can use the same heat equation to model cold and temperate ice, which is a very special case of an enthalpy gradient model (Aschwanden et al., 2012). Even though this is not likely to be realistic, we continue with this assumption for now in order to better compare our results with prior work. We will investigate the role of temperate ice formation in a forthcoming publication.~~

Importantly, the region of temperate ice in the bottom two rows of figure 3 does not form directly above the transition from subtemperate slip to free slip at the origin, but is shifted significantly (by up to one several ice thicknesses) towards the ice stream. This is the result of lateral advection of ice and of subtemperate sliding, which generates additional heat and requires less localized englacial heating (compare the local form of the temperature field in appendix A and section S2 of the supplementary material with appendix A of Schoof (2012)). This shift of the temperate region away from the slip transition suggests that the thermal physics around the transition from frozen to unfrozen bed may be relatively unaffected by the choice of temperate ice model (e.g., Aschwanden et al., 2012; Schoof and Hewitt, 2016; Hewitt and Schoof, 2017).

4 Migration velocity as a function of forcing parameters

We now turn to a systematic investigation of the dependence of the migration velocity on the ice ridge and ice stream parameters. As we have pointed out, the solution to the velocity and temperature problem is determined uniquely once we know the applied lateral shear stress τ_s , the inflow rate of cold ice q_r , the geothermal heat flux q_{geo} (or equivalently, the far-field bed temperature on the ridge side of the margin T_b) as well as ice thickness h_s , basal yield strength τ_c and the remaining material properties of ice and bed. Importantly, that solution includes the margin migration rate v_m , which is therefore a function of these physical parameters and material properties: defining the far field basal temperature through

$$T_b = T_s + \frac{q_{\text{geo}} h_s}{k}$$

we can write

$$v_m = \hat{f}(h_s, q_r, \tau_c, \tau_s, T_b; A, c_p, c_{p,\text{bed}}, g, k, k_{\text{bed}}, n, \rho, \rho_{\text{bed}}, T_m, T_s). \quad (18)$$

We emphasize ice thickness h_s , inflow rate q_r , lateral shear stress τ_s , and far-field bed temperature under the ridge T_b in particular as these are parameters that reflect the coupling of the margin to dynamics of ice ridge and ice stream. Conceivably, one might want to run a simulation that relies on simplified models of ridge and stream without having to resolve the margin region itself. The goal of a systematic solution of our margin model in that case is precisely to compute the migration rate v_m as a function of parameters that are controlled by the ridge and the stream: doing so allows the margin to be treated as a free boundary in a larger-scale model. We also emphasize the role of basal yield stress τ_c as we are interested in how allowing for varying degrees of subtemperate sliding changes the relationship between margin migration rate and the forcing the margin experiences from ridge and stream.

It is clear that v_m in (18) depends on a large number of physical parameters, and the computational effort required to find the function appears to be intractable. However, we can reduce the parameter space to a minimum by non-dimensionalising the model: doing so demonstrates that many combinations of parameter values actually correspond to scaled versions of the same calculation, which we then have to do only once. This is done in section 4.1. An additional advantage is that non-dimensionalisation allows us to identify systematically which processes dominate the temperature field and migration rate (see section 4.2). This leads to further simplification that allows us to give semi-analytical versions of (18) in a number of parameter regimes, which we study subsequently in sections 4.3–4.4.

4.1 Non-dimensionalisation

The goal of this section is to express the model in the most succinct form possible. To do so we introduce

$$[z] = h_s, \quad [s'] = \frac{n+2}{n+1} \frac{q_r}{A\tau_s^n h_s^2} \frac{\tau_s}{\rho g}, \quad [u] = Ah_s\tau_s^n, \quad [v] = \frac{n+2}{n+1} \frac{q_r}{h_s}, \quad [T] = T_m - T_s \quad (19)$$

and put $(y, z) = [z](Y, Z)$, $u = [u]U$, $(v, w) = [v](V, W)$, $s' = [s']S'$, $p = \rho g[s']P$, and $T = [T]\mathcal{T} + T_m$. This allows us to absorb quantities such as the ice thickness, inflow rate, and dimensionless lateral shear stress in the ice stream margin into five dimensionless parameters,

$$\alpha = \frac{A\tau_s^{n+1}h_s^2}{k(T_m - T_b)}, \quad \text{Pe} = \frac{(n+2)}{(n+1)} \frac{\rho c_p q_r}{k}, \quad \nu = \frac{T_b - T_s}{T_m - T_s}, \quad \tau = \frac{\tau_c}{\tau_s}, \quad \varepsilon = \frac{n+2}{n+1} \frac{q_r}{A\tau_s^n h_s^2}. \quad (20)$$

Note that our parameter α is defined slightly differently from its counterpart in Schoof (2012) and Haseloff et al. (2015): if we replace T_m by T_s in the denominator of (20)₁, we obtain the version of α used in the latter two papers. We can interpret the parameters above as a dimensionless shear heating rate α , a Péclet number (or measure of advection versus conduction) Pe , a dimensionless measure of the far field bed temperature ν (ν is between 0 and 1 for a ridge bed temperature T_b below the melting point T_m , as we assume here), a dimensionless basal yield stress τ , and a ratio of transverse to downstream velocities ε . Using the values in table 1, we get $\text{Pe} = 314$, $\alpha = 86592$, $\nu = 0.9$, $\varepsilon = 0.0402$.

Note that a large Péclet number is what we would expect in a spatially confined region like an ice stream margin: conduction of heat is relatively ineffective, and advection mostly dominates. Large α reflects the strength of heat production, which must balance the fast rates of advection of cold ice implied by large Pe. Note that ε remains small as long as the across-margin flow is significantly smaller than the downstream flow, which we assume to be the case. Terms of $O(\varepsilon)$ are retained only in order to regularize the viscosity in the ice ridge, where gradients in u vanish. In the numerical solutions presented in this study, we use $\varepsilon = 0.01$, and we have confirmed that smaller values of ε do not change our results. $O(1)$ values of ε would imply that there is significant englacial heat production in the ridge, see Haseloff (2015) and Haseloff et al. (2015). This heat production should prevent the ice ridge bed from remaining frozen, contradicting our basic assumption that the shear margin is co-located with a thermal transition at the bed. τ is poorly constrained, and we will consider different parameter regimes of τ below. Additionally, we have the following ratios of material properties

$$\gamma = \frac{\rho_{\text{bed}} c_{p,\text{bed}}}{\rho c_p}, \quad \kappa = \frac{k_{\text{bed}}}{k}. \quad (21)$$

With the definitions above, the velocity in the downstream direction is determined by the scaled version of (2)

$$\frac{\partial}{\partial Y} \left(\mu \frac{\partial U}{\partial Y} \right) + \frac{\partial}{\partial Z} \left(\mu \frac{\partial U}{\partial Z} \right) = 0, \quad (22)$$

and the across-stream flow is described by (3)

$$15 \quad \frac{\partial}{\partial Y} \left(2\mu \frac{\partial V}{\partial Y} \right) + \frac{\partial}{\partial Z} \left[\mu \left(\frac{\partial V}{\partial Z} + \frac{\partial W}{\partial Y} \right) \right] - \frac{\partial P}{\partial Y} = 0, \quad (23a)$$

$$\frac{\partial}{\partial Y} \left[\mu \left(\frac{\partial V}{\partial Z} + \frac{\partial W}{\partial Y} \right) \right] + \frac{\partial}{\partial Z} \left(\mu \frac{\partial W}{\partial Z} \right) - \frac{\partial P}{\partial Z} = 0, \quad (23b)$$

$$\frac{\partial V}{\partial Y} + \frac{\partial W}{\partial Z} = 0. \quad (23c)$$

μ is the non-dimensional viscosity

$$\mu = \frac{1}{2^{1/n}} \left[\left| \frac{\partial U}{\partial Y} \right|^2 + \left| \frac{\partial U}{\partial Z} \right|^2 + \varepsilon^2 \left(\left| \frac{\partial V}{\partial Z} + \frac{\partial W}{\partial Y} \right|^2 + 2 \left| \frac{\partial V}{\partial Y} \right|^2 + 2 \left| \frac{\partial W}{\partial Z} \right|^2 \right) \right]^{\frac{1-n}{2n}}. \quad (24)$$

20 The boundary conditions in the ice stream far field (5) are now

$$\mu \frac{\partial U}{\partial Y} \rightarrow 1, \quad \frac{\partial V}{\partial Z} \rightarrow 0, \quad W \rightarrow 0 \quad \text{for } Y \rightarrow \infty. \quad (25)$$

Towards the ice ridge, we obtain from (6)

$$U \rightarrow 0, \quad V \rightarrow 1 - (1 - Z)^{n+1}, \quad W \rightarrow 0 \quad \text{for } Y \rightarrow -\infty. \quad (26)$$

At the ice surface, we have from the boundary conditions (10)

$$25 \quad \mu \frac{\partial U}{\partial Z} = \mu \frac{\partial V}{\partial Z} = W = 0, \quad 2\mu \frac{\partial W}{\partial Z} - P + S' = 0 \quad \text{at } Z = 1. \quad (27)$$

As before, basal melting has a negligible effect on ice velocities, and (7) becomes

$$W = 0 \quad \text{at} \quad Z = 0. \quad (28)$$

On the temperate side of the bed, we have free slip from (8):

$$\mu \frac{\partial U}{\partial Z} = \mu \frac{\partial V}{\partial Z} = 0 \quad \text{at} \quad Z = 0, \quad Y > 0. \quad (29)$$

5 On the frozen side of the bed, we can either have no slip (9a):

$$U = V = 0 \quad \text{at} \quad Z = 0, \quad Y < 0, \quad (30a)$$

or we allow subtemperate slip, requiring from (9b)

$$\left. \begin{array}{l} \text{either} \quad \mu \frac{\partial U}{\partial Z} = \tau \frac{U}{\sqrt{U^2 + \varepsilon^2 V^2}}, \quad \mu \frac{\partial V}{\partial Z} = \tau \frac{V}{\sqrt{U^2 + \varepsilon^2 V^2}}, \quad \sqrt{U^2 + \varepsilon^2 V^2} > 0 \\ \text{or} \quad \sqrt{\left(\mu \frac{\partial U}{\partial Z}\right)^2 + \varepsilon^2 \left(\mu \frac{\partial V}{\partial Z}\right)^2} \leq \tau, \quad \sqrt{U^2 + \varepsilon^2 V^2} = 0 \end{array} \right\} \text{for } Y < 0, Z = 0. \quad (30b)$$

Note that the ice flow problem depends only on n , ε , and τ .

10 For later convenience, we write the thermal problem in terms of a reduced temperature Θ through $\mathcal{T} = (1 - \nu)\Theta - (1 - \nu) - \nu Z$: Θ is the deviation from the linear temperature field that would result from geothermal heat flux and conduction alone, given the imposed surface boundary value. Writing the heat equation (12a)–(12b) in terms of Θ yields

$$V_m \frac{\partial \Theta}{\partial Y} + \text{Pe} \left(V \frac{\partial \Theta}{\partial Y} + W \frac{\partial \Theta}{\partial Z} + \frac{\nu}{1 - \nu} W \right) - \left(\frac{\partial^2 \Theta}{\partial Y^2} + \frac{\partial^2 \Theta}{\partial Z^2} \right) = \alpha \mathcal{A} \quad \text{for } 0 < Z < 1, \quad (31a)$$

$$V_m \gamma \frac{\partial \Theta}{\partial Y} - \kappa \left(\frac{\partial^2 \Theta}{\partial Y^2} + \frac{\partial^2 \Theta}{\partial Z^2} \right) = 0 \quad \text{for } Z < 0, \quad (31b)$$

15 with the heat production term

$$\mathcal{A} = \frac{1}{2^{1/n}} \left[\left| \frac{\partial U}{\partial Y} \right|^2 + \left| \frac{\partial U}{\partial Z} \right|^2 + \varepsilon^2 \left(\left| \frac{\partial V}{\partial Z} + \frac{\partial W}{\partial Y} \right|^2 + 2 \left| \frac{\partial V}{\partial Y} \right|^2 + 2 \left| \frac{\partial W}{\partial Z} \right|^2 \right) \right]^{\frac{1+n}{2n}} \quad (32)$$

where we have retained the small $O(\varepsilon^2)$ term, analogous to that in (24). We have also introduced V_m as the dimensionless speed with which the ice stream margin migrates outwards. It is related to the dimensional migration speed through

$$v_m = \frac{k}{\rho c_p h_s} V_m. \quad (33)$$

20 The boundary conditions (14)–(16) are:

$$\Theta = 0 \quad \text{at} \quad Z = 1, \quad (34a)$$

$$\frac{\partial \Theta}{\partial Z} \rightarrow 0 \quad \text{for} \quad Z \rightarrow -\infty, \quad (34b)$$

$$\Theta \rightarrow 0 \quad \text{for} \quad Y \rightarrow -\infty, \quad (34c)$$

$$\frac{\partial \Theta}{\partial Y} \rightarrow 0 \quad \text{for} \quad Y \rightarrow \infty. \quad (34d)$$




Figure 4. Dependence of non-dimensional migration velocity V_m on lateral advection (parameterized by Pe , panel a), far-field bed temperature (parameterized by ν , panel b) and basal shear stress on subtemperate side of the bed (parameterized by τ , panel c). Unless indicated otherwise, we used $Pe = 10$, $\nu = 0.5$, and $\tau = \infty$.

Finally, we have the inequality constraints determining the migration velocity, which are

$$\Theta < 1 \quad \text{and} \quad -\left.\frac{\partial\Theta}{\partial Z}\right|^+ + \kappa \left.\frac{\partial\Theta}{\partial Z}\right|^- = \begin{cases} 0, & \text{if } \tau = \infty \\ \alpha\tau\sqrt{U^2 + \varepsilon V^2}, & \text{if } \tau < \infty \end{cases} \quad \text{for } Y < 0, Z = 0, \quad (35a)$$

$$\Theta = 1 \quad \text{and} \quad -\left.\frac{\partial\Theta}{\partial Z}\right|^+ + \kappa \left.\frac{\partial\Theta}{\partial Z}\right|^- \leq \begin{cases} 0, & \text{if } \tau = \infty \\ \infty, & \text{if } \tau < \infty \end{cases} \quad \text{for } Y > 0, Z = 0. \quad (35b)$$

We have now arrived at a model in which a (unique) dimensionless margin migration velocity V_m is defined by four dimensionless groups that depend on forcing from the ice stream and ridge (α , Pe , ν , and τ) and on the material constants γ , κ and n :

$$V_m = f(\alpha, Pe, \nu, \tau, n, \gamma, \kappa). \quad (36)$$

The remainder of this paper focuses on determining the form of the function f . For comparison with previous work (Schoof, 2012; Haseloff et al., 2015), we additionally assume $\gamma = \kappa = 1$ and $n = 3$, so that V_m can only depend on Pe , α , ν , and τ . We start by treating α and Pe as $O(1)$ parameters in the next section to build intuition for the dependence of V_m on Pe , α , ν , and τ and then investigate the physically more realistic case in which both Pe and α are large in sections 4.3–4.4.

4.2 Migration velocities at $O(1)$ values of α and Pe

In figure 4 we plot V_m against α at fixed values of the other parameters. We see a qualitatively similar picture to the cases described in Schoof (2012) and Haseloff et al. (2015) which assumed a constant viscosity ($n = 1$): outward migration of the margin requires a minimum value of α . Once that is reached, the migration speed increases with α (figure 4a). Increasing advection ($Pe > 0$) in the heat balance reduces the corresponding migration speed V_m . This is because heat production is now not only balanced by migration into the cold ice of the ridge, but also by the influx of cold ice into the boundary layer

(represented by Pe). In addition, for every fixed Péclet number there is a minimum nonzero value of α that generates a positive migration rate, as already found by Haseloff et al. (2015).

Solutions of V_m for different ν between 0.1 and 0.9 are shown in figure 4b. Note that the migration velocity is not very sensitive to changes in ν : for increasing ν , there appears to be a small α -independent shift of V_m to smaller values. Consequently, relative differences in the migration rate should become smaller for increasing values of α . If there are additional dependencies of V_m on ν , these are small enough to be invisible. This behavior is consistent with the analysis for large α that we present later in section 4.3.

We have seen in the discussion of the mechanical fields in section 3.1 that subtemperate slip ($\tau < \infty$) introduces dissipation along the ice–bed interface. Decreasing τ leads to more subtemperate slip and therefore to more dissipation at the bed on the cold side of the margin. Consequently we expect the migration velocity to increase with decreasing τ . Figure 4c confirms this. However, relatively small values of $\tau \lesssim 1$ are needed before there is a noticeable effect on the migration velocity.

A noticeable feature of figure 4 is not only that V_m depends in qualitatively expected ways on α , Pe and τ . We also notice that the dependence of V_m on α often appears to be nearly linear, and that V_m is insensitive to changes in ν . This suggests that, despite f being a function of α , Pe , τ and ν , it may be possible to find parameter regimes in which simple representations of f are available. In the remainder of the paper, we show that it is possible to derive such representations when the dimensionless heating rate α is large. Our estimates in section 4.1 have already indicated that this is the relevant regime that real ice stream margins should find themselves in. We begin by focusing on the case of no slip on the cold side, for which the relationship between V_m and α in figure 4 appears to be nearly linear: we are able to demonstrate that this is the case, and give a formula for the resulting dependence on not only α , but also Pe . Subsequently, we address the more complicated case of finite τ , concluding with formulae for V_m in the parameter regimes of large and moderate subtemperate slip.

4.3 Large heat production without subtemperate slip

We initially restrict ourselves to the case of no subtemperate slip, and consider the case of large α : our estimates in section 4.1 indicate that this limit is likely to apply in practice. Combined with a large heat production rate, the same estimates lead us to expect a large Péclet number: ice is a relatively poor thermal conductor, and advection dominates conduction at the scale of a single ice thickness. Mathematically, this corresponds to advection and heating terms in the heat equation dominating over diffusion in (31) over most of the domain. However, this is no longer true close to the transition from no slip to free slip where there is a small region in which conduction also contributes to the local energy balance. The physics in this region determines the migration rate: conduction is an essential part of how the margin migrates, as it controls how heat production causes the cold part of the bed to warm, and how much heat is extracted from the temperate part of the bed. The analysis below therefore focuses on this small region (known technically as a ‘conductive boundary layer’, see figure 5a).

In what follows we give a brief description of how we can derive a model that ties migration velocity to heat production and transport in the conductive boundary layer. The reader not concerned with the technical details will find the result of this analysis in equation (43).




Figure 5. Boundary layer structure for asymptotics with large α and Pe , without subtemperate slip (panel *a*) and with subtemperate slip (panel *b*). The background temperature profiles are enlargements of typical profiles in this asymptotic limit, with same color scale as in figure 3. In the outer advective boundary layer the temperature field close to the bed is advected from the ice ridge towards the inner conductive boundary layer. In case of a no slip/free slip transition at the bed the conductive boundary layer consists of a small region around the slip transition (panel *a*). In case of subtemperate slip, the conductive boundary layer is a region of small vertical extent which stretches along the length of the subtemperate slip region (panel *b*). Within the conductive boundary layer, heat dissipation is balanced by diffusion.

The non-dimensional mechanical problem (22)–(30b) is parameter-free in the absence of subtemperate slip (the $\tau = \infty$ case above). However, to analyze the temperature field in the boundary layer, we need to know the behavior of flow velocity and heat production near the transition from no slip to slip. In the supplementary section S3, we show that \mathcal{A} , U and (V, W) exhibit power-law behavior near the origin:

$$\left. \begin{aligned}
 \mathcal{A} &\sim A_\vartheta^{-1} R^{-1} \\
 U &\sim \sqrt{\frac{2n}{n+1} A_\vartheta^{\frac{2}{n+1}} + \cos \vartheta A_\vartheta^{\frac{1-n}{1+n}} R^{\frac{1}{n+1}}} \\
 (V, W) &\sim (V_\vartheta, W_\vartheta) R^\beta
 \end{aligned} \right\} \text{ as } R = \sqrt{Y^2 + Z^2} \rightarrow 0 \quad (37)$$

with $\beta = 0.5$ for $n = 1$ and $\beta \approx 0.271$ for $n = 3$, and A_ϑ , V_ϑ , and W_ϑ functions of the angle ϑ between the vector (Y, Z) and the Y -axis and independent of any other model parameters. Here, $R = \sqrt{Y^2 + Z^2}$ is distance from the origin. Knowledge of (37) enables us to study the behavior of the temperature field close to $R = 0$.

For large α , heating near the origin behaves as $\alpha \mathcal{A} \sim \alpha R^{-1}$. As described above, we are looking for a region in which this heating rate is partially balanced by conduction. This happens at distances from the origin that scale as $R_\alpha = \alpha^{-1}$. To resolve this region, we set $(Y, Z) = R_\alpha (\tilde{Y}, \tilde{Z})$, $\mathcal{A} = R_\alpha^{-1} \tilde{\mathcal{A}}$ and $(V, W) = R_\alpha^\beta (\tilde{V}, \tilde{W})$ using (37)₁ and (37)₃, and put $\tilde{\Theta} = \Theta$. If the boundary layer sets the migration rate, then the effect of the margin migrating into colder ice must also enter into the energy balance of the boundary layer at leading order. In order for this to happen, we need a large migration velocity with $V_m \sim \alpha$, which we capture by rescaling the migration velocity as

$$\tilde{V}_m = \alpha^{-1} V_m. \quad (38)$$

We can simultaneously consider conditions under which advection due to motion of the ice also contributes to the cooling of the conductive boundary layer, in addition to migration into cold ice. It turns out that this requires $\alpha^{-(1+\beta)}\text{Pe}$ to be of $O(1)$, i.e., the Péclet number must scale as $\text{Pe} \sim \alpha^{1+\beta}$. We therefore put

$$\Lambda = \text{Pe}^{\frac{1}{1+\beta}} \frac{1}{\alpha} \quad (39)$$

5 and consider the case of $\Lambda \sim O(1)$ (known technically as a ‘distinguished limit’). The case of slow advection is captured by taking the limit of small Λ . Fast advection of cold ice into the margin ($\Lambda \gg 1$) does not permit a widening of the ice stream as we are assuming here, and we exclude the case of large Λ from consideration.

With these changes of variables and parameter definitions in place, (31) becomes (neglecting an $O(\alpha^{-1})$ term):

$$\tilde{V}_m \frac{\partial \tilde{\Theta}}{\partial \tilde{Y}} + \Lambda^{(1+\beta)} \left(\tilde{W} \frac{\partial \tilde{\Theta}}{\partial \tilde{Y}} + \tilde{W} \frac{\partial \tilde{\Theta}}{\partial \tilde{Z}} \right) - \left(\frac{\partial^2 \tilde{\Theta}}{\partial \tilde{Y}^2} + \frac{\partial^2 \tilde{\Theta}}{\partial \tilde{Z}^2} \right) = \tilde{\mathcal{A}} \quad \text{for } 0 < \tilde{Z}, \quad (40a)$$

$$10 \quad \tilde{V}_m \frac{\partial \tilde{\Theta}}{\partial \tilde{Y}} - \left(\frac{\partial^2 \tilde{\Theta}}{\partial \tilde{Y}^2} + \frac{\partial^2 \tilde{\Theta}}{\partial \tilde{Z}^2} \right) = 0 \quad \text{for } \tilde{Z} < 0, \quad (40b)$$

with boundary conditions given by the $\tau = \infty$ case in (35) and each variable being replaced by its rescaled version (i.e., Θ being replaced by $\tilde{\Theta}$, Y by \tilde{Y} and Z by \tilde{Z}). This boundary layer model contains only \tilde{V}_m and Λ as parameters; if (as we expect) there is a unique migration velocity \tilde{V}_m which solves this problem for given Λ , then we have $\tilde{V}_m = \tilde{f}(\Lambda)$.

15 However, we are still missing conditions on $\tilde{\Theta}$ at large distances from the origin, as we exit the conductive boundary layer and enter a region in which diffusion does not play the same leading order role (see figure 5a). These far-field conditions dictate how cold the ice that is advected into the conductive boundary layer is and therefore control in part the strength of conductive heat loss. In order to conclude that \tilde{V}_m depends on the parameters in the original scaled model (31) only through Λ , we need to be certain that these far-field conditions on $\tilde{\Theta}$ also depend only on Λ . It turns out that these far-field conditions are determined by heat transport in a slender region near the bed. This region is marked with ‘advective boundary layer’ in figure 5a; it extends
20 above the origin and towards the cold ice ridge. In this region, shear heating is balanced predominantly by advection of cold ice into the margin, and the effect of having to warm up ice towards the melting point as the margin migrates into the ridge. It can be shown that, at leading order, the heat equation in this region again contains only \tilde{V}_m and Λ as parameters, and consequently, that the far-field conditions to (40) depend only on \tilde{V}_m and Λ as required. This is somewhat tedious, and we give details in the supplementary material (section S4). Ultimately, we are able to confirm theoretically that

$$25 \quad \tilde{V}_m = \tilde{f}(\Lambda). \quad (41)$$

Our goal now is to check numerically that this relationship is obtained from direct solutions of (22)–(35) when α and Pe are made sufficiently large, and to find the approximate form of the function \tilde{f} . Note that we have gone from having a complicated function of 16 variables in equation (18) to being able to express the migration rate as a function of a single variable Λ (which in turn depends on α and Pe). Approximating a function of a single variable numerically, for instance in the form of a look-up
30 table, is obviously much simpler than having to solve numerically for a large number of independent variables, justifying the




Figure 6. Asymptotic behavior of $\tilde{V}_m = V_m \alpha^{-1}$ for the case without subtemperate slip. In this case we expect a limiting behavior of $\tilde{V}_m = \tilde{f}(\Lambda)$ for $\alpha \gg 1$ and $\Lambda = Pe^{1/(1+\beta)} \alpha^{-1}$, see (41). Panel a: \tilde{V}_m against α for different values of constant Pe . Convergence to a constant value confirms the limiting behavior predicted by (41). Panel b: \tilde{V}_m against Pe for different values of Λ . Note that holding Λ constant as Pe changes implies that α changes proportional to $Pe^{1/(1+\beta)}$. Panel c: \tilde{V}_m against Λ at constant $Pe = 10^{10}$, corresponding to different values of $\tilde{f}(\Lambda)$. Filled markers show the same data as in panel b. The open triangles and squares additionally show results for two ~~Shown are results for~~ different temperature fields in the ridge far field with same non-dimensional bed-temperature $\Theta = 0$ at $Z = 0$, as indicated in the legend. ($\Theta_1 = 0, \Theta_2 = Z^2 - Z, \Theta_3 = \nu(1-\nu)^{-1}[(1-Z)^4 + Z - 1]$).

perhaps somewhat obscure procedure that has led us to this point (and its equivalent forms for other parameter regimes to follow later).

The limiting behavior (41) can also be written as

$$V_m = \alpha \tilde{f} \left(\frac{Pe^{1/(1+\beta)}}{\alpha} \right).$$

Immediately, we see that for no advection ($Pe = 0$) we expect a linear relationship between migration rate V_m and heating rate α , which the results in Schoof (2012) and Haseloff et al. (2015) already hinted at for large α . In fact, a linear relationship does not require vanishing Pe : it suffices that $Pe^{1/(1+\beta)}/\alpha \rightarrow 0$. We confirm this behavior numerically in figure 6a, where \tilde{V}_m is plotted against α for different fixed values of Pe . \tilde{V}_m converges to approximately 1.68 for each value of Pe , with the rate of convergence dependent on Pe (figure 6a).

A more general scenario is to consider $\alpha \rightarrow \infty$ and $Pe \rightarrow \infty$ in such a way that Λ is finite, in which case we cannot neglect the effect of advection. Figure 6b shows the convergence of \tilde{V}_m to its limiting form $f(\Lambda)$ as we make Pe (and hence α) large while holding Λ fixed. Note that holding Λ fixed means that α must grow in lock-step with $Pe^{1/(1+\beta)}$. The approach to the limit can be relatively slow, though the limiting value gives a good order-of-magnitude estimate of the actual migration rate even for smaller values of Pe .

Finally, by plotting the converged values of \tilde{V}_m at large Pe against Λ , we can find the function $\tilde{f}(\Lambda)$ in (41). Figure 6c shows \tilde{V}_m plotted against Λ for a fixed value of $Pe = 10^{10}$, which is large enough for the limiting value to have been approached closely in all the examples shown in figure 6b. We can fit a linear relationship to the computational data, of the form

$$\tilde{f}(\Lambda) = C_\alpha - C_{Pe}\Lambda, \quad (42)$$

with $C_\alpha \approx 1.68$ and $C_{Pe} \approx 0.19$, preserving the limiting value of $\tilde{f}(0)$ identified above. Written in terms of the original migration velocity $V_m = \alpha\tilde{f}(\Lambda)$, this is the same as

$$V_m = C_\alpha\alpha - C_{Pe}Pe^{\frac{1}{1+\beta}}, \quad (43)$$

for $\alpha \gg 1$ and $Pe \gg 1$. As previously noted (see also Schoof, 2012; Haseloff et al., 2015), a finite heating rate α is required in order to cause outward migration of the margin, and the formula above is only valid for arguments α and Pe that ensure $V_m > 0$.

The migration rate in the limit of large α and Pe is set by heat generation and transport in a small conductive boundary layer near the no-slip-to-free-slip transition. As we have discussed above, the conductive boundary layer is subject to the advection of cold ice from the far field. That advection takes place from the ice ridge towards the margin, and crucially, the ice that eventually enters the conductive boundary layer always remains close to the bed. As a result, the conductive boundary layer is not sensitive to the details of the temperature profile with which ice enters the margin from the ridge, except for the basal temperature of ice in the ridge: ice at higher elevations simply passes over the boundary layer, and does not affect the energy balance that controls the migration rate at leading order. In technical terms, the far field boundary conditions on $\tilde{\Theta}$ come from matching with the advective boundary layer alluded to above, which itself occupies only a small region near the bed and therefore has inflow boundary conditions dictated by the near-bed temperature prescribed in the limit $Y \rightarrow -\infty$, see the supplementary material, section S4 for details.

We can confirm computationally from solutions to (31) that V_m is insensitive to the temperature profile imposed on the left-hand boundary. Still using large values of α and Pe , we solve (31) with the purely diffusive temperature profile prescribed in (34c) replaced by several nonlinear ones that have the same temperature at the bed as (34c), but with a steeper temperature




Figure 7. Panel *a*: asymptotic behavior for $\tau \gg 1$ where we expect a limiting behavior of $\tilde{V}_m = \tilde{g}(\Lambda, \Gamma)$ for $\alpha \gg 1$, see (46). Convergence to constant values confirms the limiting behavior. Note that α is plotted on a logarithmic scale. Panel *b*: velocities against $\Gamma = \tau^{-(n+1)}\alpha$ at constant values of $\Lambda = Pe^{1/(1+\beta)}\alpha^{-1}$ as indicated by color and $\alpha = 10^{7.25}$. Note that $\Gamma \rightarrow 0$ corresponds to $\tau \rightarrow \infty$, the limit without subtemperate slip.

gradient near the bed. The corresponding migration rates are displayed as open (uncolored empty) markers in figure 6c, and we find close agreement between results obtained from different far-field temperature profiles.

4.4 Large heat production with subtemperate slip

In the last section, we have considered large dissipation rates and rapid advection of ice, but no subtemperate slip. The migration velocity V_m is then determined by heat production and transport in a small conductive layer around the no-slip-to-slip transition. The extent of that boundary layer scales as $R_\alpha = \alpha^{-1}$. Here, we extend the analysis for large α and Pe to account for subtemperate slip. When we allow for subtemperate slip in our model, sliding occurs on a patch of bed of finite size R_c , and the size of that patch relative to the size of the diffusive boundary layer R_α becomes a key consideration (see figure 5b). We need to distinguish two basic cases, $R_\alpha \sim R_c$ and $R_c \sim O(1)$. We treat the former case first. Our modus operandi also remains the same as in the previous section: by rescaling the dimensionless temperature model (31) to capture the leading order behavior in the conductive boundary layer that determines the migration velocity, we derive a simplified form for the migration rate V_m as a function of the dimensionless parameters α , Pe , ν and τ , and test that relationship by solving (31) directly in the appropriate parameter regime.

4.4.1 A small slip region: $\tau \sim \alpha^{1/(n+1)} \gg 1$

Consider a slip region that is similar in size to the conductive boundary layer of section 4.3. To have such a small slip region, the dimensionless yield stress τ must be large. τ scales as $\tau \sim |\partial U / \partial Y|^{1/n}$ and with from (37) we find that dimensionless stresses in the ice scale as $R_\alpha^{-1/(n+1)} = \alpha^{1/(n+1)}$ in the conductive boundary layer of section 4.3. These must now be comparable to

the dimensionless yield stress of the bed, hence

$$\tau \sim \alpha^{1/(n+1)}.$$

As we reduce τ from an effectively infinite value (so there is no slip, as in section 4.3) until it is comparable to $\alpha^{1/(n+1)}$, the magnitudes of velocity components and heat production rate in the conductive boundary layer remain the same as in section 4.3, but the dependence of velocity and heat production on position starts to change, so the analytical formulae (37) no longer apply directly. (Technically, these formulae remain valid at distances r from the origin for which $R_\alpha \ll r \ll 1$.) Knowing that the magnitudes remain the same, we can however use the same rescaling as in section 4.3, put $R_\alpha = \alpha^{-1}$ and set $(Y, Z) = R_\alpha(\tilde{Y}, \tilde{Z})$, $\mathcal{A} = R_\alpha^{-1}\tilde{\mathcal{A}}$, $U = R_\alpha^{1/(n+1)}\tilde{U}$, $(V, W) = R_\alpha^\beta(\tilde{V}, \tilde{W})$, and $\tilde{\Theta} = \Theta$.

The resulting mechanical problem is detailed in the supplementary material, section S5. We do not go into detail here: the point is that the velocity $(\tilde{U}, \tilde{V}, \tilde{W})$ and hence the heat production rate $\tilde{\mathcal{A}}$ are fully determined if the ratio of yield stress τ to typical stress level $\alpha^{1/(n+1)}$ in the conductive boundary layer is given. We write that ratio (effectively, a dimensionless slip parameter) here in the form

$$\Gamma = \tau^{-(1+n)}\alpha. \quad (44)$$

Note that section 4.3 effectively treated the case of no slip, $\Gamma = 0$, and we seek to generalize this here. The corresponding problem for temperature $\tilde{\Theta}$ again takes the form (40). In addition to $\tilde{\mathcal{A}}$ now being dependent on the slip parameter Γ , the boundary conditions at the bed also depend on Γ : we have from (35a)–(35b) that

$$-\left.\frac{\partial\tilde{\Theta}}{\partial\tilde{Z}}\right|^+ + \left.\frac{\partial\tilde{\Theta}}{\partial\tilde{Z}}\right|^- = \Gamma^{\frac{-1}{n+1}}|\tilde{U}| \quad \text{on } \tilde{Z} = 0, \tilde{Y} < 0 \quad (45a)$$

$$\tilde{\Theta} = 1 \quad \text{on } \tilde{Z} = 0, \tilde{Y} > 0 \quad (45b)$$

with the inequality constraints on flux and temperature still taking the same form as in (35a)–(35b).

In other words, the conductive boundary layer problem now depends on an additional parameter through Γ : with the abrupt transition from no slip to free slip (section 4.3), we had $\tilde{V}_m = \tilde{f}(\Lambda) \approx C_\alpha - C_{\text{Pe}}\Lambda$ while now we have

$$\tilde{V}_m = \tilde{g}(\Lambda, \Gamma), \quad (46)$$

with $\tilde{f}(\Lambda) = \tilde{g}(\Lambda, 0)$. To confirm that (46) holds, we fix Λ and Γ to specific values and increase α . This implies that $\text{Pe} = (\alpha/\Lambda)^{1+\beta}$ and $\tau = (\alpha/\Gamma)^{1/(1+n)}$ both increase in lock-step with α . Convergence of \tilde{V}_m to a value that depends only on Λ and Γ for $\alpha \rightarrow \infty$ then confirms (46).

Owing to the high computational cost of solving (31), especially in the limits of large Pe and α (when advection dominates and the conductive boundary layer requires high mesh resolution), we test for convergence of \tilde{V}_m in this parameter limit for a total of 39 combinations of Λ and Γ , where we have restricted ourselves to only three values of Λ and focused on the effect of changing the slip parameter Γ . Figure 7a shows the expected convergence in the limit of large α . As in section 4.3, convergence often requires quite large values of α . The limiting value of \tilde{V}_m is plotted against Γ for the three different values of Λ used in figure 7b.

As already observed in section 4.2, decreasing the basal shear stress increases the migration velocity due to increased dissipation on the cold side of the bed. We observe the same here, in the sense that increasing $\Gamma \propto \tau^{-(n+1)}$ increases the migration velocity. We also reproduce the limiting behavior for $\Gamma \rightarrow 0$ ($\tau \rightarrow \infty$), in which case we expect to reproduce the migration rate predicted for the no-slip-to-free-slip transition case of section 4.3. In fact, figure 7b shows that relatively large values of $\Gamma \approx 10$ are needed to see a significant departure of migration velocity \tilde{V}_m from its limiting value for $\Gamma = 0$. Unfortunately, computational constraints make it impossible for us to find a simple closed-form approximation for \tilde{g} analogous to (42): we simply do not have enough data to construct such an approximation. However, we will present a solution to this issue in section 4.4.3.

4.4.2 An $O(1)$ slip region: $\tau \sim O(1)$

We now turn to the case of $\tau \sim 1$, in which the lateral shear stress τ_s exerted by the ice stream on the margin is comparable with the yield strength τ_c of the frozen bed. In this limit, we expect the subtemperate slip length scale to be comparable with ice thickness, so $R_c \sim O(1)$ (see also figure 2). The region in which there is significant dissipation along the bed is now much larger than in the previous section. As a result, the region in which dissipation is balanced substantially by conduction now has a horizontal extent comparable with ice thickness, too. For large α , we however still have a conductive boundary layer whose vertical extent remains small: large temperature gradients are needed in order to account for the large amounts of dissipation, and such temperature gradients have to correspond to $O(1)$ temperature changes occurring over small vertical distances. In fact, that vertical distance still scales as $R_\alpha = \alpha^{-1}$. The primary difference is therefore that the boundary layer now has an $O(1)$ extent in the horizontal, equal to the size of the slip region.

With an $O(1)$ region of slip at the bed, there are no simplifications to the mechanical problem (22)–(30b): we are no longer confining our attention to a small region around the origin. The solution to the mechanical problem is fully specified if we know τ , so (U, V, W) are functions of τ only. The horizontal velocity components (U, V) are of $O(1)$. If we are concerned with the conductive boundary layer near the bed, then we only need the vertical velocity component near the bed. Since $W = 0$ at the bed itself, we find that $W \sim Z$ in the boundary layer. This allows us to rescale as $(Y, Z) = (Y^*, R_\alpha Z^*)$, $(U, V, W) = (U^*, V^*, R_\alpha W^*)$, $\mathcal{A} = \mathcal{A}^*$, and $\Theta = \Theta^*$. The leading order version of the heat equation (31) is thus (neglecting terms of $O(\alpha^{-1})$)

$$V_m^* \frac{\partial \Theta^*}{\partial Y^*} + \Omega \left(V^* \frac{\partial \Theta^*}{\partial Y^*} + W^* \frac{\partial \Theta^*}{\partial Z^*} \right) - \frac{\partial^2 \Theta^*}{\partial Z^{*2}} = 0 \quad (47a)$$

$$V_m^* \frac{\partial \Theta^*}{\partial Y^*} - \frac{\partial^2 \Theta^*}{\partial Z^{*2}} = 0 \quad (47b)$$

where we defined

$$V_m^* = \frac{V_m}{\alpha^2}, \quad \Omega = \frac{\text{Pe}}{\alpha^2} \quad (48)$$




Figure 8. Panel *a*: asymptotic behavior for $\tau \sim 1$ where we expect a limiting behavior of $V_m^* = V_m \alpha^{-2} = g^*(\Omega, \tau)$ for $\alpha \gg 1$, see (49). The limiting behavior is confirmed by convergence to constant values for $\alpha \gg 1$. Panel *b*: asymptotic behavior of V_m^* against $\Omega = \text{Pe} \alpha^{-2}$ along constant values of τ at values of α where convergence is observed. Dashed lines show the analytical solution (50) which is valid for $\tau \ll 1$.

and retained Ω as an $O(1)$ quantity: doing so with $\alpha \gg 1$ is again to look at a distinguished limit, analogous to treating Λ as $O(1)$ in section 4.3. The rescaled version of the heat flux constraint (35a)₂ is

$$-\left. \frac{\partial \Theta^*}{\partial Z^*} \right|^+ + \left. \frac{\partial \Theta^*}{\partial Z^*} \right|^- = \tau |U^*|.$$

Again, there are far field boundary conditions on Θ^* . These arise purely by advection into the boundary layer, and that advection occurs from the ice ridge, which fixes the far field temperature at $\Theta^* = 0$ as $Y^* \rightarrow -\infty$, so there is no additional parameter dependence through the far-field conditions. The thermal problem (47) contains only the dimensionless parameters Ω and τ (the latter both explicitly and through the velocity field). This indicates that the rescaled migration velocity V_m^* only depends on Ω and τ :

$$V_m^* = g^*(\Omega, \tau) \tag{49}$$

for $\alpha \gg 1$, $\text{Pe} \sim \alpha^2$, and $\tau \sim 1$. Expressed in terms of the original migration velocity V_m , we have

$$V_m = \alpha^2 g^* \left(\frac{\text{Pe}}{\alpha^2}, \tau \right).$$

Once more we confirm that the formula (49) holds by computing V_m^* for different fixed Ω and τ while increasing α . In this case, we can observe convergence at moderately large values of $\alpha \approx 10^2$ to 10^3 (figure 8a). Again, smaller values of τ lead to larger values of V_m^* and increasing Pe (and hence Ω) leads to decreasing V_m^* (figure 8b). Note that the solutions shown are computationally expensive: the conductive boundary layer invariably requires local mesh refinement, and in the parameter regime considered here, it extends over a larger part of the domain, with the size of the region that requires mesh refinement depending on τ . As a result, we are not able to compute solutions for very large values of α at all values of τ . Therefore, computational constraints once more mean that we are unable to sample a large enough region of the two-dimensional parameter space to give a simple formula for the function g^* .

However, for small values of $\tau \ll 1$, it is possible to solve the boundary layer problem analytically, as shown in the supplementary material, section S6. Effectively, this corresponds to finding the limiting behavior of g^* as $\tau \rightarrow 0$, for which we obtain with $n = 3$ that

$$g^*(\Omega, \tau) \sim \frac{1}{\tau} \left[\frac{64}{315\sqrt{\pi}} - \frac{63\sqrt{\pi}}{64} \Omega \tau \right]^2 \quad (50)$$

5 where the solution is valid only when the term in square brackets is positive: as before, a minimum value of α (equivalent to a maximum value of $\Omega \propto \alpha^{-2}$) is required to ensure outward migration of the margin. This limiting form is displayed in figure 8b along with the computationally obtained migration velocities for nominally small values of $\tau = 1$, $\tau = 0.5$ and $\tau = 0.25$ (note that $V_m \sim \tau^{-1}$, so V_m does not approach a finite value, but diverges in a predictable fashion). For $\tau = 1$ and $\tau = 0.5$, there is still a notable difference between the numerical solution and the analytical solution (50), implying that these values of
 10 τ do not yet satisfy $\tau \ll 1$. However, for $\tau = 0.25$, the difference between the numerical solution and the analytical solution is negligibly small.

4.4.3 Moderate slip: $1 \ll \tau \ll \alpha^{1/(1+n)}$

One of the difficulties we still face in making our results directly applicable to large scale models is that we have a closed-form approximation for the migration rate V_m for only two parameter regimes. Both assume large dissipation rates $\alpha \gg 1$, which is
 15 realistic for abrupt ice stream margins. One applies to the case of no subtemperate slip (equation 43), while the other applies to extensive subtemperate slip (equation 50). The obstacle to dealing with more moderate amounts of subtemperate slip is that the functions \tilde{g} and g^* in (46) and (49) are both functions of two independent variables, and that they are computationally expensive to evaluate.

There is one regime in which we can do better and reduce \tilde{g} and g^* effectively to functions of a single variable: both functions
 20 have to be valid representations of the migration velocity V_m in the limit $1 \ll \tau \ll \alpha^{1/(1+n)}$, in which $\Gamma = \tau^{-(1+n)}\alpha$ and τ are both large. In other words, in this parameter regime, we expect that \tilde{g} and g^* give the same answer:

$$V_m \sim \alpha \tilde{g}(\Lambda, \Gamma) \sim \alpha^2 g^*(\Omega, \tau).$$

If we denote the limiting forms of \tilde{g} and g^* in the parameter limit under consideration with a subscript 0, then eliminating Ω and τ in favor of Γ , Λ and α leads to

$$25 \quad \alpha \tilde{g}_0(\Lambda, \Gamma) = g_0^* \left(\Lambda^{1+\beta} \alpha^{\beta-1}, \alpha^{1/(1+n)} \Gamma^{-1/(1+n)} \right).$$

By choosing Pe and τ , we can vary Λ and Γ independently of α . Hence, for any given Γ and Λ , we can now pick $\alpha = \Gamma$ and are left with

$$\tilde{g}_0(\Lambda, \Gamma) = \Gamma^{-1} g_0^* \left(\Lambda^{1+\beta} \Gamma^{\beta-1}, 1 \right), \quad (51)$$

where we only need to evaluate g_0^* at a fixed value of its first argument in order to compute V_m/Γ . Define $g(\chi) = g_0^*(\chi^{1-\beta}, 1)$
 30 for arbitrary χ . It then follows that we can use (51) and $V_m = \alpha \tilde{g}_0(\Lambda, \Gamma)$ to write

$$V_m = \tau^{-(1+n)} \alpha^2 g \left(\tau^{(1+n)} \text{Pe}^{1/(1-\beta)} \alpha^{-2/(1-\beta)} \right). \quad (52)$$




Figure 9. Asymptotic behavior for large α and Pe for the case of intermediate slip ($1 \ll \tau \lesssim \alpha^{1/(1+n)}$) where we expect a limiting behavior of $V_m \tau_c^{1+n} \alpha^{-2} = g(\chi)$, see (52). Panel a: $V_m \tau_c^{1+n} / \alpha^2$ against α for different values of $\chi = \tau_c^{(1+n)} Pe^{1/(1-\beta)} \alpha^{-2/(1-\beta)}$ with Pe fixed between 0 and 10^7 and τ varying between 0.1 and 3.2. Convergence to constant values for $\alpha \gg 1$ confirms (52). Panel b: values of $V_m \tau_c^{1+n} / \alpha^2$ at different χ for $\alpha = 10^5$ and $\tau = 3.2$.

As promised, the migration rate once more reduces to a function g of a single variable.

Finding an approximation to g requires significantly fewer function evaluations. Instead of varying τ , Pe , and α , we only need to vary the argument $\chi = \tau_c^{(1+n)} Pe^{1/(1-\beta)} \alpha^{-2/(1-\beta)}$ of the function g . As before, we first confirm that (52) indeed holds by holding χ fixed and increasing α (figure 9a). Subsequently, we plot the limiting value of $V_m \tau_c^{1+n} \alpha^{-2}$ against χ in figure

5 9b. A simple polynomial fit

$$g(\chi) \approx \left[c_2 (\chi - \chi_0)^2 + c_4 (\chi - \chi_0)^4 \right] \quad (53)$$

with $c_2 = 0.8$, $c_4 = 125$ and $\chi_0 = 0.07$ provides a good representation of the computed migration rates. The expression is again only valid for $0 < \chi \leq \chi_0$: the maximum allowed value of $\chi \propto \alpha^{-2/(1-\beta)}$ corresponds to the minimum dissipation rate α that ensures outward margin migration.

10 5 Discussion and Conclusions

In this study, we have investigated how different physical processes determine the migration widening of ice streams margins that are not topographically confined. We have considered the case in which the transition from fast to slow or no sliding that characterizes a typical ice stream margin is co-located with a thermal transition at the bed. In this scenario, the often intense dissipation of heat generated by the change in sliding behavior can cause the corresponding transition from a temperate to a cold bed to move, and the main objective of this study is to determine the corresponding rate of margin migration into the cold region. This ice stream widening relies on a delicate balance between heat dissipation, heat transport by advection, and conduction to warm the initially cold bed outside the ice stream. We have specifically excluded the case where heat loss

dominates and the margin migrates into the ice stream from consideration here, although similar physics would allow inwards migration to be modeled (Schoof, 2012; Haseloff, 2015).

How the margin location is determined here differs from existing studies of heat transfer processes in Schoof (2004), Suckale et al. (2014), and Elsworth and Suckale (2016). In all of these, the mechanical transition and the thermal transition are not
 5 co-located. Instead, these studies appeal to spatial contrasts in basal friction caused by a heterogeneous drainage system as a mechanism for fixing the margin location independently of the thermal transition at the bed. These approaches and our approach likely represent end-members of how actual ice stream margins migrate, because hydrologically driven margin migration is excluded in our model: freezing in the bed leads to the formation of a thermal barrier which we assume subglacial water cannot penetrate (Haseloff, 2015). Observations suggest that the beds of ice ridges are indeed frozen in some parts (Bentley
 10 et al., 1998; Catania et al., 2003), but that widening of ice streams into these regions is nevertheless possible (Stephenson and Bindschadler, 1988; Fahnestock et al., 2000; Conway et al., 2002; Catania et al., 2012). It is therefore conceivable that drainage-driven ice stream widening and thermally driven ice stream widening are operating in different regions of the Siple Coast ice streams and future work should investigate the interplay between these different processes.

To model the migration of ice stream margins, we solve a coupled model for ice flow and heat transport in the margin. In this
 15 model, the migration rate is determined by imposing constraints on the temperature and heat flux on the cold and warm side of the margin. The migration rate depends on material properties, ice geometry, lateral shear stress in the ice stream margin, and the velocity with which the ice enters the margin from the ridge. These dependencies can be expressed in terms of a small number of non-dimensional combinations of these parameters, although there is no closed form solution and the migration rate is expensive to evaluate on a case-by-case basis through the use of our model. In general, we have been able to establish that
 20 larger lateral shear stresses and less inflow of cold ice favor margin migration, as does a lower basal yield stress on the cold side of the margin.

To go further and provide quantitative parameterizations of the migration rate, we have exploited the fact that heat dissipation is generally large. This has allowed us to construct a number of approximate solutions for migration velocities that we can give
 25 in closed form. Where the different parameterizations we have derived apply depends on the amounts of subtemperate slip, controlled in our model by basal yield stress τ_c . Note that all formulae given below are valid only where they predict $v_m > 0$: in general, there is a minimum value of dissipation rate required to produce any outward migration at all. In all cases, a Glen's law parameter $n = 3$ has been assumed.

For an infinite yield stress, which is equivalent to a sharp transition from no slip to free slip, the migration rate is (43):

$$v_m = 1.68 \frac{A \tau_s^4 h_s}{\rho c_p (T_m - T_b)} - 0.19 \frac{k}{\rho c_p h_s} \left(\frac{5}{4} \frac{\rho c_p q_r}{k} \right)^{0.79} \quad \text{if } v_m \geq 0. \quad (54)$$

30 The parameter A is the viscosity of ice, τ_s is the lateral shear stress in the ice stream margin, h_s is the ice stream thickness, q_r is the ice flux from the ice ridge into the ice stream, T_b is the bed temperature in the ice ridge, T_m is the melting point temperature, c_p is the specific heat capacity of ice, ρ is the density of ice, and k is the thermal conductivity of ice. For an




Figure 10. Margin migration velocities v_m against ice stream width W_s . Equation (54) is valid for no subtemperate slip, $\tau_c = \infty$. Equation (55) is valid for intermediate values of the subtemperate yield stress, formally when $\tau_s \ll \tau_c \ll \tau_s^2 [A h_s^2 / (k T_m - k T_b)]^{1/4}$. Equation (56) is valid for a small subtemperate yield stress (formally when $\tau_c \ll \tau_s$). Violation of the upper limit of validity of equation (55) leads to migration velocities that are less than the migration velocities without subtemperate slip (marked with thin blue line), which is unphysical, as subtemperate slip always increases migration velocities compared to no subtemperate slip.

intermediate yield stress ($\tau_s \ll \tau_c < \infty$), we have from (53)

$$v_m = \frac{k}{\rho c_p h_s} \left(\frac{A \tau_s^4 h_s^2}{k(T_m - T_b)} \right)^2 \left(\frac{\tau_s}{\tau_c} \right)^4 \left[0.8(\chi - 0.07)^2 + 125(\chi - 0.07)^4 \right] \quad \text{if } v_m \geq 0 \quad (55)$$

with χ given by

$$\chi = \left(\frac{\tau_c}{\tau_s} \right)^4 \left[\frac{5}{4} \rho c_p q_r \frac{k(T_m - T_b)^2}{A^2 \tau_s^8 h_s^4} \right]^{1.4};$$

- 5 additionally, we must have $0 \leq \chi \leq 0.07$. **The upper limit on χ corresponds to the lower limit on heat dissipation which is required for outwards migration.** Finally, for a very small yield stress ($\tau_c \ll \tau_s$), we have equation (50)

$$v_m = \frac{k}{\rho c_p h_s} \left(\frac{A \tau_s^4 h_s^2}{k(T_m - T_b)} \right)^2 \frac{\tau_s}{\tau_c} \left[\frac{64}{315\sqrt{\pi}} - \frac{315\sqrt{\pi}}{256} \frac{\rho c_p q_r}{k} \left(\frac{k(T_m - T_b)}{A \tau_s^4 h_s^2} \right)^2 \frac{\tau_c}{\tau_s} \right]^2 \quad \text{if } v_m \geq 0. \quad (56)$$

There are also parameter regimes for which we cannot provide such succinct formulae, but in these regimes the migration rate still increases with τ_s and decreases with increasing lateral inflow of ice q_r .

- 10 Equations (54)–(56) describe margin migration as the result of a balance between englacial heat dissipation, heat dissipation along the ice–bed interface and advective cooling through the inflow of cold ice from the sides. The thermally active region where these processes operate is a small, conductive boundary layer around either the no slip/free slip boundary or along the region of subtemperate slip. While the migration velocity is determined by a balance between heat production and heat transport processes on very small scales, equations (54)–(56) for v_m only require knowledge of dynamic quantities that can be
- 15 obtained in large scale ice sheet models (τ_s , h_s , q_r , T_b , and τ_c).

To illustrate how different geometric conditions alter the migration rate, we assume that the lateral shear stress is determined by $\tau_s = \rho g h_s W_s / L$, assume $L = 1000$ km (the ratio h_s / L represents the ice stream surface slope, which is generally small)

$\tau_s = \rho g \sin \alpha W_s / 2$ with $\sin \alpha = 10^{-3}$ the ice stream surface slope, and plot the migration velocity as a function of ice stream width W_s , figure 10. As one would expect, the migration velocity increases with increasing ice stream width W_s , because wider ice streams are faster and therefore produce higher lateral shear stresses in the margin. All three parameterizations predict migration velocities that are of the order of the real-world migration velocities of 7 m/year to 17–30 m/year established from relatively sparse observations (Hamilton et al., 1998; Harrison et al., 1998; Echelmeyer and Harrison, 1999). Note that the curves representing solutions with subtemperate slip dip below the curve for the non-slip case. These parts of the solution curves corresponds to parameter regimes for which the approximate solutions are not expected to be valid: subtemperate sliding always facilitates margin migration relative to the no-slip case.

Since the parameter ranges where equations (54)–(56) are applicable for a given ice stream depend on the yield strength of the subtemperate region, knowledge of the basal properties is necessary. There are only very few observations which would allow an estimate of the subtemperate basal yield strength (Holdsworth, 1974; Echelmeyer and Zhongxiang, 1987; Cuffey et al., 1999). Using the values reported in Cuffey et al. (1999) for Meserve Glacier, Antarctica, gives an approximate basal yield stress of 380 kPa (neglecting the dependence on the thickness of the interfacial water layers). As the typical lateral shear stress of an ice stream margin can be estimated from $\tau_s = \rho g \sin \alpha W_s / 2$ to be in the range of 180 kPa to 310 kPa (Joughin et al., 2002, table 3), this suggests that equation (55) will be appropriate in most circumstances.

Naturally, migration rate increases with ice stream width, hinting at the possibility of runaway widening of ice streams in a positive feedback. Note however that plotting v_m against W_s is somewhat misleading, as we have assumed the same ice thickness for different ice stream widths. In reality, we expect a negative feedback, where wider ice streams discharge more mass, thereby lowering the ice surface. This would lead to a decrease in englacial and subglacial dissipation, and slow the widening of the ice stream. Investigating this feedback requires combination of the parameterization of v_m with a large-scale ice sheet model.

In principle, incorporating ice stream widening with equation (54)–(56) in large scale models should be possible; depending on the nature of the large scale model, the formula of the migration rate needs to be supplemented explicitly with the continuity conditions on ice thickness and lateral inflow of mass from Haseloff et al. (2015). There are some practical challenges, however: to calculate the migration velocity, the dynamic parameters h_s , τ_s and q_r must be determined at the boundary between ice stream and ice ridge, which might not necessarily align with the mesh or grid of the ice sheet model. Additionally, the migration of this boundary at rates of a few meters per year will likely be below the mesh resolution of continental-scale models. In fact, high resolution is required to resolve ice streams to begin with. Use of equation (54)–(56) will therefore require methods that can adapt to moving ice stream boundaries, similar to methods for grounding lines of marine ice sheets (see e.g., Durand et al., 2009; Gladstone et al., 2012; Pattyn et al., 2012).

Moreover, we have only considered the evolution of an ice stream that is already fully evolved, but the physics governing the position of ice stream tributaries and ice stream inception remains unclear. The position of ice stream tributaries might be determined by geological factors (Peters et al., 2006), but it is conceivable that thermal transitions at the bed might play a role as well and that the pattern of ice stream tributaries is the result of an instability (Hindmarsh, 2009; Brinkerhoff and Johnson, 2015).

Our model as stated in section 2 makes several simplifying assumptions. We allow for subtemperate slip if the basal shear stress exceeds a constant yield stress τ_c to avoid the singular transition from no slip to free slip. However, an abrupt transition from a constant yield stress to free slip (corresponding to a zero yield stress) is still unlikely to occur in realistic situations. If subtemperate slip is facilitated by interfacial films, then the basal yield stress will depend on the temperature of the bed (Gilpin, 1979) and different temperature-dependent sliding laws have been suggested (Shreve, 1984; Fowler, 1986; Wolovick et al., 2014). Such a sliding law requires a two-way coupling between the solution of the mechanical model and the thermal model, potentially introducing feedbacks between these two. We will investigate this in a separate publication.

We also assume that the viscosity in the ice is independent of the ice temperature (i.e., $A = \text{constant}$). Recent studies of the velocity field and temperature in the margin of Whillans ice stream have found that matching observed profiles with numerical model results requires incorporating the temperature-dependence of viscosity (Suckale et al., 2014). However, in contrast to these studies, our boundary layer model is not intended to reproduce velocity and temperature profiles over the entirety of an ice stream. Instead, we focus on the processes in the ice stream margin that to leading order control margin migration. The asymptotic analysis of sections 4.2-4.4 would remain structurally the same if we accounted for a temperature-dependent viscosity, though the form of the englacial dissipation term would change, as would, in most cases, the velocity. The most robust result in this regard is likely to be the $\tau \sim O(1)$ result of section 4.4, in particular the result (50) for small τ and a wide subtemperate slip region. For $\tau \sim O(1)$ or smaller, englacial dissipation affected by ice temperatures does not enter into the leading order basal energy balance that determines the margin migration rate, and the advection velocity that appears in that energy balance problem is set by flow at the ice thickness scale. In other words, the thermal boundary layer described by equation (47) is not changed by having a temperature-dependent viscosity. For a wide subtemperate slip region, where lateral flow takes the form of a plug flow, it then turns out that the derivation of equation (50) in the present paper (see section S6 of the supplementary material) remains the same for a temperature-dependent viscosity. As advection above the basal thermal boundary layer simply preserves the vertical temperature profile imposed by far field conditions in the ridge, the same depth-integrated calculation as in the supplementary material can be applied. ~~and we find that the leading order processes determining the migration velocity are concentrated in a very small, thermally active layer, in which the assumption of a constant temperature is at least better justified.~~

Finally, we have assumed that the dynamics of temperate ice can be represented by a particular version of an enthalpy gradient model (Aschwanden et al., 2012). We expect future iterations of our model to incorporate ice dynamics which can account for gravity-driven moisture transport in temperate ice (e.g., Schoof and Hewitt, 2016). A more sophisticated treatment of temperate ice is likely to be particularly relevant when temperate ice forms near the transition from a cold to a temperate bed: as we have seen, the Peclet number in a shear margin is large, and temperate ice formation down-flow from the cold-temperate transition is consequently unlikely to affect the temperature field close to the transition, as T is dominated by advection. This makes our results with moderate to small τ_c (or, more accurately, with moderate to small τ) likely to be the most robust to changes in the temperate ice model, since temperate ice forms some distance inside the ice stream in that case (see for instance fig. 3). We leave a deeper investigation to future work. ~~Finally, we have assumed that the dynamics of temperate ice can be represented by a stripped-down version of an enthalpy gradient model (Aschwanden et al., 2012). We expect future iterations~~

of our model to incorporate ice dynamics which can account for gravity-driven moisture transport in temperate ice (e.g., Schoof and Hewitt, 2016).

Appendix A: Temperature close to the cold–temperate transition

Here we summarize the behavior of the velocity and temperature fields close a transition from frictional to free slip, based on calculations given in full detail in supplementary section S2. We assume a constant viscosity η , i.e., we assume $n = 1$, and we only consider the flow problem in the downstream direction (parallel to the margin), corresponding to very small distances from the origin, at which diffusion dominates heat transport, and deviations from the sliding velocity are small. In this case we can treat the velocity as the sum of a constant sliding velocity \bar{u}_b at the transition from frictional to free slip, and a correction $\tilde{u}(y, z)$, i.e., $u = \bar{u} + \tilde{u}$. The correction velocity then satisfies for $z > 0$

$$10 \quad \eta \nabla^2 \tilde{u} = 0 \tag{A1}$$

with ∇ the gradient operator in the transverse y - z -plane. \tilde{u} has to satisfy the boundary conditions

$$\eta \frac{\partial \tilde{u}}{\partial z} = \begin{cases} \tau_c & \text{at } z = 0, y < 0 \\ 0 & \text{at } z = 0, y > 0. \end{cases} \tag{A2}$$

In the supplementary section S2, we show that this leads to a leading order heat dissipation term $a = \eta |\nabla \tilde{u}|^2 \sim \tau_c^2 / (\pi^2 \eta) [\log(r/r_0)^2 + \vartheta^2]$ with r and ϑ polar coordinates (i.e., $y = r \cos \vartheta$ and $z = r \sin \vartheta$). Note that the heat production rate has only a logarithmic singularity in the present case, while there is a $1/r$ singularity for a no-slip to free slip transition, see (37).

With the heat dissipation a given, the temperature field close to the transition point satisfies at leading order

$$-k \nabla^2 T = \begin{cases} \frac{\tau_c^2}{\pi^2 \eta} [\log(r/r_0)^2 + \vartheta^2] & \text{for } z > 0 \\ 0 & \text{for } z < 0 \end{cases} \tag{A3}$$

with the boundary conditions

$$T(y, 0) = 0 \quad \text{for } z = 0, y > 0 \tag{A4a}$$

$$20 \quad -k \left[\frac{\partial T}{\partial z} \right]_{-}^{+} = \tau_c \bar{u}_b \quad \text{and} \quad [T(y, 0)]_{-}^{+} = 0 \quad \text{for } z = 0, y < 0. \tag{A4b}$$

In the ice ($0 < \vartheta < \pi$), the leading order solution of this is (see supplementary section S2)

$$T(r, \vartheta) = a_0 r^{1/2} \sin\left(\frac{\vartheta}{2}\right) + b_1 r \sin(\vartheta) + a_1 r^{3/2} \sin\left(\frac{3\vartheta}{2}\right) + b_2 r^2 \sin(2\vartheta) - \frac{\tau_c \bar{u}_b}{k} r \sin(\vartheta) + O\left(\frac{\tau_c^2}{4\pi^2 k \eta} r^2 \log(r)^2\right) \tag{A5a}$$

and in the bed ($\pi < \vartheta < 2\pi$), the leading order solution is

$$T(r, \vartheta) = a_0 r^{1/2} \sin\left(\frac{\vartheta}{2}\right) + b_1 r \sin(\vartheta) + a_1 r^{3/2} \sin\left(\frac{3\vartheta}{2}\right) + b_2 r^2 \sin(2\vartheta). \tag{A5b}$$

The term $(\tau_c \bar{u}_b/k)r \sin(\vartheta)$ results from the interfacial heating along the ice–bed contact where subtemperate slip is possible. The $O(\tau_c^2/(4\pi^2 k\eta)r^2 \log(r)^2)$ -term describes the contribution from the englacial heating, which is small in comparison to the contribution from subtemperate slip. Note that this is consistent with our results in section 4.4.2, where we found that the leading order heat equation (47) in the conductive boundary layer does not feature the englacial heat dissipation term, either.

- 5 If $a_0 \neq 0$, temperatures below the melting point for $y < 0$, $z = 0$ (on $\vartheta = \pi$) require $a_0 < 0$. However, this leads to a singular heat flux on the warm side ($z = 0$, $y > 0$)

$$-k \left. \frac{\partial T}{\partial z} \right|_{-}^{+} = -k \frac{1}{r} \left. \frac{\partial T}{\partial \vartheta} \right|_{\vartheta=2\pi}^{\vartheta=0} = -k \frac{a_0}{r^{1/2}} + O(1) \quad (\text{A6})$$

corresponding to a singular rate of freezing there. If we assume, as we do here, that a singular rate of freezing is not viable for a widening ice stream (see also Schoof, 2012), we must have $a_0 = 0$. With this choice, the temperature field at leading order is

- 10 determined by the $O(r)$ terms:

$$T(z) \sim b_1 z - \begin{cases} \frac{\tau_c \bar{u}_b}{k} z & \text{for } z > 0 \\ 0 & \text{for } z < 0. \end{cases} \quad (\text{A7})$$

- Consequently, at leading order a finite net negative heat flux out of the bed $-k[\partial T/\partial z]_{-}^{+} = \tau_c \bar{u}_b$ is possible for $y > 0$, corresponding to finite (i.e., non-singular) freezing. This heat flux is independent of the across-stream coordinate y and continuous along y , in contrast to the case without subtemperate slip (see Schoof, 2012). Physically, this means that the existence of
- 15 subtemperate slip on the cold side requires the removal of heat there and at the temperate side simultaneously. At the temperate side, where no heat is dissipated along the bed, this heat must be supplied in different form, most likely as latent heat transported by subglacial drainage.

Competing interests. The authors declare that they have no conflict of interest.

- Acknowledgements.* MH was supported by a Four Year Fellowship at the University of British Columbia, NSERC grant 357193-13, and
- 20 the Princeton AOS Postdoctoral and Visiting Scientist Program. CS acknowledges NSERC grants 357193-13 and 446042-13. Numerical calculations performed on WestGrid facilities were supported by Compute Canada. [We thank the editor Eric Larour and two anonymous referees for their thorough reviews which have helped to improve the manuscript.](#)

References

- Aschwanden, A., Bueler, E., Khroulev, C., and Blatter, H.: An enthalpy formulation for glaciers and ice sheets, *Journal of Glaciology*, 58, 441–457, doi:10.3189/2012JoG11J088, 2012.
- Bamber, J. L., Vaughan, D. G., and Joughin, I.: Widespread Complex Flow in the Interior of the Antarctic Ice Sheet, *Science*, 287, 1248–1250, doi:10.1126/science.287.5456.1248, 2000.
- 5 Bentley, C. R., Lord, N., and Liu, C.: Radar reflections reveal a wet bed beneath stagnant Ice Stream C and a frozen bed beneath ridge BC, West Antarctica, *Journal of Glaciology*, 44, 149–156, 1998.
- Bindschadler, R., Chen, X., and Vornberger, P.: The onset area of Ice Stream D, West Antarctica, *Journal of Glaciology*, 46, 95–101, doi:10.3189/172756500781833377, 2000.
- 10 Blatter, H.: Velocity and stress fields in grounded glaciers: a simple algorithm for including deviatoric stress gradients, *Journal of Glaciology*, 41, 333–344, 1995.
- Brinkerhoff, D. J. and Johnson, J. V.: Dynamics of thermally induced ice streams simulated with a higher-order flow model, *Journal of Geophysical Research*, 120, 1743–1770, doi:10.1002/2015JF003499, <http://dx.doi.org/10.1002/2015JF003499>, 2015JF003499, 2015.
- Catania, G., Hulbe, C., Conway, H., Scambos, T. A., and Raymond, C. F.: Variability in the mass flux of the Ross ice streams, West Antarctica, over the last millennium, *Journal of Glaciology*, 58, 741–752, doi:10.3189/2012JoG11J219, 2012.
- 15 Catania, G. A., Conway, H. B., Gades, A. M., Raymond, C. F., and Engelhardt, H.: Bed reflectivity beneath inactive ice streams in West Antarctica, *Annals of Glaciology*, 36, 287–291, 2003.
- Catania, G. A., Scambos, T. A., Conway, H., and Raymond, C. F.: Sequential stagnation of Kamb Ice Stream, West Antarctica, *Geophysical Research Letters*, 33, L14502, doi:10.1029/2006GL026430, 2006.
- 20 Conway, H., Catania, G., Raymond, C. F., Gades, A. M., Scambos, T. A., and Engelhardt, H.: Switch of flow direction in an Antarctic ice stream, *Nature*, 419, 465–467, doi:10.1038/nature01081, 2002.
- Cuffey, K. M., Conway, H., Hallet, B., Gades, A. M., and Raymond, C. F.: Interfacial water in polar glaciers and glacier sliding at -17°C , *Geophysical Research Letters*, 26, 751–754, 1999.
- Durand, G., Gagliardini, O., Zwinger, T., Le Meur, E., and Hindmarsh, R. C. A.: Full Stokes modeling of marine ice sheets: influence of the grid size, *Annals of Glaciology*, 50, 109–114, 2009.
- 25 Echelmeyer, K. and Zhongxiang, W.: Direct observation of basal sliding and deformation of basal drift at sub-freezing temperatures, *Journal of Glaciology*, 33, 83–98, 1987.
- Echelmeyer, K. A. and Harrison, W. D.: Ongoing margin migration of Ice Stream B, Antarctica, *Journal of Glaciology*, 45, 361–369, 1999.
- Elsworth, C. W. and Suckale, J.: Rapid ice flow rearrangement induced by subglacial drainage in West Antarctica, *Geophysical Research Letters*, 43, 11–697, 2016.
- 30 Fahnestock, M. A., Scambos, T. A., Bindschadler, R. A., and Kvaran, G.: A millennium of variable ice flow recorded by the Ross Ice Shelf, Antarctica, *Journal of Glaciology*, 46, 652–664, doi:10.3189/172756500781832693, 2000.
- Fowler, A. C.: Sub-temperate basal sliding, *Journal of Glaciology*, 32, 3–5, 1986.
- Fowler, A. C.: The motion of ice stream margins, *Journal of Fluid Mechanics*, 714, 1–4, 2013.
- 35 Fowler, A. C. and Larson, D. A.: On the Flow of Polythermal Glaciers. I. Model and Preliminary Analysis, *Proceedings of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 363, 217–242, <http://www.jstor.org/stable/79748>, 1978.

- Gagliardini, O., Zwinger, T., Gillet-Chaulet, F., Durand, G., Favier, L., Fleurian, B. d., Greve, R., Malinen, M., Martín, C., Råback, P., et al.: Capabilities and performance of Elmer/Ice, a new generation ice-sheet model, *Geoscientific Model Development*, 6, 1299–1318, doi:10.5194/gmd-6-1299-2013, 2013.
- Gilpin, R. R.: A model of the “liquid-like” layer between ice and a substrate with applications to wire regelation and particle migration, *Journal of colloid and interface science*, 68, 235–251, 1979.
- Gladstone, R. M., Payne, A. J., and Cornford, S. L.: Resolution requirements for grounding-line modelling: sensitivity to basal drag and ice-shelf buttressing, *Annals of Glaciology*, 53, 97–105, 2012.
- Hamilton, G. S., Whillans, I. M., and Morgan, P. J.: First point measurements of ice-sheet thickness change in Antarctica, *Annals of Glaciology*, 27, 125–129, 1998.
- Harrison, W. D., Echelmeyer, K. A., and Larsen, C. F.: Measurement of temperature in a margin of Ice Stream B, Antarctica: implications for margin migration and lateral drag, *Journal of Glaciology*, 44, 615–624, 1998.
- Haseloff, M.: Modelling the migration of ice stream margins, Ph.D. thesis, The University of British Columbia, Retrieved from <http://hdl.handle.net/2429/54268>, 2015.
- Haseloff, M., Schoof, C., and Gagliardini, O.: A boundary layer model for ice stream margins, *Journal of Fluid Mechanics*, 781, 353–387, doi:10.1017/jfm.2015.503, 2015.
- Hewitt, I. J. and Schoof, C.: Models for polythermal ice sheets and glaciers, *The Cryosphere*, 11, 541–551, doi:10.5194/tc-11-541-2017, <https://www.the-cryosphere.net/11/541/2017/>, 2017.
- Hindmarsh, R. C. A.: Consistent generation of ice-streams via thermo-viscous instabilities modulated by membrane stresses, *Geophysical Research Letters*, 36, L06 502, doi:10.1029/2008GL036 877, 2009.
- Holdsworth, G.: Meserve Glacier Wright Valley Antarctica: Part 1. Basal Processes, <http://hdl.handle.net/1811/47277>, 1974.
- Hulbe, C. and Fahnestock, M.: Century-scale discharge stagnation and reactivation of the Ross ice streams, West Antarctica, *Journal of Geophysical Research*, 112, F03S27, doi:10.1029/2006JF000603, 2007.
- Hutter, K.: *Theoretical Glaciology*, D. Reidel Publishing Company/ Terra Scientific Publishing Company, 1983.
- Jacobson, H. P. and Raymond, C. F.: Thermal effects on the location of ice stream margins, *Journal of Geophysical Research*, 103, 12 111–12 122, doi:10.1029/98JB00574, 1998.
- Joughin, I., Tulaczyk, S., Bindschadler, R., and Price, S. F.: Changes in west Antarctic ice stream velocities: Observation and analysis, *Journal of Geophysical Research*, 107, 2289, doi:10.1029/2001JB001 029, 2002.
- Kyrke-Smith, T. M., Katz, R. F., and Fowler, A. C.: Subglacial hydrology and the formation of ice streams, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Science*, 470, 20130 494, doi:10.1098/rspa.2013.0494, 2014.
- MacAyeal, D. R.: Large-scale ice flow over a viscous basal sediment - Theory and application to ice stream B, Antarctica, *Journal of Geophysical Research*, 94, 4071–4087, doi:10.1029/JB094iB04p04071, 1989.
- Morland, L. W. and Johnson, I. R.: Steady motion of ice sheets, *Journal of Glaciology*, 25, 229–246, 1980.
- Muszynski, I. and Birchfield, G. E.: A coupled marine ice-stream-ice-shelf model, *Journal of Glaciology*, 33, 3–15, 1987.
- Paterson, W. S. B.: *The Physics of Glaciers*, Elsevier, Oxford, 1994.
- Pattyn, F.: A new three-dimensional higher-order thermomechanical ice sheet model: Basic sensitivity, ice stream development, and ice flow across subglacial lakes, *Journal of Geophysical Research*, 108, 2382, doi:10.1029/2002JB002 329, 2003.
- Pattyn, F., Schoof, C., Perichon, L., Hindmarsh, R. C. A., Bueler, E., de Fleurian, B., Durand, G., Gagliardini, O., Gladstone, R., Goldberg, D., Gudmundsson, G. H., Huybrechts, P., Lee, V., Nick, F. M., Payne, A. J., Pollard, D., Rybak, O., Saito, F., and Vieli, A.: Results of

- the Marine Ice Sheet Model Intercomparison Project, MISIMIP, *The Cryosphere*, 6, 573–588, doi:10.5194/tc-6-573-2012, <http://www.the-cryosphere.net/6/573/2012/>, 2012.
- Perol, T. and Rice, J. R.: Shear heating and weakening of the margins of West Antarctic ice streams, *Geophysical Research Letters*, 42, 3406–3413, doi:10.1002/2015GL063638, 2015.
- 5 Perol, T., Rice, J. R., Platt, J. D., and Suckale, J.: Subglacial hydrology and ice stream margin locations, *Journal of Geophysical Research*, 120, 1–17, doi:10.1002/2015JF003542, 2015.
- Peters, L. E., Anandakrishnan, S., Alley, R. B., Winberry, J. P., Voigt, D. E., Smith, A. M., and Morse, D. L.: Subglacial sediments as a control on the onset and location of two Siple Coast ice streams, West Antarctica, *Journal of Geophysical Research*, 111, B01 302, doi:10.1029/2005JB003 766, doi:10.1029/2005JB003766, 2006.
- 10 Platt, J. D., Perol, T., Suckale, J., and Rice, J. R.: Determining conditions that allow a shear margin to coincide with a R othlisberger channel, *Journal of Geophysical Research*, 121, 1273–1294, 2016.
- Pralong, A. and Funk, M.: Dynamic damage model of crevasse opening and application to glacier calving, *Journal of Geophysical Research*, 110, B01 309, doi:10.1029/2004JB003104, 2005.
- Raymond, C.: Shear margins in glaciers and ice sheets, *Journal of Glaciology*, 42, 90–102, 1996.
- 15 Raymond, C. F.: Energy balance of ice streams, *Journal of Glaciology*, 46, 665–674, doi:10.3189/172756500781832701, 2000.
- Retzlaff, R. and Bentley, C. R.: Timing of stagnation of Ice Stream C, West Antarctica, from short-pulse radar studies of buried surface crevasses, *Journal of Glaciology*, 39, 553–561, 1993.
- Rice, J.: Stresses due to a sharp notch in a work-hardening elastic-plastic material loaded by longitudinal shear, *Journal of Applied Mechanics*, 34, 287–298, 1967.
- 20 Schoof, C.: On the mechanics of ice-stream shear margins, *Journal of Glaciology*, 50, 208–218, doi:10.3189/172756504781830024, 2004.
- Schoof, C.: Variational methods for glacier flow over plastic till, *Journal of Fluid Mechanics*, 555, 299–320, doi:10.1017/S0022112006009104, 2006.
- Schoof, C.: Coulomb friction and other sliding laws in a higher-order glacier flow model, *Mathematical Models and Methods in Applied Sciences*, 20, 157–189, 2010.
- 25 Schoof, C.: Thermally driven migration of ice-stream shear margins, *Journal of Fluid Mechanics*, 712, 552–578, 2012.
- Schoof, C. and Hewitt, I.: A model for polythermal ice incorporating gravity-driven moisture transport, *Journal of Fluid Mechanics*, 797, 504–535, doi:10.1017/jfm.2016.251, 2016.
- Shreve, R. L.: Glacier sliding at subfreezing temperatures, *Journal of Glaciology*, 30, 341–347, 1984.
- Stearns, L. A., Jezek, K. C., and Van der Veen, C.: Decadal-scale variations in ice flow along Whillans Ice Stream and its tributaries, West Antarctica, *Journal of Glaciology*, 51, 147–157, 2005.
- 30 Stephenson, S. N. and Bindshadler, R. A.: Observed velocity fluctuations on a major Antarctic ice stream, *Nature*, 334, 695–697, doi:10.1038/334695a0, 1988.
- Suckale, J., Platt, J. D., Perol, T., and Rice, J. R.: Deformation-induced melting in the margins of the West Antarctic ice streams, *Journal of Geophysical Research*, 119, 1004–1025, doi:10.1002/2013JF003008, 2014.
- 35 Wolovick, M. J., Creyts, T. T., Buck, W. R., and Bell, R. E.: Traveling slippery patches produce thickness-scale folds in ice sheets, *Geophysical Research Letters*, 41, 8895–8901, 2014.