

Dear Editor:

We are grateful for the constructive comments by the two reviewers which helped us improve the scientific quality and clarity of this manuscript. We
5 agree to most comments and suggestions by the reviewers, which are provided below.

Thank you for receiving our revised manuscript and we look forward to hearing from you.

10 Sincerely,
Juri Palmtag

Anonymous Referee #1

15

Comment: P 4. L 22. “time or”. Isn’t time part of the logistical problem?

Response: We agree that it is part of logistical problems and remove “time or”
from the text.

20 Comment: P 6. L 7-15. Materials in 2 glacial moraines (lateral and end) deeper than 100 was considered as “till”. Since soils sampled from 0-100 cm in both landform formed in moraine, the parent material of these soils is moraine. Then, why call the portion below 100 cm till?

Response: We refer to the deeper deposits as till from a sedimentological
25 perspective and not as a landform, based on the analyzed coring material which was unsorted glacial sediment

Comment: P 6. L. 31-32. C stores reported for 0-300 cm in small lakes. Not clear if the whole 300 cm includes both water and sediment?

30 Response: This is a good point so we added additional clarification in the text “from sediment surface”.

Comment: P 7. L 37. “importance of”. Would it be better say “effects of” or “important role of”?

35 Response: We changed as suggested to “important role of”.

Comment: P 8. L 25-27. Consider change “surface wetness” to “drainage” and permafrost table” to “depth of permafrost” or “active layer thickness”.

Response: As suggested we have exchanged surface wetness” to “drainage” and permafrost table” to “active layer thickness” as suggested.

5

Anonymous Referee #2

Comment: Page 1 L1 (title): Very long title, but the nitrogen is missing. What about “Deep storage of organic matter in Zackenberg, Greenland”. The
5 methodical approach (“geomorphological landform approach”) is not needed here.

Response: We agree that the title was very long and used partly used the suggestion by the reviewer. But since the actual aim of this paper is the used approach, we think that landform partitioning should be still part of the title.

10

Comment: L13: Please add an introductory sentence before stating the paper aims. General comment: are all the included samples soils in the sense of soil science? If yes, SOC is an appropriate abbreviation, otherwise please change to OC. Please add to the abstract a short discussion sentence including “what is
15 deeper than 3 m” (you did this in the paper text already for fan/delta) and “how representative” could this study area be for Greenland/or other Arctic areas (missing right now).

Response: As asked we added an introductory sentence: “Soils in the northern high latitudes are a key component in the global carbon cycle, with potential
20 feedbacks on climate.” We also changed for the deeper deposits the abbreviation to OC and added which type of sediment it is “alluvial and deltaic deposits”. We also added in the discussion section rough estimations on SOC storage in deposits below 3m.

25 Comment: L30: Please add a reference here

Response: Thank you for your comment. We have added additional reference of Schuur et al., 2015.

Comment: L31: Please concretize “most regions”. Permafrost regions, Arctic
30 regions? Despite IPCC, please add a primary reference here.

Response: Thank you again for your comment. Primary reference added and the regions are now concretized with: “northern circumpolar regions”.

Comment: Page 2 L4: As you know first assessment was not Tarnocai et al 2009, there have been a lot more before, like Post et al 1982, Tarnocai 2003, than Zimov 2006.

Response: We agree and thankful for the comment. This part is now corrected
5 with the proper references regarding the first assessment: “In 2009, Tarnocai et al. linked circumpolar SOC data (e.g. Kuhry et al., 2002; Zimov et al., 2006; Tarnocai et al, 2007; Ping et al, 2008) and presented a new total estimate of 1674 Petagram C (PgC) stored in soils and deep deposits of the northern circumpolar permafrost region.”.

10

Comment: L16: add a dot after 2016)...

Response: Typo corrected and dot added.

Comment: L29 and following: I like the concise presentation of the specific
15 aims. Please be consistent and use this order and number for the conclusion sections

Response: The conclusion section is now restructured and the aims are following the same order.

20 Comment: L36: please delete the c. or introduce this abbreviation for the coordinates.

Response: Agree that c is unnecessary and gladly removed from the text.

Comment: L38: Please add the rain/snow percentages of the precipitation (if
25 available)

Response: Asked information on rain/snow added to the site description.

Comment: Page 3 L15: Please add the GPS uncertainty here

Response: Approximate GPS uncertainty added.

30

Comment: L19: Not the motor head, but the core barrel system is of relevance here. Please add this information here.

Response: As asked we added the additional information on the core barrel in the methods section.

Comment: L32: How many samples were measured for TOC/TN. TOC and TN
5 in one run and same device? Or did you decarbonise the samples before to get rid of the TIC? Did you calibrate your 950 TIC measurements as well?

Response: Added and explained in more detail the questions raised by the reviewer #2 on the number of samples, device and sample preparation.

10 Comment: Page 4 Line 9: Please add the units to the formula, this would explain the percentage /100 and g to kg as well as the cm³ to m² conversion factor

Response: The units are explained just above the equation which is a common way to do it. It seems very unnecessary to repeat it again.

15

Comment: L16: Please explain the mentioned “data limitations”

Response: The text part is now reworked explaining the data limitation for the lack of deeper TN values.

20 Comment: L22: please delete “time or”

Response: Same comment as from reviewer 1, so we removed time from the text.

Comment: Page 5 L3: Is your data Gaussian distributed? Did you test this (e.g.
25 Shapiro-Wilk test). Why not using more robust median and interquartile ranges? Please explain and justify your decisions here.

Response: Thank you for your comment. Yes, in most cases there are too few sites to do a proper test for the normal distribution. However, yes I did test this using Shapiro-Wilk for example in the alluvial fans based on 15 sites.

30 Regarding the usage of standard deviation: We follow a standard protocol used by our group and others. For that reason and for comparability between sites we use SD.

Comment: L16: Same for CI, which requires Gaussian distributed data

Response: See response above, where tested data is normal distributed.

General comments results section: I understand you n is the number of sites, but could you add a n for the number of measurements as well? Is the percentage a result of Cable 2017 or this study? I recommend transforming the major information of Table 2 into a boxplot figure (which is allowed for a n<5 for n (measurements, nit sites) and include table 2 into a supplementary chapter. Is the data available (embargoed, e.g. PANGAEA). Then you can add a doi of your data.

Response: As asked by the reviewer, we added the number of measurements in Table 2. Question regarding the percentage landform cover. As stated in the text, we amalgamated small classes from Cable et al., where we didn't have any sampling pedons into larger landform classes. However we use the same aerial extend of the study area. Table 1 shows the geomorphological landforms and their proportion of total surface area in Cable et al., into the amalgamated classes. As stated in a former response, we would like to keep the table with all the data as it is which enables a direct comparison with other sites dealing with SOC storage. Also, the data is not yet available in e.g. PANGAEA but is planned in the future.

20

Comment: Page 7 L7: First estimated for Zackenberg or Greenland or similar landscapes?

Response: Additional information added to clarify this point.

Comment: L24: please delete c., which is not coordinates for which this abbreviation was used before

Response: c which was used for circa now removed from several places.

General comments discussion: Please include and discuss a number of how much is perennially frozen/seasonally unfrozen of you carbon and nitrogen stocks. Please include comparisons of your and other comparable case studies, which you all know because of author overlap (e.g. Fuchs et al. Biogeosciences, Discussions, Hugelius et al. 2009, etc.) To my opinion

references (4 actually, as Palmtag 2015 is mentioned twice) for a discussion are not enough. I do not care about the number of references, but this shows that the authors should discuss their findings a bit bigger context. Frozen/unfrozen percentages, what are the consequences for the carbon pool → modelling...any
5 back on the envelope calculation of the <3m pool of the fans etc. Discuss the satellite product comparisons (Bartsch and other ESA DUE or GlobPermafrost-related publications). Rough discussion of carbon qualities (using C/N) and compare this to literature concepts (for C/N Schaedel et al 2014) Maybe an additional “discussion figure” (right now just 1 intro and 1 results figure)
10 would be helpful.

Response: Thank you for these comments. Several more references were now added and their findings discussed and compared to our results. We also included comments on the perennially frozen proportion. However, there are too few sites at all available with comparable mountainous permafrost
15 environment and to my knowledge none of those which deals with carbon stocks deeper than 1m of depth. We also included the publication from Bartsch et al. using SOC estimated based on synthetic aperture radar. But since the soil penetration is only a few cm, it has the similar problem as LCC and cannot capture the long-term depositional history. And we believe that a discussion of
20 carbon qualities is beyond the scope of this paper.

Comment: Page 8 L18: Keep using introduced abbreviations (here CI) and do not introduce them twice.

Response: Mistake corrected and the secondary introduction of abbreviation
25 removed.

Comment: Page 9 L6: Making an rough estimation by stating the assumption of a <3m pool could be a nice first guess for future work

Response: We added a rough estimation for SOC storage in deltaic deposits
30 below 3m.

General comments for the conclusions: Please repeat the paper's aims here and answer this in the same order like in introduction. Please include your nitrogen calculations here as well.

5 Response: The comment implemented and the aims reordered following the introductions order. Also, the nitrogen was added in this section.

Comment: L9: "new additional" sounds strange

Response: The "additional" is now removed from the text.

10 Comment: Figures and Table: Figure 1: hard to read the site labels. Maybe a hillshade and transparent colours could improve the geomorphological understanding for the reader. According to figure 2: use A and B instead of "top right"

Response: Both comments implemented.

15

Comment: Figure 2: A is redundant with Table 1, right? An option could be deleting a and put the spotlight on B-C.

20 Response: Figure 2 A is redundant with Table 1 but we would still like to keep is as it, since it illustrates graphically the proportional contribution of each landform in comparison to the SOC storage for each landform.

Comment: Table 2: please add n measurements to the table. Moreover switching from CI (in the manuscript text) to SD could be puzzling for the readers.

25 Response: As suggested we added in the table 2 the number of samples measured for each landform. Regarding the use of SD and CI: In the manuscript text and Table 2 it is clearly stated if we used SD (only for the landform means) or CI which was used for the whole Zackenberg area. The decision to use CI is because it account for the relative spatial coverage,
30 storage variability and degree of replication of each upscaling class and therefore much more confident.

Improved Landform landscape partitioning and estimates of deep storage of soil organic carbon matter in the Zackenberg, Greenland area (NE Greenland) using a geomorphological landform approach

Juri Palmtag¹, Stefanie Cable^{2, 3}, Hanne H. Christiansen^{2, 3}, Gustaf Hugelius¹ and Peter Kuhry¹

¹ Department of Physical Geography, Stockholm University, 106 91 Stockholm, Sweden

² Center for Permafrost (CENPERM), Department of Geosciences and Natural Resource Management, University of Copenhagen, 1350 Copenhagen, Denmark

³ Arctic Geology Department, The University Centre in Svalbard (UNIS), 9170 Longyearbyen, Norway

Correspondence to: Juri Palmtag (juri.palmtag@natgeo.su.se)

Abstract. Soils in the northern high latitudes are a key component in the global carbon cycle, with potential feedbacks on climate. This study aims to improve the previous soil organic carbon (SOC) and total nitrogen (TN) storage estimates for the Zackenberg area (NE Greenland) that were based on a land cover classification (LCC) approach, by using geomorphological upscaling. In addition, novel SOC-organic carbon (OC) estimates for deeper alluvial and deltaic deposits (down to 300 cm depth) are presented. We hypothesize that landforms will better represent the long-term slope and depositional processes that result in deep SOC burial in this type of mountain permafrost environments. The updated mean SOC storage for the 0–100 cm soil depth is 4.8 kg C m⁻², which is 42% lower than the previous estimate of 8.3 kg C m⁻² based on land cover upscaling. Similarly, the mean soil TN storage in the 0–100 cm depth decreased with 44% from 0.50 kg (± 0.1 CI) to 0.28 (± 0.1 CI) kg TN m⁻². We ascribe the differences to a previous areal overestimate of SOC and TN-rich vegetated land cover classes. The landform-based approach more correctly constrains the depositional areas in alluvial fans and deltas with high SOC and TN storage. These are also areas of deep carbon storage with an additional 2.4 kg C m⁻² in the 100–300 cm depth interval. This research emphasizes the need to consider geomorphology when assessing SOC pools in mountain permafrost landscapes.

1 Introduction

Permafrost soils in the northern circumpolar region are sensitive to climate change (Schuur et al., 2015). In addition, they store large amounts of soil organic carbon (SOC) that has accumulated under low ground temperatures over decadal to millennial timescales. In most northern circumpolar regions, permafrost temperatures have increased since the 1980s (Romanovsky et al., 2010) and increased global mean surface temperatures are projected to decrease the near-surface permafrost extent by 37% to 81% (with RCP2.6 to RCP8.5,

respectively) by the end of the 21st Century (Slater & Lawrence, 2013; IPCC 2013). The observed permafrost degradation could intensify microbial activity and increase decomposition of organic matter formerly stored in permafrost, releasing more greenhouse gases into the atmosphere and providing a positive feedback on global warming (Schuur et al., 2015). Over a decade ago, permafrost was identified as a major vulnerable terrestrial carbon pool (Gruber et al., 2004). In 2009, Tarnocai et al. linked circumpolar SOC data (e.g. Kuhry et al., 2002; Zimov et al., 2006; Tarnocai et al., 2007; Ping et al., 2008) and presented a new total estimate of 1674 Petagram C (PgC) stored in soils and deep deposits of the northern circumpolar permafrost region. The first assessment of the magnitude of the Northern Permafrost Region SOC pool was made by Tarnocai et al. (2009) who estimated it to be as large as ca. 1700 Petagram C (PgC). Since then, availability of more data has constrained this estimate to ca. 1300±200 PgC (Hugelius et al., 2014). However, substantial uncertainties and data-gaps remain, particularly for high-Arctic and mountainous landscapes. This limited knowledge on the spatial distribution of SOC in permafrost landscapes remains a strong constraint on the ability to predict the vulnerability of the permafrost SOC pools from local to landscape to pan-Arctic scales.

Most landscape to regional-scale estimates of permafrost SOC stocks have used thematic maps to stratify and scale point observations to full spatial coverage (Hugelius, 2012). There are numerous examples of studies applying land cover classifications (LCC) for SOC upscaling in permafrost landscapes (e.g. Kuhry et al., 2002; Hugelius and Kuhry, 2009; Palmtag et al., 2015, 2016). In Palmtag et al. (2015), a LCC was applied for upscaling SOC for two lowland sites in NE Siberia and a mountainous site, Zackenberg Valley, in NE-Greenland. However, limitations have been pointed out especially for the mountainous Zackenberg Valley site where the geomorphology and cryostratigraphy are highly heterogeneous (Cable et al., 2017; 2018). In this area, relatively shallow deposits occur on steeper slopes, while the foothills accumulate massive deposits over millennial time-scales, involving carbon burial with syngenetic permafrost aggradation (Palmtag et al., 2015; Cable et al., 2018). Thus, the application of LCC-scaling for SOC upscaling in high relief landscapes can involve larger uncertainties, particularly when estimating deeper SOC of colluvial and alluvial deposits because the long-term depositional history, controlling SOC burial, cannot be captured by the current vegetation cover that is primarily reflected in LCC. Combining a geomorphology-based landscape classification (GLC), now available for the Zackenberg Valley (Cable et al., 2017; 2018), with high quality data from detailed field studies may, therefore, improve SOC upscaling in high relief permafrost landscapes.

The overall aim of this study is to improve the SOC and total nitrogen (TN) storage estimates for the Zackenberg area (NE Greenland). Specific objectives are (1) to use largely the same dataset as in Palmtag et al. (2015) to upscale SOC and TN at 0–100 cm depth to landscape scale, based on geomorphological mapping, (2) to compare the results with the previous LCC upscaling approach, (3) to present the first SOC estimates for deposits deeper than 1 m based

on newly collected deep pedons, and (4) to evaluate the importance of geomorphology for assessing landscape level SOC storage.

2 Study area

The Zackenberg valley (e. 74°28'N, 20°34'W) is a mountainous high-arctic tundra area with a mean annual temperature of -9.2°C located within the continuous permafrost zone of NE Greenland (Fig. 1). The mean annual precipitation is 261 mm of which 10% falls as rain during the summer months from June to September (Hansen et al., 2008). The study area extends from sea level at the shores of the Young Sound up to 1372 m at the top of Zackenberg Mountain. A large fault system divides the weathering resistant Caledonian gneiss/granite bedrock in the West from the Cretaceous-Tertiary sedimentary rocks in the east (Escher and Watt, 1976). According to Bennike et al. (2008), the valley deglaciated prior to 11 300 calendar years Before Present (cal yr BP) (~~calendar years Before Present~~). The parent material in the low-lying central valley is dominated mostly by glacial, periglacial, alluvial, fluvial and deltaic deposits, while on the slopes either boulder-fields or colluvial sediment predominate (Christiansen et al., 2008; Cable et al., 2018). A weakly developed Typic Psammenturbel is the prevailing soil type of the central valley, with Gelorthents on the slopes (terminology following Soil Survey Staff, 2014). Small areas were occupied by peaty soils, mainly Histoturbels.

3 Methods

3.1 Soil sampling

Field work was conducted during late summer in 2009, 2012 and 2013. In 2009 and 2012, 38 sampling sites were selected using a semi-random, stratified transect sampling approach with predefined equidistant pedon intervals of 100 to 500 m using a handheld GPS device (accuracy of approximately 10 meters). Mineral soil samples were collected using a steel pipe, manually hammered into the soil. A more detailed sampling procedure is described in Palmtag et al. (2015). In 2013, additional deeper core pedons (down to 455 cm) were sampled by drilling in alluvial fans (Cable et al., 20172018), using a handheld motorized rotational Earth Auger with a 50cm core barrel with outside diameter of 52mm and a cutting edge (STIHL BT 121). Out of the total of 48 sites used in this study (Fig. 1), eight sites were sampled to depths of more than 200 cm, 19 sites to between 100 and 200 cm depth, and the remaining sites to less than 1m depth, primarily due to shallow mineral soils overlying the bedrock. There were 648 samples collected in total throughout the three field seasons consisting of on average 10 cm long increments of organic layer, active layer and permafrost samples.

3.2 Soil chemical analyses and SOC/TN calculations

Each sample had a known field volume and was oven-dried, weighed and sieved to determine the dry bulk density (DBD, g cm⁻³) and the amount of coarse fragments (CF, >2mm, %).

Subsequently, each sample was individually homogenized and burned to obtain loss on ignition (LOI; Heiri et al., 2001) at 550°C and, about every second sample, at 950°C to determine, respectively, its organic matter and inorganic carbonate content through weight loss (for details, see Palmtag et al., 2015).

5 | To determine total organic carbon and nitrogen (TOC/TN, %), ca. 70% (n=284) of the samples (all samples from fieldwork in 2009) across all sites and horizons were analysed using an EA 1110 Elemental Analyzer that measures %C and %N in the same runs (CE Instruments, Italy). To calculate the soil organic carbon content for the remaining samples, we used a fifth order polynomial regression ($R^2 = 0.97$) between LOI at 550 °C and TOC values from the elemental
10 | analyser. This high order polynomial regression was necessary to correctly represent %C at very low LOI 550 °C. Results for the inorganic carbon content were very low. The latter measurements were based on 297 samples with an average LOI 950 °C weight loss of only 0.994%. Therefore, samples were not decarbonized and we did not calculate separately the contribution of inorganic carbonate in the samples.

15 | The SOC and TN storage (kg m^{-2}) was calculated for each collected sample using the available data on DBD (g cm^{-3}), %C or %TN, 1-CF (the remaining proportion of the sample after excluding the coarse mineral fractions (>2 mm)), thickness T (cm) of the sample, multiplied by 10 for unit conversion (see equation 1 and 2):

$$20 \quad SOC = (DBD * \%C * (1 - CF) * T) * 10 \quad (1)$$

$$TN = (DBD * \%TN * (1 - CF) * T) * 10 \quad (2)$$

Then, the total SOC and TN (kg m^{-2}) storage was calculated by summing all the individual samples from the same profile to the different reference depths of 0–30 cm, 0–100 cm, 0–200
25 | cm, etc. In cases when a sample was missing, the gap was interpolated from samples above and below by taking into account field notes on texture, ice content, coarse fraction and any presence of buried C-enriched soil layers. Because of ~~Due~~ too few TN values available for the deeper deposits, data limitations, the TN storage was calculated for the 0–100 cm depth interval only.

30 | In total, we used 48 sampling sites to estimate SOC storage. If bedrock was hit at any point (n=8 within 0–100 cm depth; n=10 within 100–200 cm depth; n=5 within 200–300 cm depth), we used a SOC content of 0 kg C m^{-2} for the remaining bedrock part down to 300 cm depth. In the 25 remaining sites within the depth range of 0–100 cm, 100–200 cm and 200–300 cm,
35 | where bedrock was not reached but sampling was stopped because of ~~time or~~ logistical constraints, the SOC content was extrapolated to the full 100, 200 or 300 cm depth intervals by taking into account the location, topography, geomorphologic landform, sediment type and information from other similar sites.

3.3 Upscaling procedure

The upscaling from the field measurements to landscape scale was performed in ArcGIS 10.2 (ESRI, Redlands, CA, USA) by multiplying arithmetic means of SOC from all sites belonging to the same landform class with the extent of that same class in the digital geomorphology map at a 1:10 000 scale from Cable et al. (2017~~8~~). The geomorphological map is based on geomorphological mapping using ortho-rectified panchromatic aerial images of 0.2 m resolution, and field validation. The 48 SOC sampling sites covered most, but not all of the originally recognized 28 landforms, however, 12 of these occupied only negligible areas of <3% of the total study area (Table 1). To achieve full coverage across the study area, the mapped landforms were merged into larger geomorphological classes based on topographic position and overall geomorphological characteristics. The adjusted map, consisting of 10 geomorphological classes, was then used for SOC upscaling. The map has a terrestrial coverage of 111 km² with an additional 17 km² of sea area (Young Sound/Tyrolerfjord), which is not included in the upscaling, but visible on the map (Fig. 1). The extent of this geomorphological map nearly completely overlaps with that of the LCC map used by Palmtag et al. (2015).

The arithmetic mean SOC storage with standard deviations (SD) for different depth intervals in the active layer and permafrost (and TN 0–100 cm storage) were calculated for each landform based on all study sites in each landform. One landform (bedrock) consists of only one sampling site but, since SOC in rock walls is considered negligible, any assumed small within-class variability barely affects the SOC estimate at landscape scale. Subsequently, the landscape mean SOC storage for the whole study area was calculated from the mean values of each geomorphological class multiplied with the proportion of the area occupied by that class in the simplified geomorphological map.

3.4 Statistical analyses

To provide reasonable error estimates for landscape SOC, which vary naturally in the environment, a spatially weighted 95% confidence interval (CI) was calculated following Thompson (1992). This CI is calculated to account for the relative spatial coverage, storage variability and degree of replication of each upscaling class using equation 3:

30

$$CI = t * \sqrt{\sum \left((a_i^2 * SD_i^2) / n_i \right)} \quad (3)$$

Where ‘t’ is the upper $\alpha/2$ of a normal distribution (1.96); ‘a’ the percentage of the total area per class; ‘SD’ the standard deviation of the storage estimate per class; ‘n’ the number of replicates per class; and ‘i’ refers to the specific land form classes. The applied upscaling procedure assumes that the available sample is sufficiently replicated to accurately reflect the natural variability within a class (Hugelius, 2012; Palmtag et al., 2016). It is important to note

that the presented CI ranges do not account for any spatial errors in landform upscaling products. Error estimates for landscape SOC as well as the analyses of variances with means and SD were performed using the software MS Excel.

4 Results

5 The Zackenberg study area consists of several main geomorphological classes (Figs. 1 and 2). Exposed bedrock occupies about 8% of the study area, containing negligible SOC storage (Fig. 2 and Table 2). The most widespread landform class (30% of the study area) is ‘allochthonous weathered bedrock’ (n=2). This class is predominantly located at higher elevation on steep hillslopes and consist of coarse-grained colluvium deposits with very little soil and vegetation
10 development, deposited by either downslope creep and/or gravity depositional sorting. Only 7% of the surface of this class is actually vegetated, leaving 93% bare ground and exposed bedrock. This landform is rather active leading to relative shallow sediments with very low SOC content (Fig. 2 and Table 2).

The landform ‘solifluction sheets’ (n=4) covers 14% of the study area, is located on
15 intermediate hillslopes and consists mainly of fine-grained colluvium deposits, in general loose unconsolidated weathered sediments deposited by slow downslope movement of water-saturated sediment due to recurrent freezing and thawing of the ground and driven by gravity. Only 18% of the area is vegetated, with the remaining 82% occupied by boulders. SOC content is low (Fig. 2 and Table 2).

20 The class ‘lateral and end moraines’ (n=4) occupies 18% of the area. More than 60% of its surface area is occupied by boulders. This landform is inactive (not eroding or having sedimentation) but sparsely vegetated and consists largely of coarse diamictons with shallow soil depths (<40 cm) and low SOC contents (Table 2). The ‘ground moraine’ class occupies about 2% of the study area in the lower parts of the central valley (Figs. 1 and 2). The surfaces
25 have been stable since the early Holocene and are largely vegetated with only 1–5 % boulders. Soils have developed in the top meter, with signs of cryoturbation, leading to high SOC contents in the 0–100 cm depth interval (Table 2). Deposits below 100 cm depth in these two glacially deposited classes are considered tills, with very low SOC contents (Fig. 2 and Table 2).

30 Another widespread landform (15% of the study area) is referred to as ‘alluvial fans’ (n=15), with additional small areas of peaty fens (n=5) and bogs (n=3) on alluvial fans (0.4%). These are areas of high SOC storage with on average 19.8, 29.8 and 42.7 kg C m⁻² for the top 100 cm, respectively (Fig. 2 and Table 2). The alluvial fans are generally located in the foothills of the sedimentary northeastern slopes of the study area. Wetlands are common in the lowermost
35 reaches of these foothills, where the slope decreases and water accumulates. Alluvial fan deposits have fine-grained sediments, often containing thin buried C-enriched layers, which sometimes reach depths of >300 cm (maximum observed depth of 370 cm). As a result, this is the landform with some of the highest 100–200 and 200–300 cm SOC stocks (Fig. 2 and Table

2), contributing most to the deeper SOC storage in the study area. For calculation purposes bog hummocks (isolated palsas and pounus), with very high SOC stocks but occupying only 3% of the total wetland area, were separated from peaty fen areas to not overestimate total SOC storage in this class. Furthermore, wetlands (including bogs) on alluvial fans occupy only 0.4% of the total study area and their high SOC storage increases the weighed mean SOC storage for the entire study area by only 0.1 kg C m⁻². The small area of freshwater lakes (<1%) has intermediate SOC values down to 300 cm of depth from sediment surface (Fig. 2 and Table 2). Relict fluvial and (raised) deltaic landforms occupy about 4% of the study area, in the lower reaches of the central Zackenberg valley. These landform classes have high SOC values down to depths of 300 cm and more, contributing significantly to the overall deep SOC storage in the study area. Recent fluvial stream beds occupy about 8% of the area, but have low SOC storage values at all considered soil depth intervals (Fig. 2 and Table 2).

For the entire study area, the estimated weighed mean SOC storage is 4.8 kg C m⁻² in the top 100 cm (Table 2). When comparing the mean SOC 0–100 cm distribution among different layers, 13% is stored in the organic layer (87% in the mineral part), with 77% in the active layer (23% in upper permafrost). From a SOC 0–100 cm storage perspective, the alluvial fan is the most important landform, occupying 15% of the area with a mean SOC storage of 21.3 kg C m⁻² and holding ~ 60% of the total SOC 0–100 cm in the study area (Fig. 2). In contrast, landforms at higher elevation (bedrock, allochthonous weathered bedrock, solifluction sheets, and lateral/end moraine) occupy 70% of the study area but store only 15% of the total SOC 0–100 cm stocks, with mean SOC ranges between 0–2 kg C m⁻².

This study provides first SOC estimates for the second and third meter depth intervals in the Zackenberg study area. However, estimates for deeper layers are based on fewer sites (see subsection 3.2). Mean SOC storage from 100 to 200 cm depth decreased with 65% compared to the top meter to 1.7 kg C m⁻². The ‘alluvial fan class’ is the dominant landform also regarding SOC at this depth, with 8.6 kg C m⁻², contributing with 76% to the total SOC in the study area (Table 2 and Fig. 2). From 200 to 300 cm depth we estimated an additional SOC storage of 0.69 kg C m⁻², with 71% of the total SOC located in alluvial fans (Table 2 and Fig. 2). Thus, the estimated mean SOC storage for the top 300 cm soil depth is 7.2 kg C m⁻². However, two of our sampling sites (alluvial fan and delta) had fine-grained deposits that were deeper than 300 cm indicating that buried SOC is present beneath 300 cm depth (Cable et al., 20172018). Additionally, Cable et al. (20172018) report low C densities from two cores relatively close to the coast with 7 and 11 m thick deltaic sediments overlying glacial sediments. ~~However, the areal extent and the average sediment thickness of these deposits are not known.~~

The mean soil TN storage in the 0–100 cm depth interval in the Zackenberg study area is 0.28 kg TN m⁻² according to the geomorphological upscaling. The highest values are found in the alluvial fans including the wetlands, with a mean storage of 1.1 kg N m⁻² in alluvial fans and up to 2.9 kg N m⁻² in bogs (Table 2). However, because of the small spatial extent occupied by

wetlands their contribution to the total TN storage is only 2.5%, while alluvial fans are storing 59% of the soil TN at 0–100 cm in the active layer and top permafrost in the study area.

5 Discussion

This study presents new estimates of total storage and landscape partitioning of SOC and soil TN in the Zackenberg study area based on detailed geomorphological map upscaling. In comparison with the previous land cover classification (LCC) approach performed for the same area (Palmtag et al., 2015), the geomorphologically-based upscaling shows a 42% reduced weighed mean SOC 0–100 cm storage from 8.3 (\pm 1.8 CI) to 4.8 (\pm 1.0 CI) kg C m⁻². While the SOC 0–100 cm estimates for the organic and permafrost layers deviated little in comparison with the estimates of the previous LCC approach, ~~most-some~~ deviation occurred within the mineral active layer. We estimate that 77% of SOC 0–100 cm is located in the active layer. Likewise, the mean soil TN storage at 0–100 cm depth decreased with 44% from 0.50 kg (\pm 0.1 CI) to 0.28 (\pm 0.1 CI) kg TN m⁻².

Bartsch et al. (2016), who used a new approach based on synthetic aperture radar (SAR) which determined the values directly from backscatter intensity, reported SOC 0–100 cm values for the Zackenberg area of 15.0 kg C m⁻² (based on points where pedon data (n=24) was available). In comparison, the Northern Circumpolar Soil Carbon Database (NCSCD) reports even higher values for this region of 17.8 kg C m⁻² in the 0–100 cm depth interval. In contrast, Fuchs et al. (2015) reported a mean SOC 0–100 cm storage of 0.9 (\pm 0.2 CI) kg C m⁻² with 100% located in seasonal frost and/or active layer in a mountainous area of northern Sweden.

~~Theis~~ relatively large ~~difference-decrease~~ in SOC (and TN) 0–100 cm storage for Zackenberg using the landform approach is mainly due to the important role of geomorphological processes in redistributing sediments in mountainous areas, which is to some extent neglected when using LCCs primarily based on vegetation cover classification from satellite observations. For example, in the LCC-upscaling (Palmtag et al., 2015) the SOC-rich vegetated classes ‘grasslands’ and, to a lesser degree, ‘fens’ occupied relatively large proportions of the total study area (20% and 3% coverage, respectively). These included areas on slopes at mid-elevation with patchy grassland cover and wet areas along streambeds. However, most of the pedons for these classes were located in the foothills and central valley characterized by higher SOC and TN storage. This resulted in a pedon dataset that was not truly representative for its thematic class and a high-biased mean SOC and TN storage was applied to relatively large areas. The GLC-approach in the current study better identifies areas of high SOC and TN storage in depositional environments such as alluvial fans (including wetlands on alluvial fans) occupying more limited proportions of the total study area (15% and 0.4%, respectively).

Therefore, the substantial decrease of SOC and TN has occurred because the areal extent of the SOC and TN-rich vegetated classes (grasslands 20%, *Salix* snow bed 7%, *Dryas* heath 6%, *Cassiope* heath 4% and fen 3%) has been reduced from 40% in the LCC to 22% (alluvial fans 15.4%, delta (raised, relict) 2.5%, fluvial stream bed (relict) 1.2%, and ground moraine till

2.4%) using the landform upscaling. Also in non-alpine environments, such as the Lena River Delta, geomorphological setting better explained SOC variability than vegetation cover (Siewert et al., 2016).

When comparing the ~~confidence intervals (CI)~~ of the weighed mean SOC 0–100 cm estimates for the entire study area using LCC ($8.3 \pm 1.8 \text{ kg C m}^{-2}$) and GLC ($4.8 \pm 1.0 \text{ kg C m}^{-2}$) there is an absolute decrease in the uncertainty range, but in relative terms it remains similar (20%, also for TN). Comparable results are obtained for the mean and standard deviations (SD) of the dominant classes in both upscaling approaches (despite a marked difference in sample size), with ‘grasslands’ (n=6) in the LCC ($19.1 \pm 8.3 \text{ kg C m}^{-2}$) and ‘alluvial fans’ (n=15) in the GLC ($19.8 \pm 9.3 \text{ kg C m}^{-2}$), or a coefficient of variation of 45% in both cases. This might point to the fact that there is an intrinsic variability in SOC storage within these classes, related to microtopography, drainage, plant productivity, SOC burial, coarse fraction content, active layer thickness and ground ice volume and type, among others.

The geomorphological approach is particularly important in identifying areas of deep SOC storage related to depositional environments such as alluvial, fluvial and deltaic landforms, for which the (cryo)stratigraphy (including excess ground ice) should be taken into account. Alluvial fans can consist of up to several meters ($\leq 4 \text{ m}$) thick fine-grained laminated deposits accumulated during the Holocene due to the downslope sediment transport of materials and their subsequent deposition in the foothills by nivation processes (Christiansen, 1998; [Cable et al., 2018](#)). Intercalated SOC-enriched layers up to 8600 cal yr BP old (Cable et al., 2018) indicate the repeated burial of stable vegetated surfaces and/or organic material eroded by nival meltwater higher up. Relict deltaic areas in the lower central valley are another area of deep SOC storage, with deltaic deposits OSL-dated to 11–13 ka reaching depths of 7–11 m overlying a glacial till unit (Gilbert et al., 2017).

This study includes the first SOC storage estimates down to 300 cm depth with quantitative uncertainty ranges for the Zackenberg study area. We estimate that $1.7 \pm 0.8 \text{ kg C m}^{-2}$ and $0.7 \pm 0.4 \text{ kg C m}^{-2}$ is being stored in the second and third meter of deposits, respectively. This is a considerable amount, representing an additional 50% SOC compared to the storage in the top meter, mainly located in alluvial fan deposits. Deltaic deposits further contribute to the 100–300 cm SOC storage but locally reach depths of at least 13 m. The low carbon densities of these deepest deposits are similar to those reported for the 100–300 cm depth interval (Cable et al., 2018). Assuming these deltaic deposits have the same areal extent as today (2.48%) and a mean depth of 8 m based on new data published in Gilbert et al. (2017). Using the low carbon densities reported above, we can arrive at a preliminary additional storage in deltaic deposits >300 cm of 32 kg C m^{-2} , which would increase the study area weighed mean by a further 0.8 kg C m^{-2} . However, additional work is required to increase the accuracy of these values. However, their areal extent and average depth is unknown, which does not permit an accurate calculation of their total SOC storage.

6 Conclusions

This study presents new ~~additional~~ sampling sites and improved estimates of SOC and TN from 0 to 100 cm depth in Zackenberg (NE Greenland), based on upscaling using geomorphological landforms. Moreover, we report first SOC estimates to a depth of 300 cm.

5 Following our listed aims from the introductory section, we conclude: (1) The updated weighed landscape-level mean SOC and TN 0–100 cm storage in the Zackenberg study area based on geomorphological mapping ~~is~~ are 4.8 kg C m⁻² and 0.28 kg TN m⁻², respectively. (2) The new mean SOC and TN estimates to 100 cm depth are 44% and 42% less than previously reported estimates of 8.3 kg C m⁻² and 0.50 kg TN m⁻², when using a LCC upscaling approach. ~~42% less than previously reported, when using a LCC upscaling approach. The new mean SOC estimate is 4.8 kg C m⁻² to 100 cm depth, compared to the original estimate of 8.3 kg C m⁻². (3) Our deep pedon dataset indicates that an additional 1.7 and 0.7 kg C m⁻² are stored in the second and third meter soil depth, respectively. (4) A previous areal overestimate of SOC-rich vegetated land cover classes on slopes was the main reason for this large difference in SOC~~
10 ~~storage. Downslope creep constantly transport material downslope resulting in relatively shallow soil depths and low SOC storage. Slope materials have accumulated at the foot of the slopes in alluvial fans during the entire Holocene, primarily by nivation processes creating thick, fine-grained deposits with buried SOC-enriched layers throughout their depth. Whereas the LCC recognized vegetated SOC-rich classes on the slopes, in the foothills and in the central valley, the GLC upscaling more correctly restricts the SOC-rich classes to areas with deposition. The use of LCC upscaling in these mountainous settings can introduce large uncertainties since it is based on recent land cover and vegetation only that do not necessarily reflect the long-term geomorphic processes leading to SOC burial. To the contrary, the landform-based approach identifies hotspots of SOC burial in the landscape such as alluvial fans and (to a minor extent) deltas. The GLC approach is, therefore, also highly relevant when identifying areas of deep carbon storage (between 100–300 cm, and more). Our deep pedon dataset indicates that an additional 1.7 and 0.7 kg C m⁻² are stored in the second and third meter soil depth, respectively.~~ Deltaic deposits extend below 300 cm depth, which implies that there are additional SOC stocks at greater depth but these remain poorly constrained at this time. The results emphasize the importance of geomorphology, rather than land cover, controlling SOC storage in high relief permafrost environments.

Acknowledgements

Fieldwork in Zackenberg was financed through the Nordforsk Nordic Centre of Excellence DEFROST project (grant number 23001), the EU FP7 PAGE21 project (grant number 282700)
35 and the ESF CryoCARB project (Swedish Research Council project to P. Kuhry). Additional funding for fieldwork in 2013 and subsequent core analyses were supported by the University Centre in Svalbard (UNIS) and the Centre for Permafrost (CENPERM) at the University of

Copenhagen, funded by the Danish National Research Foundation (CENPERM DNR100). We gratefully acknowledge the support with drilling and core analyses by co-leader Professor Bo Elberling (CENPERM, University of Copenhagen) and students participating in the UNIS course AG-833 'High Arctic Permafrost Landscape Dynamics in Svalbard and Greenland' in 2013, funded by the University Centre in Svalbard (UNIS), the Norden Perma-Nordnet project and the Centre for Permafrost (CENPERM), University of Copenhagen. We would like to thank the GeoBasis Programme of the Zackenberg research station for providing data (overview map and DEM data).

References

- 10 | [Bartsch, A., Widhalm, B., Kuhry, P., Hugelius, G., Palmtag, J., and Siewert, M. B.: Can C-band synthetic aperture radar be used to estimate soil organic carbon storage in tundra?, *Biogeosciences*, 13, 5453-5470, doi.org/10.5194/bg-13-5453-2016, 2016.](#)
- Bennike, O., Sørensen, M., Fredskild, B., Jacobsen, B.H., Böcher, J., Amsinck, S.L., Jeppesen, E., Andreasen, C., Christiansen, H.H. and Humlum, O.: Late Quaternary environment and cultural changes in the Wollaston Forland region, northeast Greenland. *Advances in Ecological Research*, 40, 45–79, doi:10.1016/S0065-2504(07)00003-7, 2008.
- 15 | [Cable, S.; Christiansen, H.H.; Westergaard-Nielsen, A.; Kroon, A.; Elberling, B \(2017\): Geomorphological and geocryological maps of the Zackenberg Valley, NE Greenland \(ArcGIS Map Package\). PANGAEA, Supplement to: Cable et al., 2017, doi.org/10.1594/PANGAEA.883484, 2017.](#)
- 20 | Cable, S., Christiansen, H.H., Westergaard-Nielsen, A., Kroon, A. and Elberling, B.: Geomorphological and cryostratigraphical analyses of the Zackenberg Valley, NE Greenland and significance of Holocene alluvial fans, *Geomorphology*, 303, 504–523, doi.org/10.1016/j.geomorph.2017.11.003, 2018.
- 25 | Christiansen, H.H.: Nivation forms and processes in unconsolidated sediments, NE Greenland. *Earth Surface Processes and Landforms*, 23, 751-760, 1998.
- Christiansen, H.H., Sigsgaard, C., Humlum, O., Rasch, M. and Hansen, B.U.: Permafrost and periglacial geomorphology at Zackenberg. *Advances in Ecological Research*, 40, 151–174, doi:10.1016/S0065-2504(07)00007-4, 2008.
- 30 | Escher, A. and Watt, W.S.: The geology of Greenland. The Geological Survey of Greenland, pp. 603, 1976.
- [Fuchs, M., Kuhry, P., and Hugelius, G.: Low below-ground organic carbon storage in a subarctic Alpine permafrost environment, *The Cryosphere*, 9, 427-438, doi.org/10.5194/tc-9-427-2015, 2015.](#)
- 35 | Gilbert G.L., Cable, S., Thiel, C., Christiansen, H.H. and Elberling, B.: Cryostratigraphy, sedimentology, and the late Quaternary evolution of the Zackenberg Delta, Northeast Greenland. *The Cryosphere*, 11, 1265-1282, doi:10.5194/tc-11-1265, 2017.

- Gruber, N, Friedlingstein, P., Field, C.B., Valentini, R., Heimann, M., Richey, J.E., Romero-Lankao, P., Schulze, D. and Chen, C-T.A.: The vulnerability of the carbon cycle in the 21st century: An assessment of carbon-climate-human interactions. In: *The Global Carbon Cycle: pp 45–76. Integrating Humans, Climate and the Natural World*, Field C, Raupach M (Eds.). 5 Island Press, Washington D.C, 2004.
- Hansen, B.U., Sigsgaard, C., Rasmussen, L., Cappelen, J., Hinkler, J., Mernild, S.H., Petersen, D., Tamstorf, M.P., Rasch, M. and Hasholt, B.: Present-day climate at Zackenberg. *Advances in Ecological Research*, 40, 111–149, doi:10.1016/S0065-2504(07)00006-2, 2008.
- Heiri, O., Lotter, A.F. and Lemcke, G.: Loss on ignition as a method for estimating organic 10 carbon and carbonate content in sediments: reproduction and comparability of results. *Journal of Paleolimnology*, 25, 101–110, doi:10.1023/a:1008119611481, 2001.
- Hugelius, G.: Spatial upscaling using thematic maps: An analysis of uncertainties in permafrost soil carbon estimates, *Global Biogeochemical Cycles*, 26, GB2026, doi:10.1029/2011GB004154, 2012.
- 15 Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J.W., Schuur, E.A.G., Ping, C.-L., Schirmermeister, L., Grosse, G., Michaelson, G.J., Koven, C.D., O'Donnell, J.A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J. and Kuhry, P.: Estimated storages of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps. *Biogeosciences*, 11, 6573–6593, doi:10.5194/bg-11-6573-2014, 2014.
- 20 IPCC: Summary for Policymakers. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, 2013.
- 25 Kuhry, P., Mazhitova, G.G., Forest, P.-A., Deneva, S.V., Virtanen, T. and Kultti, S.: Upscaling soil organic carbon estimates for the Usa Basin (Northeast European Russia) using GIS-based landcover and soil classification schemes. *Danish Journal of Geography*, 102, 11–25, 2002.
- Palmtag, J., Hugelius, G., Lashchinskiy, N., Tamstorf, M.P., Richter, A., Elberling, B. and 30 Kuhry, P.: Storage, landscape distribution and burial history of soil organic matter in contrasting areas of continuous permafrost. *Arctic, Antarctic, and Alpine Research*, 47(1), 71–88, doi.org/10.1657/AAAR0014-027, 2015.
- Palmtag, J., Ramage, J., Hugelius, G., Gentsch, N., Lashchinskiy, N., Richter, A., and Kuhry, P.: Controls on the storage of organic carbon in permafrost soils in northern Siberia. *European Journal of Soil Science*, 67, 478–491, doi:10.1111/ejss.12357, 2016.
- 35 [Ping, C.-L., Michaelson, G.J., Jorgenson, M.T., Kimble, J.M., Epstein, H., Romanovsky, V.E. and Walker, D.A.: High stocks of soil organic carbon in North American Arctic region. *Nature Geoscience*, 1, 615-619, doi:10.1038/ngeo284, 2008.](#)
- [Romanovsky, V. E., Smith, S. L. and Christiansen, H. H.: Permafrost thermal state in the polar Northern Hemisphere during the international polar year 2007–2009: a synthesis. *Permafrost Periglac. Process.*, 21: 106–116, doi:10.1002/ppp.689, 2010.](#)
- 40

Schuur, E.A.G., McGuire, A.D., Schädel, C., Grosse, G., Harden, J.W., Hayes D.J. Hugelius, G., Koven, C.D., Kuhry, P., Lawrence, D.M., Natali, S.M., Olefeldt, D., Romanovsky, V.E., Schaefer, K., Turetsky, M.R., Treat, C.C. and Vonk, J.E.: Climate change and the permafrost carbon feedback. *Nature*, 20, 171–179, doi:10.1038/nature14338, 2015.

5 Siewert, M.B., Hanisch, J., Weiss, N., Kuhry, P., Maximov, T.C. and Hugelius, G.: Comparing carbon storage of Siberian tundra and taiga permafrost ecosystems at very high spatial resolution. *Journal of Geophysical Research: Biogeosciences*, 120, 1973–1994, doi:10.1002/2015JG002999, 2015.

10 [Slater, A.G. and Lawrence, D.M.: Diagnosing Present and Future Permafrost from Climate Models. *Journal of Climate*, 26, 5608-5623, doi.org/10.1175/JCLI-D-12-00341.1.](#)



Soil Survey Staff: Keys to Soil Taxonomy. Twelfth Edition. USDA-Natural Resources Conservation Service, Washington, D.C., 2014.

15 [Tarnocai, C., Swanson, D., Kimble, J. and Broll, G.: Northern Circumpolar Soil Carbon Database, Digital Database, Research Branch, Agriculture and Agri-Food Canada, Ottawa, Canada, 2007.](#)


Tarnocai, C., Canadell, J.G., Schuur, E.A.G., Kuhry, P., Mazhitova, G. and Zimov, S.: Soil organic carbon pools in the northern circumpolar permafrost region. *Global Biogeochemical Cycles*, 23, GB2023, doi:10.1029/2008GB003327, 2009.

Thompson, S.K.: Sampling. John Wiley, New York, 343pp, 1992.

20 [Zimov, S. A., E. A. G. Schuur, and F. S. Chapin III.: Permafrost and the global carbon budget, *Science*, 312, 1612–1613, doi:10.1126/science.1128908, 2006.](#)



5 **Figure 1. A) Simplified geomorphological landform map of the Zackenberg study area (based on Cable et al., 2017). Top-right B): Location of Zackenberg in NE Greenland.**



5 Figure 2. Proportional contribution of each landform to: A) Areal coverage of the total study area; B) total SOC storage for 0–100 cm; C) total SOC storage for 100–200 cm; and D) total SOC storage for 200–300 cm. Landform and colors from chart A apply also to charts B-D.

Table 1. Geomorphological landforms and their proportion of total surface area as presented in Cable et al. (2017), and the amalgamated larger landform classes used in this study (excluding Young Sound).

Geomorphological landforms	Area %	Overall geomorphological classes	Area %
Bedrock, exposed	7.62	Bedrock	7.62
Bedrock, weathered, allochthonous	26.0	Allochthonous weathered bedrock	30.4
Debris cone, debris fan	2.50		
Rock glacier	0.34		
Perennial snow patch ¹	1.53		
Slide scar	0.03	Solifluction sheets	14.3
Solifluction sheet, solifluction lobe	14.3		
Moraine, ice-cored	4.78	Lateral / end moraines	18.1
Lateral moraine	14.3		
Ground moraine	2.39	Ground moraine	2.39
Alluvial fan, coarse-grained, inactive	0.47	Alluvial fans	15.0
Nivation hollow	3.55		
Alluvial fan	10.9		
Wetland ²	0.38	Wetlands (alluvial fan) ²	0.38
River, channel, relict	1.20	Fluvial stream bed (relict)	1.20
Delta, relict	2.43	Delta (raised, relict)	2.48
Sandbar, relict	0.05		
River, channel	3.48	Fluvial stream bed (recent)	7.97
Braided river	1.94		
Alluvial fan, coarse-grained	1.32		
Delta pre-recent	0.54		
Spit	0.05		
Beach	0.12		
Delta, active	0.53		
Lacustrine	0.00	Lakes	0.25
Lake	0.25		
Total	100		100

5

¹ Snow patches found at higher elevations among the other landform units belonging to the same amalgamated landform

² Wetland areas developed on top of alluvial fan deposits

Table 2. Mean SOC and TN estimates (kg m⁻²) for the Zackenberg study area by landforms.

Landforms	Type of deposits	Area %	n sites / samples	Organic layer kg C m ⁻²	Mean ± SD SOC/TN storage					
					0–30 cm kg C m ⁻²	0–100 cm kg C m ⁻²	Permafrost in 0–100 cm kg C m ⁻²	100–200 cm kg C m ⁻²	200–300 cm kg C m ⁻²	0–100 cm kg N m ⁻²
Bedrock		7.62	1 / 1	0.0	0.0	0.0	0.0	0.0	0.0	0.0
Allochthonous weathered bedrock	colluvial deposits	30.4	2 / 14	0.0 ± 0.0	0.1 ± 0.1	0.3 ± 0.4	0.2 ± 0.3	0.1 ± 0.1	0.0	0.0
Solifluction sheets	colluvial deposits	14.3	3 / 32	0.1 ± 0.6	0.1 ± 0.4	1.7 ± 0.9	0.0	0.2 ± 0.3	0.0	0.1 ± 0.1
Lateral / end moraines	glacial deposits	18.1	4 / 21	0.1 ± 0.1	0.9 ± 1.1	2.1 ± 3.2	1.1 ± 1.3	0.0	0.0	0.1 ± 0.1
Ground moraine	glacial deposits	2.39	3 / 33	1.6 ± 0.2	6.7 ± 3.1	15.9 ± 4.0	1.7 ± 2.8	2.1 ± 1.8	0.0	1.1 ± 0.3
Alluvial fans	alluvial deposits	15.0	15 / 260	2.5 ± 3.0	7.3 ± 3.6	19.8 ± 9.3	3.9 ± 4.1	8.6 ± 10.0	3.3 ± 5.8	1.1 ± 0.6
Wetlands (alluvial fan)	peat/alluvial deposits	0.36	5 / 89	16.0 ± 17.7	11.0 ± 3.4	29.8 ± 12.5	14.4 ± 10.1	9.1 ± 10.0	2.4 ± 2.2	1.9 ± 0.7
Bog (alluvial fan)	peat/alluvial deposits	0.01	3 / 48	36.5 ± 8.8	16.0 ± 9.6	42.7 ± 7.0	24.6 ± 17.2	6.2 ± 2.6	6.2 ± 2.6	2.9 ± 0.5
Lakes	lacustrine deposits	0.25	4 / 43	11.4 ± 1.3	5.6 ± 2.2	14.5 ± 2.8	1.0 ± 1.2	5.0 ± 2.5	2.1 ± 1.8	0.4 ± 0.0
Fluvial stream bed (relict)	fluvial deposits	1.20	2 / 24	2.3 ± 0.8	6.2 ± 0.2	20.9 ± 4.5	8.4 ± 6.3	4.4 ± 1.3	4.7 ± 1.7	1.4 ± 0.4
Delta (raised, relict)	deltaic deposits	2.48	4 / 78	1.2 ± 0.8	4.4 ± 1.6	9.1 ± 3.2	2.2 ± 2.3	6.7 ± 4.4	4.0 ± 1.4	0.7 ± 0.3
Fluvial stream bed (recent)	fluvial deposits	7.97	2 / 5	0.1 ± 0.2	0.8 ± 0.8	1.0 ± 0.5	0.0	0.3 ± 0.5	0.3 ± 0.5	0.0
Mean ± CI SOC/TN storage (weighed by landform proportion)		100	48 / 648	0.6 ± 0.3	1.9 ± 0.4	4.8 ± 1.0	1.1 ± 0.4	1.7 ± 0.8	0.7 ± 0.4	0.3 ± 0.1