

Response Letter

We appreciate two reviewers for investing their time and providing constructive comments on our manuscript. Below, we explain changes we made and present our opinions to the points that we did not follow. We hope that these revisions are satisfactory, and the revised manuscript would meet journal's criteria.

Some major changes to the manuscript are –

- The introduction is reorganized/rephrased to emphasize more on the motivations behind the study.
- The manuscript structure is reorganized to a more conventional structure as recommended by reviewer #2. The changes in the structure are shown in the figure below

–

Previous

1. Introduction
2. Blåskimen Island
3. Field measurements
 1. Kinematic and static GPS surveys
 2. Ice-penetrating radar profiling
 3. Firn cores and borehole-temperature measurements
4. Topography and surface flow field
5. Surface mass balance
 1. SMB derived from stake heights
 2. SMB derived from shallow-sounding radar stratigraphy
6. Mass balance of the ice rise
 1. Methods
 2. Estimate setups
 - i. Polygon setup
 - ii. Grid setup
 - iii. Flowband setup
 3. Uncertainties
 4. Results
7. Discussion
 1. Topographic characteristics
 2. Surface mass balance
 3. Present day mass balance
 4. Long-term evolution
 5. Impact on the adjacent ice shelves
8. Conclusions

New

1. Introduction
2. Blåskimen Island
3. Field measurements and data processing
 1. Kinematic and static GPS surveys
 2. Ice-penetrating radar profiling
 3. Firn cores and borehole-temperature measurements
4. Analytical methods
 1. Surface mass balance
 1. Deriving SMB using stake heights
 2. Deriving SMB using shallow-sounding radar stratigraphy
 2. Mass balance of the ice rise
 1. Constraining γ
 2. Estimate setups
 1. Polygon setup
 2. Grid setup
 3. Flowband setup
5. Results
 1. Topography and surface flow field
 2. Surface mass balance
 3. Mass balance
6. Discussion
 1. Topographic characteristics
 2. Surface mass balance
 3. Present day mass balance
 4. Long-term evolution and impact on the adjacent ice shelves
7. Conclusions

- Based on recommendation from reviewer #1, we have used the first radar method to derive SMB accounting for the vertical variability in density (not lateral variability across the whole ice rise). Consequently, the mass balance estimate is updated based on this SMB as input. Figure 2, 4, 6 and 7 were updated with the new estimate.

Below, we quote reviewers comments first, and then our responses in blue italic fonts follow. Otherwise stated, page/line numbers in this letter refer the numbers in the marked manuscript. Due to a technical issue with the software we use (Latexdiff) to prepare the marked manuscript, changes in section names are not highlighted in the marked manuscript. Also, long citations overflow the page on a few occasions. We have listed these citations at the end of this response letter.

Responses to Reviewer #1

Review of “Glaciological settings and recent mass balance of the Blåskimen Island in Dronning Maud Land.”

General Comments

Goel and coauthors report field observations from a relatively large ice rise that sits between Jelbert and Fimbul Ice Shelves in Dronning Maud Land. They use two ice-penetrating radars with different frequencies, stake measurements and surface GPS, to constrain the topography, thickness and internal stratigraphy of the ice rise as well as obtain several spatially-distributed estimates of mass balance.

The observations are well reported and are likely to be of interest to the glaciological community. Specifically, they may interest those interested in the surface mass balance and long-term evolution of the ice rises in Dronning Maud Land. As the authors mention, ice rises are important because they influence regional ice dynamics and encode information about the stability or otherwise of these dynamics in their internal stratigraphy. Ice rises also influence atmospheric circulation and therefore surface mass balance and are useful locations for drilling ice cores. Overall, any information we can gain on the glaciological setting and mass balance of ice rises, as well as their long-term stability is useful and worth publishing. In my opinion, this work should be published in the Cryosphere after revision to address the comments below.

Overall, the manuscript is written well, with some exception. There are many grammatical and other errors, as well as some passages that I found it difficult to understand. These difficult-to-understand phrases, various errors and other places where I propose that rewording could improve the paper, are indicated below in “Technical Corrections”. Note that due to the high density of errors, there are likely to be many that I missed and I suggest that any revised manuscript is read carefully for this kind of error.

We thank all of your efforts. We regret that the previous manuscript includes typos and grammatical errors, which we tried to fix as much as we can in this version.

Individual scientific comments are described below in the “Specific Comments” section.

Specific Comments

Section 5.2 describes how shallow radar layers are used to derive SMB. This involves multiplying the depths of three dated layers by the snow/firn density to compute the mass accumulated since each layer was laid down. Spatial variations in density are taken account of using two methods. Very little detail is

provided of the second method. An in preparation manuscript is cited and one sentence (L3-5, P7) describes the inverse method. Because (1) very little detail is provided (certainly too little to be able to reproduce the results) and (2) this method is apparently used to take account of a spatial variation in nearsurface density of only $\pm 2\%$, I suggest this alternative method is removed from the manuscript. The first method takes account of the much larger vertical variation in density and seems adequate. I would suggest removing the second method and simply stating that in this case the $\pm 2\%$ variation in density is small and that it is possible to alternatively take account of this at the expense of taking into account the vertical variations and that a manuscript is in preparation that will report ongoing work related to doing this with an inverse method.

We agree with the reviewer and use the SMB estimated with the first radar method to derive mass balance of the ice rise. However, we briefly mention the second radar method and results obtained with this method to give an approximate idea on errors in SMB estimates related to the uncertainties of density.

Page 5, L14-15: A value for the SMB of the whole island is quoted and an uncertainty is estimated based on measurement errors and uncertainty accounting for snow densification. Does this uncertainty estimate also take into account uncertainty associated with the interpolation? This is difficult (as the authors point out later in the manuscript), so if not, perhaps it is better to quote the SMB as the average of the stakes measurements, rather than as the average for the whole island, as this reads to me.

No, uncertainty associated with the interpolation is not included in the error estimate. As the reviewer said, it is difficult to do. In the revised manuscript, the mean SMB only used the stake measurements. We have rephrased the text at P12L2 to make it clear.

“The mean SMB from 90 stake-height measurements across Blåskimen Island is 0.78 m a^{-1} , for the period between January 2013 and January 2014 (Fig. 5a).”

Conclusions, page 13, L21-22: As sastrugi patterns can change from one day to the next, I am not sure that this a robust conclusion to draw about SMB patterns from unquantified observations of sastrugi in the field. I agree that in general it is tempting to conclude that there must be lower accumulation where you see sastrugi in the field (I have seen this myself on lower accumulation sides of ice rises), so I think it is reasonable to mention this earlier in the manuscript, but I don't think it belongs here as one of your main conclusions.

We agree with the reviewer and removed this sentence from the conclusion.

Technical Comments

Title: To my reading the ‘the’ before ‘Blåskimen Island’ is unnecessary. Indeed, in the abstract the name of the ice rise appears without the ‘the’. *Page 1*

Corrected.

L7: Suggest replacing the sentence starting with “Radar stratigraphy...” with “Arches in radar stratigraphy observed with radar suggest that the summit of the ice rise has been stable for ~600 km.” This avoids introducing the concept of characteristic time in the abstract, which may be confusing to those not familiar with it.

Corrected.

L11: Is there a reference for the statement about 74% of the coastline being surrounded by ice shelves?

It is mentioned in Bindschadler et al. (2011). We cite this reference in the revised manuscript:

Bindschadler, R., Choi, H., Wichlacz, A., Bingham, R., Bohlander, J., Brunt, K., Corr, H., Drews, R., Fricker, H., Hall, M., Hindmarsh, R., Kohler, J., Padman, L., Rack, W., Rotschky, G., Urbini, S., Vornberger, P. and Young, N.: Getting around Antarctica: new high-resolution mappings of the grounded and freely-floating boundaries of the Antarctic ice sheet created for the International Polar Year, The Cryosphere, 5(3), 569–588, doi:10.5194/tc-5-569-2011, 2011.

L14: ‘are’ → ‘is’.

Corrected.

L15: Suggest delete ‘eventual’.

The sentence was removed in the revised introduction.

L16: ‘the ice shelf’ → ‘ice shelves’.

Corrected.

L16-17: this sentence about buttressing seems to repeat what was said in the previous paragraph.

Agreed. We make this point only once in the revised introduction. (PIL19) in the marked manuscript.

L23: Two more recent papers that use isochrones to constrain millennial scale ice-rise evolution that could be referenced here are:

Drews, R., Matsuoka, K., Martín, C., Callens, D., Bergeot, N. and Pattyn, F., 2015. Evolution of Derwael ice rise in Dronning Maud Land, Antarctica, over the last millennia. *Journal of Geophysical Research: Earth Surface*, 120(3), pp.564-579.

and

Kingslake, J., Martín, C., Arthern, R.J., Corr, H.F. and King, E.C., 2016. Ice flow reorganization in West Antarctica 2.5 kyr ago dated using radar-derived englacial flow velocities. *Geophysical Research Letters*, 43(17), pp.9103-9112.

Good point. We are aware of these references, but didn't include them in the first manuscript. Now these references are cited.

Page 2

L1: Suggest replace ‘using an ice core from the summit’ with ‘using ice cores drilled through ice rises’.

Corrected.

L4: insert comma after ‘(Fig. 1a)’.

Corrected.

L12: Is it narrower by a factor of two and slower by a factor of two or does the factor of two only apply to the 'slower'? Suggest switch order to '...slower by a factor of two and narrower...' if appropriate, to avoid confusion.

We meant that ice flows slower by a factor of two and narrower. We corrected as suggested.

L16: Reword the last sentence. 'Whereas towards east...' doesn't read well to me.

We revised the last two sentences in this paragraph. The revised ones are (P2L32):

"Thus towards east, Blåskimen Island is surrounded by much slowly moving ice, whereas it abuts the eastern shear margin of the fast flowing part of the Jelbart Ice Shelf."

L19: 'summer' → 'summers'.

Corrected.

Page 3

L3: 'were' → 'was'

Not changed. As stakes are plural.

L4: How many were lost and why were they lost? Were they buried? Also did you have a criterion for rejecting stakes which were tilted too much? Surely all stakes are tilted slightly. If so, it's not correct to say here that you rejected stakes that were tilted.

We lost seven stakes out of 97 stakes. Six of them were in the southeastern slope. Stakes nearby these lost stakes show that this area has highest SMB in our study area. Stakes were 2 m high above the snow surface when they were installed and they were nearly buried one year after. So we speculate that these stakes were likely buried. One tilted stake was found in the northwestern slope, where snow surface is rough due to sastrugi. They were tilted more than ~20°, whereas the rest of stakes were tilted much smaller (only degrees or less). We then rejected this tilted stake from further analysis.

We have rephrased the sentence to be clearer; the revised sentence is (P3L18):

"The position of each of the stakes was measured in January 2013. In January 2014 we remeasured the positions of the 90 of these stakes. Six stakes were buried and one found heavily tilted (more than ~20°)."

L5: Suggest move 'relative' to between 'stakes' and 'to'.

Corrected.

L6: Suggest 'infer approximate' → 'estimate'.

Corrected.

L13: Suggest 'to the sea level' → 'to heights above local sea level'.

Corrected.

L13: This reads as if you used the gravity product to subtract the value. Suggest reword to say that you subtracted the value that was supplied by the gravity product.

The new sentence is (P3L28):

“To convert heights above the WGS84 ellipsoid to the sea level, we subtracted 13 m of geoid height uniformly provided by GOCE gravity product (<https://earth.esa.int/web/guest/dataaccess/browse-data-products/-/article/goce-gravity-fields-5777>).”

L19: ‘profiles’ → ‘surveys’. It looks like you did more than two profiles.

Corrected.

L21: Delete ‘a’.

Corrected.

L22: Delete commas after ‘radar’.

Corrected.

L23: Insert ‘a’ before dewow and Ormsby. Also ‘Post processing was made...’ does not read well.

Corrected. The revised sentence is (P4L9):

“Post processing included using a dewow filter, an Ormsby band-pass filter, and depth-variable gain functions.”

L24: ‘the’ → ‘a’ and insert ‘of’ after ‘speed’.

Corrected.

L25: How is the firn correction computed? Using the firn cores presumably, but this is not mentioned here.

We added the following to text to explain this point (P4L11):

“We used an optimization inversion routine (Brown and Matsuoka, in prep) to model depth profiles of density along shallow radar profiles using (1) surface densities measured at 9 locations, and (2) depth profile of the density measured along the 23-m-long firn core. We then used the depth profiles of the density along the radar profile to estimate laterally variable firn corrections. The magnitudes of the firn correction vary between 4–6 m.”

L29: Transported the samples where?

We transported the frozen samples to laboratories for further analysis. We rephrased the sentence as (P4L18):

“We drilled a 23 m long firn core near the summit of Blåskimen Island and transported the frozen samples to laboratory to develop visual stratigraphy as well as chemical, isotope and dielectric stratigraphy.”

L30: Suggest delete 'backed'.

Replaced 'backed' with 'back'.

Page 4

L1: 'to' → 'in', insert 'the' between 'from' and 'surface'.

Corrected.

L5: delete 'of'.

Corrected.

L9-10: suggest delete 'giving a relative dome-shaped topography to the ice rise'.

We did not delete this sentence, as we wanted to emphasize the overall shape of the ice rise.

L11: Not clear what 'a flatter basin northeast' means and it is grammatically incorrect. I cannot see a clear distinction between slopes in the northeast and in the other directions. Also I can't see clear distinctions between basins in figure 2a.

To clarify these issues, we revised the sentences as (P11L7):

"The eastern flank shows locally steep slopes and a basin in northeast with overall lower, less-tilted surface. The eastern steep slopes and the southwest ridge are consistent with lineations observed in satellite imagery (light gray feature over dark gray in Fig. 1b)."

L12: The line 2-2' isn't strictly a flow line as it passes through the ice divide. I guess it is two flow lines connected together, but describing it as a flow line I don't think is correct.

Replaced flowline to profile.

L17: The estimate of the vertical uncertainty of $\pm 5\text{m}$ could be more fully explained here. The center frequency of the deep radar was 2 MHz, corresponding (I think) to a wavelength of $c/n_i/2 \times 10^6 = 84.2\text{ m}$, where c is the speed of light and n_i is the refractive index of ice. This wavelength is considerably more than your estimated uncertainty in digitizing the bed reflector. Is the higher precision achieved due to the signal being quite broadband? Perhaps this can be explained in more detail, as one might expect a bed reflector imaged with an 84 m wavelength radar to manifest as a layer thicker than 5 m.

The center frequency of the deep-sounding radar is 2 MHz, giving the wavelength in ice approximately 84 m. Therefore, this radar is not capable to distinguish two objects that are separated less than 42 m (half of the wavelength). Nevertheless, this radar is capable to detect the range to the target more precisely. We sampled returned wave at 100 MHz, or every 10 ns. Over this period, radio wave travels about 2 m.

L27: Suggest delete 'below'.

Corrected.

L28: The surface velocity measurements are described as a surface velocity field here, when they really appear as just point measurements in figure 2c, rather than a field.

We used large number of point measurements of the ice-flow vector to estimate the flow field. We think that the manuscript is clear enough.

L28: Suggest 'from' → 'of'.

Corrected.

Page 5

L9: Is this surface density or the density averaged over the top 3 m of all the cores? It is confusing because the previous sentence mentions "surface density variations", but this sentence only says 'firm density'.

We now define the density averaged over the top 3 m (measured using 3-m-long firm cores) surface density at P4L28 and consistently use it throughout the manuscript to avoid confusion.

L10: Suggest replace 'To estimate..' with 'When estimating SMB below...', or similar.

Corrected.

L17: It is stated here that *most* SMB values lie between the first and third quartiles, but doesn't this by definition mean that half the measurements lie between these two values (i.e. half ≠ most)?

The reviewer is correct. The sentences are improved

from

SMB varies by a factor of 3.3 over the study area of ~20 km by ~20 km (0.28–1.03 m a⁻¹), although most SMB values calculated with this method range between 0.73 and 0.87 m a⁻¹ (first and third quartiles, respectively).

to

SMB varies by a factor of 3.3 over the study area of ~20 km by ~20 km (0.28–1.03 m a⁻¹), with 80% of values ranging between 0.69 and 1.03 m a⁻¹.

L22: I am not sure I see the relevance here of the parentheses about wind direction.

Previous studies show that SMB is largely related to the local slope (e.g. King (2004)), as wind is faster over a steeper slope. We didn't see this relationship clearly in our data (Fig. 2a) but we think that this is because the wind direction is oblique to the profile we sampled. With the text in parenthesis we want to clarify that as the slope in discussion is not aligned along the prevailing wind direction. Therefore, the observed result of 'no clear relationship' does not necessarily disagree with previous studies. No change was made in the manuscript.

Page 6

L6: Is it not more precise to say that it is the larger vertical strain rates near the surface at the divide (rather than strong horizontal variations in vertical strain rates) due to the Raymond effect that could mean that the shallow-layer approximation may not be valid at the divide. Gillet-Chaulet et al. (2013) and

Kingslake et al. (2014) and probably others have measured higher magnitude vertical strain rates near the surface on ice divides and could be cited here to support the point.

Gillet Chaulet, F., Hindmarsh, R.C., Corr, H.F., King, E.C. and Jenkins, A., 2011. In situ quantification of ice rheology and direct measurement of the Raymond Effect at Summit, Greenland using a phase sensitive radar. *Geophysical Research Letters*, 38(24).

Kingslake, J., Hindmarsh, R.C., Aðalgeirsdóttir, G., Conway, H., Corr, H.F., Gillet Chaulet, F., Martin, C., King, E.C., Mulvaney, R. and Pritchard, H.D., 2014. Full depth englacial vertical ice sheet velocities measured using phase sensitive radar. *Journal of Geophysical Research: Earth Surface*, 119(12), pp.2604-2618.

We agree with the reviewer. We included the suggested references and changed the text accordingly. The revised text are from P6L29-P6L34:

“However, this assumption may not be valid in areas where vertical strain rates are large, such as the region near an ice-flow divide (Gillet et al., 2011; Kingslake et al., 2014). In this region, accumulated effects of variable vertical strain can result in upward arches in isochrones, so-called Raymond arches (Raymond, 1983), which were found at many other ice rises (e.g. Vaughan et al., 1999; Conway et al., 1999). Such upward arches can also be caused by anomalously low SMB near the summit possibly due to wind erosion (Drews et al., 2013, 2015).

Eqn 2: This equation appears incorrect. The z on the top of the fraction in the integrand on the right side shouldn't be there.

We apologize for the typo in the equation. The correct equation is

$$t(z) = \int_0^z \frac{2}{v(z)} dz$$

L30: I do not see why using a constant density (when in fact the density varies with space) would make the problem ill-posed. It may make the MB inaccurate, but why ill-posed?

With this sentence, we wanted to say that the constant density assumption will make the MB calculation inaccurate. The sentence has been corrected as (P12L23):

“It implies that the use of a uniform density could make the SMB estimates less accurate, and variable density should be accounted for.”

L32: Is there a word missing from near the start of this sentence?

We added “for” so that the sentence starts with

“To account for density variations.”

Page 7

L19-20: There is a comparison made here between SMB derived with two different methods. In my specific comments above I suggest that the second of these methods be removed from the manuscript, but here I would suggest that if both methods must be included that the comparison is made clearer here. As figure 4 stands, we have to compare spatial patterns in the two fields, but much of the structure in these

fields appears to depend on the details of the interpolation, i.e. how the surface is interpolated across the large gaps between data points. The statement here that the two methods ‘give nearly identical spatial patterns in SMB’ doesn’t appear to be correct, but the differences that stand out in figure 4 are due to the interpolations. So, I suggest that you include a plot that compares the two datasets without including an interpolated field. For example, a profile along the line 2-2’ of the SMB estimates, or remove the spatial dimension entirely and include a scatter plot of the estimates from each method on each axis.

While making the comparison between the two SMB estimates derived from radar stratigraphy we only used the data along the radar profiles and did not use interpolation data. Hence the comparison is free of any interpolation effects. After considering your comment above, we have decided to use the first radar SMB estimate for the mass balance calculation. Although we still mention the second radar method for discussion purposes. Based on your suggestion we also included a comparison of the two methods in Fig 3a, as well as the stake SMB measurements along the profile 2 – 2’.

L30-31: There is a missing link here. Is the point that if there were basal melting then the Raymond arches would be smaller? As the arch heights have not been compared to arch heights expected from modelled (Martin et al., 2006) or measured (Kingslake et al., 2016) vertical velocities, I am not sure that their size can be used to support the statement about melting. Note that I think it is fine to assume that the ice rise is cold based if the 1D modelling shows that it is thin enough.

Kingslake, J., Martín, C., Arthern, R.J., Corr, H.F. and King, E.C., 2016. Ice flow reorganization in West Antarctica 2.5 kyr ago dated using radar-derived englacial flow velocities. *Geophysical Research Letters*, 43(17), pp.9103-9112.

Yes, this was the point. The presence of arches infer that there has not been any large amount of ice melt below the divide. However, as you pointed out, past small scale melting cannot be ruled out without further analysis. So, as per your suggestion we have removed the statement.

Page 8

The term laminar flow is used at least three times here to distinguish the flow of the ice rise flanks from flow within a region close to the divide where the Raymond effect acts. All glacial ice flow is laminar (as opposed to turbulent), so this seems doesn’t seem like the correct term to use. Perhaps a better way to make the distinction is to say that in the flanks the shallow-ice approximation is valid whereas in the divide region it is not. The approximation of γ in line 7 comes from the SIA and Martin et al. (2009) showed that the full-stokes models are required to describe the Raymond effect and discuss how the SIA is incapable of this. So perhaps this works better.

Martin, C., Hindmarsh, R.C. and Navarro, F.J., 2009. On the effects of divide migration, along-ridge flow, and basal sliding on isochrones near an ice divide. *Journal of Geophysical Research: Earth Surface*, 114(F2).

We agree and have made relevant changes to the paragraphs explaining the range of γ , and cite Martin et al. (2009): For changes please refer the paragraphs (P8L31 to P9L5) of the marked manuscript.

L26-27: Can you expand on the statement that a more accurate determination of γ requires us to know ice flow history? Is this because ice rheology has a memory through ice temperature and crystal fabric?

Yes. We have rephrased the sentence to (P9L19):

“More accurate determination of γ requires the knowledge of ice-rise evolution in the past millennia, because ice rheology has a memory through ice temperature and crystal fabric, and the use of detailed ice-flow modelling, which is beyond the scope of this study.”

Page 9

L1-2: This sentence, starting ‘This is because...’ is very unclear.

We rephrased the sentences:

We divide the ice rise into 19 polygons in respect to the slope direction (Fig. 6a). This is because the SMB has a contrasting feature between southeast and northwest sides (Fig. 4), and the corresponding difference in the mass balance can happen.

With (P9L30):

“As ice rises are expected to show slope dependent SMB features (King (2004), Lenaerts et al., 2014), it is probable that mass balance could also have similar features. To account for this, in this method we divide the ice rise into 19 polygons in respect to the surface slope direction (Fig. 6a) and data availability.”

I found section 6.2.3 very difficult to understand. For example, in line 21, is ‘this estimate’ the estimate of SMB using the flowband setup or the estimate of the variation in flowband width. Also is the second paragraph (which is just one sentence) missing more material? I suggest that the description of the flowband setup is re-written and expanded to make this clearer.

We rephrased the paragraph (P10L17-P10L28) as:

“In this method, we calculate mass balance along ice-flow bands of varying widths in three different slopes of the ice rise. We define a flowband width to account for flow divergence and convergence along the flowband, assuming that ice flows along the steepest descent path on the surface. Flowband width at the downslope end is taken as 1 km, and for each flowband, steepest ascent paths are determined from two points 0.5 km away from the most downstream GPS stake. We used ascent because the surface topography near the summit is much less distinct and consequently the divergence estimate is more sensitive to small topographic changes. We rejected three flow profiles out of six, because the GPS markers were not within the defined flowband. Along the three flowbands shown in Fig 6c, the flowband widths vary by a factor of 1.4–3.6 along each flowband. This variation depends on the initial band width (1 km used here), but over the range of the initial band width between 0.9 and 2.5 km, the band width estimates vary only ~3%. We further divided the flowbands into three columns based on the available data and calculated mass balance similarly to the polygon setup..”

L6: I am confused by this statement that for a given γ the polygon setup gives the largest estimate of mass balance. According to Fig 7, when $\gamma > 0.75$ the flowband setup (dashed line) is higher than the polygon setup (solid line).

Here we are comparing the mass balance values averaged over all columns in a setup for a given γ . To clarify this point, the sentence is revised to (P13L15):

“We averaged mean mass balance values of all ice columns for each setup and for each γ (shown with point symbols in Fig. 7). For a given γ , the polygon setup give the largest estimate, whereas the grid-setup estimate is smaller by 0.02- 0.03 m a⁻¹.”

L7: Suggest remove ‘much’. I am not sure that I would describe 0.05-0.07 m/a as *much* smaller than 0.1 in this context. They are the same order of magnitude.

You are right. We removed “much”.

L8-9: I am not sure that I understand the sentence starting ‘Higher γ ...’. Are you saying that the dependence of the mass balance on γ is nearly linear. If so, I suggest you replace ‘uniform’ with ‘linear’.

The sensitivity of mass balance to γ is represented by the slope of the curves in Figure 7, which are essentially uniform for all columns . The sentence is left unchanged. (P13L20).

L11: Delete ‘as it goes’.

Corrected.

L12: Rephrase this sentence to avoid the phrase ‘varies in a more variable way’.

Sentence rephrased as (P13L23):

“Along slopes C, E and F, mass balance increases monotonically as it goes downstream, whereas mass balance of polygons along A, B and D slopes is more variable.”

Page 11

The discussion contains many typos and ungrammatical sentences, that I have not listed in detail.

We apologize that the manuscript has many typos. We carefully read the manuscript to remove typos and grammatical errors in these sections.

L15-26: Do these two paragraphs belong in the introduction? They do not discuss any of the new results and I think they can be shortened without losing substance.

We think that these paragraphs better fit to Discussion, rather than Introduction. This is because this knowledge (mechanisms to determine SMB) is not required as background information to understand the contents of this paper. Rather, this knowledge is more useful to discuss and interpret our results in a larger context. We have shortened them considerably as (P14L28 – P15L12).

Page 12

L20: Another recent relevant reference is:

Kingslake, J., Martín, C., Arthern, R.J., Corr, H.F. and King, E.C., 2016. Ice-flow reorganization in West Antarctica 2.5 kyr ago dated using radar-derived englacial flow velocities. *Geophysical Research Letters*, 43(17), pp.9103-9112.

Thanks for drawing our attention to this work. We have added this reference at P15L18.

L26: ‘inferring’ → ‘implying’.

Corrected.

L27: ‘(1000s m)’ is confusing as throughout the manuscript spaces between values and units have not been consistent. Suggest replace with ‘1000’s of meters’.

Corrected.

L29: Delete ‘the’ before ‘Raymond’.

Corrected.

L31: Is a word missing from near the beginning of this sentence?

The original sentences

If the summit position is steady for longer time, then the Raymond arches are further developed into double-peaked arches (Martín et al., 2009), which are not clearly observed here.

are changed to (P16L15):

If the summit position stays stable for longer time, then the Raymond arches are further developed into double-peaked arches (Martín et al., 2009), which are not clearly observed here.

Page 13

Section 7.5 contains no discussion of your results, only the location of the ice rise. Can any of your findings (for example bed topography) contribute to this discussion of the impact of the ice rise on the adjacent ice shelves? If not, perhaps this belongs in the introduction or the section in which you introduce the ice rise.

We want to discuss the role of this ice rise in current setting for completion. Discussion seems to be the right place, as the reader is now well informed of the ice rise. To make it fit better we point out how the questions raised in this paragraph will be answered in future modelling study of Blåskimen Island. The new paragraph with the added sentence (underlined) is (P16L20-P16L29):

Roles of ice rises vary largely in terms of its settings, and thus can change during the evolution (Matsuoka et al., 2015). Blåskimen Island is currently situated at the calving front of the local ice shelves and in the ice-flow shadow of Novyy Island ice rise upstream (Fig 1). This setting implies that currently Blåskimen Island alone has limited impact on the continental grounding line and ice flux from the ice sheet. However, it seems likely that Blåskimen Island plays a more significant role than the Novyy Island to maintain the current calving front position. Favier and Pattyn (2015) demonstrated that an ice shelf landward of the ice rise is thicker than the seaward ice shelf facilitating formation of rifts and ice-shelf breakups just seaward of the ice rise. To explore the dynamics of this surrounding region better we will use the datasets presented in this study to model the ice evolution of the ice rise.

L17: Insert 'a' between 'over' and 'flat'.

Corrected.

Figures and captions

Figure 4 caption: The second sentence appears to be missing something.

Replaced.

Circles show the locations of 90 stakes.

With

Circles show the locations of the installed stakes.

Figure 5 caption: The end of second sentence is ungrammatical.

We replaced the sentence:

Dots show the arch amplitudes and dashed lines show their linear fits to the depth.

With

"Dots show the arch amplitudes and dashed lines show the linear fits of the arch amplitudes to depth."

Figure 7 caption, lines 3: Replace 'as' with 'to'.

Corrected.

Responses to reviewer #2

General Comments

This manuscript provides exhaustive details on the field measurements and data analysis over an ice rise in Dronning Maud Land, Antarctica from radar and GPS. While the manuscript provides sufficient data to support their conclusions it falls short of providing a compelling story. I think the main problem may be more due to writing however, than mis-interpretation of the data. Indeed, it seems to be more of a field report than a scientific paper that tells a story. I would prefer for the authors to follow the more traditional style of writing with sections called: introduction, data and methods, results, discussion, conclusions. While these sections are in the text, there are numerous other sections and subsections with detailed headings making it difficult to follow the logical order of the manuscript. Further, the writing could be tightened by combining very short paragraphs and reducing redundant information. For example, instead of presenting info on the ice core records in section 3, simply refer to the reference in your discussion of your measured SMB. Same for temperature info - why is this relevant to the current study? This kind of thing led to my inability to understand what the story is here.

Thank you for your constructive comments. To emphasize more on the story we have reworked the introduction section of the manuscript. We have also reorganized the manuscript to follow a more conventional structure as suggested. Please see the new structure presented at the beginning of our response letter.

We keep descriptions of the 23-m-firn core in the revised manuscript because this core is used to develop depth profiles of density, which is used to derive SMB. Amongst all field measurements we made, only firn temperature is not discussed in this paper. Rather than leaving a single measurement out from this paper, we prefer to include all field measurements together in this paper so that this paper can be a comprehensive reference point when we write following papers including the one currently under development to decipher evolution of this ice rise using numerical modeling.

Technical Comments

Page 3, Line 21: delete “a”

Corrected.

Page 3, Line 30: If you’re using the firn cores to get estimates of density variability it might be nice to include a figure of these data.

Core locations are shown in Fig. 4a (wheel spoke markers).

Page 5, Line 8: This sentence is not needed as it was already said in a previous section.

Agreed. We have removed this sentence.

Page 6, Line 6-7: Don’t forget the work of Nereson and Raymond, 2001

Thank you for this suggestion. We have included this reference (P6L32).

Page 7, Line 7: delete “a”

Deleted.

Page 7, Line 25: Rignot and Kanagaratnam is a better reference since they (I believe) pioneered the IO method.

Thank you for clarifying. We have included the reference you suggested (P8L18).

Page 7, Line 28: Not sure you can assume no melting based on the study by Neumann et al., (2008) which is on the other side of Antarctica.

We cite Neumann et al. (2008) as a reference of the modeling method used here. This is why Neumann et al. (2008) is mentioned immediately after “one-dimensional thermomechanical model”, rather than at the end of the sentence.

Page 11, Lines 3-9: The authors make an offhand assertion that current DEMs of Antarctica do not properly resolve this ice rise elevation, nor the details contained within their DEM. I think they would have to demonstrate that the added detail is necessary in order to “get the precip right” according to

Lenaerts (2014). I might be convinced that gross differences between DEMs are important but I'm not convinced that we need to include every small detail as obtained by a ground survey.

We regret that this paragraph can be read in this way. We intended to say that topography of ice rises is much steeper than the other parts of Antarctica and thus in general ice rises are difficult places to obtain precise elevations using satellite altimetry techniques, though they are most useful to generate continent-wide DEM. With the best of our knowledge, there is no validation study to examine continent-wide DEMs in the coastal Antarctica, focusing to ice rises. It is quite interesting to compare climate model results for different topography to test required accuracy of topography to precisely model SMB, density, etc over the ice rises and ice shelves. However, such modeling work is beyond the scope of our paper. Below, we show a revised paragraph with which we believe that our point is clearer.

“Distinct topographic features of the ice surface revealed with our DEM are not fully represented in continent-wide DEMs (e.g. Bamber et al., 2009; Fretwell et al., 2013, in which spatial resolution is 1 km). Also, the summit heights in those products are 24–40 m lower than our measurements. It remains unclear how much this inaccurate description of topography affects modeling SMB and surface density. Lenaerts et al. (2014) demonstrated that elevated topography associated with ice rises causes orographic precipitations and corresponding precipitation shadow not only over ice rises but also on adjacent ice shelves. Such variations could result in anomalous firn density over the ice shelves, which would result in ill-posed estimate of freeboard thickness and its long-term changes.”

Page 11, Line 31-32: your reported slope differences are within the expected range of the model prediction by Lenaerts et al, (2014) not “smaller than”

Corrected. Rephrased the sentence as (P15L13):

On Blåskimen Island, we found that upwind slopes have 2–3 times the SMB on the downwind slopes, which is within the model prediction (2–4 times; Lenaerts et al. (2014)).

Page 12, Line 19: I don't see the need for these references here - we already know what Raymond arches are.

You are right. The reference has been removed.

Page 12, Line 21: refer to Nereson's work on this

Added Nereson et al. (1998) at (P16L5).

Page 12, Line 30-32: I can clearly see double-peaked arches in Fig. 3c.

We argue that the second arch on the side is made by a small bed undulation there. Our preliminary results of ice-flow modeling show that this arch is reproduced even if ice is assumed isotropic, which will be reported in another paper that we are developing on this manuscript. To clarify this point, we revised the sentences to (P16L15):

“If the summit position is stable for longer time, then Raymond arches are further developed into double-peaked arches (Martín et al., 2009), which are not clearly observed here (small side arch is caused by bed bump nearby, according to our initial ice-flow modeling).”

Below we list the citations not shown properly in the marked manuscript due to an issue with Latexdiff –

P1L18 - (Dupont and Alley, 2005; Matsuoka et al., 2015; Fürst et al., 2016)

P6L32 - (e.g. Vaughan et al., 1999; Conway et al., 1999; Nereson and Raymond, 2001)

P8L18 - (e.g., Rignot and Kanagaratnam, 2006; Conway and Rasmussen, 2009; Zwally and Giovinetto, 2011)

P9L2 - (Martín et al., 2009a, b; Gillet-Chaulet et al., 2011; Drews et al., 2015)

P16L2 - (~~Vaughan et al., 1999, Drews et al., 2015~~)

Glaciological settings and recent mass balance of ~~the~~ Blåskimen Island in Dronning Maud Land, Antarctica

Vikram Goel¹, Joel Brown^{1,2}, and Kenichi Matsuoka¹

¹Norwegian Polar Institute, Tromsø, Norway

²Aesir Consulting LLC, Missoula, Montana, USA

Correspondence to: Vikram Goel (vikram.goel@outlook.com)

Abstract. The Dronning Maud Land coast in East Antarctica has numerous ice rises that very likely control the dynamics and mass balance of this region. However, only a few of these ice rises have been investigated in detail. Here, we report field measurements of Blåskimen Island, an isle-type ice rise adjacent to the Fimbul Ice Shelf. Blåskimen Island is largely dome shaped, with a pronounced ridge extending to the southwest from its summit (410 m a.s.l.). Its bed is mostly flat and about 100 m below the current sea level. Shallow radar-detected isochrones dated with a firn core reveal that the surface mass balance is higher on the southeastern slope than the northwestern slope by $\sim 37\%$, and this pattern has persisted for at least the past decade. ~~Radar stratigraphy shows upward arches underneath the summit, indicating~~ Arches in radar stratigraphy suggest that the summit ~~position has been stable over at least one characteristic time of this ice rise (of the ice rise has been stable for~~ ~ 600 years). Ensemble estimates of the mass balance using the input-output method show that this ice rise has thickened by ~~0.07~~ 0.12–0.35 ~~0.37~~ 0.37 m ice equivalent per year over the past decade.

1 Introduction

Around 74% of the Antarctic coastline consists of floating ice shelves fed by outlet glaciers and ice streams (Bindschadler et al., 2011). These ice shelves together with numerous pinning points (ice rises and rumpled) ~~restrain~~ regulate the outflow of the grounded ice ~~by buttressing the ice upstream (Dupont and Alley, 2005; Matsuoka et al., 2015; Fürst et al., 2016).~~ Therefore, understanding the settings of ice rises and rumpled and their evolution under the changing climate are crucial to better understand the stability of the Antarctic Ice Sheet as well as eventual Antarctic contributions to the sea level.

~~The 2000km long coastline of Dronning Maud land (DML), East Antarctica, consists of numerous outlet glaciers and ice shelves punctuated by some 30 ice rises (Matsuoka et al., 2015). Embedded into an~~ (Dupont and Alley, 2005; Matsuoka et al., 2015; Fürst et al., 2016) Embedded into the ice shelf, ~~an ice rise creates ice rises~~ create a zone of compression upstream of the ice rise which buttresses the ice shelf (Borstad et al., 2013). However, downstream of the ice rise the tensile forces leave a weak region subject to crevasses and thinner ice shelves (Favier and Pattyn, 2015). ~~Therefore, ice rises and rumpled most likely contribute significantly to the shape and mass balance of coastal DML.~~ Ice rises strongly influence regional surface mass balance (SMB) (Lenaerts et al., 2014) and can significantly alter the timing of deglaciation of the ice sheet (Favier and Pattyn, 2015; Favier et al., 2016). Although relatively small in footprint, ice rises can have far-reaching effects on the Antarctic ice sheet dynamics.

Ice rises are also a useful resource for investigating evolution and past climate in the coastal region. Englacial (isochronous) stratigraphy detected using radar has been widely used to constrain the evolution of the ice rise and adjacent ice body over the past millennia (Conway et al., 1999; Nereson and Waddington, 2002; Hindmarsh et al., 2011; Siegert et al., 2013; Drews et al., 2015; Kingslake et al., 2016). The knowledge of the evolution of ~~the-an~~ ice rise is crucial to retrieve reliable past regional climatic changes, using ~~an~~-ice cores drilled through ~~the-ice-rise-it~~ (Mulvaney et al., 2002, 2014).

The 2000 km long coastline of Dronning Maud land (DML, 20°W–45°E), East Antarctica, consists of numerous outlet glaciers and ice shelves punctuated by some 30 ice rises (Matsuoka et al., 2015). These ice rises most likely contribute significantly to the shape and mass balance of this region. In addition, they present an opportunity to better understand the evolution of this data sparse region (Mackintosh et al., 2014). So far, only two ice rises have been investigated in DML: Derwael Ice Rise (26°E) in Roi Baudouin Ice Shelf (Drews et al., 2015) and Halvfarryggen Ice Dome (6°W) between Jelbartisen and Ekstromisen ice shelves (Drews et al., 2013). Both ice rises are grounded on flat bed ~200 m below the current sea level, show large SMB contrast across upwind/downwind slopes and have dynamic characteristic times of hundreds of years. Stratigraphic evidence shows that these ice rises have been in nearly steady state over the last several millennia (3000–5000 years). These ice rises are separated by ~1200 km along the coast, where glaciological settings are variable. For example, along the DML coast between the two ice rises, SMB can vary by at least a factor of two, depending on surface topography, storm tracks and wind direction (King, 2004; Lenaerts et al., 2014). This region also consists of several ice shelves and outlet glaciers with flow speeds varying by a factor of four (Rignot et al., 2011). Ice rises in these varying settings can evolve and impact adjacent ice shelves differently and hence needs to be investigated.

We carried out field measurements of Blåskimen Island, an isle-type ice rise located west of the Fimbul Ice Shelf at the calving front (Fig. 1a), to elucidate the current status and past evolution of this coastal region. Here, we present surface and bed topography, surface flow field, and surface mass balance of Blåskimen Island. Our analysis of these data implies that the ice rise has thickened by ~~0.07~~0.12~~-0.35~~-0.37 m a⁻¹ (ice equivalent) over the past decade.

2 Blåskimen Island

Blåskimen Island has a total area of 651 km² (Fig. 1a; Moholdt and Matsuoka, 2015). It is located between the Fimbul and Jelbart Ice Shelves near the calving front. Both of these ice shelves flow around several isle-type ice rises (isolated from the ice sheet by an ice shelf) and promontory-type ice rises (elongated extension of the ice sheet into an ice shelf). The Fimbul Ice Shelf is mainly fed by the Jutulstraumen Glacier, one of the largest outlet glaciers in DML (Høydal, 1996). The Jelbart Ice Shelf is fed by the Schytt Glacier, which is slower by a factor of two and narrower, than the Jutulstraumen Glacier (Rignot et al., 2011). Blåskimen Island, along with two more ice rises upstream, Novyy Island and Lejtenanta Smidta, separate these two ice shelves. The ice between Blåskimen Island and these ice rises moves very slowly ($\ll 10$ m a⁻¹ Rignot et al., 2011). In addition, there are four smaller ice rises and rumples near the calving front of the western shear margin of the fast-flowing portion of the Fimbul Ice Shelf (Moholdt and Matsuoka, 2015). Thus towards east, Blåskimen Island ~~abuts a part of~~is surrounded by much

slowly moving ice whereas it abuts the eastern shear margin of the fast flowing part of the Jelbart Ice Shelf ~~Whereas towards east, it is surrounded by much slowly moving ice towards west.~~

3 Field measurements and data processing

We carried out field surveys of Blåskimen Island during the austral summers of 2012–2013 and 2013–2014. It included kinematic and static GPS surveys (Section 3.1), shallow- and deep-sounding radar profiling (Section 3.2), as well as firn coring and borehole temperature measurements (Section 3.3). The location of these measurements is shown in Fig. 1b.

3.1 Kinematic and static GPS surveys

To develop digital elevation models (DEM) of the ice rise surface, we conducted kinematic GPS surveys using Trimble dual-frequency receivers. Two units were installed near the ice-rise summit, one acting as a base station, with one for redundancy. Five rover stations were mounted on snowmobiles, which moved at a speed of $\sim 15 \text{ km h}^{-1}$. Our survey resulted in surface elevation measurements at the average interval of $\sim 4 \text{ m}$ along the survey transects spaced 0.8–1 km from each other. Additional surveys were carried out in the vicinity of the summit to precisely determine the summit position of the ice rise and in the eastern part of the ice rise where satellite imagery shows surface lineations (light grey feature over dark grey in Fig. 1b).

To measure the surface-flow field, we installed 3m-3 m long hollow aluminium stakes at 90-97 locations on Blåskimen Island. The stakes were installed $\sim 1 \text{ m}$ into the snowpack without any anchor resisting vertical motion. Out of these, 49-55 stakes were installed along 6 steepest descent paths determined with the surface DEM. The other 41 stakes were installed in the vicinity ($< 2.5 \text{ km}$) of the summit to better resolve small ice-flow speeds there. The stakes were occupied for ~ 20 minutes to determine their lateral positions (e.g., Conway and Rasmussen, 2009; Matsuoka et al., 2012b). The position of each of the stakes ~~were~~ was measured in January 2013. In January 2014 we remeasured the positions of the ~~stakes that were not lost or tilted~~ 90 of these stakes. Six stakes were buried and one found heavily tilted (more than $\sim 20^\circ$). We did not use GPS-measured vertical positions to determine ice-thickness changes, because of possible motion of the stakes relative to the firn and firn densification. Nevertheless, we measured their heights to the snow surface to estimate surface mass balance (SMB) over the year.

Instantaneous kinematic, and average static-rover station locations were determined relative to the base stations for each field season using TRACK software package, part of the GAMIT/GLOBK GPS positioning software (Herring et al., 2010). Base-station positions for each field season were determined using Canadian geodetic Precise Point Positioning system (CSRS-PPP; <https://webapp.geod.nrcan.gc.ca/geod/tools-outils/ppp.php>). These base stations moved negligibly over the 5 days of GPS campaigns each year; less than the error of each GPS location ($\sim 1 \text{ cm}$ laterally). To convert heights above the WGS84 ellipsoid to heights above local sea level, we subtracted 13 m uniformly using a of geoid height uniformly provided by GOCE gravity product (<https://earth.esa.int/web/guest/data-access/browse-data-products/-/article/goce-gravity-fields-5777>).

3.2 Ice-penetrating radar profiling

We collected common-offset radar transects along four of the steepest-decent paths concurrent with the GPS [stake](#) locations. These radar transects were collected with a 2-MHz ground-based radar system with resistively-loaded dipole antennas (Matsuoka et al., 2012a) to reveal the ice thickness and englacial isochronous stratigraphy. We also operated GSSI/SIR3000 radar with 400-MHz antenna to detect stratigraphy within the top ~ 50 m of the ice rise (Hawley et al., 2014). These two radar surveys were collected with snowmobiles moving at $8\text{--}10$ km h⁻¹ towing the antennas. The antenna positions were determined using kinematic GPS attached to the snowmobiles. This resulted in an average radar trace spacing of ~ 5 m for the deep-sounding radar and ~ 0.25 m for the shallow-sounding radar.

Post processing included using a dewow filter, [an](#) Ormsby band-pass filter, and depth-variable gain functions. To calculate ice thickness, we assumed a radio-wave propagation speed of 169 m μs^{-1} and added a firn correction to account for faster propagation in the firn. [We used an optimization inversion routine \(Brown and Matsuoka, in prep\) to model depth profiles of density along shallow radar profiles using \(1\) surface densities measured at 13 locations, and \(2\) depth profile of the density measured along the 23 m long firn core. We then used the depth profiles of the density along the radar profile to estimate laterally variable firn corrections.](#) The magnitudes of the firn correction ~~depend on firn density and~~ vary between $4\text{--}6$ m. To calculate depths of englacial reflectors, we used variable propagation speeds ~~. We discuss spatially variable density, firn correction, and propagation speed in Section 5.2~~ [discussed later in Section 4.1.](#)

3.3 Firn cores and borehole-temperature measurements

We drilled a ~~23-m-long~~ [23 m long](#) firn core near the summit of Blåskimen Island and transported the frozen samples to ~~labs to develop both laboratories to develop~~ visual stratigraphy as well as chemical, isotope, and dielectric ~~profiles~~ [stratigraphy](#). The core was dated back to 1996 by counting annual cycles of oxygen isotopes, and by identifying volcanic horizons using non-sea-salt sulphate data (Vega et al., 2016). They found that SMB in the past 17 years ranges between 0.44 m a⁻¹ in 2004 and 1.32 m a⁻¹ in 2011, giving the mean SMB in this period 0.76 m a⁻¹; in this paper, mass balance and SMB are always shown as meters of ice equivalent.

We installed a thermistor string in the borehole and measured temperature profiles from ~~the~~ surface to 20 m depth. Within 25 hours after the completion of drilling, the temperature became steady with ± 0.1 °C of variation at each depth. At depths between 8 and 12 m the temperature was ~ -16.2 °C.

Separately, we drilled nine 3 m long firn cores ([locations are shown in Fig. 1b](#)) to measure spatial variations of surface density. ~~Firn-core density~~ [Hereafter we call mean density in the top 3 m as surface density. It](#) was determined by measuring core volume and weight.

~~Figure 2a shows surface elevations derived from kinematic GPS surveys using bilinear interpolation. The summit is 410 m a.s.l. and is~~ [Measured surface density through firn cores varies by \$\pm \sim 350\$ m higher than the surface of the ice shelf on the southern side \(ice-shelf elevation is taken from Fretwell et al. \(2013\)\). From the summit, the elevation drops gradually towards the edges of our survey region in all directions, giving a relatively dome-shaped topography to the ice rise. A pronounced ridge](#)

extends from the summit to the southwest, whereas the eastern flank shows locally steep slopes and a flatter basin northeast. The eastern steep slopes and the southwest ridge are consistent with lineations observed in satellite imagery (light gray feature over dark gray in Fig. 1b). Along a profile through the summit (2–2' in Fig. 1b), the absolute surface slope smoothed over 500 m long segments ranges between 0.02 and 0.04, except for the summit vicinity of 1 km where the surface is virtually flat (Fig. 3a).

The measured ice thickness of Blåskimen Island ranges between 374 m (first quartile) and 444 m (third quartile), with a 2.5% over the ice rise, with a mean value of 400 m. The ice becomes thinner gradually in all directions away from the summit. Considering uncertainty associated with digitization of the bed reflector, data sampling and firn correction, the uncertainty in the ice thickness is ± 5 m 453 kg m^{-3} (uncertainty: 3% or 14 kg m^{-3}). However, no distinct pattern in surface density variation was observed in terms of elevation or slope direction. When estimating SMB below, we bilinearly interpolated the surface density.

We derived the bed elevations by subtracting the ice thicknesses from the surface elevations. At locations where radar data are available, the bed elevation is on average 110 m below the current sea level (-110 m a.s.l.), ranging between -68 m a.s.l. and -125 m a.s.l. (first and third quartiles, respectively). The highest point (-22 m a.s.l.) on the bed was observed about 6 km northeast of the summit (along the 4–4' profile, see Fig. 1b). We developed a bed DEM using bilinear interpolation (Fig 2b). The bed of the central part of ice rise is very flat; in this region bed elevations vary only by ~ 50 m within an area of $\sim 100 \text{ km}^2$. Also, individual radar profiles show that this region is smooth (Fig. 3c). This low, flat, and smooth region extends from the summit vicinity towards north and northwest, and constitutes the majority of the ice-rise bed. However, towards the southern end of the survey domain, the bed elevation decreases by ~ 200 m over a horizontal range of ~ 5 km, resulting in a mean bed slope of ~ 0.04 . Another steep bed (0.03–0.04) is found in the northeastern slope. These steeper regions in the bed are associated with steeper regions on the surface, although the surface is not as steep as the bed.

The surface-flow field is shown in Fig. 2c. The GPS stakes within 2 km of the summit moved only negligibly ($< 0.1 \text{ m a}^{-1}$). The displacement of stakes outside the summit region is larger and increases downstream; ice flows less than 3 m a^{-1} within 4 km from the summit and 10 – 15 m a^{-1} downslope. Ice flows slowest along the ridge towards southwest, whereas the fastest flow is along the south section of flowline 2–2'. The estimated positions have a mean precision of 4.9 cm and 5.1 cm for the east/west and north/south components leading to a processing uncertainty on the velocity of $\pm 7 \text{ cm a}^{-1}$. This does not include uncertainty associated with any tilt of the stakes. Nevertheless, as the velocities outside the summit area range between 4 m a^{-1} and 15 m a^{-1} , and the observed tilts were small, we consider this uncertainty to be negligible.

4 Analytical methods

4.1 Surface mass balance

We estimated surface mass balance (SMB) using two different methods. The first method uses the heights of GPS stakes above the snow surface. ~~With this method, we assume that relative motion of the stakes into the snow is negligible and thus the height differences over the two occupations are caused only by SMB.~~ The second method for determining SMB uses isochronous

radar reflectors profiled with the shallow-sounding radar. Both methods require ~~firm-density~~ surface-density distribution measurements.

~~We collected 3-m-long cores at 9 locations to measure firm density variations (locations are shown in Fig. 4a). Measured firm density varies by $\pm \sim 2.5\%$ over the ice rise, with a mean value of 453 kg m^{-3} (uncertainty: 3% or 14 kg m^{-3}). However, no distinct pattern in firm density variation was observed in terms of elevation or slope direction. While estimating SMB below, we bilinearly interpolated the firm density.~~

We estimated SMB at each stake by multiplying the measured stake-height differences by the measured surface density (Fig. 4a). ~~Using this method, we find that the mean SMB for Blåskimen Island is 0.78 m a^{-1} . $723 \text{ kg m}^{-2} \text{ a}^{-1}$~~ Considering measurement errors and uncertainty accounting snow densification (Eisen et al., 2008), this estimate has an overall uncertainty of $\pm 6\%$. We consider the sinking of the stake under its own weight to be minimal, as the observed surface densities are high. ~~SMB varies by a factor of 3.3 over the study area of $\sim 20 \text{ km}$ by $\sim 20 \text{ km}$ ($0.28\text{--}1.03 \text{ m a}^{-1}$), $\text{kg m}^{-2} \text{ a}^{-1}$ with the first and third quartiles of 0.73 and $0.87 \text{ m a}^{-1} \text{ kg m}^{-2} \text{ a}^{-1}$ respectively. The SMB shows a distinct spatial pattern; it is larger in the southern slope and smaller in the northern slope. The surface is rougher (sastrugi) in the low SMB region whereas smoother and softer in the high SMB region.~~

~~The summit vicinity has a large number of stakes, which show small variations of SMB without any distinct pattern. We compared the SMB values to local maximum surface slope (not necessarily along the prevailing wind direction) but found no clear relationship between them.~~

~~The shallow-sounding radar visualizes continuous reflectors within the firm (Fig. 3b). No major disruptions are observed that can be associated with surface melt or strong wind scour. Therefore, we assume these continuous reflectors are isochrones (Richardson et al., 1997). With this assumption, we can associate firm core ages to radar reflectors. We tracked three reflectors to almost the full extent of the radar surveys; at the 23m-long core site, they are at 8.4 m (actual depth or 2.15 m ice-equivalent depth; dated 2011 or 3 years before the survey by Vega et al. (2016)), 11.9 m (4.2 m; 2009), and 12.8 m (6.9 m; 2005).~~

~~To derive the SMB~~ Provided that radar reflectors are isochronous (Richardson et al., 1997), SMB can be derived from dated radar reflectors. In this analysis, we assume that the effects of vertical strain (thinning after the deposition of snow) on reflector depths are negligible so that thickness of an ice layer bounded by the radar reflectors are solely controlled by the differences in SMB (Waddington et al., 2007). ~~This shallow-layer~~ Thus shallow-ice approximation can hold, when depth h (ice equivalent) of a radar reflector is much smaller than the local ice thickness H ($h \ll H$). For our case, h/H is less than 0.04 and thus the ~~shallow-layer~~ shallow-ice approximation is in most cases valid.

However, ~~this assumption~~ the shallow-ice approximation may not be valid in areas where vertical strain ~~varies over a short horizontal distance~~ rates are large, such as the region near an ice-flow divide (Gillet-Chaulet et al., 2011; Kingslake et al., 2014). In this region, accumulated effects of variable vertical strain can result in upward arches in isochrones, so-called Raymond arches (Raymond, 1983), which were found at many other ice rises (e.g. Vaughan et al., 1999; Conway et al., 1999) (e.g. Vaughan et al., 1999). Such upward arches can also be caused by anomalously low SMB near the summit possibly due to wind erosion (Drews et al., 2013, 2015).

Vaughan et al. (1999) demonstrated that amplitudes of upward arches induced by anomalous SMB increases linearly with ice-equivalent depth, whereas the Raymond effect makes the amplitude increase quadratically. We used this criterion to diagnose the origin of shallow upward arches near the current summit (Fig. 3b2b). We first derived ice-equivalent depths of reflectors, assuming that firm density does not vary laterally. For this purpose, we used the depth profile of density $\rho(z)$ at the core site.

5 We estimated local propagation speed $v(z)$ at a depth z using a relationship between density and refraction index $n(z)$ (Kovacs et al., 1995):

$$v(z) = \frac{c}{n(z)} = \frac{c}{(1 + 0.851\rho(z)) \frac{c}{1 + 0.851\rho(z)}} \quad (1)$$

Here, c is the propagation speed of light in the vacuum ($300 \text{ m } \mu\text{s}^{-1}$). Then, we estimated two-way propagation time $t(z)$ to each depth z :

$$10 \quad t(z) = \int_0^z \frac{2z}{v(z)} \frac{2}{v(z)} dz \quad (2)$$

Second, using these ice-equivalent depths z , we measured arch amplitudes from the arch top to the baseline defined with reflector depths 1 km away to both sides of the arch.

~~We measured arch amplitudes of two deeper reflectors of three that we used for SMB estimates, as the arch amplitude for the shallowest reflector is insignificant. In addition, we measured arch amplitudes of six more reflectors at greater depths. These reflectors range between ~ 4 m and ~ 35 m ice-equivalent depth (Fig. 5). We analyzed the arch amplitudes in this depth range to better resolve their depth variations. All four radar profiles across the summit (Fig. 1b) show that the arch amplitude increases linearly with depth. Therefore, we conclude that the shallow-layer approximation can be used all along the radar profiles and thus the three radar reflectors within the top ~ 7 m ice-equivalent represents spatial patterns of SMB.~~

~~To precisely calculate SMB using the radar reflectors, we need to consider density variations. The 23m long core shows that the density varies $\sim 36\%$ (450 – 655 kg m^{-3}) vertically along its length and the 9 shallow cores show that the surface density varies $\pm \sim 2.5\%$ horizontally. Amongst three reflectors we analysed, the deepest reflector (12.8 m actual depth at the core site) has the largest depth range between ~ 8 and ~ 15 m. It implies that the use of a uniform density could ill-pose the SMB estimates, and variable density should be accounted for.~~

~~To account density variations, we used~~ Under the shallow-ice approximation, we accounted for density variations using two methods. The first method is to account only for vertical variations in density and ignore its lateral variations. For this purpose, we used the measured densities of the ~~23m~~ 23 m long core. The second method is to account for spatial variations in density, temperature, and SMB altogether using an inversion method (Brown and Matsuoka, in prep). The former method is not strictly valid, because the surface density varies by $\pm \sim 2.5\%$ and possible variations in SMB add more complexities in density at depths away from the core site. The latter method is also not strictly valid as it solves for the best fit at all locations to a steady-state firm densification model. Firm-core analysis shows no significant temporal trend but large year-to-year variations by a factor of three over the past 17 years (Vega et al., 2016). Nevertheless, the model fits the measured density well (within

95% confidence bounds of the fit). Although these two methods use distinct assumptions, we show that they give consistent results and thus a more confidence in SMB estimates using inaccurate data representations in these two methods.

Factors determining the uncertainty in SMB derived from dated radar isochrones can be broadly categorized as: (1) error in determining the depth of the reflector, (2) error in dating the firn core, (3) error in estimating the cumulative mass above the reflector and its spatial variability. For the first source we assess the uncertainty to be within ± 10 cm. We consider the combined errors in depth-age scale and error in linking it to radar reflector to be ± 1 year. The density model used to fit the observations has an uncertainty of $\pm 3\%$, whereas we see $\pm 3\%$ variability in the surface density. Using standard error propagation, this results in an uncertainty of $\pm 11\%$.

Figure 4b shows the SMB averaged over the past 9 years (2005–2014) estimated using the second radar method. It gives the spatially mean value of $0.75 \text{ m a}^{-1}, 692 \text{ kg m}^{-2} \text{ a}^{-1}$ with the first and third quartiles of 0.67 and $0.85 \text{ m a}^{-1}, 611$ and $784 \text{ kg m}^{-2} \text{ a}^{-1}$ respectively. The first radar method gives a slightly higher number than the second radar method by 5–10%; the mean value for the first method is $0.81 \text{ m a}^{-1}, 744 \text{ kg m}^{-2} \text{ a}^{-1}$ with first and third quartiles of 0.71 and $0.93 \text{ m a}^{-1}, 650$ and $860 \text{ kg m}^{-2} \text{ a}^{-1}$. The difference between the SMB estimated with these two methods is localized to the region in 1–6 km northeast of the summit. Except for this northeast region, these two methods give nearly identical spatial patterns in SMB.

Using the data presented in the previous sections, we estimate mass balance of Blåskimen Island for the past nine years, the longest period for which our SMB estimates are available.

4.2 Mass balance of the ice rise

We applied the Input–Output (I–O) Method (e.g., Conway and Rasmussen, 2009; Zwally and Giovinetto, 2011) (e.g., Rignot and Kanagaratnam, 2015) to individual columns over the ice rise (Section 6.2.4.2.1). The I–O Method calculates the mass balance as a difference between incoming Q_{in} and outgoing Q_{out} fluxes from all the sides of a column, with SMB (M_{SMB}) over the column area added. Here, we ignore basal melting, as one-dimensional thermomechanical model (Neumann et al., 2008) shows no basal melt for the geothermal flux estimated in this region (57 mW m^{-2} ; Maule et al., 2005). Radargrams show no anomalous features in the radar reflection from the bed that could indicate basal melting (Fig. 3e). Also, basal melting diminishes Raymond effect, which is often the cause of upward arches at large depths near the divide (?). 2c).

The ice-flow fluxes through an ice column are calculated as:

$$Q = \rho \gamma u_{\perp} h \quad (3)$$

Here, ρ is the density of a column, γ is a dimensionless factor which scales the measured surface flow speed u_{\perp} normal to the gate to depth averaged speed $u_{av\perp}$; $u_{av\perp} = \gamma u_{\perp}$, with $\gamma \leq 1$. This implicitly assumes that ice-flow direction does not change with depth, which seems valid over relatively flat bed terrain underneath the ice rise (Fig. 2e Section 5.1).

4.2.1 Constraining γ

The parameter γ is a function of local surface slope, ice thickness, ice temperature and ice rheology (Cuffey and Paterson, 2010). For isothermal, laminar ice flow over non-sliding bed, it can be approximated and assuming shallow-ice approximation

γ can be stated as $\gamma = \frac{(n+1)}{(n+2)}$, where n is the creep exponent of Glen's flow law (Cuffey and Paterson, 2010, p.310). Previous studies on ice-flow divides suggest that n can be between 3 and 5 (Martín et al., 2009a; Gillet-Chaulet et al., 2011; Drews et al., 2015) (Martín et al., 2009a; Gillet-Chaulet et al., 2011; Drews et al., 2015). Then γ ranges between 0.8 ($n = 3$), and 0.86 ($n = 5$).

The actual range of γ can be different, because the isothermal approximation is hardly valid. Also, near the ice-flow divide the laminar-flow shallow-ice approximation is invalid as longitudinal stresses are significant (Raymond, 1983).

Reeh (1988) showed that, near the flow divide, γ can be close to 0.5, when $n = 3$ and ice is isothermal. As ice becomes warmer at greater depths, the deeper ice is presumably softer than the shallower ice, implying that γ is larger when ice-temperature variations are considered. Hence, it is reasonable to assume that $0.5 \leq \gamma \leq 1$. Raymond (1983) used an isothermal model, and Hvidberg (1996) used a thermomechanical model to constrain the range of γ near the divide. Both showed that γ is smallest at the divide, and varies largely near the divide region.

~~As it transitions to the laminar-flow regime beyond~~ Out of the divide region several ice thicknesses from the divide, γ becomes less variable. For isothermal two-dimensional (divide) flow, $0.61 \leq \gamma \leq 0.75$ ($< 8H$ from the divide, Raymond (1983)) and $0.56 \leq \gamma \leq 0.77$ ($< 10H$ from the divide, Hvidberg (1996)). For thermo-mechanical case, $0.69 \leq \gamma \leq 0.86$ ($< 10H$ from the divide, Hvidberg (1996)) and for isothermal axisymmetric radial flow $0.54 \leq \gamma \leq 0.76$ ($< 10H$ from the divide, Hvidberg (1996)). Although radial flow leads to an increase in the divide region, $\sim 70\%$ of the changes in γ still happen within 4 ice thicknesses from the divide. Therefore, for the setups discussed below (with an average extent of ~ 9 km or 18 ice thicknesses from the divide), γ remains virtually uniform.

In the following analysis study, we consider the spatially uniform γ in each estimate and examine ensemble results for within the plausible range (0.7–0.9). More accurate determination of γ requires the full-knowledge of ice-rise evolution in the past millennia and, because ice rheology has a memory through ice temperature and crystal fabric. This requires detailed ice-flow modelling, which is beyond the scope of this study.

4.2.2 Estimate setups

We estimated mass balance for three different setups. (Fig. 6): (i) polygons bounded by GPS stakes, (ii) uniformly-distributed square columns (grid) on the ice rise and average them over individual polygons for (i), and (iii) several flowbands along GPS stakes and radar profiles. The polygon and grid setups have good spatial coverages, but they require data interpolations to large degrees. The flowband setup relies more on direct field measurements, but has very limited spatial coverage. Therefore, we use these three mass balance estimates as an ensemble.

We

4.2.2.1 Polygon setup

As ice rises are expected to show slope dependent SMB features (King, 2004; Lenaerts et al., 2014), it is probable that mass balance could also have similar features. To account for this, in this method we divide the ice rise into 19 polygons in respect to the slope direction (Fig. 6a). ~~This is because the SMB has a contrasting feature between southeast and northwest sides (Fig. 4),~~

~~and the corresponding difference in the mass balance can happen.~~ surface slope direction and data availability. All four sides of these polygons act as a flux gate. Ice thickness and surface-flow velocities are available at each corner of these polygons. We observe some cases that ice is thicker at one corner but ice flows faster at the other corner. Therefore, to better account for these variations along the gate, we divided each gate into ten subgates and estimated the flux at each subgate, rather than calculating
5 mass flux using single values averaged over an entire gate.

4.2.2.2 Grid setup

We divide the ice rise into grid with ~~200m~~ 200 m long square columns and estimated mass balance for each column. The mass balance values are then averaged over each polygon used in the polygon setup. We adopt this setup, together with the polygon setup, to test how data interpolations affect the mass balance estimate. We calculated mass balance MB using the continuity
10 equation to each grid element:

$$MB = \frac{\partial h}{\partial t} = M_{SMB} + \left(\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y}\right)H + \left(u\frac{\partial h}{\partial x} + v\frac{\partial h}{\partial y}\right) \quad (4)$$

Where u and v are the components of ice-flow vectors in the rectangular (local coordinate) directions x and y , and h is the variable for ice thickness. We bilinearly interpolate measured u , v , M_{SMB} , and H into grids.

~~This setup takes much less data interpolation than the previous two setups, and is more based on data collected along
15 flowlines. To define divergence of flow, we tracked the steepest ascent-~~

4.2.2.3 Flowband setup

In this method, we calculate mass balance along ice-flow bands of varying widths in three different slopes of the ice rise. We
define a flowband width to account for flow divergence and convergence along the flowband, assuming that ice flows along
the steepest descent path on the surface. Flowband width at the downslope end is taken as 1 km, and for each flowband,
20 steepest ascent paths are determined from two points 0.5 km away from the ~~GPS stake most downstream of each flowline (i.
e. the flowband width is 1 km).~~ most downstream GPS stake. We used ascent because the surface topography near the summit
is much less distinct and consequently the divergence estimate is more sensitive to small topographic changes. ~~Along three
flowbands shown in Fig. 6c, the band~~ We rejected three flow profiles out of six, because the GPS markers were not within
the defined flowband. Along the three flowbands, the flowband widths vary by a factor of 1.4–3.6. This estimate variation
25 depends on the initial band width (1 km used here), but over the range of the initial band width between 0.9 and 2.5 km, the
band width estimates vary only ~3%. ~~The other flowlines are excluded from further analysis, because GPS-measured velocity
directions are not aligned along the steepest path.~~ %. We further divided the flowbands into three columns based on the available
data and calculated mass balance similarly to the polygon setup.

5 Results

5.1 Topography and surface-flow field

Figure 3a shows surface elevations derived from kinematic GPS surveys using bilinear interpolation. The summit is 410 m a.s.l. and is ~ 350 m higher than the surface of the ice shelf on the southern side (ice-shelf elevation is taken from Fretwell et al. (2013)). From the summit, the elevation drops gradually towards the edges of our survey region in all directions, giving a relatively dome-shaped topography to the ice rise. A pronounced ridge extends from the summit to the southwest. The eastern flank shows locally steep slopes and a basin in northeast with overall lower, less-tilted surface. The eastern steep slopes and the southwest ridge are consistent with lineations observed in satellite imagery (light gray feature over dark gray in Fig. 1b). Along a profile through the summit (2–2' in Fig. 1b), the absolute surface slope smoothed over 500 m long segments ranges between 0.02 and 0.04, except for the summit vicinity of ~ 1 km where the surface is virtually flat (Fig. 2a).

~~We define three columns along each flowline from the summit~~ The radar profiles visualized the bed as well as ice stratigraphy (Figs. 2b and ~~calculated mass balance similarly to the polygon setup~~ 2c). The measured ice thickness of Blåskimen Island ranges between 374 m (first quartile) and 444 m (third quartile), with a mean value of 400 m. The ice becomes thinner gradually in all directions away from the summit. Considering uncertainty associated with digitization of the bed reflector, data sampling and firn correction, the uncertainty in the ice thickness is ± 5 m.

~~Using~~ We derived the bed elevations by subtracting the ice thicknesses from the surface elevations. At locations where radar data are available, the bed elevation is on average 110 m below the current sea level (-110 m a.s.l.), ranging between -68 m a.s.l. and -125 m a.s.l. (first and third quartiles, respectively). The highest point (-22 m a.s.l.) on the bed was observed about 6 km northeast of the summit (along the 4–4' profile, see Fig. 1b). We developed a bed DEM using bilinear interpolation (Fig. 3b). The bed of the central part of ice rise is very flat; in this region bed elevations vary only by ~ 50 m within an area of ~ 100 km². Also, individual radar profiles show that this region is smooth (Fig. 2c). This low, flat, and smooth region extends from the summit vicinity towards north and northwest, and constitutes the majority of the ice-rise bed. However, towards the southern end of the survey domain, the bed elevation decreases by ~ 200 m over a horizontal range of ~ 5 km, resulting in a mean bed slope of ~ 0.04 . Another steep bed (0.03–0.04) is found in the northeastern slope. These steeper regions in the bed are associated with steeper regions on the surface, although the surface is not as steep as the bed.

The surface-flow field is shown in Fig. 3c. The GPS stakes within 2 km of the summit moved only negligibly (< 0.1 m a⁻¹). The displacement of stakes outside the summit region is larger and increases downstream; ice flows less than 3 m a⁻¹ within 4 km from the summit and 10–15 m a⁻¹ downslope. Ice flows slowest along the ridge towards southwest, whereas the fastest flow is along the south section of flowline 2–2'. The estimated positions have a mean precision of 4.9 cm and 5.1 cm for the east/west and north/south components leading to a processing uncertainty on the velocity of ± 7 cm a⁻¹. This does not include uncertainty associated with any tilt of the stakes. Nevertheless, as the velocities outside the summit area range between 4 m a⁻¹ and 15 m a⁻¹, and the observed tilts were small, we consider this uncertainty to be negligible.

5.2 Surface mass balance

The mean SMB from 90 stake-height measurements across Blåskimen Island is 0.78 m a^{-1} , for the period between January 2013 and January 2014 (Fig. 4a). SMB varies by a factor of 3.3 over the study area of $\sim 20 \text{ km}$ by $\sim 20 \text{ km}$ ($0.28\text{--}1.03 \text{ m a}^{-1}$), with 80% of values ranging between 0.69 and 1.03 m a^{-1} . The SMB shows a distinct spatial pattern; it is larger in the southern slope and smaller in the northern slope. The surface is rougher (sastrugi) in the low SMB region whereas smoother and softer in the high SMB region, indicating strong impact of the wind. The summit vicinity has a large number of stakes, which show small variations of SMB without any distinct pattern.

The shallow-sounding radar visualizes continuous reflectors within the firn (Fig. 2b). No major disruptions are observed that can be associated with surface melt or strong wind scour. Therefore, we assume these continuous reflectors are isochrones (Richardson et al., 1997). With this assumption, we can associate firn-core ages to radar reflectors. We tracked three reflectors to almost the full extent of the radar surveys; at the 23 m long core site, they are at 8.4 m (actual depth or 2.15 m ice-equivalent depth; dated 2011 or 3 years before the survey by Vega et al. (2016)), 11.9 m (4.2 m; 2009), and 12.8 m (6.9 m; 2005).

To judge whether the reflector depths are controlled by SMB or Raymond effect (Section 4.1), we measured amplitudes of arches near the current ice-flow divide. This analysis was made for two deeper reflectors of three that we used for SMB estimates, as the arch amplitude for the shallowest reflector is insignificant. In addition, we measured arch amplitudes of six more reflectors at greater depths. These reflectors range between $\sim 4 \text{ m}$ and $\sim 35 \text{ m}$ ice-equivalent depth (Fig. 2b). We analyzed the arch amplitudes in this depth range to better resolve their depth variations. All four radar profiles across the summit (Fig. 1b) show that the arch amplitude increases linearly with depth (Fig. 5). Therefore, we conclude that the shallow-ice approximation can be used all along the radar profiles and thus the three radar reflectors within the top $\sim 7 \text{ m}$ ice-equivalent represents spatial patterns of SMB.

The 23 m long core shows that the density varies $\sim 36\%$ ($450\text{--}655 \text{ kg m}^{-3}$) vertically along its length and the 13 shallow cores show that the surface density varies $\pm \sim 2.5\%$ horizontally. Amongst three reflectors we analysed, the deepest reflector (12.8 m actual depth at the core site) has the largest depth range between ~ 8 and $\sim 15 \text{ m}$. It implies that the use of a uniform density could make the SMB estimates less accurate, and variable density should be accounted for.

Figure 2a shows the SMB averaged over the past 9 years (2005–2014) estimated using the two radar methods along the profile 2–2'. The differences between the SMB estimated with these two methods are localized to the region in 1–6 km north (profile 2–2', Fig. 2a) and northeast (along profile 4–4') of the summit. Except for this region, these two methods give nearly identical SMB spatial patterns along the radar profiles. Figure 4b shows the SMB estimated using the first radar method (accounting for the vertical density variations only). It gives the spatially mean value of 0.81 m a^{-1} , with the first and third quartiles of 0.71 and 0.93 m a^{-1} respectively. The second radar method accounting for both lateral and vertical density variations gives a slightly lower number than the second radar method by 5–10%; the mean value for the second radar method is 0.75 m a^{-1} , with first and third quartiles of 0.65 and 0.85 m a^{-1} respectively.

Factors determining the uncertainty in SMB derived from dated radar isochrones can be broadly categorized as: (1) error in determining the depth of the reflector, (2) error in dating the firn core, (3) error in estimating the cumulative mass above the

reflector and its spatial variability. For the first source we assess the uncertainty to be within ± 10 cm. We consider the combined errors in depth-age scale and error in linking it to radar reflector to be ± 1 year. The density model used to fit the observations has an uncertainty of $\pm 3\%$, whereas we see $\pm 3\%$ variability in the surface density. Using standard error propagation, this results in an uncertainty of $\pm 11\%$. It is larger than an uncertainty of $\pm 6\%$ for the stake method.

5 5.3 Mass balance

We estimated mass balance over the nine-year period between 2005 and 2014 (Fig. 6). The flowband setup shows a mean mass balance of $+0.12 \pm 0.10$ and $+0.27 \pm 0.10 \text{ m a}^{-1}$ over the range of γ . The uncertainty ($\pm 0.10 \text{ m a}^{-1}$) is estimated using the uncertainties in ice thickness (± 5 m), flow speed ($\pm 7 \text{ cm a}^{-1}$), ice density after correcting for firm ($\pm 2\%$) and SMB ($\pm 11\%$) with propagation of errors, we found the overall uncertainty for the mass balance of the flowband estimate at each column to be $\pm 0.1 \text{ m a}^{-1}$. For the other, the polygon and grid setups give very similar spatial patterns to each other. For these two setups, in addition to these measurement errors the measurements errors above, the mass balance estimate is largely affected by errors associated with data interpolations. Because it is difficult to accurately determine the net uncertainties associated with data interpolations, we used two setups (polygon and grid) with distinct data interpolations and consider the differences between them as a guide of representative uncertainties associated with the data interpolations.

We estimated mass balance over the nine-year period between 2005 and 2014. The flowband setup shows a We averaged mean mass balance of $+0.07 \pm 0.10$ and $+0.23 \pm 0.10 \text{ m a}^{-1}$ over the range of values of all ice columns for each setup and for each γ . The polygon and grid setups give very similar spatial patterns (Fig. 6) (shown with point symbols in Fig. 7). For a given γ , the polygon setup gives the largest overall estimate (Fig. 7) give the largest estimate, whereas the grid-setup estimate is smaller by $0.05-0.07-0.02-0.03 \text{ m a}^{-1}$. Because this difference is smaller than the uncertainty for the flowband-setup estimate ($\pm 0.1 \text{ m a}^{-1}$), we argue that the interpolation errors are not dominating these results. Higher γ values correspond to lower mass balance, and the sensitivity of mass balance to γ is nearly uniform for individual columns.

All the polygons show positive mass balance over the full γ range, except for southernmost downstream polygon A3 (the slope-direction codes are shown in Fig. 6a). Along slopes C, E and F, mass balance increases monotonically as it goes downstream, whereas mass balance of polygons along A, B and D slopes varies in a more variable way is more variable. Adjacent southern slopes A and F show contrasting features; the slope A shows the least thickening, whereas the slope F shows the most thickening. For the flowband setup, one column has negative mass balance in the northwest downstream, where the estimated flow divergence is anomalously large.

In conclusion, overall, Blåskimen Island has positive mass balance in the past nine years. Thickening rates vary depending on the setups and the choice of γ . Over the range of γ used here ($0.7 \leq \gamma \leq 0.9$), the mean mass balance varies between $+0.22-0.25$ and $+0.35-0.37 \text{ m a}^{-1}$ for the polygon setup, between $+0.15-0.21$ and $+0.29-0.35 \text{ m a}^{-1}$ for the grid setup, and between $+0.07-0.12 \pm 0.10$ and $+0.23-0.27 \pm 0.10 \text{ m a}^{-1}$ for the flowband setup. Outside the divide region, gamma tend to be higher (0.8–0.9; Section 6.1.4.2); consequently, the thickening rates lean towards the lower end of the estimate above.

6 Discussion

6.1 Topographic characteristics

According to a recent inventory of ice rises and rumples (Moholdt and Matsuoka, 2015; Matsuoka et al., 2015), Blåskimen Island (651 km²) is larger than 91% of the isle-type ice rises (mean: 151 km²). Its summit is 410 m a.s.l., which is higher than 89% of the others (mean: 168 m a.s.l.). Maximum measured ice flow speed (15 m a⁻¹) is above the mean of maximum ice flow speed (13 m a⁻¹) for isle-type ice rises. Also, the mean bed elevation at -110 m a.s.l. is higher than 86% of the others (-178 m a.s.l.). Overall, Blåskimen Island is one of the larger isle-type ice rises.

Detailed surface DEM developed with the kinematic GPS survey (Fig. 2a3a) revealed many distinct topographic features including the southwestern ridge and steep slopes in the east (Section 4). In addition, it confirms that lineations in satellite imagery are associated with surface undulations (Goodwin and Vaughan, 1995). The lineations appear where the surface slope varies largely and most of them are associated with uneven bed topography. This supports the use of satellite imagery as a mean to explore first-order surface and bed topography, when an ice rise remains un-surveyed.

Distinct topographic features of the ice surface revealed with our DEM are not well-fully represented in continent-wide DEMs (e.g. Bamber et al., 2009; Fretwell et al., 2013, in which spatial resolution is 1 km). Also, the summit heights in those products are 24–40 m lower than our measurements. These discrepancies highlight the need of more accurate DEMs in the coastal Antarctica. Elevated-It remains unclear how much this inaccurate description of topography affects modeling SMB and surface density. Lenaerts et al. (2014) demonstrated that elevated topography associated with ice rises causes orographic precipitations and corresponding precipitation shadow not only over ice rises but also on adjacent ice shelves (Lenaerts et al., 2014). Such variations could result in anomalous firn density over the ice shelves, which would result in ill-posed estimates of free-board thickness and its long-term changes of adjacent ice shelves.

6.2 Surface mass balance

We found good agreement in the spatial patterns of stake-measured SMB between 2013 and 2014, and radar-measured SMB between 2005 and 2014 (Figs. 4a and 4b). Relative thicknesses of three layers bounded by radar isochrones (and the surface) vary similarly along the profiles (Fig. 3b2b), inferring that spatial patterns of SMB have remained similar over this 9 year period, 2005 – 2014. SMB averaged over this period vary largely along the radar profiles between 0.66–0.71 m a⁻¹ 611 kg m⁻² a⁻¹ (0.66 m i.e. a⁻¹), (first quartile) and 0.85–0.93 m a⁻¹ 784 kg m⁻² a⁻¹ (0.85 m i.e. a⁻¹), (third quartile), with its mean of 0.75–0.81 m a⁻¹. 692 kg m⁻² a⁻¹ (0.75 m i.e. a⁻¹)

Large spatial variability in SMB was found on other ice rises as well. King (2004) showed that SMB on the Lydden Ice Rise, Brunt Ice Shelf, is highly variable both at large (10s-tens of kilometers) and small spatial scales (100s-hundreds of meters). They demonstrated that large-scale variations are a result of orographic precipitation, whereas small-scale variations are a result of snow redistribution. On both scales, the contribution from sublimation was found to be relatively small. They highlighted

that SMB is sensitive to surface topography; small variations in surface topography cause small changes in wind speed, which could result in large SMB variations, due to the nonlinear relationship between wind speed and snow transport.

~~Ice rises in DML receive most of their precipitation during sporadic synoptic events occurring 0–5 times per year (?). In such events, moisture-rich air from the Southern Ocean approaches the ice rises from a northeasterly direction. The orographic lifting of air on the ice rise leads to precipitation on the northeast slope and a precipitation shadow on the southwesterly slope. During no-precipitation periods, this region is dominated with southeasterly katabatic wind (Lenaerts et al., 2014). Consequently, the observed SMB distribution is a result of precipitation events with the northeasterly wind and erosion/redistribution with the southeasterly wind.~~

Net impact of these mechanisms on SMB in the DML coast was examined using RACMO2 regional climate model (Lenaerts et al., 2014). Among all the ice rises included in their study, SMB varies by a factor of 2–6 between ice rises. ~~Within an ice rise, SMB is higher in the upwind slopes than downwind slopes by a factor of 2–4. This feature is subject to ice-rise topography with regard to the prevailing wind direction, resulting in large SMB variability between ice rises.~~

On Blåskimen Island, we found that upwind slopes have 2–3 times the SMB on the downwind slopes, which is ~~slightly smaller than~~ within the model prediction (2–4 times; Lenaerts et al. (2014)). Similar SMB gradients are found over other ice rises in DML: 2–3 times over the Halvfarryggen Ice Dome (Drews et al., 2013) and about 2 times over the Derwael Ice Rise (Drews et al., 2015). We compared the SMB values to local maximum surface slope (not necessarily along the prevailing wind direction) but found no clear relationship between them. We observed numerous sastrugi and harder snow on the northwestern downwind slopes, a clear indication of snow erosion/redistribution processes.

~~In addition to these large-scale variations, we observed small-scale SMB variations at several locations. One such feature is likely associated with steeper slopes (e.g. -5 km and +8 km in Fig. 3a).~~ Another feature is found near the summit where the surface is virtually flat; here, SMB is lowered by 10% over 0.5 km. Similar low SMB near the summit has been observed in the Halvfarryggen Ice Dome and the Derwael Ice Rise as well, which ~~was~~ were attributed to wind erosion (Drews et al., 2013, 2015).

6.3 Present-day mass balance

Amongst all factors that affect the mass balance, ice-flow speed varies most widely (57%), as compared to changes in SMB (17%) and ice thickness (11%) about their mean values. Consequently, the mass balance distribution is more sensitive to the flow-speed distribution than the other factors. This also explains the low mass balance in the A slope ~~(Fig. 6).~~ Downstream region of the A slope has a lower bed than the central flat basin. As the ice surface is steeper, ice flows faster in this region. Overall, despite the large upwind/downwind contrast in SMB (Fig. 4), differential mass flux compensates for the difference in SMB so that no distinct mass balance patterns were found (Fig. 6). Our mass balance estimates show that Blåskimen Island is thickening almost everywhere, but the thickening rate is smaller near the summit than the flank. If this pattern is persistent for a long period, it would result in flattening of the ice rise initially ~~but~~. But ice-flow fields would probably be adjusted to the new topography, so the net impact of the ongoing differential thickening on ice topography remains unknown.

6.4 Long-term evolution and impact on the adjacent ice shelves

Distinct upward arches ~~of in~~ the ice stratigraphy ~~at least~~ up to ~ 40 m depth are caused by low SMB near the summit (~~Fig. 5~~). ~~There are.~~ We observed upward arches below this depth (Fig. ~~3e2c~~), which are most likely Raymond arches (~~Vaughan et al., 1999; Drews et al., 2003~~) ~~and we,~~ and are conducting ice-flow modelling experiments to interpret them. Regardless of its cause, the upward arch locations can be a proxy of the summit position in the past (Nereson et al., 1998).

Since the arches in the top ~ 40 m are aligned vertically below the summit (Fig. ~~3b2b~~), it indicates that the summit position has been stable over the past several decades. In contrast, the deeper arches (~ 300 m a.s.l. and below) show more offset towards the southeast as it goes deeper (radar profile 2–2' shown in Fig. ~~3e2c~~). This trend is also found in other radar profiles. A possible interpretation of this arch inclination is that the summit has migrated towards northwest in the past. We do not see clear signs of present-day mass imbalance implying such divide migration (Fig. 6). It may indicate recent changes in mass balance and/or a limitation of our mass balance estimate due to a coarse resolution (~~1000's~~ thousands of meters), compared to the observed arch offset (100–400 m).

If the upward arches at greater depths are ~~the~~ Raymond arches, it indicates that the summit position ~~was steady~~ has been stable within several ice thicknesses from the current summit over one characteristic time $T (= H/SMB, \sim 610$ years at Blåskimen Island) or longer. If the summit position ~~is steady~~ stays stable for longer time, then ~~the~~ Raymond arches are further developed into double-peaked arches (Martín et al., 2009a), which are not clearly observed here (small side arch is caused by bed bump nearby, according to our initial ice-flow modeling). According to Martín et al. (2009a), the double-peaked arches appear after several T , though its mature shape is reached only after $\sim 10T$ or so. Therefore, we speculate that the summit of Blåskimen Island has been stable within several kilometers at least in the past ~ 600 years but no longer than several millennia.

Roles of ice rises vary largely in terms of its settings, and thus can change during the evolution (Matsuoka et al., 2015). Blåskimen Island is currently situated at the calving front of the local ice shelves and in the ice-flow shadow of Novyy Island ice rise upstream (Fig. 1). This setting implies that currently Blåskimen Island alone has limited impact on the continental grounding line and ice flux from the ice sheet. However, it seems likely that Blåskimen Island plays a more significant role than the Novyy Island to maintain the current calving front position. Favier and Pattyn (2015) demonstrated that an ice shelf landward of the ice rise is thicker than the seaward ice shelf facilitating formation of rifts and ice-shelf breakups just seaward of the ice rise; ~~an observational example is shown in Fig. 2 of Matsuoka et al. (2015). It can practically mean that the ice shelf cannot extend beyond an ice rise without additional buttressing. Similarly, Blåskimen Island maintains the calving fronts of Jelbert and Fimbul ice shelves to their present positions.~~ To explore the dynamics of this surrounding region better we will use the datasets presented in this study to model the evolution of the ice rise.

30 7 Conclusions

Ice rises are a useful resource to investigate evolution and past climate of the DML coastal region. We investigate Blåskimen Island ice rise, one of the larger isle-type ice rises at the calving front of the intersection of Fimbul and Jelbart Ice Shelves,

using geophysical methods. It has an overall dome shape and at its summit it is ~ 350 m above the adjacent ice shelf. It stands over a flat bed with a mean elevation of 110 m below the sea level. The ice flows from the summit towards the flank with speeds up to 15 m a^{-1} . We found a good agreement in the spatial patterns of stake-measured Surface Mass Balance (SMB) between 2013 and 2014, and radar-measured SMB between 2005 and 2014. Both show higher SMB on the upwind slopes (southeast), in comparison to downwind slope (northwest) by $\sim 37\%$. This variation is likely a result of orographic precipitation during storms ~~from the northeast. We observed numerous sastrugi and harder snow on the northwestern slopes, indicating strong impact of wind.~~ Using the Input-Output method for a range of parameters and column setups, we conclude that Blåskimen Island has been thickening over the past ~~nine years~~ decade. Thickening rates cannot be determined precisely, but ensemble results show that thickening rate averaged over the ice rise ~~varies between 0.07~~ can be between 0.12 m a^{-1} and ~~0.35~~ 0.37 m a^{-1} . On longer timescales, we speculate that the summit of Blåskimen Island has been stable within several kilometers at least in the past ~ 600 years but no longer than several millennia.

8 Data availability

Field data (GPR, GPS, borehole thermistor data) and the derived datasets (ice thickness, flow speed, and surface mass balance) will be released at: <http://data.npolar.no> on the completion of the review process. They are available for the editor and reviewers upon request. The ~~23-m~~ 23 m long firn-core data and their availability are described in Vega et al. (2016).

Author contributions. Matsuoka and Brown designed the study. All three authors conducted fieldwork. Goel led the overall data analysis and interpretations. Brown prepared the GPS and GPR processing workflow, which Goel adapted. Brown also produced inversion SMB estimates. Goel and Matsuoka prepared the manuscript, and Brown contributed to finalize it.

Competing interests. Author KM is a member of the editorial board of the journal.

Acknowledgements. This work was funded by grants from Norwegian Antarctic Research Expeditions (NARE) and the Center for Ice Climate and Ecosystems (ICE) of the Norwegian Polar Institute. Vikram Goel received a PhD studentship from National Centre for Antarctic and Ocean Research (NCAOR), through a financial support from the Ministry of Earth Sciences (MoES), Government of India. We thank Harvey Goodwin, Kjetil Bakkland, Ørjan Karlsen, and Peter Leopold for their contributions to fieldwork. Troll Station of NARE and SANAE Station of the South African National Antarctic Programs provided field support. Tor Ivan Karlsen developed the low-frequency radar. Carmen P. Vega, Elisabeth Isaksson, and their team provided the depth-age data of the core. We thank our two reviewers for their constructive comments, which improved the manuscript significantly. Figure 1 was developed using a free GIS data package Quantarctica (quantarctica.npolar.no).

References

- Bamber, J. L., Gomez-Dans, J. L., and Griggs, J. A.: A new 1 km digital elevation model of the Antarctic derived from combined satellite radar and laser data–Part 1: Data and methods, *The Cryosphere*, 3, 101–111, 2009.
- Bindschadler, R., Choi, H., Wichlacz, A., Bingham, R., Bohlander, J., Brunt, K., Corr, H., Drews, R., Fricker, H., Hall, M., Hindmarsh, R., Kohler, J., Padman, L., Rack, W., Rotschky, G., Urbini, S., Vornberger, P., and Young, N.: Getting around Antarctica: new high-resolution mappings of the grounded and freely-floating boundaries of the Antarctic ice sheet created for the International Polar Year, *The Cryosphere*, 5, 569–588, doi:10.5194/tc-5-569-2011, 2011.
- Borstad, C. P., Rignot, E., Mouginit, J., and Schodlok, M. P.: Creep deformation and buttressing capacity of damaged ice shelves: theory and application to Larsen C ice shelf, *The Cryosphere*, 7, 1931–1947, doi:10.5194/tc-7-1931-2013, 2013.
- 10 Conway, H. and Rasmussen, L. A.: Recent thinning and migration of the Western Divide, central West Antarctica, *Geophys. Res. Lett.*, 36, L12 502, doi:10.1029/2009GL038072, 2009.
- Conway, H., Hall, B. L., Denton, G. H., Gades, A. M., and Waddington, E. D.: Past and future grounding-line retreat of the West Antarctic Ice Sheet, *Science*, 286, 280–283, doi:10.1126/science.286.5438.280, 1999.
- Cuffey, K. M. and Paterson, W. S. B.: *The Physics of Glaciers*, Academic Press, 2010.
- 15 Drews, R., Martín, C., Steinhage, D., and Eisen, O.: Characterizing the glaciological conditions at Halvfarryggen ice dome, Dronning Maud Land, Antarctica, *J. Glaciol.*, 59, 9–20, doi:10.3189/2013JoG12J134, 2013.
- Drews, R., Matsuoka, K., Martín, C., Callens, D., Bergeot, N., and Pattyn, F.: Evolution of Derwael Ice Rise in Dronning Maud Land, Antarctica, over the last millennia, *J. Geophys. Res.-Earth*, 120, 2014JF003 246, doi:10.1002/2014JF003246, 2015.
- Dupont, T. K. and Alley, R. B.: Assessment of the importance of ice-shelf buttressing to ice-sheet flow, *Geophys. Res. Lett.*, 32, L04 503, doi:10.1029/2004GL022024, 2005.
- 20 Favier, L. and Pattyn, F.: Antarctic ice rise formation, evolution, and stability, *Geophys. Res. Lett.*, 42, 2015GL064 195, doi:10.1002/2015GL064195, 2015.
- [Favier, L., Pattyn, F., Berger, S., and Drews, R.: Dynamic influence of pinning points on marine ice-sheet stability: a numerical study in Dronning Maud Land, East Antarctica, *The Cryosphere*, 10, 2623–2635, doi:10.5194/tc-10-2623-2016, 2016.](#)
- 25 Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., Bianchi, C., Bingham, R. G., Blankenship, D. D., Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita, S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W., Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langley, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., Matsuoka, K., Mouginit, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippin, D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C., and Zirizzotti, A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, *The Cryosphere*, 7, 375–393, doi:10.5194/tc-7-375-2013, 2013.
- 30 Fürst, J. J., Durand, G., Gillet-Chaulet, F., Tavard, L., Rankl, M., Braun, M., and Gagliardini, O.: The safety band of Antarctic ice shelves, *Nature Climate Change*, 6, 479–482, doi:10.1038/nclimate2912, 2016.
- Gillet-Chaulet, F., Hindmarsh, R. C. A., Corr, H. F. J., King, E. C., and Jenkins, A.: In-situ quantification of ice rheology and direct measurement of the Raymond Effect at Summit, Greenland using a phase-sensitive radar, *Geophys. Res. Lett.*, 38, L24 503, doi:10.1029/2011GL049843, 2011.

- Goodwin, A. H. and Vaughan, D. G.: A topographic origin for double-ridge features in visible imagery of ice divides in Antarctica, *J. Glaciol.*, 41, 483–489, doi:10.3198/1995JoG41-139-483-489, 1995.
- Hawley, R. L., Courville, Z. R., Kehrl, L. M., Lutz, E. R., Osterberg, E. C., Overly, T. B., and Wong, G. J.: Recent accumulation variability in northwest Greenland from ground-penetrating radar and shallow cores along the Greenland Inland Traverse, *J. Glaciol.*, 60, 375–382, doi:10.3189/2014JoG13J141, 2014.
- Herring, T. A., King, R. W., and McClusky, S. C.: *Introduction to GAMIT/GLOBK*, Massachusetts Institute of Technology, Cambridge, Massachusetts, 2010.
- Hindmarsh, R. C. A., King, E. C., Mulvaney, R., Corr, H. F. J., Hiess, G., and Gillet-Chaulet, F.: Flow at ice-divide triple junctions: 2. Three-dimensional views of isochrone architecture from ice-penetrating radar surveys, *J. Geophys. Res.-Earth*, 116, F02024, doi:10.1029/2009JF001622, 2011.
- Hvidberg, C. S.: Steady-state thermomechanical modelling of ice flow near the centre of large ice sheets with the finite-element technique, *Ann. Glaciol.*, 23, 116–123, doi:10.3198/1996AoG23-116-123, 1996.
- Høydal, Ø. A.: A force-balance study of ice flow and basal conditions of Jutulstraumen, Antarctica, *J. Glaciol.*, 42, 413–425, doi:10.3198/1996JoG42-142-413-425, 1996.
- Jezeq, K. C., Curlander, J. C., Carsey, F., Wales, C., and Barry, R.: RAMP AMM-1 SAR Image Mosaic of Antarctica, Version 1, Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center, doi:10.5067/dhc0hdhzq184, accessed: 2012-09-31, 2002.
- King, J. C.: Wind-borne redistribution of snow across an Antarctic ice rise, *J. Geophys. Res.*, 109, doi:10.1029/2003JD004361, 2004.
- [Kingslake, J., Hindmarsh, R. C. A., Aðalgeirsdóttir, G., Conway, H., Corr, H. F. J., Gillet-Chaulet, F., Martín, C., King, E. C., Mulvaney, R., and Pritchard, H. D.: Full-depth englacial vertical ice sheet velocities measured using phase-sensitive radar, *J. Geophys. Res. Earth Surf.*, 119, 2014JF003275, doi:10.1002/2014JF003275, 00003, 2014.](#)
- [Kingslake, J., Martín, C., Arthern, R. J., Corr, H. F. J., and King, E. C.: Ice-flow reorganization in West Antarctica 2.5 kyr ago dated using radar-derived englacial flow velocities: WEST ANTARCTIC ICE-FLOW REORGANIZATION, *grl*, 43, 9103–9112, doi:10.1002/2016GL070278, 2016.](#)
- Kovacs, A., Gow, A. J., and Morey, R. M.: The in-situ dielectric constant of polar firn revisited, *Cold Reg. Sci. Technol.*, 23, 245–256, doi:10.1016/0165-232X(94)00016-Q, 1995.
- Lenaerts, J. T., Brown, J., Van Den Broeke, M. R., Matsuoka, K., Drews, R., Callens, D., Philippe, M., Gorodetskaya, I. V., Van Meijgaard, E., Reijmer, C. H., Pattyn, F., and Van Lipzig, N. P.: High variability of climate and surface mass balance induced by Antarctic ice rises, *J. Glaciol.*, 60, 1101–1110, doi:10.3189/2014JoG14J040, 2014.
- [Mackintosh, A. N., Verleyen, E., O'Brien, P. E., White, D. A., Jones, R. S., McKay, R., Dunbar, R., Gore, D. B., Fink, D., Post, A. L., Miura, H., Leventer, A., Goodwin, I., Hodgson, D. A., Lilly, K., Crosta, X., Golledge, N. R., Wagner, B., Berg, S., van Ommen, T., Zwart, D., Roberts, S. J., Vyverman, W., and Masse, G.: Retreat history of the East Antarctic Ice Sheet since the Last Glacial Maximum, *Quaternary Science Reviews*, 100, 10–30, doi:10.1016/j.quascirev.2013.07.024, 2014.](#)
- Martín, C., Gudmundsson, G. H., Pritchard, H. D., and Gagliardini, O.: On the effects of anisotropic rheology on ice flow, internal structure, and the age-depth relationship at ice divides, *J. Geophys. Res.*, 114, doi:10.1029/2008JF001204, ~~2009~~, 2009a.
- [Martín, C., Hindmarsh, R. C. A., and Navarro, F. J.: On the effects of divide migration, along-ridge flow, and basal sliding on isochrones near an ice divide, *J. Geophys. Res.*, 114, doi:10.1029/2008JF001025, 2009b.](#)
- Matsuoka, K., Pattyn, F., Callens, D., and Conway, H.: Radar characterization of the basal interface across the grounding zone of an ice-rise promontory in East Antarctica, *Ann. Glaciol.*, 53, 29–34, doi:10.3189/2012AoG60A106, 2012a.

- Matsuoka, K., Power, D., Fujita, S., and Raymond, C. F.: Rapid development of anisotropic ice-crystal-alignment fabrics inferred from englacial radar polarimetry, central West Antarctica, *J. Geophys. Res.*, 117, F03 029, doi:10.1029/2012JF002440, 2012b.
- Matsuoka, K., Hindmarsh, R. C. A., Moholdt, G., Bentley, M. J., Pritchard, H. D., Brown, J., Conway, H., Drews, R., Durand, G., Goldberg, D., Hattermann, T., Kingslake, J., Lenaerts, J. T. M., Martín, C., Mulvaney, R., Nicholls, K. W., Pattyn, F., Ross, N., Scambos, T., and Whitehouse, P. L.: Antarctic ice rises and rumples: Their properties and significance for ice-sheet dynamics and evolution, *Earth-Sci. Rev.*, 150, 724–745, doi:10.1016/j.earscirev.2015.09.004, 2015.
- Maule, C. F., Purucker, M. E., Olsen, N., and Mosegaard, K.: Heat Flux Anomalies in Antarctica Revealed by Satellite Magnetic Data, *Science*, 309, 464–467, doi:10.1126/science.1106888, 2005.
- Moholdt, G. and Matsuoka, K.: Inventory of Antarctic ice rises and rumples (version 1), Norwegian Polar Institute, doi:10.21334/npolar.2015.9174e644, 2015.
- Mulvaney, R., Oerter, H., Peel, D. A., Graf, W., Arrowsmith, C., Pasteur, E. C., Knight, B., Littot, G. C., and Miners, W. D.: 1000 year ice-core records from Berkner Island, Antarctica, *Ann. Glaciol.*, 35, 45–51, doi:10.3189/172756402781817176, 2002.
- Mulvaney, R., Triest, J., and Alemany, O.: The James Ross Island and the Fletcher Promontory ice-core drilling projects, *Ann. Glaciol.*, 55, 179–188, doi:10.3189/2014AoG68A044, 2014.
- 15 [Nereson, N. A. and Raymond, C. F.: The elevation history of ice streams and the spatial accumulation pattern along the Siple Coast of West Antarctica inferred from ground-based radar data from three inter-ice-stream ridges, *J. Glaciol.*, 47, 303–313, doi:10.3189/172756501781832197, 2001.](#)
- Nereson, N. A. and Waddington, E. D.: Isochrones and isotherms beneath migrating ice divides, *J. Glaciol.*, 48, 95–108, doi:10.3189/172756502781831647, 2002.
- 20 [Nereson, N. A., Raymond, C. F., Waddington, E. D., and Jacobel, R. W.: Migration of the Siple Dome ice divide, West Antarctica, *J. Glaciol.*, 44, 643–652, doi:10.1017/S0022143000002148, 1998.](#)
- Neumann, T. A., Conway, H., Price, S. F., Waddington, E. D., Catania, G. A., and Morse, D. L.: Holocene accumulation and ice sheet dynamics in central West Antarctica, *J. Geophys. Res.-Earth*, 113, F02 018, doi:10.1029/2007JF000764, 2008.
- ~~Noone, D., Turner, J., and Mulvaney, R.: Atmospheric signals and characteristics of accumulation in Dronning Maud Land, Antarctica, *J. Geophys. Res.-Atmos.*, 104, 19 191–19 211, 1999.~~
- 25 ~~Pettit, E. C., Jacobson, H. P., and Waddington, E. D.: Effects of basal sliding on isochrones and flow near an ice divide, *Ann. Glaciol.*, 37, 370–376, 2003.~~
- Raymond, C. F.: Deformation in the Vicinity of Ice Divides, *J. Glaciol.*, 29, 357–373, doi:10.3198/1983JoG29-103-357-373, 1983.
- Reeh, N.: A flow-line model for calculating the surface profile and the velocity, strain-rate, and stress fields in an ice sheet, *J. Glaciol.*, 34, 46–55, doi:10.3198/1988JoG34-116-46-54, 1988.
- Richardson, C., Aarholt, E., Hamran, S.-E., Holmlund, P., and Isaksson, E.: Spatial distribution of snow in western Dronning Maud Land, East Antarctica, mapped by a ground-based snow radar, *J. Geophys. Res.-Sol. Ea.*, 102, 20 343–20 353, doi:10.1029/97JB01441, 1997.
- [Rignot, E. and Kanagaratnam, P.: Changes in the Velocity Structure of the Greenland Ice Sheet, *Science*, 311, 986–990, doi:10.1126/science.1121381, 2006.](#)
- 35 Rignot, E., Mouginot, J., and Scheuchl, B.: Ice Flow of the Antarctic Ice Sheet, *Science*, 333, 1427–1430, doi:10.1126/science.1208336, 2011.
- Siegert, M., Ross, N., Corr, H., Kingslake, J., and Hindmarsh, R.: Late Holocene ice-flow reconfiguration in the Weddell Sea sector of West Antarctica, *Quaternary Sci. Rev.*, 78, 98–107, doi:10.1016/j.quascirev.2013.08.003, 2013.

Vaughan, D. G., Corr, H. F. J., Doake, C. S. M., and Waddington, E. D.: Distortion of isochronous layers in ice revealed by ground-penetrating radar, *Nature*, 398, 323–326, doi:10.1038/18653, 1999.

Vega, C. P., Schlosser, E., Divine, D. V., Kohler, J., Martma, T., Eichler, A., Schwikowski, M., and Isaksson, E.: Surface mass balance and water stable isotopes derived from firn cores on three ice rises, Fimbul Ice Shelf, Antarctica, *The Cryosphere*, 10, 2763–2777, doi:10.5194/tc-10-2763-2016, 2016.

5

Waddington, E. D., Neumann, T. A., Koutnik, M. R., Marshall, H.-P., and Morse, D. L.: Inference of accumulation-rate patterns from deep layers in glaciers and ice sheets, *J. Glaciol.*, 53, 694–712, doi:10.3189/002214307784409351, 2007.

Zwally, H. J. and Giovinetto, M. B.: Overview and Assessment of Antarctic Ice-Sheet Mass Balance Estimates: 1992–2009, *Surv. Geophys.*, 32, 351–376, doi:10.1007/s10712-011-9123-5, 2011.

Figures




Figure 1. Blåskimen Island ice rise, Western DML. (a) Blaskimen Island (squared) located between the Fimbul and Jelbart ice shelves. Inset shows the coverage of this map. Ice rises are outlined in red (Moholdt and Matsuoka, 2015), and the grounding zone of the ice sheet is illustrated in green (Bindschadler et al., 2011). Color shows flow speed of the ice sheet and ice shelf (Rignot et al., 2011). The background of the both panels is Radarsat-1 satellite imagery (Jezek et al., 2002). Acronyms stand for [BI: Blåskimen Island](#), [JG: Jutulstraumen Glacier](#), [KM: Kupol Moskovskij](#), [KC: Kupol Ciolkovskogo](#) and [NI: Novyy Island](#). (b) Close-up view of Blåskimen Island. Pink curves show kinematic GPS profiles used to determine the surface topography and green curves show radar profiles. Yellow circles show GPS stake positions for ice-flow measurements. [Red circles show GPS stake position where 3 m long firn core was also drilled](#). The white triangle shows the location where the [23-m-long-23 m long](#) firn core was drilled. Maps are projected to the Antarctic Polar Stereographic view (EPSG3031).




Figure 2. Cross sections of the ice rise along the flowline 2–2' (Fig. 1b). (a) Surface slope (absolute number, left axis) and SMB (right axis). '+' markers show SMB derived from stake heights. Red curves show the two SMB estimates derived from radar data, with solid curve assuming only vertical variability in density, and dashed curve accounting for both vertical and lateral variability in density. (b) 400-MHz radargram. Three englacial reflectors are highlighted, which are dated with the 23 m long firn core (vertical bar) and used to determine SMB. (c) 2-MHz radargram. Data are shifted using the GPS-measured surface elevations to show the topography.




Figure 3. Ice surface (a), bed topography (b) and ice-flow field (c) of the ice rise. The local coordinates are parallel to the polar stereographic coordinates EPSG3031 and centered at the ice-rise summit. (a, b) Elevation contours are ~~20m-20 m~~ intervals. In panel b, radar profiles used to derive the bed topography are highlighted in yellow to show the data availability. Kinematic GPS survey locations are shown in Fig. 1b. (c) Surface ice-flow velocities (blue arrows originated from orange circles) overlaid on surface topography (~~30-m-interval-30 m interval~~ contours). There are more GPS stakes near the summit (Fig. 1b), which did not give significant results.




Figure 4. Surface mass balance (SMB). (a) SMB estimated with stake methods over the [2012-13](#) [2013-14](#). Circles show the locations of the installed stakes. Wheel spokes inside of the circles show the locations of [9 firn density](#) [13 surface density](#) measurements. (b) SMB estimated with radar over the past decade, [by accounting only the vertical variability of density](#). White curves show radar profiles. The [23-m-long](#) [23 m long](#) firn core was drilled near the crossover of all profiles at the summit.




Figure 5. Depth variations of upward-arch amplitudes observed near the summit (Fig. 3b2b) along four radar profiles (Fig. 1b). Dots show the arch amplitudes and dashed lines show the linear fits of the arch amplitudes to depth.




Figure 6. Recent mass balance derived with three different setups when the fraction γ of depth-averaged flow speed to the surface flow speed is equal to 0.8. (a) Polygon setup. The ice rise is divided into 6 slopes (A–F) with 3 polygons (1–3, 1 for most upstream in each slope) and a summit polygon. (b) Grid setup. (c) Flowband setup. The dashed lines show the extent of polygon and grid setups for comparison.




Figure 7. Mass balance estimates of individual polygonal column (Polygon setup, solid lines) and flowband columns (Flowband setup, dashed lines) in terms of γ , the fraction of the depth averaged flow speed to the surface flow speed. Those derived with the grid setup (not shown) are very similar as those with the polygon setup. The ensemble-mean mass balance estimates for each γ are shown for all the polygon setup (orange diamonds), grid setup (red squares), and flowband setup (green triangles) setups.