

Dear Editor,

We are grateful for the acceptance of our manuscript for publication. Again we thank the Editor Andreas Vieli for the handling of our manuscript, and the three anonymous reviewers whose comments greatly improved the manuscript. We address the final minor points before publication below.

Best wishes,

Emily Hill

Minor points to be considered:

p. 2 line 33 and p. 12 line 4: at both instances the term 'dynamic ice discharge' is used which I find awkward (and I have not seen before) as 'discharge' already includes 'dynamic', maybe 'dynamic mass loss' is more appropriate
[In both cases we have amended this to 'dynamic mass loss.'](#)

p.18 line 12: is 'correlated' here really the appropriate term. Correlation mostly refers to a statistical relationship but no such statistical correlation has been undertaken. Maybe 'associated' would be more appropriate.
A similar issue occurs on p. 22 line 12.
[In both cases we have amended this to 'associated.'](#)

p. 22 line 27: there is an 'in' missing: '... The switch to terminus retreat IN the 1990s...'
[Added in the missing 'in.'](#)

p. 23 line 6: '...the initial forcing that accelerated...', I would not use the term 'forcing' here as the thinning is rather the initial PROCESS that causes the acceleration. Forcing would rather be climate or ocean...
[Changed 'forcing' to 'process'](#)

p. 23 lines 14-15: maybe I am picky here but strictly speaking your data and analysis does not say anything directly about calving type, so saying '... we OBSERVE two dominant calving behaviours...' is not correct. You can only indirectly INFER anything on calving so I would replace 'observe' by 'infer'.
[Changed 'observe' to 'infer'](#)

p. 24 line 8: there is an 'of' missing: '...due to the calving OF large tabular...'
[Added in the missing 'of'.](#)

p. 24 line 16: '...substantial PARTS OF floating ice'
[Added in 'parts of' before 'floating ice'](#)

p. 27 line 19: awkward formulation '..too..also...' one of the two is probably enough.
[Removed 'also'](#)

Dynamic changes in outlet glaciers in northern Greenland from 1948 to 2015

Emily A. Hill¹, J. Rachel Carr¹, Chris R. Stokes², G. Hilmar Gudmundsson³

¹School of Geography, Politics, and Sociology, Newcastle University, Newcastle-upon-Tyne, NE1 7RU, UK

5 ²Department of Geography, Durham University, Durham, DH1 3LE, UK

³Department of Geography and Environmental Sciences, Northumbria University, Newcastle-upon-Tyne, NE1 8ST, UK

Correspondence to: Emily A. Hill (e.hill3@newcastle.ac.uk)

Abstract

The Greenland Ice Sheet (GrIS) is losing mass in response to recent climatic and oceanic warming. Since the mid-1990s, 10 tidewater outlet glaciers across the ice sheet have thinned, retreated, and accelerated, but recent changes in northern Greenland have been comparatively understudied. Consequently, the dynamic response (i.e. changes in surface elevation and velocity) of these outlet glaciers to changes at their termini, particularly calving from floating ice tongues, is poorly constrained. Here we use satellite imagery and historical maps to produce an unprecedented 68-year record of terminus change across 18 major outlet glaciers and combine this with previously published surface elevation and velocity datasets. Overall, recent (1995–2015) 15 retreat rates were higher than at any time in the previous 47 years (since 1948). Despite increased retreat rates from the 1990s, there was distinct variability in dynamic glacier behaviour depending on whether the terminus was grounded or floating. Grounded glaciers accelerated and thinned in response to retreat over the last two decades, while most glaciers terminating in ice tongues appeared dynamically insensitive to recent ice tongue retreat and/or total collapse. We also identify glacier geometry (e.g. fjord width, basal topography, and ice tongue confinement) as an important influence on the dynamic adjustment 20 of glaciers to changes at their termini. Recent grounded-outlet glacier retreat and ice tongue loss across northern Greenland suggests that the region is undergoing rapid change and could soon contribute substantially to sea level rise via the loss of grounded ice.

1 Introduction

Mass loss from the Greenland Ice Sheet (GrIS) has accelerated since the early 2000s, compared to the 1970s and 80s (Kjeldsen 25 et al., 2015; Rignot et al., 2008), and could contribute 0.45–0.82 m of sea level rise by the end of the 21st century (Church et al., 2013). Recent mass loss has been attributed to both a negative surface mass balance and increased ice discharge from marine-terminating glaciers (van den Broeke et al., 2016; Enderlin et al., 2014). The latter contributed ~40% of total mass loss across the GrIS since 1991 (van den Broeke et al., 2016), and increased mass loss was synchronous with widespread glacier acceleration from 1996 to 2010 (Carr et al., 2017b; Joughin et al., 2010; Moon et al., 2012; Rignot and Kanagaratnam, 2006).

Coincident with glacier acceleration, dynamic thinning has occurred at elevations <2000 m on fast flowing marine-terminating outlet glaciers (Abdalati et al., 2001; Krabill et al., 2000), particularly in the south-east and north-west of the GrIS (Csatho et al., 2014; Pritchard et al., 2009). Alongside thinning and acceleration, terminus retreat has been widespread since the 1990s (e.g. Box and Decker, 2011; Carr et al., 2017b; Jensen et al., 2016; Moon and Joughin, 2008), and several studies have identified terminus retreat as a key control on inland ice flow acceleration and dynamic thinning (Howat et al., 2005; Joughin et al., 2004, 2010; Nick et al., 2009; Thomas, 2004; Vieli and Nick, 2011).

Ice sheet wide dynamic changes have been linked to 21st century atmospheric/ocean warming, and the loss of sea-ice (e.g. Bevan et al., 2012; Cook et al., 2014; Holland et al., 2008; McFadden et al., 2011; Moon and Joughin, 2008). However, tidewater glaciers can also behave in a cyclic manner, which is not always directly related to climate forcing (Meier and Post, 1987; Pfeffer, 2007), but instead relates to their fjord geometry (Carr et al., 2013; Enderlin et al., 2013; Howat et al., 2007; Powell, 1990). These glacier cycles are characterised by slow periods of advance (up to centuries) followed by rapid unstable retreat (Meier and Post, 1987; Post, 1975; Post et al., 2011). Once initiated, terminus retreat can initiate dynamic adjustments independent of climate and instead modulated by local outlet glacier geometry and associated resistive stresses. However, differences in the nature of calving and basal/lateral resistive stresses acting at tidewater glaciers with either grounded or floating termini can alter their dynamic response to retreat. Continuous small magnitude calving events and the loss of basal and lateral resistance at grounded-termini, can prolong the dynamic readjustment at the terminus. In contrast to continuous calving, large episodic calving events often occur at floating ice tongues, which can decrease buttressing forces on grounded ice, and also increase driving stress and accelerate ice flow (e.g. MacGregor et al., 2012). In the case of floating ice tongues, the response of inland ice to large calving events depends on the amount of lateral resistive stress provided by the tongue prior to calving: the loss of portions of ice tongues that are highly fractured and/or have limited contact with the fjord margins are unlikely to substantially influence inland ice dynamics.

Most previous work at tidewater glaciers in Greenland has concentrated on the central-west and south-east regions, and notably at Jakobshavn Isbræ, Helheim, and Kangerdlugssuaq Glaciers (e.g. Howat et al., 2005, 2007; Joughin et al., 2004; Nick et al., 2009). Observations at all three glaciers showed acceleration and surface thinning following terminus retreat. Dynamic changes at Jakobshavn, have been linked to the gradual collapse of its floating ice tongue (Amundson et al., 2010; Joughin et al., 2008; Krabill et al., 2004). In northern Greenland, several glaciers have thinned (Rignot et al., 1997), accelerated (Joughin et al., 2010), and retreated, losing large sections of their floating ice tongues between 1990 and 2010 (Box and Decker, 2011; Carr et al., 2017b; Jensen et al., 2016; Moon and Joughin, 2008; Murray et al., 2015). Floating ice tongue retreat does not directly contribute to sea level rise. However, with amplified warming and surface melt forecast in northern Greenland (Fettweis et al., 2012; Franco et al., 2011; Mernild et al., 2010), floating ice tongues across the region could soon collapse entirely. As a consequence, northern Greenland could become a more substantial contributor to dynamic ice discharge mass loss and sea level

rise. However, fewer studies have focussed specifically on northern Greenland, with the exception of more detailed work at Petermann and the Northeast Greenland Ice Stream (NEGIS) (e.g. Khan et al., 2015; Nick et al., 2012). As such, few long-term records of frontal positions exist in the region, and their potential impact on inland ice flow remains unclear.

5 Here we present changes in frontal position, ice velocity and surface elevation over the last 68 years (1948 to 2015) in northern Greenland. We couple a multi-decadal annual terminus position record between 1948 and 2015 with recently published surface elevation and ice velocity datasets. We then use these datasets to evaluate dynamic responses (i.e. acceleration and thinning) to frontal position change and examine disparities in the context of glaciers with floating or grounded termini. Finally, we assess local topographic setting (i.e. fjord width and depth) as a control on glacier behaviour.

10 2 Methods

2.1 Study region

We define Northern Greenland as the region of the Greenland Ice Sheet located north of 77°N (Figure 1). This region drains ~40% of the ice sheet by area (Hill et al., 2017; Rignot and Kanagaratnam, 2006) and includes 18 major marine-terminating outlet glaciers, which emanate from 14 major catchments (Figure 1). Aside from at Petermann Glacier and the NEGIS, little
15 work has focused on the other glaciers in northern Greenland, where the presence of floating ice tongues could alter the dynamic response of inland ice to calving events. Here, we use the ice-ocean mask from the Operation IceBridge BedMachine v3 product (nsidc.org/data/IDBMG4) to categorise glaciers in northern Greenland based on either grounded or floating termini (Howat et al., 2014; Morlighem et al., 2017). We also use the grounding line in this dataset to assess the location of past ice
20 tongues. Currently, five glaciers in northern Greenland terminate in floating ice tongues (Figure 1), which range between 0.5 and 70 km long (Hill et al., 2017). An additional four glaciers have lost their ice tongues entirely over the last two decades (1995 to 2015). Our study region includes a further nine outlet glaciers, which are grounded at their termini. We note that Humboldt glacier is classified as grounded as the majority of the ~100 km long terminus is grounded, despite a small floating ice tongue in the northern section (Carr et al., 2015).




Figure 1: Study region of northern Greenland. Green circles show the location of each of 18 northern Greenland study outlet glaciers. Average glacier velocities (m a^{-1}) are shown between 1993 and 2015 derived from the multi-year mosaic dataset (Joughin et al., 2010). Black outlines show glacier drainage catchments. Symbols represent the state of the glacier terminus. Stars show glaciers which currently have floating ice tongues, circles represent glaciers which lost their ice tongues (during 1995 to 2015), squares denote glaciers which have some previous literature record of a floating ice tongue, and triangles are glaciers which are grounded at their termini and have been throughout the study record.

2.2 Terminus change

2.2.1 Data sources

- 10 The terminus positions of 18 study glaciers in northern Greenland (Figure 1) were manually digitised from a combination of satellite imagery and historical topographic navigational charts between 1948 and 2015 (Table S1). From 1975 to 2015 we used Landsat 1–5 MSS (1975–1994), Landsat 7 TM (2000–2013) and Landsat 8 (2013–2015). These scenes were acquired from the United States Geological Survey (USGS) Earth Explorer website (earthexplorer.usgs.gov). To reduce the influence of seasonal changes in terminus position, one scene per year was selected from late summer, and 70% were within one month of the 31st August. Several Landsat MSS images required additional georeferencing and were georeferenced to 2015 Landsat 8 images, as these have the most accurate georeferencing. Early Landsat scenes (1970–1980s) were supplemented with SPOT–1 imagery from the European Space Agency (ESA) (intelligence-airbusds.com). These scenes covered 8 of 18 study glaciers in 1986/87, and were also selected from late August. SPOT–1 scenes were also georeferenced to 2015 Landsat Imagery. Additionally, we used aerial photographs (2 m resolution), which were provided orthorectified by Korsgaard et al. (2016).

These covered all study glaciers between Humboldt east to L. Bistrup Bræ in 1978, and Harald Moltke Bræ, Heilprin and Tracy Glaciers in NW Greenland in 1985 (Korsgaard et al., 2016).

To extend the record of glacier terminus positions further back in time, declassified spy images from the Corona satellite were acquired from the USGS Earth Explorer website (Table S1), which covered 5 of 18 glaciers in 1962/63 and Petermann and Ryder Glaciers in 1966. These images were georeferenced to a Landsat 8 scene from 2015, with total RMSE errors of 105 to 360 m. Frontal position changes smaller than this error value were discounted from the assessment. To further assess the historical terminus positions of the glaciers we used navigational map charts from the United States Air Force 1:1,000,000 Operational Navigation Charts from 1968/69 (lib.utexas.edu/maps/onc/). These were made available through the Perry-Castañeda Library, courtesy of the University of Texas Libraries, Austin. Data from 1948 comes from AMS C501 Greenland 1: 250,000 Topographic Series maps distributed by the Polar Geospatial Centre (pgc.umn.edu/data/maps/). All maps were georeferenced to 2015 Landsat 8 imagery using a minimum of 10 ground control points (GCPs), which were tied to recognisable stationary features such as on nunataks and fjord walls. RMSE errors across all glaciers ranged between 150 and 510 m.

2.2.2 Front position mapping

Changes in glacier frontal positions were measured using the commonly adopted box method, which accounts for uneven calving front retreat (e.g. Carr et al., 2013; Howat and Eddy, 2012; Moon and Joughin, 2008). For each glacier, a rectilinear box was drawn parallel to the direction of glacier flow (Figure 2), and extending further inland than the minimum frontal position. Due to Steensby Glacier's sinuous fjord, a curvilinear box was used (see Lea et al., 2014). Glacier frontal positions were digitised in sequential images and the difference between successive terminus polygons give area changes over time within the box. Dividing these areas by the width of the reference box derives width-averaged relative glacier front positions.

Aside from georeferencing errors outlined in the previous section, the main source of error was attributed to manual digitisation (e.g. Carr et al., 2013; Howat and Eddy, 2012; Moon and Joughin, 2008). We quantified this by repeatedly digitising a ~3 km section of rock coastline 20 times for each image type or map source. The resultant total mean errors were: 3.6 m for Landsat 8, 19 m for Landsat 7 ETM, 17 m for Landsat MSS, 20 m for SPOT-1, 16 m for Orthophotographs, 21 m for Corona, and 27 m for historical maps. Overall, the mean total error associated with manual digitising was 19 m, which is below the pixel resolution of all imagery sources except the 15-m panchromatic Landsat band. The presence of sea ice and highly fractured glacier termini made terminus picking at Steensby, C. H. Ostenfeld and glaciers draining the NEGIS more difficult (Bevan et al., 2012; Howat and Eddy, 2012; Murray et al., 2015). Re-digitising all 1999-2015 Landsat terminus positions yielded additional errors of $\pm 13\%$ for these glaciers. At these glaciers, similar inaccuracies in identifying the true glacier terminus

may have occurred by the authors of the earliest map charts (1948 and 1969), and we therefore consider these to be a broad estimate of the past location of glacier termini rather than exact frontal positions.




Figure 2: Rectilinear box method used to measure glacier terminus positions. An example at Harald Moltke Bræ, NW Greenland. This includes: reference box (pink), and roughly decadal terminus positions (green to red). The glacier centreline profile is shown in white and the location of 500 m sample points (white circles). Background image is Landsat 8 band 8 from the USGS Earth Explorer.

2.2.3 Change point analysis

‘Change point’ analysis can be used to objectively identify significant breaks in time series data (Bunce et al., 2018; Carr et al., 2017a). Here we used change point analysis to detect statistically significant breaks in the time-series (1948 to 2015) of terminus position change for each of our 18 study glaciers in northern Greenland. We then compare the timing and duration of these breaks to determine if distinct patterns of terminus change behaviour exist based on terminus type (grounded or floating). We employ a similar method introduced in more detail by Carr et al. (2017a). To do this we use the ‘findchangepts’ function in MATLAB software which employs the methodology of Killick et al. (2012) and Lavielle (2005). Linear regression was used to detect significant breaks in the frontal position time series: a change point was identified where there was a significant change in the mean and regression coefficients (slope and intercept) of the linear regression equation on either side of a data point. In addition, to allow for the fully automatic estimation of the number of change points, we included a minimum threshold penalty value, which we set as the mean terminus position for each glacier. This penalty only allows a change point to occur when the time-series deviates significantly from the threshold value. Using these automatically-identified change points we can determine if statistically different changes in the rate of terminus position change exists for study glaciers in northern Greenland according to their terminus type.

2.3 Ice velocity and surface elevation

Previously published datasets of annual ice velocity and surface elevation change were compiled to assess dynamic glacier changes in northern Greenland. Velocity and surface elevation change datasets are generally only available from 1990 onwards. The earliest velocity maps from winters 1991/92 and 1995/96 were acquired from the European Remote Sensing (ERS) satellites (1 and 2), as part of the ESA Greenland Ice Sheet CCI (Climate Change Initiative) project (Nagler et al., 2016). The earlier (1991/92) covers northern Greenland drainage basins from Humboldt and then east to Hagen Bræ, and the later (1995/96) covers all 18 study glaciers. Using dataset error maps we estimated average errors in velocity magnitude across all northern Greenland drainage basins, which were 2.5 m a^{-1} for 1991/92 and 10 m a^{-1} for 1995/96.

Subsequent velocity datasets were primarily acquired from the NASA MEaSUREs program (Joughin et al., 2010). These velocity maps were derived from 500 m resolution Interferometric Synthetic Aperture Radar (InSAR) pairs from the RADARSAT satellite in winter 2000/01, and then annually from winter 2005/06 to 2009/10 (Joughin et al., 2010). Using the dataset error values (Joughin et al., 2010), we estimate mean velocity errors across all years and study drainage catchments to be 6.3 m a^{-1} . For 7 study glaciers, additional annual velocity data, derived from ERS1, ERS2 and Envisat satellites, were available annually between 1991/92 to 1997/98 and between 2003/04 to 2009/10 from the ESA Greenland CCI project (Nagler et al., 2016). Winter velocities from these data were calculated from October to April.

For the winters of 2010/11, 2011/12 and 2012/13, glacier velocity maps were also acquired from InSAR (TerraSAR-X image pairs) for 11 of 18 study glaciers (Joughin et al., 2010). Despite higher spatial resolution (100 m), these maps are limited to the grounding line and extend 27–56 km inland. Mean error for these data is 23 m a^{-1} across all years (Joughin et al., 2010). Winter velocities for 2013/14 were derived from intensity tracking of RADARSAT-2 satellite data, and from offset tracking of Sentinel-1 radar data for 2014/15 and 2015/16, as part of the ESA CCI project (Nagler et al., 2016). The published mean error of these data from a central section of northern Greenland is 7.3 m a^{-1} (Nagler et al., 2015). Using the earliest full regional velocity map (1995/96) and the most recent record (2015/16), the rate of annual velocity change was calculated over this 20-year period.

We use surface elevation change data from ERS-1, ERS-2, Envisat, and Cryosat-2 radar altimetry for 1992 to 2015, which were made available by the ESA's Greenland Ice Sheet CCI project (Khvorostovsky, 2012; Simonsen and Sørensen, 2017; Sørensen et al., 2015). Data from 1992 to 2011 were acquired from the ERS-1, ERS-2 and Envisat satellites, using a combination of cross-over and repeat track analysis, which have then been merged to create a continuous dataset across satellites (Khvorostovsky, 2012). These data are provided in 5-year running means from 1992–1996 to 2007–2011 and at a resolution of 5 km. For the most recent elevation change (2011 to 2015), we used Cryosat-2 satellite elevation change which are provided in 2-year means (Simonsen and Sørensen, 2017). These data were generated using the Least Mean Squares

method, where grid cells were subtracted from the Greenland Ice Mapping Project (GIMP) DEM (Howat et al., 2014) and corrected for backscatter and leading edge width (Simonsen and Sørensen, 2017). Calculations were made at a 1 km grid resolution and resampled to 5 km to conform with 1992–2011 datasets (Simonsen and Sørensen, 2017). Using error estimates (Simonsen and Sørensen, 2017), we calculated mean errors across all years and across all northern Greenland drainage basins to be $\pm 0.14 \text{ m a}^{-1}$. Elevation changes from 1992–1996 were compared to elevation changes for 2014–2015 to assess how changes in surface elevation have evolved during the study period.

Velocity and surface elevation time series were extracted along each glacier centreline, which were drawn following Lea et al. (2014). To draw these centrelines we first calculated euclidean distance between parallel fjord walls that were digitised in 2015 Landsat 8 imagery. The maximum distance line (i.e. centreline) was then traced from the furthest terminus extent back to the inland end of the glacier fjord. Annual average velocities were calculated within 5 km inland of the grounding line of each glacier, and elevation change rates were averaged across the entire centreline profile due to poorer/coarser data resolution (Figure 2).

2.4 Fjord width and basal topography

To assess the control of fjord geometry on outlet glacier behaviour we calculate fjord width and depth. Fjord width was measured perpendicular to glacier centrelines following Carr et al. (2014). Points were extracted at 500 m intervals along each fjord wall and joined by lines that crossed the fjord. The length of these lines is the width between the fjord walls, and changes along each fjord were fitted with a linear regression model to determine if the fjord widens or narrows with distance inland. To determine the fjord bathymetry of each study glacier in northern Greenland, regional basal topography was taken from the Operation IceBridge BedMachine v3 dataset which is derived from ice thickness and mass conservation (Morlighem et al., 2017). Basal topography was sampled at 500 m points along glacier centrelines. Using the error map from BedMachine v3 (Figure S1), we calculated errors along the grounded and non-grounded portions of each glacier centreline profile (Table S2). Mean grounded bed topography errors at 14 of 18 study glaciers range between 25 and 87 metres. These glaciers are well constrained by the mass conservation method, which works best for fast flowing areas near the glacier terminus (Morlighem et al., 2014, 2017). The remaining four glaciers (Storstrømmen, L. Bistrup Bræ, Kofoed-Hansen Bræ, and Brikkerne Glacier) have higher errors (from 112 to 215 m), owing to poor data coverage and kriging interpolation (Morlighem et al., 2017). Mean errors in bathymetry data are greater at all glaciers, averaging 156 metres and ranging from 15 to 283 m. To assess bed slope direction, we fit each glacier profile from the grounding line to 20 km inland with a linear regression model. These sections of each bed profile and model fit are presented in Figure S2, while entire bed profiles, and landward/seaward direction are presented in the results. Errors in basal topography do not significantly affect our assessment of bed slope direction, and we only use topography along the grounded portion of the glacier where errors are lowest. We treat basal profiles in the far east of the study region with caution due to their higher errors. Finally, to estimate drainage catchment areas and the percentage of

each catchment below present sea level for each study glacier, surface drainage catchments (Table 3) were delineated using the GIMP surface DEM (Howat et al., 2014) and topographic analysis functions within TopoToolbox in MATLAB (Schwanghart and Kuhn, 2010). We calculated each drainage area using catchments constrained by gradients in the surface DEM.

3. Results

3.1 Changes in glacier frontal position (1948–2015)

Across northern Greenland, 13 of the 18 study glaciers underwent net retreat between 1948 and 2015, while the remaining five advanced (Figure 3). However, long-term frontal retreat rates (1948–2015) varied between glaciers across northern Greenland (Table 1), and ranged from -15 m a^{-1} at Marie-Sophie Glacier, to twenty times greater at Petermann Glacier (-311 m a^{-1}). At outlet glaciers in northern Greenland, we expect terminus changes and dynamic response to be different dependent on terminus type (i.e. floating or grounded). Nine outlet glaciers were grounded at their terminus throughout the study period, while at the end of the study period another nine still had ice tongues or lost them during the last two decades (1996 to 2015: Figure 1, Table 1). Our statistical changepoint analysis compared the duration and magnitude of frontal position changes at all study glaciers: this confirms that, in general, there are two different types of frontal position behaviour and dynamic response to calving based on terminus type (grounded or floating).




Figure 3: Overall mean rate of terminus change (m a^{-1}) at 18 outlet glaciers in northern Greenland from 1948 to 2015. Green circles represent glaciers which have undergone overall advance during the record, while yellow to red circles represent increasing retreat rates from 0 to larger than -300 m a^{-1}

Table 1: Summary data for 18 northern Greenland outlet glaciers, ordered according to terminus type and by frontal position rate (from high retreat to high advance). Annual rates of terminus change are given for the entire record (1948 to 2015). Average velocity change along each glacier centreline from winter 1995/96 to 2015/2016. Average surface elevation change rates along each glacier centreline were differenced from the earliest record (1992-1996) and the most recent (2014/15).

	Northern Greenland Outlet Glaciers	Terminus Change (1948–2015) (m a ⁻¹)	Velocity Change (1995/96– 2015/16) (m a ⁻¹)	Difference in surface elevation change rates (1992-1996 and 2014-2015) (m a ⁻¹)
Category 1: Grounded terminus	Tracy	-173	36.8	-0.11
	Kofoed-Hansen Bræ	-169	-0.06	0.12
	Harald Moltke Bræ	-156	22.6	
	Humboldt	-111	0.32	-0.51
	Heilprin	-45	7.16	-0.15
	Academy	-31	-4.87	-0.97
	Harder	-25	0.58	-0.89
	Marie-Sophie	-15	1.03	-0.43
	Brikkerne	82	-2.56	
Category 2: Floating ice tongue	Petermann	-311	3.78	-1.34
	Zachariae Isstrøm	-282	20.3	-2.98
	Hagen Bræ	-162	6.45	-0.83
	C. H. Ostenfeld	-58	2.96	-1.26
	Nioghalvfjærdsfjorden	-28	1.62	-1.99
	Steensby	2	2.59	-0.33
	L. Bistrup Bræ	39	-3.89	0.57
	Storstrømmen	41	-1.11	-0.18
	Ryder	43	-0.08	0.47

5

In addition to the long-term record of frontal position change (1948–2015), we further assess the variability of retreat rates across northern Greenland, by presenting mean retreat rates across five decadal time periods (1976–1985, 1986–1995, 1996–2005, 2006–2015) in Figure 4 (a-e), except for the earliest epoch (1948–1975) which spans 27 years due to image availability. We also present mean decadal frontal position change based on terminus type in Table 2. During the first epoch (1948 to 1975) small advances and retreats took place across the region (< 500 m a⁻¹ magnitude). This was followed by a decade (1976–85) dominated by glacier advance, in particular at glaciers with floating ice tongues (Table 2), which had high retreat rates for the entire study period (e.g. Hagen Bræ, Zachariae Isstrøm, Petermann: Table 1). In the subsequent epoch (1986 to 1995), a mixture of advance and retreat occurred and the range of frontal position changes was great, from -780 m a⁻¹ retreat at C. H. Ostenfeld to 750 m a⁻¹ advance at Storstrømmen (Figure 4c). During the last two decades of the study period (1996 to 2015), retreat rates were substantially higher than in the previous three epochs. However, variability in the magnitude of frontal position change over this period appears to be particularly related to terminus type (grounded or floating). Decadal mean retreat rates at glaciers with ice tongues (-745 to -835 m a⁻¹) were substantially higher than at grounded-outlet glaciers (-99 to 165 m a⁻¹: Table 2). Based on the differences in terminus behaviour observed over the long-term and decadal time series based on terminus type

10

15

(grounded or floating), we now treat these as separate categories for the remainder of the results, during which we describe short-term trends derived from our changepoint analysis (Figure 5).




Figure 4: Mean decadal rates of terminus change across northern Greenland. These are shown for five epochs between 1948 and 2015. Increasing red circles represent glacier retreat rates between 0 and exceeding -1000 m a^{-1} . Increasing blue circles represent advance rates between 0 and exceeding 1000 m a^{-1} .

Table 2: Mean decadal frontal position change for all study outlet glaciers in northern Greenland, and split based on our two glacier categories of terminus type: grounded-terminus or terminating in a floating ice tongue

Mean terminus change (m a ⁻¹)	1948-1975	1976-1985	1986-1995	1996-2005	2006-2015
All (n =18)	-63.65	503.36	-7.83	-454.99	-467.06
Grounded-terminus (n =9)	-167.35	93.87	-112.07	-164.50	-99.23
Floating-terminus (n =9)	40.05	912.84	126.19	-745.49	-834.88

3.1.1 Grounded-terminus outlet glaciers

Calving-induced retreat of grounded marine-terminating outlet glaciers, contributes to dynamic ~~ice discharge~~mass loss and directly to sea level rise. At grounded outlet glaciers in northern Greenland (nine of our study glaciers: Figure 5), retreat rates increased substantially during the last two decades (1996 to 2015) of the study period. At the beginning of our record, grounded outlet glaciers went through an initial period of minimal frontal position change averaging -26 m a⁻¹ and ranging from 24 m a⁻¹ advance at Kofoed-Hansen Bræ to -105 m a⁻¹ retreat at Tracy Glacier (Figure 5). Frontal position change then switched to a period of higher magnitude retreat at eight glaciers (excluding Brikkerne), which lasted for an average of 26 years. During this period, frontal position change averaged -150 m a⁻¹, and net retreats ranged from -0.6 to 8 km. The greatest total terminus changes took place at Tracy Glacier (8 km retreat: 1981–2015), Harald Moltke Bræ (5 km retreat: 1988–2015), and Kofoed-Hansen Bræ (4.6 km: 1973–2015: Figure 6a-c). The timing of this switch from minimal change to steady retreat was not uniform, but most glaciers began steadily retreating from the 1990s to 2000s and continued at the same rate thereafter (Figure 5). The exception to this pattern of behaviour is Brikkerne glacier, which instead advanced by 9 km between 1968 and 1978 before returning to minimal terminus change (Figure 6i).

3.1.2 Glaciers with floating ice tongues

The retreat of floating glacier termini can reduce the resistive stresses at the terminus and increase the dynamic glacier response to calving. Rapid retreat in the form of large episodic calving events removed substantial floating ice sections (11.6–26 km net retreat: Figure 7) from several northern Greenland outlet glaciers (Zachariae Isstrøm, Petermann and Steensby, C. H. Ostfeld, and Hagen Bræ) between 1995 and 2015. However, frontal position changes at some glaciers with floating ice tongues were instead cyclic. At the beginning of the study period, six of the nine study glaciers with floating ice tongues showed minimal terminus change/advance (93 m a⁻¹), followed by short-lived rapid retreat, lasting less than 6 years on average (Figure 5). During the phases of rapid retreat, rates ranged between -700 m a⁻¹ at Nioghalvfjærdsfjorden to -8997 m a⁻¹ at Petermann Glacier (Figure 5), and were on average 40 times greater (-4536 m a⁻¹) than during the steady retreat phases at glaciers grounded at their terminus (-150 m a⁻¹). During this period, large calving events led to complete ice tongue loss at Zachariae Isstrøm by 2011/12, and at C. H. Ostfeld, Steensby and Hagen Bræ by 2016 (Figure 7). Rapid retreat was often followed by another period of relative minimal terminus change (-437 m a⁻¹) compared to order of magnitude earlier retreat (e.g. Petermann Glacier

and Hagen Bræ: Figure 5). Similar to glaciers with grounded-termini, the timing of the switch to rapid retreat was not synchronous, but mainly occurred after 1990 (Figure 5). At most glaciers, the duration of rapid retreat was short-lived (< 5 years) in comparison to the duration of steady retreat (> 13 years) at grounded-glaciers.




Figure 5: Retreat rates during identified changepoint time periods for each pre-defined category of glacier based on either grounded or floating termini. Glaciers are then ordered based on their overall (1948-2015: Table 1) frontal position change rates within each of these categories. Grey bars show their periods of minimal/variable terminus change (in some cases advance) and turquoise bars show the period of higher magnitude frontal position change.

Several glaciers with floating ice tongues (Storstrømmen, L. Bistrup Bræ, and Ryder) have instead shown cyclic periods of advance and retreat between 1948 and 2015 (Figure 7g-i). Periods of terminus advance at these glaciers averaged $\sim 420 \text{ m a}^{-1}$ and lasted for an average of 18 years (Figure 5). Adjacent glaciers Storstrømmen and L. Bistrup Bræ advanced during a similar period (from 1973 to 1990), and for $\sim 13\text{--}17$ years. After this, both glaciers underwent relatively limited terminus change from

2000 onwards (Figure 5). Despite synchronous advance, their advance rates differed by almost an order of magnitude (89 m a^{-1} at L. Bistrup Bræ, and 725 m a^{-1} at Storstrømmen, Figure 5). At Ryder Glacier, there were four main cycles of glacier advance and retreat during the record. These took place between 1948–1996, 1968–1986, 1999–2006, and 2008–2015 and advance rates ranged from 183 to 750 m a^{-1} (Figure 5). Periods of advance (7–48 years) were separated by shorter periods (2–13 years) of higher magnitude retreat (ranging from -960 to -1950 m a^{-1}) (Figures 5).




Figure 6: Front position, velocity and elevation change at nine outlet glaciers grounded at their terminus in northern Greenland. Left axes show relative front position (black line) between 1948 and 2015 relative to their initial position in 1948. Grounding line velocities (orange) on right axes one between 1996 and 2015. Surface elevation changes averaged across the glacier centreline profile (green) for 1996 to 2015.

3.2 Ice velocity change

3.2.1 Grounded-terminus outlet glaciers

Overall, most grounded outlet glaciers in northern Greenland accelerated along their centreline profiles (ranging from 0.32 to 37 m a^{-1}) from 1996 to 2016 (Table 1). Acceleration at these glaciers was also greatest (averaging 27%), during periods of

steady retreat (Figure 5). This was particularly the case in northwest Greenland at Heilprin, Tracy, and Harald Moltke Brae. At Heilprin Glacier this resulted in a 49% increase (from 458 to 681 m a⁻¹) in grounding line velocity from 2001 to 2016 (Figure 6e), during which the glacier retreated at -110 m a⁻¹ (Figure 5). Substantially greater acceleration (89%) took place at Tracy Glacier from 1996 to 2016 (Figure 6a), which was associated with higher magnitude retreat rates (-263 m a⁻¹: Figure 5). Harald Moltke Bræ also accelerated between 1990 and 2016 (22 m a⁻¹: Table 1), and retreated at -196 m a⁻¹ (Figure 5). However, it underwent two very large velocity increases (> 1000 m a⁻¹) between 2001 and 2006 and again during winter 2013/14, both of which coincided with short-lived glacier advance (0.5–0.8 km: Figure 6c). Slower flowing grounded outlet glaciers in the region (< 400 m a⁻¹: Humboldt, Harder, and Marie-Sophie Glaciers) also accelerated by 27–108% during steady retreat (Figure 6). Alternatively, some grounded-terminus outlet glaciers did not show substantial acceleration following retreat: Kofoed-Hansen Bræ and Academy Glacier had sustained periods of steady retreat, but showed no net trend in velocity and high variability (Figure 6b,f), which did not coincide with periods of increased retreat rates (Figure 5). Brikkerne Glacier decelerated from 1996 to 2016, while the terminus position changed little (Figure 6i).

3.2.2 Glaciers with floating ice tongues

Despite major retreat episodes and ice tongue disintegration, most glaciers in northern Greenland that terminate in floating ice tongues (except Zachariae Isstrøm) showed minimal net velocity change between 1996 and 2016 (Table 1). In contrast to grounded-outlet glaciers, the velocity response to retreat was also more variable between individual glaciers. However, we identify two main patterns in velocity change following periods of rapid retreat: 1) short-lived, minimal glacier acceleration, followed by some deceleration, 2) continuous acceleration following initial terminus retreat. The first pattern of velocity change encompasses seven of nine glaciers with floating ice tongues, the clearest examples of which are at C. H. Ostenfeld, Hagen Bræ, Petermann Glacier and Steensby Glacier. In all cases, rapid ice tongue retreat was followed by short-lived (< 3 year) low magnitude (<8% acceleration, but ~25% at Steensby) grounding-line acceleration. After retreat ceased, ice flow decelerated, ranging from 2% at Petermann to 28% at Hagen Bræ. In addition, prior to rapid retreat some glaciers advanced. In the year preceding rapid retreat (2005 to 2006), Hagen Brae showed higher magnitude acceleration (~52%) alongside some glacier advance. This was also the case at Ryder Glacier, which showed cyclic behaviour, of grounding line acceleration (~8%: 4.7–5.5 m a⁻¹) during both 7-year periods of terminus advance, followed by deceleration (11%) during high magnitude retreat (~2 years) in-between periods of advance. Storstrømmen and L. Bistrup Bræ also show evidence of some acceleration immediately following retreat, later followed by deceleration (Figure 7h,i) from 2010 to 2016. However, in contrast to most glaciers which flow fastest at their terminus, velocities at both glaciers are fastest inland, and decrease with distance towards the terminus (Figure 8). Grounding line terminus velocities accelerated by 350% and 150% at Storstrømmen and L. Bistrup Bræ throughout the record (1996 to 2016: Figure 7); and velocities ~20–40 km inland decelerated by 10–15 m a⁻¹ (> 54%).

The second pattern of velocity change occurred at both glaciers draining the NEGIS where ice tongue retreat was followed by gradual glacier acceleration in the subsequent decade (2006 to 2016: 43% at Zachariae Isstrøm and 10% at Nioghalvfjærdsfjorden). This prolonged glacier acceleration following retreat, is more similar to patterns observed on grounded termini, rather than the other floating tongues. Further, the removal of the entire ice tongue at Zachariae Isstrøm in 2011/12 was followed by glacier acceleration (125 m a^{-2} : 2012 to 2016, Figure 7g), whereas other glaciers (e.g. C. H. Ostenfeld and Hagen Bræ) underwent a similar collapse, but changes in velocities were limited. Despite this behaviour in the northeast of the study region, the majority of glaciers in northern Greenland showed negligible acceleration in response to retreat and/or collapse of their floating ice tongues.




Figure 7: Front position, velocity and elevation change at nine outlet glaciers which terminate in floating ice tongues in northern Greenland. Left axes show relative front position (black line) between 1948 and 2015 relative to their initial position in 1948. Grounding line velocities (orange) on right axes one between 1996 and 2015. Surface elevation changes averaged across the glacier centreline profile (green) for 1996 to 2015.

3.3 Surface elevation change

3.3.1 Grounded-terminus outlet glaciers

Alongside continuous retreat and acceleration at outlet glaciers with grounded-termini, thinning rates increased between 1992–1996 and 2014–2015 (Table 1). In most cases, surface lowering was synchronous with the start of their steady retreat (Figure 6). Examples of this were at Marie-Sophie and Academy Glaciers, where small increased thinning or reduced thickening rates (1999 to 2000: Figure 6f,h), were followed by high retreat rates in the following years at both Marie-Sophie (-130 m a^{-1} : 2001 to 2004) and Academy Glacier (-205 m a^{-1} : 2001 to 2003). Periods of greater retreat (2001 to 2003/04) were followed by dramatically increased thinning rates at both glaciers to -0.3 m a^{-1} (Academy) and -0.16 m a^{-1} (Marie-Sophie: Figure 6f,h). Thinning rates similarly increased strongly from -0.19 m a^{-1} to -0.78 m a^{-1} at Humboldt Glacier from 1996–2005 to 2005–2012, which coincided with increased retreat rates (-98 to -160 m a^{-1} : Figure 6d). Limited data prevent us from commenting in depth on elevation changes at glaciers in NW Greenland. However, the few years of data available at Harald Moltke Bræ show increased thinning between 2012 and 2015, coincident with retreat (Figure 6c). Within this record lies an anomalous year of reduced thinning rates (2013 to 2014), which were coincident with an order of magnitude increase in velocity ($\sim 1000 \text{ m a}^{-1}$) and 0.8 km terminus advance.

3.3.2 Glaciers with floating ice tongues

In general, glaciers with floating ice tongues had higher thinning rates than grounded termini from 1992–1996 to 2014–2015 (Table 1), and were characterised by short-lived increases in thinning rates following ice tongue retreat. In several cases (e.g. Petermann, Hagen Bræ, and Zachariae Isstrøm), slight thickening occurred before ice tongue retreat/collapse, followed by a switch to thinning immediately before large calving events (Figure 7). For example, Petermann Glacier switched from negligible thickening in 2008 (0.03 m a^{-1}) to thinning (-0.22 m a^{-1}) in 2009, before the removal of 27 km of floating ice in the following three years (2010 to 2013: Figure 7a). At Zachariae Isstrøm a switch to thinning was synchronous with the onset of rapid retreat in 2003 (Figure 7b), although thinning rates were greater once the entire ice tongue was lost (2011 to 2012). During or immediately after floating ice tongue retreat, thinning rates increased from minimal change ($< -0.2 \text{ m a}^{-1}$ thinning) to -0.8 m a^{-1} at Petermann Glacier (2010 to 2013), -1.7 m a^{-1} at Hagen Bræ (2007/11 to 2012/13), and -2 m a^{-1} at Zachariae Isstrøm (2011/12 to 2012/13: Figure 7). This increased thinning was also coincident with acceleration during the years following ice tongue removal (Figure 7). Other glaciers showed more gradual and smaller increases in thinning rates (Figure 7). For example, at C. H. Ostenfeld the removal of 21 km of floating ice between 2002 and 2003 was followed by a steady and low magnitude increased thinning rates at a rate of -0.15 m a^{-1} from 2006 to 2014 (Figure 7d). In this case, velocity increases alongside increased thinning rates were also gradual, but minimal in comparison to other glaciers. Ryder Glacier also showed increased thinning rates prior to retreat (2005–2006) but was followed by a rapid switch to thickening as ice flow accelerated, and the calving front advanced (Figure 7i).

Two glaciers with floating ice tongues in northeast Greenland showed a different pattern of elevation change to the rest of the region: Storstrømmen and L. Bistrup Bræ thinned at the glacier terminus and thickened inland from 1996 to 2015 (Figure 8). Periods of glacier advance (~1970s–80s) at both Storstrømmen and L. Bistrup Bræ preceded the earliest record of elevation change and, following this, their terminus positions underwent minimal change (Figure 5). Between 1996 and 2015, inland elevation change was minimal (Figure 8), whereas greater thinning took place at the terminus. Large retreat events of 2.1 km at Storstrømmen and 0.7 km at L. Bistrup Bræ between 2011 and 2013 coincided with increased terminus thinning rates of -0.8 m a^{-1} at Storstrømmen (2011 to 2012) and -1.76 m a^{-1} at L. Bistrup Bræ (2011 to 2013: Figure 8). These spatial patterns of elevation change were synchronous with velocity variations: deceleration and thickening occurred inland, while acceleration, thinning, and retreat were synchronous at the terminus (Figure 8).








Figure 8: Annual surface elevation change, annual velocity and surface/bed topography for two outlet glaciers in northeast Greenland: Storstrømmen (a) and L. Bistrup Bræ (b). Blue to green coloured lines represent annual surface elevation through time (1992–96 to 2014–15) and yellow through to red lines represent annual winter velocity from 1991/92 to 2015/16.

3.4 Topographic factors

3.4.1 Grounded-terminus outlet glaciers

Distinct variability in glacier geometry exists between individual outlet glaciers in northern Greenland. At many outlet glaciers with grounded termini (Harald Moltke Bræ, Heilprin, Tracy and Humboldt), deep inland sloping beds are ~~correlated~~associated with higher retreat rates (averaging -121 m a^{-1}), and greater increases in velocity (Table 1). These glaciers rest between -33 and -370 m below sea level (Figure 9) and appear to have been retreating down steep bed-slopes (Figure S2) away from topographic ridges at the end of their fjords (Figure 9). However, variability in fjord width (widening/narrowing) does not appear to be a main determinant of higher retreat rates (Table 3). Additionally, some grounded outlets (Harder, Academy, and Marie-Sophie) have shallower seaward sloping topography which correlated with lower magnitude retreat rates (-24 m a^{-1} : Table 1).

Table 3: Glacier-specific factors at 18 northern Greenland study glaciers. This includes: the size and percentage of each surface drainage basin below sea level, the direction of the bed slope 20 km inland of the grounding line (inland or seaward), and whether the fjord widens or narrows with distance inland. Red and blue shading for bed-slope and fjord width represent expected instability, and stability respectively for each parameter.

Northern Greenland Outlet Glaciers		Drainage Basin Size (km ²)	% of drainage basin below sea level	Inland bed-slope	Seaward bed-slope	Widening Fjord	Narrowing Fjord
							
Category 1: Grounded terminus	Tracy	3,176 _b	3.6 _b	X			X
	Kofoed-Hansen Bræ			X		X	
	Harald Moltke Bræ	666	17	X			X
	Humboldt	51,815	27	X		Does not terminate in fjord	
	Heilprin	6,593	2.9	X		X	
	Academy	_a	_a		X	X	
	Harder	792	0.2		X		X
	Marie-Sophie	2,567	6.8		X	X	
	Brikkerne	929	2.3		X		X
Category 2: Floating ice tongue	Petermann	60,093	67	X		X	
	Zachariae Isstrøm	257,542 _b	54	X		X	
	Hagen Bræ	30,250 _a	20		X	X	
	C. H. Ostenfeld	11,013	1.5	X			X
	Nioghalvfjærdsfjorden	_b	_b	X			X
	Steensby	3,356	4.2	X			X
	L. Bistrup Bræ	26,660	4.4	X		X	
	Storstrømmen	_b	_b	X		X	
	Ryder	36,384	40		X	X	




Figure 9: Basal topography from Operation IceBridge BedMachine v3 (Morlighem et al., 2017) beneath nine study glaciers with grounded termini in northern Greenland. Red points represent the position of the terminus/grounding line at each glacier from our most recent record of their terminus position (2015). Black lines are glacier centreline profiles. Profile plots show basal elevations along each glacier centreline profile and solid red lines nearest to zero show the terminus location.




Figure 10: Basal topography from Operation IceBridge BedMachine v3 (Morlighem et al., 2017) beneath nine study glaciers which terminate in floating ice tongues in northern Greenland. Red points represent the most recent recorded terminus position (2015) from this study. Green points represent the location of the grounding line along the centreline profile from the GIMP DEM mask (Howat et al., 2014). Profile plots show basal elevations along each glacier centreline profile, where closest to zero red lines show the terminus locations, and further inland green lines shown the grounding line.

3.4.2 Glaciers with floating ice tongues

Basal topography beneath floating ice tongues does not impact on their dynamic response to retreat. Instead, we examine bed topography inland of the grounding line, which may have implications for grounding line retreat and associated instability once ice tongues collapse. Deeper bed topography (-73 to -1000 m below sea level: Figure 10) exists at floating ice tongues, and most (seven of nine) also have inland sloping topography within 20 km of their grounding lines (Table 3). Unlike grounded outlet glaciers there is no obvious link to higher retreat rates or fjord widths between these glaciers. However, the current grounding line position along glacier bed profiles appear to be ~~correlated-associated~~ with differences in the dynamic response of glaciers to either large calving events or entire ice tongue collapse. Grounding line positions at Petermann, C. H. Ostenfeld and Hagen Bræ, rest on relatively flat topography (Figure 10a,c,d), rather than retrograde slopes. In all cases, either large calving events at Petermann Glacier or entire ice tongue collapse at C. H. Ostenfeld and Hagen Brae, coincided with limited glacier acceleration. In contrast, the grounding lines of glaciers draining the NEGIS (Nioghalvfjærdsfjorden and Zachariae Isstrøm), rest on steeper inland sloping beds (Figure 10), which correlates with their gradual ice tongue retreat and prolonged glacier acceleration. Since losing its ice tongue in 2011/12, Zachariae Isstrøm retreated down its steep basal trough, past the recorded (nominal date of 2007 in BedMachine dataset) grounding line position (Figure 10b). A possible exception to this pattern of grounding line instability on retrograde slopes is Ryder Glacier. It too has a deep basal trough (~800 m below sea level) 20 km inland of the grounding line, but further inland (~50 km from the terminus) it has a steep seaward sloping bed, and a large topographic ridge immediately seaward of the current grounding line position (Figure 10c).

4. Discussion

4.1 Timing of glacier change between 1948 and 2016

Decadal terminus changes at all 18 study glaciers (Figure 4), showed a transition from slow low magnitude advance and retreat (averaging $+72 \text{ m a}^{-1}$) between 1948 and 1995 to rapid high magnitude retreat (averaging -445 m a^{-1}) between 1996 and 2015. The latter period included the onset of steady retreat at most grounded outlet glaciers in northern Greenland, and the occurrence of large, rapid retreat events at floating ice tongue glaciers (Figure 5). While this switch from minimal terminus change/advance to more rapid retreat is perhaps similar to the cyclic behaviour of tidewater glaciers (Meier and Post, 1987; Pfeffer, 2007), it is unlikely that this pattern of widespread retreat is driven by internal factors alone (e.g. Nick et al., 2007). Instead, climate-induced dynamic thinning at outlet glacier margins (Khan et al., 2014; Pritchard et al., 2009), may have been the initial forcing for increased glacier retreat rates in northern Greenland. The switch to terminus retreat in the 1990s was coincident with: increased air and ocean temperatures across the GrIS (e.g. Box et al., 2009; Hanna et al., 2008; Luckman et al., 2006), ice marginal thinning (< 2000 m elevation) since the 1990s (Abdalati et al., 2001; van den Broeke et al., 2016; Krabill et al., 2000), and with Arctic-wide increased retreat rates (Carr et al., 2017b; Moon and Joughin, 2008; Jensen et al., 2016). Previous studies in northern Greenland have suggested that climate-ocean forcing at glacier termini i.e. increased air temperatures, reduced sea

ice concentration, and ocean warming induced basal melt, triggered dynamic thinning, retreat, and mass loss from outlet glacier margins (Khan et al., 2014; Reeh et al., 2001; Rignot et al., 2001; Rignot and Steffen, 2008). Here, we do not assess in detail the climate-ocean forcing mechanisms that may have influenced recent terminus change behaviour in northern Greenland, partly due to lack of data and partly because the focus of this paper is on glacier dynamics and their interaction with topography.

5 However, we note that surface thinning preceded rapid terminus retreat at many northern Greenland glaciers (Figures 6 and 7). Therefore, we suggest that dynamic thinning could have been the initial ~~processforcing~~ that accelerated glacier retreat in northern Greenland since the 1990s (cf. Felikson et al., 2017; Nick et al., 2009; Price et al., 2011). However, in line with tidewater glacier cyclic behaviour, it is likely that after an initial change in dynamics at the terminus triggered by climate forcing, fjord width and depth become more important controls (see Section 4.3) on the duration and magnitude of retreat at

10 individual glaciers (Benn et al., 2007; Catania et al., 2018; MacGregor et al., 2012).

4.2 Dynamic glacier response to terminus change

Our analysis of termini behaviour shows that the dynamic glacier response (acceleration and thinning) to a frontal position change is highly dependent on whether the terminus is grounded or the glacier terminates in a floating ice tongue (Benn et al., 2007). Informed by our changepoint analysis, we ~~observe-infer~~ two dominant calving behaviours in northern Greenland based

15 on these terminus types: 1) low magnitude continuous calving events/terminus retreat at grounded outlet glaciers, 2) large episodic tabular calving events at glaciers with floating ice tongues. Independent of these styles (continuous vs episodic), calving at both categories of terminus type is influenced by the velocity structure of the glacier, and ice velocity itself is sensitive to changes in terminus position and alterations to the force balance, i.e. decreased basal/lateral resistance and increased driving stress (Benn et al., 2007). In the previous section, we hypothesised that increased thinning at the glacier

20 terminus (~1990s), may have initiated enhanced retreat and accelerated terminus velocities in the following two decades (1996 to 2015), similar to other regions of the ice sheet (e.g. Luckman et al., 2006; McFadden et al., 2011; Moon and Joughin, 2008). Such thinning could cause downstream increases in velocity, which stretches the ice, promotes crevasse propagation induced calving, and propagates the dynamic response (i.e. acceleration) up-glacier. However, the coarse resolution of our elevation change datasets limits our ability to deduce the initial cause of accelerated retreat, so instead we focus on the trends in dynamic

25 glacier changes (acceleration and thinning), after the onset of retreat.

Following an assumed initial change in terminus conditions (~1990s), outlet glaciers in northern Greenland that are grounded at their terminus, underwent prolonged periods of steady terminus retreat (on average -150 m a^{-1}), that usually lasted for two to three decades (Figure 5). Like grounded-outlet glaciers elsewhere, e.g. Helheim and Kangerdlugssuaq (Howat et al., 2005,

30 2007, 2008) and in west Greenland (McFadden et al., 2011), periods of steady and continuous retreat at grounded-terminus outlet glaciers in northern Greenland were accompanied by increased annual ice velocities (27-110%), and dynamic thinning (Figure 6). From this dynamic response we suggest that continuous calving and retreat, and the associated reduction in resistive

stresses at the terminus, substantially altered the force balance by increasing longitudinal stretching and driving stress. This prolonged stress perturbation at the terminus of most grounded outlet glaciers in northern Greenland, allowed acceleration and thinning to propagate inland and continue for a longer period as most glaciers may have not reached a stable geometry (McFadden et al., 2011; Nick et al., 2009: Section 4.3).

5

In contrast, terminus changes at most glaciers with floating ice tongues were characterised by short-lived (<6 years), significantly higher magnitude retreat events that averaged -4536 m a^{-1} (after ~1990s). These high magnitude retreat events were often due to the calving of large tabular icebergs, initiated by rift propagation (e.g. MacGregor et al., 2012). However, these large calving events were followed by minimal/and or short-lived increases in annual velocity, and short-term increases in ice surface thinning rates (Figure 7). From these velocity records we infer that in most cases large calving events, appeared not to substantially perturb the glacier force balance by neither increasing longitudinal stretching, nor driving stresses on grounded ice, which would lead to annual acceleration (Figure 7). This was particularly the case at Petermann, Hagen Bræ and C. H. Ostenfeld, in response to ice tongue collapse or large calving events. This contrasts with the behaviour of ice-tongue terminating glaciers elsewhere in Greenland (e.g. Joughin et al., 2004, 2008) and glaciers draining into Antarctic ice shelves (e.g. Scambos et al., 2004), which instead showed prolonged acceleration and dynamic thinning following the loss of substantial parts of floating ice. At some glaciers short-lived acceleration was followed by reduced retreat, and deceleration (e.g. Hagen Bræ), which represents a rapid re-adjustment at the terminus, and that calving at floating ice tongue glaciers in northern Greenland appear to limit the dynamic glacier response to large calving events. This could be due to limited lateral resistance provided by floating ice tongues (Section 4.3).

10

15

20

However, Zachariae Isstrøm was a notable exception to this pattern. At Zachariae Isstrøm, sustained annual calving was accompanied by a longer period of glacier acceleration and thinning (Figure 7b). This is comparable to the behaviour of grounded outlet glaciers in northern Greenland, and ice-tongue terminating glaciers elsewhere (e.g. Jakobshavn Isbræ: Joughin et al., 2004, 2008). In this case, continuous retreat is likely to have gradually reduced resistive forces (i.e. backstress) acting on inland grounded ice, causing higher magnitude and prolonged flow acceleration. Apart from Zachariae Isstrøm, our data show outlet glaciers in northern Greenland have been largely insensitive to either entire ice tongue loss (C. H. Ostenfeld, Steensby and Hagen Bræ), or large iceberg calving events (Petermann, Nioghalvfjærdsfjorden). Thus, despite some similarities (e.g. Zachariae Isstrøm to grounded-behaviour), region wide glacier behaviour in northern Greenland appears dependent on whether the terminus is grounded or floating, due to their calving nature and dynamic response to perturbations of their termini. This highlights the need to consider terminus type when assessing the long-term response of outlet glaciers to changes at their terminus.

25

30

4.3 Influence of glacier geometry

While climate-ocean forcing may have triggered a change in glacier dynamics at the terminus of outlet glaciers in northern Greenland (e.g. Khan et al., 2014; Reeh et al., 2001), glacier geometry (e.g. width and depth of fjords) may have determined the duration and extent of the resultant retreat. Indeed, variations in basal topography and fjord width have been previously identified as an important control on the dynamic response of glaciers in many regions of the GrIS (e.g. Carr et al., 2013, 2017b; Howat and Eddy, 2011; McFadden et al., 2011; Millan et al., 2018; Thomas et al., 2009). Collectively these factors could explain differences between grounded-terminus and floating ice-tongue glaciers (McFadden et al., 2011), as well as individual glacier variability.

- Calving from grounded outlet margins is controlled by both basal and lateral drag, and both reduce as a glacier retreats into a deeper and wider fjord (Benn et al., 2007). At grounded outlet glaciers in northern Greenland, prolonged acceleration and thinning following retreat suggests that these glaciers were still adjusting to terminus change by the end of the study period in 2015. This is likely due to deep basal topography (> 200 m below sea level), and retrograde bed slopes (~ 20 km of their grounding zones) beneath most grounded-terminus glaciers (Figure 9). We suggest grounded-terminus retreat into deeper water contributed to: (i) buoyancy driven feedbacks, as the ice thinned to flotation (van der Veen, 1996), (ii) the penetration of basal crevasses through the full ice thickness (van der Veen, 1998, 2007), and (iii) subsequent enhanced rates of calving and continued retreat (e.g. Joughin et al., 2008). Our results showed grounded-outlet glaciers which retreated into deeper fjords, had higher retreat rates (e.g. Tracy, Harald Moltke Bræ, and Heilprin), than those with shallower basal troughs (e.g. Academy and Marie-Sophie). The former three glaciers also appear to be retreating downslope from topographic highs at the edge of their fjords (Figure 9a,c,e).

- Unlike grounded-termini, floating ice tongues predominantly provide resistive stresses through their contact with the lateral fjord margins. Consequently, lateral resistive stresses are the main control on the glacier force balance and driving stresses, and hence the impact of terminus retreat on inland ice dynamics. Our data have shown variability in glacier response to ice tongues loss (Figure 7), and we suggest that this could be due to differences in the lateral resistive the floating ice tongue provides when it is in place. Once the ice tongue has entirely collapsed, the terminus becomes grounded, at which point basal drag becomes an important control, and basal topography at and immediately inland of the grounding line becomes more significant.

- At most glaciers with floating ice tongues in northern Greenland, the minimal dynamic response to ice tongue retreat and/or collapse (Figure 7), may be due to limited lateral resistance provided by their floating ice tongues. In particular, C. H. Ostfeld and Hagen Bræ, have heavy rifting along their shear margins, appear relatively un-confined by their fjord walls, and weakly attached to the grounded terminus (Figure 11b,c). Indeed, both glaciers showed no significant increase in flow speeds following

large calving events. This suggests that, in both cases, the buttressing provide by the tongues was minimal, and large ice tongue retreats caused a limited change in the inland force balance. Alternatively, Steensby Glacier showed some acceleration ($\sim 25\%$) following ice tongue retreat, which could be due to both a greater loss of lateral resistive stresses from a well-confined ice tongue, and retreat past a narrower sinuous section of the fjord (Figure 11a).




Figure 11: Landsat imagery of three glaciers which terminate in floating ice tongues in northern Greenland before their ice tongue collapse. (a) Steensby Glacier in 2013, (b) C. H. Ostenfeld Glacier in 2002, (c) Hagen Bræ in 2005. Purple lines denote the location of the grounding line.

As well as the lack of resistive stress provided by their ice tongues, the limited response of Hagen Bræ and C. H. Ostenfeld to terminus retreat (Figure 7) may result from their basal topography: following retreat, both grounding lines retreated into shallow water (Figure 10). This may have suppressed retreat rates, as it reduces grounding line thickness and therefore discharge. In turn, this would reduce the impact on inland ice velocities and surface thinning rates (Vieli and Nick, 2011). The flat sections of basal topography beneath the grounding lines of Petermann Glacier and Nioghalvfjærdsfjorden may also control their future response to ice tongue collapse, as their grounding lines would need to retreat ~ 20 km inland to sit on a retrograde slope (Figure 8b, g). In contrast, ice tongue collapse at Zachariae Isstrøm, was followed by continued acceleration, retreat, and more dramatic thinning (Figure 7b). Here, the deep retrograde bed-slope that extends ~ 20 km inland of the grounding line, is likely responsible for continued retreat (Khan et al., 2014; Mouginot et al., 2015). Retreat into deeper water, gradually reduced buttressing forces, and caused continuous glacier acceleration and surface thinning following ice tongue collapse, similar to Jakobshavn Isbræ (Vieli and Nick, 2011).

4.4 Glacier surging

Surge-type behaviour has been previously documented at several outlet glaciers in northern Greenland (e.g. Hill et al., 2017; Rignot et al., 2001; Weidick et al., 1994), but detailed evidence for surging is rare. Here we briefly discuss how the surge-nature of several glaciers in northern Greenland may have altered their dynamic behaviour independent of their terminus type.

Our results provide further evidence for the presence of three surge-type glaciers in northern Greenland (Storstrømmen, L. Bistrup Bræ, and Harald Moltke Bræ). This is based on the following characteristics: 1) substantial periods of glacier advance ($> 90 \text{ m a}^{-1}$) followed by retreat during the study period, 2) accelerated ice flow coincident with periods of advance, and 3) surface thickening inland and thinning at the terminus position indicative of a quiescent surge-phase. Both Storstrømmen and L. Bistrup Bræ have floating ice tongues which began advancing in the 1970s (Reeh et al., 1999), and continued until 1985 at Storstrømmen, and 1998 at L. Bistrup Bræ (Figure 5). While velocity and surface elevation change datasets do not cover this period, dynamic changes later on (1992 to 2016: Figure 8) were indicative of quiescence (Abdalati et al., 2001; Csatho et al., 2014; Thomas et al., 2009). In this respect, these glaciers have behaved differently to the majority of glaciers in the region with floating ice tongues i.e. thickening and deceleration inland, and thinning and acceleration at the terminus (Figure 8). In northwest Greenland, Harald Moltke Bræ has been previously considered surge-type. Here we record an additional surge event from 2013 to 2014, based on high magnitude acceleration ($\sim 1000 \text{ m a}^{-1}$) and glacier advance (0.8 km: Figure 6c). This glacier fits the conventional definition of surging, i.e. a short active phase, which included an order of magnitude increase in velocity. However, it has a short surge-cycle (< 10 years) compared to most other glaciers in the Arctic, and underwent overall retreat from the late 1980s to 2015 (Figure 5). This suggests that despite short-lived surge events, recent climate-ocean forcing may be altering its cyclical behaviour.

Finally, we draw attention to Ryder Glacier as it appears to be behaving dissimilarly to the rest of our study glaciers in northern Greenland. It too has ~~also~~ been referred to as surge-type in the past (Joughin et al., 1996, 1999; Rignot et al., 2001), and has shown some surge-like behaviour during our study period (1948–2015): several cycles of advance (~ 7 -years) and retreat (2-years), and some acceleration during advance. Additionally, previous studies also identified near-terminus thinning ($2\text{--}4 \text{ m a}^{-1}$: 1997 to 1999) and, at $\sim 50 \text{ km}$ inland, a similar magnitude of thickening (Abdalati et al., 2001), which is indicative of the quiescent phase of surge-type glaciers (e.g. Kamb et al., 1985; Meier and Post, 1969; Sharp, 1988). However, the behaviour of Ryder Glacier appears to be more characteristic of a tidewater glacier cycle, which may be controlled by basal topography. In contrast to the role of basal topography at most other outlet glaciers in northern Greenland (i.e. unstable retreat down inland sloping beds), a large basal ridge (Figure 10i) or terminal moraine/moraine shoal at Ryder Glacier may have promoted periods of glacier advance (Alley, 1991; Nick et al., 2007; Powell, 1990). Nevertheless, further investigation on the cyclic nature and precise controls on Ryder Glaciers' dynamic behaviour is needed.

5. Conclusions

Outlet glaciers in northern Greenland drain $\sim 40\%$ of the ice sheet by area but remain understudied compared to other regions of the ice sheet. We have analysed the dynamics of 18 major marine-terminating outlet glaciers in northern Greenland between 1948 and 2015. Overall, long-term glacier retreat rates ranged from -15 to -311 m a^{-1} over the entire study period. Between

1948 and 1995 glaciers exhibited generally low magnitude advance and retreat, with an average frontal position change of +72 m a⁻¹ (advance) across the 18 study glaciers. Following this, there was a regional transition to more rapid and widespread retreat, when average frontal position change was -445 m a⁻¹ (1995 to 2015). This was coincident with accelerated retreat in other regions of the ice sheet (e.g. Carr et al., 2013; Howat et al., 2008; Howat and Eddy, 2011). From 1996 to 2015, most glaciers also experienced accelerated ice flow and increased dynamic thinning.

While increased retreat rates from the mid-1990s were near-ubiquitous, we observe distinct differences in glacier behaviour depending on whether the terminus is grounded or floating. Three factors play a role in the dynamic behaviour of these two types of glacier; i) different methods of calving (i.e. continuous small magnitude calving vs large episodic calving); ii) differences in resistive stresses at the terminus; iii) glacier geometry. Continuous retreat into deep fjords at grounded-terminus glaciers led to a greater reduction in basal/lateral resistive stresses, and caused high magnitude acceleration and dynamic thinning. In contrast, large episodic calving events, from unconfined ice tongues that provided little lateral resistance meant that most glaciers with floating ice tongues appear dynamically insensitive to the retreat of their terminus. We note there are exceptions; continuous ice tongue retreat at Zachariae Isstrøm caused prolonged acceleration and thinning, and several glaciers with ice tongues went through cycles of advance and retreat during the study record (e.g. Ryder Glacier). At Zachariae Isstrøm this can be explained by the method of glacier calving (continuous rather than episodic), and a deep wide fjord that promoted unstable retreat. Glacier advance can be explained by surging, or topographic controls which allow cyclic advance and retreat. We provide further evidence for surging at three glaciers (Harald Moltke Bræ, Storstrømmen and L. Bistrup Bræ) in northern Greenland, and an explanation for the cyclic behaviour of Ryder Glacier, which is likely related to topographic controls (e.g. moraine shoal), that allowed the re-advance of the terminus. While we have shown that northern Greenland has begun to undergo rapid dynamic change over the last two decades (1996 to 2015), we highlight variability between individual glaciers and the importance of considering terminus type and glacier geometry (basal topography, fjord width and ice tongue confinement) when considering future glacier response to climate change across this region of the ice sheet. Currently, ice tongue retreat does not appear to substantially affect inland ice dynamics. However, once these glaciers become grounded, they may accelerate, thin, and increase the volume of grounded ice discharge into the ocean.

Data availability

Shapefiles of frontal positions for all 18 outlet glaciers in this study between 1948 and 2015 are freely available on request to the corresponding author. All other data sources, including: satellite imagery, historical maps, surface elevation change, annual velocity, climate and ocean, and topographic data, are already available online. The sources of each of these datasets are given in the text and the supplementary information.

Author contribution

The initial project was designed by all authors, and E. A. Hill led the data analysis and interpretation, with comments throughout from all authors. E. A. Hill led the manuscript writing, and all authors contributed towards the editing of the manuscript and figures.

10 Competing interests

The authors declare that they have no conflict of interest.

Acknowledgements

This work was supported by a Doctoral Studentship award to E. A. Hill at Newcastle University, UK, from the IAPETUS Natural Environment Research Council Doctoral Training Partnership (grant number: NE/L002590/1). We thank the ESA for granting access to SPOT-1 data (project ID: 32435). We are also grateful for Corona imagery which was digitised on request to the US Geological Survey. We acknowledge several free datasets used in this work. Historical map charts were made available from the Polar Geospatial Centre and the Perry-Castañeda Library at the University of Texas, Austin. We are grateful for Cryosat and ESA surface elevation change data, and Sentinel annual Greenland wide velocity maps made freely available as part of the ESA Greenland Climate Change Initiative. MEaSURES velocity maps (Joughin et al., 2010), IceBridge BedMachine v3 (Morlighem et al., 2017), is available from the National Snow and Ice Data Centre (NSIDC). Landsat imagery were acquired from the US Geological Survey. We are extremely grateful to the Editor of this manuscript - Andreas Vieli - and to three anonymous reviews whose comments led to improvements in both the content and clarity.

References

- Abdalati, W., Krabill, W., Frederick, E., Manizade, S., Martin, C., Sonntag, J., Swift, R., Thomas, R., Wright, W. and Yungel, J.: Outlet glacier and margin elevation changes: Near-coastal thinning of the Greenland ice sheet, *J. Geophys. Res.*, 106(D24), 33729, doi:10.1029/2001JD900192, 2001.
- 5 Alley, R. B.: Sedimentary processes may cause fluctuations of tidewater glaciers, *Ann. Glaciol.*, 15, 119–124, 1991.
- Amundson, J. M., Fahnestock, M., Truffer, M., Brown, J., Lüthi, M. P. and Motyka, R. J.: Ice mélange dynamics and implications for terminus stability, Jakobshavn Isbræ, Greenland, *J. Geophys. Res. Earth Surf.*, 115(1), 1–12, doi:10.1029/2009JF001405, 2010.
- Benn, D. I., Warren, C. R. and Mottram, R. H.: Calving processes and the dynamics of calving glaciers, *Earth-Science Rev.*, 10
82(3–4), 143–179, doi:10.1016/j.earscirev.2007.02.002, 2007.
- Bevan, S. L., Luckman, A. J. and Murray, T.: Glacier dynamics over the last quarter of a century at Helheim, Kangerdlugssuaq and 14 other major Greenland outlet glaciers, *Cryosphere*, 6(5), 923–937, doi:10.5194/tc-6-923-2012, 2012.
- Box, J. E. and Decker, D. T.: Greenland marine-terminating glacier area changes: 2000–2010, *Ann. Glaciol.*, 52(59), 91–98,
15 doi:10.3189/172756411799096312, 2011.
- Box, J. E., Yang, L., Bromwich, D. H. and Bai, L.-S. L.-S.: Greenland Ice Sheet Surface Air Temperature Variability: 1840–2007, *J. Clim.*, 22(14), 4029–4049, doi:10.1175/2009jcli2816.1, 2009.
- van den Broeke, M. R., Enderlin, E. M., Howat, I. M., Kuipers Munneke, P., Noël, B. P. Y., Jan Van De Berg, W., Van Meijgaard, E. and Wouters, B.: On the recent contribution of the Greenland ice sheet to sea level change, *Cryosphere*, 10(5),
20 1933–1946, doi:10.5194/tc-10-1933-2016, 2016.
- Bunce, C., Carr, J. R., Nienow, P. W., Ross, N. and Killick, R.: Ice front change of marine-terminating outlet glaciers in northwest and southeast Greenland during the 21st century, *J. Glaciol.*, 1–13, doi:10.1017/jog.2018.44, 2018.
- Carr, J. R., Vieli, A. and Stokes, C.: Influence of sea ice decline, atmospheric warming, and glacier width on marine-terminating outlet glacier behavior in northwest Greenland at seasonal to interannual timescales, *J. Geophys. Res. Earth
25 Surf.*, 118(3), 1210–1226, doi:10.1002/jgrf.20088, 2013.
- Carr, J. R., Stokes, C. and Vieli, A.: Recent retreat of major outlet glaciers on Novaya Zemlya, Russian Arctic, influenced by fjord geometry and sea-ice conditions, *J. Glaciol.*, 60(219), 155–170, doi:10.3189/2014JoG13J122, 2014.
- Carr, J. R., Vieli, A., Stokes, C. R., Jamieson, S. S. R., Palmer, S. J., Christoffersen, P., Dowdeswell, J. A., Nick, F. M., Blankenship, D. D. and Young, D. A.: Basal topographic controls on rapid retreat of Humboldt Glacier, northern Greenland,
30 *J. Glaciol.*, 61(225), 137–150, doi:10.3189/2015JoG14J128, 2015.
- Carr, J. R., Bell, H., Killick, R. and Holt, T.: Exceptional retreat of Novaya Zemlya’s marine-terminating outlet glaciers between 2000 and 2013, *Cryosphere*, 11(5), 2149–2174, doi:10.5194/tc-11-2149-2017, 2017a.

- Carr, J. R., Stokes, C. R. and Vieli, A.: Threefold increase in marine-terminating outlet glacier retreat rates across the Atlantic Arctic: 1992–2010, in *Annals of Glaciology*, vol. 58, pp. 72–91., 2017b.
- Catania, G. A., Stearns, L. A., Sutherland, D. A., Fried, M. J., Bartholomaeus, T. C., Morlighem, M., Shroyer, E. and Nash, J.: Geometric Controls on Tidewater Glacier Retreat in Central Western Greenland, *JGR Earth Sci.*,
5 doi:10.1029/2017JF004499, 2018.
- Church, J. a., Clark, P. U., Cazenave, a., Gregory, J. M., Jevrejeva, S., Levermann, a., Merrifield, M. a., Milne, G. a., Nerem, R. ., Nunn, P. D., Payne, a. J., Pfeffer, W. T., Stammer, D. and Unnikrishnan, a. S.: Sea level change, *Clim. Chang.* 2013 Phys. Sci. Basis. Contrib. Work. Gr. I to Fifth Assess. Rep. Intergov. Panel Clim. Chang., 1137–1216, doi:10.1017/CB09781107415315.026, 2013.
- 10 Cook, S., Rutt, I. C., Murray, T., Luckman, A., Zwinger, T., Selmes, N., Goldsack, A. and James, T. D.: Modelling environmental influences on calving at Helheim Glacier in eastern Greenland, *Cryosphere*, 8(3), 827–841, doi:10.5194/tc-8-827-2014, 2014.
- Csatho, B. M., Schenk, A. F., van der Veen, C. J., Babonis, G., Duncan, K., Rezvanbehhahani, S., van den Broeke, M. R., Simonsen, S. B., Nagarajan, S. and van Angelen, J. H.: Laser altimetry reveals complex pattern of Greenland Ice Sheet
15 dynamics., *Proc. Natl. Acad. Sci. U. S. A.*, 111(52), 18478–83, doi:10.1073/pnas.1411680112, 2014.
- Dowdeswell, A., Hamilton, G. S. and Hagen, O. V. E.: The duration of the active phase on surge-type glaciers: contrasts between Svalbard and other regions, *J. Glaciol.*, 37(127), 388–400, doi:10.3189/S0022143000005827, 1991.
- Enderlin, E. M., Howat, I. M. and Vieli, A.: High sensitivity of tidewater outlet glacier dynamics to shape, *Cryosphere*, 7(3), 1007–1015, doi:10.5194/tc-7-1007-2013, 2013.
- 20 Enderlin, E. M., Howat, I. M., Jeong, S., Noh, M. J., Van Angelen, J. H. and Van Den Broeke, M. R.: An improved mass budget for the Greenland ice sheet, *Geophys. Res. Lett.*, 41(3), 866–872, doi:10.1002/2013GL059010, 2014.
- Felikson, D., Bartholomaeus, T. C., Catania, G. A., Korsgaard, N. J., Kjær, K. H., Morlighem, M., Noël, B., Van Den Broeke, M., Stearns, L. A., Shroyer, E. L., Sutherland, D. A. and Nash, J. D.: Inland thinning on the Greenland ice sheet controlled by outlet glacier geometry, *Nat. Geosci.*, 10(5), 366–369, doi:10.1038/ngeo2934, 2017.
- 25 Fettweis, X., Franco, B., Tedesco, M., van Angelen, J. H., Lenaerts, J. T. M., van den Broeke, M. R. and Gallée, H.: Estimating Greenland ice sheet surface mass balance contribution to future sea level rise using the regional atmospheric climate model MAR, *Cryosph.*, 7, 469–489, doi:10.5194/tcd-6-3101-2012, 2012.
- Franco, B., Fettweis, X., Erpicum, M. and Nicolay, S.: Present and future climates of the Greenland ice sheet according to the IPCC AR4 models, *Clim. Dyn.*, 36(9–10), 1897–1918, doi:10.1007/s00382-010-0779-1, 2011.
- 30 Hanna, E., Huybrechts, P., Steffen, K., Cappelen, J., Huff, R., Shuman, C., Irvine-Fynn, T., Wise, S. and Griffiths, M.: Increased runoff from melt from the Greenland Ice Sheet: A response to global warming, *J. Clim.*, 21(2), 331–341, doi:10.1175/2007JCLI1964.1, 2008.
- Hill, E. A., Carr, J. R. and Stokes, C. R.: A Review of Recent Changes in Major Marine-Terminating Outlet Glaciers in

- Northern Greenland, *Front. Earth Sci.*, 4(111), 1–23, doi:10.3389/feart.2016.00111, 2017.
- Hill, E. A., Carr, J. R., Stokes, C. R. and Gudmundsson, G. H.: Dynamic changes in outlet glaciers in northern Greenland from 1948 to 2015, *Cryosph. Discuss.*, in review, doi:10.5194/tc-2018-17, 2018.
- Holland, D. M., Thomas, R. H., de Young, B., Ribergaard, M. H. and Lyberth, B.: Acceleration of Jakobshavn Isbræ triggered by warm subsurface ocean waters, *Nat. Geosci.*, 1(10), 659–664, doi:10.1038/ngeo316, 2008.
- Howat, I. M. and Eddy, A.: Multi-decadal retreat of Greenland’s marine-terminating glaciers, *J. Glaciol.*, 57(203), 389–396, doi:10.3189/002214311796905631, 2011.
- Howat, I. M., Joughin, I., Tulaczyk, S. and Gogineni, S.: Rapid retreat and acceleration of Helheim Glacier, east Greenland, *Geophys. Res. Lett.*, 32(22), 1–4, doi:10.1029/2005GL024737, 2005.
- Howat, I. M., Joughin, I. and Scambos, T. A.: Rapid changes in ice discharge from Greenland outlet glaciers., *Science*, 315(5818), 1559–61, doi:10.1126/science.1138478, 2007.
- Howat, I. M., Joughin, I., Fahnestock, M., Smith, B. E. and Scambos, T. A.: Synchronous retreat and acceleration of southeast Greenland outlet glaciers 2000–06: Ice dynamics and coupling to climate, *J. Glaciol.*, 54(187), 646–660, doi:10.3189/002214308786570908, 2008.
- Howat, I. M., Negrete, A. and Smith, B. E.: The Greenland Ice Mapping Project (GIMP) land classification and surface elevation data sets, *Cryosphere*, 8(4), 1509–1518, doi:10.5194/tc-8-1509-2014, 2014.
- Jensen, T. S., Box, J. E. and Hvidberg, C. S.: A sensitivity study of annual area change for Greenland ice sheet marine terminating outlet glaciers : 1999 – 2013, *J. Glaciol.*, 62(231), 72–81, doi:10.1017/jog.2016.12, 2016.
- Joughin, I., Tulaczyk, S., Fahnestock, M. and Kwok, R.: A mini-surge on the Ryder Glacier, Greenland, observed by satellite radar interferometry, *Science (80-.)*, 274(5285), 228–230, doi:10.1126/science.274.5285.228, 1996.
- Joughin, I., Fahnestock, M., Kwok, R., Gogineni, P. and Allen, C.: Ice flow of Humboldt, Petermann and Ryder Gletscher, northern Greenland, *J. Glaciol.*, 45(150), 231–241, doi:10.3189/002214399793377284, 1999.
- Joughin, I., Abdalati, W. and Fahnestock, M. a.: Large fluctuations in speed on Greenland’s Jakobshavn Isbrae glacier, *Nature*, 432, 608–610, doi:10.1038/nature03130, 2004.
- Joughin, I., Howat, I. M., Fahnestock, M., Smith, B., Krabill, W., Alley, R. B., Stern, H. and Truffer, M.: Continued evolution of Jakobshavn Isbrae following its rapid speedup, *J. Geophys. Res. Earth Surf.*, 113(4), 1–14, doi:10.1029/2008JF001023, 2008.
- Joughin, I., Smith, B. E., Howat, I. M., Scambos, T. and Moon, T.: Greenland flow variability from ice-sheet-wide velocity mapping, *J. Glaciol.*, 56(197), 415–430, doi:10.3189/002214310792447734, 2010.
- Kamb, B., Raymond, C. F., Harrison, W. D., Engelhardt, H., Echelmeyer, K. A., Humphrey, N., Brugman, M. M. and Pfeffer, T.: Glacier surge mechanism: 1982–1983 surge of Variegated glacier, Alaska, *Science (80-.)*, 227(4686), 469–479, doi:10.1126/science.227.4686.469, 1985.
- Khan, S. A., Aschwanden, A., Bjørk, A. A., Wahr, J., Kjeldsen, K. K. and Kjaer, K. H.: Greenland ice sheet mass balance: a

- review, *Reports Prog. Phys.*, 046801, 1–26, doi:10.1088/0034-4885/78/4/046801, 2015.
- Khan, S. a, Kjaer, K. H., Bevis, M., Bamber, J. L., Wahr, J., Kjeldsen, K. K., Bjork, A. a, Korsgaard, N. J., Stearns, L. a, van den Broeke, M. R., Liu, L., Larsen, N. K. and Muresan, I. S.: Sustained mass loss of the northeast Greenland ice sheet triggered by regional warming, *Nat. Clim. Chang.*, 4(4), 292–299, doi:10.1038/nclimate2161, 2014.
- 5 Khvorostovsky, K. S.: Merging and Analysis of Elevation Time Series Over Greenland Ice Sheet From Satellite Radar Altimetry, *IEEE Trans. Geosci. Remote Sens.*, 50(1), 23–36, 2012.
- Killick, R., Fearnhead, P. and Eckley, I. A.: Optimal detection of changepoints with a linear computational cost, *J. Am. Stat. Assoc.*, 107(500), 1590–1598, doi:10.1080/01621459.2012.737745, 2012.
- Kjeldsen, K. K., Korsgaard, N. J., Bjørk, A. A., Khan, S. A., Box, J. E., Funder, S., Larsen, N. K., Bamber, J. L., Colgan, W.,
 10 van den Broeke, M., Siggaard-Andersen, M.-L., Nuth, C., Schomacker, A., Andresen, C. S., Willerslev, E. and Kjær, K. H.: Spatial and temporal distribution of mass loss from the Greenland Ice Sheet since AD 1900, *Nature*, 528(7582), 396–400, doi:10.1038/nature16183, 2015.
- Korsgaard, N. J., Nuth, C., Khan, S. A., Kjeldsen, K. K., Bjørk, A. A., Schomacker, A. and Kjær, K. H.: Digital elevation model and orthophotographs of Greenland based on aerial photographs from 1978-1987, *Sci. Data*, 3, 1–15,
 15 doi:10.1038/sdata.2016.32, 2016.
- Krabill, W., Abdalati, W., Frederick, E., Manizade, S., Martin, C., Sonntag, J., Swift, R., Thomas, R., Wright, W. and Yungel, J.: Greenland Ice Sheet: High-elevation balance and peripheral thinning, *Science* (80-.), 289(5478), 428–430, doi:10.1126/science.289.5478.428, 2000.
- Krabill, W., Hanna, E., Huybrechts, P., Abdalati, W., Cappelen, J., Csatho, B., Frederick, E., Manizade, S., Martin, C.,
 20 Sonntag, J., Swift, R., Thomas, R. and Yungel, J.: Greenland Ice Sheet: Increased coastal thinning, *Geophys. Res. Lett.*, 31(24), 1–4, doi:10.1029/2004GL021533, 2004.
- Lavielle, M.: Using penalized contrasts for the change-point problem, *Signal Processing*, 85(8), 1501–1510, doi:10.1016/j.sigpro.2005.01.012, 2005.
- Lea, J. M., Mair, D. W. F. and Rea, B. R.: Instruments and Methods :Evaluation of existing and new methods of tracking
 25 glacier terminus change, *J. Glaciol.*, 60(220), 323–332, doi:10.3189/2014JoG13J061, 2014.
- Luckman, A., Murray, T., de Lange, R. and Hanna, E.: Rapid and synchronous ice-dynamic changes in East Greenland, *Geophys. Res. Lett.*, 33(3), 1–4, doi:10.1029/2005GL025428, 2006.
- MacGregor, J. A., Catania, G. A., Markowski, M. S. and Andrews, A. G.: Widespread rifting and retreat of ice-shelf margins in the eastern Amundsen Sea Embayment between 1972 and 2011, *J. Glaciol.*, 58(209), 458–466,
 30 doi:10.3189/2012JoG11J262, 2012.
- McFadden, E. M., Howat, I. M., Joughin, I., Smith, B. E. and Ahn, Y.: Changes in the dynamics of marine terminating outlet glaciers in west Greenland (2000-2009), *J. Geophys. Res. Earth Surf.*, 116(2), 1–16, doi:10.1029/2010JF001757, 2011.
- Meier, M. F. and Post, A.: What are glacier surges?, *Can. J. Earth Sci.*, 6(4), 807–817, doi:10.1139/e69-081, 1969.

- Meier, M. F. and Post, A.: Fast tidewater glaciers, *J. Geophys. Res.*, 92(B9), 9051, doi:10.1029/JB092iB09p09051, 1987.
- Mernild, S. H., Liston, G. E., Hiemstra, C. a and Christensen, J. H.: Greenland Ice Sheet Surface Mass-Balance Modeling in a 131-Yr Perspective, 1950-2080, *J. Hydrometeorol.*, 11, 3–25, doi:10.1175/2009JHM1140.1, 2010.
- Millan, R., Rignot, E., Mouginot, J., Wood, M., Bjørk, A. A. and Morlighem, M.: Vulnerability of Southeast Greenland
5 Glaciers to Warm Atlantic Water From Operation IceBridge and Ocean Melting Greenland Data, *Geophys. Res. Lett.*, 45(6), 2688–2696, doi:10.1002/2017GL076561, 2018.
- Moon, T. and Joughin, I.: Changes in ice front position on Greenland’s outlet glaciers from 1992 to 2007, *J. Geophys. Res. Earth Surf.*, 113(2), 1–10, doi:10.1029/2007JF000927, 2008.
- Moon, T., Joughin, I., Smith, B. and Howat, I.: 21st-Century Evolution of Greenland Outlet Glacier Velocities, *Science* (80-
10), 336(6081), 576–578, doi:10.1126/science.1219985, 2012.
- Morlighem, M., Rignot, E., Mouginot, J., Seroussi, H. and Larour, E.: Deeply incised submarine glacial valleys beneath the Greenland ice sheet, *Nat. Geosci.*, 7(6), 18–22, doi:10.1038/ngeo2167, 2014.
- Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R.,
15 Mayer, L., Mouginot, J., Noël, B. P. Y., O’Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M. and Zinglensen, K. B.: BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of Greenland From Multibeam Echo Sounding Combined With Mass Conservation, *Geophys. Res. Lett.*, 44(21), 11,051–11,061, doi:10.1002/2017GL074954, 2017.
- Mouginot, J., Rignot, E., Scheuchl, B., Fenty, I., Khazendar, A., Morlighem, M., Buzzi, A. and Paden, J.: Fast retreat of
20 Zachariæ Isstrøm, northeast Greenland, *Science* (80-), 350, 1357–1361, doi:10.1126/science.aac7111, 2015.
- Murray, T., Scharrer, K., Selmes, N., Booth, A. D., James, T. D., Bevan, S. L., Bradley, J., Cook, S., Cordero Lana, L., Drocourt, Y., Dyke, L. M., Goldsack, A., Hughes, A. L., Luckman, A. J. and McGowan, J.: Extensive retreat of Greenland tidewater glaciers, 2000-2010, *Arctic, Antarct. Alp. Res.*, 47(3), 1–36, doi:10.1657/AAAR0014-049, 2015.
- Nagler, T., Rott, H., Hetzenecker, M., Wuite, J. and Potin, P.: The Sentinel-1 Mission: New Opportunities for Ice Sheet
25 Observations, *Remote Sens.*, 7, 9371–9389, doi:10.3390/rs70709371, 2015.
- Nagler, T., Hauglund, K., Forsberg, R. and Engdahl, M.: Product User Guide (PUG). [online] Available from: <http://www.esa-icesheets-cci.org/%0Ahttp://maps.elie.ucl.ac.be/CCI/viewer/download/ESACCI-LC-PUG-v2.5.pdf>, 2016.
- Nick, F. M., van der Veen, C. J. and Oerlemans, J.: Controls on advance of tidewater glaciers: Results from numerical modeling applied to Columbia Glacier, *J. Geophys. Res. Earth Surf.*, 112(3), 1–11, doi:10.1029/2006JF000551, 2007.
- 30 Nick, F. M., Vieli, A., Howat, I. M. and Joughin, I.: Large-scale changes in Greenland outlet glacier dynamics triggered at the terminus., *Nat. Geosci.*, 2(2), 110–114, doi:10.1038/ngeo394, 2009.
- Nick, F. M., Luckman, A., Vieli, A., Van Der Veen, C. J., Van As, D., Van De Wal, R. S. W., Pattyn, F., Hubbard, A. L. and Floricioiu, D.: The response of Petermann Glacier, Greenland, to large calving events, and its future stability in the context

- of atmospheric and oceanic warming, *J. Glaciol.*, 58(208), 229–239, doi:10.3189/2012JoG11J242, 2012.
- Pfeffer, W. T.: A simple mechanism for irreversible tidewater glacier retreat, *J. Geophys. Res. Earth Surf.*, 112(3), 1–12, doi:10.1029/2006JF000590, 2007.
- Post, A.: Preliminary hydrography and historic terminal changes of Columbia Glacier, Alaska, *US Geol. Surv. Hydrol. Atlas*, 5 559, 1975.
- Post, A., O’Neel, S., Motyka, R. J. and Streveler, G.: A complex relationship between calving glaciers and climate, *Eos* (Washington. DC), 92(37), 305–307, doi:10.1029/2011EO370001, 2011.
- Powell, R. D.: Glacimarine processes at grounding-line fans and their growth to ice-contact deltas, *Geol. Soc. London, Spec. Publ.*, 53(1), 53–73, doi:10.1144/GSL.SP.1990.053.01.03, 1990.
- 10 Price, S. F., Payne, A. J., Howat, I. M. and Smith, B. E.: Committed sea-level rise for the next century from Greenland ice sheet dynamics during the past decade., *Proc. Natl. Acad. Sci. U. S. A.*, 108(22), 8978–83, doi:10.1073/pnas.1017313108, 2011.
- Pritchard, H. D., Arthern, R. J., Vaughan, D. G. and Edwards, L. A.: Extensive dynamic thinning on the margins of the Greenland and Antarctic ice sheets, *Nature*, 461(7266), 971–975, doi:10.1038/nature08471, 2009.
- 15 Reeh, N., Mayer, C. and Miller, H.: Present and past climate control on fjord glaciations in Greenland: Implications for IRD-deposition in the sea, *Geophys. Res. Lett.*, 26(8), 1039–1042, doi:10.1029/1999GL900065, 1999.
- Reeh, N., Thomsen, H. H., Higgins, A. K. and Weidick, A.: Sea ice and the stability of north and northeast Greenland floating glaciers, *Ann. Glaciol.*, 33, 474–480, doi:10.3189/172756401781818554, 2001.
- Rignot, E. and Kanagaratnam, P.: Changes in the velocity structure of the Greenland Ice Sheet, *Science* (80-.), 311(5763), 20 986–990, doi:10.1126/science.1121381, 2006.
- Rignot, E. and Steffen, K.: Channelized bottom melting and stability of floating ice shelves, *Geophys. Res. Lett.*, 35(2), 2–6, doi:10.1029/2007GL031765, 2008.
- Rignot, E., Gogineni, S., Joughin, I. and Krabill, W.: Contribution to the glaciology of northern Greenland from satellite radar interferometry, *J. Geophys. Res. Atmos.*, 106(D24), 34,007–34,019, doi:10.1029/2001JD900071, 2001.
- 25 Rignot, E., Box, J. E., Burgess, E. and Hanna, E.: Mass balance of the Greenland ice sheet from 1958 to 2007, *Geophys. Res. Lett.*, 35(20), 1–5, doi:10.1029/2008GL035417, 2008.
- Rignot, E. J., Gogineni, S. P., Krabill, W. B. and S, E.: North and Northeast Greenland Ice Discharge from Satellite Radar Interferometry, *Science* (80-.), 276(5314), 934–937, doi:10.1126/science.276.5314.934, 1997.
- Scambos, T. A., Bohlander, J. A., Shuman, C. A. and Skvarca, P.: Glacier acceleration and thinning after ice shelf collapse 30 in the Larsen B embayment, Antarctica, *Geophys. Res. Lett.*, 31(18), 2001–2004, doi:10.1029/2004GL020670, 2004.
- Schwanghart, W. and Kuhn, N. J.: TopoToolbox: A set of Matlab functions for topographic analysis, *Environ. Model. Softw.*, 25(6), 770–781, doi:10.1016/j.envsoft.2009.12.002, 2010.
- Sharp, M.: Surging glaciers: behaviour and mechanisms, *Prog. Phys. Geogr.*, 12(3), 349–370,

doi:10.1177/030913338801200302, 1988.

Simonsen, S. B. and Sørensen, L. S.: Implications of changing scattering properties on Greenland ice sheet volume change from Cryosat-2 altimetry, *Remote Sens. Environ.*, 190, 207–216, doi:10.1016/j.rse.2016.12.012, 2017.

5 Sørensen, L. S., Simonsen, S. B., Meister, R., Forsberg, R., Levinsen, J. F. and Flament, T.: Envisat-derived elevation changes of the Greenland ice sheet, and a comparison with ICESat results in the accumulation area, *Remote Sens. Environ.*, 160, 56–62, doi:10.1016/j.rse.2014.12.022, 2015.

Thomas, R.: Force perturbation analysis of recent thinning and acceleration of Jakobshavn Isbrae, Greenland., *J. Glaciol.*, 50(168), 57–66, doi:10.3189/172756504781830321, 2004.

10 Thomas, R., Frederick, E., Krabill, W., Manizade, S. and Martin, C.: Recent changes on greenland outlet glaciers, *J. Glaciol.*, 55(189), 147–162, doi:10.3189/002214309788608958, 2009.

van der Veen, C. J.: Tidewater calving, *J. Glaciol.*, 42(141), 375–385, doi:10.1179/102453311X13127324303399, 1996.

van der Veen, C. J.: Fracture mechanics approach to penetration of surface crevasses on glaciers, *Cold Reg. Sci. Technol.*, 27(1), 31–47, doi:10.1016/S0165-232X(97)00022-0, 1998.

15 van der Veen, C. J.: Fracture propagation as means of rapidly transferring surface meltwater to the base of glaciers, *Geophys. Res. Lett.*, 34(1), 1–5, doi:10.1029/2006GL028385, 2007.

Vieli, A. and Nick, F. M.: Understanding and Modelling Rapid Dynamic Changes of Tidewater Outlet Glaciers: Issues and Implications, *Surv. Geophys.*, 32(4–5), 437–458, doi:10.1007/s10712-011-9132-4, 2011.

Weidick, A., Andreasen, C., Oerter, H. and Reeh, N.: Neoglacial Glacier Changes around Storstrømmen, North-East Greenland, *Polarforschung*, 64(3), 95–108, 1994.