

Interactive comment on “Impact of assimilating sea ice concentration, sea ice thickness and snow depth in a coupled ocean-sea ice modeling system” by Sindre Fritzner et al.

Anonymous Referee #1

Received and published: 16 October 2018

Review on “Impact of assimilating sea ice concentration, sea ice thickness and snow depth in a coupled ocean-sea ice modeling system”, by Fritzner et al., submitted for publication in The Cryosphere Discussion.

General comments :

The paper shows the results of sea ice data assimilation experiments into a coupled ocean and sea ice model using an ensemble Kalman filter (EnKF). Sea ice concentration, sea ice thickness and snow depth are assimilated in different combinations and verifications are performed using assimilated and independent observations. The impact of assimilation is measured on the analysis, 7-day forecasts and 5-month seasonal forecasts. The paper is generally well written. The assimilation experiments and verifications are well designed. The assimilation of snow depth is particularly original as it has not been done in other studies, as far as I know. However, some aspects could be clarified.

We thank the reviewer for the kind words and the careful and constructive feedback.

For examples, the observation-error used for the assimilation could have been explicitly specified. Information regarding the observation uncertainty for AMSR sea-ice concentration, SMOS sea-ice thickness and the snow depth product is now provided in the observations section.

In some cases it is not clear whether the verification has been done on the ensemble mean or on individual ensemble members (and then calculating and average).

Verification is always done on the ensemble mean. This has been clarified by writing ensemble mean instead of ensemble average when applicable.

Also the error of the ensemble mean and ensemble spread relationship could have been shown, as this is usually considered a requirement for an EnKF.

We agree that this would be an interesting result to show, but we think that for sea-ice this is less interesting than it likely is for other applications. We think it would be difficult to present in a useful manner in our case. In a normal EnKF system one would expect the ensemble spread to be of the same order as the ensemble mean error, but this is not the case here. To generate the ensemble spread we use perturbation amplitudes we find to be physically reasonable, we set this independent of model error. This creates an ensemble spread significantly lower than the model error. In addition, too large ensemble spread with only 20 ensemble members would give an assimilation result skewed towards the observations.

Specific comments :

1. A couple of sentences at the end of the abstract around line 15 are confusing to me. It seems that the conclusions about the assimilation of snow depth are contradictory: “. . . while the snow

observations have a positive effect on snow thickness and ice concentration. In our study, the seasonal forecast showed that assimilating snow depth lead to a worse estimation of sea-ice extent compared to the other assimilation systems, the other three gave similar results.” How come the assimilation of snow depth have a positive effect on ice concentration but lead to a worse estimation of sea-ice extent ?

We agree that this is not well formulated. The point we make here is that there is a difference between long- and short-term effect of the assimilation. Where a positive effect is seen at the shorter timescales, immediately after assimilation and for the one-week forecasts. While for the seasonal forecast over several months there is a negative effect of assimilating the snow depth observations. The text has been updated for clarification: «It is found that the assimilation of ice thickness strongly improves ice concentration, ice thickness and snow depth, while the snow observations have a smaller but still positive short-term effect on snow thickness and ice concentration. In our study, the seasonal forecast showed that assimilating snow depth lead to a less accurate long-term estimation of sea-ice extent compared to the other assimilation systems, the other three gave similar results.»

2. In section 4.1, page 7: First P_b is defined as the background-error covariance matrix. A couple of lines later, it is referred as the model-error covariance matrix. I think you should stick to background-error covariance matrix because model-error covariance matrix is usually reserved for the errors accumulated during model integration.

This has been changed.

3. In section 4.3, it is mentioned that there is 5 thickness categories; I assume they are the partial concentrations for each thickness categories and that the total ice concentration can be calculated from them. Later it is mentioned that the assimilation can result into a positive SIC but no volume. Does that mean that the 5 thickness categories and the SIC are all independent analysis variables ?

The 5 partial SICs and the total SIC is 6 different parameters in the analysis, but they are not independent, the total SIC is only a sum of the 5 partial SICs. The model uses the 5 partial SICs, while the total SIC is a dummy parameter used for assimilation. When assimilating, the total SIC is the parameter corresponding to the observations, while the partial SICs are updated based on correlation and these are the ones used in the model afterwards.

If that is the case, wouldn't it be better to only have the 5 partial concentrations as analysis variable and calculate SIC ? It seems that it would avoid the problem of having positive SIC but no volume.

If we understand you correctly this is what is already done. What we mean is that we can have a partial SIC larger than one, but the corresponding partial SIT zero or less than zero. New text: where some areas have a positive partial SIC but the corresponding partial SIT is zero or less than zero.

4. In Figure2, what are the observation uncertainty of AMSR-E/2 ice concentration used in the calculation of the RMSE ? Are they included in the product and are they constant values or are they specified for each points ?

The AMSR observation uncertainty is included in the product and specified in each grid point. This is now specified in section 3: Observations. New text: «The AMSR-E/2 SIC observation product includes individual uncertainty estimates for all grid points. This uncertainty is based on the sum of algorithm uncertainty and smearing uncertainty. Where smearing uncertainty is related to the location of the observation compared to the grid.»

5. Figure 5: Over which year(s) ? Is it against IceBridge observations ? Also in the text under section 5.4, please specify what is “observed satellite snow depth”, is this IceBridge ?

No, this is against the observed snow depth product used for assimilation. New caption: «RMSE of monthly averaged model SIT and snow depth averaged over all ensemble members for the years 2011-2013 calculated against the a) combined SMOS-Cryosat SIT product and b) observed snow-depth product. These are observations also used for assimilation.»

New text in section 5.4: «In Fig. \ref{fig:Snow_Thickness}b the RMSE of monthly averaged modelled snow depth over all ensembles validated against the observed satellite snow depth \citep{Rostosky_2018} used for assimilation is plotted.»

6. Page 16, line 5: “This lack of improvement can be an indication of a too simple snow component in our coupled system, only one snow layer is used.” I think that is pure speculation, unless the authors can show evidence to convince the readers. Could the reason be simply that there are large discrepancies between IceBridge and the assimilated snow depth products ? Same comment on page 21, line 25 and on page 22, line 30.

We agree that this is pure speculation and is mentioned as a suggestion to what might cause the lack of consistency between model and observations. As you mention there are large errors in the snow depth observation product too as compared to icebridge, and we agree this is a more likely reason for the large errors. We have changed the text to highlight this: «It is seen that within one grid cell, there are huge variations in the IceBridge snow observations. Such variations cannot be provided with a coarse resolution model. Hence large errors are found for the RMSE against IceBridge observations for experiments where the snow observation are assimilated, even though IceBridge is used to «tune» the assimilated product \citep{Rostosky_2018}. In addition, the snow component used in our coupled system is likely too simple, having only one snow layer, which may effect the snow cover accuracy.»

7. Table 3: It is hard to understand from the caption what are the numbers in the table. Is the averaged snow depth over a grid cell compared to the model ensemble mean ?

There was an error here, the ensemble mean is validated by observations averaged over all grid cells.

It would help rephrase the caption, maybe removing one of the 3 averages and using “ensemble mean”, if that is appropriate.

We agree, new caption: «The annual-mean RMSE of the ensemble-mean snow depth averaged over all grid cells. The five experiments and the snow-depth satellite observations are compared to the IceBridge airborne snow-depth observations.»

8. Figure 8: Are these monthly averages over the 3 years ? The caption is hard to read because the “seven day” and “forecast” are too far apart.

Yes, they are, new caption: «RMSE of monthly averaged (over three years) ensemble mean of seven-day forecast SIC validated against a) AMSR-E/2 SIC observations and b) OSISAF SIC observations.»

9. Figure 9: Please specify in the caption that (b) is the monthly averaged RMSE over the three years.

This is now corrected, new text: «The figures show RMSE of the ensemble mean SIC averaged over 3 years and verified against the assimilated OSISAF SIC.»

10. Figure 10: I think it is unrealistic to use re-analysed forcing for the seasonal forecasts, as the re-analysed forcing would not be available in an operational real time context.

Agreed, this is of course not very realistic, but more a simplification since the focus of this study is on the assimilation. Since we compare our assimilation results with a control run using the same reanalysed forcing we think that the comparison is fair.

Technical corrections :

Page 5, line 8: Change “. . .observations where given...” to “. . .observations were given...”

This is corrected

Caption of Figure 3 : Change “low concentration ice (> 50 %)” to “low concentration ice (< 50 %)”

This is corrected

Caption of Figure 4 : Change “the blue stars the OSISAF” to “the red stars the OSISAF”.

This is corrected

Caption of Figure 4 : For ice volume the units are km. Is it the volume per unit area ?

No, this is an error. The figures has been updated.

Caption of Figure 4 : It would be easier for the readers to mention that the x-labels are month-year.

Text added: The xlabel is given as month-year.

Page 14, line : Change “sea-ice extent being too large and the ice is too thick” to “sea-ice extent being too large and the ice being too thick”.

This is corrected

Page 18, line 18: “The figures show that that . . .”

This is corrected

Figure 10: Change “The blue line represents a forecast using a climatological forcing made from an over of atmospheric data for 2000-2014 with assimilation” to “The blue line represents a forecast using a climatological forcing made from atmospheric data over 2000-2014 with assimilation”

This is corrected

Page 21, line 31: Change “The main parameters analysed in this study snow depth, SIT and SIC all vary on longer time scales than one week for the spatial resolution in our model” to “The main parameters analysed in this study, snow depth, SIT and SIC, all vary on longer time scales than one week for the spatial resolution in our model”

This is corrected

Response comment Reviewer 2

Anonymous Referee #2 Received and published: 4 December 2018

General Comments

In this paper, the authors perform 5 experiments with a 20 km coupled ROMS-CICE model forced with ERA-Interim forcing for 3 full years for the period of 2011-2013. The five experiments are 1) assimilation of OSISAF sea ice concentration (SIC) only, 2) assimilation of OSISAF SIC and CryoSat-2 sea ice thickness (SIT), 3) assimilation of OSISAF SIC and SMOS SIT, 4) assimilation of OSISAF SIC and AMSR-E/2 snow depth observations, and 5) control run without any data assimilation. The Ensemble Kalman Filter (EnKF) is the data assimilation technique used in this study. Ocean boundary conditions are provided by the FOAM ocean model. Two sets of experiments are performed: 1) assimilation experiments with 20 ensemble members with a 7-day assimilation time step, 2) seasonal forecasts with 20 ensemble members for the five-month period beginning in April/May to examine the skill in predicting the September sea ice minimum extent.

The authors computed the annual RMSE of the ensemble mean SIC over the three-year period and found that from January – August, the SIT experiments performed similarly and outperformed the SIC-only run during that period when using the weighted AMSR-E/2 data. From September – November, the SIC experiment had the lowest error. This could be related to no IT data during the summer months. The authors speculate the model has difficulty in transitioning from the melt to growing season. When comparing against the OSISAF ice concentration (which was assimilated into the model), the SIT experiment using SMOS showed the lowest RMSE from January – July. The snow depth experiment showed a lower RMSE than the SIC-only experiment for the period of January – June.

The authors examined “hit rates” to determine which experiment led to the most accurate number of grid cells classified as open water (concentration < 10%), low (<50%) or high concentration (>50%) and found that the experiments with the assimilation of ice thickness performed best. Total ice volume is examined for all 5 experiments and they find that except for the control run, the volume steadily decreases from year to year. The authors need to better address why this is happening, and propose future studies to investigate this further.

The decrease in sea-ice volume is not a model problem, but a response to the assimilation where the model, in general, has too much and too thick ice. An attempt at a discussion of this case was given on page 11. line 26-32 in the old manuscript. This section has now been modified to make this more clear, new text is added: « The control model is found to have too thick ice compared to the observations, while the experiments assimilating SIT are much closer to the observations, though largely biased. This can be used to explain the drastic decrease in sea-ice volume found in Fig.

\ref{fig:extent_volume}b. The model SIT is adjusting towards the observations by rapidly thinning the sea-ice.»

Comparisons are performed with the annual mean ice thickness and snow depth from all 5 experiments versus data from NASA Operation IceBridge. Since IceBridge data is only available for typically 10 transects for March/April each year; this is not a very compelling analysis. While Arctic snow depth data is difficult to obtain, it is recommended that the authors examine additional sources of ice thickness data, such as ice mass balance data (see comment below) which has much better temporal and spatial resolution.

Thank you for this valuable suggestion. We were not aware of these observations, and they are now included in the validation of the experiments.

Seasonal forecasts are evaluated by performing 5-month experiments for all five cases beginning in April of 2011, 2012 and 2013 to examine the SIC RMSE. When averaged for all three years, the SIT experiments perform best. Through mid-June, the snow depth experiment is very similar to the CryoSat-2 (SITI), but afterward the error increased significant and mirrors the control runs high error from August through September.

With the exception to the Lisæter (2007) reference, throughout the paper you should consistently refer to CryoSat as CryoSat-2.

This is corrected

Why aren't ice mass balance buoys used in your study? Look at available data at: <http://imb-crrrel-dartmouth.org/results/>. During the period of your study, there is drifting buoy data available.

They are now included.

Are melt ponds used in your CICE simulations? If yes, state that in Section 4.3.

Yes, the model use melt pond parametrization. Information regarding this is now added to the description of the model in section 2: «The model has a thermodynamic component calculating the local growth rate of snow and ice, ice dynamics component calculating ice drift based on the material ice strength, a transport component, a melt pond parametrization and a ridging parametrization used to distribute ice in thickness categories \citep{Hunke_2015}.»

Why didn't you evaluate model ice drift errors using the International Arctic Buoy Programme buoy data? See <http://iabp.apl.washington.edu/>

We did not have in mind to include ice drift in the study, thus unfortunately, the model drift output was not saved.

This is a very well written paper with clear tables and complementary graphics. I recommend publication after my comments are addressed.

We thank the reviewer for the kind words and the careful and constructive feedback.

Specific Comments

Page 2 lines 15-25: Suggest you add the following reference to this section when discussing operational system assimilating SIC:

Posey, P. G., Metzger, E.J., Wallcraft, A.J., Hebert, D.A., Allard, R.A., Smedstad, O.M., Phelps, M.W, Fetterer, F., Stewart, J.S., Meier, W.N., Helfrich, S.R., 2015. Assimilating high horizontal resolution sea ice concentration data into the US Navy's ice forecast systems: Arctic Cap Nowcast/Forecast System (ACNFS) and the Global Ocean Forecast System (GOFS 3.1). *The Cryosphere* 9 2339-2365. doi: 10.5194/tcd-9-2339- 2015.

Thank you for the advice, this has now been added: «\cite{posey2015assimilating} assimilated high-resolution SIC observations (~4 km) into a coupled ocean sea-ice model, the Arctic Cap Nowcast/Forecast System (ACNFS) using the 3DVAR assimilation method. In this study, they showed that increased observation resolution has a significant impact on the ice-edge forecast.»

Page 3 first paragraph: Consider adding the following recent references when discussing the use of CryoSat-2 data:

Allard, R. A., Farrell, S. L., Hebert, D. H., Johnston, W. F., Li, L., Kurtz, N. T., Phelps, M. W., Posey, P. G., Tilling, R., Ridout, A., and Wallcraft, A. L.: Utilizing CryoSat-2 sea ice thickness to initialize a coupled ice-ocean modeling system, *Advances in Space Research*, 62, doi:10.1016/j.asr.2017.12.030, 2018.

Blockley, E. W. and K. A. Peterson: Improving Met Office seasonal predictions of Arctic sea ice using assimilation of CryoSat-2 thickness, *Cryosphere*, 12, 3419–3438, doi:10.5194/tc-12-3419-2018.

Xie, J., F. Countillon, and L. Bertino: Impact of assimilating a merged sea-ice thickness from CryoSat-2 and SMOS in the Arctic reanalysis, *Cryosphere*, 12, 3671-3691, doi:10.5194/tc-12-3671-2018.

Thank you, we were not aware of the recent papers. We have improved the text to include these studies: «In the last couple of years, there has also been an increase in the use of Cryosat-2 observations in various forms for assimilation. \cite{Chen_2017} assimilated both the SMOS thin SIT and the CryoSat-2 thick SIT into the National Centers for Environmental Prediction's (NCEP) Climate Forecast System version 2 \cite{Saha_2014} using the localized error subspace transform

ensemble Kalman filter \citep{Nerger_2013}. This study showed improved sea-ice prediction with SIT assimilation, thus verifying the importance of SIT observations to achieve accurate sea-ice forecasts. \cite{xie2018impact} assimilated a blended SMOS CryoSat-2 product into TOPAZ. They showed that these observations provide the primary source of observational information in the central Arctic, and when assimilating this product the model SIT was improved. \cite{blockley2018improving} argued that by assimilating Cryosat-2 observations, the Arctic summer prediction of ice extent and location were significantly improved. \cite{allard2018utilizing} used CryoSat-2 observations for initialization in the coupled ocean sea-ice ACNFS model. The study showed improved model thickness with CryoSat-2 initialization when compared to independent ice thickness observations.»

Page 4 line 12: Please state the horizontal resolution of the ERA-Interim dataset

This is now added: «The coupled model is forced by atmospheric data from the ERA-Interim dataset from the European Centre for Medium Ranged Weather Forecast \citep[ECMWF;]{Dee_2011}. The ERA-Interim dataset has a horizontal resolution of approximately 0.75° , corresponding to T255 spectral truncation.»

Page 4 lines 13-14: You use FOAM for prescribed ocean boundary conditions. What do you use for the CICE boundary conditions?

At the moment no boundary conditions are used for the sea-ice. For the most part of the year, this is not a problem because the sea-ice is surrounded by ocean. While during winter when the sea-ice extends beyond the Behring straight this might be a problem for this area, but this is not included here, and we do not believe this to be an issue regarding the results.

Page 6 last paragraph: What is the accuracy of the AMSR-E/2 snow depth data?

This has now been added: «For the snow depth product uncertainty estimates exist for every grid point. There are two main sources of uncertainties in this observation product: The first is that the number of IceBridge observations used to develop the empirical relationship between brightness temperatures and snow depths is small compared to the coverage of the product. The second uncertainty is in the input parameters (brightness temperature, ice concentration etc.). More on how the uncertainties are explicitly calculated can be found in \citep{Rostosky_2018}.»

Page 8: You state the coupled modeling system is run for 1 year as an initial state. Was it spun-up from rest? How was ice initialized? Uniform everywhere from a particular thickness?

This has now been clarified in the manuscript: «The initial ensemble is generated from ice states from January 1, based on 20 different years of a standalone ice model run without assimilation. The standalone model was initialised without ice in 1979. All initial ocean states are model output at initial date January 1, 2010. This output is taken from a model spin-up over 1993-2010.»

Page 8 last paragraph: Why didn't you include another experiment which included a blended CryoSat-2/SMOS ice thickness?

The focus of this study was on the individual observation products. , we could have tested all sorts of combination between the products, but this would become quite messy. In addition, the blended product provided an alternative dataset for model verification.

Page 13 Figure 4b: Please explain your views on why the ice volume (except for control run) steadily decreases. I suggest in your conclusions section to include to some possible follow-on studies to better investigate this issue.

This is already mentioned on p.11 lines 26-32 in the previous version of the manuscript. The decrease is related to too much ice in the control model, and due to assimilation, the thickness is slowly going towards the observed values which are thinner. This section has been updated to further clarify this result, see comment above.. Thus no further studies of this effect should be necessary.

Page 14 lines 20-24: Please include figures and discussion on comparison for April 2012 and 2013?

Figure 7. is only meant as an illustration of the curves shown in figure 4b. Even though the figures could be added for 2012 and 2013 we do not think this would provide any additional information from figure 4b. Also, 8 more figures would take a lot of space in the manuscript and would be too messy as they are providing only little new information.

Page 14 last paragraph: Have you looked at Dartmouth (formerly CRREL) IMB data for an additional source of ice thickness data? These data sets have much more temporal coverage than just Mar/Apr from NASA IceBridge.

Yes, as mentioned previously a new section including verification of these data have been added: « Another independent dataset of SIT observations complementing the IceBridge observations by observations throughout the year is the IMB buoy dataset. The result of model validation with the IMB is shown in table $\{tab:IMB_th\}$. For these observations, a slightly different method than that applied for IceBridge is performed. This is because IceBridge temporarily only covered March-April, while the IMB data span the entire year. The buoy observations are converted to daily averages on the model grid. From these values, the RMSE is calculated on the 7-day ensemble mean and averaged for each year. Since SIT is a relatively slow varying variable, for each 7-day output, observations from +/-3 days are used to increase the number of observations. The IMB observations do not include an uncertainty estimation, hence the RMSE is not normalised was the case for other other RMSE values in this work. The results show that over the three study years, the SMOS internal SIT assimilation system has the lowest RMSE values, followed by the CryoSat-2 internal SIT assimilation. The other three show similar results, indicating the positive impact of assimilating ice thickness in the model.» A table has also been added illustrating the yearly averaged results.

Page 16 last paragraph: Table 3 shows yearly averaged RMSE values of ensemble average of snow depth compared to NASA IceBridge. Explain how you can do this when NASA IceBridge is only available for Mar/Apr each year.

The yearly average is a Mar/Apr average. This has now been made clearer by adding it into the figure text: «The Mar/Apr-mean RMSE of the ensemble-mean snow depth averaged over all grid cells.» In addition, new text has been added: The same method as for the SIT in table \ref{tab:IceBridge} was used, where Mar/Apr model values are compared to the IceBridge observations and averaged.

Page 18 lines 15-16: You state five-month forecasts, but experiments are performed April – September What are the actual dates? Apr 30 – Sept 30 would be 5 months; April 1 – Sept 30 would be 6 months.

We mean it is an approximately 5-month forecast. The start date varied a bit because of the 7-day assimilation cycle. Information regarding this is now added to the text: «This is done by running each of the experiments from mid-April to mid-September each year without assimilation and validating against the OSISAF SIC observations. The actual start date varied slightly from year to year because of the 7-day assimilation cycle, but the start date was the same for all experiments.»

Technical Corrections:

Page 1 line 7: replace “asses” to “assess”

done

Page 1 line 12: should be CryoSat-2 (and throughout the paper)

done

Page 1 line 16: replace “lead” to “led”

done

Page 2 line 14: add comma after “later”

done

Page 8 line 30: reword “Five assimilation experiments” to “Five experiments”

done

Page 12: Fig 3 caption: first line should read “low concentration ice <50%” (not >50)

done

Page 13 line 8: replace “to much ice” to “too much ice”

done

Page 26 line 24: Provide more complete info for Sakov EnKF-C user guide (2015) reference

Added arXiv: «Sakov, P.: EnKF-C user guide. arXiv:1410.1233., 2015.

Interactive comment on The Cryosphere Discuss., <https://doi.org/10.5194/tc-2018-171>, 2018. C5

Impact of assimilating sea ice concentration, sea ice thickness and snow depth in a coupled ocean-sea ice modeling system

Sindre Fritzner¹, Rune Graversen¹, Kai H. Christensen², Philip Rostosky³, and Keguang Wang⁴

¹UiT The Arctic University of Norway, Tromsø, Norway

²The Norwegian Meteorological Institute, Oslo, Norway

³Institute of Environmental Physics, University of Bremen, Germany

⁴The Norwegian Meteorological Institute, Tromsø, Norway

Correspondence to: Sindre Fritzner (sindre.m.fritzner@uit.no)

Copyright statement. TEXT

Abstract. The accuracy of the initial state is very important for the quality of a forecast, and data assimilation is crucial for obtaining ~~a~~the best possible initial state. For many years, sea-ice concentration was the only parameter used for assimilation into numerical sea-ice models. Sea-ice concentration can easily be observed by satellites, and satellite observations provide a full Arctic coverage. During the last decade, an increasing number of sea-ice related variables have become available, these include sea-ice thickness and snow depth, which are both important parameters in the numerical sea-ice models. In the present study, a coupled ocean-sea-ice model is used to ~~asses~~assess the assimilation impact of sea-ice thickness and snow depth on the model. The model system with the assimilation of these parameters is verified by comparison with a system assimilating only ice concentration and a system having no assimilation. The observations assimilated are sea ice concentration from the Ocean and Sea Ice Satellite Application facility, thin sea ice thickness from the European Space Agency's (ESA) Soil Moisture and Ocean Salinity mission, thick sea ice thickness from ESA's ~~CryoSat~~CryoSat-2 satellite, and a new snow depth product derived from the National Space Agency's Advanced Microwave Scanning Radiometers (AMSR-E/AMSR-2) satellites. The model results are verified by comparing assimilated observations and independent observations of ice concentration from AMSR-E/AMSR-2, and ice thickness and snow depth from the IceBridge campaign. It is found that the assimilation of ice thickness strongly improves ice concentration, ice thickness and snow depth, while the snow observations have a ~~positive~~smaller but still positive short-term effect on snow ~~thickness and ice~~depth and sea-ice concentration. In our study, the seasonal forecast showed that assimilating snow depth ~~lead to a worse~~led to a less accurate long-term estimation of sea-ice extent compared to the other assimilation systems, the other three gave similar results. The improvements due to assimilation were found to last for at least 3-4 months, possibly even longer.

1 Introduction

Observations show that for the last 50 years there has been a decline in both Arctic sea-ice extent (Stroeve et al., 2007; Perovich et al., 2017) and sea-ice thickness (Kwok and Rothrock, 2009), in addition, models show that the sea-ice decline is likely to continue (Zhang and Walsh, 2006). Wang and Overland (2012) estimate the Arctic ocean to be nearly ice-free within the 5 2030s. This large change in the global climate system leads to a need for improved models and forecasting systems due to more variable and mobile Arctic sea ice (Eicken, 2013). In addition, a decreased amount of sea ice will lead to increased Arctic ship traffic (Smith and Stephenson, 2013). Safe travel in the Arctic is dependent on accurate knowledge of weather and sea ice. The Arctic is characterized by harsh conditions involving for instance sea ice, icebergs, and polar low storms. The numerical weather prediction models are becoming more complex and detailed, but still, the vital part of an accurate forecast is the model 10 initial state. Accurate initial states can be achieved by assimilating observations into the model system.

For sea-ice modelling in the Arctic, observations are sparse. The sea-ice concentration (SIC), defined as the fraction of the total area covered by sea-ice, has been available since the start of the satellite era in 1979, but observations of other parameters such as sea ice thickness (SIT) are more difficult to obtain because of the remote location, and satellites cannot easily be used to extract information about the SIT. The passive microwave satellites derive SIC from brightness temperatures, but many of 15 the earth observing satellites do not have sufficient wavelength to observe changes in the brightness temperature as a function of the SIT. Thus acquiring SIT from satellites are significantly more difficult than SIC, but as will be described later, satellites using the L-band frequency can, to some degree, be used to measure the SIT as a function of brightness temperature.

During the last 15 years, there have been various studies of SIC assimilation, using several different models and assimilation methods. Lisæter et al. (2003) assimilated SIC obtained from passive microwave satellite into a coupled ocean-ice model using 20 the Ensemble Kalman Filter (EnKF; Evensen, 1994; Burgers et al., 1998). In the study of Lisæter et al. (2003), the assimilation was found to have a strong effect on the modelled SIC, and also small effects on other model parameters due to the multivariate properties of the EnKF. The multivariate properties of the EnKF consist of a model update for all model variables based on correlation with the observed variables. A similar SIC assimilation study using the 3D-Variational (3D-Var) assimilation method was done by Caya et al. (2010). In this study, both ice charts from the Canadian east coast and Radarsat 2 SIC observations 25 were assimilated. Significant improvements to the short-term forecast were found for the assimilation system. Studies with the coupled ocean-ice model TOPAZ (Sakov et al., 2012) have shown improvements to SIT and multivariate effects on SIT for assimilation of SIC (Sakov et al., 2012). Other SIC studies have been done by Lindsay and Zhang (2006) and Wang et al. (2013) both using nudging methods to show model improvements for SIC assimilation. [Posey et al. \(2015\) assimilated high-resolution SIC observations \(4 km\) into a coupled ocean sea-ice model, the Arctic Cap Nowcast/Forecast System \(ACNFS\) using the 30 3DVAR assimilation method. In this study, they showed that increased observation resolution has a significant impact on the ice edge forecast.](#)

In recent years there has been a focus on increasing the number of observable ice parameters, especially accurate knowledge of the Arctic SIT is important in order to quantify changes in the total Arctic sea-ice volume and to elucidate changes related to for instance global warming. Dedicated satellite altimeters like ICESat (Forsberg and Skourup, 2005) and CryoSat-2 (Laxon

et al., 2013) have been prepared for SIT measurements. These satellites use measurements of the ice freeboard to calculate the SIT (Kurtz and Harbeck, 2017; Kurtz et al., 2014b). Another source of satellite SIT observations is the European Space Agency's (ESA) Soil Moisture and Ocean Salinity (SMOS) mission. The SMOS mission uses L-band passive microwave measurements utilizing long penetration depth and a relationship between observed brightness temperature and ice thickness (Tian-Kunze et al., 2016). However, in general, the uncertainties of the ~~CryoSat~~CryoSat-2 and SMOS SIT observations are high (Zygmuntowska et al., 2014; Xie et al., 2016), which result in reduced, though still valuable, observational information available for assimilation into the model system. The SIT observations are limited to winter conditions when the snow and ice are dry.

One of the first studies with SIT assimilation was done by Lisæter et al. (2007). In this study, computer-generated SIT observations simulating CryoSat observations were assimilated into a coupled ice-ocean model using the EnKF. The assimilation showed significant effects on the model state; both improvements to the modelled SIT and multivariate effects on SIC, ocean temperature and ocean salinity were found. Yang et al. (2014) used the localized singular evolutive interpolated Kalman filter (Pham, 2001) to assimilate the SMOS SIT observations into the Massachusetts Institute of Technology general circulation model (Marshall et al., 1997). In this study, an improved thickness forecast when assimilating SMOS observations and some improvements to the SIC forecasts were found. Similar to Yang et al. (2014), Xie et al. (2016) used the EnKF to assimilate SMOS SIT observations into the TOPAZ system (Sakov et al., 2012). In this study it was found that assimilation of SMOS observations showed improvements for the ice thickness along the ice edge, both compared to SIT observations not assimilated and compared to the SMOS observations themselves. In general, similar to that found by Yang et al. (2014) the SMOS observations were found to have a relatively small impact on the SIC, and the SIT far from the ice edge.

Fritzner et al. (2018) assimilated SMOS observations into a standalone sea-ice model with the EnKF. This study showed that due to the correlation between SIC and SIT, the SMOS observations were found to have a positive effect on the modelled SIC. In the last couple of years, there has also been an increase in the use of Cryosat-2 observations in various forms for assimilation. Chen et al. (2017) assimilated both the SMOS thin SIT and the ~~CryoSat~~CryoSat-2 thick SIT into the National Centers for Environmental Prediction's (NCEP) Climate Forecast System version 2 (Saha et al., 2014) using the localized error subspace transform ensemble Kalman filter (Nerger and Hiller, 2013). This study showed improved sea-ice prediction with SIT assimilation. Thus verifying the importance of SIT observations to achieve accurate sea-ice forecasts. Xie et al. (2018) assimilated a blended SMOS CryoSat-2 product into TOPAZ. They showed that these observations provided the primary source of observational information in the central Arctic, and when assimilating this product the model SIT was improved. Blockley and Peterson (2018) showed that by assimilating Cryosat-2 observations the Arctic summer prediction of ice extent and location was significantly improved. Allard et al. (2018) used CryoSat-2 observations for initialization in the coupled ocean sea-ice ACNFS model. The study showed improved model thickness with CryoSat-2 initialization when compared to independent ice thickness observations.

Recent attempts have proved that it might be possible to observe snow depth from satellite (Markus and Cavalieri, 1998; Maaß et al., 2013; Rostosky et al., 2018). Both Maaß et al. (2013) and Rostosky et al. (2018) used a relationship between observed brightness temperature and snow depth to calculate the latter variable. Due to the close connection between snow,

albedo and ice melting, accurately modelled snow depths are expected to have a large impact on the snow and ice models. Snow observations are limited to the winter season when the ice and snow are dry.

In our study, a coupled ocean sea-ice model (Kristensen et al., 2017) is used. The coupled model is prepared for improved sea-ice representation compared to previous coupled ocean sea-ice models. This improvement will give a deeper insight into how sea-ice is affecting both the ocean and atmosphere. The assimilation system will be tested with different kinds of observations to analyse both long-term and short-term effects. Observations of SIC, SIT and snow depth are assimilated. The results will be verified with independent and semi-independent data, in addition to forecasts both in summer and winter.

This study is important in order to elucidate the effect of different sea-ice observations and reveal the most important observations for an improved sea-ice forecast. Even though some studies have looked into the assimilation of different SIT observations, as far as we know this is the first study to compare the effect of the different observations on the assimilation system. In addition, as far as we know, this is the first study to present the assimilation of snow depth observations in a coupled ocean sea-ice model.

2 The coupled ocean sea-ice model

The coupled model (Kristensen et al., 2017) is based on the Regional Ocean Modelling System (ROMS; Shechetkin and McWilliams, 2005; Moore et al., 2011) version 3.6 as the ocean component and the Los Alamos sea-ice model version 5.1.2 (CICE; Hunke and Dukowicz, 1997; Hunke et al., 2015a) as the ice component. The ROMS model is a state-of-the-art ocean model, which in our study is configured with 35 terrain-following vertical layers. The eddy viscosity and eddy diffusivity are parametrized using a second-order turbulence closure model.

The CICE model is a state-of-the-art sea-ice model with 5 thickness categories, 7 ice layers and one snow layer. The model has a thermodynamic component calculating the local growth rate of snow and ice, ice dynamics component calculating ice drift based on the material ice strength, a transport component, [a melt pond parametrization](#) and a ridging parametrization used to distribute ice in thickness categories (Hunke et al., 2015a). The model has a horizontal resolution of 20 km with 242x322 grid cells covering the entire Arctic ocean. The model domain covering the Arctic sea is shown in Fig. 1.

The coupled model is forced by atmospheric data from the ERA-Interim dataset from the European Centre for Medium Ranged Weather Forecast (ECMWF; Dee et al., 2011). [The ERA-Interim dataset has a horizontal resolution of approximately 0.7°, corresponding to a T255 spectral truncation.](#) In addition, the model has prescribed ocean boundary and climatic forcing from the Fast Ocean Atmosphere Model (FOAM; Bell et al., 2003). The assimilation system used in the model is the Ensemble Kalman Filter. The code used for assimilation is the EnKF-c code (Sakov, 2015). The EnKF-c is an easy-to-implement and efficient framework for off-line data assimilation for use in geophysical models.




Figure 1. The model domain, the blue area is covered by the model, grey area indicate land areas

3 Observations

In the present study, observations related to the Arctic sea-ice are used for assimilation, these include SIC, SIT and snow depth. The SIC observations used for assimilation are from the European Organisation for the Exploitation of Meteorological Satellites (EUMETSAT) Ocean and Sea Ice Satellite Application Facility (OSISAF; Tonboe et al., 2016). The SIC product is the near-real-time global sea-ice concentration product. This dataset contains SIC observations calculated from brightness temperatures measured by SSMI/S passive microwave. The SSMI/S brightness temperatures are corrected for air temperature, wind roughening over open water and water vapour in the atmosphere by the ECMWF numerical weather prediction (NWP) model (Andersen et al., 2006). To convert from brightness temperatures to SIC a combination of the Bootstrap and the Bristol Algorithm is used (Tonboe et al., 2016). The Bootstrap algorithm is primarily used for observations with low SIC, and the Bristol algorithm for high SIC. The older OSISAF products do not include an error estimate, but an estimate of the observation confidence. The observation confidences are a simple measure of the observations quality, where 5 is excellent quality, 2 indicate poor quality, 1 indicates computation failure, and 0 no data. In the more recent OSISAF observations, a total uncertainty parameter is associated with each observation. In our study, the observation uncertainty of the OSISAF observations ~~where were~~ given by the following formula:

$$15 \quad TU = a + b * (5 - C), \quad (1)$$

where C is the confidence and TU is the total uncertainty, $a = 0.06$ and $b = 0.1$ are estimated based on the relationship between confidence and uncertainty in the more recent OSISAF observations. Observations flagged with a confidence of 0 or 1 are not used in our study. For verification of the modelled SIC, the ~~AMSR-E/Aqua Daily L3 12.5km-ESA Sea Ice Climate Change Initiative, Sea Ice Concentration product was used (Cavalieri et al., 2014), consisting Climate Data Record from the AMSR-E and AMSR-2 Instruments at 25 km Grid Spacing, Version 2.0 (Toudal Pedersen et al., 2017). The dataset consists of~~ satellite observations from the National Space Agency's Advanced Microwave Scanning Radiometers (AMSR-E/AMSR-2).

The AMSR-E/2 observations are, as the OSISAF SIC observations, also based on measurements from a passive microwave measuring the brightness temperature, and the observations are structured on a 12.525 km grid. ~~For the AMSR-E/2 dataset, the Enhanced NASA Team algorithm (Markus and Cavalieri, 2000) is used to convert from brightness temperatures to SIC.~~ The OSISAF and AMSR-E/2 datasets are different data products, but are in many cases tuned to give similar results and cannot be viewed as true independent datasets. The AMSR-E/2 product has a gap from October 2011, when AMSR-E failed, until AMSR-2 became operational in July 2012, this is in the middle of our analysis period resulting in less data for verification. The AMSR-E/2 SIC observation product includes individual uncertainty estimates for all grid points. This uncertainty is based on the sum of algorithm uncertainty and smearing uncertainty. Where smearing uncertainty is related to the location of the observation compared to the grid.

Two different SIT products are assimilated. For thick SIT observations, the CryoSat-2 Level-4 Sea Ice Thickness product is used (Kurtz and Harbeck, 2017). The ~~CryoSat-CryoSat-2~~ observations are based on radar altimeter measurements of sea ice freeboard. The SIT is derived assuming nominal densities for ice, snow and water, and only valid for high concentration ice (> 70 %; Kurtz et al., 2014b), thus they are assumed to be observations of thick ice relative to the SMOS observations. The snow depth used to calculate sea-ice elevation is constructed from the Warren climatology of snow depth (Warren et al., 1999), modified to account for the loss of multi-year ice in recent years (Kurtz and Farrell, 2011). The dataset has a spatial resolution of 25 km and a 30-day average temporal resolution covering the entire Arctic. For the ~~CryoSat-CryoSat-2~~ dataset, no uncertainty estimates are provided, thus following Zygmontowska et al. (2014) an uncertainty of 0.5 m was used for all ~~CryoSat-CryoSat-2~~ observations. Due to the low temporal coverage, this is most likely an underestimation of the uncertainty, and other publications have suggested higher uncertainties (Xie et al., 2016; Chen et al., 2017). In our study, the main focus is on the impact of the observations on the assimilation system and thus a low error is applied in order to elucidate the model impact of the observations. Since the ~~Cryosat-CryoSat-2~~ dataset is only valid for high concentration ice, all observations are in the internal part of the Arctic sea ice, and will in future reference also be referred to as internal ice thickness. The ~~Cryosat-CryoSat-2~~ observations are only available in the cold season from October to April.

For thin SIT observations, the daily L3C SMOS Sea Ice Thickness version 3.1 is used (Tian-Kunze et al., 2016). These SIT observations are acquired from a satellite using a passive microwave with L-band frequency. Measurements of brightness temperatures are converted into SIT using a radiation and thermodynamic model based on penetration depth (Tian-Kunze et al., 2014). Xie et al. (2016) found that observations thinner than around 0.4 m were the most realistic to use in the analysis, hence in this study observations thicker than 0.5 m have not been used. For the SMOS observations it is assumed that all observations are acquired at 100 % SIC, thus the observations are assimilated as normalised ice volume. The SMOS dataset has a resolution of 12.5 km and is structured on a stereographic grid. Since all SMOS observations are thinner than 0.5 m they are all located in the vicinity of the Arctic ice rim, and will in future reference also be referred to as rim ice thickness. As for the internal ice thickness observations, the SMOS SIT are only available in the cold season from October to April. The SMOS observations include individual uncertainty estimates for each grid point. These uncertainty estimates are a combination of uncertainties of measured brightness temperature, auxiliary data sets and assumptions made in the radiation and thermodynamic models, in general thicker ice has higher uncertainty (Kaleschke et al., 2017).

For verification of the modelled SIT, the weekly combined ~~SMOS-CryoSat~~ [SMOS-CryoSat-2](#) dataset version 1.3 was used (Ricker et al., 2017). This observation product provides SIT observations covering the whole Arctic during the cold season. In addition, the IceBridge L4 Sea Ice Thickness observations are used for verification (Kurtz et al., 2013, 2014a). This dataset consists of SIT and snow depth measurements from an aeroplane, using a radar altimeter measuring the ice freeboard.

5 The IceBridge observations are limited temporally to March-April, and spatially to parts of the Beaufort Sea, the Canadian Archipelago, and north of Greenland.

The snow depth observations are derived from AMSR-E/2 observed brightness temperatures (Rostosky et al., 2018). The data are available on a daily basis with a resolution of 25 km x 25 km. The algorithm uses the same technique which was developed by Markus and Cavalieri (1998) to retrieve snow depth over Antarctic sea ice. Their product is based on an empirical
10 relationship between the gradient ratio of the 37 GHz and 19 GHz brightness temperature observations and Antarctic snow depth. It was adapted to retrieve snow depth on Arctic sea ice (Comiso et al., 2003), but due to the radiometric properties of Arctic multi-year ice, the retrieval is limited to first-year ice only. The new product by Rostosky et al. (2018) makes use of lower frequency channels (i.e. brightness temperature observations at 6.9 GHz) which are less sensitive to the Arctic multi-year ice and thus the retrieval can be, with some exceptions (Rostosky et al., 2018), applied over the whole Arctic sea ice. The new
15 snow depth retrieval was trained and evaluated using NASA's Operation Icebridge airborne snow depth observations (Newman et al., 2014). Those observations are, however, mainly limited to March and April and, so far, no evaluation of the snow depth product exists for the remaining winter season. We, therefore, limit our analysis to snow depth observations in March and April. [For the snow depth product, uncertainty estimates exist for every grid point. There are two main sources of uncertainties in this observation product: The first is that the number of IceBridge observations used to develop the empirical relationship between brightness temperatures and snow depths is small compared to the coverage of the product. The second uncertainty is in the input parameters \(brightness temperature, ice concentration etc.\). More on how the uncertainties are explicitly calculated can be found in \(Rostosky et al., 2018\).](#) When the model simulations were performed the snow depth product was in its early development state. By now, a slightly updated version of the snow depth product exists, but since the overall differences between the updated version and the early state version are small we do not expect the updated data set to yield substantially
20 different results.
25

[In addition to the radar observations, Ice Mass Balance \(IMB\) buoy observations of SIT and snow depth \(Perovich et al., 2018\) are used for model verification. These data include measurements of SIT and snow depth from drifting buoys in the Arctic at multiple time intervals and different location every year. The measurements are performed by sounders Polashenski et al. \(2011\).](#)

4 Methods and model setup

4.1 The Ensemble kalman filter

The Ensemble Kalman Filter (EnKF) is a sequential data-assimilation method used in a wide variety of geophysical systems (Evensen, 1994, 2009; Houtekamer and Zhang, 2016). The analysis equation for the EnKF is given by (Jazwinski, 1970; Evensen, 2003),

$$\mathbf{x}_a = \mathbf{x}_b + \mathbf{P}_b \mathbf{H}^T (\mathbf{H} \mathbf{P}_b \mathbf{H}^T + \mathbf{R})^{-1} (\mathbf{y} - \mathbf{H} \mathbf{x}_b). \quad (2)$$

The model background and analysis state vectors are matrices given by, $\mathbf{x}_b \in \mathbb{R}^{n \times N}$ and $\mathbf{x}_a \in \mathbb{R}^{n \times N}$, respectively. Here n is the number of variables (that will become updated) times number of grid cells, and N is the number of ensemble members. The covariance of the observations is given by $\mathbf{R} \in \mathbb{R}^{m \times m}$, where m is the number of observations, $\mathbf{H} \in \mathbb{R}^{m \times n}$ is the observation operator, which is a transformation operator between model and observations space, and $\mathbf{y} \in \mathbb{R}^{m \times N}$ is the observation matrix. For the EnKF the background error covariance matrix, \mathbf{P}_b , is estimated based on the covariance of an ensemble of model states. The ensemble is generated by either perturbing the forcing, the model parameters, the observations or a combination of the three. The estimator for model-background error covariance, $\mathbf{P}_b \in \mathbb{R}^{n \times n}$, is

$$\mathbf{P}_b = \overline{(\mathbf{x}_b - \overline{\mathbf{x}_b})(\mathbf{x}_b - \overline{\mathbf{x}_b})^T}. \quad (3)$$

The overbars indicate an ensemble average. In our study, the Deterministic Ensemble Kalman Filter (DEnKF) proposed by Sakov and Oke (2008) is used. This method solves the analysis equation without the use of perturbed observations.

When using the EnKF spurious co-variances might occur due to distant state vector elements and insufficient model rank when small ensemble sizes are used. These artefacts can be reduced by using a method for localization (Evensen, 2003; Sakov and Bertino, 2011), limiting the assimilation to affect a smaller area. There are several methods for localization, and in this study, the polynomial taper function (Gaspari and Cohn, 1999) is used. The taper function is a bell-shaped function providing stronger influence to nearby grid cells.

4.2 Ensemble spread

Sufficient ensemble spread is essential for a robust and well-functioning EnKF assimilation system. In general, this is maintained by the Kalman Filter equations, but it is important to also take into account the uncertainty in the model and the atmospheric forcing. The atmospheric forcing is perturbed to account for uncertainty in the forcing. The atmospheric forcing is perturbed using smooth pseudo-random fields (Evensen, 2003) with zero mean and standard deviation based on perturbation values applied also in the more tested and robust TOPAZ system (Sakov et al., 2012). For the 2-m temperature, the standard deviation is 3 K, cloud cover is 20 %, per-area precipitation flux is 4×10^{-9} m, and for wind, 1 m s^{-1} in both horizontal directions is applied. To account for model uncertainty the ice strength parameter, P , is perturbed. This is done by perturbing

the model parameter C_f which is the frictional energy dissipation parameter. In CICE, C_f is proportional to the ice strength (Hunke et al., 2015b),

$$P \propto C_f. \tag{4}$$

The default value of C_f is 17, but according to Flato and Hibler (1995) this is not a well-known parameter. In our study, this parameter is modelled as a stochastic variable with a mean value of 17 and a standard deviation of 10, the different values are chosen based on values found during model tuning using observations by Flato and Hibler (1995). Since only one value less than 10 was found in their study, values less than 10 for C_f is not used.

4.3 Experimental design

The assimilation model system consists of 20 ensemble members, with an assimilation time step of seven days. Similar to Sakov et al. (2012) a localization radius of 300 km is used. The initial ensemble is generated from ice states from ~~the 1. January from January 1. based on~~ 20 different years ~~-, while all of a standalone sea-ice model run without assimilation. The standalone model was initialised without ice in 1979. All initial~~ ocean states are ~~from model output at~~ the initial date ~~1. January January 1. 2010. This output is taken from a model spin-up over 1993-2010.~~ Before performing the experiments, a model system assimilating ice concentration and sea-surface temperature (SST) from OSTIA (Donlon et al., 2012) is run for one year until 1. January 2011, to use as an initial state.

In the CICE model, the sea-ice variables are distributed in 5 thickness categories, while all observations are single category values. This discrepancy was solved by assimilating the aggregated category values and using the EnKF correlation properties to update each category individually. After assimilations, the analysis results are post-processed before running new forecasts. During post-processing, it is verified that the consistency of the different ice variables is maintained during assimilation as the analysis can lead to for instance situations where some areas have a positive ~~SIC but no volume~~ partial SIC but the corresponding partial SIT is zero or less than zero, in this example the SIC is set to zero. In addition, all variable bounds are checked during post-processing. Due to linear correlation effects of the EnKF, locations with non-physical concentrations can occur, for instance, SIC values both above one and below zero.

For the ocean parameters, only ocean temperature and ocean salinity are updated during the assimilation. Experiments have shown that large instantaneous changes to the ocean parameters lead to model instability. These large changes are especially seen in the marginal ice zone (MIZ) where the ensemble spread is the largest and the update to the ensemble is the strongest. To prevent these instabilities in the ocean the magnitude of the ocean update during an assimilation step is limited. In this work, a maximum temperature update step of 0.2 K for the ocean surface layer and 0.1 K for all other ocean layers is chosen. Similarly for the salinity 0.2 for the surface layer and 0.1 for all other layers is chosen. The limits are chosen crudely, based on values where the model did not immediately crash after assimilation. Although this is a crude simplification, almost omitting ocean update, it is believed to be sound, because the focus in this research is on the sea ice, and because it is implemented consistently for all the model experiments.

Table 1. Overview of the 5 experiments used to ~~asses~~assess observation impact, the X marks if the given observation is assimilated in the experiment.

	OSISAF	Cryosat <u>CryoSat-2</u>	SMOS	Snow depth
Exp1 (SIC)	X			
Exp2 (SIC + SIT _I)	X	X		
Exp3 (SIC + SIT _R)	X		X	
Exp4 (SIC + SD)	X			X
Exp5 (Control)				

Five ~~assimilation~~ experiments assimilating different observations are used to investigate the observations effect on the model. In experiment 1 only OSISAF SIC is assimilated, in exp. 2 both OSISAF SIC and ~~CryoSat~~CryoSat-2 SIT, in exp. 3 both OSISAF SIC and SMOS SIT, in exp 4. OSISAF SIC and snow depth (SD) observations, exp 5. is a control run without assimilation. All assimilation systems are initialized after one year of initial assimilation on 1. January 2011 and run for three
5 years. A summary of the different experiments is shown in table 1.

5 Results

In this section, the output of the five ensemble experiments is compared. All results are based on the output from 2011-2013. As mentioned, the first year of modelling, 2010, is only used to spin-up the experiments, generating a stable and consistent ice-ocean model state.

10 Many of the results shown in this section will be based on the root mean squared error (RMSE). In this study, the RMSE is weighted by the observation uncertainty, $\sigma_{Obs(i)}$,

$$RMSE = \sqrt{\frac{\sum_{i=1}^N \frac{(Mod(i) - Obs(i))^2}{\sigma_{Obs(i)}^2}}{N}}, \quad (5)$$

where N is the number of grid cells, Obs and Mod are the observations and model values, respectively. Thus an RMSE of one indicates that the difference between model and observations are on average of the same order as the observation uncertainty.

15 5.1 Validation against concentration observations

In Figure 2 the monthly averaged ensemble mean of the five experiments validated against two different SIC observation products, one assimilated and one independent is plotted. In Fig 2a the RMSE values of the ensemble mean of the modelled SIC validated with the observed AMSR-E/2 product are plotted after assimilation. All four assimilation experiments are found to be significant improvements compared to the control experiment without assimilation. Using a one-sided paired sample student t-test over all 36 months of simulation both the ~~Cryosat~~CryoSat-2 internal SIT and SMOS rim SIT experiments show significant improvements compared to the OSISAF SIC only experiment on a 5 % level, but the differences are relatively small.




Figure 2. The monthly averaged RMSE of the ensemble mean SIC over three years. In a) the model is validated against AMSR-E/2 SIC observations and in b) OSISAF SIC observations. The lines represent different experiments, black: only SIC assimilation, blue: SIC and [CryoSat-CryoSat-2](#) thick internal SIT assimilation, red: SIC and SMOS and thin rim SIT assimilation, yellow: SIC and snow depth (SD) assimilation, magenta dotted: without assimilation.

The significance is derived using monthly data, but not yearly averaged as in the figures. However, the snow experiment is not found to be significantly better than the OSISAF SIC only experiment on a 5 % level, a p-value of 0.23 is found. The difference between the SIT experiments and the SIC only experiment is largest during the first half of the year, while in the second half of the year all experiments give similar results with a peak in the RMSE in October-November. This peak in RMSE is also seen in the control model, indicating a possible model problem related to the transition from melt season to growing season.

In Fig. 2b the monthly averaged RMSE of the model SIC ensemble mean versus the assimilated OSISAF SIC observations is plotted. The result in Fig. 2b is similar to that of 2a, but the differences between the models are larger when verified against OSISAF, even though the OSISAF observations are assimilated in all experiments. This is partly related to lower observation error in the MIZ for the OSISAF dataset than the AMSR-E/2 dataset, and that the OSISAF includes almost an extra year of observations, due to the AMSR-E/2 gap. Since the RMSE values are weighted by the observation error the differences in the MIZ are more pronounced when verified against OSISAF SIC observations. In addition, evidence that there are small differences between the two observation products is seen by different shapes on the graphs, even though the curves follow the same trends. As mentioned, the [CryoSat-CryoSat-2](#) and SMOS SIT experiments are significantly better than the OSISAF SIC only experiment. When compared to the OSISAF observations also snow depth assimilation experiment is found to be significantly better than the OSISAF SIC only experiment, especially during the first half of the year there are significant differences. In conclusion, assimilating SIT and to some degree, snow depth has a significant effect on the SIC RMSE, and the effect is largest for the first half of the year. In the transition from melting ice to freezing ice, all four experiments give similar high RMSE values.

RMSE estimates are sensitive to individual measurements contributing to large portions of the total RMSE, thus a small area with large errors will obscure the overall model results. Another assimilation quality measure is hit rates, where all grid cells are given equal weight in the analysis. In our work, the hit rate is analysed by classifying the SIC in three categories, open water (concentration less than 10 %), low concentration (< 50 %), and high concentration (> 50 %). The hit rate is defined as the number of grid cells correctly classified. The independent AMSR-E/2 observations are used for verification. In Fig. 3a the number of grid cells correctly classified is shown; in Fig. 3b the number of grid cells with modelled ice and observed water; in Fig. 3c the number of grid cells with modelled water and observed ice; in Fig. 3d the number of grid cells with wrong concentration category, high SIC classified as low SIC and vice versa. All assimilation experiments outperform the control run in terms of hit rate. The control run has a large number of false positives, indicating too much ice. Among the experiments, the variations are small in spring, fall and winter, while summer shows significant differences. In summer all experiments have a minimum, this minimum is related to an under-prediction of sea ice and wrong classification of concentration in observations due to melt ponds on ice which leads to an underestimation of SIC in the observations (Kern et al., 2016). In summer the CryoSat-CryoSat-2 assimilation has the highest number of hits, closely followed by the SMOS and snow experiments.

5.2 Total extent and volume

In figure 4, the total sea ice extent (4a), the total sea-ice volume (4b), and the total sea-ice volume overlapping the area and period covered by the CryoSat-CryoSat-2 internal SIT observations (4c) are shown for the five experiments. Figure 4a shows that the control experiment has a too large sea-ice extent both in summer and winter, while the assimilation experiments have a slightly too large ice extent in winter.

The total sea-ice volume shown in Fig. 4b indicates large differences between the five experiments. Snow depth assimilation generally leads to thicker ice. The model has a lower amount of snow than the observations and due to positive correlation, the ice thickness is also increased during the assimilation of snow depth. The increased thickness can be seen by the fact that the snow depth experiment has about the same extent as the other experiments, but show a significantly larger ice volume, both in summer and winter for all three years. Both the SMOS and CryoSat-CryoSat-2 ice thickness experiments lead to thinner sea ice compared to the control experiment. Especially the SMOS assimilation model shows much thinner sea-ice than do the other assimilation experiments. The thin SIT observations have a very strong effect on the modelled SIT, seen by an abrupt update of sea-ice volume during assimilation in winter. A concerning effect of the assimilation experiments is the strong decrease in the Arctic sea-ice volume throughout the period of study. The sea-ice volume maximum in winter is decreasing for every year of assimilation, this is not seen in the control run.

In Fig. 4c the modelled sea-ice volume is compared to the sea-ice volume in the combined CryoSat-SMOS-CryoSat-2-SMOS product. The control model is found to have too thick ice compared to the observations, while the experiments assimilating SIT are much closer to the observations, though largely biased. ~~It was seen that the~~ This can be used to explain the drastic decrease in sea-ice volume found in Fig. 4b can be explained by the volume adjusting to the observed SIT. The model SIT is adjusting towards the observations by rapidly thinning the sea-ice. For the OSISAF SIC only assimilation experiment, the volume is also slowly diverging towards the observed volume, even though SIT is not assimilated. This is likely related to a more accurate




Figure 3. Classification of the model result based on three classes, high concentration ice ($> 50\%$), low concentration ice ($> 50\% < 50\%$) or water and compared to AMSR-E/2 SIC observations. The figures show a) the fraction of correctly classified grid cells, b) the fraction of grid cells with modelled ice while water is observed, c) fraction of grid cells with modelled water while ice is observed, and d) fraction of grid cells where the model and observations have different SIC classification. The colour coding in the figure is the same as that of Fig. 2. These panels cover all possible classifications, thus the sum of them equals to one.




Figure 4. The evolution of a) total sea-ice extent, b) total sea-ice volume, and c) total sea-ice volume for the area covered by the [Cryosat-SMOS-CryoSat-2-SMOS](#) SIT observation product. The coloured lines represent the same as in Fig. 2. In a) the black stars represent the AMSR-E/2 SIC observations and the [blue-red](#) stars the OSISAF SIC observations. In b) same as a) without observations. In c) the black stars represent the [Cryosat-SMOS-CryoSat-2-SMOS](#) observation product. [The xlabel is given as month-year.](#)

sea-ice extent also leads to improved ice thickness in the marginal ice zone. However, the improvements are obtained at a slower pace than when assimilating SIT directly.

5.3 Validation against thickness observations

In Fig. 5a the SIT RMSE of the ensemble [averaged-mean](#) modelled SIT is verified with the combined [SMOS-CryoSat](#)
 5 [SMOS-CryoSat-2](#) SIT product. The experiment assimilating SMOS thin SIT has significantly lower RMSE values than the other three assimilation experiments. The other three experiments are more similar, all showing high RMSE values. It is found using a one-sided paired student t-test that only the SMOS SIT experiments are significantly better than the SIC only assimilation, with p-values less than 5 %. Due to the high RMSE values, only small improvements compared to the control run is seen.

The result is consistent with what was found for the sea-ice volume (Fig. 4c), regarding the SMOS SIT assimilation having the strongest effect on the modelled SIT. The reason for the high RMSE values is that in general, the model has ~~to~~too much ice, leading to too thick ice in the MIZ. For the ~~SMOS-CryoSat~~SMOS-CryoSat-2 SIT product the uncertainties provided are very small, especially in the MIZ where the SMOS observations are used, thus when calculating the RMSE these values have a huge effect on the result. Thus it is also reasonable that the assimilation system assimilating these MIZ thickness observations also give the lowest RMSE values. For the other assimilation systems, the ice extent is updated in the MIZ, but the thickness reduction takes longer because this has to evolve over time.

As for the SIC observations, the RMSE values are biased by locations showing large differences. Particularly for thickness which is not bounded upwards, a few grid cells in the MIZ can contribute to a large total RMSE. As for concentration, an alternative measure where correctly classified model thickness hit rates are used. The model is separated into four thickness categories, less than 0.5 m, between 0.5 m and 1.5 m, between 1.5 m and 3 m, and above 3 m. In Fig. 6a the number of correctly classified ice thicknesses grid cells is plotted for each experiment. The figure shows that the ~~CryoSat-CryoSat-2~~CryoSat-CryoSat-2 internal SIT experiment is the model which has the highest number of correctly classified grid cells. The other experiments are more similar, except in spring where the SMOS rim SIT assimilation is equally good as the ~~CryoSat-CryoSat-2~~CryoSat-CryoSat-2 internal SIT assimilation, and both much better than the other three. In spring the SIC only and snow depth assimilations are not improved compared to the control case. In general, the model shows too much ice. This can be seen by a large number of grid cells having too thick ice in the control model (Fig. 6b). This is a combination of the sea-ice extent being too large and the ice ~~is~~being too thick. By assimilating observations the ice volume is reduced, not only for the SIT assimilations, but also for the snow depth and SIC only assimilations, but to a lower degree for the latter. This is an effect of a lower sea-ice extent (Fig. 4a). In Fig. 6c the number of grid cells with too thin ice compared to the observations is shown. It was found that this problem is large in early winter for all experiments but reduces during winter for all experiments except the SMOS experiment. During SMOS assimilation, only thin ice is assimilated which might lead to a bias towards the thinner ice, causing a relatively high number of grid cells with too thin ice.

As an example, in Fig. 7 the absolute differences between the experiments and the combined ~~CryoSat-CryoSat-2~~CryoSat-CryoSat-2 SMOS ice thickness observations are plotted for 1. April 2011. Figure 7 is consistent with Fig. 6a showing that the ~~CryoSat-CryoSat-2~~CryoSat-CryoSat-2 experiment has the smallest differences compared to the observations in the internal Arctic, affecting a large area, however, large differences can be seen in MIZ. While for the SMOS rim SIT assimilation the effect is the opposite, with large impact in the MIZ, and low impact in the ice interior. This shows that assimilating SIT is important both in the interior and in the MIZ.

In addition to the satellite observations, the independent airborne IceBridge dataset is used for verification of the modelled SIT (Kurtz et al., 2013, 2014a). This dataset has low temporal and spatial distribution, but is believed to have higher accuracy and has a much higher spatial and temporal resolution. All observations occurring in March and April are gathered as a yearly averaged observation as a function of space. These yearly observations are then compared to modelled SIT averaged over the same period for the observed IceBridge locations. Since the IceBridge resolution is much higher than that of the model, all IceBridge observations within one model grid cell are averaged and used for verification, the average is done by a weighted mean based on the observation uncertainty. The validation results are shown in table 2. On average the ~~CryoSat-CryoSat-2~~CryoSat-CryoSat-2 SIT




Figure 5. RMSE of monthly averaged model SIT and snow depth averaged over all ensemble members [for the years 2011-2013](#) calculated against [observed-the](#) a) [ice thickness-combined SMOS-CryoSat-2 SIT product](#) and b) [snow depth](#)[observed snow-depth product](#). [These are observations also used for assimilation.](#) The colour coding is as in Fig. 2.

Table 2. The [yearly-Mar/Apr](#) averaged RMSE values of the five experiments compared to the IceBridge aerial SIT observations. Bold values represent the model with the lowest RMSE values for that year.

	2011	2012	2013	2011-2013
SIC	0.88	0.87	1.11	0.94
SIC+SIT _I	0.63	0.86	0.72	0.80
SIC+SIT _R	0.74	1.14	0.87	0.96
SIC+SD	0.93	1.38	1.64	1.51
Control	0.82	1.25	2.31	1.38
Cryo Obs	0.67	0.95	0.84	0.84

experiment has the best SIT estimation as compared to IceBridge. Both the SMOS and the [CryoSat-CryoSat-2](#) SIT experiments give on average thinner SIT than the IceBridge observations, consistent with the findings of Chen et al. (2017). The last line in the table shows the RMSE between the [CryoSat-CryoSat-2](#) observations and the IceBridge observations and the results show that the error is comparable to the model errors.

- For all three years, the [CryoSat-CryoSat-2](#) assimilation has lower RMSE values than the [CryoSat-CryoSat-2](#) observations, indicating a well-balanced assimilation, with appropriate observation error and ensemble spread. It should also be mentioned that the [CryoSat-CryoSat-2](#) observations have less spatial coverage than the model and not all IceBridge observations are covered, thus the number of useful observations for the [CryoSat-CryoSat-2](#) RMSE calculation is smaller than for the validation of the experiments.




Figure 6. The monthly mean SIT averaged over all ensemble members is classified into four thickness categories and compared to the [CryoSat-SMOS-CryoSat-2-SMOS](#) SIT observation product. The fraction of grid cells are shown with a) correctly classified thickness category, b) too thick ice, and c) too thin ice. As in Fig. 2 the coloured lines represent different experiments.

Another independent dataset of SIT observations complementing the IceBridge observations by a temporal resolution spanning the entire year is the IMB buoy dataset. The result of model validation with the IMB product is shown in table 3. For these observations, a slightly different method than that applied for IceBridge is performed. This is because IceBridge temporarily only covered March-April, while the IMB data span the entire year. The buoy observations are converted to daily averages on the model grid. From these values, the RMSE is calculated on the 7-day ensemble mean and averaged for each year. Since SIT is a relatively slow varying variable, for each 7-day output, observations from +/-3 days are used to increase the number of observations. The IMB observations do not include an uncertainty estimation, hence the RMSE is not normalised as was the case for other RMSE estimates in this work. The results show that over the three study years, the SMOS internal SIT assimilation system has the lowest RMSE values, followed by the CryoSat-2 internal SIT assimilation. The other three show similar results, again indicating a positive impact of assimilating ice thickness in the model.




Figure 7. Absolute differences between the experiments and the combined [SMOS-CryoSat](#) [SMOS-CryoSat-2](#) observation product are given on 1. April 2011. The experiments are assimilating a) OSISAF SIC, b) OSISAF SIC and [CryoSat](#) [CryoSat-2](#) SIT, c) OSISAF SIC and SMOS SIT and d) OSISAF SIC and snow depth.

Table 3. [The yearly averaged RMSE values of the five experiments compared to the IMB SIT buoy observations. Bold values represent the model with the lowest RMSE values for that year. No uncertainties are used to normalise the RMSE values.](#)

	2011	2012	2013	2011-2013
SIC	0.99	1.45	1.32	1.27
SIC+SIT_I	1.08	1.28	1.00	1.13
SIC+SIT_R	1.09	1.07	0.99	1.05
SIC+SD	1.02	1.40	1.28	1.25
Control	1.46	1.27	1.23	1.26

Table 4. The ~~yearly-averaged-Mar/Apr-mean~~ RMSE ~~values-of-an-ensemble-average-of~~ ~~the-ensemble-mean~~ snow depth averaged over all ~~the-ensemble-members-for-the-grid-cells.~~ The five experiments ~~and the snow-depth satellite observations are~~ compared to the IceBridge airborne ~~snow-depth-snow-depth~~ observations. Bold values represent the model with the lowest RMSE values for that year.

	2011	2012	2013	<u>2011-2013</u>
SIC	0.79	1.38	2.64	<u>1.63</u>
SIC+SIT _I	0.79	1.15	1.44	<u>1.06</u>
SIC+SIT _R	0.78	0.83	1.73	<u>1.17</u>
SIC+SD	0.74	1.22	1.46	<u>1.13</u>
Control	0.77	2.49	1.85	<u>1.33</u>
Snow Obs	1.46	NA	1.17	<u>1.16</u>

5.4 Validation against Snow observations

In Fig. 5b the RMSE of monthly averaged modelled snow depth over all ensembles validated against the observed satellite snow depth (Rostosky et al., 2018) used for assimilation is plotted. The control experiment is found to have the lowest RMSE values. This is most likely an effect of sea-ice extent being different compared to the assimilation experiments, rather than the assimilation decline the accuracy of the modelled snow depth. In addition, the control experiment has an increasing RMSE during the period, while the assimilation experiments show the effect of assimilation by decreasing the RMSE. For the assimilation experiments, the snow experiment has the lowest RMSE values followed by the ~~CryoSat~~ CryoSat-2 experiment, indicating that the thick ice observations have a correlation effect on the snow depth. These two observation products also cover a similar area of the Arctic ocean.

10 A verification of the modelled snow depth using the independent IceBridge dataset is given in table 4. The same method as for the SIT in table 2 was used, where Mar/Apr model values are compared to the IceBridge observations and averaged. It is found that none of the experiments is particularly better than any of others when verified against IceBridge snow depth observations. ~~This lack of improvement can be an indication of a too simple snow component in our coupled system, only one snow layer is used. It was also~~ It is seen that within one grid cell ~~there where~~, there are huge variations in the IceBridge snow observations, ~~indicating the difficulty of modelling snow.~~ Such variations cannot be provided with a coarse resolution model. Hence, even though IceBridge is used to "tune" the snow observations (Rostosky et al., 2018), large RMSE values are estimated for the experiment assimilating snow depth. In addition, the snow component used in our coupled system is likely too simple, having only one snow layer, which may effect the snow cover accuracy. It is also important to mention that the snow observations are in ~~the-an~~ an early development stage and might have larger uncertainties than what is used in this study.

20 Additional model verification is performed with the independent IMB buoy snow depth observations. The method of validation is performed in a similar manner as for SIT validation with IMB buoy data, the results are shown in table 5. As the IMB data does not include an uncertainty these RMSE values are not normalised, thus they are significant lower than the error estimates from the ice bridge validation in table 4. From the table it is clear that the differences between the assimilation

Table 5. The yearly averaged RMSE values of the five experiments compared to the IMB snow depth buoy observations. Bold values represent the model with the lowest RMSE values for that year. No uncertainties are used to normalise the RMSE values.

	<u>2011</u>	<u>2012</u>	<u>2013</u>	<u>2011-2013</u>	
<u>SIC</u>	<u>0.06</u>	<u>0.15</u>	<u>0.2</u>	<u>0.15</u>	=11
<u>SIC+SIT_I</u>	<u>0.08</u>	<u>0.14</u>	<u>0.17</u>	<u>0.13</u>	
<u>SIC+SIT_B</u>	<u>0.09</u>	<u>0.15</u>	<u>0.17</u>	<u>0.14</u>	
<u>SIC+SD</u>	<u>0.09</u>	<u>0.14</u>	<u>0.16</u>	<u>0.13</u>	
<u>Control</u>	<u>0.09</u>	<u>0.16</u>	<u>0.19</u>	<u>0.15</u>	

systems are small, the assimilation systems assimilating snow depth and CryoSat-2 internal SIT are slightly better than the others, but the differences are too small to conclude.

5.5 One week forecasts

Figure 8 shows the RMSE of the monthly averaged modelled SIC over all ensemble members before assimilation validated against the AMSR-E/2 and OSISAF SIC observations. Since the assimilation time step is seven days, this gives the accuracy of a seven-day forecast. The comparison against AMSR-E/2 SIC observations (Fig. 8a) shows that the differences between the experiments are small, and the differences are similar to those found after assimilation (Fig. 2a). In general, the system with the most accurate initial state also gives the most accurate forecasts. Thus the ~~CryoSat~~ CryoSat-2 and SMOS SIT assimilation experiments have a better seven-day forecast from January to June than SIC only, and snow depth assimilation shows improvements from January to April. Using the OSISAF SIC observations (Fig. 8b) gave the same result as found for AMSR-E/2: the best initial states also provide the best forecast, indicating that the sea-ice overall does not change much in a week. The same experiments were also done for ice thickness and snow depth and similar results were encountered. A reason for the small differences between the different experiments is the coarse model resolution. Large-scale variations as seen by a 20 km model are not expected within a week.

6 Seasonal forecast

In the previous section, it was found that our coarse resolution model only exhibits small changes during a one week forecast. Thus a more interesting forecast would be one of seasonal length. A five-month forecast of the September sea-ice extent is performed. This is done by running each of the experiments from ~~April to September without assimilation~~ mid-April to mid-September each year without assimilation, and validating against the OSISAF SIC observations. The actual start date varied slightly from year to year because of the 7-day assimilation cycle, but the start date was the same for all experiments. In Fig. 9a, the RMSE of three five-month forecasts are shown sequentially, and a monthly averaged RMSE over the three years is shown in Fig. 9b. The figures show that ~~that~~ the experiments have very similar seasonal forecasts, with some differences in late summer. In general, the model error is gradually increasing towards the level of the control run, and in summer they have




Figure 8. RMSE of a seven-day-monthly averaged (over three years) ensemble average mean of modelled SIC seven-day forecast SIC validated by against a) AMSR-E/2 SIC observations and b) OSISAF SIC observations.




Figure 9. Seasonal forecast of summer sea-ice extent. Each forecast started at the beginning of April every year. The figures show SIC-RMSE of the ensemble mean SIC averaged over all ensemble members 3 years and verified against the assimilated OSISAF SIC. The coloured lines represent the same as in figure 2. In a) Full period, in b) monthly averaged values.

similar error levels. In August/September the experiments assimilating thickness and concentration seems to be improvements compared to without assimilation and assimilating snow depth observations. All experiments show an increased RMSE in 2013, this is related to a too low sea-ice extent. The low sea-ice extent is caused by a weaker modelled ice growth compared to observations in the first months of 2013.




Figure 10. Seasonal forecast of summer sea-ice extent both with climatological forcing and re-analysed forcing. Each forecast started at the beginning of April every year. The figure describes monthly averaged RMSE SIC averaged over all ensemble members. The blue line represents a forecast using a climatological forcing made from ~~an over of~~ atmospheric data ~~for over~~ 2000-2014 with assimilation, the black line using re-analysed atmospheric forcing and assimilation, and the dotted magenta line use re-analysed forcing only.

The seasonal forecast is compared to a climatological seasonal forecast in Fig. 10. This provides an estimate of the expected sea-ice forecast accuracy. The climatological experiment is done by running the model with averaged atmospheric forcing data over the years from 2000 to 2014. The result shows that the forecast skill of the model is rapidly decreasing and that a correct atmospheric forecast is very important for an accurate sea-ice forecast. But still, skills are evident on much longer time-scales
 5 that can be obtained with numerical weather prediction models.

7 Discussion

Significant differences in modelled SIC after assimilation was found, especially in the first half of the year. The SMOS and ~~CryoSat-CryoSat-2~~ SIT assimilations gave the lowest RMSE values, significantly better than assimilating OSISAF SIC only. The snow depth experiment showed improvements during the first half of the year compared to the experiment assimilating
 10 OSISAF SIC observations only. In addition, assimilating SIT and snow depth lead to an improved model of SIC in summer, where the ~~CryoSat-CryoSat-2~~ internal SIT assimilation gave the highest number of correctly classified grid cells, closely followed by the SMOS rim SIT and snow depth assimilations. The reason for these differences in summer is that the pace in which the ocean becomes ice-free is dependent on the ice thickness and the snow depth. In the second half of the year, fall and
 15 season to growing season not well represented in the model. The observed transition is faster than the modelled, leading to an extended period with more open water in the model than in the observations.

In the control model without assimilation, the ice extent both in summer and winter was found to be larger than observed. However, with assimilation, the experiments are closer to the observed extent, even though a slight overestimation of extent

in winter was found for the first two years. The sea-ice extent overestimation in winter is a result of a lower effect of SIC assimilation in winter due to lower ensemble spread. When the ensemble spread is low the EnKF assimilation result is closer to the model values, because the estimated model errors become small.

It is found that originally the sea-ice volume is too large compared to the observations, and over the three years, the sea-ice volume in the assimilation experiments are gradually decreasing towards the observed values in the [SMOS-CryoSat](#) [SMOS-CryoSat-2](#) SIT product. The effect is much stronger for the SMOS rim SIT assimilation, indicating that a large portion of the original sea-ice volume overestimation is located in the MIZ. This is a consequence of too much ice in the control model causing the observed MIZ to be located deeper into the Arctic as compared to the model, as noted by Fritzner et al. (2018).

In the verification of modelled SIT (Fig. 5a), the SMOS rim SIT assimilation was found to give the lowest RMSE values, while the [CryoSat-CryoSat-2](#) internal SIT assimilation had the largest amount of correctly classified thickness grid cells. This is as expected since even though the [CryoSat-CryoSat-2](#) observations cover a larger area, they are 30-day averaged observations with much larger uncertainties than the SMOS observations. In addition, the non-updated grid cells in the MIZ leads to larger RMSE values than non-updated grid cells in the internal Arctic, where the model, in general, is more accurate and less sensitive to changes. When verified by IceBridge observations which only covers the central Arctic, the [CryoSat-CryoSat-2](#) SIT assimilation experiment was found to give the lowest SIT RMSE values. The [CryoSat-CryoSat-2](#) SIT observations are in general thinner than the SIT values for the SIC only experiment. In comparison with the IceBridge observations, the [CryoSat-CryoSat-2](#) SIT is biased low, which was also found by Chen et al. (2017). When assimilating snow depth, it was found that snow depth observations, in general, were thicker than those modelled, resulting in increased snow depth during assimilation. Due to the correlation nature of the EnKF, a positive correlation between snow depth and SIT resulted in increased SIT in the snow depth assimilation experiment compared to the other assimilation experiments.

Validating our experiments with snow observations proved the control run to have the lowest RMSE values, this can be an effect of a different sea-ice extent in the control run than in the assimilation experiments. For the control model, the ice extent is too large, thus collecting more snow on the ice than the assimilation experiments. When the ice concentration is reduced during assimilation, the accumulated snow is also removed, this can result in removal of too much snow if the ice extent is less than it should be. A verification of the impact of assimilation on the snow depth is that the RMSE is decreasing throughout the observation period for the assimilation experiments, while for the control run the RMSE is increasing. Between the assimilation experiments, the snow depth assimilation was found to give the lowest snow depth RMSE values, not unexpected since the same dataset is used for assimilation and verification. More interestingly the [CryoSat-CryoSat-2](#) internal SIT experiment has significantly lower RMSE values than the SMOS rim SIT and OSISAF SIC only assimilations, indicating a close correlation between SIT and snow depth. A curiosity here is that the SIT assimilation has a positive effect on the snow depth, while it was found previously that the snow depth assimilation had a negative effect on the SIT. This is likely an effect of more SIT observations than snow depth observations, and SIT is assimilated throughout the whole winter. It could be the case that the correlation relationship between snow depth and SIT changes throughout the winter. This results in a better snow depth estimate, while for snow depth assimilation the assimilation period is limited to Mars-April. In addition, when the assimilation is only in two months of the year, the model error is larger when the assimilation period starts, thus the assimilation update

has a large effect both on snow depth and SIT. An indication of the relation between SIT and snow depth is also seen by lower snow depth RMSE value for the SMOS rim SIT assimilation than the OSISAF SIC only assimilation. Since the SMOS system covers a much smaller area and has less overlap with the snow dataset than the [CryoSat-CryoSat-2](#) internal SIT, the effect of assimilation on the modelled snow is smaller.

5 When validating our experiments with the IceBridge snow depth observations, none of the experiments showed any improvements compared to the others. This can be related to an underestimated uncertainty in the snow observations, or that the snow representation in the model is too simplistic, only utilizing a single layer. Another problem is local variations, the coupled model is coarse with a resolution of 20 km, but as can be seen from the IceBridge observations the snow depths have large spatial variations in this range. This causes high RMSE values, both compared to satellite observations (on the model grid) and the modelled snow depth values. In addition only four months of snow depth observations were available for assimilation during the three years.

For sea ice, the model drift is in general small, the model system with the best initial state provides the best short-term forecast. The main parameters analysed in this study, snow depth, SIT and SIC, all vary on longer time scales than one week for the spatial resolution in our model. Thus the correlation between day-one and day-seven is too strong, as also shown by Chen et al. (2017) the sea-ice drift is low.

Several five-month seasonal forecasts of the September sea-ice extent showed small differences between the assimilation experiments. All experiments showed a steady increase in RMSE with time. This is likely caused by the model overgrowth of ice. The seasonal forecasts showed that after 3-4 months the RMSE values were found to be of the same order as those in the control run. Thus assimilation gives at least an improvement over 3-4 months, and the September result suggests that with the assimilation of SIC and SIT there are improvements in the Arctic sea-ice extent compared to the control run on even longer timescales, this was not seen for the snow depth experiment. The seasonal forecast was compared to a climatological run, and it was found that without accurate forcing the forecast degenerates fast.

In this work, four different observation products have been used for assimilation. The different products differ widely in both temporal and spatial coverage in addition to accuracy. There is no doubt that it is preferable to have as much coverage and as accurate observations as possible. Where a realistic observation error is a necessity for the assimilation, without a realistic observation error the observation is not useful. E.g. the [Cryosat-CryoSat-2](#) product does not provide an observation error and a uniform error was chosen, which will lead to some observations given too much weight and others with too little. In this study, the spatial resolution of the observations is not a problem, because the model resolution is coarse, but in the future when the model resolution increases, there will be an increasing demand for high-resolution SIT observations. Both SIT products are only available in winter, and temporal coverage of the snow depth observations are limited to four months out of a three-year experiment. Thus for these products to be even more useful, there is a strong need for increased seasonal coverage, especially in summer when the Arctic sea-ice extent is at a minimum and there is ship traffic there. Observations can then help to improve the models, thereby helping planning operations and decrease the risks. In addition, because there are few snow observations available for assimilation, there are large unknown aspects regarding the assimilation effect. Finally, it should be remembered

that the model itself has in general too much ice and that the forcing is known to contain biases and errors (Jakobson et al., 2012).

8 Conclusions

In conclusion, we have found that assimilation of more observation types than SIC into coupled sea-ice ocean models can lead to significant model improvements. We show that especially the assimilation of SIT leads to improvements in modelled SIC, SIT and snow depth, for long-term model improvement. There is a clear evidence that assimilation of SIT gives a better representation of the full ice state and we recommend that they are assimilated into models when available. Even though SIT seems to be an important variable for improving sea-ice modelling, it still has several limitations in terms of spatial and temporal resolution and realistic observation errors. It is important to emphasize that assimilation of SIC is vital to the assimilation update, these observations give important information with full Arctic coverage about where the ice is located. In addition the fractional ice in the MIZ is important for the forecast in terms of how the short-term changes of the sea-ice will look like, and also and indication of the thickness.

Assimilation of snow depth was found to have a weaker effect on the model than assimilating SIT, but improvements to modelled SIC and modelled snow depth were found. In addition, we found a strong correlation between SIT and snow depth which should be analysed further when more observations from other months become available. The low efficiency of snow depth observations can be related to ~~a too simple snow component with only a single layer used or the~~ low model resolution, the IceBridge observations show large snow depth variations within a model grid cell. It is also important to keep in mind that the snow depth observations are in an early development stage, and ~~we should expect improvements in the future~~ the uncertainty estimations might not be correct. Possibly inaccurate observations or a wrong uncertainty estimation can have a huge impact on the assimilation result. Due to the small temporal coverage in our study more investigation has to be done on the effect of assimilating snow depth observations.

As mentioned the assimilation of SIT leads to an improved model, which leads to improved short-term forecasts over time, because the initial states are better represented. For seasonal forecast, we found that the model improvements due to assimilating observations have a memory of at least 3-4 months, and possibly even longer. Assimilating SIC and SIT showed improvements of the September ice forecasts compared to assimilating snow depth and no assimilation.

Comparing the two SIT products, SMOS thin SIT and CryoSat-2 thick SIT, we see that in general the CryoSat-2 observations gives the best long-term model improvement, especially decreased RMSE values for the model in summer. It is expected that SMOS observation assimilation should be better for the short-term forecast, but we were not able to provide any results on this. This could be an effect of too low model resolution, with 20 km resolution the ice state does not change much during a week of simulation.

Competing interests. No competing interests.

Acknowledgements. We would like to thank Pavel Sakov for help using and implementing the EnKF-c code and for helpful discussions regarding the EnKF. We thank the EUMETSAT OSISAF centre and the NASA National Snow and Ice Data Center for providing the sea-ice concentration data, and the Integrated Climate Data Center at the University of Hamburg for the SMOS ice thickness observations. We would like to thank NASA National Snow and Ice Data Center for providing the Cryosat sea-ice thickness observation, and the Icebridge snow and sea-ice thickness observations.

This work was funded through the Center for Integrated Remote sensing and Forecast for Arctic Operations through the Norwegian Research Council grant no. 237906. Rostosky was supported by the Transregional Collaborative Research Center (TR 172) "Arctic Amplification: Climate Relevant Atmospheric and Surface Processes, and Feedback Mechanisms (AC)³", which is funded by the German Research Foundation (DFG). Two supercomputers provided by the Norwegian Metacenter for Computational Science (NOTUR) was used for the computational work, the Vilje computer at the Norwegian University of Science and Technology was used with project and the Stallo computer at the University of Tromsø both under project NN9348K.

References

- Allard, R. A., Farrell, S. L., Hebert, D. A., Johnston, W. F., Li, L., Kurtz, N. T., Phelps, M. W., Posey, P. G., Tilling, R., Ridout, A., et al.: Utilizing CryoSat-2 sea ice thickness to initialize a coupled ice-ocean modeling system, *Advances in Space Research*, 2018.
- Andersen, S., Tonboe, R., Kern, S., and Schyberg, H.: Improved retrieval of sea ice total concentration from spaceborne passive microwave observations using Numerical Weather Prediction model fields: An intercomparison of nine algorithms., *Remote Sens. Environ.*, 104, 374–392, 2006.
- Bell, M., Barciela, R., Hines, A., Martin, M., McCulloch, M., and Storkey, D.: The forecasting ocean assimilation model (FOAM) system, in: Elsevier oceanography series, vol. 69, pp. 197–202, Elsevier, 2003.
- Blockley, E. W. and Peterson, K. A.: Improving Met Office seasonal predictions of Arctic sea ice using assimilation of CryoSat-2 thickness, *The Cryosphere*, 12, 3419–3438, 2018.
- Burgers, G., van Leeuwen, P., and Evensen, G.: Analysis Scheme in the Ensemble Kalman Filter, *Mon. Weather Rev.*, 126, 1719–1791, [https://doi.org/10.1175/1520-0493\(1998\)126<1719:ASITEK>2.0.CO;2](https://doi.org/10.1175/1520-0493(1998)126<1719:ASITEK>2.0.CO;2), 1998.
- Cavalieri, D., Markus, J., and Comiso, J.: AMSR-E/Aqua Daily L3 12.5 km Brightness Temperature, Sea Ice Concentration, & Snow Depth Polar Grids, Version 3, Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center., https://doi.org/http://dx.doi.org/10.5067/AMSR-E/AE_SI12.003, 2014.
- Caya, A., Buehner, M., and Carrieres, T.: Analysis and Forecasting of Sea Ice Conditions with Three-Dimensional Variational Data Assimilation and a Coupled Ice-Ocean Model, *J. Atmos. Oceanic Technol.*, 27, 353–369, <https://doi.org/10.1175/2009JTECHO701.1>, 2010.
- Chen, Z., Liu, J., Song, M., Yang, Q., and Xu, S.: Impacts of Assimilating Satellite Sea Ice Concentration and Thickness on Arctic Sea Ice Prediction in the NCEP Climate Forecast System, *J. Climate*, 30, 8429–8446, 2017.
- Comiso, J., Cavalieri, D., and Markus, T.: Sea Ice Concentration, Ice Temperature, and Snow Depth Using AMSR-E Data., *IEEE Trans. On Geosci. And Rem. Sens.*, 41, 2003.
- Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G., Bauer, P., Bechtold, P., Beljaars, A. C. M., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, I., Kållberg, P., Köhler, M., Matricardi, M., McNally, A. P., Monge-Sanz, B. M., Morcrette, J.-J., Park, B.-K., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J.-N., and Vitart, F.: The ERA-Interim reanalysis: configuration and performance of the data assimilation system, *Quart. J. Roy. Meteor. Soc.*, 137, 553–597, <https://doi.org/10.1002/qj.828>, 2011.
- Donlon, C. J., Martin, M., Stark, J., Roberts-Jones, J., Fiedler, E., and Wimmer, W.: The operational sea surface temperature and sea ice analysis (OSTIA) system, *Remote Sensing of Environment*, 116, 140–158, 2012.
- Eicken, H.: Ocean science: Arctic sea ice needs better forecasts, *Nature*, 497, 431–433, 2013.
- Evensen, G.: Sequential data assimilation with a nonlinear quasi-geostrophic model using Monte Carlo methods to forecast error statistics, *J. Geophys. Res.*, 99, 10 143–10 162, <https://doi.org/10.1029/94JC00572>, 1994.
- Evensen, G.: The Ensemble Kalman Filter: Theoretical Formulation and Practical Implementation, *Ocean Dyn.*, 53, 343–367, <https://doi.org/10.1007/s10236-003-0036-9>, 2003.
- Evensen, G.: The Ensemble Kalman Filter for Combined State and Parameter Estimation, *IEEE Control Syst. Mag.*, pp. 83–104, <https://doi.org/10.1109/MCS.2009.932223>, 2009.
- Flato, G. M. and Hibler, W. D.: Ridging and strength in modeling the thickness distribution of Arctic sea ice, *J. Geophys. Res-Oceans*, 100, 18 611–18 626, 1995.

- Forsberg, R. and Skourup, H.: Arctic Ocean gravity, geoid and sea-ice freeboard heights from ICESat and GRACE, *Geophysical Research Letters*, 32, 2005.
- Fritzner, S., Graverson, R., Wang, K., and Christensen, K.: Comparison between a multi-variate nudging method and the ensemble Kalman filter for sea-ice data assimilation, *J. Glaciol.*, pp. n/a–n/a, 2018.
- 5 Gaspari, G. and Cohn, S. E.: Construction of correlation functions in two and three dimensions, *Q. J. R. Meteorol. Soc.*, 125, 723–757, <https://doi.org/10.1002/qj.49712555417>, 1999.
- Houtekamer, P. L. and Zhang, F.: Review of the Ensemble Kalman Filter for Atmospheric Data Assimilation, *Mon. Weather Rev.*, 144, 4489–4532, <https://doi.org/10.1175/MWR-D-15-0440.1>, 2016.
- Hunke, E. and Dukowicz, J.: An elastic-viscous-plastic model for sea ice dynamics, *J. Phys. Oceanogr.*, 27, 1849–1867, 1997.
- 10 Hunke, E., Lipscomb, W., Turner, A., Jeffery, N., and Elliott, S.: CICE: the Los Alamos sea Ice Model Documentation and Software User’s Manual Version 5.1, Los Alamos National Laboratory, 2015a.
- Hunke, E., Lipscomb, W., Turner, A., Jeffery, N., and Elliott, S.: CICE: the Los Alamos Sea Ice Model Documentation and Software User’s Manual, 5.1, 2015b.
- Jakobson, E., Vihma, T., Palo, T., Jakobson, L., Keernik, H., and Jaagus, J.: Validation of atmospheric reanalyses over the central Arctic Ocean, *Geophys. Res. Lett.*, 39, 2012.
- 15 Jazwinski, A.: *Stochastic processes and filtering theory*, Academic, Sand Diego, California, 1970.
- Kaleschke, L., Tian-Kunze, X., Heygster, G., Patilea, C., Hendricks, S., Ricker, R., Tonboe, R., Mäkynen, M., Bertino, L., and Xie, J.: SMOS+SeaIce Final Report, ESA Support To Science Element (STSE) Contract No.: 4000112022/14/I-AM, version: August 28, Univ. Hamburg, Institute of Oceanography, 2017.
- 20 Kern, S., Rösel, A., Pedersen, L. T., Ivanova, N., Saldo, R., and Tonboe, R. T.: The impact of melt ponds on summertime microwave brightness temperatures and sea-ice concentrations, *The Cryosphere*, 10, 2217–2239, 2016.
- Kristensen, N., Debernard, J., Maartensson, S., Wans, K., and Hedstrom, K.: metno/metroms, <https://doi.org/10.5281/zenodo.1046114>, 2017.
- Kurtz, N. and Harbeck, J.: CryoSat-2 Level-4 Sea Ice Elevation, Freeboard, and Thickness, Version 1, Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center, [https://doi.org/10.1016/S1463-5003\(01\)00012-9](https://doi.org/10.1016/S1463-5003(01)00012-9), 2017.
- 25 Kurtz, N., Farrell, S., Studinger, M., Galin, N., Harbeck, J., Lindsay, R., Onana, V., Panzer, B., and Sonntag, J.: Sea ice thickness, freeboard, and snow depth products from Operation IceBridge airborne data, *The Cryosphere*, 7, 1035–1056, <https://doi.org/10.5194/tc-7-1035-2013>, 2013.
- Kurtz, N., Studinger, M., Harbeck, J., Onana, V., and Yi, D.: IceBridge L4 Sea Ice Freeboard, Snow Depth, and Thickness, Version 1, Boulder, Colorado USA. NASA National Snow and Ice Data Center Distributed Active Archive Center, <https://doi.org/https://doi.org/10.5067/G519SHCKWQV6>, 2014a.
- 30 Kurtz, N. T. and Farrell, S. L.: Large-scale surveys of snow depth on Arctic sea ice from Operation IceBridge, *Geophys. Res. Lett.*, 38, 2011.
- Kurtz, N. T., Galin, N., and Studinger, M.: An improved CryoSat-2 sea ice freeboard retrieval algorithm through the use of waveform fitting, *The Cryosphere*, 8, 1217–1237, 2014b.
- Kwok, R. and Rothrock, D.: Decline in Arctic sea ice thickness from submarine and ICESat records: 1958–2008, *Geophys. Res. Lett.*, 36, 2009.
- 35 Laxon, S. W., Giles, K. A., Ridout, A. L., Wingham, D. J., Willatt, R., Cullen, R., Kwok, R., Schweiger, A., Zhang, J., Haas, C., et al.: CryoSat-2 estimates of Arctic sea ice thickness and volume, *Geophysical Research Letters*, 40, 732–737, 2013.

- Lindsay, R. W. and Zhang, J.: Assimilation of Ice Concentration in an Ice-Ocean Model, *J. Atmos. Oceanic Technol.*, 23, 742–749, <https://doi.org/10.1175/JTECH1871.1>, 2006.
- Lisæter, K. A., Rosanova, J., and Evensen, G.: Assimilation of ice concentration in a coupled ice-ocean model, using the Ensemble Kalman filter, *Ocean Dynam.*, 53, 368–388, <https://doi.org/10.1007/s10236-003-0049-4>, 2003.
- 5 Lisæter, K. A., Evensen, G., and Laxon, S.: Assimilating synthetic CryoSat sea ice thickness in a coupled ice-ocean model, *J. Geophys. Res.*, 112, <https://doi.org/10.1029/2006JC003786>, 2007.
- Maaß, N., Kaleschke, L., Tian-Kunze, X., and Drusch, M.: Snow thickness retrieval over thick Arctic sea ice using SMOS satellite data, *The Cryosphere*, 7, 1971–1989, 2013.
- Markus, T. and Cavalieri, D.: Snow depth distribution over sea ice in the southern ocean from satellite passive microwave data, *Antarctic Research Ser.*, 74, 19–39, 1998.
- 10 Markus, T. and Cavalieri, D. J.: An enhancement of the NASA Team sea ice algorithm, *IEEE Transactions on Geoscience and Remote Sensing*, 38, 1387–1398, 2000.
- Marshall, J., Adcroft, A., Hill, C., Perelman, L., and Heisey, C.: A finite-volume, incompressible Navier Stokes model for studies of the ocean on parallel computers, *J. Geophys. Res.*, 102, 5753–5766, <https://doi.org/10.1029/96JC02775>, 1997.
- 15 Moore, A. M., Arango, H. G., Broquet, G., Powell, B. S., Weaver, A. T., and Zavala-Garay, J.: The Regional Ocean Modeling System (ROMS) 4-dimensional variational data assimilation systems: Part I—System overview and formulation, *Prog. Oceanogr.*, 91, 34–49, 2011.
- Nerger, L. and Hiller, W.: Software for ensemble-based data assimilation systems—Implementation strategies and scalability, *Computers & Geosciences*, 55, 110–118, 2013.
- Newman, T., Farrel, S., Richter-Menge, J., Elder, B., Connor, L., Kutz, N., and McAdoo, D.: Assessment of Radar-derived Snow Depth Measurements over Arctic sea ice., *J. Geophys. Res.*, 119, 8578–8602, <https://doi.org/https://doi.org/10.1002/2014JC010284>, 2014.
- 20 Perovich, D., Meier, W., Tschudi, M., Farrell, S., Hendricks, S., S. Gerland, C. H., Krumpen, T., Polashenski, C., Ricker, R., and Webster, M.: Sea Ice, <https://www.arctic.noaa.gov/Report-Card/Report-Card-2017/ArtMID/7798/ArticleID/699/Sea-Ice>, 2017.
- Perovich, D., Richter-Menge, J., and Polashenski, C.: Observing and understanding climate change: Monitoring the mass balance, motion, and thickness of Arctic sea ice, <http://imb-crrel-dartmouth.org>, 2018.
- 25 Pham, D. T.: Stochastic methods for sequential data assimilation in strongly nonlinear systems, *Monthly weather review*, 129, 1194–1207, 2001.
- Polashenski, C., Perovich, D., Richter-Menge, J., and Elder, B.: Seasonal ice mass-balance buoys: Adapting tools to the changing Arctic, *Annals of Glaciology*, 52, 18–26, 2011.
- Posey, P., Metzger, E., Wallcraft, A., Hebert, D., Allard, R., Smedstad, O., Phelps, M., Fetterer, F., Stewart, J., Meier, W., et al.: Assimilating high horizontal resolution sea ice concentration data into the US Navy’s ice forecast systems: Arctic Cap Nowcast/Forecast System (ACNFS) and the Global Ocean Forecast System (GOFS 3.1)., *Cryosphere Discussions*, 9, 2015.
- 30 Ricker, R., Hendricks, S., Kaleschke, L., Tian-Kunze, X., King, J., and Haas, C.: A weekly arctic sea-ice thickness data record from merged cryosat-2 and smos satellite data, *Cryosphere Discuss.*, pp. 1–27, 2017.
- Rostosky, P., Spreen, G., Farrell, S., Frost, S., Heygster, G., and Melsheimer, C.: Snow Depth Retrieval on Arctic Sea Ice from Passive Microwave Radiometers - Improvements and Extensions to Multiyear Ice Using Lower Frequencies, *JGR:Ocean*, <https://doi.org/https://doi.org/10.1029/2018JC014028>, 2018.
- Saha, S., Moorthi, S., Wu, X., Wang, J., Nadiga, S., Tripp, P., Behringer, D., Hou, Y.-T., Chuang, H.-y., Iredell, M., et al.: The NCEP climate forecast system version 2, *J. Climate*, 27, 2185–2208, 2014.

- Sakov, P.: EnKF-C user guide. arXiv:1410.1233., 2015.
- Sakov, P. and Bertino, L.: Relation between two common localisation methods for the EnKF, *Comput. Geosci.*, 15, 225–237, <https://doi.org/10.1007/s10596-010-9202-6>, 2011.
- Sakov, P. and Oke, P.: A deterministic formulation of the ensemble Kalman filter: an alternative to ensemble square root filters, *Tellus*, 60A, 361–371, <https://doi.org/10.1111/j.1600-0870.2007.00299.x>, 2008.
- Sakov, P., Counillon, F., Bertino, L., Lisæter, K., Oke, P., and Korablev, A.: TOPAZ4: an ocean-sea ice data assimilation system for the North Atlantic and Arctic, *Ocean Sci.*, 8, 633–656, <https://doi.org/10.5194/os-8-633-2012>, 2012.
- Shchepetkin, A. and McWilliams, J.: The regional oceanic modeling system (ROMS): a split-explicit, free-surface, topography-following-coordinate oceanic model, *Ocean Model.*, 9, 347 – 404, <https://doi.org/10.1016/j.ocemod.2004.08.002>, 2005.
- 10 Smith, L. C. and Stephenson, S. R.: New Trans-Arctic shipping routes navigable by midcentury, *Proceedings of the National Academy of Sciences*, 110, E1191–E1195, 2013.
- Stroeve, J., Holland, M. M., Meier, W., Scambos, T., and Serreze, M.: Arctic sea ice decline: Faster than forecast, *Geophys. Res. Lett.*, 34, <https://doi.org/10.1029/2007GL029703>, 2007.
- Tian-Kunze, X., Kaleschke, L., Maaß, N., Mäkynen, M., Serra, N., Drusch, M., and Krumpfen, T.: SMOS-derived thin sea ice thickness: algorithm baseline, product specifications and initial verification, *Cryosphere*, 8, 997–1018, <https://doi.org/10.5194/tc-8-997-2014>, 2014.
- 15 Tian-Kunze, X., Kaleschke, L., and Maaß: SMOS Daily sea ice thickness version 3, updated current year., CDC, icdc.cen.uni-hamburg.de, University of Hamburg, Germany, Digital media, 2016.
- Tonboe, R., Lavelle, J., Pfeiffer, R., and Howe, E.: Product User Manual for OSI SAF Global Sea Ice Concentration, http://osisa.f.met.no/docs/osisa_f_cdop3_ss2_pum_ice-conc_v1p6.pdf, 1.4, 2016.
- 20 Toudal Pedersen, L., Dybkjær, G., Eastwood, S., Heygster, G., Ivanova, N., Kern, S., Lavergne, T., Saldo, R., Sandven, S., Sørensen, A., and Tonboe, R.: ESA Sea Ice Climate Change Initiative (*Sea_Ice_cci*): Sea Ice Concentration Climate Data Record from the AMSR-E and AMSR-2 instruments at 25km grid spacing, version 2.0., Centre for Environmental Data Analysis, <https://doi.org/http://dx.doi.org/10.5285/c61bfe88-873b-44d8-9b0e-6a0ee884ad95>, 2017.
- Wang, K., Debernard, J., Sperrevik, A., Isachsen, P., and Lavergne, T.: A combined optimal interpolation and nudging scheme to assimilate OSISAF sea-ice concentration into ROMS, *Ann. Glaciol.*, 54, 8–12, <https://doi.org/10.3189/2013AoG62A138>, 2013.
- 25 Wang, M. and Overland, J. E.: A sea ice free summer Arctic within 30 years: An update from CMIP5 models, *Geophysical Research Letters*, 39, 2012.
- Warren, S. G., Rigor, I. G., Untersteiner, N., Radionov, V. F., Bryazgin, N. N., Aleksandrov, Y. I., and Colony, R.: Snow depth on Arctic sea ice, *J. Climate*, 12, 1814–1829, 1999.
- 30 Xie, J., Counillon, F., Bertino, L., Tian-Kunze, X., and Kaleschke, L.: Benefits of assimilating thin sea ice thickness from SMOS into the TOPAZ system, *Cryosphere*, 10, 2745–2761, <https://doi.org/10.5194/tc-10-2745-2016>, 2016.
- Xie, J., Counillon, F., and Bertino, L.: Impact of assimilating a merged sea-ice thickness from CryoSat-2 and SMOS in the Arctic reanalysis, *The Cryosphere*, 12, 3671–3691, 2018.
- Yang, Q., Losa, S., Losch, M., Tian-Kunze, X., Nerger, L., Liu, J., Kaleschke, L., and Zhang, Z.: Assimilating SMOS sea ice thickness into a coupled ice-ocean model using a local SEIK filter, *J. Geophys. Res.*, 119, 6680–6692, <https://doi.org/10.1002/2014JC009963>, 2014.
- 35 Zhang, X. and Walsh, J. E.: Toward a seasonally ice-covered Arctic Ocean: Scenarios from the IPCC AR4 model simulations, *J. Climate*, 19, 1730–1747, 2006.

Zygmuntowska, M., Rampal, P., Ivanova, N., and Smedsrud, L. H.: Uncertainties in Arctic sea ice thickness and volume: new estimates and implications for trends, *The Cryosphere*, 8, 705, 2014.