

# **Brief communication: Candidate sites of 1.5-Ma ice 37 km southwest of the Dome C summit, East Antarctica**

Olivier Passalacqua<sup>1</sup>, Marie Cavitte<sup>2,3</sup>, Olivier Gagliardini<sup>1</sup>,  
Fabien Gillet-Chaulet<sup>1</sup>, Frédéric Parrenin<sup>1</sup>, Catherine Ritz<sup>1</sup>, and Duncan Young<sup>2</sup>

<sup>1</sup>Univ. Grenoble Alpes, CNRS, IRD, IGE, F-38000 Grenoble, France

<sup>2</sup>University of Texas Institute for Geophysics, Jackson School of Geosciences, University of Texas  
at Austin, Austin, Texas, USA

<sup>3</sup>Department of Geological Sciences, Jackson School of Geosciences, University of Texas at  
Austin, Austin, Texas, USA

*Correspondence to:* Olivier Passalacqua (olivier.passalacqua@univ-grenoble-alpes.fr)

## **Abstract.**

The search for ice as old as 1.5 Ma requires the identification of places that maximize our chances to retrieve old, well-resolved, undisturbed and datable ice. One of these locations is very likely southwest of the Dome C summit, where elevated bedrock makes the ice thin enough to limit basal melting. A 3D ice flow simulation is used to calculate five selection criteria, which spatial variability is used to locate areas that have better glaciological properties than elsewhere. These selected areas (a few km<sup>2</sup>) lie on the flanks of the Dome C bedrock high, where a balance is found between risks of basal melting and sufficient age resolution. Within these areas, several sites of potential 1.5 Ma-old ice are proposed, situated on local bedrock summits or ridges. The trajectories of the ice particles towards these sites are short, and the ice flows over a smoothly-undulating bedrock. These sites will help to choose where new dense ground radar surveys should be conducted in upcoming field seasons.

## **1 Introduction**

Antarctic ice is an exceptional archive of the Earth's paleoclimates across all the glacial/interglacial periods, and the only one that contains direct samples of ancient atmospheres. The oldest available ice archive goes back 0.8 Ma in time (EPICA Dome C ice core, Jouzel et al., 2007), but is not old enough to study a main climatic transition that occurred between 1.2 Ma and 0.9 Ma, known from the temporal variations of the isotopic composition of benthic sediments (mid-Pleistocene transition, MPT, Lisiecki and Raymo, 2005). The main climatic periodicity and the amplitude of climate cycles seem to radically change during the MPT. There is presently no general agreement on the processes responsible for this transition and its origin, and the influence of the trend of atmospheric CO<sub>2</sub>, the

dynamics of ice sheets in the northern hemisphere, the sea ice extent, or the dust content of the atmosphere have been proposed. Considering these scientific issues, locating a future 1.5 Ma-old ice drill site was identified as one of the main goal of the ice-core community (see an overview in Jouzel  
25 and Masson-Delmotte, 2010).

Thermal and mechanical ice flow simulations have to be carried out to assess the potential of possible drill sites (Fischer et al., 2013). The present study is part of a set of three modelling exercises led by the IGE (Institut des Géosciences de l'Environnement, Grenoble) group and collaborators around Dome C, and more specifically on the bedrock high lying  $\sim 40$  km southwest of the Dome C  
30 summit, suspected to be a good old-ice candidate (Van Liefferinge and Pattyn, 2013), under the ice ridge linking Dome C to the Vostok region (for a precise description, see Young et al., 2017). First, a 1D thermal model was run over the last 0.8 Ma to determine the present state (temperate or cold) of basal ice, which was compared to the reflectivity map in the region of Dome C (Passalacqua et al.,  
35 2017). The value of the local geothermal flux was inferred, and the spatial distribution of interpreted subglacial water was explained. The results of these transient simulations show that the top of this bedrock high is likely to be melt-free on long timescales, or to undergo very limited basal melting. Hence, this subregion is thermally compatible with the archiving process on glacial/interglacial timescales. Second, a kinematic 1D ice flow model was used to evaluate the age and age resolution of the deepest portion of the ice sheet (Parrenin et al., 2017). The distance between the dated isochrones  
40 and the modelled ones was minimized to infer a thinning parameter that characterizes the vertical deformation through the ice column. We showed that the observed isochrones are compatible with high basal age and provide sufficient resolution. However, these two previous studies neglect horizontal motion of ice, whereas the trajectories of the ice particles are not only vertical. The travel time of these particles, and consequently their age, is strongly influenced by the shape of the bedrock and  
45 the ice surface. These prior studies lacked the description of the local 3D state of stress of the ice, where the geometry of the terrain and the strain history of the ice particles can be properly taken into account.

To evaluate the quality and position of deep old ice, we here proceed to a steady state 3D ice flow simulation, at a regional scale. Whereas these two previous 1D modelling results alone could  
50 not help us determine the better oldest-ice targets, we here intend to provide objective criteria that together delimit kilometer-scale areas of old, well-resolved, undisturbed basal ice. Note that the goal of this study is not to assess the best age estimation, but to evaluate which sites have better glaciological characteristics than others. The bottom-most ice recovered should be older than the MPT, ideally as old as 1.5 Ma such that several climate cycles pre-MPT can be recorded. The vertical  
55 age resolution has to be better than  $10 \text{ ka m}^{-1}$  to detect high-frequency climatic variability in the ice core (J. Chappellaz, personal communication). Moreover, basal ice will probably be disturbed, similarly to the deepest 60 m of the EPICA Dome C ice core (Jouzel et al., 2007; Tison et al., 2015). We have no better evaluation of the height of disturbed ice in the region, and we will similarly

take 60 m as a safety distance for the drilling minimum distance from the bedrock, but we should  
60 keep in mind this cutoff height could be an underestimation. One should look for places where the  
mechanisms responsible for stratigraphy disturbances (cumulated basal shear, bedrock roughness)  
should be minimal. Convergent flow should be avoided as well, because it tends to thicken basal  
layers. This is unfavourable to recovering oldest ice as it will shift older layers downwards, and  
makes dating process more complex (Tison et al., 2015). Finally, the location of the future drill  
65 site should be above the highest subglacial lake detected by the radar survey (Young et al., 2017),  
otherwise the risk of basal melting could be drastically increased. We will call this threshold "water  
limit", above which there is no evidence of the presence of water in the radargrams.


Another modelling study has been recently led at a continental scale to locate a future drill site  
(Van Liefferinge et al., 2018). They used a transient thermodynamical model to compute the highest  
70 geothermal flux value that keep basal ice frozen. By comparison with available continental geother-  
mal flux datasets, they locate areas where basal ice has likely remained frozen over 1.5 Ma. The  
authors also included a further mechanical constraint representing limited impact of bedrock rough-  
ness on the preservation of the bottom ice stratigraphy. Our study and the one of Van Liefferinge  
et al. (2018) differ in the way they account for the heat budget of ice and for the mechanical con-  
75 straints on the basal ice. We should consider these differences as an interesting source of information  
for the decision-making process, and they will be discussed in § 3.5.

The decision-making process of a drill site needs both field survey and modelling, the former  
feeding the latter with geophysical constraints, and the latter reducing the areas of interest for new  
field surveys, focusing more and more on promising sites. The information brought by the present  
80 study should be sufficient for a last dense radar survey to be led on promising sites during the next  
field seasons, at a scale of a few hundred of meters. Then the community should be able to take a  
decision for a drill site in the Dome C region.

## 2 Ice flow model

### 2.1 Model description

85 The Stokes equations are solved on a  $83 \times 114$  km domain, approximately centered on the Dome C  
(Fig. 1), using the finite element model Elmer/Ice. The surface and bed geometries are provided by  
the Bedmap2 data set (Fretwell et al., 2013), except on the bedrock high southwest of Dome C,  
where a dense airborne radar survey has been recently collected (Young et al., 2017). The model  
works in ice-equivalent, and we adjust the surface height by assuming that the density profile of the  
90 firn is the one of Dome C on the whole domain (Schwander et al., 2001). Horizontal resolution of the  
corresponding mesh is 1 km. The resolution of the 20 vertical elements of the mesh evolves linearly,  
so that the deepest one being 25 times finer than the upper one, and to ensure that the velocity field  
is better resolved near the bottom.



**Figure 1.** Mesh, bedrock dataset (Fretwell et al., 2013; Young et al., 2017) and basal melt rate (Passalacqua et al., 2017) used for the simulation on a  $83 \times 114$  km domain. The red patch on the context map (bottom left) shows the domain used for the calculation, and the blue rectangle at the top of the image is the location of Fig 2.

Our domain sits in the center of the Antarctic plateau, and lateral boundary conditions do not  
 95 correspond to physical boundaries, but to virtual vertical surfaces. As the domain contains the dome  
 summit, the ice flow is divergent and no input flow should be considered. We impose velocity bound-  
 ary conditions corresponding to the shallow-ice approximation (SIA) (Hutter, 1983). The ice surface  
 as observed today may not correspond to the chosen rheology and is relaxed for 50 years such that  
 the present surface slope does not induce an unrealistic velocity field. The ice surface is very flat,  
 100 and 50 years is enough to accommodate the surface altitude to the ice rheology up to  $\sim 1$  m, with-  
 out radically changing the orientation of the ice ridge, on which we have little control. The part of  
 ice motion attributed to basal sliding is not known precisely in the Dome C region, and depends on  
 water circulation. We here focus on a region where basal melting is probably not present or limited,  
 and horizontal velocities are very small, so that, for the sake of simplicity, a no-sliding condition is  
 105 imposed at the bottom of the ice column. Vertical velocities at the base are equal to the basal melt  
 rate output from previous modelling work (Fig. 7 in Passalacqua et al., 2017).

As we know that the basal ice in the Dome C region is at or near the melting point (Passalacqua  
 et al., 2017; Van Liefferinge and Pattyn, 2013), the temperature profile measured in the EPICA  
 Dome C borehole is a good representation of the thermal structure of the ice in the Dome C vicinity.  
 110 Hence, we account for ice temperature by using the same normalized temperature profile on all  
 the domain. Solving the coupled thermo-mechanical equations would require excessive computing  
 resources, without radically changing the ice fluidity – which is mainly controlled by temperature.  
 Similarly, we do not account here for long-term evolutions of the ice sheet surface, but are aware  
 that this assumption strongly affects the trajectories of the ice particles.

115 As the main interest of this work focuses on what happens for deepest ice, mechanical anisotropy  
of the ice has to be accounted for. The relation between strain rates and stresses are described by the  
generalized orthotropic linear flow law (GOLF, Gillet-Chaulet et al., 2005), given a certain vertical  
profile of ice fabric. By introducing a dependence on the second invariant of the deviatoric stress, this  
law can be extended to the non-linear case (Ma et al., 2010). The fabric profile is only known at the  
120 dome summit (Durand et al., 2009), and shows that the ice mainly undergoes vertical compression,  
but also longitudinal extension in the deep layers (Tison et al., 2015). However, there is no reason  
to use the very same fabric profile everywhere else, where shearing has more influence or the bed  
shape is different. In this short study we will not discuss the influence of the chosen rheology, but we  
first made sure that the computed surface velocities correctly simulated the horizontal velocity field  
125 measured at the surface by Vittuari et al. (2004) for different  $n$  values and fabric profiles. Hence, we  
decided to use the widely-accepted value of 3 for  $n$ , and a synthetic vertical fabric profile, for which  
the eigenvalues of the second order orientation tensor evolve linearly with normalized depth, from  
isotropy at the surface to a single maximum fabric at the bottom.

## 2.2 Model outputs

130 Back trajectories are computed from the 3D velocity field using a Lagrangian scheme such that the  
age is known along the forward trajectory of the ice particles, and an age field is generated. The age  
resolution could be calculated from the vertical derivative of this age field, but we found it more  
accurate to track the annual layer thickness  $\lambda$  from the ice surface (Whillans, 1979) using

$$\frac{d\lambda}{dt} = \lambda \dot{\epsilon}_{zz} \quad (1)$$

135 where  $\dot{\epsilon}_{zz}$  is the vertical strain rate. This formulation neglects vertical rotation effects that tends to  
overturn internal layers. This assumption is reasonable if internal layers are mainly horizontal, which  
is the case over the studied bedrock high. The age resolution is then stated as  $1/\lambda$ , considered as the  
layer thickness for a single year.

The way ice strains by flowing over a rough bed differs depending on the shape of the bedrock  
140 underneath. Similarly, a given bedrock shape can be a convergence or a divergence area, depending  
on the orientation of horizontal ice flow. Once the velocity field is known, a local coordinate system  
 $(X, Y, Z)$  can be defined at each point, for which the  $X$ -axis is oriented along flow. The curvature of  
the bed perpendicular to the flow is then computed everywhere, and convergence areas are identified  
where the bed curvature is positive.

145 Beyond the computation of ages and age resolutions, the 3D simulation is also useful to detect  
where deep ice is more likely to be folded. Shearing will tend to amplify small wrinkles in the ice  
layers and so disturb the ice's basal stratigraphy, whereas longitudinal extension will tend to flatten

them. Competition between shear and longitudinal stresses can be represented by a dimensionless shear number (Waddington et al., 2001)

$$150 \quad S = \frac{2\dot{\epsilon}_{XZ}}{\dot{\epsilon}_{XX} - \dot{\epsilon}_{ZZ}} \quad (2)$$

where  $\dot{\epsilon}_{XZ}$  is the shear strain rates along ice flow,  $\dot{\epsilon}_{XX}$  and  $\dot{\epsilon}_{ZZ}$ , the local longitudinal and vertical strain rates. Waddington et al. (2001) used this shear number as a criterium to detect if a given wrinkle can be amplified by shear. More simply here, we use it to predict the presence of undisturbed ice, whereby the smallest shear number is best.

### 155 **2.3 Selection of favourable locations**

If the absolute value of the age, age resolution, or strain rates can be discussed regarding the choices of the model parameters, we reckon their spatial variabilities are robust since they mainly depend on the shape of the bedrock and of the ice surface. As a consequence, this study focuses on comparing in-between promising locations rather than on discussing the influence of model parameters.

160 The five selection criteria (age, age resolution, bed curvature, shear number and bed height) are used to compute five masks, thresholded as follows. For bed curvature,  $0 \text{ m}^{-1}$  would have been the natural threshold but it was too much restrictive and we decided for a slightly higher value ( $2 \cdot 10^{-5} \text{ m}^{-1}$ ). The shear number threshold appeared naturally by studying its spatial evolution (see §3.3). The bed height threshold correspond to the "water limit", at 480 m. Furthermore, our results  
165 show that most of the subglacial elevated bed high southwest of Dome C is favourable to the existence of 1.5 Ma-old ice, so we adopted more conservative age and age resolution thresholds for the selection of smaller suitable locations (1.8 Ma for the age and  $8.5 \text{ ka m}^{-1}$  for the age resolution, 60 m above the bed). A boolean combination of the five masks delineate the areas fulfilling all our five selection criteria.

## 170 **3 Results and interpretation**

### **3.1 Age at 60 m above the bed**

The area identified as possibly hosting oldest ice is elongated along the bedrock high, and stands at an intermediate bed elevation (mainly between 400 m and 550 m above seal level, Fig. 2a). Neither the very top of the bedrock high, nor its lowest foothills appear to be suitable for the archiving  
175 process of very old ice. The imposed basal melt rates are zero on the upper part of the bedrock high, therefore infinite age are calculated for the very basal ice. In that case, the age of the ice standing 60 m above the bed is strongly dependent on the ice thickness. On the top of the bedrock high, the ice is at its thinnest and the old ice is at less than 60 m above the bed. On the foothills of the bedrock

high, the ice is thicker, basal melt rates are above zero, and the basal ice is therefore continuously  
180 being melted from the bottom.

Some places may host ice even older than 2 Ma, but they all stand below Young et al's (2017)  
water limit (Fig. 2, yellow line). The presence of very old ice in those areas is not impossible, but  
may also be the consequence of insufficient imposed basal melting. The transition between melting  
and frozen ice should stand somewhere on the flanks of the bedrock high, but it is difficult to pinpoint  
185 the precise location of this threshold. Despite the promising thick ice in this region that could allow  
old ages and sufficient age resolution, the risk of basal melting is real.

### 3.2 Age resolution at 1.5 Ma


Age resolution in the deepest part of the ice column is influenced by two factors. First and obviously,  
the thicker the ice, the better the age resolution. As a consequence, the tops of the bedrock high are  
190 rarely compatible with a sufficient age resolution of oldest ice. Bedrock flanks should be preferred,  
but some of the thickest ice areas on the flanks will be discarded as well because of an increased  
risk of basal melting. Second, for ice positioned close to the ice divide, age resolution may benefit  
from a thickening effect of the deeper layers (so-called Raymond effect, Raymond, 1983). This  
results in a band of well-resolved ice, oriented along the ice ridge and perpendicular to the bedrock  
195 high. However, no Raymond arch is visible in the radargrams that would indicate a strong Raymond  
effect here. One explanation is that the shape of the ice surface at these sites is much more rounded  
than at the Dome C summit, and the produced along-ridge flow tend to dampen the amplitude of  
the Raymond arches (Martín et al., 2009). Moreover, the characteristic time for a Raymond arch to  
form here would be several 100 ka (Martín et al., 2009), during which the surface ridge may have  
200 moved, smoothing out the developing arches. Unfortunately, the past position and shape of the ridge  
are unknown, and drilling far from its present position would not guarantee a better resolution.

### 3.3 Stratigraphic disturbances

At a divide, the shear stress perpendicular to the divide is zero, so that in general the shear number  $S$   
of the deep ice close to the ridge is low ( $\sim 10$ ). However, a serie of subglacial peaks lie immediately  
205 west of the ice divide. As a consequence the shear number increases very sharply and reaches much  
higher values ( $\sim 100$ ), and this zone of higher shear should be discarded for a future drill site (Fig. 2d,  
wavy area). All the areas for which the basal ice crossed this zone of high shear should be discarded  
as well, so the trajectories of the ice particles need to be evaluated.


### 3.4 Trajectories for oldest-ice spots

210 The best combination of age, age resolution, folding, convergence and melting criteria is shown in  
Fig. 2 e, revealing several spots of ice meeting the five criteria. The areas for which the trajectories  
are the shorter should be preferred, highlighted here by three magenta boxes. For boxes A, B and C



**Figure 2.** Selection criteria thresholded as follows (yellow areas): a) age  $> 1.8$  Ma, b) age resolution  $< 8.5 \text{ ka m}^{-1}$ , c) bed curvature  $< 2 \cdot 10^{-5} \text{ m}^{-1}$ , d) shear number  $S < 40$ , e) boolean combination of the selection criteria. The water limit correspond to a bed height of 480 m and is shown here by a thick yellow contour. Magenta boxes A, B and C correspond to areas that could be considered as our best oldest-ice targets. Dashed lines show trajectories of ice particles, the red one correspond to the profile presented in Fig. 3. Colored points locate possible drill sites, discussed in the text. Hatched areas show the best areas of Van Liefferinge et al. (2018). All the panels cover the same area. This figure focuses on the bedrock high designed as such in Fig.1, located  $\sim 40$  km southwest of the Dome C summit. Projection: WGS84/Antarctic Polar Stereographic – EPSG:3031 (kilometers).





**Figure 3.** Trajectories of the ice particles from the ice surface towards the black point in Fig. 2 (red trajectory). Red numbers indicate the age of the ice, in Ma. The bed profile is shown in grey. The thin red line corresponds to one possible drill site (red point in Fig. 2).

the ice originates from the divide, guided by the strong lateral divergence resulting from the shape of the ice surface. Locations within boxes C and A are closer to the ice ridge and should be considered first for a future Oldest Ice drilling because of shorter trajectories, and less risk of stratigraphy 215 disturbances. Box B in particular lies on a relatively flat bedrock platform (black point in Fig. 2e), that may ensure stability of ice flow. Figure 3 shows the paths taken by ice from the surface to this site, demonstrating that the horizontal distance travelled does not exceed 10 km from the surface. However, as there is probably no basal melting here, deep ice would closely follow the bed for 220 several kilometers, in a depth range dominated by strong vertical shear, enhanced by an undulating bed underneath. To minimize the bed influence, we could also consider a drill site located 3 km upstream, where the ice age would still exceed 1.5 Ma (red point in Fig. 2e, red dashed line in Fig. 3).

Of course, locating a unique "best" drill site within one of these three boxes is not possible with this 3D-modelling approach only. However, it allows a restricted area to be defined where a new set 225 of observations will be the most valuable. We should focus on local bedrock summits or crest lines, because local troughs make the ice flow converge, but also heat flow (Van der Veen et al., 2007), increasing the possibility of insufficient age and positive basal melt. Considering that, only a few set of favourable drill sites remain in boxes A, B and C (blue, orange, pink and yellow points in Fig 2e). Pink and blue points have less risks of basal melting, while yellow and orange have less risks of 230 stratigraphic disturbances. The best choice between these sites should be now guided by local radar surveys characterizing the internal layering of the ice, and the vertical strain rate profile (Nicholls et al., 2015).

### 3.5 Comparison with results based on thermodynamical modelling

Van Liefferinge et al. (2018) identified a 8 km-long area covering the SE upper part of the bedrock  
235 high, that crosses our box A, but both do not overlap perfectly (Fig. 2e, hatched blue and yellow  
areas). They also identified a site within our magenta box B, but no site in the box C. These compar-  
ative results highlights the complementarity of the two approaches. The 1D thermodynamical model  
of Van Liefferinge et al. (2018) has a better control on the thermal aspect of the problem than on  
its mechanical aspect, and selects sites that are more conservative from a heat budget point of view,  
240 i.e. preferentially local bedrock heights. In contrast, our 3D approach accounts for the horizontal  
strain of the ice, and select sites that are more conservative from a mechanical point of view. The  
upper part of the box A or the left part of the box B validate the constraints of both approaches. In  
our approach, the bedrock summit in box C is the safest mechanically; however, it was not selected  
by Van Liefferinge et al. (2018) because of a local bedrock roughness exceeding their threshold of  
245 20 m, despite the fact that their thermal criterion was fulfilled.

## 4 Conclusions

The three-dimensional ice flow simulation presented here aims at defining and calculating several  
objective criteria, which represent ideal conditions for the retrieval of old, well-resolved, undisturbed  
ice. The influence of the bedrock high and of the position of the ice ridge allows us to define only  
250 few sites compatible with all our selection criteria. The ages calculated at the base by our model  
simulations predict ice older than 1.5 Ma old high enough above the bedrock, which gives us con-  
fidence that the community's target of 1.5 Ma years should be attainable, with the required age  
resolution. However, the modelling approach implicitly assumes that the ice flow is regular down  
to the bedrock, but there is no guarantee it is actually the case. A ground radar survey focusing on a  
255 few hundred-meter scale areas presented here is essential to explore the structure of the deep layers.  
Finally, a rapid-access drill (Grilli et al., 2014) is currently planned to be deployed in the 2018/19  
season to assess ice quality and age for a chosen target site, before the final site is decided upon.

*Acknowledgements.* We would like to thank Jean-Louis Tison for the editing process and we appreciate the  
helpful comments of Michelle Koutnik, Tas Van Ommen and Bryn Hubbard, as referees. The Australian Antarc-  
260 tic Division provided funding and logistical support (AAS 3103, 4077, 4346). This work was supported by the  
Australian Government Cooperative Research Centres Programme through the Antarctic Climate and Ecosys-  
tems Cooperative Research Centre (ACE CRC); support for UTIG came from the G. Unger Vetlesen Foundation  
and NSF grant PLR-1443690. This paper is UTIG contribution number 3269. This publication was generated in  
the frame of Beyond EPICA-Oldest Ice (BE-OI). The project has received funding from the European Union's  
265 Horizon 2020 research and innovation programme under grant agreement No. 730258 (BE-OI CSA). It has re-  
ceived funding from the Swiss State Secretariate for Education, Research and Innovation (SERI) under contract

number 16.0144. It is furthermore supported by national partners and funding agencies in Belgium, Denmark, France, Germany, Italy, Norway, Sweden, Switzerland, The Netherlands and the United Kingdom. Logistic support is mainly provided by AWI, BAS, ENEA and IPEV. The opinions expressed and arguments employed  
270 herein do not necessarily reflect the official views of the European Union funding agency, the Swiss Government or other national funding bodies. This is BE-OI publication number ... This publication also benefitted from support by the Université Grenoble Alpes in the framework of the proposal called Grenoble Innovation Recherche AGIR. This work was granted access to the HPC resources of CINES under the allocation A0020106066 made by GENCI. We thank Brice Van Liefferinge for making its datasets available.

## 275 References

- Durand, G., Svensson, A., Persson, A., Gagliardini, O., Gillet-Chaulet, F., Sjolte, J., Montagnat, M., and Dahl-Jensen, D.: Evolution of the texture along the EPICA Dome C ice core, *Low Temperature Science*, 68, 91–105, <http://133.87.26.249/dspace/handle/2115/45436>, 2009.
- 280 Fischer, H., Severinghaus, J., Brook, E., Wolff, E., Albert, M., Alemany, O., Arthern, R., Bentley, C., Blankenship, D., Chappellaz, J., Creyts, T., Dahl-Jensen, D., Dinn, M., Frezzotti, M., Fujita, S., Galée, H., Hindmarsh, R., Hudspeth, D., Jugie, G., Kawamura, K., Lipenkov, V., Miller, H., Mulvaney, R., Parrenin, F., Patryn, F., Ritz, C., Schwander, J., Steinhage, D., van Ommen, T., and Wilhelms, F.: Where to find 1.5 million yr old ice for the IPICS "Oldest-Ice" ice core, *Climate of the Past*, 9, 2489–2505, doi:10.5194/cp-9-2489-2013, <http://www.clim-past.net/9/2489/2013/>, 2013.
- 285 Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., Bianchi, C., Bingham, R. G., Blankenship, D. D., Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita, S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W., Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langley, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., 290 Matsuoka, K., Mouginot, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippin, D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C., and Zirizzotti, A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, *The Cryosphere*, 7, 375–393, doi:10.5194/tc-7-375-2013, <http://www.the-cryosphere.net/7/375/2013/>, 2013.
- 295 Gillet-Chaulet, F., Gagliardini, O., Meyssonnier, J., Montagnat, M., and Castelnau, O.: A user-friendly anisotropic flow law for ice-sheet modelling, *Journal of Glaciology*, 51, 3–14, 2005.
- Grilli, R., Marrocco, N., Desbois, T., Guillerm, C., Triest, J., Kerstel, E., and Romanini, D.: Invited Article: SUBGLACIOR: An optical analyzer embedded in an Antarctic ice probe for exploring the past climate, *Review of Scientific Instruments*, 85, 111 301, 2014.
- 300 Hutter, K.: *Theoretical glaciology: material science of ice and the mechanics of glaciers and ice sheets*, vol. 1, Springer, 1983.
- Jouzel, J. and Masson-Delmotte, V.: Deep ice cores: the need for going back in time, *Quaternary Science Reviews*, 29, 3683–3689, doi:10.1016/j.quascirev.2010.10.002, <http://linkinghub.elsevier.com/retrieve/pii/S0277379110003562>, 2010.
- 305 Jouzel, J., Masson-Delmotte, V., Cattani, O., Dreyfus, G., Falourd, S., Hoffmann, G., Minster, B., Nouet, J., Barnola, J. M., Chappellaz, J., Fischer, H., Gallet, J. C., Johnsen, S., Leuenberger, M., Loulergue, L., Luethi, D., Oerter, H., Parrenin, F., Raisbeck, G., Raynaud, D., Schilt, A., Schwander, J., Selmo, E., Souchez, R., Spahni, R., Stauffer, B., Steffensen, J. P., Stenni, B., Stocker, T. F., Tison, J. L., Werner, M., and Wolff, E. W.: Orbital and Millennial Antarctic Climate Variability over the Past 800,000 Years, *Science*, 317, 793– 310 796, doi:10.1126/science.1141038, <http://www.sciencemag.org/cgi/doi/10.1126/science.1141038>, 2007.
- Lisiecki, L. E. and Raymo, M. E.: A Pliocene-Pleistocene stack of 57 globally distributed benthic  $\delta^{18}\text{O}$  records, *Paleoceanography*, 20, 2005.

- Ma, Y., Gagliardini, O., Ritz, C., Gillet-Chaulet, F., Durand, G., and Montagnat, M.: Enhancement factors for grounded ice and ice shelves inferred from an anisotropic ice-flow model, *Journal of Glaciology*, 56, 805–812, <http://www.ingentaconnect.com/content/igsoc/jog/2010/00000056/00000199/art00006>, 2010.
- 315
- Martín, C., Hindmarsh, R. C., and Navarro, F. J.: On the effects of divide migration, along-ridge flow, and basal sliding on isochrones near an ice divide, *Journal of Geophysical Research: Earth Surface*, 114, 2009.
- Martín, C., Gudmundsson, G. H., Pritchard, H. D., and Gagliardini, O.: On the effects of anisotropic rheology on ice flow, internal structure, and the age-depth relationship at ice divides, *Journal of Geophysical Research: Earth Surface*, 114, doi:10.1029/2008JF001204, <http://dx.doi.org/10.1029/2008JF001204>, F04001, 2009.
- 320
- Nicholls, K. W., Corr, H. F., Stewart, C. L., Lok, L. B., Brennan, P. V., and Vaughan, D. G.: A ground-based radar for measuring vertical strain rates and time-varying basal melt rates in ice sheets and shelves, *Journal of Glaciology*, 61, 1079–1087, 2015.
- Parrenin, F., Cavitte, M. G. P., Blankenship, D. D., Chappellaz, J., Fischer, H., Gagliardini, O., Masson-Delmotte, V., Passalacqua, O., Ritz, C., Roberts, J., Siegert, M. J., and Young, D. A.: Is there 1.5 million-year old ice near Dome C, Antarctica?, *The Cryosphere Discussions*, 2017, 1–16, doi:10.5194/tc-2017-69, <http://www.the-cryosphere-discuss.net/tc-2017-69/>, 2017.
- 325
- Passalacqua, O., Ritz, C., Parrenin, F., Urbini, S., and Frezzotti, M.: Geothermal flux and basal melt rate in the Dome C region inferred from radar reflectivity and heat modelling, *The Cryosphere*, 11, 2231–2246, doi:10.5194/tc-11-2231-2017, <https://www.the-cryosphere.net/11/2231/2017/>, 2017.
- 330
- Raymond, C. F.: Deformation in the vicinity of ice divides, *Journal of Glaciology*, 29, 357–373, 1983.
- Schwander, J., Jouzel, J., Hammer, C. U., Petit, J.-R., Udasti, R., and Wolff, E.: A tentative chronology for the EPICA Dome Concordia ice core, *Geophysical research letters*, 28, 4243–4246, 2001.
- 335
- Tison, J.-L., de Angelis, M., Littot, G., Wolff, E., Fischer, H., Hansson, M., Bigler, M., Udasti, R., Wegner, A., Jouzel, J., et al.: Retrieving the paleoclimatic signal from the deeper part of the EPICA Dome C ice core, *The Cryosphere*, 9, 1633–1648, 2015.
- Van der Veen, C. J., Leftwich, T., von Frese, R., Csatho, B., and Li, J.: Subglacial topography and geothermal heat flux: Potential interactions with drainage of the Greenland ice sheet, *Geophysical Research Letters*, 34, 2007.
- 340
- Van Liefferinge, B. and Pattyn, F.: Using ice-flow models to evaluate potential sites of million year-old ice in Antarctica, *Climate of the Past Discussions*, 9, 2859–2887, doi:10.5194/cpd-9-2859-2013, <http://www.clim-past-discuss.net/9/2859/2013/>, 2013.
- Van Liefferinge, B., Pattyn, F., Cavitte, M. G. P., Karlsson, N. B., Young, D. A., Sutter, J., and Eisen, O.: Promising Oldest Ice sites in East Antarctica based on thermodynamical modelling, *The Cryosphere Discussions*, 2018, 1–22, doi:10.5194/tc-2017-276, <https://www.the-cryosphere-discuss.net/tc-2017-276/>, 2018.
- 345
- Vittuari, L., Vincent, C., Frezzotti, M., Mancini, F., Gandolfi, S., Bitelli, G., and Capra, A.: Space geodesy as a tool for measuring ice surface velocity in the Dome C region and along the ITASE traverse, *Annals of Glaciology*, 39, 402–408, 2004.
- 350
- Waddington, E. D., Bolzan, J. F., and Alley, R. B.: Potential for stratigraphic folding near ice-sheet centers, *Journal of Glaciology*, 47, 639–648, 2001.

Whillans, I. M.: Ice flow along the Byrd Station strain network, Antarctica, *Journal of Glaciology*, 24, 15–28, 1979.

355 Young, D. A., Roberts, J. L., Ritz, C., Frezzotti, M., Quartini, E., Cavitte, M. G. P., Tozer, C. R., Steinhage, D., Urbini, S., Corr, H. F. J., van Ommen, T., and Blankenship, D. D.: High-resolution boundary conditions of an old ice target near Dome C, Antarctica, *The Cryosphere*, 11, 1897–1911, doi:10.5194/tc-11-1897-2017, <https://www.the-cryosphere.net/11/1897/2017/>, 2017.