

# Quantifying the snowmelt-albedo feedback at Neumayer Station, East Antarctica

Constantijn L. Jakobs<sup>1</sup>, Carleen H. Reijmer<sup>1</sup>, Peter Kuipers Munneke<sup>1</sup>, Gert König-Langlo<sup>2</sup>, and Michiel R. van den Broeke<sup>1</sup>

<sup>1</sup>Institute for Marine and Atmospheric Research Utrecht, Utrecht University, Utrecht, The Netherlands

<sup>2</sup>Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven, Germany

**Correspondence:** Stan Jakobs (c.l.jakobs@uu.nl)

**Abstract.** We use 24 years (1992–2016) of high-quality meteorological observations at Neumayer Station, East Antarctica, to force a surface energy balance model. The modelled 24-year cumulative surface melt at Neumayer amounts to 1154 mm water equivalent (w.e.), with only a small uncertainty ( $\pm 3$  mm w.e.) from random measurement errors. Results are more sensitive to the chosen value for the surface momentum roughness length and new snow density, yielding a range of 900–1220 mm w.e. Melt at Neumayer occurs only in the months November to February, with a summer average of 50 mm w.e. and large interannual variability ( $\sigma = 42$  mm w.e.). This is a small value compared to an annual average (1992–2016) accumulation of  $415 \pm 86$  mm w.e. Absorbed shortwave radiation is the dominant driver of temporal melt variability at Neumayer. To assess the importance of the snowmelt-albedo feedback we include and calibrate an albedo parameterisation in the surface energy balance model. We show that, without the snowmelt-albedo feedback, surface melt at Neumayer would be approximately three times weaker, demonstrating how important it is to correctly represent this feedback in model simulations of surface melt in Antarctica.

## 1 Introduction

The Antarctic ice sheet (AIS) contains more than 25 million km<sup>3</sup> of ice, sufficient to raise global mean sea level by almost 60 m if melted completely (Fretwell et al., 2013). Between 1992 and 2017, the AIS lost mass at an accelerated rate, contributing  $7.6 \pm 3.9$  mm to global sea level (Shepherd et al., 2018). This mass loss is mainly observed in coastal West Antarctica and the Antarctic Peninsula (AP), and is caused by glaciers that accelerated after their buttressing ice shelves had thinned or disintegrated (Wouters et al., 2015; Turner et al., 2017). The interaction between meltwater and firn, the intermediate product between snow and glacier ice, is hypothesised to play an important role in ice shelf disintegration (Kuipers Munneke et al., 2014). If the firn layer contains enough air, as is the case for most of the AIS, meltwater can percolate downwards and re-freeze (Ligtenberg et al., 2014). If the storage capacity of the firn layer is reduced, surface meltwater can flow laterally towards the ice shelf edge (Bell et al., 2017), be stored englacially (Lenaerts et al., 2017) or form ponds on the ice shelf surface (Kingslake et al., 2017). In all cases, meltwater can accumulate in crevasses, thereby increasing the hydrostatic pressure in the crevasse tip, causing it to penetrate farther down. When a crevasse reaches the bottom of the ice shelf or a basal crevasse, part

of the ice shelf disintegrates, a process called hydrofracturing (Van der Veen, 2007). Hydrofracturing has been identified as a potential precursor to rapid loss of Antarctic ice, accelerating sea level rise (DeConto and Pollard, 2016). In combination with enhanced ocean swell under low sea-ice conditions (Massom et al., 2018), hydrofracturing likely caused the disintegration of Larsen B ice shelf in the AP in 2002 (Rignot et al., 2004; Scambos et al., 2004). In July 2017, a large iceberg calved from Larsen C ice shelf, but it is unclear whether this signifies a further southward progression of ice shelf destabilisation in the AP (Hogg and Gudmundsson, 2017).

Improving our predictive capabilities of future ice shelf stability, AIS mass loss and associated sea level rise, thus requires a thorough understanding of the surface melt process on Antarctic ice shelves. In contrast to meltwater occurrence, which is readily observed from space (Picard et al., 2007; Tedesco, 2009; Luckman et al., 2014), observational estimates of surface melt rates on Antarctic ice shelves are rare; they have been obtained locally through explicit modelling of the surface energy balance (SEB) (Van den Broeke et al., 2010; Kuipers Munneke et al., 2012, 2018). In turn, these enabled continent-wide melt rate estimates using calibrated satellite products based on backscatter strength of radio waves (Trusel et al., 2013, 2015). These studies invariably demonstrate that in most parts of Antarctica, melt currently is a weak and intermittent process. In this melt regime, the positive snowmelt-albedo feedback plays a decisive role: when snow melts, meltwater may refreeze in the cold snowpack, resulting in considerably larger grains ( $\sim 1$  mm) than new snow or snow that has been subjected to dry compaction ( $\sim 0.1$  mm). Larger snow grains reduce backward scattering of photons into the snowpack, increasing the probability of absorption, reducing the surface albedo especially in the near-infrared (Wiscombe and Warren, 1980; Gardner and Sharp, 2010). This further enhances absorption of solar radiation and melt. For pure, uncontaminated snow, the strength of the snowmelt-albedo feedback depends on multiple factors, e.g. the intensity and duration of the melt and the frequency and intensity of snowfall events, which provide new snow consisting of smaller grains. We therefore expect the snowmelt-albedo feedback to be spatially and temporally variable on Antarctic ice shelves.

Most studies on the snowmelt-albedo feedback address the removal of (seasonal) snow and the appearance of dark soil or water (Perovich et al., 2002; Hall, 2004; Flanner et al., 2007; Qu and Hall, 2007), leading to further warming of the air/water. These studies commonly express the melt-albedo feedback in terms of air/water temperature sensitivity. Our aim is to quantify the impact on the melt rate of the darkening but not the disappearance of snow, a process addressed by far fewer studies (Box et al., 2012; Van As et al., 2013). To that end, we implement a snow albedo parameterization (Gardner and Sharp, 2010; Kuipers Munneke et al., 2011b) in an SEB model, which is then calibrated using observations and used to study the sensitivity of melt rates to snow properties that influence snow albedo. We use 24 years of high-quality in situ observations (König-Langlo, 2017) from the German research station Neumayer (Fig. 1) to calculate the SEB and melt rate. We investigate the effects of measurement uncertainties and model settings on the calculated cumulative amount of surface melt. We then analyse the main drivers of surface melt and the magnitude of the snowmelt-albedo feedback at Neumayer by switching on/off the feedback process in the albedo parameterisation.

The SEB model is explained in Sect. 2.1, followed by a description of the albedo parameterisation in Sect. 2.2. The meteorological data used to force the SEB model are described in Sect. 2.3. The results section is split into two parts; in Sect. 3 we

present and discuss the SEB and melt rate that are obtained using the observed albedo. In Sect. 4 the albedo parameterisation is used instead and the snowmelt-albedo feedback is quantified and discussed. Finally, the results are briefly discussed in Sect. 5.

## 2 Methods

### 2.1 Surface energy balance model

- 5 The **one-dimensional** energy balance model is a further development of the models presented by Reijmer et al. (1999), Reijmer and Oerlema (2002), Van den Broeke et al. (2005) and Kuipers Munneke et al. (2012); here only the main features are described. The energy balance of an infinitesimally thin surface layer (the ‘skin’ layer) is defined as:

$$M = SW \downarrow + SW \uparrow + LW \downarrow + LW \uparrow + Q_S + Q_L + Q_G \quad (1)$$

- where positive fluxes are defined to be directed towards the surface.  $SW \downarrow$  and  $SW \uparrow$  are the incoming and reflected shortwave radiation,  $LW \downarrow$  and  $LW \uparrow$  are the downward and upward longwave radiation,  $Q_S$  and  $Q_L$  the turbulent sensible and latent heat fluxes and  $Q_G$  is the conductive subsurface heat flux. We neglect latent energy from rain.  $M$  is the energy used to melt snow or ice, and is non-zero only when the surface has reached the melting point of ice ( $T_s = 273.15$  K). Throughout this paper, melt and accumulation amounts are expressed in terms of mm water equivalent (mm w.e.), which equals  $\text{kg m}^{-2}$ . In order to calculate  $Q_G$  and allow for densification, meltwater percolation and refreezing, a snow/firn model is used, initialised with 70 layers. The layer thickness varies from 1 cm at the top to 2 m at the bottom (25 m depth). We impose a no-energy flux boundary condition at the lowermost model level. New snow density is parameterised following the expression of Lenaerts et al. (2012), which relates it to the prevailing surface temperature ( $T_s$ ) and 10 m wind speed ( $V_{10m}$ ) and imposes a lower limit of new snow density  $\rho_{s,0}$ . Meltwater percolation is based on the tipping-bucket method (e.g. Ligtenberg et al., 2011), allowing for immediate downward transport (within a single timestep of 10 s) of remaining water if a layer has attained its maximum capillary retention, as modelled using the expressions of Schneider and Jansson (2004). Meltwater refreezing increases the density and temperature of a layer. At the bottom of the firn layer, the meltwater is assumed to run off immediately, i.e. the model does not allow for slush/superimposed ice formation or lateral water movement. The calculation of the turbulent fluxes is based on Monin-Obukhov similarity theory between a single measurement level (2 m for temperature and humidity, 10 m for wind) and the surface, assuming the latter to be saturated with respect to ice and using the stability functions according to Dyer (1974) for unstable and Holtslag and De Bruin (1988) for stable conditions.

Subsurface penetration of shortwave radiation is calculated using a spectral model (Kuipers Munneke et al., 2009), based on the parameterisation by Brandt and Warren (1993), which is in turn based on the two-stream radiation model of Schlatter (1972). The impact on modelled melt and the quantification of the snowmelt-albedo feedback is discussed in the relevant sections.

- 30 The terms in Eq. (1) are either based on observations or can be expressed as a function of the skin temperature  $T_s$ . The SEB is solved iteratively by looking for a value of  $T_s$  that closes the SEB to within 0.005 K between iterations: if  $T_s > 273.15$  K, it is reset to 273.15 K and excess energy is used for surface melt  $M$ . To evaluate model performance, the modelled value of  $T_s$  is

compared to observed  $T_s$  calculated from  $LW \uparrow$ , using Stefan-Boltzmann's law for a longwave emissivity  $\epsilon = 1$ :

$$LW \uparrow = \sigma T_s^4, \quad (2)$$

where  $\sigma = 5.67 \cdot 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$  is the Stefan-Boltzmann constant.

Surface roughness lengths for momentum, heat and moisture are related through the expression of Andreas (1987):

$$\ln \left( \frac{z_{0*}}{z_{0,m}} \right) = a_1 + a_2 \ln(\text{Re}_*) + a_3 \ln(\text{Re}_*)^2, \quad (3)$$

where  $z_{0*}$  represents either  $z_{0,h}$  or  $z_{0,q}$ , the roughness lengths for heat and moisture respectively,  $a_1$ ,  $a_2$  and  $a_3$  are coefficients determined by Andreas (1987) for various regimes of the roughness Reynolds number  $\text{Re}_* = \frac{u_* z_{0,m}}{\nu}$  with kinematic viscosity  $\nu$  and friction velocity  $u_*$ .

## 2.2 Albedo parameterisation

- 10 Because the shortwave radiation sensor faces the sky and includes a significant direct component, measured  $SW \downarrow$  suffers from relatively large uncertainties owing to poor sensor cosine response, sensor tilt and/or rime formation (Smeets et al., 2018). In order to improve the accuracy of observed net shortwave radiation used in the SEB calculations (Sect. 3), we calculate  $SW_{net}$  based on  $SW \uparrow$ , which is diffuse and hence much less sensitive to these errors, in combination with a 24-hour moving average albedo, as described in Van den Broeke et al. (2004). In Sect. 4, in which albedo is parameterised to study melt-albedo
- 15 feedbacks, for consistency we use measured  $SW \uparrow$  in combination with parameterised albedo to estimate  $SW_{net}$ .

In Sect. 4, the parameterised surface albedo  $\alpha$  is described as a base albedo  $\alpha_S$ , modified by perturbations describing the effect of changing solar zenith angle  $\theta$  ( $d\alpha_u$ ), the cloud optical thickness  $\tau$  ( $d\alpha_\tau$ ) and the concentration of black carbon in the snow ( $d\alpha_c$ ) (Gardner and Sharp, 2010; Kuipers Munneke et al., 2011b):

$$\alpha = \alpha_S + d\alpha_u + d\alpha_\tau + d\alpha_c. \quad (4)$$

- 20 For Antarctica, we neglect the impact of impurities in the snow ( $d\alpha_c = 0$ );  $d\alpha_u$  and  $d\alpha_\tau$  both depend on the base albedo  $\alpha_S$ ,  $d\alpha_u$  in addition depends on the solar zenith angle ( $u = \cos\theta$ ), and  $d\alpha_\tau$  on the cloud optical thickness  $\tau$ :

$$d\alpha_u = 0.53\alpha_S(1 - \alpha_S)(1 - 0.64x - (1 - x)u)^{1.2}, \quad (5)$$

$$d\alpha_\tau = \frac{0.1\tau(\alpha_S + d\alpha_c)^{1.3}}{(1 + 1.5\tau)^{\alpha_S}}, \quad (6)$$

where  $x = \min\left(\frac{\sqrt{\tau}}{3u}, 1\right)$ . The base albedo depends on the snow grain size  $r_e$  (in m):

$$\alpha_S = 1.48 - 1.27048r_e^{0.07}, \quad (7)$$

in which the snow grain size  $r_e$  on time step  $t$  is parameterised as

$$r_e(t) = \left[ r_e(t-1) + dr_{e,dry} + dr_{e,wet} \right] f_o + r_{e,0} f_n + r_{e,r} f_r. \quad (8)$$

Here,  $dr_{e,dry}$  and  $dr_{e,wet}$  describe the metamorphism of dry and wet snow respectively,  $f_o$ ,  $f_n$  and  $f_r$  are the fractions of old, new and refrozen snow, and  $r_{e,0}$  and  $r_{e,r}$  are the grain sizes of **new** and refrozen snow. **Dry snow metamorphism is parameterised following Kuipers Munneke et al. (2011b)**

$$\frac{dr_{e,dry}}{dt} = \left( \frac{dr_e}{dt} \right)_0 \left( \frac{\eta}{(r_e - r_{e,0}) + \eta} \right)^{1/\kappa}, \quad (9)$$

- 5 where  $r_{e,0}$  is the new snow grain size, and the coefficients  $\left( \frac{dr_e}{dt} \right)_0$ ,  $\eta$  and  $\kappa$  are obtained from a look-up table. This look-up table is compiled based on simulations with the SNICAR model (Flanner and Zender, 2006), which calculates the snow metamorphism resulting from temperature gradient metamorphism.  $dr_{e,wet}$  is a function of the snow grain size  $r_e$  itself and the liquid water content  $f_{liq}$  (Brun et al., 1989):

$$\frac{dr_{e,wet}}{dt} = \frac{C f_{liq}^3}{4\pi r_e^2}, \quad (10)$$

- 10 where  $C$  is a constant ( $4.22 \cdot 10^{-13} \text{ m}^3 \text{ s}^{-1}$ ).

The fractions  $f_o$ ,  $f_n$  and  $f_r$  are derived from the snow/firn model, and the grain sizes of **new** and refrozen snow are constants; the method for determining their values from a tuning exercise is described in Sect. 4.1.

To determine cloud optical thickness  $\tau$ , an empirical relation between  $\tau$  and the longwave-equivalent cloud cover  $N_\epsilon$  is used following Kuipers Munneke et al. (2011a):

$$15 \quad \tau = c_1 (\exp(c_2 N_\epsilon) - 1), \quad (11)$$

with **fitting** parameters  $c_1$  and  $c_2$ .  $N_\epsilon$  is determined using a method described by Kuipers Munneke et al. (2011a) which relates hourly values of downward longwave radiation  $LW \downarrow$  to near-surface air temperature  $T_{2m}$  as illustrated in Fig. 2a. Red lines indicate quadratic fits through the upper and lower 5 percentile of the data, assuming to represent fully cloudy and clear conditions, respectively.  $N_\epsilon$  is obtained by linearly interpolating between these upper and lower bounds, yielding values between

20 0 and 1. Hourly values for cloud cover are then used to obtain values for  $\tau$  (Fig. 2b). The values used for the fit parameters  $c_1 = 5.404$  and  $c_2 = 2.207$  (both dimensionless) differ somewhat from Kuipers Munneke et al. (2011a), who used daily values for the fit.

### 2.3 Observational data

- The SEB model is forced with data from the meteorological observatory at the German research station Neumayer, situated
- 25 on the Ekström ice shelf (König-Langlo, 2017). The observatory has been operational since 1981, and was relocated in 1992 and 2009. **In 2016, its location was  $70^\circ 40'S$ ,  $8^\circ 16'W$**  (Fig. 1). The observatory is one of only four Antarctic stations –and the only one situated on an ice shelf– that is part of the Baseline Surface Radiation Network (BSRN), a global network of stations with high-quality **(artificially ventilated)** radiation observations, coordinated by the Alfred Wegener Institute (AWI). We use hourly averages of 2 m temperature ( $T_{2m}$ ) and specific humidity ( $q_{2m}$ ), 10 m wind speed ( $V_{10m}$ ), surface pressure ( $p$ )
- 30 and radiation fluxes for the period April 1992–January 2016 (24 years) to force the SEB model; **their uncertainty ranges are**

provided in Table 1. Approximately 4.1 % of the data points contained at least one missing variable, which mostly come from daily performed visual inspection of the data. To obtain a continuous data set, all missing data were replaced: pressure, relative humidity, wind speed, temperature and longwave radiation were simply linearly interpolated. In case of shortwave radiation, the missing value was replaced by imitating the average daily cycle of the two preceding days. As the measurement station is visited and maintained every day, the impact of rime formation is limited, as is the tilt of the observation mast, resulting in a high-quality meteorological data set.

Accumulation observations are only available from stake measurements, provided by AWI, which were performed weekly for the period April 1992–January 2009. As timing of precipitation is important for correctly simulating the effects of new snow on snow albedo, we combined the stake observations with precipitation predicted by the regional atmospheric climate model RACMO2.3p2 (Van Wessem et al., 2018) to obtain realistic timing of precipitation in between stake observations, as well as for the post-2009 period. The amount of precipitation modelled by RACMO2 was scaled such that the modelled surface height changes agree with stake measurements; this required a 15.3 % upward adjustment of the modelled precipitation flux.

### 2.3.1 Local near-surface climate

Neumayer station is located on an ice shelf  $\sim 20$  km from Halvfarryggen ice rise to the southeast,  $\sim 100$  km from the ice shelf break (grounding line) to the south,  $\sim 20$  km from open water/sea ice to the north and  $\sim 5$  km to open water/sea ice to the east. As a result, Neumayer experiences relatively mild conditions without significant impact from katabatic winds but with a pronounced influence of synoptic low-pressure systems passing mainly from west to east in the south Atlantic Ocean to the north of the station. The seasonal cycles of 2 m temperature, 10 m wind and 2 m specific humidity are presented in Fig. 3a. Summer temperatures around  $-4^{\circ}\text{C}$  and winter temperatures around  $-25^{\circ}\text{C}$  imply a substantial ( $>20$  K) seasonal temperature amplitude based on monthly mean values. This is in line with the formation of a surface-based temperature inversion in winter, a phenomenon that is representative for the flat ice shelves as well as the interior ice domes and in contrast to the topographically steeper escarpment zone, where the quasi-continuous mixing by katabatic flow limits the formation of such an inversion (Van den Broeke, 1998). As expected from the strong link to the air temperature through the Clausius-Clapeyron relation and a high annual mean relative humidity of 82 % (relative to either water or ice, depending on the air temperature) because of the proximity of a saturated snow surface and the ocean, the seasonal cycle of  $q_{2m}$  closely follows that of temperature.

## 3 Results: surface energy balance and melt

### 3.1 SEB model performance and uncertainties

There are several SEB model parameters for which the exact values or formulations are unknown, e.g. the surface roughness length for momentum  $z_{0,m}$ , the density of new snow  $\rho_s$ , the stability functions (required to calculate the turbulent scales) and the effective conductivity, which couples the magnitude of  $Q_G$  to the temperature gradient in the snow. We estimated the impact of observational and model uncertainties on modelled melt by running the model 600 times while randomly varying all

hourly observations within the specified measurement uncertainty ranges (Table 1) and using multiple expressions for the heat conductivity and stability functions. Model performance is quantified by comparing modelled with observed  $T_s$  and assessing the changes in 24-year total cumulative melt. **Note that in this section, the albedo based on observations is used to obtain  $SW_{net}$ .**

5 The choice of expressions for the stability functions and heat conductivity did not significantly impact the modelled amount of melt (total within 30 mm w.e. or 2.7 %, not shown). The model outcomes are more sensitive to the choice of surface roughness length for momentum  $z_{0,m}$  and the lower limit of density of **new** snow  $\rho_{s,0}$ : when  $z_{0,m}$  is varied between 0.5 mm and 50 mm and  $\rho_{s,0}$  between 150 and 500 kg m<sup>-3</sup>, the cumulative amount of surface melt over the 24-year period varies between **900–1220 mm w.e.**, with higher melt values for smaller values of  $z_{0,m}$  and  $\rho_{s,0}$ . Optimal values in terms of simulated  $T_s$  are  
10  $z_{0,m} = 1.65 \text{ mm}$  and  $\rho_{s,0} = 280 \text{ kg m}^{-3}$ , resulting in a  $T_s$  bias of 0.01 K and an RMSD of **0.79 K** (Fig. 4). We use these values in the remainder of this study. Figure 5a and b show modelled 24-year cumulative melt **and annual melt (Mar–Feb)** at Neumayer, combined with uncertainties associated with model parameters. The annual mean values for year  $X$  are obtained by averaging monthly values for March of year  $X$  until February of year  $X + 1$ . The total melt amounts to **1154 mm w.e.**, with a small uncertainty associated with measurement uncertainties ( $1\sigma \approx 3 \text{ mm w.e.}$ , i.e. 0.3 %). The adopted method to estimate this  
15 uncertainty has its limitations, as measurement errors are probably autocorrelated: if a measurement at one time is disturbed in some way, it is probably disturbed in a similar way at the next time step. Therefore, this result could be interpreted as a lower bound of the uncertainty range, which is supported by the larger uncertainty estimates ( $\sim 15 \%$ ) by Van den Broeke et al. (2010), who applied a constant systematic error which can be interpreted as an upper bound on the modelled uncertainty range. This also explains why the uncertainties deriving from the choice of  $z_{0,m}$  are so much larger: these runs represent prescribing  
20 a systematic error between the true (unknown) value and the chosen value, assuming the true value to be constant, which likely is an oversimplification (Smeets and Van den Broeke, 2008).

The sensitivity to  $z_{0,m}$  is somewhat unexpected; **following Eq. (3)** both  $z_{0,h}$  and  $z_{0,q}$  decrease for increasing  $z_{0,m}$ , which acts to dampen the effect on the magnitude of the turbulent fluxes. Our interpretation of this result is that decreasing  $z_{0,m}$  and  $\rho_{s,0}$  decreases the turbulent fluxes as well as  $Q_G$ , reducing the efficiency **with** which heat is removed from the surface,  
25 allowing more energy to be invested in melt. The **obtained** value for  $z_{0,m} = 1.65 \text{ mm}$  is high compared to the **average value of  $z_{0,m} = 0.1 \text{ mm}$**  found during a field campaign **at Neumayer** in 1982 (König, 1985) **but it is not uncommon for snow surfaces (Amory et al., 2017).** Measured values of  $T_s$  in excess of the melting point in Fig. 4 **only occurred in the first 6 seasons; from 1998–99 onwards they were removed by additional post-processing. These measurements** mainly reflect uncertainties in the adopted unit value of longwave emissivity and in measured  $LW \uparrow$ , e.g. from sensor window heating (Smeets et al., 2018) **and**  
30 **the fact that the downward facing radiation sensor also measures longwave radiation emitted by the relatively warm air between the surface and the sensor.**

### 3.2 Surface energy balance

Annual (Mar–Feb) mean **values of** near-surface meteorological quantities and SEB components are presented in **Table 2, with seasonal cycles of SEB components presented in Fig. 3b. These** show that the **summertime** SEB is dominated by the radiation



fluxes; **despite** the high albedo of the snow surface,  $SW_{net}$  is the dominant heat source for the skin layer, **whereas**  $LW_{net}$  extracts energy from the surface, most efficiently so in summer when the surface is heated by the sun. In summer,  $Q_L$  becomes a significant heat loss in the SEB (sublimation), preventing strong negative  $Q_S$  (convection). The seasonal cycle of  $Q_G$  is small, indicating a small net transport of heat away from the surface in summer and towards the surface in winter. The net annually integrated amount is less than zero as a result of the refreezing of meltwater, warming the subsurface snow layers.

Significant and previously unreported trends (**not shown**) are detected in  $LW \uparrow (-0.28 \pm 0.14 \text{ W m}^{-2} \text{ yr}^{-1})$  and  $Q_S (+0.21 \pm 0.07 \text{ W m}^{-2} \text{ yr}^{-1})$ , both a result of wintertime trends.  $LW \uparrow$  is linked directly to  $T_s$ , which **shows an insignificant** negative trend ( $-0.029 \pm 0.026 \text{ K yr}^{-1}$ ), which in magnitude exceeds the negative trend in  $T_{2m}$  ( $-0.0045 \pm 0.02 \text{ K yr}^{-1}$ , the probability that the negative trend in  $T_s$  is greater in magnitude than the trend in  $T_{2m}$  is **0.76**). As a result, the temperature gradient near the surface has increased, enhancing  $Q_S$ . The negative trend in  $T_s$  originates from a decrease in  $LW \downarrow (-0.26 \pm 0.17 \text{ W m}^{-2} \text{ yr}^{-1})$ , which is in turn driven by a slight decrease in cloud cover ( $-0.003 \pm 0.001 \text{ yr}^{-1}$ ). This is suggested independently by the decrease in average winter humidity ( $-0.004 \pm 0.002 \text{ g kg}^{-1} \text{ yr}^{-1}$ ). These findings agree with Herman et al. (2013) **and Kuipers Munneke et al. (2011a)**, who determined from satellite observations that summer cloud cover has decreased over that part of coastal Antarctica in the period 1979–2011.

### 3.3 Melt season

Melt occurs at Neumayer from November until February (Fig. 6), but is highly variable from year to year. The mean annual amount of melt is **50 mm w.e.** with an interannual variability of **42 mm w.e.** and a range of 2 mm w.e. in 1999–2000 to **176 mm w.e.** in 2012–13. Most melt occurs in December and January and the surface only sporadically reaches the melting point in February. Only in 2007 did melt occur in November, and no melt occurs outside these four months. The cumulative melt occurring at Neumayer shows **step-wise** increases (Fig. 5a), which **represent** the peaked melt seasons, in which melt occurs on average on **18 ± 10 days**. The uncertainty in the number of melt days due to the chosen values of  $z_{0,m}$  and  $\rho_{s,0}$  is relatively small compared to the interannual variability in melt totals (Fig. 6), implying that this choice does not significantly affect the modelled melt duration, but it does affect the total melt.

To investigate the link between melt and climate, we compare the two summers with the highest (2003–04 and 2012–13, on average **145 mm w.e.**) and lowest (1999–2000 and 2014–15, on average **4 mm w.e.**) melt amounts. Figure 7 shows the meteorological and SEB components for these years, averaged over December and January. The largest differences are found in  $T_{2m}$  (**+2.3 K**) and  $SW_{net}$  ( $+17 \text{ W m}^{-2}$ ); based on the measurement uncertainties (**Table 1**), these differences are significant. In cold summers, the low  $T_{2m}$  corresponds to a stronger temperature inversion ( $T_{2m} - T_s$ ), more longwave cooling, less sublimation and a larger  $Q_S$ .  **$SW \downarrow$  and  $LW \downarrow$  show almost no difference between high and low melt seasons; therefore, the difference in  $SW_{net}$  cannot be caused by a change in cloud cover and is likely** caused solely by surface albedo, which suggests an important role for the snowmelt-albedo feedback. This will be elaborated upon in the next section. Finally, the direction of  $Q_G$  is reversed; in high melt years, the surface is warmed from below while in low melt years the surface loses heat to the subsurface. More refreezing of meltwater in high melt years warms the near surface snow layers, which in turn leads to a conductive heat flux towards the surface.



Using the subsurface radiation model of Kuipers Munneke et al. (2009), the influence of subsurface penetration of shortwave radiation is estimated. Its inclusion increases the cumulative amount of melt by 13 %, from 1154 mm w.e. to 1326 mm w.e. The absorbed shortwave radiation heats the subsurface layers, but the heat cannot be transported away as effectively as would happen at the surface by turbulent fluxes and longwave radiation. This leads to an increase in total melt.

- 5 The findings presented in this section are in good agreement with Van den Broeke et al. (2010), who used a similar approach to calculate the SEB at Neumayer, but used a lower value for  $z_{0,m} = 0.32$  mm and a higher snow density that was assumed constant with depth ( $420 \text{ kg m}^{-3}$  cf.  $320 \text{ kg m}^{-3}$ ). Compared to melt estimates from Larsen C ice shelf, obtained through a similar modelling approach by Kuipers Munneke et al. (2012), melt at Neumayer is weak. Owing to its more northerly location, on Larsen C ice shelf an annual (2009–2011) average melt energy of  $2.8 \text{ W m}^{-2}$  is obtained, compared to the 2009–2011 annual  
10 average of  $0.7 \text{ W m}^{-2}$  obtained at Neumayer; furthermore, in November and February melt occurs much more frequently on Larsen C ice shelf.

#### 4 Results: the snowmelt-albedo feedback

- The snowmelt-albedo feedback is a well-known phenomenon, but has not before been quantified for Antarctica. The feedback occurs after the rapid growth of snow grains when meltwater penetrates into the subsurface and refreezes. Because a photon  
15 on average travels farther through snow with large particles than in new snow with smaller particles, the probability of it being absorbed is increased, effectively lowering the surface albedo (Gardner and Sharp, 2010). Even without melt, albedo decreases when snow ages, following grain growth from dry snow metamorphism, but this is a much slower process which mainly depends on temperature gradients in the snow, favouring moisture transport onto larger grains. Precipitation of new, fine-grained snow has been shown to inhibit the albedo decrease by metamorphism on the Antarctic plateau (Picard et al.,  
20 2012).

- To quantify the snowmelt-albedo feedback at Neumayer, we need to be able to switch on and off the albedo dependency on melt-driven grain growth. To that end, we implemented an albedo parameterisation in the SEB model, as described in Sect. 2.2. Because no data on grain size are available from Neumayer, we optimised the albedo model performance by maximising the correspondence between 1) modelled and observed hourly  $SW \uparrow$ , and 2) the total melt obtained from the calculations based on  
25 observed albedo (Sect. 4.1). We compare  $SW \uparrow$  instead of the albedo itself because by doing so the hourly values are naturally weighted with its contribution to  $S_{net}$  and hence its importance for the SEB. We then perform several runs with different processes switched on and off affecting the surface albedo to investigate the importance of the snowmelt-albedo feedback for melt at Neumayer (Sect. 4.2).

##### 4.1 Optimising the albedo parameterisation

- 30 The albedo parameterisation, and especially the expression for snow grain size (Eq. (8)) contains several parameters that are not well constrained, such as new snow grain size  $r_{e,0}$  and refrozen snow grain size  $r_{e,r}$ . These parameters were varied within reasonable ranges to optimise the results: new snow grain size between 0.04 mm and 0.3 mm, refrozen snow grain size between

0.1 mm and 10 mm. The best comparison with observed albedo was achieved when using the look-up table for dry snow metamorphism  $dr_{e,dry}$  corresponding to a grain size of 0.055 mm.

The first step in optimising the parameterisation was to split the summer season into two parts, the ‘dry’ and the ‘wet’ season. The respective starts of the dry and wet seasons are the first day on which the sun rises more than 15° above the horizon and the first day that surface melt occurs. The wet season ends when the sun no longer rises higher than 15°. For the dry season, we varied the dry snow metamorphism factor and the new snow grain size to best match observed  $SW \uparrow$ . This resulted in a new snow grain size of 0.25 mm. This value is then used in the second step, in which the refrozen snow grain size  $r_{e,r}$  is varied to best match the cumulative melt using observed albedo. This was achieved for a refrozen snow grain size of 1.45 mm.

This value for refrozen snow grain size is compatible with the typical largest grains in dry metamorphosed snow of O(1 mm), and which Kuipers Munneke et al. (2011b) used as a lower limit for refrozen snow grains. Libois et al. (2015) and Picard et al. (2016) present observations of snow grain sizes on the Antarctic plateau during field campaigns in 2012–13 and 2013–14 as well as estimates from satellite observations. On the plateau, summer temperatures are comparable to Neumayer winter temperatures. Libois et al. (2015) report summertime snow grain size estimates of approximately 0.11 mm (Fig. 6 in their study, reported as a specific surface area  $SSA = \frac{3}{\rho_i r_e}$ , where  $\rho_i$  is the density of ice and  $r_e$  is the snow grain size). In our study, wintertime snow grain sizes approach 0.21 mm. The difference is expected as the plateau is generally much colder than Neumayer. The seasonal cycle of modelled average snow grain size in the upper 7 cm (Fig. 8) is comparable to the one presented in Libois et al. (2015).

When the adopted albedo values are combined with the observations of  $SW \uparrow$ , the model adequately reproduces the incoming shortwave radiation (Fig. 9, bias = +0.93 W m<sup>-2</sup>, RMSD = 7.3 W m<sup>-2</sup>), providing confidence in the modelled albedo.

## 4.2 Magnitude of the snowmelt-albedo feedback

Three experiments with the SEB model were carried out in addition to the original run ( $R_0$ ) which uses the measured albedo:

- $R_1$ : the average measured albedo (0.84, determined by adding all  $SW \downarrow$  and  $SW \uparrow$  for all measurements when the Sun is higher than 15° above the horizon and taking the ratio between the two) is prescribed for the entire period;
- $R_2$ : the full albedo parameterisation is used;
- $R_3$ : refrozen snow does not contribute to the changing snow characteristics, i.e.  $f_r = 0$  in Eq. (8).

Figure 10a and b show time series of cumulative and seasonal surface melt for the four experiments. Experiment  $R_1$  underpredicts melt in most seasons, yielding a mean annual amount of surface melt of  $39 \pm 27$  mm w.e. yr<sup>-1</sup> (c.f.  $50 \pm 42$  mm w.e. yr<sup>-1</sup> for experiment  $R_0$ ). More melt was modelled in the 1995–96 melt season, which was characterised by frequent precipitation events and cloudy conditions, keeping observed albedo higher than the long-term mean. The experiment using the full albedo parameterisation ( $R_2$ ) adequately reproduces the amount of seasonal melt ( $50 \pm 34$  mm w.e. yr<sup>-1</sup>), although melt in e.g. the 2012 melt season is underestimated. Run  $R_3$  represents the situation in which the snowmelt-albedo feedback has been switched off, leading to significantly underpredicted melt ( $21 \pm 16$  mm w.e. yr<sup>-1</sup>).

Defining the strength of the snowmelt-albedo feedback (SMAF) as the ratio between the total seasonal surface melt in experiments  $R_2$  and  $R_3$ , we obtain an average value of 2.6, with a range of 1.3 (1996–97) to 4.8 (1993–94, see Fig. 10c). The effect of subsurface penetration of shortwave radiation on this result is estimated by repeating the above experiments with inclusion of the radiation penetration model of Kuipers Munneke et al. (2009). This yielded an average SMAF of 2.3, ranging from 1.5 (2005–06) to 3.2 (2002–03). The main difference between the two experiments is the reduced interannual variability: including penetration of shortwave radiation does not yield SMAF values larger than 3.5. Shortwave radiation penetration heats the subsurface, causing subsurface melt which is less affected by the snowmelt-albedo feedback because the radiative flux is smaller in the subsurface. Therefore, the ‘extreme’ years in the sense of SMAF are less distinct in the experiment with shortwave radiation penetration. The effect of shortwave radiation penetration is included in the uncertainties indicated in Fig. 10c. Combining this with the uncertainties in observed  $SW \uparrow$  and the determination of  $\tau$  (Fig. 2b) leads to uncertainties in the determination of the SMAF of typically 15 %, with a range of 4 % (1995–96) to 32 % (1993–94).

A weak positive correlation was found between SMAF and  $SW \downarrow$  ( $R^2 = 0.15$ ,  $p = 0.07$ ); if  $SW \downarrow$  increases, more energy is available at the surface for melting, which is then in turn further intensified by SMAF. Another weak negative correlation was found between SMAF and summer precipitation ( $R^2 = 0.13$ ,  $p = 0.09$ ); snowfall inhibits SMAF as it effectively ‘resets’ the surface albedo as was also shown by Picard et al. (2012). Alternatively, SMAF could be defined as the ratio between  $R_0$  and  $R_3$ , or the ratio between  $R_0$  and  $R'_1$ , where  $R'_1$  uses the average albedo at the end of the winter. Using the former definition, the results become more prone to noise due to the performance of the albedo parameterisation itself. The latter definition neglects the dependency on solar zenith angle and the impact of precipitation. Therefore, we believe defining SMAF as the ratio between  $R_2$  and  $R_3$  is more consistent.

Only few studies report on the snowmelt-albedo feedback concerning the darkening of snow rather than disappearance of it. Box et al. (2012) provide relationships between anomalies of seasonal  $T_{2m}$  and  $SW_{net}$  (Fig. 5 and 12 of Box et al. (2012)). They find a negative relationship for accumulation regions, i.e. lower 2 m temperatures are associated with smaller  $SW_{net}$ . No such relationship is found for Neumayer (not shown).

## 5 Conclusions

In this study, we used 24 years of high-quality meteorological and radiation observations from the BSRN station Neumayer, situated on Ekström ice shelf, East Antarctica, to force a surface energy balance model. The primary goal was to calculate the amount of melt at Neumayer and to investigate the importance of the snowmelt-albedo feedback. Model performance was evaluated based on the difference between modelled and measured surface temperature, and the calculated melt was tested for measurement and model parameter uncertainties. We found that measurement uncertainties, when considered random in time, do not significantly impact modelled melt at Neumayer (<0.5 % difference). However, melt amount and model performance are sensitive to the values chosen for the surface roughness length for momentum  $z_{0,m}$  and lower limit of new snow density  $\rho_{s,0}$ , thus accurate measurements of these values would further improve future modelling studies. Our results confirm that melt at Neumayer is an intermittent process, occurring on average on only 18 days each summer, totalling 50 mm w.e. and

with an interannual variability of 42 mm w.e. Melt occurs mainly in December and January, sporadically in February and only once melt was modelled in November. Significant and previously unreported trends were found in the net longwave radiation (decreasing) and the sensible heat flux (increasing), but these are unrelated to the melt at Neumayer as they mainly occur in winter and are attributed to a decrease in cloud cover.

- 5 The main difference between high and low melt years was found to be surface albedo, implying an important role for the snowmelt-albedo feedback (SMAF). We quantified SMAF by implementing and tuning an albedo parameterisation in the SEB model, which includes the effects of snowfall and wet and dry snow metamorphism on albedo. The albedo parameterisation adequately reproduces the seasonal variability in snow grain size, compared to measurements on the Antarctic Plateau (Libois et al., 2015). Our derived wintertime snow grain sizes at Neumayer are somewhat smaller than the satellite-derived
- 10 summertime snow grain sizes at the Antarctic Plateau (Libois et al., 2015) owing to the lower temperatures on the plateau. Our main finding is that SMAF on average enhances surface melt at Neumayer by a factor of  $2.6 \pm 0.8$ .

Weak correlations were found of SMAF with summertime  $SW \downarrow$  and precipitation ( $0.1 < R^2 < 0.2$ ). To assess how the importance of the snowmelt-albedo feedback varies spatially and temporally, the next step in this research will be applying this method to other sites in Antarctica and a regional climate model (Van Wessem et al., 2018).

- 15 *Code and data availability.* The Neumayer data are available upon request via the website of AWI (<https://bsrn.awi.de/data/data-retrieval-via-pangaea/>). The model output is available upon request by the authors.

*Author contributions.* CLJ performed the study and wrote the manuscript. PKM assisted with the implementation of the albedo parameterisation. GKL was in charge of the Neumayer data. CHR, PKM, GKL and MRvdB have commented on the manuscript,

*Competing interests.* The authors declare no competing interests.

- 20 *Acknowledgements.* We would like to thank the AWI for maintaining the station and the Baseline Surface Radiation Network (BSRN) for providing the data, with special thanks to Amelie Driemel for creating a citation reference and Holger Schmithüsen for helping to interpret the data. C.L. Jakobs and C.H. Reijmer acknowledge support from the Netherlands Polar Programme under project number 866.15.204. M.R. van den Broeke acknowledges support from the Netherlands Earth System Science Centre (NESSC). We would like to thank Ghislain Picard and one anonymous referee for their constructive comments.

## References


- Amory, C., Gallée, H., Naaim-Bouvet, F., Favier, V., Vignon, E., Picard, G., Trouvilliez, A., Piard, L., Genthon, C., and Bellot, H.: Seasonal Variations in Drag Coefficient over a Sastrugi-Covered Snowfield in Coastal East Antarctica, *Boundary-Layer Meteorology*, 164, 107–133, <https://doi.org/10.1007/s10546-017-0242-5>, 2017.
- 5 Andreas, E. L.: A theory for the scalar roughness and the scalar transfer coefficients over snow and sea ice, *Boundary-Layer Meteorology*, 38, 159–184, <https://doi.org/10.1007/BF00121562>, 1987.
- Bell, R. E., Chu, W., Kingslake, J., Das, I., Tedesco, M., Tinto, K. J., Zappa, C. J., Frezzotti, M., Boghosian, A., and Lee, W. S.: Antarctic ice shelf potentially stabilized by export of meltwater in surface river, *Nature*, 544, 344–348, <https://doi.org/10.1038/nature22048>, 2017.
- Box, J. E., Fettweis, X., Stroeve, J. C., Tedesco, M., Hall, D. K., and Steffen, K.: Greenland ice sheet albedo feedback: thermodynamics and  
10 atmospheric drivers, *The Cryosphere*, 6, 821–839, <https://doi.org/10.5194/tc-6-821-2012>, 2012.
- Brandt, R. E. and Warren, S. G.: Solar-heating rates and temperature profiles in Antarctic snow and ice, *Journal of Glaciology*, 39, 99–110, <https://doi.org/10.3198/1993JoG39-131-99-110>, 1993.
- Brun, E., Martin, E., Simon, V., Gendre, C., and Coléou, C.: An Energy and Mass Model of Snow Cover Suitable for Operational Avalanche Forecasting, *Journal of Glaciology*, 35, 333–342, <https://doi.org/10.3189/S0022143000009254>, 1989.
- 15 DeConto, R. M. and Pollard, D.: Contribution of Antarctica to past and future sea-level rise, *Nature*, 531, 591–597, <https://doi.org/10.1038/nature17145>, 2016.
- Dyer, A. J.: A review of flux-profile relationships, *Boundary-Layer Meteorology*, 7, 363–372, <https://doi.org/10.1007/BF00240838>, 1974.
- Flanner, M. G. and Zender, C. S.: Linking snowpack microphysics and albedo evolution, *Journal of Geophysical Research*, 111, <https://doi.org/10.1029/2005JD006834>, 2006.
- 20 Flanner, M. G., Zender, C. S., Randerson, J. T., and Rasch, P. J.: Present-day climate forcing and response from black carbon in snow, *Journal of Geophysical Research*, 112, <https://doi.org/10.1029/2006JD008003>, 2007.
- Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R. E., Bianchi, C., Bingham, R. G., Blankenship, D. D., Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita, S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W., Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langley, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P.,  
25 Matsuoka, K., Mouginot, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippin, D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C., and Zirizzotti, A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, *The Cryosphere*, 7, 375–393, <https://doi.org/10.5194/tc-7-375-2013>, 2013.
- 30 Gardner, A. S. and Sharp, M. J.: A review of snow and ice albedo and the development of a new physically based broadband albedo parameterization, *Journal of Geophysical Research: Earth Surface*, 115, <https://doi.org/10.1029/2009JF001444>, 2010.
- Hall, A.: The Role of Surface Albedo Feedback in Climate, *Journal of Climate*, 17, 1550–1568, [https://doi.org/10.1175/1520-0442\(2004\)017<1550:TROSAF>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<1550:TROSAF>2.0.CO;2), 2004.
- Herman, J., DeLand, M. T., Huang, L.-K., Labow, G., Larko, D., Lloyd, S. A., Mao, J., Qin, W., and Weaver, C.: A net decrease in the Earth’s  
35 cloud, aerosol, and surface 340 nm reflectivity during the past 33yr (1979–2011), *Atmospheric Chemistry and Physics*, 13, 8505–8524, <https://doi.org/10.5194/acp-13-8505-2013>, 2013.

- Hogg, A. E. and Gudmundsson, G. H.: Impacts of the Larsen-C Ice Shelf calving event, *Nature Climate Change*, 7, 540–542, <https://doi.org/10.1038/nclimate3359>, 2017.
- Holtslag, A. A. M. and De Bruin, H. A. R.: Applied Modeling of the Nighttime Surface Energy Balance over Land, *Journal of Applied Meteorology*, 27, 689–704, [https://doi.org/10.1175/1520-0450\(1988\)027<0689:AMOTNS>2.0.CO;2](https://doi.org/10.1175/1520-0450(1988)027<0689:AMOTNS>2.0.CO;2), 1988.
- 5 Kingslake, J., Ely, J. C., Das, I., and Bell, R. E.: Widespread movement of meltwater onto and across Antarctic ice shelves, *Nature*, 544, 349–352, <https://doi.org/10.1038/nature22049>, 2017.
- König, G.: Roughness Length of an Antarctic Ice Shelf, *Polarforschung*, 55, 27–32, <https://doi.org/10.2312/polarforschung.55.1.27>, 1985.
- König-Langlo, G.: Basic and other measurements, and meteorological synoptical observations from Neumayer Station, 1992-04 to 2016-01, reference list of 572 datasets, <https://doi.pangaea.de/10.1594/PANGAEA.874984>, 2017.
- 10 Kuipers Munneke, P., Van den Broeke, M. R., Reijmer, C. H., Helsen, M. M., Boot, W., Schneebeli, M., and Steffen, K.: The role of radiation penetration in the energy budget of the snowpack at Summit, Greenland, *The Cryosphere*, 3, 155–165, <https://doi.org/10.5194/tc-3-155-2009>, 2009.
- Kuipers Munneke, P., Reijmer, C. H., and Van den Broeke, M. R.: Assessing the retrieval of cloud properties from radiation measurements over snow and ice, *International Journal of Climatology*, 31, 756–769, <https://doi.org/10.1002/joc.2114>, 2011a.
- 15 Kuipers Munneke, P., Van den Broeke, M. R., Lenaerts, J. T. M., Flanner, M. G., Gardner, A. S., and Van de Berg, W. J.: A new albedo parameterization for use in climate models over the Antarctic ice sheet, *Journal of Geophysical Research: Atmospheres*, 116, <https://doi.org/10.1029/2010JD015113>, 2011b.
- Kuipers Munneke, P., Van den Broeke, M. R., King, J. C., Gray, T., and Reijmer, C. H.: Near-surface climate and surface energy budget of Larsen C ice shelf, Antarctic Peninsula, *The Cryosphere*, 6, 353–363, <https://doi.org/10.5194/tc-6-353-2012>, 2012.
- 20 Kuipers Munneke, P., Ligtenberg, S. R. M., Van den Broeke, M. R., and Vaughan, D. G.: Firn air depletion as a precursor of Antarctic ice-shelf collapse, *Journal of Glaciology*, 60, 205–214, <https://doi.org/10.3189/2014JoG13J183>, 2014.
- Kuipers Munneke, P., Luckman, A. J., Bevan, S. L., Smeets, C. J. P. P., Gilbert, E., Van den Broeke, M. R., Wang, W., Zender, C. S., Hubbard, B., Ashmore, D., Orr, A., King, J. C., and Kulesa, B.: Intense Winter Surface Melt on an Antarctic Ice Shelf, *Geophysical Research Letters*, 45, <https://doi.org/10.1029/2018GL077899>, 2018.
- 25 Lenaerts, J. T. M., Van den Broeke, M. R., Déry, S. J., Van Meijgaard, E., Van de Berg, W. J., Palm, S. P., and Sanz Rodrigo, J.: Modeling drifting snow in Antarctica with a regional climate model: 1. Methods and model evaluation, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/10.1029/2011JD016145>, 2012.
- Lenaerts, J. T. M., Lhermitte, S., Drews, R., Ligtenberg, S. R. M., Berger, S., Helm, V., Smeets, C. J. P. P., Van den Broeke, M. R., Van de Berg, W. J., Van Meijgaard, E., Eijkelboom, M., Eisen, O., and Pattyn, F.: Meltwater produced by wind-albedo interaction stored in an East Antarctic ice shelf, *Nature Climate Change*, 7, 58–62, <https://doi.org/10.1038/nclimate3180>, 2017.
- 30 Libois, Q., Picard, G., Arnaud, L., Dumont, M., Lafaysse, M., Morin, S., and Lefebvre, E.: Summertime evolution of snow specific surface area close to the surface on the Antarctic Plateau, *The Cryosphere*, 9, 2383–2398, <https://doi.org/10.5194/tc-9-2383-2015>, 2015.
- Ligtenberg, S. R. M., Helsen, M. M., and Van den Broeke, M. R.: An improved semi-empirical model for the densification of Antarctic firn, *The Cryosphere*, 5, 809–819, <https://doi.org/10.5194/tc-5-809-2011>, 2011.
- 35 Ligtenberg, S. R. M., Kuipers Munneke, P., and Van den Broeke, M. R.: Present and future variations in Antarctic firn air content, *The Cryosphere*, 8, 1711–1723, <https://doi.org/10.5194/tc-8-1711-2014>, 2014.
- Luckman, A., A., E., Jansen, D., Kulesa, B., Kuipers Munneke, P., King, J. C., and Barrand, N. E.: Surface melt and ponding on Larsen C Ice Shelf and the impact of föhn winds, *Antarctic Science*, 26, 625–635, <https://doi.org/10.1017/S0954102014000339>, 2014.


- Massom, R. A., Scambos, T. A., Bennetts, L. G., Reid, P. A., Squire, V. A., and Stammerjohn, S. E.: Antarctic ice shelf disintegration triggered by sea ice loss and ocean swell, *Nature*, 558, 383–389, <https://doi.org/10.1038/s41586-018-0212-1>, 2018.
- Perovich, D. K., Grenfell, T. C., Light, B., and Hobbs, P. V.: Seasonal evolution of the albedo of multiyear Arctic sea ice, *Journal of Geophysical Research*, 107, <https://doi.org/10.1029/2000JC000438>, 2002.
- 5 Picard, G., Fily, M., and Gallée, H.: Surface melting derived from microwave radiometers: a climatic indicator in Antarctica, *Annals of Glaciology*, 46, 29–34, <https://doi.org/10.3189/172756407782871684>, 2007.
- Picard, G., Domine, F., Krinner, G., Arnaud, L., and Lefebvre, E.: Inhibition of the positive snow-albedo feedback by precipitation in interior Antarctica, *Nature Climate Change*, 2, 795–798, <https://doi.org/10.1038/nclimate1590>, 2012.
- Picard, G., Libois, Q., Arnaud, L., Verin, G., and Dumont, M.: Development and calibration of an automatic spectral albedometer to estimate near-surface snow SSA time series, *The Cryosphere*, 10, 1297–1316, <https://doi.org/10.5194/tc-10-1297-2016>, 2016.
- 10 Qu, X. and Hall, A.: What Controls the Strength of Snow-Albedo Feedback?, *Journal of Climate*, 20, 3971–3981, <https://doi.org/10.1175/JCLI4186.1>, 2007.
- Reijmer, C. H. and Oerlemans, J.: Temporal and spatial variability of the surface energy balance in Dronning Maud Land, East Antarctica, *Journal of Geophysical Research*, 107, <https://doi.org/10.1029/2000JD000110>, 2002.
- 15 Reijmer, C. H., Greuell, W., and Oerlemans, J.: The annual cycle of meteorological variables and the surface energy balance on Berkner Island, Antarctica, *Annals of Glaciology*, 29, 49–54, <https://doi.org/10.3189/172756499781821166>, 1999.
- Rignot, E., Casassa, G., Gogineni, P., Krabill, W., Rivera, A., and Thomas, R.: Accelerated ice discharge from the Antarctic Peninsula following the collapse of Larsen B ice shelf, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004GL020697>, 2004.
- Scambos, T. A., Bohlander, J. A., Shuman, C. A., and Skvarca, P.: Glacier acceleration and thinning after ice shelf collapse in the Larsen B embayment, Antarctica, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004GL020670>, 2004.
- 20 Schlatter, T. W.: The Local Surface Energy Balance and Subsurface Temperature Regime in Antarctica, *Journal of Applied Meteorology*, 11, 1048–1062, [https://doi.org/10.1175/1520-0450\(1972\)011<1048:TLSEBA>2.0.CO;2](https://doi.org/10.1175/1520-0450(1972)011<1048:TLSEBA>2.0.CO;2), 1972.
- Schneider, T. and Jansson, P.: Internal accumulation in firn and its significance for the mass balance of Storglaciären, Sweden, *Journal of Glaciology*, 50, 25–34, <https://doi.org/10.3189/172756504781830277>, 2004.
- 25 Shepherd, A., Ivins, E. R., Rignot, E., Smith, B., Van den Broeke, M. R., Velicogna, I., Whitehouse, P. L., Briggs, K. H., Joughin, I., Krinner, G., Nowicki, S., Payne, A. J., Scambos, T. A., Schlegel, N., A. G., Agosta, C., Ahlstrøm, A. P., Babonis, G., Barletta, V. R., Blazquez, A., Bonin, J., Csatho, B., Cullather, R. I., Felikson, D., Fettweis, X., Forsberg, R., Gallée, H., Gardner, A. S., Gilbert, L., Groh, A., Gunter, B., Hanna, E., Harig, C., Helm, V., Horvath, A., Horwath, M., Khan, S., Kjeldsen, K. K., Konrad, H., Langen, P., Lecavalier, B., Loomis, B., Luthcke, S. B., McMillan, M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P., Mouginot, J., Moyano, G., Muir, A., Nagler, T., Nield, G., Nilsson, J., Noël, B. P. Y., Otsuka, I., Pattie, M. E., Peltier, W. R., Pie, N., Rietbroek, R., Rott, H., Sandberg Sørensen, L., Sasgen, I., Save, H., Scheuchl, B., Schrama, E. J. O., Schröder, L., Seo, K.-W., Simonsen, S., Slater, T., Spada, G., Sutterly, T. C., Talpe, M., Tarasov, L., Van de Berg, W. J., Van der Wal, W., Van Wessem, J. M., Vishwakarma, B. D., Wiese, D., and Wouters, B.: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558, 219–222, <https://doi.org/10.1038/s41586-018-0179-y>, 2018.
- 30 Smeets, C. J. P. P. and Van den Broeke, M. R.: Temporal and Spatial Variations of the Aerodynamic Roughness Length in the Ablation Zone of the Greenland Ice Sheet, *Boundary-Layer Meteorology*, 128, 315–338, <https://doi.org/10.1007/s10546-008-9291-0>, 2008.
- Smeets, C. J. P. P., Kuipers Munneke, P., Van As, D., Van den Broeke, M. R., Boot, W., Oerlemans, J., Snellen, H., Reijmer, C. H., and Van de Wal, R. S. W.: The K-transect in west Greenland: Automatic weather station data (1993–2016), *Arctic, Antarctic, and Alpine Research*, 50, <https://doi.org/10.1080/15230430.2017.1420954>, 2018.



- Tedesco, M.: Assessment and development of snowmelt retrieval algorithms over Antarctica from K-band spaceborn brightness temperature (1979–2008), *Remote Sensing of Environment*, 113, 979–997, <https://doi.org/10.1016/j.rse.2009.01.009>, 2009.
- Trusel, L. D., Frey, K. E., Das, S. B., Kuipers Munneke, P., and Van den Broeke, M. R.: Satellite-based estimates of Antarctic surface meltwater fluxes, *Geophysical Research Letters*, 40, 6148–6153, <https://doi.org/10.1002/2013GL058138>, 2013.
- 5 Trusel, L. D., Frey, K. E., Das, S. B., Karnauskas, K. B., Kuipers Munneke, P., Van Meijgaard, E., and Van den Broeke, M. R.: Divergent trajectories of Antarctic surface melt under two twenty-first-century climate scenarios, *Nature Geoscience*, 8, 927–934, <https://doi.org/10.1038/NGEO2563>, 2015.
- Turner, J., Orr, A., Gudmundsson, G. H., Jenkins, A., Bingham, R. G., Hillenbrand, C.-D., and Bracegirdle, T. J.: Atmosphere-ocean-ice interactions in the Amundsen Sea Embayment, West Antarctica, *Reviews of Geophysics*, 55, 235–276, <https://doi.org/10.1002/2016RG000532>,  
10 2017.
- Van As, D., Fausto, R. S., Colgan, W. T., Box, J. E., Ahlstrøm, A. P., Andersen, S. B., Andersen, M. L., Charalampidis, C., Citterio, M., Edelvang, K., Jensen, T. S., Larsen, S. H., Machguth, H., Nielsen, S., Veicherts, M., and Weidick, A.: Darkening of the Greenland ice sheet due to the melt-albedo feedback observed at PROMICE weather stations, *Geological Survey of Denmark and Greenland Bulletin*, 28, 69–72, 2013.
- 15 Van den Broeke, M. R.: The semi-annual oscillation and Antarctic climate. Part 1: influence on near surface temperatures (1957–79), *Antarctic Science*, 10, 175–183, <https://doi.org/10.1017/S0954102098000248>, 1998.
- Van den Broeke, M. R., Reijmer, C. H., and Van de Wal, R. S. W.: Surface radiation balance in Antarctica as measured with automatic weather stations, *Journal of Geophysical Research*, 109, <https://doi.org/10.1029/2003JD004394>, 2004.
- Van den Broeke, M. R., Reijmer, C. H., Van As, D., Van de Wal, R. S. W., and Oerlemans, J.: Seasonal cycles of Antarctic surface energy  
20 balance from automatic weather stations, *Annals of Glaciology*, 41, 131–139, <https://doi.org/10.3189/172756405781813168>, 2005.
- Van den Broeke, M. R., König-Langlo, G., Picard, G., Kuipers Munneke, P., and Lenaerts, J. T. M.: Surface energy balance, melt and sublimation at Neumayer Station, East Antarctica, *Antarctic Science*, 22, 87–96, <https://doi.org/10.1017/S0954102009990538>, 2010.
- Van der Veen, C. J.: Fracture propagation as means of rapidly transferring surface meltwater to the base of glaciers, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2006GL028385>, 2007.
- 25 Van Wessem, J. M., Van de Berg, W. J., Noël, B. P. Y., Van Meijgaard, E., Birnbaum, G., Jakobs, C. L., Krüger, K., Lenaerts, J. T. M., Lhermitte, S., Ligtenberg, S. R. M., Medley, B., Reijmer, C. H., Van Tricht, K., Trusel, L. D., Van Uft, L. H., Wouters, B., Wuite, J., and Van den Broeke, M. R.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2, part 2: Antarctica (1979–2016)., *The Cryosphere*, 12, 1479–1498, <https://doi.org/10.5194/tc-2017-202>, 2018.
- Wiscombe, W. J. and Warren, S. G.: A Model for the Spectral Albedo of Snow. I: Pure Snow, *Journal of the Atmospheric Sciences*, 37,  
30 2712–2733, [https://doi.org/10.1175/1520-0469\(1980\)037<2712:AMFTSA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2712:AMFTSA>2.0.CO;2), 1980.
- Wouters, B., Martin-Español, A., Helm, V., Flament, T., Van Wessem, J. M., Ligtenberg, S. R. M., Van den Broeke, M. R., and Bamber, J. L.: Dynamic thinning of glaciers on the Southern Antarctic Peninsula, *Science*, 348, 899–903, <https://doi.org/10.1126/science.aaa5727>, 2015.




**Figure 1.** Map of the Antarctic continent, the red cross indicates the location of Neumayer Station. Imagery (C) 2016 DigitalGlobe, Inc.




**Figure 2.** (a) Downward longwave radiation versus air temperature. The red lines are quadratic fits of the upper and lower 5 percentile boundaries. The longwave-equivalent cloud cover is determined by linear interpolation between these bounds. (b) Optical thickness versus cloud cover. The red line resembles the best fit to a function  $\tau = c_1 (e^{c_2 N_{\epsilon}} - 1)$ , the shaded area indicates the 95% uncertainty range.

**Table 1.** Listing of used measurement variables and their associated measurement errors.


Variable	Neumayer errors
$V_{10m}$	$\max(0.5 \text{ m/s}, 5\%)$
$SW \downarrow$	$5 \text{ W/m}^2$
$SW \uparrow$	$5 \text{ W/m}^2$
$LW \downarrow$	$5 \text{ W/m}^2$
$LW \uparrow$	$5 \text{ W/m}^2$
$T_{2m}$	$0.1^\circ\text{C}$
$RH_{2m}$	$5\%$
$p$	$0.5 \text{ hPa}$



**Figure 3.** (a) Seasonal cycles of 2 m temperature (red, left axis), 10 m wind speed (green, right axis) and 2 m specific humidity (blue, right axis). Shaded areas indicate the standard deviations of monthly means. (b) Same as (a) for melt (red), net shortwave radiation (blue), net longwave radiation (orange), sensible heat (black), latent heat (magenta) and ground heat (green).



**Figure 4.** Daily values of modelled versus measured  $T_s$  for the parameter settings used in this study:  $z_{0,m} = 1.65$  mm,  $\rho_{s,0} = 280$  kg m $^{-3}$ .




**Figure 5.** Effect of model uncertainties on (a) cumulative melt and (b) seasonal melt. The shaded area indicates the  $1\sigma$  range due to model uncertainties (changing  $z_{0,m}$  and  $\rho_{s,0}$  between their respective values) which is asymmetrical because the values that are used for the rest of the study ( $z_{0,m} = 1.65$  mm,  $\rho_{s,0} = 280$  kg m $^{-3}$ ) are not in the middle of the range that was probed. The vertical grey patches in (a) indicate Nov–Feb of each year. Note that (b) ends earlier than (a) because the observations do not cover the 2015–16 melt season entirely.




**Table 2.** Mean annual values and interannual variability (calculated as standard deviations of monthly means) of meteorological variables and SEB components. For precipitation and melt, total annual values are given.


Variable	Yearly mean	Variability
$T_{2m}$ (K)	257.1	0.7
$T_s$ (K)	256.0	0.8
$q_{2m}$ (g kg <sup>-1</sup> )	1.1	0.1
$V_{10m}$ (m s <sup>-1</sup> )	8.9	0.6
$p$ (hPa)	981.6	2.0
$SW_{net}$ (W m <sup>-2</sup> )	20	2
$SW \downarrow$ (W m <sup>-2</sup> )	127	3
$SW \uparrow$ (W m <sup>-2</sup> )	107	2
$LW_{net}$ (W m <sup>-2</sup> )	-28	3
$LW \downarrow$ (W m <sup>-2</sup> )	218	5
$LW \uparrow$ (W m <sup>-2</sup> )	246	4
$Q_S$ (W m <sup>-2</sup> )	14.5	2.7
$Q_L$ (W m <sup>-2</sup> )	-6.3	1.2
$Q_G$ (W m <sup>-2</sup> )	0.7	0.4
$M$ (W m <sup>-2</sup> )	0.5	0.4
Precipitation (mm w.e.)	415	86
Melt (mm w.e.)	50	42




**Figure 6.** Average number of melt days per month at Neumayer. The inner error bars (with larger caps) indicate the  $1\sigma$  uncertainty range resulting from the runs performed with different settings for roughness length  $z_0$  and lower limit of new snow density  $\rho_{s,0}$  (Sect. 3.1). The outer error bars (with smaller caps) indicate the  $1\sigma$  range of the interannual variability.




**Figure 7.** Average values of some SEB components (a) and (b) and some meteorological variables (c) for December and January in the years with the highest (light grey) and lowest (dark grey) amount of melt, as identified in Sect. 3.3. Note that  $SW_{\downarrow}$ ,  $SW_{\uparrow}$ ,  $LW_{\downarrow}$  and  $LW_{\uparrow}$  are scaled by a factor of 10 in (a) for clarification.



**Figure 8.** Seasonal cycle of modelled average grain size in the upper 7 cm for the period 2000–2014. The grain size is expressed in terms of specific surface area ( $SSA = \frac{3}{\rho_i r_e}$ ) rather than grain size itself, to allow for a comparison with Fig. 6 of Libois et al. (2015).



**Figure 9.** Measured versus modelled daily average incoming shortwave radiation ( $SW_{\downarrow}$ ). The modelled  $SW_{\downarrow}$  was obtained by dividing the hourly measured  $SW_{\uparrow}$  by the calculated hourly albedo.



**Figure 10.** (a) Timeseries of cumulative amount of melt for the run with measured albedo ( $R_0$ , blue), a constant albedo of 0.84 ( $R_1$ , red), a run in which refrozen snow does impact snow grain size ( $R_2$ , yellow) and a run in which snow grain size is not influenced by refrozen snow ( $R_3$ , purple). (b) Same as (a) but for seasonal amount of melt. (c) Ratio of modeled surface melt between yellow and purple lines in (a) and (b) (runs  $R_2$  and  $R_3$  respectively). The grey area indicates the uncertainty coming from the uncertainty in the determination of  $\tau$  (Fig. 2b)),  $\pm 5 \text{ W m}^{-2}$  measurement uncertainty in  $SW \uparrow$  and the inclusion of shortwave radiation penetration.