

Interactive comment on “Simulated single-layer forest canopies delay Northern Hemisphere snowmelt” **by Markus Todt et al.**

Markus Todt et al.

markus.todt@northumbria.ac.uk

Received and published: 20 June 2019

This document contains comments by reviewers (regular font), **responses to reviewers (bold)**, and *citations from the manuscript (italic)* including [additions to the manuscript \(blue\)](#).

Reviewer 2 Comments:

Technically, the demonstration is not thoroughly made, that single-layer canopy formu-

Printer-friendly version

Discussion paper



lations generate melt delay in the NH with respect to "real world" (that here would be observations). Indeed, the demonstration of this effect is only made with respect to an approximation of a 2-layer model, that may itself be heavily biased. This is all the more worrying as Todt et al., 2018, illustrate an increase in a subcanopy LW positive bias upon the use of a 2-layer model at the Seehornwald conifer site (Todt et al., 2018, Figure 5).

While the two-layer canopy model SNOWPACK did show a positive bias at the Seehornwald site, it also showed a substantial negative bias at the site of Sodankylä. These biases stem from SNOWPACK being calibrated for the site of Alptal, which features a lower vegetation density than Seehornwald but a larger vegetation density than Sodankylä. Consistent for those three sites was the substantially smaller spread in and diurnal cycle of sub-canopy longwave radiation shown by Todt et al. (2018). Furthermore, Gouttevin et al. (2015) showed that the improvement from one layer to two layers did result in a reduction of the negative bias that SNOWPACK displayed for Alptal, although this did not (just) originate from longer nights than days but (also) from larger nighttime underestimations than daytime overestimations. But it is true that the delay in meltout found between global simulations had not been shown relative to observations, instead had been inferred as a consequence from comparison of sub-canopy longwave radiation with observations at forest-stand scales. Evaluation of global simulations has been added as described in the next paragraph.

The melt delay associated with 1-layer canopy models claimed by the authors, may clearly be real, but the demonstration should be improved, for instance by confrontation of simulation results to (i) in-situ data at field sites and (ii) satellite estimate of NHsnow disappearance dates. Comparison of simulation results to observed snow-cover fractions is briefly mentioned in the Discussion while it should be an important

[Printer-friendly version](#)[Discussion paper](#)

part of the Result section (as it is already quite well advertised as a rationale for this study in the Introduction). Thackeray et al., 2015 (their Figure 3 for instance) provides a good baseline for such comparisons.

We do understand and acknowledge the value of a comparison to observed snow-off dates, and we have included comparison of meltout between simulations and a state-of-the-art snow water equivalent product in this revision. This comparison revealed a general delay of snow-off dates across boreal forests in simulations, and correction of sub-canopy longwave radiation was found to decrease this bias. However, comparison at Toy Model sites is challenging and potentially inconclusive as snow measurements are largely unavailable. Various approximations would have to be used for comparison as was done for driving the Toy Model by Todt et al. (2018), and the use of this might be limited. Therefore, we restricted evaluation to global snowmelt.

A blended data set of five global observation-based SWE products (henceforth, Blended-5) covering the period 1981 to 2010 (Mudryk et al., 2015) was used to estimate snow-off dates across the Northern Hemisphere and evaluate simulation of snowmelt in CTRL and CORR. In contrast to simulations, observations display snow persisting for physically unrealistical durations, which necessitates a SWE threshold to estimate snow-off dates (Krinner et al., 2018). While Mudryk et al. (2017) and Krinner et al. (2018) used thresholds of 4mm and 5mm, respectively, for estimates of spatial snow cover extent, a smaller SWE value was necessary to represent the precise timing of meltout within individual grid cells. A threshold of 1mm was used in this study to define meltout for the Blended-5 mean, and snow-off date was defined as the first day of a year for which SWE did not exceed this threshold. Sensitivity of snow-off dates to threshold values was tested for the range 0.5mm to 4mm, however, the overall conclusions of this study are unchanged for different thresholds.

Simulated and observed snow-off dates are compared in Fig. 10 for grid cells with

[Interactive
comment](#)

[Printer-friendly version](#)

[Discussion paper](#)



consistent snow cover throughout preceding December and coverage by evergreen needleleaf trees of at least 50%. Simulations CTRL and CORR generally feature a narrower probability density function (PDF) of snow-off dates, indicating a shorter snowmelt season, and later meltout compared to observations across the entire Northern Hemisphere (Fig. 10a). While shapes of observed PDFs are well represented by simulations over Eurasia (Fig. 10b, d), observations show a clearer, shorter peak of meltout compared to simulations over mountainous western North America (Fig. 10c). Correction of sub-canopy longwave radiation displays little impact when accumulated over the entire Northern Hemisphere, however, it systematically reduces the delay of simulated snow-off dates throughout the snowmelt season. PDFs of snow-off dates for regional subsets reflect spatial patterns seen in Fig. 7h, with minor differences between CTRL and CORR over most of western North America (Fig. 10c) and eastern Siberia (Fig. 10d) but substantial acceleration of snow-off dates over western Siberia and eastern Europe (Fig. 10b) due to correction of sub-canopy longwave radiation.

The regionally limited impact of corrected sub-canopy longwave radiation is highlighted by filtering PDFs of snow-off date for grid cells with average differences in snow-off date between CORR and CTRL of at least 3 days (Fig. 10e, f). Correction of sub-canopy longwave radiation improves timing of meltout in filtered grid cells, especially over western Siberia and eastern Europe where the filtered PDF for CORR, in contrast to CTRL, closely resembles observations. PDFs of snow-off dates derived from Blended-5 SWE display sensitivity to threshold choices, however, this uncertainty is generally smaller than differences between simulations and observations.

Secondly, for the evaluation of the delay effect against in-situ data, simulation errors coming from the meteorological forcing should be minimized. Evergreen forest sites used in Todt et al., 2018, provide appropriate observed meteorological data. They could be used in the place of erroneous large-scale meteorological forcing data (for e.g. Figure 4 and the associated results analysis).

As mentioned in the previous comment, snow cover measurements at Toy Model sites are either available only by approximation or not at all. Replicating Figure 4 and its analysis with stand-scale simulations is possible, and the impact of correction on sub-canopy longwave radiation indeed leads to improved simulations. However, this should be expected as observations of sub-canopy longwave radiation from these forest stands are used to create the correction. Therefore, we test and train correction of sub-canopy longwave radiation with the same dataset, and this comparison on forest-stand scales is a further illustration rather than an evaluation, which is why we decided against its inclusion in our manuscript.

p1L10 : the last sentence of the abstract associates "boreal forests" and "warm winters" where "snowmelt occurs early". This is not very intuitive

"Warm winters" are meant to be relative to regions where there is snow. The chain of causation is warmer winters → earlier snowmelt → snowmelt when nights are longer than days → substantial underestimation of sub-canopy longwave radiation → delay of snowmelt. We deleted the part about warmer winters as early snowmelt already indicates that these regions are warmer than other snow-covered regions.

Increasing insolation and day length change the impact of overestimated diurnal cycles on daily average sub-canopy longwave radiation throughout the snowmelt season. Consequently, delay of snowmelt in land-only simulations is more substantial where snowmelt occurs early.

p2L3 : cite Krinner et al., 2018

[Printer-friendly version](#)[Discussion paper](#)

We thank the reviewer for pointing out this paper. However, we feel that it does not relate closely to the material being discussed in that section because, while ESM-SnowMIP will include an experiment to investigate the impact of vegetation distribution, the rationale of Krinner et al. (2018) and its references are focused on surface albedo. Also, modification of vegetation distribution will not have an impact on the control of vegetation density on vegetation temperatures. We therefore elect to retain the original text.

p3L9-12 : "emissivity" is inappropriately used. What the authors call "emissivity", is an unexplained combination of emissivity and sky view factor. Please explicit and justify the approximations that you make here.

Yes, the authors absolutely agree that the parameter "vegetation emissivity" used by CLM4.5 is not an emissivity in the physical sense, i.e. as an emissivity is commonly used in the Stefan-Boltzmann equation. However, this is the nomenclature used by CLM4.5, which we decided to stick to in order for consistency with CLM4.5, and the technical description of CLM4.5 (Oleson et al., 2013) does not give any reasoning for or description of the combination of actual physical emissivity and SVF/canopy coverage.

p3L9-12 : please explain briefly how T_v is calculated.

A short description of the calculation of T_v by CLM4.5 has been added.

...using the Stefan-Boltzmann law with Stefan-Boltzmann constant $\sigma = 5.67 \cdot 10^{-8} \text{ W m}^{-2} \text{ K}^{-4}$ and vegetation temperature T_v . [Vegetation temperature is calculated based on an energy balance, net radiation minus turbulent heat fluxes. Radiative trans-](#)

[Printer-friendly version](#)[Discussion paper](#)

fer of direct and diffuse shortwave radiation is calculated via a two-stream approximation (Sellers, 1985) considering one reflection from ground to canopy. Net longwave radiation is calculated from atmospheric longwave radiation, vegetation temperature, and (ground) surface temperature and determined by vegetation emissivity and emissivity of the ground. Calculation of turbulent heat fluxes in CLM4.5 is based on Monin-Obukhov similarity theory and described by Oleson et al. (2013). Vegetation emissivity depends on Leaf Area Index (LAI) and Stem Area Index (SAI) and is calculated as...

P6L11-13 : In the current structure of the paper, calculating and exhibiting correction factors for deciduous forest regions makes in my opinion little sense, as no use is made of them, and very little analysis of their difference w/r to coefficients calculated for evergreen forests is made. If Cherskii's coefficients are similar to those from evergreen sites, then what is the added value of including this site in the calibration intended for evergreen sites ?! To me, it just undermines the calibration approach. Calculating correction coefficients for deciduous forests stands may be interesting, though, providing their difference (w/r evergreen sites) and impact (on e.g. snowmelt) is discussed in the paper.

As this paper is a continuation of Todt et al. (2018), we included all sites from that paper that could be used for multiple linear regression. The reasons why sub-canopy longwave radiation is only corrected for evergreen trees and why Cherskiy is included in calculating this correction are explained in the paper. Simulations for Cherskiy are similar to evergreen trees while Abisko is not reliable/sufficient. This similarity is then used to balance very dense (Alptal, Seehornwald) and sparser sites (Cherskiy, Sodankylä) for a more general applicability. Similarities and differences between regression coefficients for Abisko, Cherskiy, and evergreen forest stands are indeed interesting and open up a separate discussion about the influence of stand characteristics and structure on

[Printer-friendly version](#)[Discussion paper](#)

sub-canopy radiative fluxes. For the sake of clarity and to keep the focus of our paper on the impact of corrected sub-canopy longwave radiation, we (had) decided against including that discussion.

p6L20 : please add 'CLM4.5' before 'grid cell' for more clarity

Has been added. Thank you for this suggestion.

For the location of Alptal, in contrast to other forest stands used in this study, forest stand and CLM4.5 [grid cell](#) feature similarly high vegetation densities (PAIs of $4.1 \text{ m}^2 \text{ m}^{-2}$ and $3.7 \text{ m}^2 \text{ m}^{-2}$, respectively) and thus similar vegetation emissivities ϵ_v (0.983 and 0.975, respectively). This allows for a comparison of diurnal cycles of sub-canopy longwave radiation as well as longwave enhancement between stand-scale measurements and offline simulations.

P6paragraph4.1 : I suggest to carry this analysis using an observed forcing to evidence the effect of going from 1-layer to 2-layer w/r to observations in an "ideal" case. Also, it should be mentioned more clearly in the text that the selected period corresponds to the snowmelt season at Alptal and hence is relevant to assess the effect of subcanopy LW on snowmelt.

This analysis is done to highlight the (successful) effect on diurnal cycles of sub-canopy longwave radiation rather than as an evaluation. The Toy Model also does not include all calculations/parameterizations of CLM4.5, only those necessary to simulate sub-canopy longwave radiation while offline simulations model the entire land surface, so the impact on sub-canopy longwave radiation simulated by the Toy Model does not tell the entire story. Caption of Figure 4

[Interactive comment](#)

[Printer-friendly version](#)

[Discussion paper](#)



mentions that the snowmelt season at Alptal is shown, and this is added to the text.

Implementation of correction factors in CLM4.5 results in decreased sub-canopy long-wave radiation during daytime and increased sub-canopy longwave radiation during nighttime, thereby reducing diurnal cycles. For the grid cell representing Alptal, diurnal ranges decrease from about 70 W m^{-2} to about 30 W m^{-2} during snowmelt season (Fig. 4a and Fig. 4b). Observations at the forest stand show an average diurnal range of about 15 W m^{-2} during snowmelt season.

P7Paragraph4.2 : Maybe other forcings than CRUNCEP, exhibit less bias with respect to spatio-temporal in sky emissivity. Brun et al., 2013, concluded that ERA-i generally leads to improved simulations (w/r to other forcings) over large areas of the N high latitudes.

Only two datasets are options in CLM4.5 for forcing of offline simulations – the CRUNCEP dataset used in this study and a dataset described by Qian et al. (2006). Snow cover in CLM4.5 offline simulations driven with both of these datasets have been analyzed by Thackeray et al. (2015), revealing higher skill scores for the simulation driven by the CRUNCEP dataset. Because of this and because of shorter coverage by the Qian dataset, we decided to use and stick to CRUNCEP forcing data.

References:

Qian, T., A. Dai, K. E. Trenberth, and K. W. Oleson (2006), Simulation of global land surface conditions from 1948 to 2004. Part I: Forcing data and evaluations, J. Hydrometeorol., 7(5), 953–975, doi:10.1175/JHM540.1.

Thackeray, C.W., C. G. Fletcher, and C. Derksen (2015), Quantifying the

[Printer-friendly version](#)[Discussion paper](#)

skill of CMIP5 models in simulating seasonal albedo and snow cover evolution, *Journal of Geophysical Research: Atmospheres*, 120, 5831–5849, doi:10.1002/2015JD023325.

P8L3 : I suggest to outline the regions with $\text{frac_PFT} > 0.5$ on these maps, for better understanding of the effects of CORR and their magnitude.

A contour line for fractional coverage by evergreen needleleaf trees of at least 50% has been added. Thank you for this suggestion.

P8L22: maybe the glaciated areas should be masked out as they are not the focus of this study. Otherwise, the question arises as to whether cold content for these areas refers to the snow, or to the whole snow+ice columns.

The reviewer is correct that cold content in glaciated regions includes ice, which is why explicit values for these regions are not mentioned in the text and the colorbar is cropped at 5 MJ m^{-2} . The choice to include these grid cells was made for visual reasons, as many grid cells that include the landunit "glaciated" are entirely glaciated and would thus appear as empty on maps. Therefore, we decided to take no action.

P8L30 : this is not true to my understanding as melt out date is largely determined by the energy required to melt the snow (which is often higher than the one needed to raise snow temperature to 0°C)

We agree with the reviewer and the statement made at that point is not intended

[Printer-friendly version](#)[Discussion paper](#)

to suggest that spatial differences in cold content solely cause the spatial pattern in meltout. However, there is a clear spatial correlation between these two variables, especially between changes in meltout and realtive changes in cold content, and more and/or colder snow inevitably requires a higher energy input for melt. Which is why we decided to take no action.

P9L1 : maybe the delay between melt-out-date, and equinox, could be an interesting additionnal explanatory variable in Fig8. Also, an illustration of daily cycle changes before and after equinox for Southeastern regions would be a great complement to the explanations.

The difference between equinox and melt-out date is indeed one of the major governing variables of the impact of corrected sub-canopy longwave radiation. However, grid cells across Siberia generally feature meltout past the equinox, and location (latitude/insolation, elevation) appears to be the most important characteristic for this region, which is why we decided not to add a panel for difference to equinox. Difference to equinox is a more helpful metric when assessing the impact on snowmelt across regions with starker contrasts in melt-out date, e.g. Europe.

The illustration of changes in diurnal cycles only has limited explanatory value, as daytime overestimations and nighttime underestimations are fairly consistant and changes in day length are not overly striking when visualized as a diurnal cycle. In fact, Figure 4b already features diurnal cycles of sub-canopy longwave radiation prior to the boreal spring equinox and post-equinox diurnal cycles would only appear slightly different. However, we included a figure showing Northern Hemisphere maps of changes in longwave enhancement due to correction before and after the equinox (see last comment in this review).

[Printer-friendly version](#)[Discussion paper](#)

P9L15 : specify "over the study region"

The particular statement holds generally and should not be limited to our study region, as the governing factors are day length, snow cover, and presence of evergreen needleleaf trees. Therefore, we decided to stick to the original phrasing.

P9L30 : Illustrations of comparisons to observations like in Thackeray et al. 2014, 2015, is exactly what is missing in your study, and would add great value to it.

We have included comparison of meltout between simulations and a state-of-the-art snow water equivalent product in this revision, which revealed a general delay of snow-off dates across boreal forests in simulations. Correction of sub-canopy longwave radiation was found to decrease this bias. Detailed additions to the manuscript are listed underneath the first comment of this review.

Overall, there is a lack of proper illustration of the competing effects of CORR on the daily subcanopy LW radiation before vs after the equinox, and how this governs the effects of CORR at the global scale.

We added a figure showing changes in longwave enhancement by evergreen needleleaf trees across the Northern Hemisphere before and after the boreal spring equinox, which show contrasting signs in the impact of corrected sub-canopy longwave radiation.

As offline simulations lack spatial variability in ϵ_{sky} , latitude (through insolation) and duration of snow on the ground (through day length) control spatial differences in impact of correction of sub-canopy longwave radiation on snow-off date. Changes in long-

Printer-friendly version

Discussion paper



wave enhancement due to correction of sub-canopy longwave radiation before and after the boreal spring equinox, approximated by averages over February/March and April/May, display opposite signs across the Northern Hemisphere (Fig. 8), with shorter (longer) days than nights before (after) the equinox resulting in an increase (decrease) in daily average longwave enhancement. Generally, lower insolation at higher latitudes leads to a more positive impact of correction on daily average longwave enhancement, increasing (decreasing) positive (negative) changes in longwave enhancement with increasing latitude before (after) the boreal spring equinox. Across mid-latitudes, increase in daily average longwave enhancement over February and March is roughly similar to decrease in daily average longwave enhancement over April and May, while increase over February and March outweighs decrease over April and May across high latitudes including most of the regions covered by boreal forests.

Interactive comment on The Cryosphere Discuss., <https://doi.org/10.5194/tc-2018-270>, 2019.

Printer-friendly version

Discussion paper

