

Supplement of

Recent changes in pan-Antarctic surface snowmelt detected by AMSR-E and AMSR2

Lei Zheng et al.

Correspondence to: Chunxia Zhou (zhoucx@whu.edu.cn) and Tingjun Zhang (tjzhang@lzu.edu.cn)

Content

Figure 1

Figure 2




Figure 1. Comparison between AMSR-E vertically polarized 36.5 GHz Tb (Tb_{V36}) and SSM/I vertically polarized 37 GHz Tb (Tb_{V37}) during 2002-2003. (a), (b) and (c) show the comparisons for ascending passes, descending passes, and their differences. DAV37 and DAV36 denote diurnal amplitude variations (DAV) of vertically polarized SSM/I 37 GHz Tb and AMSR-E 36.5 GHz Tb, respectively.




Figure 2. Surface snowmelt detection on the Antarctic sea ice. (a) Pan-Antarctic ice cover days in 2002-2003, Point B and C show the locations of the pixels examined in (b) and (c). (b) and (c) show the comparisons of sea ice concentration (SIC), ERA-Interim T_{air} and satellite observations for a multi-year sea ice pixel (Point B) and a first-year sea ice pixel (Point C). DAV_{37ice} and DAV_{36ice} denote diurnal amplitude variations (DAV) of vertically polarized SSM/I 37 GHz Tb and AMSR-E 36.5 GHz Tb contributed by the ice portion, respectively.