

Response to reviewers:
On the suitability of the Thorpe-Mason model for
calculating sublimation of saltating snow

Varun Sharma, Francesco Comola and Michael Lehning

October 3, 2018

1 Response to Reviewer # 1

We thank Reviewer # 1 for his/her comments on the revised manuscript and our answers to the questions raised in the previous round of review. We address the remaining concerns below.

- ***A.1 : Response A.14: It is unclear whether this is the complete answer to the comment or not. Please clarify if this is a typo.***

Response A.1: The response to A.14 was indeed a typo. The question and the intended answer are reproduced below.

A.14 : P6 L5: These varying friction velocities are referred to as “low medium and high wind speeds” in line 33. What wind speeds were necessary for these values? Friction velocities do a poor job of representing turbulence in even subtly complex terrain, and as saltation is a drag-driven process, at the very least, mean wind speeds should be included in the manuscript, and extensive time series of turbulence statistics (Turbulence Intensity, TKE, shear stress, etc.) in the supplementary material. As this research is conducted to benefit those that models in natural settings, and those natural setting will be much more turbulent than the LES, and that turbulence is what is driving the ventilation rates, more information about the model is needed.

Response A.14: In the revised manuscript, wind speeds at the height of 1 m above the surface are mentioned for each the different u_* cases. We agree that this information is valuable and deserves to be added. Some turbulent statistics are presented in the revised supplement.

-
- ***A.2 : Response A.21: The authors responded to a question of the impact of surface sublimation on suppression of vertical motions. However, the question posed was actually on the potential presence of a temperature gradient in the atmosphere as could be found in a typical stable boundary layer in calm wind over snow. However, as is mentioned below, it is suggested that these comments on vertical suppression just be removed altogether. It is unclear whether it was intended to imply that there is no such temperature gradient in nature with turbulent mixing, or that such a temperature gradient only exists***

if there is actively sublimating surface snow?

Response A.2: We agree with the suggestion that comments on vertical suppression are removed from the main portion of the manuscript.

- *A.3 Author Response Page 4, First paragraph: with the air is so close Please correct.*

Response A.3: Corrected!

- *A.4 Author Response Page 4, Last Paragraph lighther.*

Response A.4: Corrected!

- *A.5 Page 5 Line 3: Please add citation that these values are typical for saltation.*

Response A.5: Some citations have been added. Note that since the work of Thorpe and Mason, not many studies explicitly talk about Nusselt and Sherwood numbers for ice grains.

- *A.6 P6 L15-16: Do you mean or*

Response A.6: Yes, the sentence has been corrected.

- *A.7 Section 3.1 The use of LES is very attractive to many people in hydrology and snow science, but the complexity of such an endeavor is often the limiting factor. Please explicitly cite the software used, be it in-house code or a modified commercial product? It is appreciated that the governing equations of motion are included, but for a model of this nature, a note on where the code was actually developed (or ideally, available) would be beneficial to the community. As well, please note in the main body of text that is LES of a channel flow.*

Response A.7: The code has been developed completely in-house over a period of 20 years with multiple dissertations contributing to its development and applications. We note this information in the text. Additionally, it is explicitly stated that the LES is indeed of a channel flow.

-
- *A.8 P8L22: to decelerate*

Response A.8: Thank you, it stands corrected!

-
- *A.9 P10L7: Fix units*

Response A.9: Thank you, it stands corrected!

-
- *A.10 P10L8: “Mean wind speeds.” It may be stationary, but it is still a turbulent flow.*

Response A.10: Thank you, the sentence is now modified.

-
- *A.11 P10L13-17: This doesnt seem to be supported by the supplement. It is not clear that there was vertical suppression because of sublimation, only vertical suppression during saltation. This claim seems like it would need further investigation, for example a comparison with saltation without sublimation to show the actual impact of the cooling. Obviously TKE drops, but that is insufficient to show SUPPRESSION of vertical motions by thermal effects. The buoyancy is a function of vertical motions, so that is a bit of a circular argument. I suggest removing comment of any causal relationship between sublimation and vertical motion suppression until the referenced future research is conducted.*

Response A.11: Similar to the answer to point A.2, we remove any statements linking vertical motions and sublimation etc.

With regards to our discussion here, we would like to state that the vertical profiles in the supplement are for simulations with the same dynamical forcing and thus, the number of particles in saltation. The only difference between the three cases is that they have different initial conditions of relative humidity. In other words, the principle difference between the three profiles is the varying amounts of sublimation. Given the initial condition of uniform relative humidity, the buoyancy comes only from the sublimation (cooling of the air) due to particles and can be thought of as a cause. The effect is the vertical motions - if we write the Reynolds averaged equation for the vertical velocity fluctuation, we can see that all other terms being equal, the vertical motions are directly related to the buoyancy forcing - thus, more sublimation, more negative buoyancy and thus reduced vertical motions.

We thank the reviewer for his/her suggestion for simulations of saltation with and without sublimation. This will address the link between sublimation and residence times definitively. This will indeed be a part of our future efforts.

-
- *A.12 Figure 1: The 40% reduction in windspeed with the presence of saltation is substantial and should be commented on. There are numerous wind tunnel (and some field) studies addressing momentum extraction from the wind by particles in saltation. This 40% should be put in context of previous work to add credibility to an inherently complex model of a simple scenario to be found in nature.*

Response A.12: Citations of some papers discussing the velocity reduction have been added. Note that for most studies, the wind velocities, whether in the field or in wind tunnels are either lower than our UM case.

-
- *A.13 It is claimed in the manuscript that turbulent mixing of different temperatures is responsible for the regeneration of warm dry air that then results in different rates of sublimation for LES. It would be illuminating to have a movie showing an animation of evidence of that process, as it is the heart of the conclusions presented here.*

Response A.13: An additional supplementary movie M2 is now added to the supplementary material. In M2, we show the evolution of the relative humidity instead of simply temperature. It can be clearly seen in the video, how eddies transport air with low relative humidity from aloft closer to the surface.

2 Response to Reviewer # 2

We thank Reviewer # 2 for agreeing for the manuscript to proceed for publication. We once again thank the Reviewer for his/her comments and suggestions during the review process that has undoubtedly improved the quality of the paper.

On the suitability of the Thorpe-Mason model for Calculating Sublimation of Saltating Snow

Varun Sharma¹, Francesco Comola¹, and Michael Lehning^{1,2}

¹School of Architecture, Civil and Environmental Engineering, Swiss Federal Institute of Technology, Lausanne, Switzerland

²WSL Institute for Snow and Avalanche Research SLF, Davos, Switzerland

Correspondence: Varun Sharma (varun.sharma@epfl.ch)

Abstract. The Thorpe and Mason (TM) model for calculating the mass lost from a sublimating snow grain is the basis of all existing small and large-scale estimates of drifting snow sublimation and the associated snow mass balance of polar and alpine regions. We revisit this model to test its validity for calculating sublimation from saltating snow grains. It is shown that numerical solutions of the unsteady mass and heat balance equations of an individual snow grain reconcile well with the steady-state solution of the TM model, albeit after a transient regime. Using large-eddy simulations (LES), it is found that the residence time of a typical saltating particle is shorter than the period of the transient regime, implying that using the steady state solution might be erroneous. For scenarios with equal initial air and particle temperatures of 263.15 K, these errors range from 26% for low-wind low saturation-rate conditions to 38% for high-wind high saturation-rate conditions. With a small temperature difference of 1 K between the air and the snow particles, the errors due to the TM model are already as high as 100% with errors increasing for larger temperature differences.

Copyright statement. TEXT

1 Introduction

Sublimation of drifting and blowing snow has been recognized as an important component of the mass budget of polar and alpine regions (Liston and Sturm, 2004; van den Broeke et al., 2006; Lenaerts et al., 2012; Vionnet et al., 2014). Field observations and modeling efforts focused on Antarctica have highlighted the fact that precipitation and sublimation losses are the dominant terms of the mass budget in the katabatic flow region as well as the coastal plains (van den Broeke et al., 2006). Even though precipitation is challenging to measure accurately, methods to measure it exist, for example, using radar (Grazioli et al., 2017) or snow depth change (Vögeli et al., 2016). In comparison, sublimation losses are even harder to measure and can only be calculated implicitly; using measurements of wind speed, temperature and humidity. Thus, in regions where sublimation loss is a dominant term of the mass balance, it is also a major source of error. This error ultimately results in errors in the mass accumulation of ice on Antarctica, which is a crucial quantity for understanding sea-level rise and climate change (Rémy and Frezzotti, 2006; Rignot et al., 2011; Lenaerts et al., 2012).

Aeolian transport of snow can be classified into three modes, namely, creeping, saltation and suspension. Creeping consists of heavy particles rolling and sliding along the surface of the snowpack either due to form drag or bombardment due to impacting particles. Saltation consists of particles being transported along the surface via short, ballistic trajectories with heights mostly less than 10 cm and involves mechanisms of aerodynamic entrainment along with rebound and splashing of ice grains (Doorschot and Lehning, 2002; Comola and Lehning, 2017). Suspension on the other hand refers to transport of small ice grains at higher elevations and over large distances without contact with the surface. Current calculations of sublimation losses are largely restricted to losses from ice grains in suspension. This is true for both field studies (Mann et al., 2000), where sublimation losses are calculated using measurements, usually at the height of $\mathcal{O}(1\text{m})$, and in mesoscale modeling studies (Xiao et al., 2000; Déry and Yau, 2002; Groot Zwaaftink et al., 2011; Vionnet et al., 2014), where the computational grids and time-steps are too large to resolve flow dynamics at saltation length and time scales. Mass loss in the saltation layer is hard to measure and is neglected based on the justification that the saltation layer is saturated. However, recent studies using high-resolution steady-flow, Reynolds-averaged Navier-Stokes (RANS) type simulations (Dai and Huang, 2014) claim that sublimation losses in the saltation layer are not negligible, particularly for wind speeds close to the threshold velocities for aeolian transport, wherein a majority of aeolian snow transport occurs via saltation rather than suspension.

The coupled heat and mass balance equations of a single ice particle immersed in turbulent flow are

$$c_i m_p \frac{dT_p}{dt} = L_s \frac{dm_p}{dt} + \pi \mathcal{K} d_p (T_{a,\infty} - T_p) \mathcal{N}u, \quad (1)$$

$$\frac{dm_p}{dt} = \pi \mathcal{D} d_p (\rho_{w,\infty} - \rho_{w,p}) \mathcal{S}h, \quad (2)$$

where, m_p , T_p and d_p are the mass, temperature and diameter of the particle respectively that vary with time, c_i is the specific heat capacity of ice, L_s is the latent heat of sublimation, \mathcal{K} is the thermal conductivity of moist air and \mathcal{D} is the mass diffusivity of water vapor in air. Transfer of heat and mass is driven by differences of temperature and vapor density between the particle surface ($T_p, \rho_{w,p}$) and the surrounding fluid ($T_{a,\infty}, \rho_{w,\infty}$). The vapor density at the surface of the ice particle is considered to be the saturation vapor density for the particle temperature. The transfer mechanisms are enhanced by turbulence, the effect of which is parameterized by the Nusselt ($\mathcal{N}u$) and Sherwood ($\mathcal{S}h$) numbers respectively. $\mathcal{N}u$ and $\mathcal{S}h$ are related to the relative speed ($|\mathbf{u}_{rel}|$) between the air and the particle via the particle Reynolds number (Re_p) as

$$Re_p = \frac{d|\mathbf{u}_{rel}|}{\nu_{air}}; \mathcal{N}u = 1.79 + 0.606 Re_p^{1/2} Pr^{1/3}; \mathcal{S}h = 1.79 + 0.606 Re_p^{1/2} Sc^{1/3}, \quad (3)$$

where ν_{air} is the kinematic viscosity of air and Pr and Sc are the Prandtl and Schmidt numbers respectively.

Thorpe and Mason (1966) solved the above coupled Eqs. (1) and (2) by, (a) neglecting the thermal inertia of the ice particle, thus effectively stating that all the heat necessary for sublimation is supplied by the air, and (b) considering the temperature difference between the particle and surrounding air to be *small*, thereby allowing for Taylor series expansion of the Clausius-Clapeyron equation and neglecting higher-order terms, resulting in their formulation for the mass loss term as,

$$\frac{dm_p}{dt} = \pi d_p (\sigma_* - 1) \left/ \left(\frac{L_s}{\mathcal{K} T_{a,\infty} \mathcal{N}u} \left(\frac{L_s M}{R T_{a,\infty}} - 1 \right) + \frac{1}{\mathcal{D} \rho_s(T_{a,\infty}) \mathcal{S}h} \right) \right., \quad (4)$$

where $\rho_s(T_{a,\infty})$ is the saturation vapor density of air surrounding the particle, saturation-rate $\sigma_* = \rho_{w,\infty}/\rho_s(T_{a,\infty})$, M is the molecular weight of water and R is the universal gas constant. The above formulation has been used extensively to analyze data collected in the field (Mann et al., 2000), wind tunnel experiments (Wever et al., 2009), and numerical simulations of drifting and blowing snow (Déry and Yau, 2002; Groot Zwaaftink et al., 2011; Vionnet et al., 2014). In the modeling studies, the mass loss term is computed using Eq. (4) and is added, with proper normalization, to the advection-diffusion equation of specific humidity while the latent heat of sublimation multiplied by the mass loss term is added to the corresponding equation for temperature (Groot Zwaaftink et al., 2011).

Two observations motivated us to investigate the suitability of the TM model for sublimation of saltating snow particles. Firstly, the TM model assumes that all the energy required for sublimation is supplied by the air. This assumption was tested by Dover (1993) who compared the potential rates of cooling of particles with that of the surrounding air due to sublimation. Using scale analysis, Dover (1993) formulated the quantity $\xi = 6\rho_{air}c_{p,air} / \pi\rho_i c_i \bar{d}_p^3 N$, where \bar{d}_p is the mean particle diameter, N is the particle number density, ρ_i is the density of ice, and showed that for $\xi \gg 1$, it can be accurately considered that the heat necessary for sublimation comes from the air. For standard values for an ice particle in suspension, $\bar{d}_p = 50 \mu\text{m}$ and $N \sim \mathcal{O}(10^6)$, this condition is easily met ($\xi \sim \mathcal{O}(10^3)$). However, if we input values typical for saltation, i.e., $\bar{d}_p = 200 \mu\text{m}$ and $N \sim \mathcal{O}(10^8)$, $\xi \sim \mathcal{O}(1)$, and the condition is not met. Thus, for sublimation of saltating particles, it is important to consider the thermal inertia of the particles. A similar conclusion was reached in other modeling studies on topics of heat and mass exchange between disperse particulate matter in turbulent flow such as small water droplets in heat exchangers (Russo et al., 2014) and sea-sprays (Helgans and Richter, 2016).

Secondly, Eq. (4) computes mass loss as being directly proportional to σ_* and neglects the temperature difference between the particle and air. Eq. (4) thus predicts a mass loss even in extremely high saturation-rate conditions, whereas immediate deposition of water vapor would occur on a particle even slightly colder than the air. Indeed, some field experiments have reported deposition as opposed to sublimation which was expected, on the basis of the measured under-saturation of the environment, particularly near coastal polar regions (Sturm et al., 2002). A simple everyday observation illustrates this fact clearly; There is immediate deposition of vapor and formation of small droplets on the surface of a cold bottle of beer even in room conditions with moderate humidity!

Motivated by the observations described above, in this article, we describe four numerical experiments where we compare differences between the fully numerical and the Thorpe and Mason (1966) solutions (referred to as NUM and TM approaches respectively). In Experiment I and II, we numerically solve Eqs. (1) and (2) and compare the results with Eq. (4) for physically plausible values of a saltating ice particle. Results of these tests are presented in Sect. 2. High-resolution large-eddy simulations (LES) of the atmospheric surface layer with saltating snow are performed for a range of environmental conditions to compute the differences between the NUM and TM approaches in realistic wind-driven saltating events. These results are presented in Sect. 3. A summary of the article is made in Sect. 4

2 Comparison between NUM and TM solutions: EXPERIMENT I and II

We consider an idealized scenario where a solitary spherical ice particle is held still in a turbulent air flow with constant mean speed, temperature and under-saturation. The evolution of the mass, diameter and temperature of the ice particle is calculated using both the NUM and TM models and an inter-comparison is made. This scenario is similar to the wind-tunnel study performed by Thorpe and Mason (1966) who measured mass loss of solitary ice grains suspended on fine fibers. In this scenario, we consider that the heat and mass transfer between the ice particle and the air changes the mass and temperature of the particle only while the mass and energy anomalies in the air are rapidly advected and mixed away. This implies that the environmental conditions subjected to the ice particle remain constant. While it can be expected that the environmental conditions will vary along the trajectory of a ice particle undergoing saltation or suspension, it is nevertheless useful to perform this analysis as it reveals important characteristics about the heat and mass evolution of a ice particle during sublimation and about the approximations used to derive the TM model.

For the NUM approach, Eqs. (1) and (2) are solved in a coupled manner using a simple first-order finite-differencing scheme for time-stepping with a time-step of $50 \mu\text{s}$. For the TM approach, Eq. (4) is used with a similar numerical setup as for the NUM approach. In the TM approach, particle temperature is not considered and the mass and energy transfer is determined only by air temperature and saturation-rate. The initial particle diameter ($d_{p,IC}$) is $200 \mu\text{m}$ and the air-flow temperature is 263.15 K for both the NUM and TM approaches. We use a constant air speed of 5 ms^{-1} resulting in $Re_p = 80$, $Nu = 6.7$ and $Sh = 6.5$ (using Eq. (3)). The values used here are typical of a saltating ice particle (Thorpe and Mason, 1966; Kok and Renno, 2009; Vionnet et al., 201

In Experiment I, we study the heat and mass output from a sublimating ice grain as a function of time. In the first case, Experiment I-A, we consider the effect of three different values of air-flow saturation-rate ($\sigma_* = 0.8, 0.9$ and 0.95) on differences between NUM and TM solutions. The NUM approach requires specification of the initial condition for the particle temperature ($T_{p,IC}$). In Experiment I-A, ($T_{p,IC}$) is taken to be the same as the air-flow temperature for the NUM approach, i.e, 263.15 K . Results for Experiment I-A are shown in Fig. 1(a-c), with subfigure (a) showing the mass output rate, F_M and subfigure (b) showing the heat output rate, F_Q . Note that in this figure and subsequent figures, +(-) signifies mass and heat gained (lost) by the air. Since we keep the temperature and under-saturation of the air constant, the solutions of the TM approach are “steady-state” solutions with constant heat and mass transfer rates as seen in Fig. 1a and b. On the other hand, since the NUM approach solves the coupled equations that consider the evolution of the particle temperature, the heat and mass transfer rates evolve with time.

It can be seen that the NUM solutions initially evolve with time and reconcile with the steady-state TM solutions after a transient regime of about 0.3 seconds. Since the initial temperature of the particle is the same as the air, there is no heat transfer between the air and the particle (see the second term of the R.H.S of Eq. 1) initially. Thus, all heat transfer rates are initially zero for the NUM case in Fig. 1b. The under-saturation of the air forces mass transfer from the ice particle to air and the energy for the phase change comes from the internal energy of the ice particle. This causes the particle temperature to drop (see Fig. 2 below). With the particle now colder than the air, heat transfer from the air to the particle commences and ultimately, the

energy for sublimation comes entirely from the heat extracted from the air. The initial dynamics of the heat and mass transfer are completely neglected by the TM approach. In subfigure (c), the errors ($Err(t) = \left(\int_0^t F_{NUM} dt / \int_0^t F_{TM} dt - 1 \right) * 100$) for mass, Err_M and heat, Err_Q are shown. The errors reduce dramatically with time (for example, 15% at 0.3 seconds) and interestingly do not depend on the saturation-rate of the air-flow.

5 In the following case, Experiment I-B, similar simulations as in Experiment I-A are performed, but with $\sigma_* = 0.95$ while the initial temperature difference between the particle and the air is varied as $T_{p,IC} - T_{Air} = -2, -1, 1, 2$ K. The results are shown in Fig. 1(d-f). It is interesting to note that for each of the four cases considered, the TM solution predicts sublimation of the particle (consistent with $\sigma_* < 1$, see numerator of R.H.S of Eq. 3). On the other hand, for cases with colder particles, the NUM solutions show that there is initially deposition on the particle, along with larger values of heat absorbed from the
 10 air. Correspondingly, in the cases with particles being warmer than the air, the mass loss is much higher in the NUM solution than that computed by the TM solution while the heat gained by the particle is also much higher. These higher differences are reflected in the Err_M and Err_Q curves in subfigure (f) where errors are found to be an order of magnitude higher than those in subfigure (c).

We define relaxation time ($\tau_{relaxation}$) as the time required for the NUM solution to reconcile with the TM solution. The
 15 importance of this quantity lies in the fact that if the residence time of a saltating ice grain in air is shorter than $\tau_{relaxation}$, the TM approach is likely to be erroneous and the NUM approach would be required. It is intuitive that $\tau_{relaxation}$ increases with d_p on account of increasing inertia and decreases with $|\mathbf{u}_{rel}|$ due to more vigorous heat and mass transfer. Experiment I was repeated for values of d_p and $|\mathbf{u}_{rel}|$ ranging between (50 – 1000 μm) and (0 – 10 ms^{-1}) respectively. The upper-bound of the wind-speed range is quite high and it is extremely unlikely to find $|\mathbf{u}_{rel}| > 10 \text{ms}^{-1}$ in naturally-occurring aeolian transport.
 20 Numerical results indeed confirm our intuition and it is found that for any given value of $|\mathbf{u}_{rel}|$, $\tau_{relaxation}$ is found to be $\propto d_p^\alpha$, where $\alpha (\sim 1.65)$. Furthermore, $\tau_{relaxation}$ decreases monotonically with increasing $|\mathbf{u}_{rel}|$ for a given value of d_p . For $d_p = 200 \mu\text{m}$, the plausible values of $\tau_{relaxation}$ are found to lie between 0.28 and 1.5 seconds (for $|\mathbf{u}_{rel}| = 10$ and 0 ms^{-1} respectively). Interestingly, $\tau_{relaxation}$ is not found to depend on either the saturation-rate of air and-or the difference between the initial particle ~~temperature and air~~ and air temperature. Plots of $\tau_{relaxation}$ are highly relevant to discussion in Sect. 3 and
 25 presented there.

In Fig. 2, evolution of particle diameter (d_p) and temperature (T_p) is presented with subfigures (a) and (b) respectively describing the evolution for simulations in Experiment I-A with (c) and (d) being the corresponding results from Experiment I-B. In Experiment I-A, the particle diameters reduce linearly with time for both the NUM and TM approaches with the more shrinking (or in other words, sublimation) in the NUM solutions. More interesting is the evolution of the particle temperature,
 30 where the particle undergoes significant cooling due to sublimation and ultimately achieves a constant temperature. For example, in the case for $\sigma_* = 0.8$, the particle temperature is ultimately 0.85 K lower than the initial particle temperature of 263.15 K. Note that for the TM approach, particle temperature is of no consequence and it is shown simply for reference.

Following results of Experiment I, in Experiment II, we explore the parameter space of (σ_* , $T_{p,IC} - T_{Air}$) and compute the total mass ($M = \int_0^t F_M dt$) and total heat ($Q = \int_0^t F_Q dt$) output by a sublimating ice grain for a finite time of $t = 0.5$ seconds.
 35 Results shown in Fig. 3 subfigures ~~a and b~~ (a) and (b) provide a comparison of the total mass lost using the NUM and TM

solutions respectively and the corresponding error is shown in subfigure e(c). Similar figures are presented for the total heat lost/gained by the air in subfigures (d-f). The inclusion of the inertial terms essentially causes the contours to be sloped for the NUM solution while the TM solutions do not depend on $T_{p,IC} - T_{Air}$ as expected. The error between the NUM and TM solutions are accentuated at high saturation-rate regimes, with errors larger than 30 % for $\sigma_* > 0.8$.

5 In summary, Experiments I and II highlight the fact that during the sublimation of an ice grain, there exists a finite, well-defined transient regime before the NUM solutions match the steady-state TM solutions. Furthermore, the NUM and TM solutions diverge rapidly with slight temperature differences between the particle and the air and with increasing σ_* (which is a cause of concern since in snow-covered environments, the air usually is highly saturated). The results described above prompt an interesting question: are the residence times of saltating ice particles comparable to $\tau_{relaxation}$? We use large-eddy
 10 simulations to answer this question in the following section.

3 Large-eddy Simulations of Saltating Snow

3.1 Experiment III and IV: Simulation Details

To further understand the implications of the differences between the NUM and the TM approach, we performed LES of the atmospheric surface layer with an erodible snow surface as the lower boundary. We describe here only the main details
 15 of the LES that are relevant to our discussion with full model description along with equations presented in Supplementary Material S1. The LES solves filtered Eulerian equations for momentum, temperature and specific humidity on a computational domain of $6.4\text{ m} \times 6.4\text{ m}$ in the horizontal directions with vertical extent of the domain being 6.4 m as well. The snow surface, which constitutes the lower boundary of the computational domain, consists of spherical snow particles with a log-normal size distribution with a mean particle diameter of $200\ \mu\text{m}$ and standard-deviation of $100\ \mu\text{m}$. The coupling between the erodible
 20 snow-bed and the atmosphere is modeled through statistical models for aerodynamic entrainment (Anderson and Haff, 1988), splashing and rebounding of particle grains (Kok and Renno, 2009), which have been updated recently by Comola and Lehning (2017) to include the effects of cohesion and heterogeneous particle sizes. The use of these models essentially allows for overcoming the immense computational cost of resolving individual grain-to-grain interactions and allow us to consider the snow-surface as a *bulk* quantity rather than a collection of millions of individual snow particles. Once the ice grains are in
 25 the flow, their equations of motion are solved in the Lagrangian frame of reference with only gravitational and turbulent form drag forces included. Since the particle velocities are known, $|\mathbf{u}_{rel}|$ is calculated explicitly and used to compute Re_p , Nu and Sh . The horizontal boundaries of the domain are periodic and the lower boundary condition (LBC) for velocity uses flux parameterizations based on Monin-Obukhov similarity theory, additionally corrected for flux partition between fluid and particles between the wall and the first flow grid point (Raupach, 1991; Shao and Li, 1999). The LBC for scalars (temperature and specific humidity) are flux-free and thus the only source/sink of heat and water vapor in the simulations is through the
 30 interaction of the flow with the saltating particles.

All simulations are performed on a grid of $64 \times 64 \times 128$ grid points with a uniform grid in the horizontal directions and a stretched grid in the vertical. A stationary turbulent flow is allowed to first develop, following which, the snow sur-




Figure 1. TM and NUM solutions for a particle of $200 \mu\text{m}$ diameter in different environmental conditions. **Experiment I-A:** (a) Rate of mass and (b) heat output with (c) corresponding errors; $T_{p,IC} - T_{a,\infty} = 0$, $\sigma_* = 0.8$ (squares), 0.9 (circles), 0.95 (triangles). **Experiment I-B:** (d-f) same as (a-c) with $\sigma_* = 0.95$; $T_{p,IC} - T_{Air} = -2 \text{ K}$ (squares), -1 K (circles), 1 K (triangles), 2 K (stars).




Figure 2. TM and NUM solutions for a particle of $200 \mu\text{m}$ diameter in different environmental conditions. **Experiment I-A:** Evolution of particle (a) diameter and (b) temperature; $T_{p,IC} - T_{Air} = 0$, $\sigma_* = 0.8$ (squares), 0.9 (circles), 0.95 (triangles). **Experiment I-B:** (c-d) same as (a-b) with $\sigma_* = 0.95$; $T_{p,IC} - T_{Air} = -2 \text{ K}$ (squares), -1 K (circles), 1 K (triangles), 2 K (stars). Note that the particle diameters are normalized by the initial diameter of the particle ($d_{p,IC}$).

face is allowed to be eroded by the air. All physical constants and parameters along with additional details of the numerical setup are provided in Supplementary Material S2. [The LES in the configuration used in this study resembles the classic case of LES of a channel flow common in computational fluid dynamics research. The LES code has been developed in-house for last many years beginning with \(Albertson and Parlange, 1999\) and has been used and validated for various atmospheric boundary layer problems such as flows over heterogeneous surfaces Bou-Zeid et al. \(2004\), hills \(Diebold et al., 2013\), diurnal cycles \(Kumar et al., 2006\), urban canopies \(Giometto et al., 2016\) and wind farms \(Calaf et al., 2010; Sharma et al., 2017\). The same code was used previously for modeling snow saltation by Groot Zwaafink et al. \(2014\).](#)




Figure 3. TM and NUM solutions for a particle of $200 \mu m$ diameter in different environmental conditions. **Experiment II:** Total mass output during 0.5 seconds by the (a) NUM and (b) TM solutions with (c) corresponding error for $\{0.3 \leq \sigma_* \leq 1.1, -5 K \leq T_p - T_{Air} \leq 5 K\}$. Similar plots for total heat output presented in (d-f).

For the TM approach, Eq. (4) is used to compute the specific humidity and (by multiplying with the latent heat of sublimation) heat forcing due to each ice grain in the flow. On the other hand, for the NUM approach, Eqs. (1) and (2) are solved and only the turbulent transfer of heat between the air and the particle (second term in R.H.S of Eq. (1)) acts as a heat forcing on the flow. An implication of the NUM approach is that the particle temperature evolves during the ice-grain's motion and this necessitates providing an initial condition for the particle temperature ($T_{p,IC}$).

The principle aims of Experiment III are to firstly quantify particle residence times (PRT) and their dependence on wind speeds and relative humidities and secondly, compute the differences in the heat and mass output between the NUM and the TM approaches during saltation of snow with complete feedback between the air and the particles. PRT is defined as the total time the particle is air-borne and in motion, including multiple hops across the surface. Note that the PRT is not computed for particles in suspension, i.e, particles that stay aloft and never return to the surface. Towards this goal, simulations are performed, each with a combination of initial surface stress, $u_* \in \{0.4, 0.6, 0.8\} \text{ ms}^{-1}$ and initial saturation-rate, $\sigma_* \in \{0.3, 0.6, 0.9\}$. These values are classified as low (L), medium (M) and high (H) and correspond to wind speed at 1 m height above the surface of 11 m/s, 16.3 m/s and 21.8 m/s respectively. Note that during fully developed snow transport, the particles in the air impart

drag on the flow causing the flow ~~the-to~~ decelerate. The wind speeds at 1 m during fully developed saltation are 8.77 m/s, 11.34 m/s and 13 m/s respectively. The simulations are named as $U\alpha\text{-R}\beta$, where $(\alpha, \beta) \in \{L, M, H\}$. Each combination is simulated independently for the NUM and TM approaches resulting in a total of eighteen simulations. Experiment III is limited to simulating the *usual* case where the initial air temperature ($T_{Air,IC}$) is the same as $T_{p,IC}$.

- 5 Experiment IV is aimed at exploring the implications of differences between the two approaches in cases where $T_{Air,IC}$ is significantly different from $T_{p,IC}$. Such conditions can occur in nature during events such as marine-air intrusions, katabatic winds, spring-season saltation events and winter flows over sea-ice floes, where significant temperature differences between the air and snow-surface are likely. We repeat the low wind case of Experiment III with $u_* = 0.4\text{ms}^{-1}$ and choose the initial saturation-rate to be 0.95, motivated by results in Experiment II where errors were found to increase with increasing saturation-
- 10 rate. Simulations (named as UL-T(γ), where $T_{Air,IC} - T_{p,IC} = \gamma$) are performed once again for each of two approaches with $\gamma \in \{\pm 1\text{K}, \pm 2.5\text{K}, \pm 5\text{K}\}$ resulting in a total of twelve simulations. In all simulations performed for Experiments III and IV, $T_{p,IC} = 263.15\text{K}$. It is important to note that the initial condition for particle temperature ($T_{p,IC}$) is fixed throughout the simulation period, which essentially means that surface temperature is kept constant. This is consistent with the imposed zero flux of heat at the surface. This imposition will be justified *a posteriori* in the following section.

15 3.2 Results

In this section, results from the LESs performed for Experiments III and IV are presented. Note that only the relevant results are presented, namely (a) particle residence times as a function of particle diameters and different forcing setups and (b) differences between the NUM and TM approaches for calculating average mass and heat transfer rates during saltation. Other results, for example, vertical profiles of mean and turbulent quantities, although interesting are relegated to the supplementary material

20 as their analysis is out of scope of the current work. Additionally, ~~a video illustration~~ [two video illustrations](#) (Supplementary Movie M1 [and M2](#), see [Supplementary Material S4](#)) of an LES is provided to help visualize and frame the context of the simulations performed.

3.2.1 Particle Residence Times versus $\tau_{relaxation}$

As mentioned in the concluding lines of the Sect. 2, the principle quantity of interest is the PRT of saltating ice grains. In Fig. 4a,

25 the mean and median PRT of five different simulations of Experiment III are shown as a function of the particle diameter. Additionally, values of $\tau_{relaxation}$ computed in Experiment I for wind speeds ranging from 0 to 10ms^{-1} are also shown in the shaded region. Recall that the shaded region represents all the plausible values of $\tau_{relaxation}$ in naturally-occurring aeolian transport. As examples, $\tau_{relaxation}$ trends for 3 wind speeds, 0, 1 and 10ms^{-1} are shown and the power-law dependence can clearly be seen. It is found that $\tau_{relaxation}$ is comparable to the PRT of saltating grains with diameters between 125 and 225

30 μm . For $200\mu\text{m}$, the mean PRT is found to be 0.6 seconds while the median PRT is 0.2 seconds, which is outside the range of admissible values of $\tau_{relaxation}$. For particles larger than $225\mu\text{m}$, the PRTs are an order of magnitude smaller than plausible values of $\tau_{relaxation}$ and therefore the TM model is likely to provide wrong values of mass loss. On the other hand, lighter

particles with diameters smaller than $100 \mu\text{m}$ have much longer PRTs and the TM model is therefore valid. This proves that while the TM model is applicable for a majority of particles in suspension, it is likely to cause errors for particles in saltation.

Results presented in Fig. 4a provides two additional insights. Firstly, it is quite interesting to note that particles larger than $100 \mu\text{m}$ have the same mean PRT irrespective of low, medium or high mean wind speeds. This means that the dynamics of the heavier particles are unaffected by different mean wind speeds simulated in Experiment III, which is consistent with the notion of self-organized saltation, which has recently been shown by Paterna et al. (2016). For particles smaller than $100 \mu\text{m}$, the mean PRTs increases with wind speed. Secondly, the initial saturation-rate does not seem to effect the PRT statistics for medium and high wind conditions and the UM- and UH- curves for different R values overlap (this is the reason only five PRTs are shown in Fig. 4a). In these cases turbulence is sufficient to rapidly mix any temperature anomaly due to sublimation throughout the surface layer. On the other hand, the mean PRTs of particles smaller than $75 \mu\text{m}$, decrease with decreasing initial σ_* . A preliminary hypothesis is that in low wind conditions (UL), low initial saturation-rate results in more sublimation and cooling near the surface, resulting in suppression of vertical motions. ~~This is reflected in the mean PRTs of particles smaller than $75 \mu\text{m}$, which decrease with decreasing initial σ_* .~~ Even though this is an interesting result, ~~we~~ further research is needed to definitively link drifting snow sublimation to lower PRTs. We leave further exploration of this phenomenon for a future study with some preliminary analysis provided in Supplementary Material S3.

The PRT distributions are found to be quasi-exponential with long tails, thus resulting in large differences in mean and median PRTs shown in Fig. 4a. These distributions are also strongly dependent on the particle diameter. As an illustration, in Fig. 4b, cumulative distributions of PRTs are shown for four particle diameters along with the corresponding range of plausible $\tau_{relaxation}$ values. For the mean particle diameter of $200 \mu\text{m}$, we find that between 65% to 85% of particles have PRTs shorter than $\tau_{relaxation}$, whereas for the $75 \mu\text{m}$ particles, at most 30% particles lie below the maximum $\tau_{relaxation}$ threshold. This reinforces the fact that applying the steady-state TM solution to sublimating ice-grains in saltation could be potentially erroneous.

3.2.2 Differences in total mass loss between NUM and TM models

We can directly assess the implications of differences in grain-scale sublimation between the two approaches on total mass loss rates during saltation at larger spatial scales as simulated using LES in Experiments III and IV. In Fig. 5, we compare the total 15-min averaged rate of mass loss computed in all cases in Experiment III (subfigure a) and Experiment IV (subfigure c) using the NUM and the TM approaches with corresponding errors shown in subfigures b and d respectively. Recalling the adopted convention of +(-) as gain(loss) of flow quantities, it can be seen in Experiment III, that sublimation increases with u_* and decreases with σ_* . The errors on the other hand increase with increasing values of both u_* and σ_* . The increase in error with u_* is mostly due to the fact that an increase in u_* proportionally increases the total mass entrained by air (see Supplementary Fig. S1). The increase in error with increasing σ_* is in accordance with analysis done in Experiment II (see Fig. 3(c,f)) where it was shown that the NUM and TM solutions diverge with increasing saturation-rate. The least error, 26% is found for case UL-RL (i.e., $u_* = 0.4$, $\sigma_* = 0.3$) while the largest error, 38% is found for UH-RH ($u_* = 0.8$, $\sigma_* = 0.9$). Overall, for all the simulation combinations, the NUM approach computes larger mass-loss than the TM approach.






Figure 4. (a) Mean and median particle residence time (PRT) as a function of particle diameter. The plausible values of $\tau_{relaxation}$ are represented by the shaded region with trends for three values of $|\mathbf{u}_{rel}|$ shown by straight lines. Note that the horizontal axis is logarithmic. (b) Cumulative Distribution Functions of PRTs for four particle diameters along with range of plausible $\tau_{relaxation}$ values marked by overlying black curves

Experiment IV highlights the effect of temperature difference between particle and air on sublimation. As shown in Fig. 5c, the mass output is found to be negative (deposition) for the NUM solutions when the air is warmer than the particles (i.e., cases UL-T(γ) with $\gamma > 0$). This is contrary to the TM solutions which indicate sublimation. In cases with $\gamma < 0$, the NUM approach shows a much higher sublimation rate than the TM solutions. This occurs firstly due to higher vapor pressure at the grain surface that results in enhanced vapor transport and secondly because the warmer particles heat the surrounding air via sensible heat exchange, causing the relative humidity to decrease. Errors increase dramatically from an already high 100% for UL-T(+1) to 800% for UL-T(+5). Simulations performed for medium and high wind cases in Experiment IV showed even higher errors, similar to results in Experiment III and are shown here.

4 Discussion and Conclusion

10 In this article, we revisit the Thorpe and Mason (1966) model used to calculate sublimation of drifting and blowing snow and check its validity for saltating ice grains. We highlight the fact that solutions to unsteady heat and mass transfer equations (NUM solutions) converge to the steady-state TM model solutions after a *relaxation* time, denoted as $\tau_{relaxation}$ that has a power-law dependence on the particle diameter and is inversely proportional to the relative wind speed. Through extensive LESs of snow

EXPERIMENT - III



EXPERIMENT - IV





Figure 5. Experiment III: (a) Average rate of mass loss during 15 minutes of saltation, (b) Error between NUM and TM solutions. Corresponding plots for Experiment IV in (c) and (d) respectively. Note that the units used for rate of mass loss are kilograms per unit area per unit year.

saltation, we compute the statistics of the PRTs as a function of their diameters and find them to be comparable to $\tau_{relaxation}$. This helps explain the difference between mass output when using the NUM model to the TM approach, also computed during the same LESs. The NUM approach computes higher sublimation losses ranging from 26% in low-wind, low saturation-rate conditions to 38% in high-wind, high saturation-rate conditions. Another set of numerical experiments explore the role of temperature differences between particle and air temperature in inducing differences between NUM and TM solutions. We find the effect to be extremely dramatic with errors of 100% for a temperature difference of 1 K with increasing errors for larger temperature perturbations. In general, the two solutions are found to diverge rapidly as the saturation-rate tends towards 1. The results showing differences of mass output between the NUM and TM approaches in the LESs in Experiments III and IV, with complete feedback between particles and the air are thus shown to be closely correlated to the results from extremely idealized simulations of heat and mass transfer from a solitary ice grain in Experiments I and II.

The LES results do come with a few important caveats. Firstly, the temperature and specific humidity fluxes at the surface are neglected. In other words, particles lying on the surface are considered to be dormant and do not exchange heat or mass with the air. A corollary to neglecting the scalar fluxes at the surface is that the initial condition for temperature of the particles entering the flow is fixed. This may be justified by considering that during drifting and blowing snow events, the friction velocity at the surface drops dramatically. This fact has been observed in both in experiments (Walter et al., 2014) and in our current LESs (see Supplementary Fig. S2). This implies that direct turbulent exchange between the surface and air is curtailed and instead, the dominant exchange occurs between air-borne particles and the air. In fully-developed snow transport events, this is most likely to be true and only in intermittent snow-transport events will the surface fluxes be relatively important. This is nevertheless an important assertion that shall be more closely examined in future studies involving a full surface energy balance model, where the evolving temperature of the saltating ice grains, prior to deposition is taken into account while calculating snow-surface temperatures.

Further work is required to make concrete improvements to modeling of sublimation of saltating snow, especially in large-scale models that do not explicitly resolve saltation dynamics. One potential approach is to modify the Monin-Obukhov based lower boundary conditions for heat and moisture to account for particle temperature during blowing snow events. An ancillary outcome of this study is the discovery that buoyancy can affect the dynamics of lighter snow particles (with diameters less than $75 \mu\text{m}$) and decrease their residence times. Investigating this phenomenon requires a detailed analysis of turbulent structure within the saltation layer and is left for future publications.

In conclusion, analogous to the role played by saltating grains in efficient momentum transfer to the underlying granular bed, the NUM approach can be considered as an efficient transfer of heat and mass between the flow and the underlying snow surface, albeit with a closer physical relationship between the thermodynamics of the snow surface and that of the air. Thus, along with momentum balance of blowing snow particles, particle temperature and its thermal balance must also be taken into account.

Acknowledgements. We acknowledge the support of the Swiss National Science Foundation (Grant n. 200021-150146) and the Swiss Super Computing Center (CSCS) for providing computational resources (Project: s633). We additionally thank Marco Giometto for providing the original version of the LES code and Hendrik Huwald, Tristan Brauchli, Annelen Kahl and Celine Labouesse for illuminating discussions and important suggestions in improving the manuscript. No new data were used in producing this manuscript.

References

- Albertson, J. D. and Parlange, M. B.: Natural integration of scalar fluxes from complex terrain, *Advances in Water Resources*, 23, 239 – 252, [https://doi.org/https://doi.org/10.1016/S0309-1708\(99\)00011-1](https://doi.org/https://doi.org/10.1016/S0309-1708(99)00011-1), <http://www.sciencedirect.com/science/article/pii/S0309170899000111>, 1999.
- 5 Anderson, R. S. and Haff, P. K.: Simulation of Eolian Saltation, *Science*, 241, 820, <https://search.proquest.com/docview/213543077?accountid=27198>, copyright - Copyright American Association for the Advancement of Science Aug 12, 1988; Last updated - 2014-05-30; CODEN - SCIEAS, 1988.
- Bou-Zeid, E., Meneveau, C., and Parlange, M. B.: Large-eddy simulation of neutral atmospheric boundary layer flow over heterogeneous surfaces: Blending height and effective surface roughness, *Water Resources Research*, 40, 2004.
- 10 Calaf, M., Meneveau, C., and Meyers, J.: Large eddy simulation study of fully developed wind-turbine array boundary layers, *Physics of fluids*, 22, 015 110, 2010.
- Comola, F. and Lehning, M.: Energy- and momentum-conserving model of splash entrainment in sand and snow saltation, *Geophysical Research Letters*, 44, 1601–1609, <https://doi.org/10.1002/2016GL071822>, <http://dx.doi.org/10.1002/2016GL071822>, 2016GL071822, 2017.
- Dai, X. and Huang, N.: Numerical simulation of drifting snow sublimation in the saltation layer., *Scientific reports*, 4, 6611, <https://doi.org/10.1038/srep06611>, <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=4196100>{&}tool=pmcentrez{&}rendertype=abstract, 2014.
- 15 Déry, S. J. and Yau, M. K.: Large-scale mass balance effects of blowing snow and surface sublimation, *Journal of Geophysical Research Atmospheres*, 107, <https://doi.org/10.1029/2001JD001251>, 2002.
- Diebold, M., Higgins, C., Fang, J., Bechmann, A., and Parlange, M. B.: Flow over hills: a large-eddy simulation of the Bolund case, *Boundary-layer meteorology*, 148, 177–194, 2013.
- 20 Doorschot, J. J. J. and Lehning, M.: Equilibrium Saltation: Mass Fluxes, Aerodynamic Entrainment, and Dependence on Grain Properties, *Boundary-Layer Meteorology*, 104, 111–130, <https://doi.org/10.1023/A:1015516420286>, <https://doi.org/10.1023/A:1015516420286>, 2002.
- Dover, S.: Numerical Modelling of Blowing Snow, Ph.D. thesis, University of Leeds, 1993.
- 25 Giometto, M., Christen, A., Meneveau, C., Fang, J., Krafczyk, M., and Parlange, M.: Spatial characteristics of roughness sublayer mean flow and turbulence over a realistic urban surface, *Boundary-layer meteorology*, 160, 425–452, 2016.
- Grazioli, J., Madeleine, J.-B., Gallée, H., Forbes, R. M., Genthon, C., Krinner, G., and Berne, A.: Katabatic winds diminish precipitation contribution to the Antarctic ice mass balance, *Proceedings of the National Academy of Sciences*, 114, 10 858–10 863, <https://doi.org/10.1073/pnas.1707633114>, <http://www.pnas.org/content/114/41/10858.abstract>, 2017.
- 30 Groot Zwaaftink, C. D., Löwe, H., Mott, R., Bavay, M., and Lehning, M.: Drifting snow sublimation: A high-resolution 3-D model with temperature and moisture feedbacks, *Journal of Geophysical Research: Atmospheres*, 116, n/a–n/a, <https://doi.org/10.1029/2011JD015754>, <http://dx.doi.org/10.1029/2011JD015754>, d16107, 2011.
- Groot Zwaaftink, C. D., Diebold, M., Horender, S., Overney, J., Lieberherr, G., Parlange, M. B., and Lehning, M.: Modelling Small-Scale Drifting Snow with a Lagrangian Stochastic Model Based on Large-Eddy Simulations, *Boundary-Layer Meteorology*, 153, 117–139, <https://doi.org/10.1007/s10546-014-9934-2>, <https://doi.org/10.1007/s10546-014-9934-2>, 2014.
- 35

- Helgans, B. and Richter, D. H.: Turbulent latent and sensible heat flux in the presence of evaporative droplets, *International Journal of Multiphase Flow*, 78, 1 – 11, <https://doi.org/https://doi.org/10.1016/j.ijmultiphaseflow.2015.09.010>, <http://www.sciencedirect.com/science/article/pii/S0301932215002086>, 2016.
- Kok, J. F. and Renno, N. O.: A comprehensive numerical model of steady state saltation (COMSALT), *Journal of Geophysical Research: Atmospheres*, 114, n/a–n/a, <https://doi.org/10.1029/2009JD011702>, <http://dx.doi.org/10.1029/2009JD011702>, d17204, 2009.
- Kumar, V., Kleissl, J., Meneveau, C., and Parlange, M. B.: Large-eddy simulation of a diurnal cycle of the atmospheric boundary layer: Atmospheric stability and scaling issues, *Water resources research*, 42, 2006.
- Lenaerts, J. T. M., van den Broeke, M. R., Déry, S. J., van Meijgaard, E., van de Berg, W. J., Palm, S. P., and Sanz Rodrigo, J.: Modeling drifting snow in Antarctica with a regional climate model: 1. Methods and model evaluation, *Journal of Geophysical Research: Atmospheres*, 117, n/a–n/a, <https://doi.org/10.1029/2011JD016145>, <http://doi.wiley.com/10.1029/2011JD016145>, 2012.
- Liston, G. E. and Sturm, M.: The role of winter sublimation in the Arctic moisture budget, *Nordic Hydrology*, 35, 325–334, <https://doi.org/10.1029/2001JD001251>, <ftp://ftp.cira.colostate.edu/ftp/Liston/papers/first{ }author/2004.liston.NH.pdf>, 2004.
- Mann, G. W., Anderson, P. S., and Mobbs, S. D.: Profile measurements of blowing snow at Halley, Antarctica, *Journal of Geophysical Research: Atmospheres*, 105, 24 491–24 508, <https://doi.org/10.1029/2000JD900247>, <http://dx.doi.org/10.1029/2000JD900247>, 2000.
- Paterna, E., Crivelli, P., and Lehning, M.: Decoupling of mass flux and turbulent wind fluctuations in drifting snow, *Geophysical Research Letters*, 43, 4441–4447, <https://doi.org/10.1002/2016GL068171>, <http://dx.doi.org/10.1002/2016GL068171>, 2016GL068171, 2016.
- Raupach, M.: Saltation layers, vegetation canopies and roughness lengths, *Acta Mechanica*, I, pp. 83–96, 1991.
- Rémy, F. and Frezzotti, M.: Antarctica ice sheet mass balance, *Comptes Rendus - Geoscience*, 338, 1084–1097, <https://doi.org/10.1016/j.crte.2006.05.009>, 2006.
- Rignot, E., Velicogna, I., van den Broeke, M. R., Monaghan, A., and Lenaerts, J. T. M.: Acceleration of the contribution of the Greenland and Antarctic ice sheets to sea level rise, *Geophysical Research Letters*, 38, n/a–n/a, <https://doi.org/10.1029/2011GL046583>, <http://dx.doi.org/10.1029/2011GL046583>, 105503, 2011.
- Russo, E., Kuerten, J. G. M., van der Geld, C. W. M., and Geurts, B. J.: Water droplet condensation and evaporation in turbulent channel flow, *Journal of Fluid Mechanics*, 749, 666–700, <https://doi.org/10.1017/jfm.2014.239>, 2014.
- Shao, Y. and Li, A.: Numerical Modelling of Saltation in the Atmospheric Surface Layer, *Boundary-Layer Meteorology*, 91, 199–225, <https://doi.org/10.1023/A:1001816013475>, <https://doi.org/10.1023/A:1001816013475>, 1999.
- Sharma, V., Parlange, M., and Calaf, M.: Perturbations to the spatial and temporal characteristics of the diurnally-varying atmospheric boundary layer due to an extensive wind farm, *Boundary-Layer Meteorology*, 162, 255–282, 2017.
- Sturm, M., Holmgren, J., and Perovich, D. K.: Winter snow cover on the sea ice of the Arctic Ocean at the Surface Heat Budget of the Arctic Ocean (SHEBA): Temporal evolution and spatial variability, *Journal of Geophysical Research: Oceans*, 107, 2002.
- Thorpe, A. D. and Mason, B. J.: The evaporation of ice spheres and ice crystals, *British Journal of Applied Physics*, 17, 541, <https://doi.org/10.1088/0508-3443/17/4/316>, <http://iopscience.iop.org/0508-3443/17/4/316>, 1966.
- van den Broeke, M., van de Berg, W. J., van Meijgaard, E., and Reijmer, C.: Identification of Antarctic ablation areas using a regional atmospheric climate model, *Journal of Geophysical Research*, 111, D18 110, <https://doi.org/10.1029/2006JD007127>, <http://doi.wiley.com/10.1029/2006JD007127>, 2006.
- Vionnet, V., Martin, E., Masson, V., Guyomarc'H, G., Naaim-Bouvet, F., Prokop, A., Durand, Y., and Lac, C.: Simulation of wind-induced snow transport and sublimation in alpine terrain using a fully coupled snowpack/atmosphere model, *Cryosphere*, 8, 395–415, <https://doi.org/10.5194/tc-8-395-2014>, 2014.

- Vögeli, C., Lehning, M., Wever, N., and Bavay, M.: Scaling Precipitation Input to Spatially Distributed Hydrological Models by Measured Snow Distribution, *Frontiers in Earth Science*, 4, 108, <https://doi.org/10.3389/feart.2016.00108>, <https://www.frontiersin.org/article/10.3389/feart.2016.00108>, 2016.
- 5 Walter, B., Horender, S., Voegeli, C., and Lehning, M.: Experimental assessment of Owen's second hypothesis on surface shear stress induced by a fluid during sediment saltation, *Geophysical Research Letters*, 41, 6298–6305, <https://doi.org/10.1002/2014GL061069>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014GL061069>, 2014.
- Wever, N., Lehning, M., Clifton, A., Rüedi, J. D., Nishimura, K., Nemoto, M., Yamaguchi, S., and Sato, A.: Verification of moisture budgets during drifting snow conditions in a cold wind tunnel, *Water Resources Research*, 45, 1–14, <https://doi.org/10.1029/2008WR007522>, 2009.
- 10 Xiao, J., Bintanja, R., Déry, S. J., Mann, G. W., and Taylor, P. A.: An Intercomparison Among Four Models Of Blowing Snow, *Boundary-Layer Meteorology*, 97, 109–135, <https://doi.org/10.1023/A:1002795531073>, <https://doi.org/10.1023/A:1002795531073>, 2000.