

Answer to the referee's comments

We would like to thank Carrie Jennings and the anonymous reviewer for the careful reading of our manuscript and the helpful comments. Replies to referee's comments are addressed below (blue colored). Broadly speaking, both reviewer agree on the novelty of our experimental approach but ask for a more clarification on the limitations of the experiment.

Referee #1 : Carrie Jennings

General comments

“I appreciate this modeling attempt. I am not aware of any other work on modeling subglacial hydrology since G. Catania and C. Paola, 2001, Braiding under glass. *Geology*, 29(3), 259-262. I believe it is relevant and should be cited. Models inform our intuition. They cannot prove anything but they can lead us to a better understanding of physical processes if we understand the limitations of the model setup. I would like to see the model and its limitations more fully described. What about model is not like real world? What are the shortcomings? How could these shortcomings affect model results and deviate from real-world processes?”

The referee is right to point out that analog modelling provides intuitions and ideas on a specific process but does not constitute a proof. Experiments produce morphologies and dynamics that, although imperfect, compare well with natural systems despite differences of spatial scale, time scale, material properties, and number of active processes. Thanks to the numerous comments of the referee, we added (i) some restrictions on the interpretation of the experiments and (ii) the limitations of the model to reproduce its natural counterpart. An entire paragraph (section 2.3) is now entirely dedicated to the limitations of the model.

Specific comments – Abstract

L20 – 21. Do they ever evolve to be efficient drainage systems? Tunnels seem very short-lived and episodic to me and ice-streaming redevelops again and again. Your experiment represents a very coarse-textured bed when scaled up, so this may be an effect related to grain size.

In our experiments, tunnel valleys stay active during the whole experiment. Meltwater routing, dynamics of tunnel formation and evolution of ice stream dynamics are intricately connected during experiments. Figure 5 in the revised version shows that tunnel valleys, once reaching a certain overall volume, reduce the silicon flow velocity until the modelled ice stream switches off. This suggests that the tunnel valley system evolves in an efficient drainage system able to drain all the subglacial water, thus reducing water pressure and enhancing basal friction. Grain size has necessarily an effect on the drainage capacity of experimental tunnel valleys. The use of a substratum with different properties would probably change tunnel valleys amount and development rates.

Specific comments – Introduction

L32-33. drainage pathway for sediment...reword ?

We suggest to reword the sentence as:

“approximately 80% of the ice discharge is focused in a finite number of ice streams, which act as preferential drainage pathways for meltwater also (Bamber et al., 2000; Bennett, 2003)”

L34. Ancient ? Is palaeo a word by itself

We suggest to replace every use of palaeo by “ancient” or “former”.

L35. Is m. correct ?

We thank the reviewer for this comment. It is true that the classical way to write any ratio is to use a slash. We suggest modifying all the “m.s⁻¹” by “m/s”.

L37. How about the evolution ?

We suggest to modify the end of the sentence in: “and the controls on their dynamics evolution remain debated”.

L54. Reference for this? I think it relates to Ice Stream C and B in WAIS

We suggest to add Vaughan et al., (2008) and Carter et al., (2013) as references for the subglacial water piracy processes.

L68. ours are much shorter but formed in segments--or at least are interrupted by ice-marginal fans.

We modify the sentence to inform the range of tunnel valleys dimensions: “These valleys are elongated and over-deepened hollows, ranging from a few kilometres to hundreds of kilometres long, from hundreds metres to several kilometres wide and from meters to hundreds of meters deep.”

L72. not by all--this feels a bit like you are setting up a straw-man type argument.

We agree with the reviewer that this sentence was perhaps misleading. We suggest to rewrite as follows:

“Indeed, ice streams commonly operate because of high basal water pressure while the development of a tunnel valley system generally leads to enhances drainage efficiency and basal water pressure reduction (*Engelhardt et al., 1990; Kyrke-Smith et al., 2014; Marciznek and Piotrowski, 2006*).”

L74-76. I would say that from the field evidence, there is a third process: 1) ice streaming; 2) drainage through tunnel valleys; 3) stagnation of the ice margin.

We suggest adding a sentence to explain that field studies have already suggested a link between outburst flood and a set of events involving ice streaming, tunnel valley formation and ice margin stagnation: “Several field studies have already suggested a connection between catastrophic glacial outburst floods at ice sheets margins and a suite of events involving ice streaming, tunnel valley development and stagnation of the ice margin. (*Bell et al., 2007; Hooke and Jennings, 2006; Jørgensen and Piotrowski, 2003; Alley et al., 2006*).”

L79. Bering glacier behavior during and after a surge comes close https://www.cambridge.org/core/services/aop-cambridge-core/content/view/08A39D0DD9EBE9C32D232B7769B55728/S0022143000202311a.pdf/lacuna_banded_surface_depressions_occurrence_and_conditions_of_formation_bering_glacier_alaska.pdf

This reference does not connect ice streaming with subglacial erosional processes so we choose to not add this reference because it is not appropriate with the meaning of our sentence.

L86. How would tunnel valleys influence the location of ice streaming if they only happen after streaming is already occurring? This may need to be more precisely worded.

We agree with the reviewer that this sentence requires clarifications. We suggest writing as follows:

“We propose that the location and initiation of ice streams might arise from subglacial meltwater pocket migration and drainage pathways and that the evolution of ice stream dynamics is latter controlled by subglacial drainage reorganization and tunnel valleys development.”

L86. This work ?

We suggest to writing as follows: “This study reconciles into a single story”

Specific comments – Experimental ice stream model

L94. partially overcome ?

We agree that this phrasing might not be the best one to explain that our model is useful to explore the connection between subglacial meltwater routing and ice dynamics without being a perfect representation of nature. We suggest to rewrite this section as follows:

“Considering all these processes and components simultaneously, together with processes of subglacial erosion, is thus a challenge for numerical computational modelling (Fowler and Johnson, 1995; Marshall, 2005; Bingham et al., 2010). Based on this statement, some attempts in analogue modelling have been made to improve our knowledge on subglacial erosional processes by meltwater (Catania and Paola, 2001) or gravity current instabilities produced by lubrication (Kowal and Worster, 2015). To combine ice flow dynamics and erosional aspects in a single model, we designed an alternative experimental approach that allows simultaneous modelling of ice flow, subglacial hydrology and sedimentary/geomorphic processes. With all the precautions of use inherent of analogue modelling, our experiments reproduce morphologies and dynamics that compare well with subglacial landforms and ice stream dynamics despite some differences of spatial and time scales and a number of active processes (e.g. Paola et al., 2009).”

L97. Paola et al., 2009--the way this is referenced now makes it seem like they simultaneously modeled these things.

We thank the reviewer for this clarification. See the modification made to answer the last comment.

L98. Use as an example? This is not the fundamental reference for the previous statement. Shreve, R.L. 1972. Movement of water in glaciers: Journal of Glaciology, 11(62), 205-214? or even earlier: Glen, J.W. 1952. The stability of ice-dammed lakes and other water-filled holes in glaciers. Journal of Glaciology, 2(15), 316-318.

We agree with the reviewer that it is not the fundamental reference and we added Shreve (1972) and Glen (1952) according to your proposition.

Line 99 : thus in part controlled by....This seems overly simplistic or at least backwards--active margins of ice sheets are an expression of mass balance, bed topography and ice surface slope

We agree with the reviewer that meltwater routing is function of many parameters. We suggest modifying this sentence so that we understand that ice slope is prevailing to control meltwater routing but that subglacial topography, and the mass balance also influence meltwater routes. We suggest to add a section dedicated to the limitations of the model (2.3 Scaling and limitations) and to rewrite this sentence as follows:

“Subglacial meltwater routing is indeed controlled by the ice surface, slope, the bed topography and the glacier mass balance (Röthlisberger and Lang, 1987). The ice surface slope controls potentiometric surfaces, generally guiding subglacial water flow parallel to ice sheet surfaces (Glen, 1952; Shreve, 1972; Fountain and Walder, 1998).”

Lines 100-101 : I am not following. It appears I need to refer to the earlier paper. Can this be avoided by providing a bit more here?

We understand the enquiry of the reviewer to understand the scaling of the experiment without referring to the first paper presenting the experiment. We propose to add explanations on the scaling and to move this section after the description of the model (cf. new section 2.3. “Scaling and limitations”):

“Considering that meltwater is here simulated by an injection of water, the rules of a classical scaling where the model is a miniaturisation of nature are not practical (*Paola et al., 2009*). Subglacial water drainage is generally controlled by fluctuations in locations of ice sheet margins. Similarly, in our experiments, the silicon putty margin controls the water pressure gradient. In this perspective, we base the scaling on the displacement of the natural ice and experimental silicon margins through time. We use a unit-free speed ratio between the silicon/ice margin velocity and the incision rate of experimental/natural tunnel valleys. The scaling is designed to ensure that the value of the ratio between margin velocity and incision rate of tunnel valleys in the experiment equals its value in natural. The projection of the minimal and maximal experimental speed ratios on the field of possible natural speed ratios highlights the field of validity of the experiments and defines the range of natural settings we can reproduce experimentally (full details in *Lelandais et al., 2016*). The main scaling limit regards the viscosity ratios between glacier ice, silicon putty and water. The size of the experimental ice stream, being partly controlled by the high silicon viscosity, may be underestimated compared to the size of modelled tunnel valleys.”

L103-106. Models at SAFL U of M often use hollow glass beads to overcome issues of density when using small models.

Producing DEM of the ice-bed interface was one of the main goal of this study and glass beads properties would have probably been less suitable for photogrammetry and 3D reconstruction (reflection problems, transparency, lack of roughness etc...). However, we think that glass beads would probably lead to the same suite of events, with a similar process of tunnel valley formation as the density of glass beads and the sand we use are similar. However the morphologies of tunnel valleys would probably differ due to changes in substratum permeability and friction coefficient.

L110-113. Comment: Having trouble visualizing where water is injected based on this description. Is water focused in one area?

The water is injected through an injector placed at the centre of the model which corresponds to the center of the silicon layer. The radial boundary of the silicon layer provide a radial flow of water so water is not constrained to flow in only one direction. For the visualization the cross-sectional profile in the Figure 1 show how water is injected in the system.

L118. again, placement makes this feel like the first time someone suggested that rheological softening was function of strain rate, T, etc.

We suggest to modify in e.g. Bingham et al., 2010.

L119. Nor can the potentiometric surface of water within the ice.

We suggest adding another restriction to our model in this sentence. Water flow is not driven by the silicon surface slope in the experiment. We suggest to rewrite as:

“This punctual injection does not simulate the mosaic of meltwater production regions existing beneath glaciers or the episodic input from supraglacial/englacial meltwater reservoirs. Experimental meltwater routing is predominantly controlled by the water discharge we inject in our system and therefore differs from parameters controlling hydrology in glacial systems. Subglacial meltwater routing is indeed controlled by the ice surface slope, the bed topography and the glacier mass balance (*Röthlisberger and Lang, 1987*). The ice surface slope controls potentiometric surfaces, generally guiding subglacial water flow parallel to ice sheet surfaces (*Glen, 1952; Shreve, 1972; Fountain and Walder, 1998*).”

L120.: Appropriateness of reference: as I recall, he speculated and modeled that it was (based on dilatancy of layer?), but others measured it in W. Ant much more recently? Reword the way the citation is used?

We agree with the reviewer that the till influence on ice stream should be mentioned. We suggest to rewrite as follows:

“This model, designed to decipher the interaction between subglacial hydrology and ice dynamics, hinders the influence of bed topography and geology (especially the influence of subglacial till) (*Winsborrow et al., 2010*). The deformation of the subglacial till and its complex rheological behavior is known to promote ice streaming (*Alley et al., 1987*), modify the subglacial hydrology and alter the size of tunnel valleys. The development of an analogue material scaled to reproduce subglacial till characteristics is extremely difficult so we did not try to include the equivalent of a till layer in the experiment.”

L120. Till would also change the behavior of water beneath the ice and potentially after the tunnel development

L123. and narrower tunnels

A till layer is extremely difficult to reproduce so we did no try to include one in our model. Doing so, we probably enhance some processes in the development of the ice stream and in the development of tunnel valleys. Hence we suggest to add some restrictions as follows :

“The deformation of the subglacial till and its complex rheological behavior is known to promote ice streaming (*Alley et al., 1987*), modify the subglacial hydrology and alter the size of tunnel valleys. The development of an analogue material scaled to reproduce subglacial till characteristics is extremely difficult so we did not try to include the equivalent of a till layer in the experiment. We thus assume that the velocity contrasts observed in the experiment are thus likely to be amplified in natural ice sheets, by the complex rheological behaviour of ice and till. This may lead to the development of narrower ice streams with higher relative velocities and sharper lateral shear margins in natural ice sheets than in the experiment (*Raymond, 1987; Perol et al., 2015*).”

L130. 3 levels? I don't understand and diagram doesn't help resolve.

The figure is not helpful to understand how we dispose the UV markers. Hence, we modified the figure 1 in the revised version of the manuscript to help the readers distinguishing the 3 levels of UV markers.

L130. this word helps and could be used in text. However, why this style of water injection is considered to be realistic escapes me. How could water be added to the center of an ice sheet?

The central position of the injection is specified in an earlier comment. A circular shape of the silicon layer was preferred to avoid any preliminary constraints on the water flow route and to avoid lateral boundary effects on silicon flow. This circular layer of silicon simulates only a portion of an ice sheet but not the whole ice sheet. The central injection of water is thus simulating an upstream source of water along an ice sheet portion that does not correspond to the centre of an ice sheet.

L139. Vertical? You are seeing the ice surface sink? I think that this needs to be better explained because I thought it was probably as a result of horizontal advection of "ice" and deformation of ice into a void that is formed as sediment is evacuated. But from the caption I see that it is also (primarily?) because of the water pocket forming and that the surface is elevated. The caption and figures help but the text is not clear and I have to work hard to figure out all the possibilities. Are you facing a word limit? If not, make it easier on your reader to follow experimental design and expectations.

As the injected water is pressurized, we can observe and monitor vertical displacements of the silicon surface due to water flow. We did not see the silicon sink properly but we could monitor a subsidence area when the water pocket was moving from one place to another. We propose to add the following sentence in the next section describing the UV device:

“The monitoring of every UV marker positions (in both horizontal vertical plans) through time was used to produce velocity and vertical displacement maps. Vertical displacement maps are interpolated from the subtraction of the DEM at time t with the DEM generated from the photographs taken a few seconds before the injection.”

L169-170. I'd like to see photos of the setup also. These may have photo backgrounds but I cannot tell with the color overlay. Seems very idealized. Look at Ginny Catania's description of her model of subglacial drainage. Catania and Paola, 2001.

We agree with the reviewer that, in order to convince the readers who may not be familiar with such models, we should propose a better explanation of our device. Thus, we have add a picture of the device in Supplementary data. We also added a new figure in the revised version of the manuscript (Fig. 2) with six raw photographs of the main experiment stages described in figure 3.

L171-172. What is the scale of these tunnel valleys? Approximate volume of the fans? If the box is 70 cm across, how well resolved are they? I would like to see them rather than just trust the drawing.

The addition of figure 2 in the revised version with raw photographs of every stages should solve this problem. We also specify the size of every tunnel valleys and fans for every stages in the revised version of the manuscript.

Specific comments – Experimental results

L181-185. How long was the experiment run? How long did it take this change to occur?

L 188-189. This experiment was repeated 12 times with identical input parameters. A six-stage ice stream lifecycle linking outburst flooding transitory ice streaming and tunnel valley.

The experiment typically lasts 30 minutes (cf. Figure 3). The silicon flow pattern changes instantly when water injection starts (cf. Figure 3). We added the duration of the experiment and the information that the silicon flow pattern evolves spontaneously in response of water injection within the text as follows:

“This experiment was repeated 12 times with identical input parameters (a 30 mm-thick silicon layer of 150 mm radius; constant water input of 1.5 dm³/h during 1800 s). After an initial identical state, a six-stage ice stream lifecycle linking outburst flooding, transitory ice streaming and tunnel valleys development has been observed for all these simulations (Fig. 3a-f, Fig. 6).”

L 197. There are earlier references that first describe and document this phenomenon: https://www.igsoc.org/journal/35/120/igs_journal_vol35_issue120_pg201-208.pdf . or https://www.cambridge.org/core/services/aop-cambridge-core/content/view/B2AD36180AEF8E3E8403A7BF2D627319/S0022143000002689a.pdf/shortterm_velocity_and_waterpressure_variations_downglacier_from_a_riegel_storglaciaren_sweden.pdf

We thank the reviewer for this reference and add it at the end of the sentence.

L 201-202. Rerword. This sentence seems to be missing a word. It doesn't make sense as written.

We agree with the reviewer that this sentence is not correctly written. We suggest to modify as follows: “The lack of channels incised in the substratum indicates that water flow occurs as a distributed drainage system without any basal erosion”.

L 208. What does this mean, to migrate by distributed drainage? Can you simplify and say the water pockets migrate? Are you saying that the water pockets stay at the ice-bed interface and migrate?

We agree with the reviewer and simplify the sentence as follows:

“The experiment suggests that the migration of water pockets at the ice-bed interface can contribute to the emergence of ice streams”.

L 211. What is channel scale and how does it evolve? Is water emerging from the ice margin through a narrow channel that grows headward and bifurcates?

At this stage, water is drained as a sheet flow at the silicon margin. This drainage is associated with widespread erosion, wider than the subsequent tunnel valley, producing the first low-angle fan described in supplement Figure 3. Once the water pocket drained, the water flow channelizes and a narrow valley starts forming by regressive erosion. Tunnel valleys are constantly growing along the experiment. Their size (length, width, depth) have been added for every stage.

L 212. What is the width of the head of this fan? Is it a lot broader than the fan that develops in the next stage? Need scales. I do not know of anything like this at a terrestrial margin. Till deltas off the coast of Antarctica might be of a similar scale. I wouldn't over-emphasize this since you just have sand, not till in the subsurface. I don't think that real-world ice margins leak this easily, especially where frozen. So instead you get thrust moraines from water pressure drop and tunnels.

The width of the head of every fan have been added for every stage. The first fan originating from the outburst is similar to the fan developing afterward apart from the angle. Indeed the first fan is a low-angle fan and the subsequent fan developing with tunnel valley is a high-angle fan. Although both fans are similar in size, the first one originating from the outburst flood forms nearly instantly, however the second one forms progressively during tunnel valley development.

L 213-216. I would expect the ice stream to evolve immediately prior to the tunnel valley formation. Why decreasing basal water pressure lead to it? Are you sure ice stream isn't when bubble reaches margin, immediately prior to TV formation? How long is all this taking and can you really resolve it? Are you slowing down the cameras? Is it video? Stop motion photography?

Silicon flow is progressively accelerating during migration of the water pocket. Obviously, the silicon flow is high when the water pocket reaches the margin but we record a peak velocity when the water pocket drains. We use a system of 7 cameras with a 5 second delay, but to be sure of our results we ran some experiments with a 1 second delay to visualize and validate the process described in the text.

L 219. Comments: e.g. Kamb? Bering Glacier surge and outburst is well documented.

As pointed out by the second reviewer this reference is not the best fit to support our result so we decide to choose Anderson et al. (2005) who documented ice flow acceleration triggered by outburst flooding.

L 221. Magnusson et al., 2007

We suggest to add e.g. before the reference

L 226-227. What is their scale in the experiment? How do they scale with the ice streams?

We agree with the reviewer that the figure itself is not sufficient to estimate the dimensions of our tunnel valleys. Consequently, we added tunnel valleys sizes at every stage in the manuscript. Compared to nature, experimental tunnel valleys are disproportionate compared to the ice stream size. The model itself constrains the size of the ice stream. The disproportion could also be the consequence of the analogue material we use in the experimental setup. The silicon putty is way too viscous to be scaled to glacier ice (this is clearly mentioned in the text). Using a less viscous material, the ice stream would have probably been wider and more scaled to tunnel valley size. However, silicon was selected as it

shares some essential characteristics with glacier ice and mainly because its perfect transparency allows DEMs of the silicon-bed interface to be reconstructed.

L 227-228. I would like to see a cleaner description of model observations because here it appears that interpretations and discussion are interwoven.

We have chosen this organization which alternates description of the model and natural examples, directly after the description of every stage, in order 1- to validate each experimental observation by natural examples, 2- to avoid many repetitions in the manuscript to facilitate the reading. We wanted to base the main discussion on more general and global implications of our modelling results.

L 230 . What is the width at the head of the fan? Is it the width of the tunnel?

We added sizes of tunnel valleys and fans for every stage in the results part. As silicon flows, the silicon layer progressively pushes the fan so the width at the head of the fan is wider than the width of the tunnel valley.

L 249. and that eskers formed in tunnel valleys represent a waning flow stage.

It is true that eskers within tunnel valleys symbolize a decrease in water flow velocity. Hence, they might represent the ultimate stage of tunnel valley development. However, we cannot simulate this final deposition stage in our experiment.

L 259-260. We also see stagnation of the ice lobe margin. Large glaciotectonic thrust masses at ice margins are located near tunnel valley fans and seem to represent the fast flow stage immediately prior to drainage.

We thank the reviewer for this useful comment. We suggest to use these field evidence in the revised version to support our experimental results: “Large glaciotectonic thrust masses at the ice margin near tunnel valleys fans are generally assumed to be a field evidence a fast ice flow stage prior to drainage through tunnel valleys (*Hooke and Jennings, 2006*).”

L 265. I do not see field evidence of two, very different scales of floods and two styles of fan formation. I suspect that the first one you observe is more a result of the unusual way you are building water pressure beneath your ice sheet (at a single point).

The first outburst flood resulting from the first water pocket drainage is more obvious and its consequence on silicon flow is more visible (cf. Figure 3). The second outburst flood consists in the drainage of a new water pocket probably originating from the inefficiency of the first tunnel valleys to drain all water. This second water pocket is probably less significant than the first one so the consequence on the silicon is less visible. In nature if tunnel valleys originate from an outburst flood, we might not find any evidence from another catastrophic drainage as they are probably occurring in the same water path.

Specific comments – Proposed lifecycle of transitory ice streams

L 274. Are these the earliest references to this phenomenon? I think not. Use as examples or cite the foundational work.

We agree with the reviewer and add some earlier references: Bentley, 1987; Blankenship et al., 1993; Anandkrishnan et al., 1998

L 284. this is a review paper. As reviewed in Kehew and Piotrowski,

We agree with the reviewer and we suggest modifying the sentence as follows: “As reviewed in *Kehew et al. (2012)* and suggested in *Ravier et al. (2015)* this relation was suspected from the occurrence of tunnel valleys on ancient ice streams beds.”

L 285-286. You may be setting up a "straw man" because I don't know how widely believed/modeled this is.

Ice streams may arise from various processes (basal decoupling, deformation of the substratum...). However most of the models, emphasizes the development of high water pressure in the bed or at the ice-bed interface. For tunnel valleys we agree that we might not be so categorical about water pressure. We suggest to modify this sentence as follows:

“However, it raised a contradiction: subglacial meltwater pressures are generally supposed to be high below ice streams (Bennett, 2003) while tunnel valleys are generally assumed to operate at lower water pressures (Marczinek and Piotrowski, 2006).”

L 288-290. I think this has been speculated before from field evidence. You may be the first to physically model it, however.

We agree with the reviewer that we might rewrite this sentence to emphasize that although several observations have already connected ice stream with outburst flood and outburst flood with tunnel valley formation we are the first to model these interactions and to propose a single model connecting outburst flooding, ice stream and tunnel valley development.

“Although speculated from field evidences, our results demonstrate that ice streaming, tunnel valley formation, release of marginal outburst floods and subglacial water drainage reorganization may be interdependent parts of a single ice stream lifecycle that involves temporal changes in subglacial meltwater pressures (Fig. 6).”

L 291-293. no comma after Approximately. But more importantly, this pocket migration is highly dependent on the focused way you introduced water into the subglacial environment. I would not make too much of it since it is highly unrealistic.

We agree with the reviewer that the way we introduce water in the model is unrealistic. However, water injection at a given discharge triggers water flow at the silicon/substratum interface. The consequences on silicon flow should be comparable to the influence of subglacial meltwater on ice flow dynamics and therefore not be so unrealistic.

L 297-300. Why would they switch on precisely when water drains? How do ice streams migrate headward in this scenario? I think the ice stream migration timescale is very different than the water drainage timescale. You refer to timescales in a vague way in the beginning of the paper. Time to return to those ideas?

In our experiment, we define the ice stream birth phase when a corridor of high silicon flow appears. Before, the silicon flow velocity only increases above the migrating water pocket. Using a more viscous material than glacier ice has a consequence on timescale involved in ice stream migration. . The silicon putty accelerates the process of ice streaming and we cannot observe the headward migration of the ice stream. The migration of the experimental ice stream in response to re-routing of the water occurs almost instantaneously. Once more, this very quick migration may be a consequence of the high viscosity of the silicon we use during modelling.

L 309-310. in contrast, we see ice stream locations and tunnel systems becoming fixed. Tunnels are reoccupied again and again as an ice sheet retreats. I'm not saying that ice streams don't migrate, just at a different time scale (or again, this might be an artifact of the way you are introducing water.)

Our study only proposes an alternative solution for ice stream migration and lateral development of tunnel valley, based on meltwater re-routing. If we consider that tunnel valley development occurs progressively, the drainage can be inefficient, possibly leading to meltwater storage. The drainage of stored meltwater could trigger a second phase of ice flow acceleration. If drainage occurs laterally to the

main drainage path represented by the pre-existing tunnel valley system, drainage may indeed trigger a lateral migration of the ice stream path and the formation of a new tunnel valley (as observed in our experiment; Fig. 3). Of course, this drainage could also occur within the pre-existing tunnel valley system, thus not modifying the position of the ice stream path.

L 324-327. This seems to be taking the results of the experiment a bit far and a slightly misrepresenting the conclusions (or at least the dire nature of them) of these papers.

We agree with the reviewer that we might have over-interpreted the references we used. We propose to rewrite this sentence as follows: “This further suggest that tunnel valley development could secure ice sheet stability as hinted by Marczynek and Piotrowski. (2006) by preventing catastrophic ice stream collapses”.

L 328. a slash is not proper punctuation (a pet peeve).

We suggest to rewrite as :

“In a global change context, phenomena of ice stream stabilisation would requires that pre-existing and newly forming tunnel valleys systems expand sufficiently fast to accommodate increased meltwater production.”

L 329-332 : Comment : You may have needed to make the conclusions seem relevant to a broad audience but to readers of The Cryosphere, this seems a bit extreme and sensational.

We agree with the reviewer that the current conclusion is a bit extreme and focused on current global warming issues. We deleted the “sensational” part of the conclusion in the revised version of the manuscript.

L 335-336. This schematic is fine as long as we understand it is an interpretation. I'd like the original model results to look more "real" like photos of a model, and less like this.

This schematic is actually an interpretation of the six-stages ice stream lifecycle we describe in the manuscript, based on true experimental data. This interpretative sketch is actually drawn using experimental DEM of the bed and flow velocity maps of the silicon. We just additionally drew an ice column and fans to the experimental data to obtain a practical model, easier to compare with nature. We decided to mix interpretational and experimental data in this final diagram of the manuscript to constitute a synthetic model, easy to understand and usable for any glacial geologists or glaciologists. We already added in the revised version a figure (Fig. 2) displaying “real” photo of the experiment that will help the reader to better apprehend the model.

Referee #2 : Anonymous reviewer

General comments

This paper describes an analog experimental model for ice flow over sediments and water, and uses the results of the experiments to describe a transitory lifecycle of an ice stream. The paper is short; it identifies some of the known features of modern and paleo ice streams, discusses the combination of conditions that are thought to play a role in the dynamics of ice streams, describes the experimental setup, the results of an experiment, and the inferred ‘lifecycle’ behaviour of an ice stream. The experimental approach is quite novel (though not without precedent; notably the paper of Catania & Paola (2001) is absent from the references and deserves comment) and I think it is welcome. You might also reference the laboratory work of Kowal & Worster (2015), which has some similar results. The setup appears to be quite sophisticated, allowing detailed mapping of elevation changes and velocities. There therefore appears to be considerable scope with this approach. However, the current manuscript is somewhat lacking in detail and I think there needs to be more scientific discussion about the extent to

which the experiment does and does not represent the real world. There also is relatively little data presented on the detailed measurements that have evidently been taken. At present, it reads like a re-hash of a submission to Nature, and I think it needs a bit of expansion to fill in some details for the more discerning reader. The paper is nevertheless well written and interesting, and I think with improvements it can be a valuable contribution to the literature.

As pointed out by both reviewers, we agree that the current version is lacking some data especially on tunnel valleys size and a section specifically dedicated to discuss the limitations of this model. This model do not and cannot simulate the whole complexity of a natural system, we therefore added some clarifications throughout the manuscript (cf. replies to referee 1's comments) on how this model works but also on how some of the experimental parameters can potentially alter the model validity.

Specific comments:

The experimental approach is advocated partly on the basis that numerical modelling and field observations are not able to include all the coupled components of the ice stream, sediment, water system. However, there is almost no discussion given to the drawbacks of an experimental approach; in particular, the issues of things that are missing (the analog 'ice' does not change phase for example), and the extent to which the processes can be scaled down. There should be more attention given to this. For example, what is the Reynolds number of the subglacial water flow? Are the dimensions of the 'tunnel valleys' that form comparable to real tunnel valleys (relative to ice thickness, say), and does the grain size of the sand not have some effect.

We agree with the reviewer that the lack of discussion on the drawbacks of an experimental approach is an issue. We suggest to add a section within the methods to mention the process that are not simulated in the model (cf. section, 2.3. "Scaling and limitations").

How was the flow-rate of water to be injected chosen, and are the results sensitive to this? Is it realistic? (In terms of water flux as compared to ice flux, say). How is it decided when to start injecting the water? Does this make a difference?

The flow rate of water is calculated so that water pressure is exceeding the combined weight of the silicon and sand layers. This is calculated beforehand to initiate water flow at the silicon-bed interface. The flow rate of water is not realistic against the silicon flux because it would require a perfect scaling which is impossible from a material point of view. We added these details in the methods part of the revised version of the manuscript:

"Water discharge is calculated beforehand so that water pressure exceeds the combined weight of the sand and silicon layers. The injection of water starts when the silicon layer reaches the dimensions we fixed for every experiment (15 cm radius and 3 cm thickness) and a perfect transparency. Once injected, water flow is divided into a Darcy flow within the substratum and a flow at the silicon/substratum interface. The water flowing at the silicon/substratum interface originates from a pipe forming at the injector once water pressure exceeds the cumulative pressure of the silicon and sand layers. The ratio between the Darcy flow and the flow at the silicon/substratum interface is inferred from computations of the water discharge flowing through the pipe based on the substratum properties and the input discharge. We estimate that 75% of the input discharge is transferred as Darcy flow in the substratum and 25% of the input discharge along the silicon/substratum interface."

How much of the water flow is through the permeable sediments and how much in a film at the sediment/silicon interface? How thick is the water layer? Are the sediments in suspension or carried as bedload?

We have estimated that 75% of the water is flowing through the permeable sediments and 25% at the interface. This ratio have been inferred from computations of the water flowing through the pipe forming over the injector. We add these details in the methods part (See the modifications to answer the last comment). Water layer thickness (over a 1 mm inside the water pocket) can be inferred from the vertical uplift maps in Figure 3.

Only one particular experiment is described in any detail. It is not clear how repeatable this is except for the comment on 1190 that the observed lifecycle is the same for 12 identical runs; but it is hard to imagine that the development of the three ‘tunnel valleys’ is exactly the same each time. Is there really always two stages of streaming? Do they always appear on the same sides of the experiment? How different are the plots in figure 3 between different experiments (in terms of peak velocity for example)? There should be more discussion of the other experiments.

We agree that it might be confusing to state that every experiment leads to the same outcome. It is true that every experiment lead to the development, migration and drainage of a water pocket that subsequently trigger ice streaming. For every experiment, the drainage phase is followed by the development of tunnel valleys that causes ice stream deceleration. However, there is variability in the amount and size of tunnel valleys we form between the different experiments. We also notice that there is not always a lateral migration of the ice stream when the drainage efficiency of the tunnel valley system is sufficiently high to prevent storage/drainage of a second water pocket.

A new paragraph (section 3.2 “Experimental reproducibility and variability”) discussing the range of experimental results has been added to the revised version of the manuscript.

Figure 2. It is not completely clear what is shown in the first column, and the color scale chosen is not particularly suited to showing elevation changes (e.g. it is quite unclear where zero is). Given that there are negative values, this is presumably an elevation change from some reference? What is taken as the reference, given that the silicon is anyway spreading (and presumably lowering?) before injection starts?

The surface elevation maps are made from a reference picture taken just before the injection (few seconds before the injection). We suggest to add the 0 on the color scale for the vertical displacement maps and to add information on how silicon flow velocities and elevation map are interpolated within the methods section:

“The monitoring of every UV marker positions (in both horizontal vertical plans) through time was used to produce velocity and vertical displacement maps. Vertical displacement maps are interpolated from the subtraction of the DEM at time t with the DEM generated from the photographs taken a few seconds before the injection. Velocity maps are interpolated from the subtraction of the position of every marker at time t with the position of the same markers at the previous stage.”

The surge of the Variegated glacier referenced on line 219 was, as I understand it, accompanied by a decrease in the outlet discharge of subglacial water rather than an increase. A subsequent increase in discharge, with the development of a more efficient drainage system, accompanied the termination of the surge. So I am not sure this is quite the same behaviour as seen in your experiments.

We agree with the reviewer that the study of Kamb (1985) on the Variegated glacier is not the best fit to compare with our results as the surge termination is associated with an outburst flood. We suggest to switch this reference with the study of Anderson et al., 2005 which observe and measure an ice flow acceleration following the outburst flood of the Hidden Creek Lake, Alaska.

The slow-down of the ice stream is attributed to a lowering of subglacial water pressure together with the growth of tunnel valleys, but presumably in the experiments there is also an influence of the changing silicon geometry which is driving the silicon flow. The surface is lowered over the central part of the dome and the driving stress is therefore reduced. What is the evidence that the ageing of the ice stream is not simply due to this effect? (which is also present in the real ice-stream problem too).

Broadly speaking it is true that the change of silicon geometry will inevitably slow down the silicon flow as the amount of silicon which is available to flow progressively decreases. However, we described an experiment where all processes of ice streaming and tunnel valley formation occur in a short period of time (30 min). Hence, the progressive decay of the ice stream related to ice thinning is negligible here. However, if we consider a context where the silicon layer thins significantly, water pressure would decrease similarly to nature. Indeed, the size of the pipe forming when we inject water within the substratum is dependent on the thickness of the sand and silicon layers. Hence, if the silicon layer thickness decreases the pipe circumference would significantly increase, which conduct to an increase of the water discharge flowing at the substratum/interface and a decrease of water pressure. Consequently, the stagnation of the ice stream in our 12 experiments is always achieved despite an increase of water flow at the silicon/substratum interface.

1 **Modelled subglacial floods and tunnel valleys control the lifecycle of transitory ice**
2 **streams**

3 T. Lelandais^(a*), É. Ravier^(a), S. Pochat^(b), O. Bourgeois^(b), C.D. Clark^(c), R. Mourgues^(a) and P.
4 Strzeczynski^(a)

5 ^(a) Laboratoire de Planétologie et Géodynamique, UMR 6112, CNRS, Le Mans Université,
6 Avenue Olivier Messiaen, 72085 Le Mans Cedex 9, France

7 ^(b) Laboratoire de Planétologie et Géodynamique, UMR 6112, CNRS, Université de Nantes, 2
8 rue de la Houssinière, BP 92208, 44322 Nantes Cedex 3, France

9 ^(c) Department of Geography, University of Sheffield, Sheffield, UK

10 ^(*) Correspondence to: thomas.lelandais@univ-lemans.fr

Code de champ modifié

11 **Ice streams are corridors of fast-flowing ice that control mass transfers from continental**
12 **ice sheets to oceans. Their flow speeds are known to accelerate and decelerate, their**
13 **activity to switch on and off, and even their locations to shift entirely. Our analogue**
14 **physical experiments reveal that a lifecycle incorporating evolving subglacial meltwater**
15 **routing and bed erosion can govern this complex transitory behaviour. The modelled ice**
16 **streams switch on and accelerate when subglacial water pockets drain as marginal**
17 **outburst floods (basal decoupling). Then they decelerate when the lubricating water**
18 **drainage system spontaneously organising itself into channels that create tunnel valleys**
19 **(partial basal recoupling). The ice streams surge or jump in location when these water**
20 **drainage systems maintain low discharge but they ultimately switch off when tunnel**
21 **valleys have expanded to develop efficient drainage systems. Beyond reconciling**
22 **previously disconnected observations of modern and ancient ice streams into a single**
23 **lifecycle, the modelling suggests that tunnel valley development may be crucial in**
24 **stabilising portions of ice sheets during periods of climate change.**

25 **Keywords:** ice streams, experimental modelling, subglacial meltwater drainage, tunnel
26 valleys, subglacial outburst floods

27 1. Introduction

28 Continental ice sheets currently store the equivalent of a 65 m thick global equivalent
29 water layer and have been major contributors to the nearly 85 mm global sea level rise measured
30 between 1993 and 2017 (Vaughan et al., 2013; Beckley et al., 2015). The mass transfer from
31 these ice sheets to the ocean is spatially heterogeneous: approximately 80% of the ice discharge
32 is focused in a finite number of ice streams, which act as preferential drainage pathways for
33 meltwater also (Bamber et al., 2000; Bennett, 2003).

34 Modern and ancient ice streams are typically hundreds of kilometres long and a few
35 kilometres to tens of kilometres wide, with ice velocities of the order 10^2 to 10^4 m/yr. ~~Despite~~
36 ~~the fact that they occurred and occur in all former and modern ice sheets, their initiation, and~~
37 ~~the controls on their dynamics and evolution remain debated~~ They occur in all known ice sheets,
38 ~~but why and where they initiate, and the controls on their dynamics and evolution remain~~
39 ~~debated.~~ Numerical modelling suggests that ice flow might self-organise into regularly-spaced

40 ice streams as a consequence of thermomechanical feedbacks within ice (Payne and
41 Dongelmans, 1997; Hindmarsh, 2009) or because of inherent instability of thin subglacial
42 meltwater films (Kyrke-Smith et al., 2014). Numerous observations however, have highlighted
43 preferential location of ice streams at sites of specific bed properties such as in topographic
44 troughs, over areas of soft sedimentary geology, zones of higher geothermal heat flux or as a
45 consequence of where subglacial meltwater is routed (Winsborrow et al., 2010; Kleiner and
46 Humbert, 2014). These viewpoints might not be mutually exclusive if self-organisation into
47 regularly-spaced streams is the primary control but that it is strongly mediated by local bed
48 templates (e.g. troughs) or events (meltwater drainage) that initiate or anchor streams in certain
49 locations. Exploring this hypothesis by numerical modelling has not yet been achieved because
50 of uncertainties in how to formulate basal ice flow in relation to bed friction, and due to
51 challenges of including all potentially relevant processes, especially so for subglacial water
52 flow (Flowers, 2015).

53 Observations of spatial and temporal variations in the activity of ice streams against
54 fluctuations in their subglacial hydrology ~~indicates~~~~suggest~~ that the style and flux of water
55 drainage is a major component driving change. Examples include: reorganisation of subglacial
56 drainage systems (Elsworth and Suckale, 2016), subglacial water piracy (Vaughan et al., 2008;
57 Carter et al., 2013), and development and migration of transient subglacial water pockets (Gray
58 et al., 2005; Peters et al., 2007; Siegfried et al., 2016). However, these variations have been
59 observed or inferred independently, at different places and on yearly timescales, thus limiting
60 our understanding of the true role of the subglacial hydrology as primary or secondary drivers
61 of ice stream changes. In this paper, we circumvent the challenge of numerically modelling ice
62 stream initiation and dynamics, including subglacial water drainage, by exploiting a physical
63 laboratory approach that simultaneously combines ~~ice~~-silicon flow, water drainage and bed
64 erosion.

65 Connections between ice stream activity and subglacial hydrology are supported by the
66 occurrence of geomorphic markers of meltwater drainage on ancient ice stream beds (e.g.
67 meltwater channels, tunnel valleys, eskers) (~~Patterson, 1997~~; Margold et al., 2015; Livingstone
68 et al., 2016; ~~Patterson, 1997~~). Among these markers, tunnel valleys deserve specific attention
69 ~~network~~-because they have high discharge capacities and, as such, may be major contributors
70 to the release of meltwater and sediment to the ocean ~~and~~; ~~they~~ may ~~also~~-promote ice sheet
71 stability by reducing the lubricating effect of high basal water pressure. These valleys are
72 elongated and over-deepened hollows, ranging from a few kilometres to hundreds of kilometres
73 long, from hundreds metres to several kilometres wide and from meters to hundreds of metres
74 deep. Their initiation is generally attributed to subglacial meltwater erosion but their
75 development processes (in time and space) and ~~on~~-their relationship to ice streaming are still
76 debated. Indeed, ice streams commonly operate because of high basal water pressure while the
77 development of a tunnel valleys system generally leads to ~~enhance~~s drainage efficiency and
78 basal water pressure reduction (Engelhardt et al., 1990; ~~Marczinek and Piotrowski, 2006~~;
79 Kyrke-Smith et al., 2014; ~~Marczinek and Piotrowski, 2006~~).

80 Several field studies have already suggested a connection between catastrophic glacial
81 outburst floods at ice sheets margins and a suite of events involving ice streaming, tunnel valley

82 development and stagnation of the ice margin. (Jørgensen and Piotrowski, 2003; Alley et al.,
83 2006; Hooke and Jennings, 2006; Bell et al., 2007; Bell et al., 2007; Hooke and Jennings, 2006;
84 Jørgensen and Piotrowski, 2003; Alley et al., 2006). Such outburst floods can profoundly and
85 rapidly alter the oceanic environment by transferring considerable amounts of ice, freshwater,
86 and sediment from continents to oceans (Evatt et al., 2006). The suspected connection between
87 ice streams, tunnel valleys and outburst floods have never been observed or modelled however.

88 Here, we describe the results of a physical experiment performed with an innovative
89 analogue modelling device that provides simultaneous constraints on ice flow, subglacial
90 meltwater drainage, subglacial sediment transport and subglacial landform development
91 (Lelandais et al., 2016; Fig. 1). We propose that the location and initiation of ice streams might
92 arise from subglacial meltwater pocket migration and drainage pathways and that the evolution
93 of ice stream dynamics is ~~subsequently~~ ~~latter~~ controlled by subglacial drainage reorganization
94 and tunnel valleys development. This study reconciles into a single story several detached
95 inferences, derived from observations at different timescales and at different places on modern
96 and ancient ice streams.

97 2. Experimental ice stream model

98 Ice stream dynamics are controlled by various processes that act at different ~~spatial~~ ~~space~~
99 and ~~temporal~~ ~~time~~ scales; they also involve several components with complex thermo-
100 mechanical behaviours (ice, water, till, bedrock) (Paterson, 1994). Considering all these
101 processes and components simultaneously, together with processes of subglacial erosion, is thus
102 a challenge for numerical computational modelling (Fowler and Johnson, 1995; Marshall, 2005;
103 Bingham et al., 2010). ~~Based on this statement,~~ ~~s~~Some attempts in analogue modelling have
104 been made to improve our knowledge on subglacial erosional processes by meltwater (Catania
105 and Paola, 2001) or gravity current instabilities produced by lubrication (Kowal and Worster,
106 2015). To combine ice flow dynamics and erosional aspects in a single model, we designed an
107 alternative experimental approach that allows simultaneous modelling of ice flow, subglacial
108 hydrology and sedimentary/geomorphic processes. With all the precautions ~~of use~~ ~~inherent~~ ~~of~~
109 ~~in using~~ analogue modelling, our experiments reproduce morphologies and dynamics that
110 compare well with subglacial landforms and ice stream dynamics despite some differences ~~of~~
111 ~~in~~ spatial and ~~time~~ ~~temporal~~ scales and a number of active processes (e.g. Paola et al., 2009).

112 2.1. Experimental apparatus

113 The model is set in a glass box (70 cm long, 70 cm wide and 5 cm deep) (Fig. 1). A 5 cm thick,
114 flat, horizontal, permeable and erodible substratum, made of sand ($d_{50}=100\ \mu\text{m}$) saturated with
115 pure water and compacted to ensure homogeneous values for its density ($\rho_{\text{bulk}} = 2000\ \text{kg/m}^3$),
116 porosity ($\Phi = 41\ \%$) and permeability ($K = 10^{-4}\ \text{m/s}$), rests on the box floor. The ice sheet
117 portion is modelled with a 3 cm thick layer of viscous ($\eta = 5 \cdot 10^4\ \text{Pa s}$) and transparent but
118 refractive ($n = 1.47$) silicon putty placed on the substratum. The model is not designed to
119 simulate an entire ice sheet. The silicon layer is circular in plan view (radius = 15 cm) to avoid
120 lateral boundary effects on silicon flow~~The model is not designed to simulate an entire ice sheet~~
121 ~~and; it is also circular in plan view (radius = 15 cm) to avoid lateral boundary effects on silicon~~
122 ~~flow~~. Subglacial meltwater production is simulated by injection of water with a punctual

123 injector, 4 mm in radius, placed at a depth of 1.8 cm in the substratum and connected to a pump
124 (Fig. 1). The injector is located below the centre of the silicon layer to be consistent with the
125 circular geometry of the experiment. The water discharge is constant ($1.5 \text{ dm}^3/\text{h}$) over the
126 duration of the experiment and generates water flow at the silicon-substratum interface and
127 within the substratum. Water discharge is calculated beforehand so that water pressure exceeds
128 the combined weight of the sand and silicon layers. The injection of water starts when the
129 silicon layer reaches the dimensions we fixed for every experiment (15 cm radius and 3 cm
130 thickness) and a perfect transparency. Once injected, water flow is divided into a Darcy flow
131 within the substratum and a flow at the silicon/substratum interface. The water flowing at the
132 silicon/substratum interface originates from a pipe forming at the injector once water pressure
133 exceeds the cumulative pressure of the silicon and sand layers. The ratio between the Darcy
134 flow and the flow at the silicon/substratum interface is inferred from computations of the water
135 discharge flowing through the pipe based on the substratum properties and the input discharge.
136 We estimate that 75% of the input discharge is transferred as Darcy flow in the substratum and
137 25% of the input discharge along the silicon/substratum interface.

138 2.2. Acquisition process and post-processing

139 In order to monitor the development of landforms on the substratum, we use six
140 synchronised cameras equidistant from the experiment centre (Fig. 1) taking photographs of the
141 experiment every 5 seconds. Two cameras (orange on Fig. 1) cover the whole extent of the
142 experiment and four cameras (blue on Fig. 1) focus on specific regions to obtain higher
143 resolution images. These cameras take simultaneous pictures with differing positions and
144 orientations. Digital elevation models of the silicon surface and of the substratum are derived
145 from these images by photogrammetry. The ultimate stage of the experiment is to remove
146 distortions due to light refraction through the silicon putty and apply corrections to the
147 substratum topography. This treatment is achieved using a custom algorithm able to evaluate
148 the gap between the measured altitude and the real altitude of each pixel of the DEM (cf detailed
149 post-treatment methods in Lelandais et al., 2016). Tests performed on previously known
150 topographies show that the vertical precision of the retrieved digital elevation models is better
151 than 10^{-1} mm .

152 The flow velocity of the silicon layer is monitored near its base (V_{base}), at mid-depth (V_{mid}) and
153 at its surface (V_{surface}), with an additional camera placed over the centre of the experiment (green
154 on Fig. 1). For that purpose, the camera records the position on pictures taken at regular time
155 intervals in ultraviolet (UV) of 180 UV paint drops (1 mm in radius) placed at 1 mm above the
156 base, at mid-depth and at the surface of the silicon layer (Figs. 1, S1 Fig. 1 and Fig. S4). The
157 monitoring of every UV marker positions (in both horizontal-vertical plans) through time was
158 used to produce velocity and vertical displacement maps. Vertical displacement maps are
159 interpolated from the subtraction of the DEM at time t with the DEM generated from the
160 photographs taken a few seconds before the injection. Velocity maps are interpolated from the
161 subtraction of the position of every marker at time t with the position of the same markers at
162 the previous stage. These passive markers are transparent at visible wavelengths and do not
163 alter pictures of the substratum taken through the silicon cap. They represent less than 0.5% of
164 the silicon layer in volume and tests have shown that they do not affect its overall rheological

165 behaviour. Uncertainties in the measured position of markers on images are less than one pixel
166 in size (i.e. less than 10^{-1} mm), thus uncertainties in the derived velocities are comprised between
167 $5 \cdot 10^{-4}$ and $2 \cdot 10^{-3}$ mm/s, depending on the time interval between photographs.

168



169 2.3. Scaling and limitations


170 In this study, we focus our attention on the relations between subglacial water flow,
171 subglacial erosion and ice flow using an experiment approach. Considering that in our model
172 meltwater is simulated by an injection of water, the rules of a classical scaling where the model
173 is a perfect miniaturisation of nature are not practical (Paola et al., 2009). In this perspective,
174 we base the scaling of our model on the displacement of the natural ice and experimental silicon
175 margins through time. We use a unit-free speed ratio between the silicon/ice margin velocity
176 and the incision rate of experimental/natural tunnel valleys. In this way, the complexity of the
177 relations between subglacial hydrology, subglacial erosion and ice flow, which is one of the
178 main issue in numerical modelling, is included in the velocity values. The scaling attest that the
179 value of the ratio between margin velocity and incision rate of tunnel valleys in the experiment
180 fall within the field validity defined by the range of natural settings (full details in Lelandais et
181 al., 2016). The main scaling limit regards the viscosity ratios between glacier ice, silicon putty
182 and water. The size of the experimental ice stream, being partly controlled by the high silicon
183 viscosity, may be underestimated compared to the size of modelled tunnel valleys.

184 ~~Considering that meltwater is here simulated by an injection of water, the rules of a~~
185 ~~classical scaling where the model is a miniaturisation of nature are not practical (Paola et al.,~~
186 ~~2009). Subglacial water drainage is generally controlled by fluctuations in locations of ice sheet~~
187 ~~margins. Similarly, in our experiments, the silicon putty margin controls the water pressure~~
188 ~~gradient. In this perspective, we base the scaling on the displacement of the natural ice and~~
189 ~~experimental silicon margins through time. We use a unit free speed ratio between the~~
190 ~~silicon/ice margin velocity and the incision rate of experimental/natural tunnel valleys. The~~
191 ~~scaling is designed to ensure that the value of the ratio between margin velocity and incision~~
192 ~~rate of tunnel valleys in the experiment equals its value in natural. The projection of the minimal~~
193 ~~and maximal experimental speed ratios on the field of possible natural speed ratios highlights~~
194 ~~the field of validity of the experiments and defines the range of natural settings we can~~
195 ~~reproduce experimentally (full details in Lelandais et al., 2016). The main scaling limit regards~~
196 ~~the viscosity ratios between glacier ice, silicon putty and water. The size of the experimental~~
197 ~~ice stream, being partly controlled by the high silicon viscosity, may be underestimated~~
198 ~~compared to the size of modelled tunnel valleys.~~

199 Considering that our model is a simplification of nature, we cannot simulate the whole
200 complexity of the nature processes~~Considering that our model is a simplification of nature, we~~
201 ~~cannot simulate its whole complexity.~~ In contrast with ice, the commercial silicon putty we use
202 (Dow Corning, SGM36) is impermeable, newtonian, isotropic, and its viscosity is nearly
203 independent of temperature between 10 and 30°C. Therefore, rheological softening ~~of ice~~ with
204 strain rate, temperature, anisotropy, and meltwater content (e.g. Bingham et al., 2010) cannot
205 be fully reproduced. The silicon putty cannot reproduce the ice/water phase transition either,
206 supporting-requiring the use of punctual water injection in the experiment. This punctual


207 injection does not simulate the mosaic of meltwater production regions existing beneath
208 glaciers or the episodic input from supraglacial/englacial meltwater reservoirs. Experimental
209 meltwater routing is predominantly controlled by the water discharge we inject in our system
210 and therefore differs from parameters controlling hydrology in glacial systems. Subglacial
211 meltwater routing is indeed controlled by the ice surface slope, the bed topography, and the
212 glacier mass balance (Röthlisberger and Lang, 1987). The ice surface slope controls
213 potentiometric surfaces, generally guiding subglacial water flow parallel to ice sheet surfaces
214 (Glen, 1954; Shreve, 1972; Fountain and Walder, 1998). Finally, the substratum we use is
215 homogeneous, flat and composed of a well-sorted mixture of sand-sized grains. This model,
216 designed to decipher the interaction between subglacial hydrology and ice dynamics, hinders
217 the influence of bed topography and geology (especially the influence of subglacial till)
218 (Winsborrow et al., 2010). The deformation of the subglacial till and its complex rheological
219 behavior is known to promote ice streaming (Alley et al., 1987), modify the subglacial
220 hydrology, and alter the size of tunnel valleys. The development of an analogue material scaled
221 to reproduce subglacial till characteristics is extremely difficult so we did not try to include the
222 equivalent of a till layer in the experiment. We thus assume that the velocity contrasts observed
223 in the experiment are ~~thus~~ likely to be amplified in natural ice sheets, by the complex
224 rheological behaviour of ice and till. This may lead to the development of narrower ice streams
225 with higher relative velocities and sharper lateral shear margins in natural ice sheets than in the
226 experiment (Raymond, 1987; Perol et al., 2015).





228

229 **Figure 1.** Description of the analogue device used in this study. a, Overview of the analogue device.
 230 The analogue device consists in a 70 cm long, 70 cm wide and 5 cm deep glass box filled with saturated
 231 and compacted sand simulating the substratum. The ice sheet portion is simulated by a circular layer of
 232 silicon putty containing 3 levels of UV markers. Meltwater production is simulated by a central and
 233 punctual injection of pure water within the substratum. Five synchronized cameras placed above the
 234 silicon putty (in blue) focus on the tunnel valley system and are used to produce digital elevation models
 235 by photogrammetry. Another camera (in orange) takes overview photographs of the analogue device to
 236 follow the progress of the whole experiment. A last camera (in green) is positioned at the vertical of the
 237 silicon layer centre and is configured to take high-resolution photographs in black light of the UV
 238 markers (illuminated with two lateral UV led lights). b, Cross-sectional profile of the analogue device
 239 displaying the position of the UV markers and the physical characteristics of both the substratum and
 240 the silicon layer.



241
 242 **Figure 2.** Temporal evolution of the experiment seen on raw photographs. a. Formation of a water
 243 pocket. b. Migration of the water pocket. c. Marginal drainage of the water pocket and onset of the
 244 silicon stream. d. Development of two tunnel valleys (TV1 and TV2). e. Drainage of a second water
 245 pocket and silicon stream migration. f. Development of a new generation of tunnel valleys (TV3)
 246 and silicon stream decay. Silicon flow velocity and silicon surface displacement maps corresponding to the
 247 six stages described here are presented in Figure 3.

248 3. Experimental results

249 3.1. Stage-by-stage experimental progress

250 This experiment was repeated 12 times with identical input parameters (a 30 mm-thick silicon
251 layer of 150 mm radius; constant water input of 1.5 dm³/h during 1800 s). After an initial
252 identical state, a six-stage ice stream lifecycle linking outburst flooding, transitory ice
253 streaming, and tunnel valleys development has been observed for all these simulations (Figs. 2,
254 3, Fig. 3a-f, Fig. 6).


255 Initial state (Fig. S2). As long as no water is injected in the substratum, the silicon layer spreads
256 under its own weight and displays the typical parabolic surface profile of an ice sheet. It
257 increases in diameter and decreases in thickness with time, thus producing a radial pattern of
258 horizontal velocities, which increase in magnitude from the centre ($V_{\text{surface}} < 3 \cdot 10^{-3}$ mm/s) to the
259 margin ($V_{\text{surface}} = 8 \cdot 10^{-3}$ mm/s) (Fig. S2). V_{base} is close to 0 over the full extent of the silicon
260 layer ($\frac{V_{\text{base}}}{V_{\text{surface}}} \sim 0\%$), indicating coupling with the substratum. The silicon flow pattern changes
261 when meltwater production is simulated by injecting water at a constant discharge (1.5 dm³/h),
262 beneath the silicon layer.

263 Stage 1 (Figs. 2a, 3a, Fig. 2a-3a). A water pocket grows below the centre of the silicon layer and
264 raises its surface by 2 mm. Above the water pocket, the silicon accelerates ($V_{\text{surface}} \geq 35 \cdot 10^{-3}$
265 mm/s), and is decoupled from the substratum ($\frac{V_{\text{base}}}{V_{\text{surface}}} = 75$ to 80%). Below the rest of the
266 silicon layer, lower velocities ($V_{\text{surface}} = 8 \cdot 10^{-3}$ mm/s, $\frac{V_{\text{base}}}{V_{\text{surface}}} = 40$ to 50%) indicate higher
267 basal friction. These results are consistent with inferences that meltwater ponding can form
268 pressurised subglacial water pockets associated with basal decoupling, surface uplift, and ice
269 flow acceleration in natural ice sheets (e.g. Hanson et al., 1998; Elsworth and Suckale, 2016;
270 Livingstone et al., 2016). In the experiment however, these effects are restricted to an
271 approximately circular region and are not sufficient to produce channelised ice streaming.

272 Stage 2 (Figs. 2b, 3b, Fig. 2b-3b). The water pocket expands and migrates towards the margin
273 of the silicon layer. The lack of channels incised in the substratum indicates that this
274 displacement occurs as distributed water drainage without any basal erosion. In the silicon
275 layer, the region of surface uplift, basal decoupling and acceleration ($V_{\text{surface}} = 18 \cdot 10^{-3}$ mm/s,
276 $\frac{V_{\text{base}}}{V_{\text{surface}}} = 75$ to 85%) expands and migrates downstream with the water pocket. Similar
277 migrations of pressurised subglacial water pockets have been observed or inferred under
278 modern and ancient ice sheets (Fricker et al., 2007; Carter et al., 2017), sometimes associated
279 with migrations of regions of ice surface uplift and ice flow acceleration (Bell et al., 2007;
280 Stearns et al., 2008; Siegfried et al., 2016). The experiment ~~suggests~~ indicates that the migration
281 of water pockets at the ice-bed interface can contribute to the emergence of ice streams.

282

283



284
 285 **Figure 3.** Temporal evolution of the experiment. a, Formation of water pocket, uplift of silicon surface
 286 uplift and acceleration. b, Migration of water pocket and overlying region of uplift and accelerated flow.
 287 c, Marginal drainage of water pocket and onset of silicon streaming. d, Tunnel valley development and
 288 silicon stream deceleration. e, Formation, migration and marginal drainage of a new water pocket,


289 development of a second silicon stream and of a new tunnel valley. f, Decay of the second silicon stream.
290 From left to right: (i) maps of vertical displacements of silicon layer surface, (ii) maps of horizontal
291 velocity at silicon ~~cap-layer~~ surface, (iii) cross-sectional velocity profiles (absolute velocity on right
292 axis, velocity normalised by background velocity on left axis, profile locations indicated by white lines
293 A-B on maps), (iv) vertical velocity profiles for silicon stream (red profiles, locations labelled ~~1-2~~
294 on maps) and for region opposed to silicon stream (black profiles, locations labelled ~~2-1~~ on maps).

295 Stage 3 (Figs. 2c, 3c Fig. 2e-3e). When the water pocket reaches the margin of the silicon layer,
296 it drains suddenly as a sheet flow. This marginal outburst flood is still fed by distributed
297 drainage and conveys sand particles eroded from the substratum towards a low-angle marginal
298 sedimentary fan (up to 40 mm long, 30 mm wide and 0.3 mm thick; Fig. S3). Simultaneously,
299 the silicon flow focuses in a stream (200 mm wide at the margin) that propagates upstream from
300 the silicon margin to the water injection area. This stream immediately peaks in velocity (V_{surface}
301 = $80 \cdot 10^{-3}$ mm/s, 16 times higher than the surrounding silicon) and is entirely decoupled from its
302 substratum ($\frac{V_{\text{base}}}{V_{\text{surface}}} > 90\%$). Although similar relations between outburst floods and ice flow
303 accelerations have been suspected in modern (Alley et al., 2006; Bell et al., 2007; Stearns et al.,
304 2008) and ~~past-former~~ (Livingstone et al., 2016) ice sheets, they have been documented for
305 valley glaciers only (e.g. Anderson et al., 2005): In these regions, they can produce sudden
306 meltwater discharges that exceed the capacity of distributed subglacial meltwater drainages and
307 promote basal decoupling and ice flow acceleration~~there, they can produce sudden meltwater~~
308 ~~discharges that exceed the capacity of distributed subglacial meltwater drainages and promote~~
309 ~~basal decoupling and ice flow acceleration~~ (e.g. Magnússon et al., 2007). The experiment
310 confirms that outburst floods can promote basal decoupling and trigger ice streaming in ice
311 sheets (Fowler and Johnson, 1995).

312 Stage 4 (Fig. 2d, 3d Fig. 2d-3d). The distributed subglacial drainage system starts to channelise:
313 two valleys (TV1 and TV2) appear below the margin of the silicon layer and gradually expand
314 by regressive erosion of the substratum. At this stage, TV1 is 30 mm long, 12 mm wide and 0.5
315 mm deep; TV2 is 80 mm long, 10 mm wide and 0.5 mm deep. These valleys, with their constant
316 widths, undulating long profiles and radial distribution, are analogue to natural tunnel valleys
317 in their dimensions, shapes, and spatial organization (Lelandais et al., 2016; Fig. 4). They are
318 fed by distributed water drainage. The sand eroded from the substratum transits through these
319 valleys and accumulates in high-angle marginal sedimentary fans, higher in elevation than the
320 valley floors (TV1 fan is up to 27 mm long, 30 mm wide and 0.5 mm thick; TV2 fan is up to
321 20 mm long, 24 mm wide and 1 mm thick; Fig. S3). In response to progressive channelisation
322 of the water drainage into the expanding valleys, the silicon stream narrows and slows down
323 (120 mm wide at the margin; $V_{\text{surface}} = 24 \cdot 10^{-3}$ mm/s). The silicon stream, still channelised, is
324 still flowing 8 times faster than the rest of the silicon layer and is still decoupled from the
325 substratum ($\frac{V_{\text{base}}}{V_{\text{surface}}} > 85\%$). These results are consistent with inferences that channelisation of
326 hitherto distributed subglacial water drainage systems can occur and reduce ice flow velocity
327 after outburst floods (Kamb, 1987; Retzlaff and Bentley, 1993; Magnússon et al.,
328 2007; Magnússon et al., 2007; Kamb, 1987; Retzlaff and Bentley, 1993), and can be responsible
329 for narrowing and deceleration of ice streams (Raymond, 1987; Retzlaff and Bentley, 1993;
330 Catania et al., 2006; Beem et al., 2014; Kim et al., 2016). At this stage of the experiment, this

331 transition, which corresponds to the initiation of tunnel valleys, is not sufficient to stop ice
332 streaming however.

333




334

335 **Figure 4.** Digital Elevation Model (DEM) of an experimental tunnel valley and its associated
336 longitudinal profile. a, Snapshot of the tunnel valley system. b, DEM of the tunnel valley corresponding
337 to the one highlighted by a dashed box in a. c, Undulating longitudinal profile of the tunnel valley bottom
338 extracted from the DEM in the dashed box shown in b.

339 Stage 5 (Figs. 2e, 3e Fig. 2e-3e). A new transient water pocket grows below the silicon layer,
340 migrates and drains as an outburst flood, thus forming a new low-angle marginal sedimentary
341 fan with a lateral offset of 4 cm with respect to TV1. This induces the activation of a second
342 stream ($V_{\text{surface}} = 40 \cdot 10^{-3} \text{ mm s}^{-1}$) decoupled from its substratum ($\frac{V_{\text{base}}}{V_{\text{surface}}} = 80\%$) and the
343 initiation of a new radial valley (TV3), in a hitherto slow-moving region of the silicon cap.
344 Simultaneously, the first silicon stream switches off ($V_{\text{surface}} = 10 \cdot 10^{-3} \text{ mm/s}$), and recouple to
345 its substratum ($\frac{V_{\text{base}}}{V_{\text{surface}}} = 30\%$), but water and sand still flow through TV1 and TV2. At this
346 stage, TV1 is 100 mm long, 8 mm wide and 0.7 mm deep and its fan is up to 21 mm long, 40
347 mm wide, and 1.1 mm thick; ~~TV2 is 80 mm long, 7.5 mm wide, and 0.6 mm deep~~
348 ~~mm long, 0.75 mm deep and 0.6 mm deep~~ and its fan is up to 20 mm long, 28 mm wide, and
349 1.6 mm thick. This result is consistent with inferences that natural ice streams can switch on
350 and off, surge or jump in location in response to changes in subglacial water drainage
351 reorganisation (Catania et al., 2012; Le Brocq et al., 2013; Beem et al., 2014; Hulbe et al.,
352 2016; Beem et al., 2014; Hulbe et al., 2016; Catania et al., 2012; Le Brocq et al., 2013). The
353 experiment further suggests indicates that this complex behaviour is controlled by the growth
354 and migration, in various possible directions, of transient pressurised subglacial water pockets
355 that form successively as long as the discharge capacity of tunnel valleys systems is not
356 sufficient to drain efficiently the available meltwater.

357 Stage 6 (Fig. 2f-3f). Since their initiation, TV1, TV2 and TV3 have progressively increased in
 358 width, depth and length. At this stage TV1 is 100 mm long, 17 mm wide and 1.2 mm deep and
 359 its fan is 28 mm long, 4 mm wide and 1.5 mm high at the maximum; TV2 is 80 mm long, 10
 360 mm wide and 0.8 mm deep and its fan is up to 16 mm long, 23 mm wide and 1.6 mm thick ;
 361 TV3 is 60 mm long, 11 mm wide and 0.55 mm deep and its fan is up to 14 mm long, 23 mm
 362 wide and 0.7 mm thick. Their overall volume and discharge capacity have thus increased (Fig.
 363 5). In response to this increased drainage efficiency, the second stream gradually decays ($V_{surface}$
 364 = $5 \cdot 10^3 \text{ mm} \cdot \text{s}^{-1}$), and recouples to its substratum ($\frac{V_{base}}{V_{surface}} = 35\%$). The silicon layer ultimately
 365 recovers a radial flow pattern (Fig. 3f). This result is consistent with the inference that ice
 366 streams may decelerate and even switch off in response to reduction of subglacial water
 367 pressures when efficient subglacial water drainage systems develop (Retzlaff and Bentley,
 368 1993; Beem et al., 2014; Livingstone et al., 2016; Kim et al., 2016). In the experiment, this
 369 development is governed by the expansion of tunnel valley networks. Large glaciotectonic
 370 thrust masses at the ice margin near tunnel valleys fans are generally assumed to be field
 371 evidences of a fast ice flow stage prior to drainage through tunnel valleys (Hooke and Jennings,
 372 2006).



373
 374 **Figure 5.** Progressive expansion of overall volume of tunnel valleys system vs. velocity of silicon
 375 margin through the experiment. The circled numbers correspond to the six-stages of the proposed ice
 376 stream lifecycle.

377 **3.2. Experimental reproducibility and variability**

378 This experiment has been reproduced 12 times with identical input parameters. We always
 379 observe the same processes and events acting in a similar chronological order : (1) water pocket
 380 forms; (2) water pocket migrates; (3) water pocket drains (outburst flood) and silicon stream
 381 switches on; (4) Tunnel valleys form in response to channelisation-; (5) silicon stream slows
 382 down (↔) and (6) finally switches off (↔) in response to the increase of drainage efficiency
 383 during tunnel valley development . However, despite this consistency in the progress of all

384 simulations we ran, some variability has been detected. We measured different migration rates
385 for the water pocket ranging from 30 s to 80 s that may result from small changes in subglacial
386 topography and in the dynamics of silicon-bed decoupling. Considering a constant water
387 discharge and the characteristics of the experiment, a longer period of migration implies: a
388 longer period of water storage and a bigger water volume released at the silicon margin during
389 the pocket drainage. We therefore recorded peak velocities for water pocket drainage ranging
390 from $6 \cdot 10^{-3}$ to $12 \cdot 10^{-3}$ mm \cdot s $^{-1}$. In response to variations of the water volume drained at the
391 margin and ~~the~~ peak discharge, the maximum width of the silicon stream varies from 120 to
392 250 mm. The magnitude of the outburst flood triggered during water pocket drainage also
393 influences the amount of tunnel valleys that ~~subsequently will later~~ forms during the
394 channelisation stage. A high magnitude outburst flood generates a wider erosion beneath the
395 silicon that will be suitable for the development of multiple tunnel valleys. Hence, the amount
396 of tunnel valleys at the end of the experiments range~~s~~ from 1 to 5 with 1 to 3 tunnel valleys
397 formed simultaneously during the initiation of the channelisation stage. These valleys range
398 from 40 to 120 mm long, 3 to 18 mm wide, and 0.3 to 1.8 mm deep. During tunnel valleys
399 development, the evolution of drainage efficiency varies between the experiments. A relatively
400 inefficient system of tunnel valleys induces upstream water pocket formation. As observed in
401 Figure 3e, the drainage of this belated water pocket may provoke water re-routing ~~behind~~
402 ~~beneath~~ the silicon and subsequent lateral migration of the silicon stream. We counted 0 to 2
403 events of silicon stream migration for single experiments. Finally, the time required to reach
404 the phase of ice stream decay highly depends on the amount of tunnel valleys formed during
405 the experiments and their progressive development. We observed a lifetime for the silicon
406 stream ranging from 500 s to 1700 s, correlated with the evolution of the drainage efficiency
407 during tunnel valleys development.

408 4. Proposed lifecycle of transitory ice streams

409 The experiment demonstrates that, on flat and homogenous beds, ice streams may arise,
410 progress, and decay in response to mechanical interactions between ice flow, subglacial water
411 drainage, and bed erosion. On uneven or heterogeneous beds (not simulated in this model),
412 these interactions may additionally be enhanced or disturbed by spatial variations in the
413 subglacial topography, geology, and geothermal heat flux (e.g. Bentley, 1987; Blankenship et
414 al., 1993; Anandakrishnan et al., 1998; Bourgeois et al., 2000; Winsborrow et al., 2010). The
415 complex rheology of glacial ice and subglacial till (both generally soften with increasing strain
416 rate, temperature, water content, and anisotropy) may also enhance these interactions by
417 increasing velocity contrasts between ice streams and their slower-moving margins. This may
418 lead to the development of narrower ice streams with higher velocities and sharper lateral shear
419 margins in natural ice sheets than in the experiment (Raymond, 1987; Perol et al., 2015).

420 Although the complexity of glacial systems cannot be fully modelled using the present
421 experimental setup, our results highlight the critical connection between ice streams and tunnel
422 valleys. As reviewed in Kehew et al., (2012) and suggested in Ravier et al., (2015) this relation
423 was suspected from the occurrence of tunnel valleys on ancient ice streams beds. However, it
424 raised a contradiction: subglacial meltwater pressures are generally supposed to be high below
425 ice streams (Bennett, 2003) while tunnel valleys are generally assumed to operate at lower water

426 pressures (Marczinek and Piotrowski, 2006). Although speculated from field evidences, our
427 results demonstrate that ice streaming, tunnel valley formation, release of marginal outburst
428 floods and subglacial water drainage reorganization may be interdependent parts of a single ice
429 stream lifecycle that involves temporal changes in subglacial meltwater pressures (Fig. 6).

430 1. Ice stream seeding. A prerequisite to the activation of ice streams is the formation of
431 pressurised subglacial pockets by meltwater ponding in ice sheet hinterlands. Approximately
432 circular regions of surface uplift and accelerated ice flow develop above these transient water
433 pockets.

434 2. Ice stream gestation. Pressurised water pockets migrate downstream by distributed water
435 flow. Regions of surface uplift and accelerated ice flow migrate accordingly.

436 3. Ice stream birth. Once water pockets reach ice sheet margins, they drain as outburst floods.
437 At that time, ice streams switch on, peak in velocity, and propagate towards ice sheet hinterlands
438 as decoupled corridors of accelerated ice flow underlain by pressurised distributed water
439 drainage.

440 4. Ice stream aging. Subglacial water drainage then channelises gradually: tunnel valleys fed
441 by pressurised distributed drainage start to form at ice stream fronts. Subsequent expansion of
442 tunnel valleys by regressive erosion progressively increases their overall discharge capacity,
443 lowers subglacial water pressures and provokes gradual ice stream recoupling and deceleration.
444 The response of ice stream dynamics to drainage channelisation and tunnel valley development
445 might be underestimated due to the high erodability of the subglacial bed used in the
446 experiment.

447 5. Ice stream rebirth (relocation or surge). As long as tunnel valley systems keep low drainage
448 capacities, successive pressurised subglacial water pockets can form, migrate, and drain as
449 marginal outburst floods. On even and homogeneous ice sheet beds, the subglacial water
450 drainage is controlled by the surface topography of ice sheets: subtle temporal changes in this
451 topography may thus be able to produce consecutive generations of ice streams and tunnel
452 valleys at different locations and with different flow directions. These jumps in locations and
453 directions may be responsible for the formation of independent, but sometimes intersecting, ice
454 streams corridors and tunnel valleys networks on some ancient ice sheet beds ([Fowler and
455 Johnson, 1995](#); [Jørgensen and Piotrowski, 2003](#); ~~[Jørgensen and Piotrowski, 2003](#)~~; ~~[Fowler and
456 Johnson, 1995](#)~~). By contrast, if subglacial water routes and ice flow are constrained by bed
457 heterogeneities, migration of successive subglacial water pockets along predetermined paths
458 may induce sequential ice stream surges (Fowler and Johnson, 1995; Hulbe et al., 2016) and
459 participate in the gradual development of complex tunnel valley systems at fixed places, like
460 the Dry Valleys “Labyrinth” in Antarctica (Lewis et al., 2006).

461 6. Ice stream senescence. Ice streams may ultimately switch off when drainage capacities of
462 tunnel valley systems are sufficient to limit subglacial water overpressures. The progressive
463 decay of an ice stream activity can be partially produced by the thinning of the ice layer and the
464 subsequent reduction of the stress driving ice flow in ice stream corridors (Robel et al., 2013).
465 Our experiments display negligible thinning prior to ice stream decay. A constant water
466 discharge being applied in experiments, we demonstrate that increased drainage efficiency

467 during tunnel valley development can solely be responsible for ice stream slowdown. Tunnel
468 valleys and ice streams are frequently found to co-exist ~~as exemplified by~~ ~~and with~~ the many
469 examples reported from the southern margin of the Laurentide Ice Sheet (Patterson, 1997;
470 Livingstone and Clark, 2016). In one case, development of tunnel valleys has been suggested
471 to have led to stagnation of ice flow at an ice stream terminus (Patterson, 1997), a process that
472 we have now demonstrated by modelling. This further ~~suggests~~ ~~indicates~~ that tunnel valleys
473 development could secure ice sheet stability as hinted by Marczynek and Piotrowski, (2006) by
474 preventing ice stream destabilisation. We apply a constant meltwater discharge to our model,
475 however meltwater production and discharge in a subglacial system fluctuates at different times
476 scales (day, year, decades). Fluctuating water production may have further implication on the
477 size of ice streams, the size and amount of tunnel valleys that develop through time, ~~and~~ ~~or~~ the
478 timescale involved in ice sheet destabilization and stabilization. The oscillation in water
479 production could strengthen and multiply the life cycles of some transitory ice streams, already
480 deciphered with a constant water discharge in this study.

481 In a global change context, phenomena of ice stream stabilisation would requires that pre-
482 existing and newly forming tunnel valley systems expand sufficiently fast to accommodate
483 increased meltwater production. Investigating the processes and rates of tunnel valley
484 development are more than ever warranted to better assess ancient and present-day ice sheets
485 behaviour.

486

487

488

489

490

491

492


493

494

495

496

497



498
 499 **Figure 6.** Chronological sequence with interpretative sketches illustrating the proposed ice
 500 stream lifecycle and the relations with tunnel valley development. Basal topography and surface
 501 flow velocity maps are derived from the experiment.

502
 503
 504

505 5. Conclusion

506 The transitory and mobile nature of ice streams may be understood in the framework of a
507 model lifecycle that involves temporal changes in subglacial meltwater pressures and arises
508 from interactions between ice flow, subglacial water drainage and bed erosion. In this model
509 lifecycle transitory ice streams arise, progress and decay in response to subglacial flooding,
510 changes in type and efficiency of subglacial drainage, and development of tunnel valleys. These
511 results are consistent with (and reconcile) a variety of otherwise detached observations
512 performed at different timescales and at different places, on modern and ancient natural ice
513 streams. One of the most novel outcomes of this study, is that subglacial tunnel valley
514 development may be crucial in controlling ice stream vanishing and perhaps, as a consequence,
515 in preventing catastrophic ice sheet collapses during periods of climate change. The processes
516 and rates of tunnel valley development are thus major issues for predicting the forthcoming
517 behaviour of present-day ice sheets and for assessing their contribution to the release of ice and
518 freshwater to the ocean. The innovative experimental approach, used here opens new
519 perspectives on the understanding of subglacial processes controlling ice sheet dynamics and
520 destabilisation.

521 **Author contributions:**

522 OB, RM, ER and SP conceived this research and gathered funding. TL designed and conducted
523 the experiments (setup, monitoring and post-treatment), with contributions by RM and PS. TL,
524 ER, OB, CDC, SP and RM contributed to the interpretation of the results and of their natural
525 implications. TL wrote the first draft of the manuscript; ER, OB, SP and CDC contributed
526 substantially to its present version.

527 **Competing interests:**

528 The authors declare that they have no conflict of interest

529 **Acknowledgements:**

530 This study is part of the DEFORm project (Deformation and Erosion by Fluid Overpressure)
531 funded by “Région Pays de la Loire”. Additional financial support was provided by the French
532 “Agence Nationale de la Recherche” through grant ANR-12-BS06-0014 ‘SEQSTRAT-ICE’
533 and the “Institut National des Sciences de l’Univers” (INSU) through the ‘Programme National
534 de Planétologie’ (PNP) and ‘Système Terre : Processus et Couplages’ (SYSTER) programs.

535 **References**

536 [Alley, R. B., Blankenship, D. D., Rooney, S. T. and Bentley, C. R.: Till beneath ice stream B:
537 4. A coupled ice-till flow model, *J. Geophys. Res. Solid Earth.*, 92, 8931–8940, doi:
538 10.1029/JB092iB09p08931, 1987.](#)
539
540 [Alley, R. B., Dupont, T. K., Parizek, B. R., Anandakrishnan, S., Lawson, D. E., Larson, G. J.,
541 and Evenson, E. B.: Outburst flooding and the initiation of ice-stream surges in response to
542 climatic cooling: A hypothesis, *Geomorphology.*, 75, 76–89,
543 doi:10.1016/j.geomorph.2004.01.011, 2006.](#)

544
545 [Anandakrishnan, S., Blankenship, D. D., Alley, R. B., and Stoffa, P. L.: Influence of subglacial](#)
546 [geology on the position of a West Antarctic ice stream from seismic observations, *Nature.*,](#)
547 [394\(6688\), 62-65, doi: 10.1038/27889, 1998.](#)

548
549 [Anderson, R. S., Walder, J. S., Anderson, S. P., Trabant, D. C. and Fountain, A. G.: The](#)
550 [dynamic response of Kennicott Glacier , Alaska , USA , to the Hidden Creek Lake outburst](#)
551 [flood, *Annals of Glaciology.*, 40, 237–242, doi: 10.3189/172756405781813438, 2005.](#)

552
553 [Bamber, J. L., Vaughan, D. G. & Joughin, I.: Widespread Complex Flow in the Interior of the](#)
554 [Antarctic Ice Sheet, *Science.*, 287, 1248–1250, doi: 10.1126/science.287.5456.1248, 2000.](#)

555
556 [Beckley, B.; Zelensky, N.P.; Holmes, S.A.; Lemoine, F.G.; Ray, R.D.; Mitchum, G.T.; Desai,](#)
557 [S.; Brown, S.T.. 2015. Global Mean Sea Level Trend from Integrated Multi-Mission Ocean](#)
558 [Altimeters TOPEX/Poseidon Jason-1 and OSTM/Jason-2 Version 3. Ver. 3. PO.DAAC, CA,](#)
559 [USA. Dataset accessed at <http://dx.doi.org/10.5067/GMSLM-TJ123>.](#)

560
561 [Beem, L. H., Tulaczyk, S. M., King, M. A., Bougamont, M., Fricker, H. A., and Christoffersen,](#)
562 [P.: Variable deceleration of Whillans Ice Stream, West Antarctica, *J. Geophys. Res. Earth Surf.*,](#)
563 [119, 212–224, doi: 10.1002/2013JF002958, 2014.](#)

564
565 [Bell, R. E., Studinger, M., Shuman, C. A., Fahnestock, M. A. and Joughin, I.: Large subglacial](#)
566 [lakes in East Antarctica at the onset of fast-flowing ice streams, *Nature.*, 445, 904–907, doi:](#)
567 [10.1038/nature05554, 2007.](#)

568
569 [Bennett, M. R.: Ice streams as the arteries of an ice sheet: their mechanics, stability and](#)
570 [significance, *Earth-Science Rev.*, 61, 309–339, doi: 10.1016/S0012-8252\(02\)00130-7, 2003.](#)

571
572 [Bentley, C. R.: Antarctic ice streams: a review. *J. Geophys. Res. Solid Earth.*, 92\(B9\), 8843–](#)
573 [8858, doi: 10.1029/JB092iB09p08843, 1987.](#)

574
575 [Bingham, R. G., King, E. C., Smith, A. M. and Pritchard, H. D.: Glacial geomorphology:](#)
576 [towards a convergence of glaciology and geomorphology, *Prog. Phys. Geogr.*, 34, 327–355,](#)
577 [doi: 10.1177/0309133309360631, 2010.](#)

578
579 [Blankenship, D. D., Bell, R. E., Hodge, S. M., Brozena, J. M., Behrendt, J. C., and Finn, C. A.:](#)
580 [Active volcanism beneath the West Antarctic ice sheet and implications for ice-sheet stability,](#)
581 [*Nature.*, 361\(6412\), 526-529, doi: 10.1038/361526a0, 1993.](#)

582
583 [Bourgeois, O., Dauteuil, O., and Van Vliet-Lanoe, B.: Geothermal control on flow patterns in](#)
584 [the Last Glacial Maximum ice sheet of Iceland, *Earth Surf. Process and Landforms.*, 25, 59–](#)
585 [76, doi: 10.1002/\(SICI\)1096-9837\(200001\)25:1<59::AID-ESP48>3.0.CO;2-T, 2000.](#)

586
587 [Carter, S. P., Fricker, H. A., and Siegfried, M. R.: Evidence of rapid subglacial water piracy](#)
588 [under Whillans Ice Stream, West Antarctica, *Journal of Glaciology.*, 59, 1147–1162, doi:](#)
[10.3189/2013JoG13J085, 2013.](#)

589

590 [Carter, S. P., Fricker, H. A., and Siegfried, M. R.: Antarctic subglacial lakes drain through](#)
591 [sediment-floored canals: Theory and model testing on real and idealized domains, *Cryosph.*](#)
592 [11, 381–405, doi: 10.5194/tc-11-381-2017, 2017.](#)

593
594 [Catania, G., and Paola, C.: Braiding under glass, *Geology*, 29, 259–262, doi: 10.1130/0091-](#)
595 [7613\(2001\)029<0259:BUG>2.0.CO;2, 2001.](#)

596
597 [Catania, G. A., Scambos, T. A., Conway, H., and Raymond, C. F.: Sequential stagnation of](#)
598 [Kamb ice stream, West Antarctica, *Geophys. Res. Lett.*, 33\(14\), L14502 doi:](#)
599 [10.1029/2006GL026430, 2006.](#)

600
601 [Catania, G., Hulbe, C., Conway, H., Scambos, T. A., and Raymond, C. F.: Variability in the](#)
602 [mass flux of the Ross ice streams, West Antarctica, over the last millennium, *J. Glaciol.* 58,](#)
603 [741–752, doi: 10.3189/2012JoG11J219, 2012.](#)

604
605 [Elsworth, C. W., and Suckale, J.: Rapid ice flow rearrangement induced by subglacial drainage](#)
606 [in West Antarctica, *Geophys. Res. Lett.*, 43, 697-707, doi: 10.1002/2016GL070430, 2016.](#)

607
608 [Engelhardt, H., Humphrey, N., Kamb, B., and Fahnestock, M.: Physical conditions at the base](#)
609 [of a fast moving antarctic ice stream, *Science*, 248\(4951\), 57-59, doi:](#)
610 [10.1126/science.248.4951.57, 1990.](#)

611
612 [Evatt, G. W., Fowler, A. C., Clark, C. D., and Hulton, N. R. J.: Subglacial floods beneath ice](#)
613 [sheets, *Philos. Trans. R. Soc. London A Math. Phys. Eng. Sci.*, 364, 1769–1794, doi:](#)
614 [10.1098/rsta.2006.1798, 2006.](#)

615
616 [Flowers, G. E.: Modelling water flow under glaciers and ice sheets, *Proc. R. Soc. A Math. Phys.*](#)
617 [Eng. Sci., 471\(2176\), 20140907, doi: 10.1098/rspa.2014.0907, 2015.](#)

618
619 [Fountain, A. G., and Walder, J. S.: Water flow through temperate glaciers, *Rev. Geophys.*, 36,](#)
620 [299–328, doi: 10.1029/97RG03579, 1998.](#)

621
622 [Fowler, A. C., and Johnson, C.: Hydraulic run-away: a mechanism for thermally regulated](#)
623 [surges of ice sheets, *J. Glaciol.*, 41, 454–461, doi: 10.3189/S002214300003478X, 1995.](#)

624
625 [Fricker, H. A., Scambos, T., Bindschadler, R., and Padman, L.: An active subglacial water](#)
626 [system in West Antarctica mapped from space, *Science*, 315, 1544–1548, doi:](#)
627 [10.1126/science.1136897, 2007.](#)

628
629 [Glen, J. W.: The stability of ice-dammed lakes and other water-filled holes in glaciers. *Journal*](#)
630 [of *Glaciology*, 2\(15\), 316-318, doi: 10.3189/S0022143000025132, 1954.](#)

631
632 [Gray, L., Joughin, I., Tulaczyk, S., Spikes, V. B., Bindschadler, R., and Jezek, K.: Evidence for](#)
633 [subglacial water transport in the West Antarctic Ice Sheet through three-dimensional satellite](#)
[radar interferometry, *Geophys. Res. Lett.*, 32, L03501, doi: 10.1029/2004GL021387, 2005.](#)

634
635 [Hanson, B., Hooke, R. L., and Grace, E. M.: Short-term velocity and water-pressure variations](#)
636 [down-glacier from a riegel, Storglaciären, Sweden. *Journal of Glaciology*, 44\(147\), 359-367,](#)
637 [doi: 10.3189/S0022143000002689, 1998\).](#)

Code de champ modifié

Code de champ modifié

634 [Hindmarsh, R. C. A.: Consistent generation of ice-streams via thermo-viscous instabilities](#)
635 [modulated by membrane stresses, *Geophys. Res. Lett.*, 36, L06502, doi:](#)
636 [10.1029/2008GL036877, 2009.](#)

637
638 [Hooke, R. L., and Jennings, C. E.: On the formation of the tunnel valleys of the southern](#)
639 [Laurentide ice sheet, *Quat. Sci. Rev.*, 25, 1364–1372, doi: 10.1016/j.quascirev.2006.01.018,](#)
640 [2006.](#)

Code de champ modifié

641
642 [Hulbe, C.: Is ice sheet collapse in West Antarctica unstoppable?, *Science.*, 356, 910–911, doi:](#)
643 [10.1126/science.aam9728, 2017.](#)

644
645 [Hulbe, C. L., Scambos, T. A., Klinger, M., and Fahnestock, M. A.: Flow variability and ongoing](#)
646 [margin shifts on Bindschadler and MacAyeal Ice Streams, West Antarctica, *J. Geophys. Res.*](#)
647 [Earth Surf., 121, 283–293, doi: 10.1002/2015JF003670, 2016.](#)

648
649 [Jørgensen, F., and Piotrowski, J. A.: Signature of the Baltic ice stream on Funen Island,](#)
650 [Denmark during the Weichselian glaciation, *Boreas.*, 32, 242–255, doi: 10.1111/j.1502-](#)
651 [3885.2003.tb01440.x, 2003.](#)

652
653 [Kamb, B.: Glacier surge mechanism based on linked cavity configuration of the basal water](#)
654 [conduit system, *J. Geophys. Res. Solid Earth.*, 92, 9083–9100, doi: 10.1029/JB092iB09p09083,](#)
655 [1987.](#)

656
657 [Kehew, A. E., Piotrowski, J. A., and Jørgensen, F.: Tunnel valleys: Concepts and controversies.](#)
658 [A review, *Earth-Science Rev.*, 113, 33–58, doi: 10.1016/j.earscirev.2012.02.002, 2012.](#)

Code de champ modifié

659
660 [Kim, B.-H., Lee, C.-K., Seo, K.-W., Lee, W. S., and Scambos, T.: Active subglacial lakes](#)
661 [beneath the stagnant trunk of Kamb Ice Stream: evidence of channelized subglacial flow,](#)
662 [Cryosph., 10, 2971-2980, https://doi.org/10.5194/tc-10-2971-2016, 2016.](#)

663
664 [Kleiner, T., and Humbert, A.: Numerical simulations of major ice streams in western](#)
665 [Dronning Maud Land, Antarctica, under wet and dry basal conditions, *J. Glaciol.*, 60, 215–](#)
666 [232, doi: 10.3189/2014JoG13J006, 2014.](#)

667 [Kowal, K. N., and Worster, M. G.: Lubricated viscous gravity currents, *Journal of Fluid*](#)
668 [Mechanics., 766, 626-655, doi: 10.1017/jfm.2015.30, 2000.](#)

669 [Kyrke-Smith, T.M., Katz, R.F., and Fowler, A.C.: Subglacial hydrology and the formation of](#)
670 [ice streams, *Proceedings of the Royal Society.*, A 470\(2161\), 20130494, doi:](#)
671 [10.1098/rspa.2013.0494, 2014.](#)

672
673 [Le Brocq, A. M., Ross, N., Griggs, J.A., Bingham, R.G., Corr, H.F.J., Ferraccioli, F., Jenkins,](#)
674 [A., Jordan, T.A., Payne, A.J., Rippin, D.M., and Siegert, M.J.: Evidence from ice shelves for](#)
675 [channelized meltwater flow beneath the Antarctic Ice Sheet, *Nat. Geosci.*, 6, 945–948, doi:](#)
676 [10.1038/ngeo1977, 2013.](#)

Code de champ modifié

677
678 [Lelandais, T., Mourgues, R., Ravier, É., Pochat, S., Strzeczynski, P., and Bourgeois, O.:](#)
679 [Experimental modeling of pressurized subglacial water flow: Implications for tunnel valley](#)
680 [formation, *J. of Geophy. Res. Earth Surface.*, 121\(11\), 2022-2041, doi: 10.1002/2016JF003957,](#)
681 [2016.](#)

682

683 [Lewis, A. R., Marchant, D. R., Kowalewski, D. E., Baldwin, S. L., and Webb, L. E.: The age](#)
684 [and origin of the Labyrinth, western Dry Valleys, Antarctica: Evidence for extensive middle](#)
685 [Miocene subglacial floods and freshwater discharge to the Southern Ocean, *Geology*, 34,](#)
686 [513–516, doi: 10.1130/G22145.1, 2006.](#)

687 [Livingstone, S. J., and Clark, C. D.: Morphological properties of tunnel valleys of the southern](#)
688 [sector of the Laurentide Ice Sheet and implications for their formation, *Earth Surface*](#)
689 [Dynamics, 4, 567–589, doi: 10.5194/esurf-4-567-2016, 2016.](#)

690

691 [Livingstone, S. J., Utting, D. J., Ruffell, A., Clark, C. D., Pawley, S., Atkinson, N., and Fowler,](#)
692 [A. C.: Discovery of relict subglacial lakes and their geometry and mechanism of drainage, *Nat.*](#)
693 [Commun., 7, 11767, doi: 10.1038/ncomms11767, 2016.](#)

694

695 [Magnússon, E., Rott, H., Björnsson, H., and Pálsson, F., The impact of jökulhlaups on basal](#)
696 [sliding observed by SAR interferometry on Vatnajökull, Iceland, *J. Glaciol.*, 53, 232–240,](#)
697 [doi: 10.3189/172756507782202810, 2007.](#)

698 [Marczinek, S., and Piotrowski, J. A.: Groundwater flow under the margin of the last](#)
699 [Scandinavian ice Sheet Around the Eckernförde Bay, Northwest Germany, *Glacier Sci.*](#)
700 [Environ. Chang., Knight, P.G., Blackwell Science Ltd 60–62, doi:](#)
701 [10.1002/9780470750636.ch10, 2006.](#)

702 [Margold, M., Stokes, C. R., Clark, C. D., and Kleman, J.: Ice streams in the Laurentide Ice](#)
703 [Sheet: a new mapping inventory, *J. Maps.*, 11, 380–395, doi: 10.1080/17445647.2014.912036,](#)
704 [2015.](#)

705 [Marshall, S. J.: Recent advances in understanding ice sheet dynamics, *Earth Planet. Sci. Lett.*,](#)
706 [240, 191–204, doi: 10.1016/j.epsl.2005.08.016, 2005.](#)

707 [Paola, C., Straub, K., Mohrig, D., and Reinhardt, L.: The unreasonable effectiveness of](#)
708 [stratigraphic and geomorphic experiments, *Earth-Science Rev.*, 97, 1–43, doi:](#)
709 [10.1080/17445647.2014.912036, 2009.](#)

710 [Paterson, W. S. B., Pergamon Kidlington, The Physics of Glaciers, Elsevier Science Ltd,](#)
711 [Great Britain, 480 pages, 1994.](#)

712 [Patterson, C. J.: Southern Laurentide ice lobes were created by ice streams: Des Moines Lobe](#)
713 [in Minnesota, USA, *Sediment. Geol.*, 111, 249–261, doi: 10.1016/S0037-0738\(97\)00018-3,](#)
714 [1997.](#)

715 [Payne, A. J., and Dongelmans, P. W.: Self-organization in the thermomechanical flow of ice](#)
716 [sheets, *J. Geophys. Res. Solid Earth.*, 102, 12219–12233, doi: 10.1029/97JB00513, 1997.](#)

717

718 [Perol, T., and Rice, J. R.: Shear heating and weakening of the margins of West Antarctic ice](#)
719 [streams, *Geophys. Res. Lett.*, 42, 3406–3413, doi: 10.1002/2015GL063638, 2015.](#)

720

Code de champ modifié

721 [Peters, L. E., Anandkrishnan, S., Alley, R. B., and Smith, A. M., Extensive storage of basal](#)
722 [meltwater in the onset region of a major West Antarctic ice stream, *Geology*, 35, 251–254,](#)
723 [doi: 10.1130/G23222A.1, 2007.](#)

724 [Ravier, E., Buoncristiani, J. F., Menzies, J., Guiraud, M., Clerc, S., and Portier, E.: Does](#)
725 [porewater or meltwater control tunnel valley genesis? Case studies from the Hirnantian of](#)
726 [Morocco, *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 418, 359–376, doi:](#)
727 [10.1016/j.palaeo.2014.12.003, 2015.](#)

728 [Raymond, C. F.: How do glaciers surge? A review, *J. Geophys. Res. Solid Earth*, 92, 9121–](#)
729 [9134, doi: 10.1029/JB092iB09p09121, 1987.](#)

730 [Retzlaff, R., and Bentley, C. R.: Timing of stagnation of Ice Stream C, West Antarctica, from](#)
731 [short-pulse radar studies of buried surface crevasses, *J. Glaciol.*, 39, 553–561, doi:](#)
732 [10.3189/S0022143000016440, 1993.](#)

733 [Robel, A. A., Degiuli, E., Schoof, C., and Tziperman, E.: Dynamics of ice stream temporal](#)
734 [variability: Modes, scales, and hysteresis, *J. of Geophys. Res. Earth Surface*, 118, 925–936,](#)
735 [doi: 10.1002/jgrf.20072, 2013.](#)

736

737 [Rothlisberger, H., and Lang, H., *Glacial hydrology. Glacio-Fluvial Sediment Transfer: An*](#)
738 [Alpine Perspective, Gurnell and Clark, John Wiley and Sons, New York New York, p 207-](#)
739 [284, 1987.](#)

740 [Shreve, R. L.: Movement of water in glaciers, *Journal of Glaciology*, 11\(62\), 205-214, doi:](#)
741 [10.3189/S002214300002219X, 1972.](#)

742 [Siegfried, M. R., Fricker, H. A., Carter, S. P., and Tulaczyk, S.: Episodic ice velocity](#)
743 [fluctuations triggered by a subglacial flood in West Antarctica, *Geophys. Res. Lett.*, 43, 2640–](#)
744 [2648, doi: 10.1002/2016GL067758, 2016.](#)

745

746 [Stearns, L. A., Smith, B. E., and Hamilton, G. S.: Increased flow speed on a large East Antarctic](#)
747 [outlet glacier caused by subglacial floods, *Nat. Geosci.*, 1, 827–831, doi: 10.1038/ngeo356,](#)
748 [2008.](#)

749

750 [Vaughan, D. G., Corr, H. F. J., Smith, A. M., Pritchard, H. D., and Shepherd, A.: Flow-](#)
751 [switching and water piracy between Rutford Ice Stream and Carlson Inlet, West Antarctica,](#)
752 [*Journal of Glaciology*, 54, 41–48, doi: 10.3189/002214308784409125, 2008.](#)

753 [Vaughan, D.G., J.C. Comiso, I. Allison, J. Carrasco, G. Kaser, R. Kwok, P. Mote, T. Murray,](#)
754 [F. Paul, J. Ren, E. Rignot, O. Solomina, K. Steffen and T. Zhang, 2013: Observations:](#)
755 [Cryosphere. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working*](#)
756 [Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change](#)
757 [\[Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia,](#)
758 [V. Bex and P.M. Midgley \(eds.\)\]. Cambridge University Press, Cambridge, United Kingdom](#)
759 [and New York, NY, USA](#)

760

761 [Winsborrow, M. C. M., Clark, C. D., and Stokes, C. R.: What controls the location of ice](#)
762 [streams?, *Earth-Science Rev.*, 103, 45–59, doi: 10.1016/j.earscirev.2010.07.003, 2010.](#)

Code de champ modifié

763 Alley, R. B., Blankenship, D. D., Rooney, S. T. and Bentley, C. R.: Till beneath ice stream B-
764 4. A coupled ice-till flow model, *J. Geophys. Res. Solid Earth.*, 92, 8931–8940, doi:
765 10.1029/JB092iB09p08931, 1987.

766
767 Alley, R. B., Dupont, T. K., Parizek, B. R., Anandakrishnan, S., Lawson, D. E., Larson, G. J.,
768 and Evenson, E. B.: Outburst flooding and the initiation of ice stream surges in response to
769 climatic cooling: A hypothesis, *Geomorphology.*, 75, 76–89,
770 doi:10.1016/j.geomorph.2004.01.011, 2006.

Code de champ modifié

771
772 Anandakrishnan, S., Blankenship, D. D., Alley, R. B., and Stoffa, P. L.: Influence of subglacial
773 geology on the position of a West Antarctic ice stream from seismic observations, *Nature.*,
774 394(6688), 62–65, doi: 10.1038/27889, 1998.

Code de champ modifié

775
776 Anderson, R. S., Walder, J. S., Anderson, S. P., Trabant, D. C. and Fountain, A. G.: The
777 dynamic response of Kennicott Glacier, Alaska, USA, to the Hidden Creek Lake outburst
778 flood, *Annals of Glaciology.*, 40, 237–242, doi: 10.3189/172756405781813438, 2005.

779
780 Bamber, J. L., Vaughan, D. G. & Joughin, I.: Widespread Complex Flow in the Interior of the
781 Antarctic Ice Sheet, *Science.*, 287, 1248–1250, doi: 10.1126/science.287.5456.1248, 2000.

782
783 Beckley, B.; Zelensky, N.P.; Holmes, S.A.; Lemoine, F.G.; Ray, R.D.; Mitchum, G.T.; Desai,
784 S.; Brown, S.T., 2015. Global Mean Sea Level Trend from Integrated Multi-Mission Ocean
785 Altimeters TOPEX/Poseidon Jason-1 and OSTM/Jason-2 Version 3, Ver. 3. PO.DAAC, CA,
786 USA. Dataset accessed [YYYY-MM-DD] at <http://dx.doi.org/10.5067/GMSLM-TJ123>.

Code de champ modifié

787
788 Beem, L. H., Tulaczyk, S. M., King, M. A., Bougamont, M., Fricker, H. A., and Christoffersen,
789 P.: Variable deceleration of Whillans Ice Stream, West Antarctica, *J. Geophys. Res. Earth Surf.*,
790 119, 212–224, doi: 10.1002/2013JF002958, 2014.

791
792 Bell, R. E., Studinger, M., Shuman, C. A., Fahnestock, M. A. and Joughin, I.: Large subglacial
793 lakes in East Antarctica at the onset of fast-flowing ice streams, *Nature.*, 445, 904–907, doi:
794 10.1038/nature05554, 2007.

Code de champ modifié

795
796 Bennett, M. R.: Ice streams as the arteries of an ice sheet: their mechanics, stability and
797 significance, *Earth Science Rev.*, 61, 309–339, doi: 10.1016/S0012-8252(02)00130-7, 2003.

Code de champ modifié

798
799 Bentley, C. R.: Antarctic ice streams: a review. *J. Geophys. Res. Solid Earth.*, 92(B9), 8843–
800 8858, doi: 10.1029/JB092iB09p08843, 1987.

801
802 Bingham, R. G., King, E. C., Smith, A. M. and Pritchard, H. D.: Glacial geomorphology:
803 towards a convergence of glaciology and geomorphology, *Prog. Phys. Geogr.*, 34, 327–355,
804 doi: 10.1177/0309133309360631, 2010.

Code de champ modifié

805
806 Blankenship, D. D., Bell, R. E., Hodge, S. M., Brozena, J. M., Behrendt, J. C., and Finn, C. A.:
807 Active volcanism beneath the West Antarctic ice sheet and implications for ice sheet stability,
808 *Nature.*, 361(6412), 526, doi: 10.1038/361526a0, 1993.

Code de champ modifié

809
810 Bourgeois, O., Dauteuil, O., and Vliet-Lanoe, B. Van.: Geothermal control on flow patterns in
811 the Last Glacial Maximum ice sheet of Iceland, *Earth Surf. Process and Landforms.*, 25, 59–
812 76, doi: 10.1002/(SICI)1096-9837(200001)25:1<59::AID-ESP48>3.0.CO;2-T, 2000.

813 Carter, S. P., Fricker, H. A., and Siegfried, M. R.: Evidence of rapid subglacial water piracy
814 under Whillans Ice Stream, West Antarctica, *Journal of Glaciology*, 59, 1147–1162, doi:
815 10.3189/2013JG13J085, 2013.

816
817 Carter, S. P., Fricker, H. A., and Siegfried, M. R.: Antarctic subglacial lakes drain through
818 sediment floored canals: Theory and model testing on real and idealized domains, *Cryosph.*,
819 11, 381–405, doi: 10.5194/te-11-381-2017, 2017.

820
821 Catania, G., and Paola, C.: Braiding under glass, *Geology*, 29, 259–262, doi: 10.1130/0091-
822 7613(2001)029<0259:BUG>2.0.CO;2, 2001.

823
824 Catania, G. A., Scambos, T. A., Conway, H., and Raymond, C. F.: Sequential stagnation of
825 Kamb ice stream, West Antarctica, *Geophys. Res. Lett.*, 33(14), doi: 10.1029/2006GL026430,
826 2006.

827
828 Catania, G., Hulbe, C., Conway, H., Scambos, T. A., and Raymond, C. F.: Variability in the
829 mass flux of the Ross ice streams, West Antarctica, over the last millennium. *J. Glaciol.* 58,
830 741–752, doi: 10.3189/2012JG11J219, 2012.

831
832 Elsworth, C. W., and Suckale, J.: Rapid ice flow rearrangement induced by subglacial drainage
833 in West Antarctica, *Geophys. Res. Lett.*, 43, doi: 10.1002/2016GL070430, 2016.

834
835 Engelhardt, H., Humphrey, N., Kamb, B., and Fahnestock, M.: Physical Conditions at the Base
836 of a Fast Moving Antarctic Ice Stream, *Science*, 248(4951), 57–59, doi:
837 10.1126/science.248.4951.57, 1990.

838
839 Evatt, G. W., Fowler, A. C., Clark, C. D., and Hulton, N. R. J.: Subglacial floods beneath ice
840 sheets, *Philos. Trans. R. Soc. London A Math. Phys. Eng. Sci.*, 364, 1769–1794, doi:
841 10.1098/rsta.2006.1798, 2006.

842
843 Flowers, G. E.: Modelling water flow under glaciers and ice sheets, *Proc. R. Soc. A Math. Phys.*
844 *Eng. Sci.*, 471(2176), doi: 10.1098/rspa.2014.0907, 2015.

845
846 Fountain, A. G., and Walder, J. S.: Water flow through temperate glaciers, *Rev. Geophys.*, 36,
299–328, doi: 10.1029/97RG03579, 1998.

847
848 Fowler, A. C., and Johnson, C.: Hydraulic run-away: a mechanism for thermally regulated
surges of ice sheets, *J. Glaciol.*, 41, 454–461, doi: 10.3189/S002214300003478X, 1995.

849
850 Fricker, H. A., Scambos, T., Bindshadler, R., and Padman, L.: An active subglacial water
851 system in West Antarctica mapped from space, *Science*, 315, 1544–1548, doi:
10.1126/science.1136897, 2007.

852
853 Glen, J. W.: Experiments on the deformation of ice, *Journal of Glaciology*, 2(12), 111–114,
doi: 10.3189/S0022143000034067, 1952.

854
855 Gray, L., Joughin, I., Tulaczyk, S., Spikes, V. B., Bindshadler, R., and Jezek, K.: Evidence for
856 subglacial water transport in the West Antarctic Ice Sheet through three-dimensional satellite
radar interferometry, *Geophys. Res. Lett.*, 32, doi: 10.1029/2004GL021387, 2005.

Code de champ modifié

857
858 Hindmarsh, R. C. A.: Consistent generation of ice streams via thermo viscous instabilities
859 modulated by membrane stresses, *Geophys. Res. Lett.*, 36, doi: 10.1029/2008GL036877, 2009.
860
861 Hooke, R. L., and Jennings, C. E.: On the formation of the tunnel valleys of the southern
862 Laurentide ice sheet, *Quat. Sci. Rev.*, 25, 1364–1372, doi: 10.1016/j.quascirev.2006.01.018,
863 2006.
864
865 Hulbe, C. L., Scambos, T. A., Klinger, M., and Fahnestock, M. A.: Flow variability and ongoing
866 margin shifts on Bindeschadler and MacAyeal Ice Streams, West Antarctica, *J. Geophys. Res.*
867 *Earth Surf.*, 121, 283–293, doi: 10.1002/2015JF003670, 2016.
868
869 Hulbe, C.: Is ice sheet collapse in West Antarctica unstoppable?, *Science.*, 356, 910–911, doi:
870 10.1126/science.aam9728, 2017.
871
872 Jørgensen, F., and Piotrowski, J. A.: Signature of the Baltic ice stream on Funen Island,
873 Denmark during the Weichselian glaciation, *Boreas.*, 32, 242–255, doi: 10.1111/j.1502-
874 3885.2003.tb01440.x, 2003.
875
876 Kamb, B., Raymond, C.F., Harrison, W.D., Engelhardt, H., Echelmeyer, K.A., Humphrey, N.,
877 Brugman, M.M., and Pfeffer, T.: Glacier surge mechanism: 1982–1983 surge of Variegated
878 Glacier, Alaska, *Science.*, 227, 469–479, doi: 10.1126/science.227.4686.469, 1985.
879
880 Kamb, B.: Glacier surge mechanism based on linked cavity configuration of the basal water
881 conduit system, *J. Geophys. Res. Solid Earth.*, 92, 9083–9100, doi: 10.1029/JB092iB09p09083,
882 1987.
883
884 Kehew, A. E., Piotrowski, J. A., and Jørgensen, F.: Tunnel valleys: Concepts and controversies:
885 A review, *Earth Science Rev.*, 113, 33–58, doi: 10.1016/j.earseirev.2012.02.002, 2012.
886
887 Kim, B. H., Lee, C. K., Seo, K. W., Lee, W. S., and Scambos, T.: Active subglacial lakes
888 beneath the stagnant trunk of Kamb Ice Stream: evidence of channelized subglacial flow,
889 *Cryosph.*, 1–15, doi: 10.5194/tc-10-2971-2016, 2016.
890
891 Kleiner, T., and Humbert, A.: Numerical simulations of major ice streams in western
892 Dronning Maud Land, Antarctica, under wet and dry basal conditions, *J. Glaciol.*, 60, 215–
893 232, doi: 10.3189/2014JG13J006, 2014.
894
895 Kowal, K. N., and Worster, M. G.: Lubricated viscous gravity currents, *Journal of Fluid*
Mechanics., 766, 626–655, doi: 10.1017/jfm.2015.30, 2000.
896
897 Kyrke Smith, T.M., Katz, R.F., and Fowler, A.C.: Subglacial hydrology and the formation of
898 ice streams, *Proceedings of the Royal Society.*, A 470(2161), 20130494, doi:
899 10.1098/rspa.2013.0494, 2014.
900
901 Hanson, B., Hooke, R. L., and Grace, E. M.: Short term velocity and water pressure variations
902 down glacier from a riegel, Storgraciären, Sweden. *Journal of Glaciology*, 44(147), 359–367,
doi: 10.3189/S0022143000002689, 1998).
903
904 Le Brocq, A. M., Ross, N., Griggs, J.A., Bingham, R.G., Corr, H.F.J., Ferraccioli, F., Jenkins,
A., Jordan, T.A., Payne, A.J., Rippin, D.M., and Siegert, M.J.: Evidence from ice shelves for

Code de champ modifié

Code de champ modifié

Code de champ modifié

905 channelized meltwater flow beneath the Antarctic Ice Sheet, *Nat. Geosci.*, 6, 945–948, doi:
906 10.1038/ngeo1977, 2013.

907

908 Lelandais, T., Mourgues, R., Ravier, É., Pochat, S., Strzeczynski, P., and Bourgeois, O.:
909 Experimental modeling of pressurized subglacial water flow: Implications for tunnel valley
910 formation, *J. of Geophys. Res. Earth Surface.*, 121(11), 2022–2041, doi: 10.1002/2016JF003957,
911 2016.

912

913 Lewis, A. R., Marchant, D. R., Kowalewski, D. E., Baldwin, S. L., and Webb, L. E.: The age
914 and origin of the Labyrinth, western Dry Valleys, Antarctica: Evidence for extensive middle
915 Miocene subglacial floods and freshwater discharge to the Southern Ocean, *Geology.*, 34,
916 513–516, doi: 10.1130/G22145.1, 2006.

917 Livingstone, S. J., Utting, D. J., Ruffell, A., Clark, C. D., Pawley, S., Atkinson, N., and Fowler,
918 A. C.: Discovery of relict subglacial lakes and their geometry and mechanism of drainage. *Nat.*
919 *Commun.*, 7, neomms11767, doi: 10.1038/neomms11767, 2016.

920

921 Livingstone, S. J., and Clark, C. D.: Morphological properties of tunnel valleys of the southern
922 sector of the Laurentide Ice Sheet and implications for their formation, *Earth Surface*
923 *Dynamics*, 4, 567–589, doi: 10.5194/esurf-4-567-2016, 2016.

924

925 Magnússon, E., Rott, H., Björnsson, H., and Pálsson, F.: The impact of jökulhlaups on basal
926 sliding observed by SAR interferometry on Vatnajökull, Iceland, *J. Glaciol.*, 53, 232–240,
927 doi: 10.3189/172756507782202810, 2007.

928

929 Marciznek, S., and Piotrowski, J. A.: Groundwater flow under the margin of the last
930 Scandinavian ice Sheet Around the Eckernförde Bay, Northwest Germany, *Glacier Sci.*
931 *Environ. Chang.*, 60–62, doi: 10.1002/9780470750636.ch10, 2006.

932

933 Marshall, S. J.: Recent advances in understanding ice sheet dynamics, *Earth Planet. Sci. Lett.*,
934 240, 191–204, doi: 10.1016/j.epsl.2005.08.016, 2005.

935

936 Margold, M., Stokes, C. R., Clark, C. D., and Kleman, J.: Ice streams in the Laurentide Ice
937 Sheet: a new mapping inventory, *J. Maps.*, 11, 380–395, doi: 10.1080/17445647.2014.912036,
938 2015.

939

940 Paola, C., Straub, K., Mohrig, D., and Reinhardt, L.: The unreasonable effectiveness of
941 stratigraphic and geomorphic experiments, *Earth Science Rev.*, 97, 1–43, doi:
942 10.1080/17445647.2014.912036, 2009.

943

944 Paterson, W. S. B., Pergamon Kidlington, *The Physics of Glaciers*, Elsevier Science Ltd,
Great Britain, 480 pages, 1994.

945

946 Patterson, C. J.: Southern Laurentide ice lobes were created by ice streams: Des Moines Lobe
947 in Minnesota, USA, *Sediment. Geol.*, 111, 249–261, doi: 10.1016/S0037-0738(97)00018-3,
1997.

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

944 Payne, A. J., and Dongelmans, P. W.: Self-organization in the thermomechanical flow of ice
 945 sheets. *J. Geophys. Res. Solid Earth.*, 102, 12219–12233, doi: 10.1029/97JB00513, 1997.

946

947 Perol, T., and Rice, J. R.: Shear heating and weakening of the margins of West Antarctic ice
 948 streams. *Geophys. Res. Lett.*, 42, 3406–3413, doi: 10.1002/2015GL063638, 2015.

949

950 Peters, L. E., Anandakrishnan, S., Alley, R. B., and Smith, A. M.: Extensive storage of basal
 951 meltwater in the onset region of a major West Antarctic ice stream. *Geology.*, 35, 251–254,
 952 doi: 10.1130/G23222A.1, 2007.

953 Ravier, E., Buoneristiani, J. F., Menzies, J., Guiraud, M., Clere, S., and Portier, E.: Does
 954 porewater or meltwater control tunnel valley genesis? Case studies from the Hirnantian of
 955 Morocco. *Palaeogeogr. Palaeoclimatol. Palaeoecol.*, 418, 359–376, doi:
 956 10.1016/j.palaeo.2014.12.003, 2015.

957 Raymond, C. F.: How do glaciers surge? A review. *J. Geophys. Res. Solid Earth.*, 92, 9121–
 958 9134, doi: 10.1029/JB092iB09p09121, 1987.

959 Retzlaff, R., and Bentley, C. R.: Timing of stagnation of Ice Stream C, West Antarctica, from
 960 short-pulse radar studies of buried surface crevasses. *J. Glaciol.*, 39, 553–561, doi:
 961 10.3189/S0022143000016440, 1993.

962 Robel, A. A., Degiuli, E., Schoof, C., and Tziperman, E.: Dynamics of ice stream temporal
 963 variability: Modes, scales, and hysteresis. *J. Geophys. Res.*, 118, 925–936, doi: 10.1002/jgrf.20072, 2013.

964

965 Rothlisberger, H., and Lang, H.: Glacial hydrology. *Glacio-Fluvial Sediment Transfer: An*
 966 *Alpine Perspective*. John Wiley and Sons, New York New York, p 207–284, 1987.

967 Shreve, R. L.: Movement of water in glaciers. *Journal of Glaciology.*, 11(62), 205–214, doi:
 968 10.3189/S002214300002219X, 1972.

969 Siegfried, M. R., Fricker, H. A., Carter, S. P., and Tulaczyk, S.: Episodic ice velocity
 970 fluctuations triggered by a subglacial flood in West Antarctica. *Geophys. Res. Lett.*, 43, 2640–
 971 2648, doi: 10.1002/2016GL067758, 2016.

972

973 Stearns, L. A., Smith, B. E., and Hamilton, G. S.: Increased flow speed on a large East Antarctic
 974 outlet glacier caused by subglacial floods. *Nat. Geosci.*, 1, 827–831, doi: 10.1038/ngeo356,
 975 2008.

976

977 Vaughan, D. G., Corr, H. F. J., Smith, A. M., Pritchard, H. D., and Shepherd, A.: Flow-
 978 switching and water piracy between Rutford Ice Stream and Carlson Inlet, West Antarctica.
 979 *Journal of Glaciology.*, 54, 41–48, doi: 10.3189/002214308784409125, 2008.

980 Vaughan, D.G., J.C. Comiso, I. Allison, J. Carrasco, G. Kaser, R. Kwok, P. Mote, T. Murray,
 981 F. Paul, J. Ren, E. Rignot, O. Solomina, K. Steffen and T. Zhang, 2013: Observations:
 982 Cryosphere. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working*
 983 *Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*
 984 [Stocker, T.F., D. Qin, G. K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia,

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

Code de champ modifié

985 ~~V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom~~
986 ~~and New York, NY, USA~~
987
988 ~~Winsborrow, M. C. M., Clark, C. D., and Stokes, C. R.: What controls the location of ice~~
989 ~~streams?, Earth Science Rev., 103, 45–59, doi: 10.1016/j.earscirev.2010.07.003, 2010.~~