

Satellite Passive Microwave Sea-Ice Concentration Data Set Intercomparison: Closed Ice and Ship-Based Observations

Stefan Kern¹, Thomas Lavergne², Dirk Notz³, Leif Toudal Pedersen⁴, Rasmus Tage Tonboe⁵, Roberto Saldo⁴, and Atle Macdonald Sørensen²

¹Integrated Climate Data Center (ICDC), Center for Earth System Research and Sustainability (CEN), University of Hamburg, Hamburg, Germany

²Research and Development Department, Norwegian Meteorological Institute, Oslo, Norway

³Institute for Marine Research, University of Hamburg and Max-Planck Institute for Meteorology, Hamburg, Germany

⁴Danish Technical University, Lyngby, Denmark

⁵Danish Meteorological Institute, Copenhagen, Denmark

Correspondence to: Stefan Kern (stefan.kern@uni-hamburg.de)

Abstract. We report on results of a systematic inter-comparison of ten global sea-ice concentration (SIC) data products at 12.5 to 50.0 km grid resolution for both the Arctic and the Antarctic. The products are compared with each other with respect to differences in SIC, sea-ice area (SIA), and sea-ice extent (SIE), and they are compared against a global winter-time near-100% reference SIC data set for closed pack ice conditions and against global year-round ship-based visual observations of the sea-ice cover. We can group the products based on the concept of their SIC retrieval algorithms. Group I consists of data sets using the self-optimizing EUMETSAT-OSISAF / ESA-CCI algorithms. Group II includes data using the Comiso-Bootstrap algorithm, and the NOAA-NSIDC sea-ice concentration climate data record (CDR). The standard NASA-Team and the ARTIST Sea Ice (ASI) algorithms are put into group III, and NASA-Team 2 is the only element of group IV. The three CDRs of group I (SICCI-25km, SICCI-50km, and OSI-450) are biased low compared to a 100% reference SIC data set with biases of -0.4% to -1.0% (Arctic) and -0.3% to -1.1% (Antarctic). Products of group II appear to be mostly biased high in the Arctic by between +1.0% and +3.5%, while their biases in the Antarctic range from -0.2% to +0.9%. Group III product biases are different for the Arctic: +0.9% (NASA-Team), -3.7% (ASI) but similar for the Antarctic: -5.4% and -5.6%, respectively. The standard deviation is smaller in the Arctic for the quoted group I products: 1.9% to 2.9% and Antarctic: 2.5% to 3.1%, than for group II and III products: Arctic: 3.6% to 5.0%, Antarctic: 4.0% to 6.5%. We refer to the paper to understand why we could not give values for group IV here. We discuss the impact of truncating the SIC distribution, as naturally retrieved by the algorithms around the 100% sea-ice concentration end. We show that evaluation studies of such truncated SIC products can result in misleading statistics and favour data sets that systematically overestimate SIC. We describe a method to re-construct the un-truncated distribution of SIC before the evaluation is performed. On the basis of this evaluation, we open a discussion about the overestimation of SIC in data products, with far-reaching consequences for, e.g., surface heat-flux estimations in winter. We also document inconsistencies in the behaviour of the weather filters used in products of group II, and suggest advancing studies about the influence of these weather filters on SIA and SIE time-series and their trends.

1 Introduction

For more than 40 years, the fraction of the polar oceans covered by sea ice, or sea-ice concentration, has been monitored by means of satellite microwave radiometry. This enabled a better understanding of ocean–sea-ice–atmosphere interactions for the polar regions where observations with other means than satellites are challenging due to remoteness, harsh environment and limited daylight. Based on the long-term satellite record, a substantial negative trend in the Arctic sea-ice area and extent has been found (e.g. Meier et al., 2014; Comiso et al., 2012, 2017a). In the Antarctic, sea-ice area and extent are highly variable

41 with a period of positive trend (Turner et al., 2013; Comiso et al., 2017b) and sea-ice extent maxima (Reid et al., 2015) being
42 followed recently by record minima (Schlosser et al., 2018; Turner et al., 2017; Stuecker et al., 2017).

43 In this contribution, we evaluate a number of satellite estimates of the sea-ice concentration from which sea-ice area
44 and extent are derived. Such detailed evaluation allows one to better estimate the uncertainties of these products, knowledge
45 of which is required for all their applications. These applications range from estimates of the future evolution of the Arctic
46 sea-ice cover, whose confidence is directly affected by observational uncertainty of sea-ice concentration (e.g., Niederrenk
47 and Notz, 2018), and short-term forecasts for ship routing (e.g., Wayand et al., 2019; Melia et al., 2017) to detailed climate-
48 model evaluation (e.g., Ivanova et al., 2017).

49 The sea-ice concentration is computed from satellite observations of the microwave brightness temperature (TB),
50 which is a measure of the Earth-leaving thermal microwave radiation received by the satellite sensor. A number of different
51 satellite sensors has been in place for sea-ice monitoring, summarized in Table 1 (see also Lavergne et al., 2019, Table 2).
52 With these sensors the polar regions are covered almost completely daily since October 1978 (every other day with Scanning
53 Multichannel Microwave Radiometer (SMMR) before July 1987).

54 A considerable number of different algorithms to compute the sea-ice concentration from microwave satellite TB
55 measurements has been developed during the past decades. All exploit the fact that under typical viewing angles (50-55
56 degrees) the difference in microwave brightness temperature, measured at horizontal (h) and vertical (v) polarization, between
57 open water and sea ice is sufficiently large to estimate sea-ice concentration. Whether or not a given algorithm is accepted by
58 the scientific community as a candidate for computing a climate data record (CDR), depends among other things on the length
59 of the available satellite raw data record, spatial and temporal resolution, quantification of uncertainties, and sensitivity to
60 noise which might introduce artificial trends (e.g., Tonboe et al., 2016; Lavergne et al., 2019).

61 Several inter-comparison studies were carried out to assess the quality of the sea-ice concentration obtained with different
62 algorithms (e.g., Andersen et al., 2007; Ivanova et al., 2014, 2015; Beitsch et al., 2015; Comiso et al., 2017a). Two different
63 kinds of such inter-comparisons exist. One kind deals with an inter-comparison of sea-ice cover products of a certain number
64 of algorithms without incorporating independent information of the sea-ice cover. Such inter-comparisons provide very
65 valuable information about inter-product consistencies in, e.g. the overall sea-ice concentration distribution, and in sea-ice area
66 and extent time series and trends. They also reveal differences, for instance, with respect to the representation of the seasonal
67 cycle or with respect to regional differences between sea-ice concentration estimates. Inter-comparisons of this kind are, e.g.,
68 Ivanova et al. (2014) and Comiso et al. (2017a). These studies, however, do not provide information about how accurate a sea-
69 ice concentration product is. The other kind of algorithm inter-comparison studies deals with the comparison of the satellite
70 sea-ice concentration with independent data. These can be ship-based observations, or sea-ice concentration estimates derived
71 from independent satellite observations, for instance, in the optical frequency range or with active microwave sensors such as
72 Synthetic Aperture Radar (SAR). Inter-comparisons of this second kind seldom involve more than one to two algorithms (e.g.,
73 Wiebe et al., 2009; Meier, 2005; Comiso et al., 1997; Comiso and Steffen, 2001; Markus and Dokken, 2002; Kern et al., 2003;
74 Cavalieri et al., 2010; Spreen et al., 2008). Exceptions to this are Andersen et al. (2007), who compared seven different
75 algorithms with ship-based sea-ice cover observations and SAR imagery for the high Arctic, and Beitsch et al. (2015), who
76 compared six different algorithms with ship-based sea-ice cover observations in the Antarctic. Both these studies each focused
77 on one hemisphere only. The work of Andersen et al. (2007) is relatively old and is based on comparably old versions of the
78 algorithms and products. In the present paper, we inter-compare the newest available versions of the sea-ice concentration
79 algorithm and products used in both studies, including three CDRs. We perform our evaluation for both hemispheres.
80 Additionally, we take advantage of a recently published new calibration / validation data package (see Section 2.2).

81 This paper is the first of a series of papers in which we are going to present and discuss results of a systematic evaluation
82 of ten sea-ice concentration products (see Sect. 2). We want to provide users and algorithm developers with new information
83 about the accuracy and precision of this suite of products, some of which are widely used in the climate research community.

84 In this paper, we present the sea-ice concentration products used. We focus on differences in sea-ice concentration, area and
85 extent, and on inter-comparisons with near-100% reference sea-ice concentrations and with a large suite of ship-based manual
86 visual observations of the sea-ice conditions. The second paper is going to focus on an inter-comparison with sea-ice
87 parameters derived from MODerate resolution Imaging Spectroradiometer (MODIS) satellite observations in the Arctic. The
88 third paper is going to focus on presenting and discussing results of an inter-comparison with sea-ice concentrations computed
89 from Landsat satellite visible imagery.

90 In the following Sect. 2, we introduce the sea-ice concentration data sets and ancillary data used as input. This section
91 further describes the preparation of the ancillary data and inter-comparison steps. Section 3 illustrates how the sea-ice
92 concentration products compare with each other in terms of multi-annual monthly average sea-ice concentration as well as
93 sea-ice area and extent. In Sect. 4 and Sect. 5, we show the results of the inter-comparison against a near-100% reference data
94 set and against ship-based sea-ice observations, respectively. Section 6 covers a discussion, an outlook, and conclusions.

95 **2 Data & Methodologies**

96 **2.1 Sea-ice concentration data sets**

97 For this study, we consider 10 different sea-ice concentration products (Table 2, with more details in Appendices A to
98 F). There are many more algorithms and products available than we are using here, see e.g. Ivanova et al. (2015). The main
99 criteria for our choice of algorithms and products are 1) length of the product time series, 2) grid resolution, 3) accessibility
100 and sustained production. We exclude products with less than ten years coverage and/or with a finer grid resolution than 12.5
101 km. Following Table 2 we comment on several specific issues that are important for the correct interpretation of sea-ice
102 concentration products, namely the grid resolution, the land spill-over correction, the weather / open water filter, and the sea-
103 ice concentration distributions around 0% and 100%.

104 We group the products according to their concept for sea-ice concentration retrieval (Table 2, column “Group”). Group
105 I contains the four EUMETSAT-OSISAF – ESA-CCI products. Group II contains the CBT-like algorithms, which are CBT-
106 SSMI; CBT-AMSRE and NOAA-CDR (the latter is a combination of CBT-SSMI and NT1-SSMI but is clearly dominated by
107 CBT-SSMI). NT1-SSMI and ASI-SSMI are assigned to group III. These algorithms follow a different concept to retrieve the
108 sea-ice concentration where the sea-ice concentration is mainly based on a brightness temperature polarization difference.
109 Finally, NT2-AMSRE is assigned to group IV; its concept to derive sea-ice concentrations via a look-up table and model
110 atmospheres is fundamentally different from the other nine algorithms.

111

112 **2.1.1 Grid resolution**

113 Given grid resolutions apply to every grid cell for group I products since their EASE grid has equal area of all grid
114 cells (App. A). For all other products which are provided on polar-stereographic grid (App. B through F), the grid resolution
115 is true at 70 degrees latitude (see also Peng et al., 2013). For the computation of sea-ice area and extent (Sect. 3), we take this
116 difference in grid-cell area into account and use the respective files of the grid-cell areas provided by NSIDC
117 (<ftp://sidacs.colorado.edu/pub/DATASETS/brightness-temperatures/polar-stereo/tools/geo-coord/grid>, last access date:
118 26/9/2018).

119

120 **2.1.2 Land spill-over correction**

121 The difference in brightness temperatures observed over open water (low) and land (high) combined with the size of
122 the field-of-view of several kilometres to a few tens of kilometres can cause spurious sea-ice concentrations to appear along
123 coasts (e.g., Lavergne et al., 2019). Various methods to reduce this so-called land spill-over effect are applied in all products
124 (Cavalieri et al., 1999; Cho et al., 1996; Maass and Kaleschke, 2010). For ASI-SSMI (App. B), reduction of land spill-over

125 effects is carried out for both the ASI algorithm as well as the NASA-Team algorithm product used over open water. For
126 NOAA-CDR (App. F), the reduction of land spill-over effects is applied separately to both input data sets before merging
127 (Meier and Windnagel, 2018). In this paper, we do not further correct potential differences between the ten products caused
128 by this effect.

129

130 **2.1.3 Weather / open water filter**

131 The two standard weather filters based on brightness temperature gradient ratios at 19 GHz, 22 GHz, and 37 GHz,
132 which mitigate noise due to atmospheric moisture and wind-induced roughening of the ocean surface (Cavalieri et al., 1995,
133 1999) are applied in the products NT1-SSMI and NT2-AMSRE. In the products CBT-SSMI and CBT-AMSRE, spurious sea-
134 ice concentrations caused by weather effects are filtered using the same frequencies as mentioned above but applying a
135 bootstrap technique (Comiso and Nishio, 2008). For NOAA-CDR (App. F), the above-mentioned weather filters are applied
136 before the merging (Meier and Windnagel, 2018). In the two NASA GSFC (App. C, D) sea-ice concentration products, i.e.,
137 NT1-SSMI version 1 and CBT-SSMI version 3 weather effects are reduced by screening of input brightness temperatures,
138 application of the above-mentioned weather filters, and some additional manual correction (Meier and Windnagel, 2018; Peng
139 et al., 2013, <https://nsidc.org/data/g02202/versions/3>, <https://nsidc.org/data/nsidc-0051>, and <https://nsidc.org/data/nsidc-0079>). In the ASI-SSMI product no specific weather filter is applied to the ASI algorithm itself. However, ASI algorithm sea-
141 ice concentrations are set to 0% where NASA-Team algorithm sea-ice concentration values are < 30% (see App. B). Because
142 the above-mentioned two weather filters are applied to the NASA-Team sea-ice concentration, the ASI-SSMI product
143 implicitly contains a weather filter as well (Ezraty et al., 2007). Note that the 5-day median filter used for the ASI-SSMI
144 product used here (Kern et al., 2010) not only removes remaining spurious sea ice over open ocean but also reduces weather-
145 induced elevated sea-ice concentrations along the ice edge. In the group I products a dynamic open water filter is applied. It is
146 based on the quoted standard weather filters but takes into account changes in filter efficiency due to changes in the frequencies
147 between the different sensors, for instance between SMMR and SSM/I. Also, it does not use the channels close to 22 GHz. All
148 weather filters may in addition to the spurious ice also remove real ice along the ice edge. All ten products apply a monthly
149 varying climatological sea-ice cover mask to erase spurious sea ice at low latitudes.

150 We investigate the temporal consistency of the weather filters. For this we focus on the sea-ice concentration interval
151]0.0%, 30.0%], i.e. exclude grid cells set to exactly 0.0% by the weather filter. Then, for each day of the month, we identify
152 the 5% percentile of all gridded sea-ice concentrations falling into the above-mentioned interval. Subsequently, we average
153 over the month. We look at two aspects. First, it is desirable that these time-series are mostly stable across the time-period
154 covered by a given data record, indicating that the weather filter cuts the sea-ice edge evenly across inter-annual variability
155 and changes of frequencies. Second, it is also desirable that the weather filter cuts “well below” the 15% SIC threshold that is
156 commonly used in the computation of the sea-ice extent (SIE) (e.g. Gloersen et al., 1992; Meier et al., 2014, Comiso et al.,
157 2017) but also of the sea-ice area (SIA) (e.g. Gloersen et al., 1992; Cavalieri and Parkinson, 2012; Ivanova et al., 2014). We
158 choose the 5% percentile (and not a minimum value) to obtain less noisy time-series. We plot examples of these time-series
159 for all ten products in Fig. 1 and Fig. 2 for the Arctic and Antarctic, respectively. For the Arctic we use March (Fig. 1 a) and
160 September (Fig. 1 g); for the Antarctic September (Fig. 2 a) and February (Fig. 2 g). In addition to the time-series, we also plot
161 the cumulative distribution of the daily sea-ice concentrations of the range]0.0% to 30.0%] for the respective month for the
162 year 2004 as an example for CBT-SSMI, OSI-450, NT1-SSMI, ASI-SSMI, and NT2-AMSRE.

163 We find little inter-annual variation of the monthly mean percentile sea-ice concentration over time particularly for
164 OSI-450 and SICCI-25km (see also Lavergne et al., 2019). Changes as caused, e.g., due to a switch in sensor remain below
165 1%. On average, the 5% percentile sea-ice concentration is < 12%, which ensures that the enhanced open water filter applied
166 in these two products barely influences computation of SIA and SIE. For group II products, these monthly mean 5% percentile
167 sea-ice concentrations are considerably larger and sometimes exceed 15%. Additionally, the time-series for CBT-SSMI and

168 NOAA-CDR reveal larger inter-annual variations than OSI-450, inter-sensor transitions (e.g. in 1987-1988 changing from
169 SMMR to SSM/I, Fig. 1 g), and in 2007-2008 changing from SSM/I to SSMIS, Fig. 1 a) leading to trends in the percentile
170 time-series. Compared to OSI-450 and SICCI-25km we find for NOAA-CDR: 1) The sea-ice concentration at which the
171 weather filter applies varies seasonally. For instance, in the year 1996, the mean 5% percentile of sea-ice concentrations within
172 the interval [0%, 30%] is for the Arctic 14% in March but 17% in September, and for the Antarctic 16% in September but 18%
173 in February. OSI-450 cuts at 10%, SICCI-25km at 11% in these months, and for both hemispheres. 2) The inter-annual
174 variation of the sea-ice concentration at which the filter applies is larger for NOAA-CDR than for OSI-450 and SICCI-25km.
175 The time-series for NT1-SSMI, ASI-SSMI, and NT2-AMSRE, in contrast, have very low (~1%) monthly mean 5% percentile
176 sea-ice concentration values with little or no inter-annual variation.

177 In the plots of the sample daily cumulative fraction for year 2004, we find a cumulative distribution with a first
178 increasing, later merely constant slope with no sea-ice concentrations below ~10% and ~8% for CBT-SSMI and OSI-450,
179 respectively (Fig. 1 b, c, h and i; Fig. 2 b, c, h and i). This agrees with the application of the open water filter presented in
180 Lavergne et al. (2019). For NT1-SSMI and NT2-AMSRE, in contrast, we find a substantial amount of near-0% sea-ice
181 concentrations (Fig. 1 d, f, j and l; Fig. 2 d, f, j and l). This suggests that while a weather filter is applied (according to the
182 documentations) there are still concentrations near 0% left. We checked this by looking at the respective daily sea-ice
183 concentration maps. Both products reveal a considerable number of grid cells with < 5% sea-ice concentration along the ice
184 edge. All but the group I products (see Table 2), only provide integer sea-ice concentration values. At the near-0% end of the
185 sea-ice concentration distribution these products have sea-ice concentrations 0%, 1%, 2%, and so forth. The number of NT2-
186 AMSRE sea-ice concentration values of 1% exceeds the 5% percentile most of the time which is the explanation why most
187 NT2-AMSRE values are missing in the time series in plots a) and g) of Fig. 1 and Fig. 2. We take the results shown in these
188 two figures into account when discussing the results presented in Sect. 3.

189

190 **2.1.4 Distribution around 100%**

191 A considerable fraction of this paper focuses on the evaluation near 100% sea-ice concentration. Sea-ice concentrations
192 are retrieved from satellite microwave brightness temperatures using a geophysical algorithm, usually involving tie points. Tie
193 points are typical signatures, e.g. brightness temperatures, or parameters derived from these, of ice (SIC: 100%) and open
194 water (Sic: 0%). Because of the natural variability of the surface properties of 100% sea ice relevant for its microwave remote
195 sensing, one fixed tie point value for 100% sea ice, even if retrieved daily, can only be an average representation of these
196 properties. In other words, ten different kinds of 100% sea ice can cause ten different brightness temperatures. As a result, a
197 retrieved sea-ice concentration naturally varies *around* 100%. This means even though the actual sea-ice concentration is
198 exactly 100% the retrieved one could be, for example, 97% or 100% or 103%. While the group I products retain the naturally
199 retrieved sea-ice concentration the others do not; in all group II to group IV products (see Table 2) sea-ice concentrations are
200 truncated at 0% and 100%, i.e. values < 0% are set to 0% and values > 100% are set to 100%. Figure 3 illustrates the sea-ice
201 concentration distribution at the locations of the near-100% sea-ice concentration reference data set (see Sect. 2.2) for SICCI-
202 25km, SICCI-50km and NOAA-CDR for the Arctic (plots a) to c)) and the Antarctic (plots d) to f)).

203 We use a Gaussian fit to reconstruct the true distribution of the sea-ice concentration retrieval around 100% for type II
204 and III product types. The methodology is tested on the group I products. This is done by finding that Gaussian curve which
205 provides the lowest root-mean-squared difference (RMSD) to the sea-ice concentration distribution for values $\leq 99\%$, i.e.,
206 basically the left hand side of the histograms shown in Fig. 3. For the fitting process, we also take into account the fraction of
207 sea-ice concentrations $\leq 99\%$ relative to the entire count of valid sea-ice concentrations: F_{99} . The difference between original
208 F_{99} and F_{99} resulting from the Gaussian fit, ΔF_{99} , has to be < 0.1 . We allow a maximum RMSD value of 0.0125. We first
209 binned SICCI-25km and SICCI-50km sea-ice concentration values to integer values to be consistent with the other products.
210 Figure 3 a, b and d, e) illustrate that the fits (red) agree well with the originally retrieved SICCI-25km and SICCI-50km sea-

211 ice concentrations (blue) with modal values slightly below 100%. Figure 3 c, f) illustrate how well the Gaussian fit matches
212 the original sea-ice concentration distribution for sea-ice concentrations $\leq 99\%$ for NOAA-CDR as an example. Here the modal
213 sea-ice concentrations of the Gaussian fit are larger than 100%: 103.5% for NH (Fig. 3 c) and 100.9% for SH (Fig. 3 f). In
214 addition, the Gaussian curve is broader than for SICCI-25km and SICCI-50km, resulting in larger values for the standard
215 deviation. We also note that F_{99} is ~ 0.6 and ~ 0.5 for SICCI-25km and SICCI-50km, respectively, but only ~ 0.2 and ~ 0.4 for
216 NOAA-CDR. We take the information from Fig. 3 into account when interpreting the results presented in Sect. 4 and Sect. 5.
217 We refer to Appendix H for the full set of sea-ice concentrations and Gaussian fits obtained for all ten products in both
218 hemispheres.

219 **2.2 The near-100% sea-ice concentration reference data set**

220 For the evaluation of the ten products at 100% sea-ice concentration (see Sect. 4), we use the Round Robin Data Package
221 version 2 (RRDP2) near-100% reference sea-ice concentration data set developed within the ESA Sea_Ice_cci and EU-SPICES
222 projects (Pedersen et al., 2019). In short, for this reference data set, areas of $\sim 100\%$ sea-ice concentration are found by
223 identifying areas of interest (AOI) of approximately 100 km x 100 km size with net convergence in the ice drift pattern on two
224 consecutive 1-day periods. Information about convergence is derived from the PolarView / MyOcean / CMEMS ice drift data
225 set derived from Envisat ASAR, RADARSAT-2 SAR and Sentinel-1 SAR imagery. By choosing AOIs for regions with high
226 concentrations, near 100% sea-ice concentration can be assured (e.g., Kwok, 2002; Andersen et al., 2007) in winter. Each AOI
227 contains up to hundred 10 km x 10 km cells for which the SAR ice drift is computed. The number of cells depends on SAR
228 image coverage. Convergence in the ice drift pattern results in a decrease in the total area of these cells. A cell is included in
229 the dataset of $\sim 100\%$ sea-ice concentration if the area reduction between day 1 and day 2 is between 0.4% and 1.5% and if
230 more than 40% of the AOI contains cells with such an area reduction. The RRDP2 near-100% sea-ice concentration reference
231 data set contains the AOI centre geographic latitude and longitude, time, total sea-ice concentration (100%) and AOI average
232 area reduction due to net ice convergence. It is available for years 2007 through 2015 for both hemispheres.

233 We cannot provide a definite uncertainty for this reference data set but for its production, we combine a suite of
234 measures to ensure high precision and close-to-zero bias (high accuracy). The drift/convergence selection is based on
235 convergence on two consecutive days of 1-day drift. During winter, i.e., November through March (Arctic) and May through
236 September (Antarctic), this is assumed to ensure that all openings existing on day 0 (prior to the two convergence days) are
237 closed by convergence or refrozen. The refreezing assumption is the reason for the product quality to be higher during winter
238 when openings rapidly refreeze than during summer when openings may not freeze up. There is no prior assumption of the
239 initial ice concentration (on day 0), but the ice-drift product requires quite high concentrations for the 2-D cross correlation to
240 work. Andersen et al. (2007) reported an accuracy of 2% for ice-water SAR image classification from ice analysts without
241 additional drift / convergence information and a sea-ice concentration standard deviation of $\sim 1\%$ for own cold-season high-
242 resolution high-quality SAR image classification. Based on our above-mentioned measures and the results of Andersen et al.
243 (2007) we can state estimated values for precision: $\sim 1\%$ and accuracy: $< 0.5\%$ for our reference data set.

244 We co-locate the sea-ice concentrations of the ten products with the selected AOI grid cells by computing the minimum
245 distance between AOI grid cell centre and grid cell centre of the respective sea-ice concentration product. For this step, we
246 convert the geographic coordinates of all data sets into Cartesian coordinates taking into account the different projections.
247 Figure 4 illustrates the spatial distribution of selected AOIs for both hemispheres for two different years by showing the co-
248 located OSI-450 sea-ice concentration. We give an example of a typical “good distribution” (Fig. 4 a, c) and for a typical “poor
249 distribution” (Fig. 4 b, d). The RRDP2 near-100% sea-ice concentration reference data set contains basically no AOIs in the
250 Eastern Antarctic because of the lack of SAR image coverage required for the ice drift product used to generate this RRDP2
251 data set.

252 We evaluate the products at their native grid resolution without applying any spatial averaging. For each product, we
253 compute the mean difference “product minus 100%” and its standard deviation as well as the cumulative distribution function
254 of the differences.

255 **2.3 Ship-based visual sea-ice cover observations**

256 According to the Antarctic Sea Ice Processes & Climate (ASPeCt) protocol <http://www.aspect.aq> (Worby and Allison,
257 1999; Worby and Dirita, 1999, see also Worby et al., 2008) and the IceWatch/ASSIST (Arctic Ship-based Sea-Ice
258 Standardization) protocol (<http://icewatch.gina.alaska.edu>), ship-based observations of the sea-ice conditions shall be carried
259 out every hour, at least every second hour, during daylight conditions while the ship is traversing the sea ice. Observations
260 shall be carried out from the ship’s bridge for an area of about one kilometre around the ship and shall report ice conditions as
261 follows: total ice concentration, type of openings, and concentration, thickness, ridge fraction and height, and snow depth and
262 type for up to three ice types. All ship-based visual observations used here result from manual, non-automated observations.

263 For our evaluation of the ten products with respect to ship-based visual observations of the sea-ice conditions (see Sect.
264 5), we use about 15 000 individual observations. About ~7000 of these were carried out in the Antarctic (ASPeCt) and ~8000
265 in the Arctic (IceWatch/ASSIST). A substantial fraction of the Antarctic observations (until 2005) is available via
266 <http://www.aspect.aq> (Worby et al., 2008). The more recent observations were collected from various sources (e.g.
267 PANGAEA, ACE-CRC, AWI, see also Beitsch et al., 2015) and merged with the existing ASPeCt data. The majority of the
268 ASSIST data is taken from the data portal <http://icewatch.gina.alaska.edu>. Additional sources for ASSIST data are PANGAEA
269 (for Polarstern cruises before IceWatch/ASSIST), the Arctic Data Center of the NSF: <https://arcticdata.io/catalog/#data>, and
270 the data archive of the Bering Sea Ecosystem Study (BEST): <https://www.eol.ucar.edu/projects/best/ice.shtml>. All data are
271 standardized, i.e. the ascii format data files containing the observations use similar formats for all variables and missing data.
272 The data are also manually quality checked for outliers. For the comparison presented in this manuscript, we use all ASPeCt
273 and IceWatch/ASSIST observations from the period June 2002 through December 2015. This data set is available via
274 <http://icdc.cen.uni.hamburg.de/1/daten/cryosphere/seaiceparameters-shipobs>, last access date: 28/3/2019.

275 Figure 5 summarizes the locations of the ship-based observations used in this manuscript, separately for the Arctic (Fig.
276 5 a, b) and the Antarctic (Fig. 5 c, d). For both hemispheres, just comparably small regions contain such observations. Figure
277 5 a, c) illustrates that some regions are visited during several years while others just once or twice during the 13-year period
278 considered. The seasonal distribution (Fig. 5 b, d) illustrates that the more central (Arctic) or southern (Antarctic) regions were
279 only visited during summer months due to harsh winter conditions and missing daylight for these regions.

280 We co-locate the sea-ice concentrations of the ten products with the selected ship-based observations by computing the
281 minimum distance between geographic location of the ship-based observation and the grid cell centre of the respective sea-ice
282 concentration product. For this step, we convert the geographic coordinates of all data sets into Cartesian coordinates taking
283 into account the different projections of the sea-ice concentration products. Following the co-location, we average over all
284 ship-based and all satellite-based sea-ice concentration values, including reports of open water, i.e. 0% concentration, of one
285 day following the approach of Beitsch et al. (2015). This results in a comparison of along ship-track daily average sea-ice
286 conditions. Data pairs with less than three ship-based observations per day are discarded. The results of the comparison
287 between ship-based and satellite-based sea-ice concentration are solely based on these daily average sea-ice concentrations.
288 Note that all satellite-based data are used at their native grid resolution.

289 The ship-based and satellite-based sea-ice concentration data sets are inter-compared (Sect. 5) by means of scatterplots
290 and linear regression analysis and statistics separately for summer data, winter data, and data of the entire year. Summer
291 comprises months May through September for the Arctic and months November through March for the Antarctic; winter
292 comprises the respective remaining months. In the scatterplots, we compare the daily average sea-ice concentrations and
293 additionally compute averages of the satellite-based sea-ice concentration for each of the above-mentioned bins applied to the

294 ship-based data and vice versa. We compute the overall average sea-ice concentration difference and its standard deviation,
295 and perform a linear regression analysis based on the daily average and the binned data.

296 **3 Inter-comparison of sea-ice area, extent, and distribution**

297 We follow Ivanova et al. (2014) and begin our inter-comparison with time-series of the sea-ice area (SIA) and sea-ice
298 extent (SIE) (Subsection 3.1 and 3.2) derived from monthly mean sea-ice concentration. The monthly mean sea-ice
299 concentration is derived for every product at the native grid and grid resolution using data of all days of a month of the entire
300 sea-ice concentration range including 0%. SIE is computed by summing over the grid-cell area of grid cells with > 15% sea-
301 ice concentration. SIA is computed by summing over the ice-covered portion of the grid-cell area of grid cells with > 15% sea-
302 ice concentration. By using this threshold, we follow Gloersen et al. (1992) and numerous SIA and SIE inter-comparison
303 studies. We compare SIA and SIE time-series for the entire period for which we have data from the respective products at the
304 time of the analysis. We exclude sea-ice concentrations estimated for lakes and other inland waters. We fill the circular area
305 with missing data around the pole caused by the satellite orbit inclination and swath width with a constant sea-ice concentration
306 value of 98%. Andersen et al. (2007) found a mean sea-ice concentration of ~98% from a comparison of cold season passive
307 microwave and synthetic aperture radar observations in the high Arctic. They noted a smaller value of ~95% in summer. Both
308 values are confirmed by own work (Kern, 2018, [http://icdc.cen.uni-hamburg.de/fileadmin/user_upload/ESA_Sea-Ice-
309 ECV_Phase2/SICCI_Phase2_SIV-Retrieval_Report_v02.pdf](http://icdc.cen.uni-hamburg.de/fileadmin/user_upload/ESA_Sea-Ice-ECV_Phase2/SICCI_Phase2_SIV-Retrieval_Report_v02.pdf), last accessed September 6, 2019). Using 98% instead of 95%
310 during summer results in an overestimation of the SIA of about 10 000 km², a small value compared to other sources of biases
311 for the SIA during summer. This filling is applied to the Arctic and only to the products at polar-stereographic projection. This
312 area is already interpolated spatially in the four group I products. As described in Lavergne et al. (2019), these products contain
313 a fully filtered and truncated to the range [0.0% ... 100.0%] version and a non-filtered, non-truncated version of the sea-ice
314 concentration. The latter contains the naturally retrieved sea-ice concentrations, i.e., also values < 0% and > 100% (see Sect.
315 2.1.3) and no weather filters are applied (see Sect. 2.1.3). We use the fully filtered and truncated version. Without the truncation
316 the SIA of the group I products increases slightly, while the SIE does not change because the number of grid cells covered
317 with > 15% sea ice is not influenced by the truncation (not shown).

318 We complement these SIA and SIE time series with maps of the multi-annual average sea-ice concentration difference
319 for selected months for the AMSR-E measurement period: June 2002 to September 2011 (Subsection 3.3 and 3.4). We choose
320 this period to be able to compare all ten products for a similarly long time-period. For these maps, we first re-grid the monthly
321 mean sea-ice concentrations of all products, except SICCI-50km, onto the EASE grid version 2.0 with 50 km grid resolution
322 using bilinear interpolation. Then we compute the multi-annual average sea-ice concentration for each month from which we
323 subsequently calculate an ensemble median and the difference product minus ensemble median.

324 **3.1 Arctic sea-ice area and extent time-series**

325 The SIA and SIE time-series for the Arctic reveal a very similar overall development for the products extending back
326 into the 1980s and 1990s (Fig. 6). This applies to: i) the overall negative trend in both quantities; ii) the inter-annual variability
327 as, for instance, during 1991-1997 and around 2007 and 2012 for SIA and SIE in September (Fig. 6 b, d), or in 2011 for the
328 relative minimum in March SIA and SIE (Fig. 6 a, c); iii) to the ranking between the products. NOAA-CDR and CBT-SSMI
329 provide largest SIA and SIE in both March and September. NT1-SSMI provides lowest SIA and SIE in September, while in
330 March we find SIA and SIE from ASI-SSMI to be even lower. It is obvious that differences between products are smaller for
331 SIE than for SIA as was shown also by Ivanova et al., (2014). For the AMSR-E period, when SIA and SIE of all ten products
332 are available, the inter-annual variation is similar for all ten products. We find ESA-CCI products, CBT-AMSRE and NT2-

333 AMSRE, fall into the ranges of SIA and SIE given by the other products. An exception to this is SICCI-50km, providing
334 clearly the lowest SIE of all products in March (Fig. 6 c). We will discuss this finding in Sect. 3.3.

335 **3.2 Antarctic sea-ice area and extent time-series**

336 The SIA and SIE time-series for the Antarctic reveal a similar overall development for products extending back into the
337 1980s and 1990s (Fig. 7). This applies to i): the overall positive trend until 2015 in both quantities; ii) the inter-annual
338 variability, for instance, during 2000-2003 for SIA and SIE in February (Fig. 7 b, d), or in 2008 for the relative minimum in
339 SIA and SIE in September (Fig. 7 a, c); iii) the ranking between products. NOAA-CDR and CBT-SSMI provide largest SIA
340 while NT1-SSMI provides smallest SIA, being ~ 1.8 million km^2 and 300 000 km^2 below the SIA of NOAA-CDR and CBT-
341 SSMI in September and February, respectively. OSI-450 and CBT-SSMI provide the largest SIE, exceeding the smallest SIE:
342 NT1-SSMI by ~ 500 000 km^2 and ~ 300 000 km^2 in September and February, respectively (Fig. 7 c, d). Inter-product differences
343 are larger for SIA than SIE in September but not in February. For the AMSR-E period, the inter-annual variation is similar for
344 all ten products. In September, SIA and SIE of the ESA-CCI products mostly fall between ASI-SSMI and OSI-450 (Fig. 7 a,
345 c). In February (Fig. 7 b, d), SICCI-50km provides smallest overall SIA and SIE while the SICCI-12km and SICCI-25km
346 products agree closely with OSI-450. In contrast to the Arctic (compare Fig. 6), NT2-AMSRE clearly provides largest SIA in
347 September and February (Fig. 7 a, b). NT2-AMSRE provides largest SIE in February as well (Fig. 7 d).

348 Figure 7 c) contains an example of discontinuities caused by the application of weather filters not adapted to sensor
349 changes. Wintertime Antarctic OSI-450 and CBT-SSMI SIE agree with each other for the SMMR period. After 1987 the CBT-
350 SSMI SIE is below the OSI-450 SIE by ~ 150 000 km^2 . This corresponds to the area of one quarter of all 25 km grid cells at a
351 latitude of 60°S, the approximate average location of the Antarctic sea-ice edge in September. This change in SIE is
352 concomitant with a jump in the weather-filter sea-ice concentration from 11% to 14% (Fig. 2 a). It is noteworthy that the
353 23.0 GHz channels of the SMMR instrument have been highly unstable since launch, and eventually ceased to function on 11
354 March 1985 (Njoku et al., 1998). Thus, the water-vapour part of the “classic” weather filter is un-reliable in the early decade
355 of the satellite data record. This is solved in the OSI-450 product by relying on explicit atmospheric correction of the brightness
356 temperatures using -among others- water vapour fields from atmosphere re-analysis (see Lavergne et al., 2019). Another
357 example of this kind (not shown), is a shift between OSI-450 SIE and NOAA-CDR SIE by 50 000 km^2 to 100 000 km^2 between
358 2007 and 2008. This shift is concomitant with a discontinuity in the weather-filter sea-ice concentration for NOAA-CDR in
359 March (Arctic, Fig. 1 a) and September (Antarctic, Fig. 2 a) during the transition from 2007 to 2008. This corresponds to when
360 SSMIS (F17) is processed instead of SSM/I (F15). OSI-450 exhibits no discontinuity here.

361 **3.3 Arctic sea-ice concentration distribution differences**

362 In March, the difference between the sea-ice concentration of an individual product and the ensemble median of all ten
363 products (Fig. 8) remains within $\pm 5\%$ over most of the Arctic Ocean, except for ASI-SSMI (Fig. 8 d) and SICCI-12km (Fig.
364 8 f). The largest differences between individual products and the ensemble median are located in the peripheral seas. Group II
365 and IV (see Table 2) products have more sea ice than the ensemble median (Fig. 8 b, c, i, j) differences can exceed 20%. NT1-
366 SSMI (Fig. 8 a) has less sea ice than the ensemble median with negative differences greater than 10% or even 15% in magnitude
367 in all peripheral seas. Differences of OSI-450 or SICCI-25km and the ensemble median are within $\pm 5\%$ almost everywhere
368 (Fig. 8 e, g). SICCI-50km exhibits negative differences greater than 20% in magnitude along some of the coastlines, e.g., the
369 Labrador Sea, the Irminger Sea or the coastlines of the Pechora Sea and Barents Sea (Fig. 8 h). A careful check of these areas
370 in daily and monthly mean maps of the SICCI-50km sea-ice concentration reveals, that for regions with a relatively narrow
371 sea-ice cover stretching along coastlines, the coarse resolution of the 6.9 GHz frequency channel entering the algorithm (see
372 Table 2), in combination with land spill-over filter and open water filter can result in an unwanted complete removal of sea ice

373 from the grid cells of the product. We are confident that this explains the particularly low SICCI-50km SIA and SIE shown in
374 Fig. 6 a) and c) for Arctic SIE in March.

375 For the matrices shown in Fig. 9 (and Fig. 12, Appendices G1 to G6), we re-grid the monthly mean sea-ice concentration
376 onto the EASE grid version 2.0 with 50 km grid resolution using bilinear interpolation and apply a common land mask (the
377 one of SICCI-50km) to all products. The differences between SIA and SIE values are computed from these gridded 50 km
378 resolution, common land mask products. Figure 9 a) agrees with the results shown in Fig. 8. In winter (March), sea-ice
379 concentration differences between members of groups I to III are < 1%. Group III members NT1-SSMI and ASI-SSMI (see
380 Table 2) exhibit less sea ice than the other groups. Group II and IV members exhibit higher sea-ice concentrations than the
381 other two groups in winter and summer (September). In summer (Fig. 9 d), sea-ice concentration differences remain < 2%
382 between members of groups I and II but not within group III where differences exceed 10%. We refer to App. G for the
383 respective results of the other months.

384 **3.4 Antarctic sea-ice concentration distribution differences**

385 In September (Fig. 10), most products show more sea ice than the ensemble median over high-concentration ice and less
386 sea ice along the marginal ice zone. NT1-SSMI (Fig. 10 a) exhibits considerably less sea ice than the ensemble median almost
387 everywhere (see also Fig. 12 a). ASI-SSMI (Fig. 10 d) exhibits a distribution of sea-ice concentration differences that is
388 reversed compared to most other products. CBT-AMSRE and especially NT2-AMSRE (Fig. 10 i, j) show more sea ice than
389 the ensemble median for most regions. This is evident in Fig. 12 a) as well.

390 In January (Fig. 11), the few, comparably small, high-concentration areas exhibit sea-ice concentration differences mostly
391 below $\pm 5\%$. Over the lower concentration areas, i.e. mainly in the Weddell Sea and the Ross Sea, most products show less sea
392 ice than the ensemble median with differences between 0 and -6%. NT1-SSMI has less sea ice while ASI-SSMI and particularly
393 NT2-AMSRE have more sea ice (see also Fig. 12 d). Important to note are the negative differences of $\sim -10\%$ along most of
394 the Antarctic coast for SICCI-50km (Fig. 11 h). Like for the Arctic, the coarse resolution of the 6.9 GHz frequency channels
395 combined with land-spill over and open water filters can result in the unwanted removal of sea ice from grid cells (compare
396 Fig. 8 h) and discussion of it) with the same influence on SICCI-50km SIA and SIE values (Fig. 7 b, d). This is also evident
397 in Fig. 12 d) where SICCI-50km exhibits the largest inter-product differences within group I.

398 For the results shown in Fig. 6 through Fig. 12 we used the truncated sea-ice concentration values as far as it concerns
399 group I products. Repeating these computations with the non-truncated values, e.g. in September and March, does not change
400 the results with respect to SIE. SIA increase by $\sim 50\,000\text{ km}^2$ in winter months and there is almost no impact in summer.

401 **3.5 Summary and discussion of sea-ice area and extent findings**

402 The partly substantial inter-product mean sea-ice concentration differences (Figures 8 through 12) are associated with a
403 notable impact on SIA and SIE (Fig. 9 and Fig. 12). For summer SIE, inter-product differences are below $\sim 200\,000\text{ km}^2$
404 (Arctic, Fig. 9 f) and below $\sim 300\,000\text{ km}^2$ (Antarctic, Fig. 12 f) for most products. NT1-SSMI (Arctic) and SICCI-50km, NT1-
405 SSMI, NT2-AMSRE (Antarctic) show the largest differences here. For winter SIE, most inter-product differences are below
406 $\sim 200\,000\text{ km}^2$ for the Antarctic (Fig. 12 c); larger differences are mostly found for NT1-SSMI (see also Fig. 7 c). For the Arctic
407 (Fig. 9 c), inter-product differences range between $100\,000\text{ km}^2$ and $600\,000\text{ km}^2$ and seem to be associated with the type of
408 algorithm and partly also the sensor.

409 For Arctic winter SIA (Fig. 9 b), group III products provide systematically smaller values (by $\sim 400\,000\text{ km}^2$) than the
410 other three groups while group II SSMI products provide systematically larger values (by $\sim 300\,000\text{ km}^2$). Group I and CBT-
411 and NT2-AMSRE exhibit the lowest inter-product SIA differences. For Arctic summer SIA (Fig. 9 e), group II and IV products
412 agree with each other within $100\,000\text{ km}^2$ but exceed group I SIA by $\sim 400\,000\text{ km}^2$. This equals to 10% of the summer
413 minimum Arctic SIA. For Antarctic summer SIA (Fig. 12 e), NT2-AMSRE exceeds SIA of all other products by $\sim 400\,000$

414 km². This equals 20% of the summer minimum Antarctic SIA. SIA differences between the other products are almost all <
415 200 000 km². For Antarctic winter SIA (Fig. 12 b), NT2-AMSRE and NT1-SSMI stand out with very large systematic
416 differences to all other products.

417 We summarize our findings from the matrices in Fig. 9 and Fig. 12 (see also App. G1 to G6) as follows: i) Absolute and
418 relative (to the respective minimum or maximum value) inter-product differences are smaller for SIE than for SIA. ii) SIA and
419 SIE derived from products of different algorithms of the same group (see Table 2) may differ considerably. iii) Inter-product
420 differences for SIA and SIE for the investigated CDRs are, on average, larger for the Arctic than the Antarctic.

421 We note that the grid resolution of the products is not necessarily compatible with the true spatial resolution because the
422 footprints of the satellite sensor channels used in some of the algorithms is coarser (Table 2). This applies to NT1-SSMI, CBT-
423 SSMI, NOAA-CDR, and OSI-450, i.e. the products at 25 km grid resolution based on SMMR, SSM/I, and SSMIS data. For
424 these products, we expect that gradients in the sea-ice concentration are more smeared than for products with a better match
425 between footprint size and grid resolution as, for instance, CBT-AMSRE or SICCI-25km. This is illustrated for OSI-450 and
426 SICCI-25km in Lavergne et al. (2019, Figure 6). For a typical wintertime Antarctic ice edge at 65°S comprising half a compact
427 and half an open sea-ice cover, this difference in the match of true resolution and grid resolution between SSM/I and AMSR-
428 E products would result in a slightly larger SIE (by ~200 000 km²) derived from the SSM/I product. This is because a compact
429 ice edge is smeared more in the SSM/I product, resulting in more grid cells with a sea-ice concentration > 15%, the threshold
430 used currently to compute SIE. In fact we find that during winter OSI-450 SIE exceeds SICCI-25km SIE by ~100 000 km² in
431 the Arctic (Fig. 9 c) and by ~200 000 km² in the Antarctic (Fig. 12 c). Since the algorithms used for the sea-ice concentration
432 retrieval for these two products are almost identical, the difference in SIE can well be attributed to the above-mentioned impact
433 of differences between true and grid resolution. The second pair of almost identical algorithms is CBT-SSMI and CBT-
434 AMSRE. In the Arctic, in winter, CBT-SSMI SIE exceeds CBT-AMSRE SIE by ~400 000 km² (Fig. 9 c) but in the Antarctic
435 CBT-SSMI SIE is smaller than CBT-AMSRE SIE by ~100 000 km² (Fig. 12 c). Differences in the algorithm itself and/or in
436 the weather filter might be the cause. We refer to the discussion at the end of Sect. 3.2 in this context.

437 **4 Comparison with near-100% SIC reference data set**

438 In this section, we present results of the evaluation of the ten products at 100% sea-ice concentration using the data
439 described in Sect. 2.2. We note upfront that caution should be exercised when reporting and interpreting evaluation statistics
440 like bias (mean value minus 100%) or root mean square error, near the 100% end of truncated sea-ice concentration products.
441 This applies to the results presented in this section but also to Sect. 5. First, the bias of truncated products will necessarily be
442 negative or zero (mean value lower than or equal to 100%), even if the bias of the product was exactly 0% before truncation.
443 Second, products whose non-truncated distribution is biased high (modal value larger than 100%) will seemingly achieve
444 better evaluation statistics after truncation, because of the accumulation of values > 100% being folded to exactly 100%. Both
445 bias and RMSE of these products are smaller than those of products that do not overestimate at 100% sea-ice concentration.
446 The larger the fraction of truncated values, the better the statistics. The values that accumulate at the 100% bin in the truncated
447 product are in majority from the above 100% range of the non-truncated distribution (note the value of F_{99} in Fig. 3), and
448 improve the evaluation statistics (bias and RMSE). In fact, under the hypothesis that the distribution of the retrieved sea-ice
449 concentration is mostly Gaussian around the modal value before truncation, products with overestimation of the non-truncated
450 distribution can be recognized by their abnormal (with respect to a Gaussian model) accumulation of values exactly at 100%.
451 For more discussion and quantification of the error distribution of sea-ice concentration products and algorithms before
452 truncation around 100% (and around 0%), see Ivanova et al. (2015).

454 The distribution of the sea-ice concentration near 100% is shown for the Arctic for each product in Fig. 13 a) in form
 455 of the cumulative fraction of the deviation (bias) from 100%. As expected, the cumulative fraction increases towards 1.0 for
 456 all products. Considerable differences between the products are evident. Group I products (see Table 2) exhibit a cumulative
 457 fraction between ~ 0.5 and ~ 0.7 in bin $-1.5\% \dots -0.5\%$. Group II products and NT1-SSMI exhibit substantially lower fractions
 458 for this bin: between ~ 0.2 (NOAA-CDR, see also F_{99} in Fig. 3 c) and ~ 0.4 (NT1-SSMI). Consequently, the change in
 459 cumulative fraction to the last bin is between 0.6 and 0.8 for group II and NT1-SSMI but only between 0.3 and 0.5 for group
 460 I. The respective F_{99} values in Fig. 3 and App. H suggest that a large portion of the cumulative fraction in the last bin is in fact
 461 due to sea-ice concentrations $> 100\%$ being set to 100% (i.e. truncated). For NT2-AMSRE, the cumulative fraction increases
 462 abruptly from < 0.1 in bin $-2.5\% \dots -1.5\%$ to ~ 0.8 in bin $-1.5\% \dots -0.5\%$. This behaviour is completely different to all other
 463 products and confirms the results of Andersen et al. (2007) and Ivanova et al. (2015). For their inter-comparison of sea-ice
 464 concentration algorithms in the high Arctic, Andersen et al. (2007) extended the range within which the NT2 algorithm permits
 465 to retrieve the sea-ice concentration to 120%, in order to be able to properly compare all algorithms with respect to their
 466 precision close to 100%. The original implementation of the NT2 algorithm is constrained to sea-ice concentrations up to
 467 100%. Ivanova et al. (2015) also aimed to compare the precision of several sea-ice concentration algorithms at 100% by means
 468 of computing the standard deviation of the sea-ice concentration at 100%. They were, however, unable to obtain standard
 469 deviations with a comparable statistics because of large positive biases for the, e.g., NT2 and ASI algorithms and because of
 470 the truncation of sea-ice concentrations at 100%. Only by constructing a reference sea-ice concentration of 75%, Ivanova et
 471 al. (2015) were able to carry out a comparison of the sea-ice concentration standard deviation based on non-truncated sea-ice
 472 concentrations.

473 These cumulative distributions suggest that an inter-comparison of the mean difference to 100%, i.e. the bias, and its
 474 standard deviation, i.e. the precision, should be carried out by means of the Gaussian fit proposed in Subsection 2.1.4.
 475 Consequently, the mean sea-ice concentrations of the ten products shown by the black symbols in Fig. 14 a) for the Arctic
 476 near-100% reference sea-ice concentration locations are the modal values of the Gaussian fits. The error bars denote one
 477 standard deviation of this fit around the modal value. The blue symbols and error bars denote the respective mean and one
 478 standard deviation computed from the non-truncated group I sea-ice concentration products. All values shown here are
 479 summarized together with the results obtained from the truncated sea-ice concentration products in Table 3.

480 Figure 14 a) confirms our hypothesis that those products where the cumulative distributions seem to contain a large
 481 fraction of sea-ice concentrations larger than 100% set to 100% (Fig. 13 a), i.e. the group II products, and NT1-SSMI, are
 482 likely to over-estimate the actual sea-ice concentration. The over-estimation is particularly high for NOAA-CDR with a modal
 483 sea-ice concentration of $\sim 103\%$ (see also Table 3). The group I products exhibit modal sea-ice concentrations that are slightly
 484 below and closest to the near-100% reference. The Gaussian fit almost perfectly matches the actually observed non-truncated
 485 sea-ice concentration for SICCI-50km also in terms of the standard deviation (compare blue and black symbols in Fig. 14 a)
 486 and Table 3). The match is less accurate for OSI-450 and SICCI-25km.

487 In addition, Fig. 14 a) and Table 3 confirm our hypothesis that the group II products and NT1-SSMI are likely to have
 488 a standard deviation that is lowered by truncating sea-ice concentrations to a maximum of 100%. This is illustrated by the
 489 standard deviations of the non-truncated and the truncated sea-ice concentration for group I products (Table 3). Under the
 490 assumption that the standard deviation of the Gaussian fit is a better measure of the spread in sea-ice concentrations near (but
 491 below) 100%, we state that the CDRs of group I: SICCI-50km, OSI-450, and SICCI-25km allow a better precision than group
 492 II products. We also confirm the findings of Lavergne et al. (2019) that the group I products are slightly – but significantly –
 493 biased low.

494 For ASI-SSMI and NT2-AMSRE, the application of a Gaussian fit is potentially not justified given the way sea-ice
 495 concentrations are retrieved. ASI-SSMI is non-linear near 100% (Kaleschke et al., 2001) while no statement about the

496 functional relationship of the input satellite data and the retrieved sea-ice concentrations can be made for NT2-AMSRE
497 (Markus and Cavalieri, 2000; Brucker et al., 2014). Our analysis, however, results in a reasonable Gaussian fit for ASI-SSMI
498 (see App. H, Fig. H1 a). We are confident therefore that the values taken from the fit and shown in Fig. 14 a) and Table 3 can
499 be used. In contrast, the NT2-AMSRE sea-ice concentration distribution does not allow reasonable application of a Gaussian
500 fit (see App. H, Fig H1 j) and we do not report it along with the others in Fig. 14 a). We note that Andersen et al. (2007)
501 reported a mean NT2 sea-ice concentration of $\sim 105\%$ with a standard deviation of $\sim 5\%$ at near-100% sea-ice concentrations
502 for their unconstrained version of this algorithm.

503 **4.2 Antarctic**

504 The cumulative fraction of the deviation (bias) from 100% fFor the Antarctic (Fig. 13 b), we find more similarities in
505 the cumulative fractions of the deviation from 100% than for the Arctic. Common to all products, except NT2-AMSRE and
506 ASI-SSMI, is a notable step change in the cumulative fraction between bin $-1.5\% \dots -0.5\%$ and the last bin. This step change
507 is least pronounced for NT1-SSMI and most pronounced for NOAA-CDR exhibiting cumulative fractions of ~ 0.8 and ~ 0.4 ,
508 respectively, in the last bin. Step changes are slightly larger for the two CBT products than for group I products (see also F_{99}
509 values in Fig. 3 d), e)). For ASI-SSMI the cumulative fraction levels off before the last bins (see also the F_{99} value App. H,
510 Fig. H2 a). 93% of the ASI-SSMI sea-ice concentrations at the near-100% reference sea-ice concentration locations are below
511 99.5%. The distribution for NT2-AMSRE is extremely narrow. About 75% of the NT2-AMSRE sea-ice concentrations at the
512 near-100% reference sea-ice concentration locations are above 99.5%. This behaviour is, like for the Arctic, completely
513 different to all other products and agrees with the findings of Ivanova et al. (2015), see also our discussion in Sect. 4.1.

514 Group I and II products provide a modal sea-ice concentration which deviates by less than $\sim 1\%$ from 100% (Fig. 14
515 b), Table 4). While group II products tend to exhibit a modal sea-ice concentration $> 100\%$, group I products exhibit modal
516 sea-ice concentrations $< 100\%$. Modal sea-ice concentrations obtained with the Gaussian fit agree within 0.5% to the actually
517 measured mean sea-ice concentration derived from the non-truncated values of group I products (compare black and blue
518 symbols in Fig. 14 b). Respective standard deviations match better in the Antarctic than the Arctic and are systematically
519 smaller for the truncated than the non-truncated results (compare black and blue bars Fig. 14 a) and b) as well as Table 3 and
520 4). For group II products using the Gaussian fit method suggests standard deviations between $\sim 4.5\%$ and $\sim 5.5\%$ while using
521 the truncated values results in a standard deviation between 2.0% and 2.5%. Following our assumption in Sect. 4.1, we state a
522 better precision for group I products: SICCI-50km, SICCI-25km, and OSI-450 than group II products. Similarly but to a lesser
523 extent than in the Arctic, group I products are slightly biased low with respect to the validation data set.

524 **4.3 Summary and discussion of the evaluation near 100%**

525 In near-100% sea-ice concentration conditions, most retrieval algorithms will naturally retrieve some distribution of
526 values around 100% sea-ice concentration, i.e. also values $> 100\%$. However, these values are almost never accessible to the
527 user, and thus generally not accessible for validation. Here, we used the availability of these “off-range” estimates in the group
528 I products to demonstrate how the entire distribution around 100% can effectively be reconstructed a-posteriori from the
529 products with truncated sea-ice concentration distributions (Fig. 3, App. H). This Gaussian-fit methodology allows us to go
530 deeper in the analysis of near-100% conditions. Indeed, if the analysis had been limited to the truncated distributions only (Fig.
531 13), algorithms that over-estimate sea-ice concentration (modal value of the non-truncated distribution larger than 100%)
532 would obtain better validation statistics (smaller bias and RMSE) than products without such over-estimation. The larger the
533 over-estimation, the better the statistics would be. Using the Gaussian fit, we unveil a possible over-estimation of several sea-
534 ice concentration products, including NT1-SSMI, CBT-SSMI, and NOAA-CDR in the Arctic, and NOAA-CDR (but only
535 slightly) in the Antarctic. This Gaussian-fit methodology also confirms that group I products are slightly low biased in the

536 Arctic (see Lavergne et al., 2019). The worst of these biases we find in the Arctic for SICCI-12km that was not openly
537 distributed at the end of the ESA CCI Sea Ice Phase 2 project, partly based on these results.

538 Our results suggest that group I products, i.e., the three CDRs OSI-450, SICCI-25km and SICCI-50km, are more
539 accurate and have a higher precision than the fourth CDR investigated: NOAA-CDR; this applies to both hemispheres. Group
540 I products can be regarded as being superior, in terms of the precision and accuracy, to NT1-SSMI and CBT-SSMI as well.
541 Our results furthermore confirm earlier work (Andersen et al., 2007; Ivanova et al., 2015) that the accuracy of NT2-algorithm
542 products near 100% cannot be quantitatively assessed. We hypothesize that this is merely caused by the fact that sea-ice
543 concentrations are constrained to a maximum of 100% by the algorithm concept. This is fundamentally different to the other
544 nine products investigated. In the iterative retrieval used in the NT2 algorithm, sea-ice concentrations are only allowed to
545 converge at 100% from one side, i.e. $< 100\%$ - in contrast to other iterative algorithms such as, e.g., the polynya signature
546 simulation method (PSSM) (Markus and Burns, 1995). Another element of the NT2-algorithm, which complicates quantitative
547 assessment of the accuracy near 100% sea-ice concentration, is the usage of model atmospheres to create look-up tables from
548 which the sea-ice concentration is retrieved. This approach likely reduces the natural variability of the obtained sea-ice
549 concentration. We refer to Brucker et al. [2014] for more details of NT2-algorithm sea-ice concentration uncertainty.

550 The results of the comparison in Sect. 4.1 and 4.2 are based on winter data (see Sect. 2.2). But also during winter
551 temperatures can get close to 0°C influencing the brightness temperatures used to compute the sea-ice concentration and
552 questioning the assumption of freezing conditions for generation of the near-100% reference sea-ice concentration data set.
553 By using the co-located air temperature from ERA-Interim reanalysis data included in the RRDP2 data set (Pedersen et al.,
554 2019), we repeated the analyses for cold cases, i.e., air temperatures below -10°C . We find that biases between satellite and
555 reference sea-ice concentrations change by less than 0.1% for all products except NT1-SSMI. For NT1-SSMI constraining the
556 analyses to cold cases yields a bias reduction by $\sim 0.4\%$ for the Arctic and $\sim 0.7\%$ for the Antarctic, indicating that this particular
557 algorithm is more sensitive than others to variability of air temperature.

558 **5 Comparison with ship-based visual sea-ice cover observations**

559 In this section, we present the results of the evaluation of the ten products at intermediate sea-ice concentrations by means
560 of an inter-comparison to the visual ship-based observations described in Sect. 2.3. Upfront we note the limitations of the
561 manual ship-based visual sea-ice cover observations used here. They were collected by a myriad of different observers with
562 different levels of experience for this task. For an untrained observer it is relatively straightforward to estimate the total sea-
563 ice concentration for closed ice conditions, i.e. $> 80\%$, or very open ice conditions, i.e. $< 30\%$. It is more difficult, however,
564 to estimate whether sea ice covers, e.g. 40% or 50% of the 1 km radius area around the ship. Therefore, we can expect a
565 reduced accuracy for ship-observations of the intermediate sea-ice concentration range from $\sim 30\%$ to $\sim 80\%$. At the same time,
566 this is possibly the sea-ice concentration range where the different spatial scales of the two kind of observations compared here
567 have the largest impact on the results. Note that Worby and Comiso [2004] reported an uncertainty estimate between 5% and
568 10% for the total sea-ice concentration based on observations of the same scene by different observers. To the best knowledge
569 of the authors, papers about a better quantification of the accuracy of these observations have not yet been published.

570 In addition, ship observations were collected under different weather and daylight conditions as well as during ship
571 transits with different speeds. The first two points influence the visibility and change the visual appearance of sea ice and
572 openings, and can result in a larger spread of an observed sea-ice concentration value around the actual value. Different weather
573 conditions also have an influence on the size of the area actually observed around the ship that is difficult to quantify. This
574 observation area is supposed to be of one kilometre radius but it can be assumed that it is smaller in case of poor visibility than
575 it is in case of clear-sky, good visibility conditions; visibility is not regularly reported along with the ice observations. A single
576 observation of the sea-ice conditions takes a certain amount of time, which is a function of the experience of the observer. The

577 observation area around the ship's track represented by a single observation is hence a function of the ships' speed and of the
578 experience of the observer. As long as ice conditions do not change for a few kilometres this does not matter but in highly
579 heterogeneous ice conditions, this can be important. Therefore, there is a variable representativity of the observed sea-ice
580 conditions around and along the ship's track (see Sect. 6.1.4).

581 Ships often tend to avoid thick and deformed sea ice and ship-track forecasts are often optimized accordingly (e.g.
582 [Pizzolato et al., 2016; Kuuliala et al., 2017]). Ship-based observations therefore often represent the thinner ice categories
583 and/or conditions encountered in leads or openings. These are frozen over with sea ice in winter but are open water in summer.
584 Therefore, particularly during summer or episodes of warmer weather, the sea-ice concentrations from the small scale ship-
585 based observations are likely lower than from the larger scale satellite microwave radiometry. According to Ivanova et al.
586 (2015), microwave radiometry tends to underestimate sea-ice concentrations over very thin (< 15 cm) ice. This suggests that
587 during winter, sea-ice concentrations from ship-based observations could be, contrary to summer, slightly higher than from
588 satellite microwave radiometry.

589 A systematic quantification of the uncertainty in ship-based sea-ice observations has not been carried out yet and is
590 beyond our scope. Even though we do not use single ship-based observations but follow Beitsch et al. (2015) and average over
591 all ship-based sea-ice observations along the ship's track of one day, discarding days with less than three observations (see
592 Sect. 2.3), we cannot rule out that some of these daily average observations are biased because of the reasons discussed above.

593 **5.1 Arctic**

594 Overall, for the Arctic, all ten products compare reasonably well to the ship-based observations (Fig. 15, Table 5). At high
595 concentrations (> ~80%), group I products and ASI-SSMI (Fig. 15 d to h) exhibit the most symmetric distributions around the
596 identity line. NT1-SSMI (Fig. 15 a) shows an asymmetric distribution with a considerable fraction of satellite sea-ice
597 concentrations even below 60%, translating into a difference between satellite- and ship-based sea-ice concentrations of up to
598 40%. Group II and IV products (Fig. 15 b, c, i, j) show an asymmetric distribution with more high satellite-based than high
599 ship-based sea-ice concentrations. Most products have more data pairs below the identity line, i.e., satellite-based are smaller
600 than ship-based sea-ice concentrations as illustrated by a negative overall bias and regression lines located below the identity
601 line for six of the ten products (Table 5). Group II and IV products provide the smallest absolute overall bias of < 1%. Biases
602 are larger for group I but at the same time the standard deviation of the difference (SDEV) is smallest for group I products.
603 Highest correlations are obtained for CBT-AMSRE and group I products, except OSI-450. Best linear fits, i.e. slopes closest
604 to the identity line, we find, however, for CBT-SSMI and NOAA-CDR.

605 How do results of the entire year compare to those obtained separately for winter or summer months (see Fig. 5 for
606 differences in the location of the ship-based observations)? For all products, the correlation is lower in winter than summer
607 and the entire year. This can be explained by fewer observations during winter in general and by substantially fewer low sea-
608 ice concentrations; most observations during winter are > 75%. Except for ASI-SSMI, the differences in the bias between
609 summer, winter and the entire year are small (mostly < 1%). All products except SICCI-12km and ASI-SSMI provide a lower
610 SDEV in winter than summer. Compared to the other groups, group I exhibits smallest SDEV in summer while group II does
611 so in winter. OSI-450 stands out with the smallest inter-seasonal change in bias and SDEV of only ~0.1% (Table 5, see also
612 Sect. 6.1.4).

613 We want to better visualize the average distribution of the two data sets and investigate the partitioning of the data into
614 sea-ice concentration bins of 10% width - the average accuracy of the ship-based sea-ice concentration observations. For this
615 purpose we bin sea-ice concentrations of one data set, e.g. ship observations, into 10% wide bins and compute the mean sea-
616 ice concentration of the other data set (Fig. 15, red symbols) and vice versa (blue symbols). The binned values and associated
617 regression lines illustrate even better the above-mentioned asymmetry in the distribution of the data pairs. For instance, NT1-
618 SSMI (Fig. 15 a) sea-ice concentrations range between 60% and 100% over a ship-based observations range of 80 to 100%.

619 Consequently, the average NT1-SSMI sea-ice concentration for ship-observation bin 95%-100% is ~85% (uppermost red
620 triangle), while the average ship-based sea-ice concentration for NT1-SSMI bin 95%-100% is ~95% (uppermost blue square).
621 For two equally well-distributed data sets, one would expect that red and blue symbols and regression lines are close to each
622 other. This is not the case and we refer to Sect. 6.1.4 for more discussion of this issue.

623 5.2 Antarctic

624 The scatterplots of the daily average along-track mean sea-ice concentrations for the Antarctic (Fig. 16) reveal, for sea-
625 ice concentrations $> \sim 80\%$, mostly symmetric distributions for seven of the ten products. NT1-SSMI has considerably more
626 low than high sea-ice concentration values (compare Fig. 15 a). NT2-AMSRE has almost no data points below the identity
627 line at $> \sim 80\%$ with data pairs concentrated at 100% satellite sea-ice concentration. A considerable drop in the count of data
628 values above the identity line at lower concentrations results in a highly asymmetric distribution at concentrations below $\sim 50\%$.
629 .Group II products have the lowest overall biases (Table 6) but slopes of the linear regression are considerably steeper than the
630 identity line (Fig. 16 b, c, j) in contrast to group III products (Fig. 16 a, d) and OSI-450 (Fig. 16 e). We find the largest bias
631 for NT1-SSMI: -11.0%. NT2-AMSRE shows a considerable positive bias: +4.5%. Highest correlations between ship- and
632 satellite-based observations we find for group I products and CBT-AMSRE, lowest SDEV values for group I products as well.

633 In contrast to the Arctic, correlations between ship- and satellite-based sea-ice concentrations are smaller in summer
634 than winter when correlations are even higher than for the entire year for most products (Table 6) For all products, except
635 group III, biases are smaller in winter than summer by mostly $< 2\%$. NT2-AMRE is the only product with a positive bias in
636 both seasons. The largest inter-seasonal bias change we find for CBT-AMSRE: 2.7% and SICCI-25km: 2.9%, respectively,
637 the smallest for CBT-SSMI: 0.4%. For all products, the SDEV is smaller by 3-5% (larger by 2-3%) compared to the entire
638 year in winter (in summer). Inter-seasonal SDEV changes are hence considerably larger in the Antarctic than in the Arctic
639 (compare Tables 5 and 6).

640 Binning the sea-ice concentrations like described in Sect. 5.1 (red and blue symbols in Fig. 16) leads to the following
641 main observations. 1) All products – except NT1-SSMI and NT2-AMSRE – have the best agreement with ship-based
642 observations in the sea-ice concentration range 60% to 80%. 2) All products under-estimate ship-based sea-ice concentrations
643 for concentrations $< \sim 50\%$. 3) The negative bias of $\sim 10\%$ observed for NT1-SSMI applies to the entire sea-ice concentration
644 range. 4) NT2-AMSRE is the only product over-estimating ship-based sea-ice concentrations considerably; this over-
645 estimation reaches 10% for the range 60% to 80%. Apart from that we find, like for the Arctic, that blue regression lines exhibit
646 a considerably steeper slope than the red ones, suggesting that also in the Antarctic the distribution of the data over the range
647 0 to 100% is asymmetric (see Sect. 6.1.4).

648 5.3 Summary and discussion of comparison against ship-based observations

649 Group II products provide the smallest overall difference to the ship-based observations: around 0% (Arctic) and around
650 -2% (Antarctic). Group I products provide differences around -7.5% (Arctic) and -3.5% (Antarctic). Group I and II products
651 share similar average correlations (R^2) in the Arctic: 0.77 and 0.76, and in the Antarctic: 0.74 and 0.72, respectively. Standard
652 deviations for group I products are smaller – by 1% in the Arctic and by 2% in the Antarctic – than for group II products. On
653 average, these results are better than those obtained for group III and IV products. We refrain from giving a ranking or
654 recommendation as to which product is the best when compared to ship-based observations.

655 We find the lowest correlation for ASI-SSMI (group III) in both hemispheres. However, it is the product with the largest
656 improvement in the inter-comparison results between winter and summer in the Arctic: correlation increases, difference and
657 standard deviation decrease. This could be attributed to the higher fraction of intermediate sea-ice concentrations during
658 summer for which the comparably fine grid resolution of 12.5 km of the ASI-SSMI product could be of advantage. There is
659 evidence that this behaviour is coupled to the usage of the near 90 GHz channels because we observed a similar, albeit less

660 pronounced behaviour for SICCI-12km, which also employs near 90 GHz data and is provided at 12.5 km grid resolution
661 (Table 2). Because in the Antarctic the sea-ice cover is more open year round it is plausible that we did not find similar
662 behaviour there for these two products.

663 There are a few points to discuss. First of all, Fig. 15 and Fig. 16 reveal quite a high relative occurrence of sea-ice
664 concentrations in the range 95% to 100%. This might have biased our results. We therefore repeated our comparison by
665 discarding all data pairs with daily mean sea-ice concentrations $\geq 95\%$. Main results (not shown) are an overall increase in the
666 differences by about 2% and of the standard deviations by $\sim 1\%$ (group I) and 2-3% (group II) and a reduction of the correlation
667 by ~ 0.08 . Ranking between groups are otherwise not changed. Secondly, the results obtained for group I products are based
668 on the truncated sea-ice concentrations. Using the non-truncated data does not considerably change our findings (not shown).
669 Differences between satellite-based and ship-based sea-ice concentrations decrease by between 0 and 0.3% with a concomitant
670 increase in the standard deviation of up to 0.2%; this applies primarily to winter when the fraction of near $\sim 100\%$ sea-ice
671 concentrations is large. Thirdly, application of the Gaussian-fit method (see Sect. 2.1.4 and Sect. 6.1.3) seems not appropriate
672 given the sea-ice concentration range considered in this inter-comparison. Even if it would be, we can assume that differences
673 and standard deviations obtained for the other products would change by less than one percent. All these changes would be
674 small compared to the accuracy of the ship-based sea-ice concentrations (see Sect. 2.3, beginning of Sect. 5, and Sect. 6.1.4).

675 In Fig. 15 and Fig. 16, several of the products exhibit points along the $y=0\%$ sea-ice concentration line, i.e. conditions
676 where sea ice is reported by the ship while the satellite estimates exactly 0%. This applies most to group II products and ASI-
677 SSMI (Fig 15 b, c, d and j). These points with zero daily mean sea-ice concentration are very likely the result of the weather
678 filters applied (Sect. 2.1.3), which besides removing false sea ice caused by atmospheric effects, also removes true sea ice
679 (Ivanova et al., 2015; Lavergne et al., 2019). The combination of explicit atmospheric correction and dynamic tuning of the
680 weather filter in group I products seems to reduce the occurrence of such cases notable (Fig. 15 e, f and g); Fig. 16 e, f, and g).

681 **6 Discussion and Conclusions**

682 **6.1 Discussion**

683 **6.1.1 Observed differences in sea-ice area and extent**

684 Time series of SIA and SIE have long been used to derive conclusions about the past development of the sea-ice cover
685 and to even extrapolate its future development. In order to do so, such time series need to be sufficiently long, consistent and
686 accurate. A long, consistent time-series is typically obtained using a fundamental climate data record of brightness
687 temperatures as input for the retrieval, as is done, e.g., for group I product OSI-450, to ensure that inter-sensor differences are
688 as small as possible. Our paper suggests that additional steps might be required, for instance, dynamic retrieval of tie points
689 and dynamic adaptation of weather filters (see Sect. 6.1.2) to reach the goal of a long-term consistent sea-ice concentration
690 data set to be used to compute long-term consistent time-series of SIA and SIE.

691 Meier and Steward (2019) suggested a method to obtain an estimate of SIE and NSIDC sea-ice index accuracy, which
692 they found to be $\sim 50\,000\text{ km}^2$ under certain circumstances for the Arctic. They also pointed out, however, that there is a clear
693 bias (or spread) of $500\,000\text{ km}^2$ to 1 million km^2 between SIE estimates from different products (see also Ivanova et al., 2014).
694 Therefore, as long as one does not know which product provides the best representation of the actual sea-ice cover, one is left
695 alone with a relatively precise estimate of the SIE, which might be biased, however, by an amount an order of magnitude
696 larger. Notz (2014) found that the SIE and its trend provide a limited metric for the performance of numerical models. Petty
697 et al. (2018) suggested that predictions of the September Arctic sea-ice minimum in area and extent would benefit from giving
698 more weight to SIA. Niederdrenk and Notz (2018) concluded that observational uncertainty is the main source of uncertainty
699 for estimating at which level of global warming the Arctic will lose its summer sea-ice cover. In the light of these findings, the
700 inter-product differences in SIA and SIE resulting from our study provide useful information about which algorithm or group

701 of algorithms is particularly well suited for investigations of SIA and SIE in both or just one hemispheres year-round or just a
702 season. Because we were able to estimate the effect of the mismatch between true and grid resolution and of the pole-hole
703 interpolation in the Arctic and could further rule out influences of different land-masks, we are confident that the inter-product
704 differences observed are mostly originating from differences in how the algorithms handle surface emissivity variations or
705 variations in the atmospheric influence.

706 **6.1.2 The role of weather filters**

707 We illustrated that the weather filters implemented in each of the ten products, have quite different behaviour, despite the fact
708 they all use the same gradient ratios of brightness temperature frequency channels. Potential users should be aware of this. We
709 confirm that the dynamic open water filters designed for group I products (see Table 2) have a stable impact on the lower part
710 of the sea-ice concentration, cutting through it at about 10% SIC. This is across the months, changes of sensors (and thus
711 frequencies, calibration, etc...) and for both hemispheres. We refer to Lavergne et al. (2019) for discussions how this
712 consistency could be improved further. The analysis sheds also light on how the other six products perform in terms of stability.
713 Here, we are interested both in the temporal consistency of the weather filter effects (e.g. jumps across satellite series, across
714 months, across climate-induced trends) and the absolute level at which they cut through the sea-ice concentration distribution,
715 especially with respect to the 15% threshold embedded in the SIE and SIA curves shown in Fig. 6 and Fig. 7. Group II products
716 all cut around 15%, sometimes below, sometimes above, but in general at higher sea-ice concentration than group I products
717 (Fig. 1 and Fig. 2). We observed evidence for an actual impact of weather-filter cut-off sea-ice concentrations variation over
718 time in the SIE time series in winter in the Antarctic (Fig. 7 c). We believe Fig. 1 and Fig. 2 are a new illustration that data
719 products can differ in many ways. Such time-series, adapted from Lavergne et al. (2019), are an effective tool for data-
720 producers and users to assess the temporal consistency of sea-ice concentration data products.

721 **6.1.3 The impact of truncating sea-ice concentrations at 100%**

722 The analysis of the validation at 100% conditions (Sect. 4) raises a critical question to be discussed in the future among data
723 producers and with the data users: “Given that sea-ice concentration products are used following application of a 100% sea-
724 ice concentration threshold, is it better to have algorithms (slightly) overshoot the sea-ice concentration distribution or should
725 data producers aim at an unbiased non-truncated distribution?” To let algorithms overshoot will return fewer below-100%
726 estimates in the product files which might be positive for large areas of the inner sea-ice cover during winter. However, one
727 consequence of this (hypothetically) shifted sea-ice concentration distribution with modal values $> 100\%$ is that the sea-ice
728 concentration in areas with a true $< 100\%$ sea ice concentration, i.e. 99% or even 98%, might be set to 100%. Such areas could
729 contain leads. According to, e.g., Marcq and Weiss (2012) about 70% of the upward ocean-atmosphere heat exchange occurs
730 through leads even though these cover only one to two percent of the Central Arctic ocean. Assuming a heat transfer through
731 thick ice of 5 W/m^2 and through a lead of 400 W/m^2 (e.g. Marcq and Weiss, 2012, near-surface air-temperature difference of
732 30K) a heat flux calculation yields 5 W/m^2 for 100% and $\sim 9 \text{ W/m}^2$ for 99% true sea-ice concentration, an increase by 80%.
733 Using sea-ice concentrations of an algorithm with a modal value at 101% or higher might therefore result in a substantial
734 under-estimation of the surface heat flux. Integrated over the sea-ice covered central Arctic Ocean (area ~ 7 million km^2) this
735 under-estimation could be as high as $2.4 * 10^{12}$ MJ per day.

736 To aim at an unbiased algorithm might help with a better sensitivity to small openings (but note the RMSE of non-
737 truncated retrievals ranges between 2% and 5% for the various algorithms studied here), however, the product grid will have
738 more below-100% estimates. Our analysis, supported by the Gaussian-fit method, helps introducing the question and opening
739 the discussion, but does not bring the answer.

740 6.1.4 Observed differences to ship-based observations

741 One of the innovations of the products of group I (see Table 2) is the self-optimizing capability of the algorithms to
742 adopt to seasonally changing sea-ice conditions, i.e. the transition between winter and summer. If ship-based sea-ice
743 observations are as reliable in summer as they are in winter then a comparison of the differences and standard deviations
744 obtained in winter and summer could reveal how well an algorithm deals with the seasonally changing sea-ice conditions. If
745 we focus on sea-ice concentrations $< 95\%$ to avoid the clustering of data pairs near 100% during winter, we find the CDRs of
746 the group I products: SICCI-25km, SICCI-50km and OSI-450 to stand out with winter-to-summer changes in the difference
747 between ship- and satellite-based sea-ice concentrations around 0.2% , compared to $\sim 3.5\%$ for group II products in the Arctic.
748 Respective changes in the standard deviation range between 0.1% and 5.2% for group II but are near $\sim 2.7\%$ for group I CDRs.
749 In the Antarctic, smallest winter-to-summer differences of $\sim 0.3\%$ are obtained for group I products SICCI-12km and OSI-450,
750 compared to $\sim 2.5\%$ for group II products. These provide a winter-to-summer change in standard deviation of $\sim 6.5\%$ while
751 SICCI-12km and OSI-450 exhibit values of $\sim 4.0\%$. These results suggest that most group I products compare with the ship-
752 based sea-ice observations more consistently across seasons than the other products.

753 It is noteworthy, however, to keep in mind the difference in scales and observational limitations between ship-based
754 and satellite-based observations of the sea-ice concentration – as illustrated in Figure 17. For the pack ice / lead case (Fig. 17
755 a), Table 7 top), the variation of the ship-based observations depends strongly on the fraction of thin ice. For leads covered by
756 open water, satellite sea-ice concentrations tend to exceed ship-based concentrations. This could explain the banana-shaped
757 distribution of data pairs for the SH (Fig. 16). For leads covered by thin ice, it is the other way round. Ship-observations would
758 reveal a total sea-ice concentration of 100% , while most of the tested algorithms underestimate the true sea-ice concentration
759 in presence of thin ice [Cavaliere, 1994; Comiso and Steffen, 2001; Ivanova et al., 2015]. This could explain the larger fraction
760 of ship-based sea-ice concentrations near 100% as is particularly pronounced, e.g., for NT1-SSMI (Fig. 15 a) and, in general,
761 the larger range of satellite versus ship-based sea-ice concentrations at comparably high concentrations. For the marginal ice
762 zone (Fig. 17 b), Table 7 bottom), it is more likely that most products provide smaller sea-ice concentrations than observed
763 from a ship. During winter, a considerable fraction of the sea ice might be thin ice – causing under-estimation as stated above.
764 During summer, a considerable fraction of the sea ice might be too wet to be recognized as ice by satellite microwave
765 radiometry – causing an under-estimation as well (see e.g. Worby and Comiso, 2004; Ozsoy-Cicek et al., 2009). In addition,
766 low ice concentrations are often filtered by the weather filters applied (see Sect. 2.1.3). This results in a substantially larger
767 range of ship-based sea-ice concentrations at comparably low satellite sea-ice concentrations. Furthermore, this also results in
768 a larger fraction of low ice-concentration values for the satellite-based than the ship-based observations. This explains why
769 mean ship-based sea-ice concentrations per binned satellite sea-ice concentration (blue symbols in Fig. 15 and 16) are shifted
770 so much to the right compared to the red symbols; this is also evident from the larger fraction of data pairs below than above
771 the identity line for sea-ice concentrations below 60 to 80% .

772 We note that another data set of a different kind of ship-based observations of Arctic sea-ice conditions is available at
773 the Arctic and Antarctic Research Institute (AARI) and has recently been used for a sea-ice concentration algorithm inter-
774 comparison study using an approach different to the one used in our paper (Alekseeva et al., 2019). We might extend our inter-
775 comparison studies to their data set in the future.

776 6.2 Conclusions

777 Recently, three new global sea-ice concentration (SIC) climate data records (CDRs) have been released. They are
778 described in Lavergne et al. (2019). These products: SICCI-25km, SICCI-50km, and OSI-450, utilize a dynamic, self-
779 optimizing hybrid sea-ice concentration algorithm, which is applied to satellite microwave brightness temperature
780 measurements of the SMMR, SSM/I and SSMIS instruments (OSI-450) or the AMSR-E and AMSR2 instruments (SICCI-
781 25km and SICCI-50km); see Table 1 for instruments and frequencies. Within this paper, these new products are evaluated by

782 means of an inter-comparison to seven other sea-ice concentration products (see Table 2 for acronyms, satellite sensors and
783 frequencies used, and assignment to groups of algorithms I to IV) and with independent sea-ice cover data.

784 We find a very good and consistent agreement in inter-annual variation of the monthly mean SIA and SIE time series for
785 both hemispheres for the overlap periods of the respective products used at their native grid resolution. We can explain
786 unexpected low SIE of SICCI-50km during Arctic winter and Antarctic summer by the coarse resolution of the 6.9 GHz
787 frequency observations combined with too aggressive filtering of near-coastal and potentially weather-influenced grid cells
788 near coastlines, which removed a substantial number of grid cells with ice. We note that SIE differences are to be expected
789 simply from the different grid resolutions. When inter-comparing products on the same grid (50km) applying the same land
790 mask, the products of group I provide quite similar values of SIC, SIA and SIE for both hemispheres, during times of both
791 maximum and minimum sea-ice cover. Overall differences are $< 1.0\%$ for SIC, and $< 100\,000\text{ km}^2$ (Arctic) and $< 200\,000\text{ km}^2$
792 (Antarctic) for both SIA and SIE. A similarly good agreement we find for the products CBT-SSMI and NOAA-CDR of group
793 II (see Table 2), as can be expected from the design of NOAA-CDR (see Appendix F and Peng et al., 2013). Largest differences
794 we find for NT1-SSMI and NT2-AMSRE: NT1-SSMI provides less sea ice than other products particularly in summer in the
795 Arctic and winter in the Antarctic; NT2-AMSRE provides more sea ice than other products in the Antarctic. Based on our
796 results we state that it matters which algorithm and/or product is used for monitoring the polar sea-ice cover as long as one is
797 interested in absolute values; similarity of trends has been shown elsewhere (e.g. Ivanova et al., 2014; Comiso et al., 2017a).

798 Results quoted in the previous paragraph rely on computations applying the often-used 15% SIC threshold (e.g. Gloersen
799 et al., 1992). Our investigations suggest that it might be worth to start reconsidering this threshold because, as illustrated in
800 our paper, the weather filters applied might have inter-sensor jumps and fail to cut at a constant sea-ice concentration across
801 the different satellite sensors used. For example, sea-ice concentrations from sensor A might be cut by the weather filter at
802 14% while these might be cut at 17% for a subsequent sensor B. As a result, more sea ice is removed for sensor B and the sea-
803 ice extent computed from sea-ice concentration data of sensor B is systematically smaller than the one computed from sensor
804 A. We observe evidence for this in our results. A more thorough analysis of trends over sub-periods observed by the ten
805 products could reveal other such impacts. This impact is likely to be particularly pronounced in the peripheral seas with a
806 comparably large fraction of the marginal ice zone, such as the Bering Sea or the Barents Sea. Note that the OSI-450 algorithm
807 provides a particularly stable weather-filter induced sea-ice concentration cut-off at $\sim 10\%$ across the sensors used.

808 Sea-ice concentrations are retrieved from satellite microwave brightness temperatures using an empirical physical
809 algorithm, which usually involves a limited set of tie points. However, the natural variability of surface properties relevant for
810 microwave remote sensing of sea ice is large and these tie points can only be an average signature associated with a significant
811 spread. As a result, sea-ice concentrations retrieved by such algorithms naturally vary *around* (below and above) 100% even
812 though the actual sea-ice concentration is exactly 100%. The natural variability of the sea-ice surface properties and the
813 linearity of most algorithms suggest that the spread of retrieved sea-ice concentrations around 100% follows a Gaussian
814 distribution. However, all ten products examined here either truncate sea-ice concentrations at 100%, i.e. fold retrieved values
815 $> 100\%$ to exactly 100%, or do simply not allow retrieval of SIC $> 100\%$ (NT2-AMSRE) and the natural variability cannot be
816 assessed. We develop a Gaussian-fit method to re-construct the full distribution of sea-ice concentrations around 100%. We
817 demonstrate its performance on the group I products (as these provide, for the first time, both the non-truncated and truncated
818 values) and subsequently use it to re-construct non-truncated sea-ice concentration distributions for the remaining six products
819 (Fig. 3, and Appendix H, Fig. H1 and Fig. H2). Based on our results we conclude that it is worthwhile to re-think the concept
820 of truncation at 100% (but also at 0% SIC) and critically re-assess evaluation results at the two ends of the sea-ice concentration
821 distribution. Indeed, we argue that direct evaluation of truncated data sets give a misleading information on the accuracy of
822 the sea-ice concentration data, and favour those data sets that overestimate sea-ice concentrations. Such overestimation has
823 direct implications on the ability of a given sea-ice concentration data set to e.g. observe small openings in an otherwise

824 complete sea-ice cover. We invite the sea-ice concentration data producers and users to take this into consideration and discuss
825 the implications towards future versions of such data products.

826 *Data availability.* All sea-ice concentration products except SICCI-12km are publicly available from the sources provided in
827 the respective sections of the supplementary material and the reference list. The SICCI-12km product is available upon request
828 from T. Lavergne. The standardized ship-based observations are available from the Integrated Climate Data Center (ICDC):
829 <http://icdc.cen.uni-hamburg.de/1/daten/cryosphere/seaiceparameter-shipobs/>. The RRDP2 data set is publicly available from
830 the source specified in the reference list.

831 7 Appendices

832 7.1 Appendix A. The EUMETSAT-OSISAF – ESA-CCI algorithm suite

833 The four products OSI-450, SICCI-12km, SICCI-25km, and SICCI-50km have in common that they are based on a
834 hybrid, self-tuning, self-optimizing sea-ice concentration algorithm (Lavergne et al., 2019). This algorithm is applied to
835 brightness temperature (TB) observations of the SMMR, SSM/I and SSMIS instruments for OSI-450, providing a fully revised
836 version of the OSI-409 CDR (Tonboe et al., 2016). This algorithm is applied to brightness temperatures measured by the
837 AMSR-E and AMSR2 instruments for the SICCI CDRs. Apart from the input satellite data the processing chains are the same
838 for these four products. The algorithm is a generalization of the Comiso Bootstrap frequency mode algorithm (App. C) and of
839 the Bristol algorithm (Smith and Barret, 1994; Smith 1996) and is described in detail in Lavergne et al. (2019). Two algorithms
840 that each combine three frequency channels (e.g. ~19 GHz at vertical polarization, and ~37 GHz at both horizontal and vertical
841 polarizations) are respectively optimized to provide best accuracy in Open Water (the B_{OW} algorithm) and Consolidated Ice
842 (the B_{CI} algorithm) conditions. The sea-ice concentrations obtained with each of the two optimized algorithms are merged
843 linearly into a hybrid sea-ice concentration SIC_{hybrid} according to the general formula:

$$844 \left\{ \begin{array}{l} w_{OW} = 1; \text{ for } B_{OW} < 0.7 \\ w_{OW} = 0; \text{ for } B_{OW} > 0.9 \\ w_{OW} = 1 - \frac{B_{OW} - 0.7}{0.2}; \text{ for } B_{OW} \in [0.7; 0.9] \end{array} \right. ; \quad SIC_{hybrid} = w_{OW} \times B_{OW} + (1 - w_{OW}) \times B_{CI} , \quad (A1)$$

845 For sea-ice concentrations below 70%, SIC_{hybrid} relies completely on B_{OW} and for sea-ice concentrations above 90% it
846 relies entirely on B_{CI} . The B_{OW} and B_{CI} algorithms can be regarded as a generalized version of the Comiso Bootstrap and Bristol
847 algorithms, in that sense that they combine the three different brightness temperature channels used by the two algorithms in
848 a 3-dimensional TB space, and optimize their data projection plane for best accuracy.

849 In this paper, we use the sea-ice concentration CDR derived from SMMR, SSM/I and SSMIS data: OSI-450
850 [https://doi.org/10.15770/EUM_SAF_OSI_0008], and two sea-ice concentration CDRs derived from AMSR-E and
851 AMSR2 data: SICCI-25km [<https://doi.org/10.5285/f17f146a31b14dfd960cde0874236ee5>] and SICCI-50km
852 [<https://doi.org/10.5285/5f75fcb0c58740d99b07953797bc041e>]. While SICCI-25km is based on brightness temperatures
853 measured at ~19 GHz and ~37 GHz, similar to OSI-450, SICCI-50km is based on brightness temperatures measured at ~7
854 GHz and ~37 GHz. OSI-450 and SICCI-25km come at 25 km grid resolution while SICCI-50km has 50 km grid resolution. In
855 addition, we use a fourth product, SICCI-12km, which is provided at 12.5 km grid resolution and is based on brightness
856 temperatures measured by AMSR-E and AMSR2 at ~19 GHz and ~90 GHz. Here, we use a prototype of SICCI-12km, which
857 was produced during the ESA CCI Sea Ice project, but was not released publicly (partly based on the results presented in this
858 manuscript). All these data sets have daily temporal resolution and are provided on polar EASE grids version 2.0 (Brodzik et
859 al. 2012, 2014).

860 7.2 Appendix B. The ARTIST sea-ice (ASI) algorithm

861 The ARTIST Sea Ice (ASI) algorithm (Kaleschke et al., 2001; Spreen et al., 2008) is a modified hybrid of the Near 90
862 GHz algorithm (Svendsen et al., 1987) and the NASA Team algorithm (see App. D). Water and ice are distinguished at high
863 resolution by the TB polarization difference (P) at ~ 90 GHz:

$$864 \quad P = TB_{90}^V - TB_{90}^H$$

865 The basic equations for the ASI algorithm are based on the Near 90 GHz algorithm of Svendsen et al. (1987):

866

$$867 \quad P = a \times (C \times \Delta\epsilon_{ice} \times T_{ice} + (1 - C) \times \Delta\epsilon_{water} \times T_{water}) \quad (B1)$$

868 with the atmospheric influence $a = (1.1 \times e^{-\tau} - 0.11) \times e^{-\tau}$

869

870 C is the total sea-ice concentration, T is the temperature, $\Delta\epsilon$ is the difference in surface emissivity between vertical and
871 horizontal polarization for the ice or water surface fraction, and τ is the total atmospheric optical depth for Arctic conditions
872 at this frequency and viewing conditions. For ice free ($C = 0$) and totally ice covered ($C = 1$) conditions, Eq. (B1) yields the
873 tie-points for open water $P_{water} = a_{water} \times \Delta\epsilon_{water} \times T_{water}$ and sea ice $P_{ice} = a_{ice} \times \Delta\epsilon_{ice} \times T_{ice}$. Taylor expansions of Eq.
874 (B1) around $C=0$ and $C=1$ lead to a pair of equations for P , in which the atmospheric influences a_{water} and a_{ice} can be substituted
875 with the aid of the tie point equations – provided that the variation of the atmospheric influence is small over water or ice (see
876 Spreen et al., 2008). After substitution one obtains

$$877 \quad C = \left(\frac{P}{P_{water}} - 1 \right) \times \left(\frac{\Delta\epsilon_{water} \times T_{water}}{\Delta\epsilon_{ice} \times T_{ice} - \Delta\epsilon_{water} \times T_{water}} \right) \text{ for } C \rightarrow 0 \quad (B2)$$

$$878 \quad C = \frac{P}{P_{ice}} + \left(\frac{P}{P_{ice}} - 1 \right) \times \left(\frac{\Delta\epsilon_{water} \times T_{water}}{\Delta\epsilon_{ice} \times T_{ice} - \Delta\epsilon_{water} \times T_{water}} \right) \text{ for } C \rightarrow 1 \quad (B3)$$

879 According to Svendsen et al. (1987) the ratio of the surface emissivity differences can be set to a constant value (-1.14).
880 With this simplification and by assuming that the atmospheric influence inherent in P is a smooth function of the sea-ice
881 concentration one can use a third order polynomial function to interpolate between the solutions of Eq. (B2) and Eq. (B3) to
882 obtain sea-ice concentrations between 0 and 1 as a function of P :

$$883 \quad C(P) = d_3 \times P^3 + d_2 \times P^2 + d_1 \times P + d_0 \quad (B4)$$

884 The coefficients d_i are derived with a linear equation system based on Eq. (B2) and Eq. (B3) and their first derivatives
885 (Spreen et al., 2008).

886 The larger, compared to the lower frequencies used in most products (see Table 2), weather influence at ~ 90 GHz
887 frequencies by atmospheric water content and surface wind speed can cause substantial over-estimation of the sea-ice
888 concentration over open water and within the ice edge (Kern, 2004; Andersen et al., 2006). Over open water, the weather
889 influence is reduced by combining sea-ice concentrations obtained with Eq. (B4) with NASA-Team algorithm (NTA, see also
890 App. D) sea-ice concentrations following:

$$891 \quad C = C_{ASI} \text{ for } C_{NTA} > 5\%; C = 0\% \text{ for } C_{NTA} \leq 5\% \quad (B5)$$

892 Hence, the ASI algorithm is a hybrid of the near-90GHz algorithm (Eq. (B1) through Eq. (B4)) and the NTA (Kaleschke
893 et al., 2001; Ezraty et al., 2007; Girard-Ardhuin, personal communication, June 3 2019).

894 We use the ASI algorithm sea-ice concentration product provided via the Integrated Climate Data Center:
895 <https://icdc.cen.uni-hamburg.de> [last access date: 27/2/2019]. This product is processed at the French Institute for Exploitation
896 of the Sea (IFREMER) from SSM/I and SSMIS data, and provided via ICDC after application of a running 5-day-median
897 filter, further reducing spurious weather-influence induced sea-ice concentration in the open water (Kern et al., 2010), on a
898 polar-stereographic grid with 12.5 km grid resolution (at 70 degrees latitude). We abbreviate this data with ASI-SSMI.

7.3 Appendix C. The Comiso-Bootstrap algorithm

The Comiso-Bootstrap algorithm (Comiso, 1986; Comiso et al, 1997; Comiso et al., 2003; Comiso and Nishio, 2008) combines TB observations at either two different frequencies (frequency mode, 37 GHz and 19 GHz, vertical polarization) or at two different polarizations (polarization mode, 37GHz, vertical and horizontal polarization). It is based on the observation that brightness temperatures measured at these frequencies / polarizations over closed sea ice tend to cluster along a line (ice line) while those over open water tend to cluster around a single point in the respective two-dimensional brightness temperature space. The total sea-ice concentration is computed using

$$C = \frac{TB_f^V - TB_{f,OW}^V}{TB_{f,I}^P - TB_{f,OW}^V} \quad (C1)$$

with the brightness temperature measured at vertical polarization and frequency $f=37$ GHz (polarization mode) or $f=19$ GHz (frequency mode): TB_f^V , the open water tie point $TB_{f,OW}^V$ at vertical polarization and the same frequency as TB_f^V , and the intersection of the ice line with a line from the open water tie point through the observed brightness temperature: $TB_{f,I}^P = A \times \frac{B-W}{Q-A} + B$. Scalars A and B are functions of the ice tie points for first-year ice (FYI) and multiyear ice (MYI) at 37 GHz at vertical and horizontal polarization (polarization mode) or at 19 GHz and 37 GHz, both vertical polarization (frequency mode). Q and W are functions of the actually observed brightness temperature and the water tie point at the respective frequencies / polarizations. The two algorithms (frequency and polarization mode) are combined so that only the polarization mode is used in high concentration conditions and the frequency mode otherwise.

We use daily gridded sea-ice concentrations derived with the Comiso-Bootstrap (CBT) algorithm from SMMR, SSM/I and SSMIS instruments, as processed at NASA Goddard Space Flight Center (GSFC), and made available at <https://nsidc.org/data/nsidc-0079>. They are on a polar-stereographic grid with 25 km grid resolution (at 70 degrees latitude). We abbreviate this data with CBT-SSMI. For practical reasons, we access these GSFC CBT-SSMI fields from the NOAA sea-ice concentration CDR (App. F) files, where they are provided as additional data (Meier and Windnagel, 2018). The GSFC CBT-SSMI sea-ice concentration data set involves manual filtering, especially at the beginning of the record (SMMR period).

In addition, we use daily gridded sea-ice concentrations derived with this algorithm from AMSR-E data (Comiso et al., 2003; Comiso and Nishio, 2008) as provided by NSIDC (AE_SI25.003, Cavalieri et al., 2014, https://nsidc.org/data/ae_si25/versions/3, last access date: 26/4/2018) on the same polar-stereographic grid. The AMSR-E Comiso Bootstrap algorithm sea-ice concentration is referred with CBT-AMSRE throughout this paper. Note that the NSIDC product AE_SI25.003 does not contain CBT-AMSRE sea-ice concentrations itself. It contains the NT2 sea-ice concentration and the difference ‘‘Comiso Bootstrap minus NT2 sea-ice concentration’’. Therefore, we needed to compute the CBT-AMSRE sea-ice concentration by adding the NT2 sea-ice concentration to that difference.

7.4 Appendix D. The NASA-Team algorithm

The NASA-Team algorithm (Cavalieri et al., 1984, 1992, 1999) combines the large difference of the normalized brightness temperature polarization difference at 19 GHz, $PR = \frac{TB_{19}^V - TB_{19}^H}{TB_{19}^V + TB_{19}^H}$, between water and ice, with the observation, that the normalized brightness temperature frequency difference between 37 and 19 GHz at vertical polarization, $GR = \frac{TB_{37}^V - TB_{19}^V}{TB_{37}^V + TB_{19}^V}$, is negative for MYI and close to zero or slightly positive for FYI and open water. The total sea-ice concentration is derived as the sum of the fractions of MYI and FYI, which is constrained to a maximum of 1:

$$C_{FYI} = \frac{F_0 + F_1 \times PR + F_2 \times GR + F_3 \times PR \times GR}{D}, \quad C_{MYI} = \frac{M_0 + M_1 \times PR + M_2 \times GR + M_3 \times PR \times GR}{D},$$

with $D = D_0 + D_1 \times PR + D_2 \times GR + D_3 \times PR \times GR$ (D1)

Where coefficients F_i , M_i and D_i include the tie point information.

We use daily gridded sea-ice concentrations derived with the NASA-Team (NT1) algorithm from SMMR, SSM/I and SSMIS instruments, as processed at NASA Goddard Space Flight Center (GSFC), and made available at <https://nsidc.org/data/nsidc-0051>. They are on a polar-stereographic grid with 25 km grid resolution (at 70 degrees latitude). We abbreviate this data with NT1-SSMI. For practical reasons, we access these GSFC NT1-SSMI fields from the NOAA sea-ice concentration CDR (App. F) files, where they are provided as additional data (Meier et al., 2017; Meier and Windnagel, 2018). The GSFC NT1-SSMI sea-ice concentration data set involves manual filtering, especially at the beginning of the record (SMMR period).

7.5 Appendix E. The enhanced NASA-Team algorithm (NT2)

Inter-comparison studies such as those of Comiso and Steffen (2001) and Comiso et al. (1997) led to the development of the enhanced NASA-Team algorithm (NT2) (Markus and Cavalieri, 2000; Comiso et al., 2003) to mitigate effects such as layering in snow on sea ice on the accuracy of the sea-ice concentrations obtained with NT1. NT2 is conceptually different from the other algorithms presented here. The three relevant parameters (see below) are modelled as a function of sea-ice concentration in steps of 1% for 12 different atmospheric states using a radiative transfer model. The sea-ice concentration resulting in the minimum cost function between modelled and observed values of these parameters is taken as the retrieved total sea-ice concentration. The three parameters used are selected such that the influence of layering in snow on sea ice is mitigated:

$$\begin{aligned} \Delta GR &= \frac{TB_{90}^H - TB_{19}^H}{TB_{90}^H + TB_{19}^H} - \frac{TB_{90}^V - TB_{19}^V}{TB_{90}^V + TB_{19}^V} \\ PR_{19}^{rotated} &= -\frac{TB_{37}^V - TB_{19}^V}{TB_{37}^V + TB_{19}^V} \times \sin \theta_{19} + \frac{TB_{19}^V - TB_{19}^H}{TB_{19}^V + TB_{19}^H} \times \cos \theta_{19} \\ PR_{90}^{rotated} &= -\frac{TB_{37}^V - TB_{19}^V}{TB_{37}^V + TB_{19}^V} \times \sin \theta_{90} + \frac{TB_{90}^V - TB_{90}^H}{TB_{90}^V + TB_{90}^H} \times \cos \theta_{90} \end{aligned} \quad (E1)$$

The rotation is done in the space given by PR_{19} and GR (see App. D) or by PR_{90} and GR for $PR_{19}^{rotated}$ and $PR_{90}^{rotated}$, respectively, at an angle θ chosen such that the ice lines in the respective space are parallel to the GR axis.

We use daily gridded NT2 sea-ice concentrations derived from AMSR-E data as provided by NSIDC: AE_SI25.003 (Cavalieri et al., 2014, https://nsidc.org/data/ae_si25/versions/3, last access date: 26/4/2018) on polar-stereographic grid with 25 km grid resolution. We abbreviate this data with NT2-AMSRE.

7.6 Appendix F. The NOAA/NSIDC sea-ice concentration CDR

The NOAA/NSIDC sea-ice concentration CDR combines sea-ice concentrations computed with the NT1 algorithm (App. D) with those computed with the CBT algorithm (App. C), via

$$C = \max(C_{NT1}, C_{CBT}) \quad (F1)$$

within the ice edge. The ice edge is defined by the CBT sea-ice concentration of 10%. The generation and characteristics of the NOAA/NSIDC CDR as well as details about filters (see also Sect. 2.1) and about the statistical uncertainty estimate provided with the product are described in Peng et al. (2013) and Meier and Windnagel (2018). We use the daily gridded sea-ice concentration data of NOAA/NSIDC CDR version 3, named NOAA-CDR in this manuscript, provided by NSIDC on polar-stereographic grid with 25 km grid resolution (Meier et al., 2017, <https://nsidc.org/data/g02202/versions/3>, last access date: 7/2/2019).

It is important to note that the data sets NT1-SSMI (App. D) and CBT-SSMI (App. C), both from GSFC are not used as input in the NOAA/NSIDC CDR. Instead, sea-ice concentrations are computed at NSIDC using re-implementations of the two algorithms, which allow for a fully-automated and transparent processing as required for a CDR, and combined with Eq. (F1). One of the key difference between the NSIDC and GSFC versions is that the NSIDC ones do not involve manual editing, and start with SSM/I in July 1987.

978 7.7 Appendix G. Matrices of sea-ice concentration, area, and extent differences

979 This subsection contains the full set of matrices of differences between all ten products of the overall hemispheric average
980 monthly mean sea-ice concentration of the AMSR-E measurement period and of the respective overall monthly mean sea-ice
981 area (SIA) and extent (SIE) in Fig. G1 through Fig. G6.

982 7.8 Appendix H. Gaussian fits for all ten products

983 This subsection contains the two sets of Gaussian fits obtained for the Arctic (Fig. H1) and Antarctic (Fig. H2) based on
984 the methodology described in Sect. 2.1.4; see also Fig. 3). These are the fits obtained from data of the overlap period between
985 the AMSR-E measurement period and the RRDP2 near-100% reference sea-ice concentration data set period, that is in winter
986 of years 2007 through 2011.

987 *Author contributions.* SK led the writing in most sections, with contributions by TL (Sect. 2.1, Sect. 6.1, Sect. 6.2), LTP (Sect.
988 1, Sect. 2.1), DN (Sect. 1, Sect. 3), and RTT (Sect. 2.1, Sect. 6.2). AMS and TL contributed to concept and work of Sect. 2.1.3.
989 TL contributed to concept and work of Sect. 2.1.4. RS and LTP produced and provided the RRDP2 data set and consulted its
990 usage (Sect. 2.2). DN and TL contributed to concept and work as well as design of figures of Sect. 3. SK performed the data
991 analysis and inter-comparison with contributions in the interpretation of the results from all co-authors.

992 *Competing interests.* The authors declare that they have no conflict of interest.

993 *Acknowledgements.* The work presented here was funded by EUMETSAT (through the 2nd Continuous Developments and
994 Operation Phase of OSI SAF) and ESA (through the Climate Change Initiative Sea_Ice_cci project), and the German Research
995 Foundation (DFG) Excellence Initiative CLISAP under Grant EXC 177/2. The publication itself is funded by the Deutsche
996 Forschungsgemeinschaft (DFG, German Research Foundation) under Germany's Excellence Strategy – EXC 2037 'CLICCS
997 – Climate, Climatic Change, and Society' – Project Number: 390683824, contribution to the Center for Earth System Research
998 and Sustainability (CEN) of the University of Hamburg.

999 8 References

- 1000 Alekseeva, T. Tikhonov, V., Frolov, S., Repina, I., Raev, M., Sokolova, J., Sharkov, E., Afanasieva, E., and Serovetnikov, S.:
1001 Comparison of Arctic sea ice concentration from the NASA Team, ASI, and VASIA2 algorithms with summer and winter ship
1002 data, *Rem. Sens.*, 11(21), 2481, <https://doi.org/10.3390/rs11212481>, 2019.
- 1003 Andersen, S., Tonboe, R. T., Kern, S., and Schyberg, H.: Improved retrieval of sea ice total concentration from spaceborne
1004 passive microwave observations using Numerical Weather Prediction model fields: An intercomparison of nine algorithms,
1005 *Rem. Sens. Environ.*, 104(4), 374-392, 2006.
- 1006 Andersen, S., Pedersen, L. T., Heygster, G., Tonboe, R. T., and Kaleschke, L.: Intercomparison of passive microwave sea ice
1007 concentration retrievals over the high concentration Arctic sea ice, *J. Geophys. Res.*, 112, C08004,
1008 <https://doi.org/10.1029/2006JC003543>, 2007.
- 1009 Beitsch, A., Kern, S., and Kaleschke, L.: Comparison of SSM/I and AMSR-E sea ice concentrations with ASPeCt ship
1010 observations around Antarctica, *IEEE Trans. Geosci. Rem. Sens.*, 53(4), 1985-1996,
1011 <https://doi.org/10.1109/TGRS.2014.2351497>, 2015.
- 1012 Brodzik, M. J., Billingsley, B., Haran, T., Raup, B., and Savoie, M. H.: EASE-Grid 2.0: Incremental but Significant
1013 Improvements for Earth-Gridded Data Sets, *ISPRS Int. Geo-Inf.*, 1(1), 32-45, <https://doi.org/10.3390/ijgi1010032>,
1014 <http://www.mdpi.com/2220-9964/1/1/32>, 2012.

1015 Brodzik, M. J., Billingsley, B., Haran, T., Raup, B., and Savoie, M. H.: Correction: Brodzik, M. J. et al. EASE-Grid 2.0:
1016 Incremental but Significant Improvements for Earth-Gridded Data Sets, *ISPRS Int. Geo-Inf.*, 1(1), 32-45, 2012. *ISPRS Int.*
1017 *Geo-Inf.*, 3(3), 1154-1156, <https://doi.org/10.3390/ijgi3031154>, <http://www.mdpi.com/2220-9964/3/3/1154>, 2014.

1018 Brucker, L., Cavalieri, D. J., Markus, T., and Ivanoff, A.: NASA Team 2 sea ice concentration algorithm retrieval uncertainty,
1019 *IEEE Trans. Geosci. Rem. Sens.*, 52(11), 7336-7352, <https://doi.org/10.1109/TGRS.2014.2311376>, 2014.

1020 Cavalieri, D. J.: A microwave technique for mapping thin sea ice, *J. Geophys. Res.*, 99(C6), 12561-12572, 1994.

1021 Cavalieri D. J., Gloersen, P., and Campbell, W. J.: Determination of Sea Ice Parameters With the NIMBUS 7 SMMR, *J.*
1022 *Geophys. Res.*, 89(D4), 5355-5369, 1984.

1023 Cavalieri, D. J., Crawford, J., Drinkwater, M., Emery, W. J., Eppler, D. T., Farmer, L. D., Goodberlet, M., Jentz, R., Milman,
1024 A., Morris, C., Onstott, R., Schweiger, A., Shuchman, R., Steffen, K., Swift, C. T., Wackerman, C., and Weaver, R. L.: NASA
1025 sea ice validation program for the DMSP SSM/I: final report. NASA Technical Memorandum 104559. National Aeronautics
1026 and Space Administration, Washington, D.C., 126 pp., 1992.

1027 Cavalieri, D. J., St. Germain, K. M., and Swift, C. T.: Reduction of weather effects in the calculation of sea ice concentration
1028 with the DMSP SSM/I, *J. Glaciol.*, 41(139), 455-464, 1995.

1029 Cavalieri, D. J., Parkinson, C. L., Gloersen, P., Comiso, J. C., and Zwally, H. J.: Deriving long-term time series of sea ice
1030 cover from satellite passive-microwave multisensor data sets, *J. Geophys. Res.*, 104(C7), 15803–15814,
1031 <https://doi.org/10.1029/1999JC900081>, 1999.

1032 Cavalieri, D. J., Markus, T., Hall, D. K., Gasiewski, A. J., Klein, M., and Ivanoff, A.: Assessment of EOS Aqua AMSR-E
1033 Arctic sea ice concentrations using Landsat-7 and airborne microwave imagery, *IEEE Trans. Geosci. Rem. Sens.*, 44(11),
1034 3057-3069, <https://doi.org/10.1109/TGRS.2006.878445>, 2006.

1035 Cavalieri, D. J., Markus, T., Hall, D. K., Ivanoff, A., and Glick, E.: Assessment of AMSR-E Antarctic winter sea-ice
1036 concentrations using Aqua MODIS, *IEEE Trans. Geosci. Rem. Sens.*, 48(9), 3331-3340,
1037 <https://doi.org/10.1109/TGRS.2010.2046495>, 2010.

1038 Cavalieri, D. J., Markus, T., and Comiso, J. C.: AMSR-E/Aqua daily L3 25km brightness temperature and sea ice concentration
1039 polar grids, version 3, Boulder, Colorado USA, NASA National Snow and Ice Data Center Distributed Arctic Archive Center,
1040 https://doi.org/10.5067/AMSR-E/AE_SI25.003, [date accessed: 2018-04-26], 2014.

1041 Cho, K., Sasaki, N., Shimoda, H., Sakata, T., and Nishio, F.: Evaluation and improvement of SSM/I sea ice concentration
1042 algorithms for the Sea of Okhotsk, *J. Rem. Sens. of Japan*, 16(2), 47-58, 1996.

1043 Comiso J. C.: Characteristics of arctic winter sea ice from satellite multispectral microwave observations, *J. Geophys. Res.*,
1044 91(C1), 975-994, 1986.

1045 Comiso, J. C.: Large Decadal Decline of the Arctic Multiyear Ice Cover. *J. Climate*, 25(4), 1176–1193,
1046 <https://doi.org/10.1175/JCLI-D-11-00113.1>, 2012.

1047 Comiso, J. C., and Nishio, F.: Trends in the sea ice cover using enhanced and compatible AMSR-E, SSM/I, and SMMR data, *J.*
1048 *Geophys. Res.*, 113, C02S07, <https://doi.org/10.1029/2007JC004257>, 2008.

1049 Comiso, J. C., and Steffen, K.: Studies of Antarctic sea ice concentrations from satellite data and their applications, *J. Geophys.*
1050 *Res.*, 106(C12), 31361-31385, 2001.

1051 Comiso, J. C., Cavalieri, D. J., Parkinson, C. L., and Gloersen, P.: Passive microwave algorithms for sea ice concentration: A
1052 comparison of two techniques, *Rem. Sens. Environ.*, 60(3), 357-384, 1997.

1053 Comiso, J. C., Cavalieri, D. J., and Markus, T.: Sea ice concentration, ice temperature, and snow depth, using AMSR-E data,
1054 *IEEE Trans. Geosci. Rem. Sens.*, 41(2), 243-252, <https://doi.org/10.1109/TGRS.2002.808317>, 2003.

1055 Comiso, J. C., Meier, W. N., and Gersten, R. A.: Variability and trends in the Arctic Sea ice cover: Results from different
1056 techniques, *J. Geophys. Res. - Oceans*, 122(8), 6883–6900, <https://doi.org/10.1002/2017JC012768>, 2017a.

1057 Comiso, J. C., Gersten, R. A., Stock, L.V., Turner, J., Perez, G. J., and Cho, K.: Positive trends in the Antarctic sea ice cover
1058 and associated changes in surface temperature, *J. Climate*, 30(6), 2251–2267, <https://doi.org/10.1175/JCLI-D-16-0408.1>,
1059 2017b.

1060 Ezraty, R., Girard-Arduin, F., Piollé, J.-F., Kaleschke, L., and Heygster, G.: Arctic and Antarctic sea ice concentration and
1061 Arctic sea ice drift estimated from special sensor microwave data – Users’s Manual, Version 2.1, IFREMER, Brest, France,
1062 February 2007.

1063 Gloersen, P., Campbell, W., Cavalieri, D. J., Comiso, J. C., Parkinson, C. L., and Zwally, H. J.: Arctic and Antarctic sea ice,
1064 1978-1987: satellite passive-microwave observations and analysis, Scientific and technical information program, vol. NASA
1065 SP-511, National Aeronautics and Space Administration (NASA), Washington D.C., 1992.

1066 Ivanova, D. P., Gleckler, P. J., Taylor, K. E., Durack, P. J., and Marvel, K. D.: Moving beyond the total sea ice extent in
1067 gauging model biases. *J. Clim.*, 29(24), 8965-8987, <https://doi.org/10.1175/JCLI-D-16-0026.1>, 2017.

1068 Ivanova, N., Johannessen, O. M., Pedersen, R. T., and Tonboe, R. T.: Retrieval of Arctic sea ice parameters by satellite passive
1069 microwave sensors: A comparison of eleven sea ice concentration algorithms, *IEEE Trans. Geosci. Rem. Sens.*, 52(11), 7233-
1070 7246, <https://doi.org/10.1109/TGRS.2014.2310136>, 2014.

1071 Ivanova, N., Pedersen, L. T., Tonboe, R. T., Kern, S., Heygster, G., Lavergne, T., Sørensen, A., Saldo, R., Dybkjær, G.,
1072 Brucker, L., and Shokr, M.: Inter-comparison and evaluation of sea ice algorithms: towards further identification of challenges
1073 and optimal approach using passive microwave observations, *The Cryosphere*, 9, 1797-1817, <https://doi.org/10.5194/tc-9-1797-2015>, 2015.

1075 Kaleschke, L., Lüpkes, C., Vihma, T., Haarpaintner, J., Bochert, A., Hartmann, J., and Heygster, G.: SSM/I sea ice remote
1076 sensing for mesoscale ocean-atmosphere interaction analysis, *Can. J. Rem. Sens.*, 27(5), 526-537, 2001.

1077 Kern, S.: A new method for medium-resolution sea ice analysis using weather-influence corrected Special Sensor
1078 Microwave/Imager 85 GHz data, *Int. J. Rem. Sens.*, 25(21), 4555-4582, <https://doi.org/10.1080/01431160410001698898>,
1079 2004.

1080 Kern, S., Kaleschke, L., and Clausi, D. A.: A comparison of two 85-GHz SSM/I ice concentration algorithms with AVHRR
1081 and ERS-2 SAR imagery, *IEEE Trans. Geosci. Rem. Sens.*, 41(10), 2294-2306, <https://doi.org/10.1109/TGRS.2003.817181>,
1082 2003.

1083 Kern, S., Kaleschke, L., and Spreen, G.: Climatology of the Nordic (Irminger, Greenland, Barents, Kara and White/Pechora)
1084 Seas ice cover based on 85 GHz satellite microwave radiometry: 1992-2008, *Tellus*, 62A(4), 411-434,
1085 <https://doi.org/10.1111/j.1600-0870.2010.00457.x>, 2010.

1086 Kuuliala, L., Kujala, P., Suominen, M., and Montewka, J.: Estimating operability of ships in ridged ice fields, *Cold Reg. Sci.*
1087 *Technol.*, 135, 51-61, <https://doi.org/10.1016/j.coldregions.2016.12.003>, 2017.

1088 Kwok, R.: Sea ice concentration estimates from satellite passive microwave radiometry and openings from SAR ice motion,
1089 *Geophys. Res. Lett.*, 29(9), 1311, <https://doi.org/10.1029/2002GL014787>, 2002.

1090 Lavergne, T., Sørensen, A. M., Kern, S., Tonboe, R., Notz, D., Aaboe, S., Bell, L., Dybkjær, G., Eastwood, S., Gabarro, C.,
1091 Heygster, G., Killie M. A., Brandt Kreiner, M., Lavelle, J., Saldo, R., Sandven, S., and Pedersen, L. T.: Version 2 of the
1092 EUMETSAT OSI SAF and ESA CCI sea-ice concentration climate data records, *The Cryosphere*, 13(1), 49-78,
1093 <https://doi.org/10.5194/tc-13-49-2019>, 2019.

1094 Maass, N., and Kaleschke, L.: Improving passive microwave sea ice concentration algorithms for coastal areas: applications
1095 to the Baltic Sea, *Tellus*, 62A(4), 393-410, <https://doi.org/10.1111/j.1600-0870.2010.00452.x>, 2010.

1096 Markus, T., and Burns, B. A.: A method to estimate subpixel-scale coastal polynyas with satellite passive microwave data, *J.*
1097 *Geophys. Res.*, 100(C3), 4473-4487, 1995.

1098 Markus, T., and Cavalieri, D. J.: An enhancement of the NASA Team sea ice algorithm, *IEEE Trans. Geosci. Rem. Sens.*,
1099 38(3), 1387-1398, 2000.

1100 Markus, T., and Dokken, S. T.: Evaluation of late summer passive microwave Arctic sea ice retrievals, *IEEE Trans. Geosci.*
1101 *Rem. Sens.*, 40(2), 348-356, 2002.

1102 Meier, W. N.: Comparison of passive microwave ice concentration algorithm retrievals with AVHRR imagery in Arctic
1103 peripheral seas, *IEEE Trans. Geosci. Rem. Sens.*, 43(6), 1324-1337, <https://doi.org/10.1109/TGRS.2005.846151>, 2005.

1104 Meier, W. N., and Steward, J. S.: Assessing uncertainties in sea ice extent climate indicators, *Environ. Res. Lett.*, 14, 035005,
1105 <https://doi.org/10.1088/1748-9326/aaf52c>, 2019.

1106 Meier, W. N., and Windnagel, A.: Sea ice concentration – climate algorithm theoretical basis document, NOAA Climate Data
1107 Record Program CDRP-ATBD-0107 Rev. 7 (03/062018), available at [https://www.ncdc.noaa.gov/cdr/oceanic/sea-ice-](https://www.ncdc.noaa.gov/cdr/oceanic/sea-ice-concentration)
1108 [concentration](https://www.ncdc.noaa.gov/cdr/oceanic/sea-ice-concentration), 2018.

1109 Meier, W. N., Hovelsrud, G. K., van Oort, B. E. H., Key, J. R., Kovacs, K. M., Michel, C., Haas, C., Granskog, M. A., Gerland,
1110 S., Perovich, D. K., Makshtas, A., and Reist, J. D.: Arctic sea ice in transformation: A review of recent observed changes and
1111 impacts on biology and human activity, *Rev. Geophys.*, 51, 185-217, <https://doi.org/10.1002/2013RG000431>, 2014.

1112 Meier, W. N., Fetterer, F., Savoie, M., Mallory, S., Duerr, R., and Stroeve, J.: *NOAA/NSIDC Climate Data Record of Passive*
1113 *Microwave Sea Ice Concentration, Version 3*. Boulder, Colorado USA. NSIDC: National Snow and Ice Data
1114 Center, <https://doi.org/10.7265/N59P2ZTG>, [Date accessed: 2019-02-07], 2017.

1115 Melia, N., Haines, K., Hawkins, E., and Day, J. J.: Towards seasonal Arctic shipping route predictions, *Environ. Res. Lett.*,
1116 12(8), 084005, <https://doi.org/10.1088/1748-9326/aa7a60>, 2017.

1117 Niederrenk, A. L., and Notz, D.: Arctic sea ice in a 1.5°C warmer world, *Geophys. Res. Lett.*, 45(4), 1963-1971,
1118 <https://doi.org/10.1002/2017GL076159>, 2018.

1119 Notz, D.: Sea-ice extent and its trend provide limited metrics of model performance, *The Cryosphere*, 8(1), 229-243,
1120 <https://doi.org/10.5194/tc-8-229-2014>, 2014.

1121 Njoku, E. G., Rague, B., and Fleming, K.: The Nimbus-7 SMMR Pathfinder Brightness Temperature Data Set, Jet Propulsion
1122 Laboratory Publication, Pasadena, USA, 98-4, 1998.

1123 Ozsoy-Cicek, B., Xie, H., Ackley, S. F., Ye, K.: Antarctic summer sea ice concentration and extent: comparison of ODEN
1124 2006 ship observations, satellite passive microwave and NICsea ice charts, *The Cryosphere*, 3, 1-9, [https://doi.org/10.5194/tc-](https://doi.org/10.5194/tc-3-1-2009)
1125 [3-1-2009](https://doi.org/10.5194/tc-3-1-2009), 2009.

1126 Pedersen, L. T., Saldo, R., Ivanova, N., Kern, S., Heygster, G., Tonboe, R. T., Huntemann, M., Ozsoy, B., Girard-Ardhuin, F.,
1127 and Kaleschke, L.: Reference dataset for sea ice concentration, <https://doi.org/10.6084/m9.figshare.6626549.v6>,
1128 https://figshare.com/articles/Reference_dataset_for_sea_ice_concentration/6626549, 2019.

1129 Peng, G., Meier, W. N., Scott, D., and Savoie, M.: A long-term and reproducible passive microwave sea ice concentration data
1130 record for climate studies and monitoring, *Earth Syst. Sci. Data*, 5, 311-318, <https://doi.org/10.5194/essd-5-311-2013>, 2013.

1131 Petty, A. A., Stroeve, J. C., Holland, P. R., Boisvert, L. N., Bliss, A. C., Kimura, N., and Meier, W. N.: The Arctic sea ice
1132 cover of 2016: a year of record-low highs and higher-than-expected lows, *The Cryosphere*, 12(2), 433-452,
1133 <https://doi.org/10.5194/tc-12-433-2018>, 2018.

1134 Pizzolato, L., Howell, S. E. L., Dawson, J., Laliberte, F., and Copland, L.: The influence of declining sea ice on shipping
1135 activity in the Canasian Arctic, *Geophys. Res. Lett.*, 43(23), 12146-12154, <https://doi.org/10.1002/2016GL071489>, 2016.

1136 Reid, P., Stammerjohn, S., Massom, R., Scambos, T., and Lieser, J. L.: The record 2013 Southern Hemisphere sea-ice extent
1137 maximum, *Ann. Glaciol.*, 56(69), 99-106, <https://doi.org/10.3189/2015AoG69A892>, 2015.

1138 Schlosser, E., Haumann, F. A., and Raphael, M. N.: Atmospheric influences on the anomalous 2016 Antarctic sea ice decay,
1139 *The Cryosphere*, 12(3), 1103-1119, <https://doi.org/10.5194/tc-12-1103-2018>, 2018.

1140 Smith, D. M.: Extraction of winter total sea ice concentration in the Greenland and Barents Seas from SSM/I data, *Int. J. Rem.*
1141 *Sens.*, 17(13), 2625-2646, 1996.

1142 Smith D. M., and Barrett, E. C.: Satellite mapping and monitoring of sea ice, CB/RAE/9/2/4/2034/113/ARE, RSU, University
1143 of Bristol, Bristol, UK, 1994.

1144 Spreen, G., Kaleschke, L., and Heygster, G., Sea ice remote sensing using AMSR-E 89-GHz channels, *J. Geophys. Res.*, 113,
1145 C02S03, <https://doi.org/10.1029/2005JC003384>, 2008.

1146 Svendsen, E., Mätzler, C., and Grenfell, T. C.: A model for retrieving total sea ice concentration from a spaceborne dual-
1147 polarized passive microwave instrument operating near 90 GHz, *Int. J. Rem. Sens.*, 8(10), 1479-1487, 1987.

1148 Stuecker, M. F., Bitz, C. M., and Armour, K. C.: Conditions leading to the unprecedented low Antarctic sea ice extent during
1149 the 2016 austral spring season, *Geophys. Res. Lett.*, 44, 9008-9019, <https://doi.org/10.1002/2017GL074691>, 2017.

1150 Tonboe, R. T., Eastwood, S., Lavergne, T., Sørensen, A. M., Rathmann, N., Dybkjær, G., Pedersen, L. T., Høyer, J. L., and
1151 Kern, S.: The EUMETSAT sea ice concentration climate data record, *The Cryosphere*, 10, 2275-2290,
1152 <https://doi.org/10.5194/tc-10-2275-2016>, 2016.

1153 Turner, J., Hosking, J. S., Phillips, T., and Marshall, G. J.: Temporal and spatial evolution of the Antarctic sea ice prior to the
1154 September 2012 record maximum extent, *Geophys. Res. Lett.*, 40, 5894-5898, <https://doi.org/10.1002/2013GL058371>, 2013.

1155 Turner, J., Phillips, T., Marshall, G. J., Hosking, J. S., Pope, J. O., Bracegirdle, T. J., and Deb, P.: Unprecedented springtime
1156 retreat of Antarctic sea ice in 2016, *Geophys. Res. Lett.*, 44, 6868-6875, <https://doi.org/10.1002/2017GL073656>, 2017.

1157 Wayand, N. E., Bitz, C. M., and Blanchard-Wrigglesworth, E.: A year-round subseasonal-to-seasonal sea ice prediction portal,
1158 *Geophys. Res. Lett.*, 46(6), 3298-3307, <https://doi.org/10.1029/2018GL081565>, 2019.

1159 Wiebe, H., Heygster, G., and Markus, T.: Comparison of the ASI ice concentration algorithm with Landsat-7 ETM+ and SAR
1160 imagery, *IEEE Trans. Geosci. Rem. Sens.*, 47(9), 3008-3015, <https://doi.org/10.1109/TGRS.2009.2026367>, 2009.

1161 Worby, A. P., and Allison, I. A.: Ship-Based Technique for Observing Antarctic Sea Ice: Part I: Observational Techniques and
1162 Results, Research Report No. 14, Antarctic Cooperative Research Centre, Hobart, TAS, Australia, 1999.

1163 Worby, A. P., and Comiso, J. C.: Studies of the Antarctic sea ice edge and ice extent from satellite and ship observations, *Rem.*
1164 *Sens. Environ.*, 92, 98-111, 2004.

1165 Worby, A. P., and Dirita, V.: A technique for making ship-based observations of Antarctic sea-ice thickness and characteristics
1166 - Part II: User Operating Manual, Research Report No. 14, Antarctic Cooperative Research Centre, Hobart, TAS, Australia,
1167 1999.

1168 Worby, A. P., Geiger, C. A., Paget, M. J., Van Woert, M. L., Ackley, S. F., and DeLiberty, T. L.: The thickness distribution
1169 of Antarctic sea ice. *J. Geophys. Res.*, 2008, 113, <https://doi.org/10.1029/2007JC004254>, 2008.

1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186

1189 **Table 1.** Overview of relevant multi-channel satellite microwave sensors.

sensor	relevant frequencies [GHz]	operation periods
Scanning Multichannel Microwave Radiometer (SMMR)	6.6, 10.7, 18.0, 21.0, 37.0	1978-10-25 – 1987-08-20
Special Sensor Microwave/Imager (SSM/I)	19.4, 22.2, 37.0, 85.5	1987-07-09 – today
Special Sensor Microwave Imager and Sounder (SSMIS)	19.4, 22.2, 37.0, 91.7	2003-10-18 – today
Advanced Microwave Scanning Radiometer on EOS (AMSR-E)	6.9, 10.7, 18.7, 23.8, 36.5, 89.0	2002-05-05 – 2011-10-04
Advanced Microwave Scanning Radiometer 2 (AMSR2)	6.9, 7.3, 10.7, 18.7, 23.8, 36.5, 89.0	2012-05-18 – today

1191 **Table 2.** Overview of the investigated sea-ice concentration products. Column “ID (Algorithm)” holds the identifier we use henceforth to
 1192 refer to the data record, and which algorithm it uses. Group is an identifier for the retrieval concept used. Column “Input data” refers to the
 1193 input satellite data for the data set. Column “Open water filter” refers to whether weather-related spurious sea-ice concentrations in open
 1194 water and low concentration areas are filtered. Weather filters do not remove weather related noise over areas with near 100% sea-ice
 1195 concentration. Column “Atmospheric correction” refers to correcting the input TBs for a potential inherent weather influence using additional
 1196 independent weather information. Column “Error” refers to provision of sea-ice concentration uncertainties, and “Period” is the time period
 1197 for which we use the data set, given as StartYearStartMonth-EndYearEndMonth.

ID (algorithm)	Group	Input data & frequencies	Grid resolution & type	Open water filter	Atmospheric correction	Error	Period
OSI-450 (SICCI2)	I	SMMR, SSM/I, SSMIS 19.35 & 37.0 GHz	25 km x 25 km EASE2.0	Yes	Yes	Yes	197901-201512
SICCI-12km (SICCI2)	I	AMSR-E, AMSR2 18.7 & 89.0 GHz	12.5 km x 12.5 km EASE2.0	Yes	Yes	Yes	200205-201705
SICCI-25km (SICCI2)	I	AMSR-E, AMSR2 18.7 & 36.5 GHz	25 km x 25 km EASE2.0	Yes	Yes	Yes	200205-201705
SICCI-50km (SICCI2)	I	AMSR-E, AMSR2 6.9 & 36.5 GHz	50 km x 50 km EASE2.0	Yes	Yes	Yes	200205-201705
ASI-SSMI (ASI)	III	SSM/I, SSMIS 85.5 GHz	12.5 km x 12.5 km PolarStereo	Yes	No	No	199201-201812
CBT-AMSRE (Comiso-Bootstrap)	II	AMSR-E 18.7 & 36.5 GHz	25 km x 25 km PolarStereo	Yes	No	No	200205-201109
CBT-SSMI (Comiso-Bootstrap)	II	SMMR, SSM/I, SSMIS 19.35 & 37.0 GHz	25 km x 25 km PolarStereo	Yes	No	No	197810-201712
NT1-SSMI (NASA-Team)	III	SMMR, SSM/I, SSMIS 19.35 & 37.0 GHz	25 km x 25 km PolarStereo	No	No	No	197810-201712
NOAA-CDR (Comiso-Bootstrap & NASA-Team)	II	SSM/I, SSMIS 19.35 & 37.0 GHz	25 km x 25 km PolarStereo	Yes	No	Yes	198708-201712
NT2-AMSRE (NASA-Team-2)	IV	AMSR-E 18.7, 36.5 & 89.0 GHz	25 km x 25 km PolarStereo	Yes	Yes	No	200205-201109

1200 **Table 3.** Inter-comparison results to near-100% SIC (RRDP2) for the Arctic (see Fig. 14 a). Rows “Gaussian”: Mean difference of modal
 1201 value of Gaussian fit to satellite SIC \leq 99% (compare Fig. 3) minus near-100% reference SIC (RRDP2 SIC) \pm one sigma of the Gaussian fit
 1202 (see Fig. 14 a). Rows “Non-truncated”: Mean difference satellite SIC minus RRDP2 SIC \pm one standard deviation of the difference for
 1203 SICCI and OSI-450 products. Rows “Truncated”: Mean difference of satellite SIC constrained to a maximum value of 100% minus RRDP2
 1204 SIC \pm one standard deviation of the difference. All values in these rows are given in percent sea-ice concentration. Values in rows denoting
 1205 the periods 2007-2015 and 2007-2011, contain the number of valid data pairs. See text for meaning of * in column “NT2-AMSRE”.

Group	I				III			II			IV
	SICCI-12km	SICCI-25km	SICCI-50km	OSI-450	ASI-SSMI	NT1-SSMI	CBT-SSMI	NOAA-CDR	CBT-AMSRE	NT2-AMSRE	
NH											
2007-2015	23262	23262	23262	23343	23343	23343	23343	23037	--	--	
Gaussian	-2.4 \pm 5.2	-1.2 \pm 3.1	-0.5 \pm 1.9	-1.0 \pm 3.0	-4.1 \pm 3.6	+0.1 \pm 5.2	+1.4 \pm 4.5	+2.7 \pm 4.6	--	--	
Non-truncated	-4.2 \pm 5.9	-2.2 \pm 3.7	-0.5 \pm 2.1	-1.9 \pm 3.6	--	--	--	--	--	--	
Truncated	-4.8 \pm 5.2	-2.7 \pm 3.1	-1.1 \pm 1.5	-2.4 \pm 3.0	-4.5 \pm 3.5	-2.6 \pm 4.5	-1.1 \pm 1.9	-0.7 \pm 1.6	--	--	
2007-2011	13351	13351	13351	13432	13432	13432	13432	13126	13344	13344	
Gaussian	-2.4 \pm 5.0	-1.0 \pm 2.9	-0.4 \pm 1.9	-0.8 \pm 2.8	-3.7 \pm 3.7	+0.9 \pm 4.6	+1.3 \pm 3.6	+3.5 \pm 5.0	+1.0 \pm 3.9	-0.7 \pm 1.7*	
Non-truncated	-4.2 \pm 5.4	-2.0 \pm 3.5	-0.6 \pm 2.0	-1.7 \pm 3.3	--	--	--	--	--	--	
Truncated	-5.0 \pm 5.0	-2.8 \pm 3.1	-1.4 \pm 1.6	-2.2 \pm 2.7	-3.9 \pm 3.1	-1.9 \pm 3.7	-0.9 \pm 1.7	-0.6 \pm 1.4	-1.1 \pm 1.8	-0.9 \pm 1.0	

1206

1207 **Table 4.** As Table 3 but for the Antarctic (see Fig. 14 b).

Group	I				III		II			IV
	SICCI-12km	SICCI-25km	SICCI-50km	OSI-450	ASI-SSMI	NT1-SSMI	CBT-SSMI	NOAA-CDR	CBT-AMSRE	NT2-AMSRE
2007-2015	6397	6397	6397	6449	6449	6449	6449	6430	--	--
Gaussian	-0.7±3.7	-1.1±3.0	-0.3±2.5	-1.1±3.1	-6.2±3.9	-5.1±5.9	+0.2±4.5	+0.8±4.4	--	--
Non-truncated	-0.7±4.0	-1.4±4.0	-0.7±2.4	-1.5±3.8	--	--	--	--	--	--
Truncated	-1.9±2.9	-2.3±3.0	-1.3±1.7	-2.3±2.9	-6.5±4.0	-6.0±5.5	-1.7±2.4	-1.3±2.1	--	--
2007-2011	5896	5896	5896	5896	5896	5896	5896	5877	5896	5896
Gaussian	-0.6±4.2	-1.1±3.0	-0.3±2.5	-1.1±3.1	-5.6±4.0	-5.4±6.5	-0.2±4.6	+0.9±4.5	+0.2±5.4	-0.2±3.2*
Non-truncated	-0.7±3.9	-1.5±3.8	-0.8±2.4	-1.5±3.7	--	--	--	--	--	--
Truncated	-2.1±2.9	-2.6±3.1	-1.6±1.9	-2.2±2.8	-6.4±4.0	-5.8±5.5	-1.7±2.4	-1.3±2.1	-1.8±2.5	-0.3±0.5

1208

1209 **Table 5.** Summary of the statistics of the comparison between daily mean ship-based and satellite SIC data (see Fig. 15, black symbols) for
1210 – from top to bottom - the entire year, only winter and only summer. DIFF is the mean difference satellite minus ship-based SIC, SDEV is
1211 the respective standard deviation; R² is the squared linear correlation coefficient. All concentration values are given in percent.

Group	I				III		II			IV
	SICCI-12km	SICCI-25km	SICCI-50km	OSI-450	ASI-SSMI	NT1-SSMI	CBT-SSMI	NOAA-CDR	CBT-AMSRE	NT2-AMSRE
All year										
DIFF	-6.9	-7.8	-7.3	-7.3	-5.4	-13.8	+0.4	+0.6	-0.7	-0.7
SDEV	12.0	12.1	12.4	12.9	16.1	14.5	13.4	13.3	12.9	13.3
R ²	0.784	0.781	0.775	0.734	0.647	0.693	0.737	0.745	0.778	0.767
Winter										
DIFF	-7.4	-7.4	-6.2	-7.4	-8.6	-14.2	< 0.1	-0.2	-1.5	-0.3
SDEV	12.6	11.8	11.8	12.8	17.4	13.8	10.9	11.6	12.6	11.5
R ²	0.558	0.594	0.606	0.591	0.429	0.507	0.595	0.587	0.551	0.595
Summer										
DIFF	-6.7	-8.0	-7.9	-7.3	-3.7	-13.6	+0.7	+0.9	-0.3	-0.9
SDEV	11.7	12.3	12.7	12.9	15.1	14.9	14.5	14.0	13.1	14.1
R ²	0.814	0.793	0.780	0.754	0.722	0.702	0.734	0.750	0.806	0.771

1212

1213 **Table 6.** As Table 5 but for the Antarctic (see Fig. 16, black symbols).

Group	I				III		II			IV
	SICCI-12km	SICCI-25km	SICCI-50km	OSI-450	ASI-SSMI	NT1-SSMI	CBT-SSMI	NOAA-CDR	CBT-AMSRE	NT2-AMSRE
All year										
DIFF	-3.0	-4.4	-3.1	-3.8	-3.3	-11.0	-1.8	-2.3	-1.4	+4.5
SDEV	13.4	13.8	14.0	13.7	15.7	14.8	15.2	15.5	14.8	16.9
R ²	0.763	0.745	0.737	0.733	0.671	0.698	0.711	0.716	0.755	0.679
Winter										
DIFF	-1.6	-2.7	-2.6	-3.2	-3.6	-11.6	-1.6	-2.0	+0.2	+3.8
SDEV	9.8	9.6	10.5	10.5	10.6	11.7	10.7	11.0	9.5	10.7
R ²	0.771	0.771	0.741	0.731	0.659	0.700	0.748	0.751	0.753	0.732
Summer										
DIFF	-3.9	-5.6	-3.4	-4.2	-3.1	-10.6	-2.0	-2.5	-2.5	+5.0
SDEV	15.3	16.1	16.0	15.6	18.4	16.6	17.7	17.9	17.4	20.0
R ²	0.698	0.666	0.675	0.667	0.614	0.640	0.643	0.651	0.693	0.621

1214

1215

1216

1217

1218

1219

1220

1221

1222


1223

1224

1225 **Table 7.** Summary of hypothetical daily average sea-ice concentrations from Figure 17 for top: the pack ice case (a) and bottom the marginal
 1226 ice zone (MIZ) (b). Left two columns: sea-ice concentrations as shown in Figure 17; middle two columns: illustration of the effect of sea-
 1227 ice concentration under-estimation for thin ice; right two columns: illustration of the effect of sea-ice concentration over-estimation, e.g.
 1228 thick ice with a wet snow cover (top), or of sea-ice concentration under-estimation, e.g. MIZ during end of summer (bottom).
 1229


Pack ice, a)	Lead is open water		Lead is thin ice; Satellite underestimates by 20%		Lead is open water; ice surface properties cause 5% over-estimation on 50% of the ice	
	Ship	Satellite	Ship	Satellite	Ship	Satellite
Day 1	48	92	100	98	48	96
Day 2	100	95	100	99	100	98
Day 3	68	90	100	97	68	95
MIZ, b)			50% of sea ice is thin ice; Satellite underestimates by 20%		50% of sea ice is soaked wet; Satellite fails to see this as ice	
	Ship	Sat	Ship	Satellite	Ship	Satellite
Day 1	35	40	35	36	35	20
Day 2	67	38	67	35	67	19
Day 3	63	63	63	57	63	44

1230
 1231
 1232
 1233
 1234
 1235
 1236
 1237
 1238
 1239
 1240
 1241
 1242
 1243
 1244
 1245
 1246
 1247
 1248
 1249
 1250
 1251
 1252
 1253
 1254
 1255
 1256
 1257
 1258
 1259
 1260
 1261
 1262
 1263




1266
 1267

1268 **Figure 1.** Time series of the monthly mean 5%-percentile sea-ice concentration of the range [0.0% to 30.0%] for the Arctic in (a) March
 1269 and (g) September for all ten products. (b) to (f) and (h) to (l) Daily cumulative sea-ice concentration distributions of five selected products
 1270 in these two months, respectively, in a sample year: 2004. See Table 2 for the time periods with data from the respective products.



1271
 1272
 1273
 1274
 1275

Figure 2. Time series of the monthly mean 5%-percentile sea-ice concentration of the range [0.0% to 30.0%] for the Antarctic in (a) September and (g) February for all ten products. (b) to (f) and (h) to (l) Daily cumulative sea-ice concentration distributions of five selected products in these two months, respectively, in a sample year: 2004. See Table 2 for the time periods with data from the respective products.



1276

1277

1278

1279


1280

1281

1282

1283

Figure 3. Examples of the sea-ice concentration distribution at near-100% reference sea-ice concentration locations. Black symbols and lines show values cut off at 100%; blue lines denote the original distribution (for SICCI-25km and SICCI-50km only); red lines denote the distribution resulting from the Gaussian fit to values of the distribution $\leq 99\%$. In each panel, the modal sea-ice concentration (= center of the Gaussian fit: C_{modal}), the standard deviation of the fit σ_c and fit parameters with respect to the fraction of the distribution $\leq 99\%$ (F_{99} , ΔF_{99} , see text for more explanation) and the root-mean-squared difference (RMSD) between original and fitted probability are given. **(a)** to **(c)** Arctic, **(d)** to **(f)** Antarctic. See Appendix H for Fig. H1 and Fig. H2 containing plots of this kind for all ten products.




1284

1285


1286

Figure 4. Illustration of the typical distribution of near-100% SIC reference data by means of the co-located OSI-450 sea-ice concentration for **(a,b)** the Arctic and **(c,d)** the Antarctic in a year with good (left) and poor (right) data coverage.



1287
 1288
 1289
 1290

Figure 5. Spatiotemporal distribution of the ship tracks for (a,b) the Arctic and (c,d) the Antarctic from which ship-based visual observations of the sea-ice cover were used. Maps on the left illustrate the years, maps on the right distinguish between winter (red) and summer (cyan) months.




1291

1292

1293

Figure 6. Arctic sea-ice area (a,b) and extent (c,d) computed in (a,c) winter (March) and (b,d) summer (September) from the sea-ice concentration data sets used. See Table 2 for start and end month of the respective time series.




1294

1295

Figure 7. As Fig. 6 but for the Antarctic.

1296




1297

1298


1299

1300

Figure 8. (a) to (j) Maps of the difference between the multi-annual average monthly SIC of the individual algorithms and the 10-algorithm ensemble median multi-annual average monthly SIC (k) for the Arctic in winter (March) 2003-2011. Differences are only computed for sea-ice concentration of both data sets > 15%. Roman numbers I to IV denote the group assigned to the respective algorithm (see text for details).



1301
 1302 **Figure 9.** Differences (row minus column) between all ten products of, from left to right, the average sea-ice concentration SIC, average
 1303 SIA, and average SIE for the Arctic (NH) in winter (a) through (c) and summer (d) through (f). The averages are computed from monthly
 1304 mean values of the respective months (MARch, SEPTember) of the AMSR-E period 06/2002 to 09/2011. All data are on EASE 2.0 grid with
 1305 50 km grid resolution. The land-mask of the SICCI-50km product is applied to all products. Roman numbers I to IV denote groups according
 1306 to Table 2. For matrices of all remaining months we refer to Fig. G1 through Fig. G3 in Appendix G.
 1307



1308

1309 **Figure 10.** (a) to (j) Maps of the difference between the multi-annual average monthly SIC of the individual algorithms and the 10-algorithm
 1310 ensemble median multi-annual average monthly SIC (k) for the Antarctic in winter (September) 2002-2011. Differences are only computed
 1311 for sea-ice concentration of both data sets > 15%. Roman numbers I to IV denote the group assigned to the respective algorithm (see text for
 1312 details).






Figure 11. As Fig. 10 but in summer (January) 2003-2011.

1313
 1314
 1315
 1316




1317
1318
1319
1320
1321
1322
1323
1324

Figure 12. Differences (row minus column) between all ten products of, from left to right, the average sea-ice concentration SIC, average SIA, and average SIE for the Antarctic (SH) in winter (a) through (c) and summer (d) through (f). The averages are computed from monthly mean values of the respective months (SEPtember, FEBruary) of the AMSR-E period 06/2002 to 09/2011. All data are on EASE 2.0 grid with 50 km grid resolution. The land-mask of the SICCI-50km product is applied to all products. Roman numbers I to IV denote groups according to Table 2. For matrices of all remaining months we refer to Fig. G4 through Fig. G6 in Appendix G.




1325
1326
1327
1328

Figure 13. Cumulative distribution of the SIC difference to 100% at the near-100% reference SIC locations for all ten algorithms based on data of years 2007 through 2011 for (a) the Arctic, and (b) the Antarctic. Roman numbers at the top denote product groups (see Table 2).



1329

1330 **Figure 14.** Summary of the results of the inter-comparison to the near-100% reference SIC (RRDP2) for (a) the Arctic and (b) the Antarctic.
 1331 Shown for each sea-ice concentration product in black is the center of the Gaussian fit of the sea-ice concentration $\leq 99\%$ at the respective
 1332 RRDP2 locations (see Fig. 4). Bars denote \pm one standard deviation of the Gaussian fit. For symbol pairs (all products but CBT-AMSRE
 1333 and NT2-AMSRE) the left symbol is based on data of years 2007-2011, the right one on data of years 2007-2015. Blue symbols denote the
 1334 mean non-truncated sea-ice concentration at the near-100% reference SIC for OSI-450 and SICCI products; blue bars denote the respective
 1335 sea-ice concentration standard deviation. Roman numerals at the top denote product groups (see Table 2). No values are shown for NT2-
 1336 AMSRE, because a Gaussian fit could not be adequately applied.



1337

1338

1339

1340

1341

1342

1343

1344

1345

1346

Figure 15. Scatterplots of co-located daily average SIC from visual ship-based observations (ASPeCt, x-axis, note that these are ASSIST/IceWatch for the Arctic) and the ten satellite SIC algorithm products (SAT, y-axes) for the Arctic during years 2002-2011. Red symbols denote the average satellite SIC binned into 10% ASPeCt SIC intervals (except 0 ... 5% and 95% ... 100%, where 5% bins are used). Blue symbols denote the average ASPeCt SIC binned into 10% satellite SIC intervals, respectively. Error bars denote one standard deviation of the average. Dotted lines denote the identity line. Solid lines denote the linear regression of the respective value pairs. The mean difference and the standard deviation, the linear regression equation, the number of valid data pairs (N), and the squared linear correlation coefficient (R^2) is given in the top left or every image for the daily SIC values. Roman numbers I to IV denote the group assigned to the respective algorithm (see text for details).






Figure 16. As Fig. 15 but for the Antarctic.

1347
 1348
 1349
 1350
 1351
 1352
 1353
 1354
 1355



1356

1357


1358

1359

Figure 17. Illustration of the representivity of ship-based observations (red ellipses and numbers) compared to gridded satellite observations (black grid and numbers) for a) close pack ice with leads and b) an open sea-ice cover in the marginal ice zone. Size of ellipses is in scale with the grid-cell size of 25 km by 25 km. Short black bars denote transitions between days. See also Table 7.

NH SIC

minus




1360
1361
1362
1363

Figure G1. Differences between all ten products of the average sea-ice concentration for the Arctic. The averages are computed from monthly mean values of the respective months of the AMSR-E period 06/2002 to 09/2011. All data are on EASE 2.0 grid with 50 km grid resolution. The land-mask of the SICCI-50km product is applied to all products.


SH SIC

minus



1364
1365


Figure G4. As Fig. G1 but for the Antarctic.



1366
1367
1368
1369

Figure G2. Differences between all ten products of the average sea-ice area (SIA) for the Arctic. The differences are computed from monthly mean SIA of the respective months of the AMSR-E period 06/2002 to 09/2011. All data are on EASE 2.0 grid with 50 km grid resolution. The land-mask of the SICCI-50km product is applied to all products.

SH SIA



1370
1371

Figure G5. As Fig. G2 but for the Antarctic. Note the larger range of the SIA differences compared to the Arctic.






Figure G3. Differences between all ten products of the average sea-ice extent (SIE) for the Arctic. The differences are computed from monthly mean SIE of the respective months of the AMSR-E period 06/2002 to 09/2011. All data are on EASE 2.0 grid with 50 km grid resolution. The land-mask of the SICCI-50km product is applied to all products.

SH SIE



1376
1377

Figure G6. As Fig. G3 but for the Antarctic. Note the larger range of the SIE differences compared to the Arctic.



1378

1379

1380

1381


1382

1383

1384

1385

Figure H1. Sea-ice concentration distribution at RRD2 near-100% reference sea-ice concentration locations in the Arctic during winter for 2007-2011. Black symbols and lines show values cut off at 100%; blue lines denote the original distribution (for OSI-450, SICCI-12km, SICCI-25km and SICCI-50km); red lines denote the distribution resulting from the Gaussian fit to values of the distribution $\leq 99\%$. In each image the modal sea-ice concentration (= center of the Gaussian fit: C_{modal}), the standard deviation of the fit σ_C and fit parameters with respect to the fraction of the distribution $\leq 99\%$ (F_{99} , ΔF_{99} , see text in Sect. 2.1.4 for more explanation) and the root-mean-squared difference (RMSD) between original and fitted probability are given.



1386
1387

Figure H2. As Fig. H1 but for the Antarctic.