

## Revisions associated with:

**Manuscript #2019-125 by Crawford, Mueller, Crocker, Mingo, Desjardins, Dumont and Babin:** *“Ice island thinning: Rates and model calibration with in situ observations from Baffin Bay, Nunavut”*

6 January 2020

The following document includes:

- Comments from Reviewer 1 and corresponding author responses
- Comments from Reviewer 2 and corresponding author responses
- Other changes made by the authors
- A revised manuscript with all changes tracked

## Reply to Reviewer 1:

Original reviewer comments are in black.

Author replies are in blue.

### General comments

The article “Ice island thinning: Rates and model calibration with in situ observations from Baffin Bay, Nunavut” presents thickness and ablation rates data collected from an ice island, and uses those measurements to calibrate a coefficient present in an equation used to calculate iceberg basal melt in numerical models. The paper is in general very well-written and brings some much-needed iceberg observations to the scientific community. My biggest concern is the definition of temperature gradient used here, which differs from the traditional  $T_{\text{ocean}} - T_{\text{ice}}$ . If ice temperature was taken into consideration, their calibrated value for  $C$  would probably be an order of magnitude smaller. Because of that, I recommend this paper for publication in The Cryosphere after major revisions.

We greatly appreciate the reviewer’s thorough and informed comments regarding our manuscript, as well as the acknowledgement that our work contributes valuable in situ observations that are largely lacking in this field of research. A complete response regarding the  $T_{\text{ice}}$  temperature assignment can be found below in our replies to the reviewer’s specific comments. Our current approach for determining  $T_{\text{ice}}$  is, in fact, appropriate as we carefully demonstrate. However, we provide additional text regarding the approach taken and state the calibration result is specific to this approach. Our paper has been strengthened by addressing this comment, as doing so provided us with the opportunity to thoroughly research regarding how  $T_{\text{ice}}$  is assigned in this model. We highlight and important divergence in approach among papers, discuss the implications and make recommendations.

### Specific comments

1. Lines 28-29: “Approximately 30 to 60 % of the freshwater flux from the Greenland Ice Sheet is in the form of solid ice discharge, i.e. iceberg calving”. Actually, solid ice discharge is not equivalent to iceberg calving. As described by Bamber et al. (2018, JGR Oceans 123(3),1827-1837), “Solid ice discharge is the product of surface velocity and ice thickness along an outlet glacier flux gate, typically located near and upstream of the grounding line”. While possibly most of it will be calved, part of this solid ice will be subjected to submarine melting as well.

The reviewer is correct, and we thank them for bringing up this important distinction. We have removed the sentence and now begin the Introduction with new subject matter regarding an anticipated future calving event from Petermann Glacier.

## 2. Section 3.1:

(a) You mention first that ice thickness was recorded daily. Then you say the ice thickness was linearly interpolated between dates used as calibration intervals (6 of them). Why do you have only 6 points for daily data spanning almost a year? Does that mean that the thickness recorded daily had step-changes from one day to another, say, measuring 105.5 m constantly from December 2015 until it suddenly changed to about 105 m in February 2016 (Fig 3a)? I imagine this is due to the instrument precision (0.67 m), but it would be good to clarify this.

Yes, we interpolated between dates when step changes in the ice thickness measurement were recorded. These step changes were due to the set-up of the stationary ice penetrating radar. We have slightly rearranged the information in the paragraph and add the term ‘step changes’ to clarify this.

(b) Did you compare the sonic ranger measurements of surface ablation to the stakes’ data? I imagine retrieving ablation rates from the stakes has a degree of uncertainty that maybe the SR data could constrain.

We have previously compared these measurements. The very similar trend in surface height and derived basal ablation associated with the sonic ranger data provides confidence in the stake measurements and basal ablation calculations. However, we do not feel that we are able to constrain uncertainty in the stake data with the sonic ranger data due to unknown differences in the surface conditions between their locations. Using the weekly photographs, we are confident that the surface conditions between the stationary ice penetrating radar and the stakes match well whereas the surface conditions were not monitored at the location of the SR50. The difference in surface conditions between the sIPR and the SR50 likely caused the unrealistic basal ablation time series derived from the SR50 data. The latter point is now added in the last paragraph of section 3.1.

## 3. Section 3.2.2:

(a) If  $M_b$  is in m/day and the other variables have units in seconds, you need to include, on the right-hand side of equation 1, a multiplying factor of 86400.

The multiplying factor has been added.

(b) You calculate  $\Delta T$  as the difference between the ocean temperature at the keel depth (ocean temperature) and the melting point of ice. Why? The original equation defines  $\Delta T$  as the difference between the ocean temperature and ice temperature. While FitzMaurice et al. (2017) assume  $T_{ice} = -15^\circ\text{C}$ , in models, ice temperature is generally assumed  $-4^\circ\text{C}$ . If you use  $T_{ice} = -4^\circ\text{C}$  and considering the temperatures you normally have at the keel (as per Fig 4a), you will have  $\Delta T = T_{ocean} - T_{ice} = -1.5 + 4 = 2.5^\circ\text{C}$ , which is an order of magnitude larger than your usual  $\Delta T$  (Table 2). If  $\Delta T$ , in turn, is one order of magnitude larger, using the same  $M_b$ , you will get a  $C$  one order of magnitude smaller ( $10^{-6}$ ) which is consistent with the value given by Weeks and Campbell (1986).

Our approach to assigning  $\Delta T$  follows that of Løset (1993a) and the iceberg and ice island deterioration modelling of Kubat et al. (2007) and Ballicater Consulting (2012). We acknowledge that another line of literature has assigned  $-4^\circ\text{C}$  to  $T_{ice}$ , and we reviewed this literature in depth after the topic of ice temperature assignment was brought forward. The foundational literature of Weeks and Campbell (1973) and White (1980) do not explicitly state how  $T_{ice}$  was assigned in their work, but it can be inferred through their  $\Delta T$  and variable descriptions that  $T_{ice}$  was assigned a value representative of the ice interface temperature. The assignment of  $-4^\circ\text{C}$  to  $T_{ice}$  references back to Løset (1993b), where the top-most temperatures ( $\sim 50$  cm depth) on several icebergs approached  $-4^\circ\text{C}$  (see Figure 4 in that paper) and 2D model output shows that this temperature is reached within a few meters of the ice-ocean interface (see figures 8 and 9 of that paper). Løset (1993b) does not explicitly mention this temperature or recommendation of its use in the paper itself. Overall, it is a challenge to determine the single correct value to assign to  $T_{ice}$ , as this should consider the evolving temperature gradient between the ocean interface and iceberg core. We use the method of Løset (1993a), as it is empirically based on work conducted by Josberger (1977) and is also supported by FitzMaurice and Stern (2018). We have added explicitly statement regarding the calibrated coefficient fitting the value assigned to, or approach used to derive,  $T_{ice}$  in the Abstract and Conclusion. We also added a paragraph to

section 5.1: Basal Ablation Model Calibration that discusses the ice temperature assignment in our work as well as in previous literature. Examples of how our calibrated value of  $C$  would differ due to the use of different  $T_{ice}$  values are also provided in this location. We appreciate the push to investigate this topic, as it opened an interesting line of discussion regarding best practices for assigning  $T_{ice}$ .

4. Lines 273-276: “While intervals 3, 4 and 5 were characterized by relatively **low**  $\Theta$  along with high  $S_A$  and  $u$  values (Fig. 4), it is difficult to draw conclusions regarding the alignment of oceanographic conditions and  $C_i$  values due to the low resolution of the sIPR thickness measurements.”

In fact, the theoretical way to calculate  $C$  is through

$$C = \frac{kPr^{\frac{1}{3}}\alpha}{\rho_i\Gamma v^m} \quad (1)$$

where  $\alpha$  and  $m$  are dependent on iceberg shape,  $k$  is the thermal conductivity,  $Pr$  is the Prandtl number (function of  $k$ ),  $\rho_i$  is the ice density,  $\Gamma$  is the latent heat of ice and  $v$  is the kinematic viscosity (see Supporting Information for FitzMaurice et al. 2017). So, although some of these variables are dependent on temperature and salinity, I would not expect a straightforward relationship between  $C$  and oceanographic parameters.

We appreciate the point and explanation that is provided. The sentence has been removed.

5. Lines 315-316: “The volume of PII-A-1-f was  $1.4 \pm 0.01 \text{ km}^3$  when it was first visited in October 2015. By September 2016, the volume and areal extent decreased by  $0.4 \pm 0.01 \text{ km}^3$  and  $3.4 \pm 0.1 \text{ km}^2$ , respectively.” - What was its area by the time of the first visit?

We have added the surface area of the ice island when it was first visited in October 2015.

6. Line 345: Take a look at FitzMaurice et al. (2017, Geophys. Res. Lett., 44, 5637-5644, doi:10.1002/2017GL073585). They discuss a new parameterization of iceberg melting due to the influence of attached/detached plumes. This paper is very relevant for your discussion.

We thank the reviewer for bringing the FitzMaurice et al. (2017) paper to our attention. It is indeed relevant to incorporate into our discussion of iceberg/ice island melt model parameterizations. We have included the following text in the relevant paragraph, “(...) FitzMaurice et al. (2017) showed to be the case when parameterizing Eq. (1) for the sidewall melt of an iceberg with different scenarios of meltwater plume and relative ocean velocities.”

7. Lines 348-350: “It is possible that the adjustment to the melting point of ice ( $M_p$ ) to account for the influence of the meltwater plume is not necessary and  $M_p$  will simply equal the far field ocean temperature ( $\Theta_f$ )” - If  $M_p$  was the far field temperature and your  $\Delta T$  is  $T_{keel} - M_p$ , then the basal melt would be close to zero! Once again, it makes more sense to use the temperature of ice. Maybe what you had in mind is to use  $M_p$  instead of  $T_{keel}$ , and do  $M_p - T_{ice}$  to calculate  $\Delta T$ .

The reviewer’s comment led us to discover an important omission in this sentence. We have updated the sentence to read, “It is possible that the adjustment to the melting point of ice ( $M_p$ ) to account for the influence of the meltwater plume is not necessary and  $M_p$  will simply equal the far field ocean **freezing** temperature ( $\Theta_f$ )”. Please see our response to point 3b regarding the ice temperature assignment.

8. Lines 350-352: “Determining this will require concerted study of the difference in the basal boundary layer conditions of grounded versus drifting ice islands. Observations of  $\Delta u$  for the drifting ice island case are rare but would be useful for this work and for correctly assigning values to this variable in Eq. (1).” - It would also be useful to have an estimate of the plume’s vertical velocity, according to FitzMaurice et al. (2017).

150

We agree that an estimate of the plume's vertical velocity would be useful for understanding sidewall boundary layer conditions and modeling sidewall melt. However, our work focuses solely on basal boundary conditions and we do not feel that mentioning this suggestion by FitzMaurice et al. (2017) is appropriate in this location of our manuscript.

155

9. Lines 354-355: It is worth mentioning that those exponents are related to the shape of the iceberg (in this case, taken as a flat plate).

The flat plate nature of tabular icebergs and ice islands is now mentioned.

160

10. Lines 386-387: "and the protection of a meltwater plume" - Again, it depends on the vertical velocity of the plume in relation to  $\Delta u$  (FitzMaurice et al., 2017). You could have a detached plume in a drifting iceberg.

We have added "and the **potential** protection of a meltwater plume" to account for the findings of FitzMaurice et al. (2017). We now also reference this work at this location.

165

### Technical corrections

Line 16: "(...) thereby increasing the risk to marine transport and infrastructure as well as [affecting, impacting] the distribution of freshwater from the polar ice sheets."

170

This line now reads: "'(...) thereby increasing the risk to marine transport and infrastructure as well as affecting the distribution of freshwater from the polar ice sheets.'" We appreciate this catch and suggestion.

175

Lines 19-20: "The majority of thinning (73 %) resulted from basal ablation, but the associated volume loss was 12 times less than that caused by areal reduction" - It is not clear to me if the "associated volume loss" refers to the total thinning or only to the thinning caused by basal ablation.

We have clarified the wording so that it is apparent that the "associated volume loss" refers to the thinning caused by basal ablation.

180

Lines 34-35: "(...) and impact the biological and physical characteristics of ocean waters in their vicinity **due to meltwater input and latent heat uptake** resulting from their deterioration"

We have incorporated the reviewer's suggested edit.

185

Lines 57-61: "PII-A-1-f was a fragment of the 130 km<sup>2</sup> PII that calved from the Petermann Glacier in northwest Greenland on 5 August 2012 (Crawford et al., 2018a). After calving, the Canadian Ice Service (CIS; Environment and Climate Change Canada) tracked the ice island with RADARSAT-2 synthetic aperture radar (SAR) acquisitions. Between August 2012 and November 2014 the ice island drifted through Nares Strait and Baffin Bay, though it also experienced periods of stagnation while grounded in Kane Basin and northern Baffin Bay (Fig. 1a)" - I assume that the one that "calved from Petermann Glacier in northwest Greenland on 5 August 2012" is the PII. In this case, which ice island are you referring to in the next sentences? PII or its fragment PII-A-1-f?

190

We agree that it was not clear if we were referring to PII and PII-A-1-f. This has been clarified by adding some new information and slightly restructuring parts of the paragraph. It now reads, "PII-A-1-f was a fragment of the 130 km<sup>2</sup> PII that calved from Petermann Glacier in northwest Greenland on 5 August 2012 (Crawford et al., 2018a). Using tracking data derived from RADARSAT-2 satellite images in the Canadian Ice Island Drift Deterioration and Detection (CI2D3) Database (Crawford et al., 2018a) and by the Canadian Ice Service (CIS; Environment and Climate Change Canada), we were able to trace the origins of PII-A-1-f as PII broke up and drifted through Nares Strait and Baffin Bay between August 2012 and

195

- 200 November 2014. As this piece drifted south, it further fragmented and experienced periods of stagnation while grounded in Kane Basin and northern Baffin Bay (Fig. 1a). The PII-A-1-f fragment entered northern Baffin Bay in late 2013. Continued monitoring with RADARSAT-2 acquisitions showed that a portion of the deterioration that PII-A-1-f experienced after 2013 was caused by sidewall notches that progressively enlarged on opposing sides of the ice island.”
- 205 Line 137: “ablation rate ( $M_b$ ;  $m\ d^{-1}$ )” - there is a minus sign missing from d exponent
- Fixed
- Line 139: “ $C\ (m^{2/5}\ s^{1/5}\ ^\circ C^{-1})$ ” - there is a minus sign missing from  $^\circ C$  exponent
- 210 Fixed
- Line 142: “melting point ( $M_p$ ;  $^\circ C$ )” - remove semicolon after  $^\circ C$
- 215 Removed
- Equation 1: using fractional exponents makes the layout of this equation confusing (it looks like  $\Delta T$  is part of the 4/5). I suggest using the typical notation:  $M_b = 86400C\Delta u^{0.8} \frac{\Delta T}{L^{0.2}}$ .
- 220 We agree and have modified the layout of the equation.
- Line 150: I would rephrase that as “Values of  $C$  were obtained for each calibration interval  $i$  ( $C_i$ )”, since up to that point you have only defined  $C$ .
- 225 We agree and have made this change.
- Line 180: I think you meant “it is the time between the recording of the air and reflected waves by the receiver”
- 230 Fixed
- Line 184: “However, an insufficient number (...)” - Since you use “However,” again on line 187, you could remove this one without losing any meaning.
- 235 The first “However” on line 184 was removed.
- Line 195: “the locations recorded by the mIPR onboard GPS were replaced with those recorded by a Hiper V dual-frequency GPS” - Remove first “GPS”
- 240 “GPS” was included in this sentence three times. We have removed the first “GPS” but have kept the latter two as they are referring to specific sensors.
- Lines 267-268: “were an order-of-magnitude larger than those that have been previously calibrated or theoretically derived (Weeks and Campbell, 1986; White et al., 1980).” - I think you should mention here what the theoretically derived value is, so the reader can readily compare it with the following  $C_i$  values.
- 245 The theoretically derived values are now reported, and the mention of ‘previously calibrated’ has been removed. We also corrected the publication year of the Weeks and Campbell work to 1973.

250 Line 308: “The magnitude of thinning (3 to 4 m) **observed** over this thin section **observed** along transect segment AB” - Perhaps replace the second “observed” with “present”

Replacement made

255 Lines 344 and 347: “(...) model skill would likely improve if Eq. (2) was calibrated for drifting vs. grounded ice islands. (...) different parameterizations required for predicting the basal ablation of drifting versus grounded ice islands” – I would change the first sentence (even remove it) to avoid repeating information a few lines below.

The portion of the first sentence that the reviewer notes has been removed to avoid repetition.

260 Line 375: improved

Fixed

265 Line 388: “(...) also contributed to the high, 13.5 m month<sup>-1</sup> basal ablation rate that **Jansen et al. (2007)** estimated (...)”

This citation has been added.

270 Line 409: “that on-ice data.” - ... were collected (?)

Correct, this is now fixed and reads, “that on-ice data were collected”.

275 Table 1: Check the “Dates” column on the first and second ablation periods. I believe it should be 2015-12-04 and 2015-12-05 instead of 2016.

Correct! We thank the reviewer for catching this and have fixed the dates.

280 Table 3 caption: “over the time period that basal ablation was derived with each incremental increase in the value assigned to a given variable.” - I found this sentence very confusing

We have edited this caption. The concerned sentence now reads, “The sensitivity of the model was assessed as the mean percent increase in predicted cumulative basal ablation following each incremental increase in the value assigned to a given variable. The sensitivity was analysed over the longer period (2015-11-15 to 2016-09-18) due to the certainty in the bulk basal ablation magnitude over this time.”

285 Figure 5 caption, line 642: There is a “(“ missing from “b)”

Fixed

290

## Reply to Reviewer 2:

295 Original reviewer comments are in black.

Author replies are in blue.

300 This paper presents measured ablation rates of a tabular iceberg which calved from the Peterman Glacier, North Greenland, which are then used to calibrate a model for melt rates. The *in situ* data set, which comprises repeated ice-penetrating radar surveys, surface mass balance measurements, as well as oceanographic data nearby is very unique, and a valuable contribution to the cryosphere community. Previously, studies on iceberg ablation rates were mostly based on modelling alone, or estimation of ablation rates from satellite altimetry, which is not straight forward because of the unknown surface processes and densification rates of a possible firn layer, which is especially important for the Antarctic icebergs. Thus, I  
305 would see it as an excellent contribution to The Cryosphere.

We thank the reviewer for their comments and their appreciation for the unique data set and basal ablation analysis presented in our manuscript.

310 I would like to rise a few points which might need addressing before publication:

1. At some point (section 3.2.2) it is stated that for the calibration of the model it is assumed that  $\Delta u$  is set equal to the ocean current velocity. As the iceberg is grounded during much of the time this might be a valid assumption. Nevertheless, as the iceberg can be easily tracked by remote sensing data, it would have been possible to look at real drift velocities, and  
315 how they compare to the ocean currents. Some studies have suggested (e.g. Lichey & Hellmer, 2001, Journal of Glaciology) that not only the ocean current is responsible for iceberg drift velocity and direction, but also the wind conditions. A comparison / discussion of these parameters would be an improvement.

We agree with the reviewer, it is the relative speed between the ocean and drifting ice island that is necessary to constrain, and this will be affected by wind speed and direction. It is indeed possible to track ice islands with remote sensing data, and members of the Water and Ice Research Lab at Carleton University made a tremendous effort to do so while creating the Canadian Ice Island Drift, Deterioration and Detection (CI2D3) Database. This database will be used in the future to take on investigations into ice island drift patterns, such as that suggested by the reviewer. We believe that it is much too big of a step to incorporate such an assessment into the current paper. Such a study would also reply upon coarse resolution data for  
325 ice island drift velocities and modelled environmental conditions, which deviates from the *in situ* measurements and ocean conditions used in our assessment. We have added a statement to section 5.1 that notes the influence of both wind and ocean currents on iceberg drift and  $\Delta u$  and reference the work by Lichey and Hellmer (2001) brought forward by the reviewer as well as Kubat et al. (2005).

330 2. In the introduction it is stated that this study is the first of its kind, for Arctic and Antarctic icebergs. However, to my knowledge there was a similar study set up for an Antarctic iceberg (Scambos et al., 2008, Journal of Glaciology) which might deserve a mentioning here.

Scambos et al. (2008) set up automated meteorological stations on two Antarctic icebergs, and paired one station with a stationary ice penetrating radar. Unfortunately, their ice thickness data was unusable as a result of interference from nearby instruments. Due to this, our dataset still includes the first long-term *in situ* ice thickness measurements from an iceberg in either Polar Region. We also present the first spatially distributed thickness change values, which were possible with our repeated mobile ice penetrating radar transects. The study by Scambos et al. (2008) incorporated novel field data and we agree that the work deserves mentioning as it is an important pre-cursor to our work. We have included text that describes  
340 this work in our Discussion (section 5.2).

3. In the discussion the big difference between the ration of basal and surface ablation rates from results of a former study is mentioned. When comparing these results it has to be considered the in case of the other study the Antarctic tabular iceberg started off with a firn column, while the Peterman iceberg did not have any snow cover, so in fact a blue-ice surface. If there is a firn column, surface melt water can percolate into the firn and refreeze, so the mass is not immediately lost. While on blue ice it is more likely to run off. The problem of refreezing melt water and firn densification is the biggest contribution to uncertainty for previous studies estimating ablation of tabular icebergs from altimeter data. For this setting it would be immensely helpful to have an in situ data set like the one presented here. This might be added to the discussion.

We agree that it is important to note the difference in surface conditions between our study subject and the large tabular icebergs in Antarctica. This is included in section 5.2. With this note, we also mention the iceberg firn observations of Scambos et al. (2008) and suggest that a follow-up study with concurrent ice thickness measurements would be of high value.



## Other Author changes:

360 Greg Crocker's affiliation was corrected to Carleton University.

In the Abstract, "The calibration of the basal ablation model, the with such field data...." has been changed to "The calibration of the basal ablation model, the first known to be conducted with field data..."

365 We changed "deployment" to "collection of a long-term *in situ* dataset of ice island thinning" in the last paragraph of the Introduction.

The reference to Mingo et al. (forthcoming) has been updated in section 3.1 and References.

370 Minor edits to language and sentence structure have been made in section 3.2.1. We have added a reference to Oziel et al. (2019) in regard to our ocean velocity measurements in the same section. We also explicitly state that keel depth was derived using the measured ice thickness.

Edits have been made to ensure that 'data' are plural

375  $T_A$  was changed to  $T_a$  in the Results to maintain consistency throughout the paper.

We explicitly state the length of time that basal ablation was estimated in section 4.1. A redundant sentence at the end of this section was removed as well.

380 We have modified the language at the end of section 4.2.1 to reflect our decision (not recommendation) to apply corrections to the oceanographic data.

Weeks et al. (1973) was corrected to Weeks and Campbell (1973) at the start of section 5.1.

385 An additional Løset (1993) reference has been added in regards to the assignment of ice temperature values in section 5.1. The two Løset (1993) reference are now assigned 'a' and 'b'. We now cite and reference FitzMaurice et al. (2016) and FitzMaurice and Stern (2018) in the same section.

390 We have corrected a reference to Ballicater (2012) to Ballicater Consulting (2012) at the end of section 5.2.

The reference to Sazidy et al. (forthcoming) has been updated in section 5.2 and References.

395 "...after an ice island stops freely drifting..." was modified to "...after an ice island stops drifting freely..." in the last paragraph of section 5.2.

A small amount of text was removed from the Conclusions as it was redundant with the preceding section.

400

# Ice island thinning: Rates and model calibration with in situ observations from Baffin Bay, Nunavut

405

Anna J. Crawford<sup>1,2</sup>, Derek Mueller<sup>1</sup>, Gregory Crocker<sup>1,2</sup>, Laurent Mingo<sup>3</sup>, Luc Desjardins<sup>1</sup>, Dany Dumont<sup>4</sup>, Marcel Babin<sup>5</sup>

<sup>1</sup>Department of Geography and Environmental Studies, Carleton University, Ottawa, Ontario, K1S 5B6, Canada

<sup>2</sup>School of Geography and Sustainable Development, University of St Andrews, St Andrews, KY16 9AJ, United Kingdom

410 <sup>3</sup>Blue System Integration Ltd., Vancouver, British Columbia, V5W 3H4, Canada

<sup>4</sup>Institut des sciences de la mer de Rimouski, Université du Québec à Rimouski, Rimouski, Québec, G5L 3A1, Canada

<sup>5</sup>Département de Biologie, Université Laval, Québec, Québec, G1V 0A6, Canada

*Correspondence to:* Anna J. Crawford (ajc44@st-andrews.ac.uk)

**Abstract.** A 130 km<sup>2</sup> tabular iceberg calved from Petermann Glacier in northwest Greenland on 5 August 2012. Subsequent  
415 fracturing generated many individual large “ice islands”, including “Petermann Ice Island (PII)-A-1-f”, that drifted between  
Nares Strait and the North Atlantic. Thinning caused by basal and surface ablation increases the likelihood that these ice  
islands will fracture and disperse further, thereby increasing the risk to marine transport and infrastructure as well as  
affecting the distribution of freshwater from the polar ice sheets. We use a unique stationary and mobile ice penetrating radar  
dataset collected over four campaigns to PII-A-1-f to quantify and contextualize ice island surface and basal ablation rates  
420 and calibrate a forced convection basal ablation model. The ice island thinned by 4.7 m over 11 months. The majority of  
thinning (73 %) resulted from basal ablation, but the ~~associated~~-volume loss associated with basal ablation was ~12 times  
less than that caused by areal reduction (e.g. wave erosion, calving, and fracture). However, localized thinning may have  
influenced a large fracture event that occurred along a section of ice that was ~40 m thinner than the remainder of the ice  
island. The calibration of the basal ablation model, the first known to be conducted with ~~such~~-field data, supports assigning  
425 the theoretically-derived value of  $1.2 \times 10^{-5} \text{ m}^{2/5} \text{ s}^{-1/5} \text{ }^{\circ}\text{C}^{-1}$  to the model’s bulk heat transfer coefficient with the use of an  
empirically estimated ice-ocean interface temperature. -Overall, this work highlights the value of systematically collecting  
ice island field data for analyzing deterioration processes, assessing their connections to ice island morphology, and  
adequately developing models for operational and research purposes.

## 1 Introduction

A nascent ‘ice island’ is currently being monitored as transverse rifts propagate and widen across the floating ice tongue of Petermann Glacier, northwest Greenland (USGS, 2019). Ice shelves and ice tongues in this region episodically calve these large, tabular icebergs, which are morphologically similar to the those common in the Antarctic although thinner and less extensive (Higgins, 1989). — Approximately 30 to 60 % of the freshwater flux from the Greenland Ice Sheet is in the form of solid ice discharge, i.e. iceberg calving (Bamber et al., 2012; Enderlin et al., 2014). Particularly large iceberg calving events that occur at the floating ice tongues in northwest Greenland produce “ice islands” that, through their drift and deterioration, these ice islands disperse freshwater between Nares Strait and the North Atlantic. Their journey to southern latitudes can be disrupted due to. These ice islands often grounding on the continental shelf of Baffin Island during their journey to southern latitudes (Crawford et al., 2018a). Such groundings can disrupt benthic ecosystems (Dowdeswell and Bamber, 2007), put seafloor infrastructure at risk (Fuglem et al., 2017), extend and stabilize landfast ice covers offshore (Fraser et al. 2012; Massom et al. 2001), and impact the biological and physical composition of ocean waters in their vicinity due to concentrated meltwater input resulting from their deterioration and latent heat uptake resulting from their deterioration (Jansen et al., 2007; Stern et al., 2015).

Ice island thinning has previously been estimated via freeboard monitoring with satellite-borne altimetry data (Bouhier et al., 2018; Jansen et al., 2007). Without field observations, the individual contribution of surface versus basal ablation to that thinning can then only be derived through modelling (Ballicater Consulting, 2012; Jansen et al., 2007), often with a relatively simple fluid-dynamics approach being used to model basal ablation in iceberg and ice island studies (Ballicater Consulting, 2012; Merino et al., 2016; Wagner and Eisenman, 2017). This forced convection model is based on the transfer of heat across a flat plate due to the turbulent flow of the underlying water. A bulk heat transfer coefficient is used to estimate an ablation rate based on the differential velocity between the ice island and the ocean current as well as the water temperature across the length of the ice island (Ballicater Consulting, 2012; Bouhier et al., 2018; Weeks and Campbell, 1973). Previous efforts to calibrate basal ablation models have relied upon remotely sensed datasets as well as potentially inaccurate modelled environmental data (Bouhier et al., 2018; Jansen et al., 2007). This is largely due to the challenges associated with collecting field data for determining the individual rates of surface and basal ablation.

We overcame these challenges to collect a unique dataset from “Petermann Ice Island (PII)-A-1-f” between November 2015 and September 2016. PII-A-1-f was grounded on the continental shelf of Baffin Island and had a surface areal extent of 13 km<sup>2</sup> when it was first visited. The field work included repeat mobile ice penetrating radar (mIPR) transects and the deployment collection of a long-term in situ dataset of ice island thinning by of a stationary IPR (sIPR). These were the first such field data to be collected from an ice island or iceberg in either the Arctic or Antarctic. The measurements are used to calibrate the forced convection basal ablation model, which had previously not been validated or calibrated with direct field observations. In addition, the IPR data was were used to assess the spatial and temporal variation in ice island thinning and

ablation rates. Using remotely sensed imagery to monitor areal reduction, the ablation magnitudes are put into context with respect to other processes (e.g. fracture) that contributed to the deterioration of PII-A-1-f.

## 2 Study site

PII-A-1-f was a fragment of the 130 km<sup>2</sup> PII that calved from the Petermann Glacier in northwest Greenland on 5 August 2012 (Crawford et al., 2018a). ~~After calving, Using tracking data derived from RADARSAT-2 satellite images in the Canadian Ice Island Drift, Deterioration and Detection (CI2D3) Database (Crawford et al., 2018a) and by the Canadian Ice Service (CIS; Environment and Climate Change Canada) we were able to trace the origins of PII-A-1-f as PII broke up and drifted through Nares Strait and Baffin Bay between August 2012 and November 2014. tracked the ice island with RADARSAT-2 synthetic aperture radar (SAR) acquisitions. Between August 2012 and November 2014 the ice island drifted~~  
through Nares Strait and Baffin Bay. As this piece drifted south, it further fragmented though it also and experienced periods of stagnation while grounded in Kane Basin and northern Baffin Bay (Fig. 1a). The PII-A-1-f fragment entered northern Baffin Bay in late 2013. Continued monitoring with RADARSAT-2 acquisitions showed that Aa portion of the deterioration that PII-A-1-f experienced after 2013 was caused by sidewall notches that progressively enlarged on opposing sides of the ice island. The notch ‘roots’, or the tips of these wedge-shaped features, were located in the vicinity of a linear surface feature that was identified in a ScanSAR acquisition in November 2012 (Fig. 1b). This deterioration was likely caused by increased wave-induced turbulent heat flux within these wedge-shaped features as described by White et al. (1980).

The linear surface feature and one sidewall notch were still apparent when PII-A-1-f became grounded in November 2014 at 67°23’N, 63°18’W, approximately 10 km from the east coast of Baffin Island and 35 km southeast from the Hamlet of Qikiqtarjuaq, Nunavut (Fig. 1a). Ice island groundings are especially common in the region immediately north of the Cumberland Peninsula due to the presence of many underwater shoals and ridges. PII-A-1-f was visited at this location by three field teams with transportation provided by the CCGS *Amundsen* icebreaker and its helicopter during the annual ArcticNet science cruises in October 2015, July 2016 and September 2016. An additional field campaign was completed in May 2016 when a field team accessed the ice island, then surrounded by sea ice, by snowmobile from Qikiqtarjuaq, Nunavut.

## 3 Methods

### 3.1 Thinning (temporal assessment)

To ascertain the magnitudes and rates of surface and basal ablation, a series of ablation stakes, a 40 MHz sIPR, and a small meteorological station with a sonic ranger and camera were installed on PII-A-1-f on 20 October 2015 (Fig. 2). The ice island position was recorded hourly with a Garmin 6X-HVS GPS (Garmin International, Inc.) and air temperature ( $T_a$ ) was measured with a 109 thermistor (Campbell Scientific Canada Corporation (CSCC)) in a radiation shield at 1-minute intervals

and logged as hourly averages. All meteorological station data were recorded on a CR1000 datalogger (CSCC) and telemetered with an Iridium L-Band modem (9522B; CSCC).

Ice thickness was recorded ~~at a resolution of 0.67 m~~ and telemetered daily by the sIPR (Blue System Integration, Ltd.) until 27 September 2016. Full details regarding this system set-up and measurement specifications can be found in S1 and Mingo et al. (forthcoming). The ice thickness was measured at a resolution of 0.67 m so the ice thickness data observations were linearly interpolated between dates when ~~measured~~ step changes in thickness were recorded (16 November 2015 - 18 September 2016). These dates were used to establish “calibration intervals” for the calibration of the forced convection basal ablation described in the following section.

Daily mean surface ablation was calculated from hourly SR50A sonic ranger (CSCC) height-above-surface (snow or ice) measurements. Surface ablation was also calculated from five ablation stakes that were marked with tape every 10 cm and placed between the meteorological station and the sIPR. A weekly image acquired with a Campbell Scientific CC5MPX camera that was controlled by the meteorological station to visually check on the positioning of the sIPR system also captured these ablation stakes. No snow was present on the date when the stakes and instruments were installed and the average weekly ablation (or accumulation) at these stakes was calculated with ImageJ (v. 10.8.0) software using the markings for scale (Abramoff et al., 2004). Increases in surface height due to the accumulation of snow are presented in this study as negative values. A linear interpolation was applied to estimate daily ablation or accumulation between these observations so that the sIPR, SR50 and ablation stake data were available over a consistent time span. Basal ablation was calculated by subtracting the interpolated ablation magnitudes derived from the stake data from the ice thickness time series. These values are therefore approximations of daily basal ablation rates, as they were based on values of ice thickness that were linearly interpolated across longer time periods. Finally, the ablation stake data were used instead of the SR50 data because an unrealistic basal ablation time series was derived from the SR50 data. This was likely caused by a difference in surface conditions between the sIPR and SR50. It was not possible to assess the surface conditions at the SR50 (30 m away from the sIPR), while the surface conditions were confirmed to be similar between the locations of the ablation stakes and the sIPR (10-15 m away) in weekly images.

## 3.2 Basal ablation model calibration

### 3.2.1 Oceanographic data collection and comparison

The derived basal ablation was used to calibrate the forced convection model described in the Introduction. In addition to the general paucity of ice island thinning measurements, there is a large dearth of in situ oceanographic data available for the validation and calibration of iceberg and ice island numerical deterioration models. Here, we use two oceanographic field datasets to validate and calibrate a time-series of modelled ocean temperature, salinity and current data that spans the full duration for which basal ablation rates were available. Using the oceanographic model data greatly extended the time span over which the model calibration could be conducted.

The first oceanographic field dataset was collected by the CCGS *Amundsen* during its annual ArcticNet science research cruise. Conductivity, temperature, depth (CTD) profiles were conducted around PII-A-1-f on 28 July (4 casts) and 29  
525 September (5 casts) 2016 (Amundsen Science Data Collection, 2016) (Fig. 1c,d). CTD profiles were also collected on 29  
July 2016 (2 casts) and 28 September 2016 (1 cast) close to the location of a sea ice camp that was situated 20 km northwest  
(67° 29' N, 63° 47' W) of the grounding location of PII-A-1-f (Fig. 1a bottom inset). This sea ice camp was operated by the  
GreenEdge project ~~which was~~ based out of Université Laval, Québec (<http://www.greenedgeproject.info/>). The second  
oceanographic field dataset was composed of 39 CTD profiles (SBE 49 FastCAT CT; Sea-Bird Electronics, Inc.) acquired at  
530 the GreenEdge sea ice camp between 20 April and 22 July 2016. Absolute salinity ( $S_A$ ; g kg<sup>-1</sup>) and conservative temperature  
( $\Theta$ ; °C) measurements were reported at 1 m depth bins for all CTD casts. Current speed ( $u$ ; m s<sup>-1</sup>) was calculated as an  
average of ~~measurements of~~ the current vectors ~~taken-measured~~ every 30 sec over 2 m depth bins and recorded every 30  
minutes at the ice camp with a Teledyne/RDI Workhorse Sentinel 300 kHz acoustic Doppler current profiler (ADCP;  
Teledyne RD Instruments; [Oziel et al. 2019](#)). A linear interpolation was applied to fill missing values before daily average  $u$   
535 was calculated for the individual depth bins.

Data from the Copernicus Marine Environment Monitoring Service (CMEMS) Global Ocean Physical Reanalysis  
(GLORYS12V1) product were used for the calibration of the forced convection basal ablation model. This reanalysis  
product, referred to as the “CMEMS data” henceforth, had a spatial resolution of 1/12° and 50 depth levels. Potential  
temperature, practical salinity and the horizontal current velocity components were extracted from the model grid cells that  
540 corresponded with the locations of PII-A-1-f and the GreenEdge sea ice camp at the model depth bin in which the keel of the  
ice island was located over the course of sIPR data collection. Keel depth was calculated [using ice thickness](#) assuming ice  
density ( $\rho_i$ ) = 873 kg m<sup>-3</sup> (Crawford et al., 2018c) and hydrostatic equilibrium. Potential temperature and practical salinity  
were converted to  $\Theta$  and  $S_A$ , respectively, using the Gibbs Sea Water functions provided in the R ‘gsw’ package (Kelly,  
2017).

545 The CMEMS data ~~was-were~~ compared against the in situ oceanographic data to justify its use for calibrating the forced  
convection basal ablation model and to identify any bias in the modelled oceanographic data. Comparisons were conducted  
between: 1) the full CCGS *Amundsen* CTD profiles collected at the location of PII-A-1-f and near the GreenEdge sea ice  
camp location in July and September 2016, 2) the full CCGS *Amundsen* profiles and the CMEMS data profiles, and 3) the  
CMEMS data time series and the mean  $S_A$ ,  $\Theta$  and  $u$  values of all CTD casts and ADCP measurements that fell within the  
550 CMEMS depth bin in which the ice island keel was located. Bias in the CMEMS data was identified as consistent over- or  
under-estimation of the GreenEdge sea ice camp time series of  $S_A$ ,  $\Theta$  and  $u$ . If bias existed in a given variable, the CMEMS  
data ~~waswere~~ corrected by the average daily difference between ~~itthem~~ and the in situ values.

### 3.2.2 Model calibration

555 The forced convection basal ablation model (Eq. (1)) estimates an ablation rate ( $M_b$ ; m ~~d~~<sup>-1</sup>) from the velocity difference  
between the iceberg and the water ( $\Delta u$ ; m s<sup>-1</sup>) and the driving temperature ( $\Delta T$ ) across the length ( $L$ ; m) of an ice face using a

bulk heat transfer coefficient,  $C$  ( $\text{m}^{2/5} \text{s}^{-1/5} \text{°C}^{-1}$ ) (Weeks and Campbell, 1973). The multiplier in Eq. 1 ( $86400 \text{ s d}^{-1}$ ) converts  $M_b$  from  $\text{m s}^{-1}$  to a daily ablation value; it is also common to scale  $C$  by this factor to produce daily melt estimates directly (e.g. Bigg et al., 1997). In the calibration, it was assumed that the ice island was stationary and therefore  $\Delta u$  is simply equivalent to  $u$ .  $\Delta T$  was calculated as the difference between the ocean temperature at the keel depth and the melting point of ice (equivalent to the freezing point of the adjacent sea water). The melting point ( $M_p$ ;  $\text{°C}$ ) was adjusted to account for the influence of meltwater near the ice-water interface with an established empirical relationship with the far field water temperature ( $\Theta$ ) and the freezing temperature of the far field ocean water ( $\Theta_f$ ;  $\text{°C}$ ; Eq. (2)) (Kubat et al., 2007; Løset, 1993a).  $\Theta_f$  was derived with  $S_A$  and pressure ( $p$ ; dbar) as per TEOS-10 conventions (IOC, SCOR and IAPSO, 2010).

$$M_b = 86400 C \Delta u^{0.8} \frac{\Delta T}{L^{0.2}} \quad (1)$$

$$M_p = \Theta_f e^{-0.19(\Theta - \Theta_f)} \quad (2)$$

Values of  $C_i$  were obtained for each calibration interval ii ( $C_i$ ), which are associated with a measured change in ice thickness. The individual  $C_i$  values were calculated by dividing the cumulative basal ablation over the respective calibration interval by the corresponding cumulative driving force (i.e.,  $\sum \Delta u^{0.8} \frac{\Delta T}{L^{0.2}}$ ). The driving force was derived from CMEMS modelled oceanographic data ( $S_A$ ,  $\Theta$ , and  $u$ ) at the ice island keel.  $p$  was set to the pressure at the keel depth.  $L$  was assigned as the median of all distances between the sIPR location and all vertices of the outline of the areal surface extent of PII-A-1-f that was digitized from RADARSAT-2 Fine-Quad (FQ; 8 m nominal resolution) SAR imagery acquired on 27 July 2016 (S2).

A second calibration for each interval was conducted with the CMEMS data after corrections were applied based on the comparisons against the in situ oceanographic data (Section 3.2.1). A final calibration of  $C$  was obtained based on the basal ablation and driving force over the *total* duration over which basal ablation was derived. This final calibration provided a single value of  $C$  as opposed to the previous calibrations where values of  $C_i$  were obtained for each calibration interval.

An analysis was conducted to determine the sensitivity of the basal melt magnitude predicted by the forced convection basal ablation model to variations in  $u$ ,  $\Theta$ , and  $C$ . The two variables and one parameter were individually perturbed across 8 equally spaced intervals that covered their observed and calculated ranges. The non-perturbed variables were held constant at their median values. All assigned values were based on the corrected CMEMS data series. The sensitivity of the forced convection basal ablation model was assessed as the average percent increase in cumulative basal ablation predicted for the duration over which basal ablation was derived with each incremental increase in the value assigned to a given variable/parameter. The sensitivity was analyzed over this longer time period due to the certainty in the bulk basal ablation derived from total ice thinning and surface ablation.

### 3.3 Thinning (spatial assessment)

A repeat transect was conducted to assess spatial variation in thickness and thinning and to verify that the sIPR data ~~were~~ representative with respect to other locations across the ice island. An initial thickness profile was collected with a 25 MHz mIPR (Blue System Integration Ltd.) that was towed by snowmobile over an approximately 3 km transect on 8 May 2016. A series of 10 ablation stakes were also installed along the profile route. Thickness data were collected over 2.4 km of the original transect when the same mIPR system was towed on foot on 28 September 2016. Eight of the stakes were also re-measured at this time. Further details of the mIPR system can be found in S1 and Mingo and Flowers (2010).

All mIPR processing was conducted with Radar Tools (release 0.4), a library of Python scripts and tools that was used to standardize, clean, visualize, and process data contained in the raw radar data (S1; Wilson, 2013). This program was also used to select the location of the air and reflected radar waves. Ice thickness was calculated with Eq. (3) (Wilson, 2012):

$$H = \sqrt{\left(\left(t + \frac{s}{v_a}\right)\frac{v}{2}\right)^2 - \frac{s^2}{4}} \quad (3)$$

where  $H$  is thickness (m),  $s$  is the distance between the transmitting and receiving antennas (m),  $t$  is the time between the recording of the air and reflected waves by the receiver, and  $v_a$  is the speed of the electromagnetic wave in air ( $3.0 \times 10^8 \text{ m s}^{-1}$ ) traveling between the transmitter and receiver (Wilson, 2012). The speed of the radar wave in ice ( $v$ ) was set to  $1.7 \times 10^8 \text{ m s}^{-1}$  (Macheret et al., 1993). The ice thickness resolution ( $\pm 0.5 \text{ m}$ ) was limited by  $v$  and the waveform sampling interval of the mIPR system (Crawford, 2013).

Snow was present on the ice surface when the mIPR transect was conducted in May 2016. ~~However, a~~An insufficient number of snow depth measurements were recorded to adequately account for the snow depth over the transect area and it was not possible to distinguish the snow/ice interface in the radargram. Thickness (ice + snow) in May was therefore calculated using a single velocity value,  $v = 1.7 \times 10^8 \text{ m s}^{-1}$ . However, an additional uncertainty was added to the May ice thickness to account for the possible presence of snow. This uncertainty was based on the mean snow depth (36 cm) recorded at nine locations along the mIPR transect and the amount of time a radar wave would travel through this layer given  $v = 2.0 \times 10^8 \text{ m s}^{-1}$  for snow (Haas and Druckenmiller, 2009). The errors associated with the resolution of the mIPR system and the average snow depth estimate were summed to determine the average amount that the ice thickness, as measured in May, could be overestimated (0.8 m). Uncertainty in thickness change calculations was determined by propagating the uncertainties in ice thickness resulting from the resolution of the mIPR system and the presence of snow.

The magnitude of thinning that occurred between May and September 2016 was calculated between the closest pairs of mIPR traces recorded during each field visit. To improve positional accuracy, the ~~GPS~~ locations recorded by the mIPR onboard GPS were replaced with those recorded by a Hiper V dual-frequency GPS (Topcon Corp.) after precise point positioning (PPP) processing (Natural Resources Canada, 2016). The September transect was adjusted relative to the



orientation of PII-A-1-f in May by matching mIPR traces known to be collected at the same location (i.e., the ablation stakes) on the ice island to correct for a small amount of movement by the ice island between the two field visits. Since it was not possible to retrace the transect exactly due to changes in topography between field visits, the thickness measurements that were compared between May and September were offset by varying distances (meters to 10s of meters). Using Eq. (4), we derived an index of comparability for these measurements from the percentage of overlap between the radar footprints at the ice-water interface. The radii ( $r$ ) of these footprints depend on the center wavelength of the 25 MHz antenna ( $\lambda = 6.8$  m), relative permittivity of ice ( $K$ ; 3), and  $H$  (Leucci et al., 2003),

$$r = \frac{\lambda}{4} + \frac{H}{\sqrt{K+1}}. \quad (4)$$

The positions of thickness change measurements were categorized as  $\geq 50$  % overlap,  $\geq 30$  % overlap,  $< 30$  % overlap, and no overlap.

### 3.4 Surface extent reduction and contributions to deterioration

The areal extent of PII-A-1-f was monitored with seven RADARSAT-2 FQ SAR scenes to determine the relative importance of surface and basal ablation to the total deterioration of the ice island. The SAR scenes were acquired between 1 November 2015 and 23 September 2017, and the image processing and areal extent digitization details are included in S2. We quantified and compared the contributions of basal ablation, surface ablation, and areal reduction processes (e.g. fracture, forced and buoyant convection, wave erosion, and calving) to the overall deterioration of PII-A-1-f over the temporal extent sIPR data collection. We also estimated the contribution of surface and basal ablation to total deterioration over the longer temporal extent that RADARSAT-2 FQ images were acquired. For this estimation, a cubic spline was applied to fill missing cumulative ablation values between 24 September 2016 and 21 October 2016 and it was assumed that the same surface and basal ablation magnitudes that occurred in the first year of observation also occurred between October 2016 and September 2017.

## 4 Results

### 4.1 Thinning (temporal assessment) and environmental conditions

PII-A-1-f decreased in thickness by  $4.7 \pm 1.4$  m over the eleven months that sIPR data ~~was~~were collected and by  $4.0 \pm 1.4$  m during 10 months ~~the length of time~~ that basal ablation was estimated and used for the basal ablation model calibration (Fig. 3a; Table 1). The latter can be divided into three periods related to surface ablation magnitudes as per the ablation stake data (Fig. 3b; Table 1). These are referred to as “ablation periods” ~~for~~in the remainder of the text. Minimal surface ablation was observed over ablation period 1 (November to mid-December 2015). No surface ablation occurred during the second period, which distinguishes it from ablation periods 1 and 3. Therefore, basal ablation was the sole contributor to thinning during ablation period 2 when the mean daily thinning rate was  $0.9 \text{ cm d}^{-1}$  (mid-December 2015 to mid-July 2016). This rate

tripled during the third ablation period, which spanned from mid-July to September 2016, due to the onset of continuous surface ablation in mid-July. Ninety-eight percent of the recorded surface ablation occurred in this period when  $T_A - T_{at}$  was consistently  $> 0$  °C.  $T_A - T_{at}$  was  $> 0$  °C for 15 % of the second ablation period, however, this contributed to melting the snow that had accumulated on the ice surface instead of ice ablation (Fig. 3b; Table 1).

The mean daily basal melt rate ( $M_b$ ) between November 2015 and September 2016 was almost three times greater than the mean rate of daily surface ablation ( $M_s$ ; cm d<sup>-1</sup>). However, this ratio varied between ablation periods. For example, the mean  $M_b$  was seven times greater than the mean  $M_s$  during ablation period 1 when only 2 cm of surface ablation was observed. In contrast, the mean  $M_s$  was approximately 50 % greater than the mean  $M_b$  over the third ablation period when over 1 m of surface ablation was observed over the two-month duration (Fig. 3b; Table 1). The mean  $M_b$  increased between successive ablation periods (Table 1) and the increase in daily mean  $M_b$  from 0.9 to 1.2 cm d<sup>-1</sup> between periods 2 and 3 coincides with an increase in CMEMS temperature and velocity. Basal ablation was responsible for 73 % of the observed thinning between November 2015 and September 2016. The mean  $M_b$  increased between successive ablation periods (Table 1) and the increase in daily mean  $M_b$  from 0.9 to 1.2 cm d<sup>-1</sup> between periods 2 and 3 coincides with an increase in CMEMS temperature and velocity.

## 4.2 Basal ablation model calibration

### 4.2.1 Oceanographic data comparisons

Figure 4 shows the CMEMS  $\Theta$ ,  $S_A$  and  $u$  for the duration that PII-A-1-f basal ablation was derived. The mean  $\Theta$  and  $S_A$  of measurements from CTD casts acquired in the vicinity of PII-A-1-f or the GreenEdge sea ice camp are plotted alongside the model data in Fig. 4a,b. These data are associated with the depth interval of the model data in which the keel of the ice island fell within for the duration of the data collection. The mean  $u$  in this depth interval, as measured by the ADCP moored near to the GreenEdge sea ice camp location, is included in Fig. 4c. The data collected at the GreenEdge sea ice camp ( $\Theta$  and  $S_A$ ) and at the nearby moored ADCP ( $u$ ) provide the longest in situ time series of oceanographic conditions and were used to assess if these variables were consistently over- or under-estimated in the CMEMS data.

The means of the absolute daily difference between  $\Theta$  and  $S_A$  at the GreenEdge sea ice camp and the CMEMS data were 0.07 °C and 0.67 g kg<sup>-1</sup>, respectively. The CMEMS  $S_A$  was consistently greater than the in situ data and was therefore corrected by subtracting 0.67 g kg<sup>-1</sup> (Fig. 4b, dotted lines). A bias in  $\Theta$  was not seen and no correction was applied; however, there was a consistent under-estimation of  $u$  in the CMEMS data (Fig 4c). The CMEMS data ~~was-were~~ corrected by the mean daily difference (0.1 m s<sup>-1</sup>) between the CMEMS and in situ values of  $u$  (Fig. 4c, dotted lines).

CTD profiles collected in the vicinity of PII-A-1-f and the GreenEdge sea ice camp on successive days by the CCGS *Amundsen* in July and September 2016 were plotted (not shown) to compare the oceanographic conditions at the two sites. In the depth interval of interest, the values of  $\Theta$  and  $S_A$  were reasonably similar and not consistently different between sites. This supported our use of the  $S_A$  correction for the CMEMS data that was derived from the difference between the CMEMS

680 data and the longer time series collected at the GreenEdge sea ice camp. This  $S_A$  correction was applied to the CMEMS data associated with the location of PII-A-1-f, which was used for the model calibration (Fig. 4b).

The full CMEMS  $\Theta$  and  $S_A$  profiles were also plotted (not shown) against those acquired by the CCGS *Amundsen* to ensure that the CMEMS data were generally representative of the water column. The CMEMS profiles were reasonable in form and, in the depth interval of interest, CMEMS consistently over-estimated the  $S_A$  recorded by the CTD casts acquired  
685 by the CCGS *Amundsen*. The average over-estimation was  $0.60 \text{ g kg}^{-1}$ . This further supports the ~~recommendation decision~~ to apply the  $0.67 \text{ g kg}^{-1}$  correction to the CMEMS  $S_A$  for the forced convection basal ablation model calibration. The CMEMS  $\Theta$  was both over- and under-estimated in the depth interval of interest, which ~~reinforces the recommendation~~ confirms that ~~to~~ not ~~applying~~ a correction to this variable was appropriate.

#### 4.2.2 Model calibration and sensitivity analysis

690  $C_i$  values were calculated with the uncorrected and corrected CMEMS data for each of the six calibration intervals. The range and mean values of  $C_i$  calculated with the uncorrected CMEMS data were an order-of-magnitude larger than those that ~~have been previously calibrated or were~~ theoretically derived by Weeks and Campbell (1973) ( $6.74 \times 10^{-6} \text{ m}^{2/5} \text{ s}^{-1/5} \text{ }^\circ\text{C}^{-1}$ ) and White et al. (1980) ( $1.2 \times 10^{-5} \text{ m}^{2/5} \text{ s}^{-1/5} \text{ }^\circ\text{C}^{-1}$ ). Due to this, plus the poor representation of uncorrected CMEMS  $S_A$  and  $u$ , it was decided to further analyze the more reasonable  $C_i$  values that were calculated with the corrected CMEMS data. These  
695 latter values of  $C_i$  are included in Table 2 with the corresponding mean  $\Delta T$  and  $u$  values found with the CMEMS data.

Using ~~this~~ these data, the mean  $C_i$  value was  $1.7 \times 10^{-5} \text{ m}^{2/5} \text{ s}^{-1/5} \text{ }^\circ\text{C}^{-1}$  and values for the individual calibration intervals ranged from  $7.8 \times 10^{-6}$  to  $2.1 \times 10^{-5} \text{ m}^{2/5} \text{ s}^{-1/5} \text{ }^\circ\text{C}^{-1}$ . The greatest values of  $C_i$  were associated with calibration intervals 3 through 6. ~~While intervals 3, 4 and 5 were characterized by relatively low  $\Theta$  along with high  $S_A$  and  $u$  values (Fig. 4), it is difficult to draw conclusions regarding the alignment of oceanographic conditions and  $C_i$  values due to the low resolution of the SIPR thickness measurements.~~ Cumulative basal ablation was over-predicted by 38% when the forced convection basal  
700 ablation model was run with the mean  $C_i$  value over the 308 days that basal ablation was calculated. Finally, when calculated with the total basal ablation and total driving force between 15 November 2015 and 18 September 2016,  $C$  was calibrated to  $1.2 \times 10^{-5} \text{ m}^{2/5} \text{ s}^{-1/5} \text{ }^\circ\text{C}^{-1}$ .

Values of  $u$ ,  $\Theta$ , and  $L$  were not normally distributed. Therefore, the median values of these variables were first assigned  
705 during the sensitivity analysis when a given variable was held constant. The calibrated  $C_i$  values were found to follow a normal distribution, though the power of this test is low due to the small sample size. For consistency, the median  $C_i$  value was also assigned when this parameter was held constant during the sensitivity analysis.

Of the environmental driving variables included in the forced convection basal ablation model,  $\Delta T$  ~~has~~ is the greatest relative range of values over all of the calibration intervals (0.10 to  $0.37 \text{ }^\circ\text{C}$ ). A mean increase of 19 % was applied to the  $\Delta T$   
710 value during the sensitivity analysis and the same increase in cumulative basal ablation was predicted over the time period that basal ablation was derived with observations. This is due to the linear relationship between  $\Delta T$  and  $M_b$  in Eq. (1) (Table 3).

-A linear relationship also exists between  $C$  and  $M_b$  in Eq. (1); however, the range in  $C_i$  was more constrained relative to the  $\Delta T$  data series. For this reason, the cumulative predicted basal ablation only increased by 16.5 % with each incremental change in the value assigned to  $C$  during the sensitivity analysis (Table 3). Finally, due to the smaller range of  $u$  (0.11 to 0.18 m s<sup>-1</sup>) and the non-linear relationship between  $u$  and  $M_b$  in Eq. (1), the relative increment adjustment (6.1 %) and corresponding change (4.8 %) to the predicted cumulative basal ablation over the time period that basal ablation was derived with observations is less than that associated with  $\Delta T$  or  $C$ .

#### 4.3 Thinning (spatial assessment)

The data collected at the main site on PII-A-1-f over 11 months provide unprecedented information regarding the temporal thinning of an ice island. However, these data are representative of a single point on a large ice island. Considering that the rate of surface ablation along the mIPR transect would have been near zero until the snow had melted around 15 July 2016, as determined from the ablation stakes at the sIPR site, the surface would have ablated at a rate of 1.5 cm d<sup>-1</sup> after this date. This is slightly less than the rate at the sIPR site over ablation period 3 (1.6 cm d<sup>-1</sup>; Table 1) and increases the confidence that the surface ablation conditions at the sIPR site were similar to those across the mIPR transect.

The ice island thickness over the mIPR transect ranged from 80.7 to 127.1 [-0.5, +0.8] m in May and 77.3 to 123.7 ± 0.5 m in September 2016 (Fig. 5a). A 50 m long section of relatively thin ice that was approximately 80 m thick was recorded along transect segment AB during both field visits (Fig. 5a); this section was approximately 40 m thinner than adjacent areas. The air and basal wave locations in the mIPR data could not be definitively selected when this thin section was passed during transect segment CD (Fig. 5a). This made it impossible to ascertain thickness at this location; however, a pair of in-line hyperbolae in both the May and September 2016 radargrams support the interpretation that this thinner section ran across both transect segments AB and CD. This section of thin ice is referred to as a ‘subsurface feature’ and corresponds to the location of the linear ‘surface feature’ described in Sect. 2. Together these are referred to as the ‘paired feature’.

The magnitude of thinning (3 to 4 m) observed over this thin section ~~observed-present~~ along transect segment AB (shown in the grey boxes in Fig. 5) was in the low to mid-range of the thinning observed along the entire transect. However, the general gradient in thinning magnitudes leading towards the paired feature shows a similar pattern in both transect segments AB and CD (shown in the pink boxes in Fig. 5). Thinning magnitudes of 3 to 4 m increased to 5 to 6 m across a distance of approximately 90 to 150 m, ending at the surface feature and where a gap in the thickness record exists. A fracture that occurred along this paired feature in September 2017 caused the areal extent of the ice island to reduce by approximately 2.7 ± 0.1 km<sup>2</sup>.

#### 4.4 Volume/area loss and contributions to deterioration

The area and volume of PII-A-1-f ~~was~~ ere 13.2 ± 0.01 km<sup>2</sup> and 1.4 ± 0.01 km<sup>3</sup>, respectively, when it was first visited in October 2015. By September 2016, the volume and areal extent decreased by 0.4 ± 0.01 km<sup>3</sup> and 3.4 ± 0.1 km<sup>2</sup>, respectively. A fracture event in September 2016 caused approximately 94 % of the areal reduction and 88 % of this volume loss. These

745 values represent the maximum possible reductions caused by the fracture. Other areal reduction processes (e.g., small-scale calving) would also have contributed to areal reduction during the time interval between RADARSAT-2 acquisitions over which the fracture occurred.

The 23 month RADARSAT-2 monitoring period (October 2015 to September 2017) captured a longer period of areal change. During the latter half of this time span (i.e., September 2016 and September 2017), the ice island twice un-grounded  
750 and re-grounded in the same vicinity and the large September 2017 fracture event occurred (Fig. 1e). The ice island volume decreased by  $0.6 \pm 0.01 \text{ km}^3$  over this period, with the vast majority of this volume loss (94 %) being caused by processes that decreased the areal extent of the ice island. The remaining volume was lost to basal and surface ablation, with basal ablation causing three times more volume loss than surface ablation. These ablation magnitudes were extrapolated from the on-ice data collection period. Over the entire two years that PII-A-1-f was monitored via RADARSAT-2 image acquisition,  
755 areal reduction processes reduced the volume of the ice island by approximately 67 %. Combined, surface and basal ablation resulted in an approximate 7 % reduction the ice island volume, with basal ablation again causing three times more loss than surface ablation.

## 5 Discussion

### 5.1 Basal ablation model calibrations

760 Two approaches are predominately used to model ice island and iceberg basal ablation (Bouhier et al., 2018). The first is the semi-empirical fluid-dynamics approach that is calibrated in this study; Eq. (1) approximates melt resulting from the bulk energy transfer occurring within the complex boundary conditions that are present at the ice-water interface (Weeks ~~et al., and Campbell~~ 1973; White et al., 1980). This model has been widely utilized for iceberg and ice island ~~deterioration-basal ablation~~ modelling in both the Arctic (~~Ballicater Consulting, 2012; Bigg et al., 1997; -Ballicater Consulting, 2012;~~  
765 ~~Keghouche et al., 2010; Wagner and Eisenman, 2017)~~ and Antarctic (Gladstone et al., 2001; Martin and Adcroft, 2010; Merino et al., 2016). The second approach is based on thermodynamic principles (Bouhier et al., 2018). Holland and Jenkins (1999) and Hellmer and Obers (1989) document a more complex three-equation thermodynamic model for ice shelf basal ablation, which represents both the salt and temperature flux across the ice-ocean interface.

The forced convection basal ablation model is advantageous due to its computational simplicity and the direct  
770 incorporation of  $\Delta u$ . In the thermodynamic approach, the values assigned to the turbulent heat and salt exchange parameters must be adjusted for varying values of  $\Delta u$  in iceberg and ice island applications (Jansen et al. 2007). Jansen et al. (2007) provide calibrated values for three stages of drift, each with a unique range of  $\Delta u$  values for an Antarctic ice island. ~~-W~~while Eq. (1) in the forced convection model directly incorporates  $\Delta u$ , ~~-model skill would likely improve if Eq. (2) was calibrated for drifting vs. grounded ice islands.~~ Equation (2) was developed to account for the plume of relatively cold and fresh  
775 iceberg meltwater that has been observed to surround icebergs and inhibit further melt (Foldvik et al., 1980). This meltwater plume will be stripped from the keel as  $\Delta u$  increases, as is the case when an ice island is grounded (Jansen et al., 2007).

Therefore, different parameterizations of Eq. (2) are likely required for predicting the basal ablation of drifting versus grounded ice islands, which FitzMaurice et al. (2017) showed to be the case when parameterizing Eq. (1) for the sidewall melt of an iceberg with different scenarios of meltwater plume and relative ocean velocities. It is possible that an the adjustment to the melting point of ice ( $M_p$ ) to account for the influence of the meltwater plume is not necessary under some conditions and  $M_p$  will simply equal the far field ocean freezing temperature ~~temperature~~ ( $\Theta_f$ ). Determining this will require concerted study of the difference in the basal boundary layer conditions of grounded versus drifting ice islands. The  $\Delta u$  for a drifting iceberg or ice island will be influenced by both ocean and wind currents (Kubat et al., 2005; Lichey and Hellmer, 2001). Observations of  $\Delta u$  for the drifting ice island case are rare but would be useful for this work and for correctly assigning values to this variable in Eq. (1).

This study used a unique dataset of observed ice island thinning, as well as in situ and modelled oceanographic conditions to present a calibration of the forced convection basal ablation model. It is noted that the exponents within Eq. (1), which were derived for the flat plate nature of a tabular iceberg or ice island, could also be calibrated. However, these have remained constant in previous iceberg and ice island literature while the bulk heat transfer coefficient,  $C$ , has typically been assigned one of two values that differ by an order of magnitude. Bigg et al. (1997), Bouhier et al. (2018), Martin and Adcroft (2010), Wagner and Eisenman (2017), and Weeks and Campbell (1973) assign a value of  $6.74 \times 10^{-6} \text{ m}^{2/5} \text{ s}^{-1/5} \text{ }^\circ\text{C}^{-1}$ . However, the confidence interval of the normal distribution of our calibrated  $C$  values does not overlap with this theoretically-derived value of the bulk heat transfer coefficient. We recommend assigning the  $C$  parameter a value of  $1.2 \times 10^{-5} \text{ m}^{2/5} \text{ s}^{-1/5} \text{ }^\circ\text{C}^{-1}$  when modelling ice island basal ablation in the future. This value was calculated from the calibration conducted over the full time span that basal ablation data was-were available and not the individual calibration intervals associated with the measured changes in ice thickness. In addition, the confidence interval of the normal distribution of  $C_i$  values found in Table 2 does overlap with this value. The value also matches, remarkably, that which was theoretically derived by White et al. (1980) and was assigned in the ice island deterioration model used by the CIS (Ballicater Consulting, 2012). We believe that it is more appropriate to use this value instead of the mean of the individually calibrated  $C_i$  values reported in Table 2 ( $1.7 \times 10^{-5} \text{ m}^{2/5} \text{ s}^{-1/5} \text{ }^\circ\text{C}^{-1}$ ), as we cannot be sure of the alignment between changes in oceanographic conditions, basal melt rates, and  $C_i$  values due to the low resolution of data collected by the sIPR. It is noted that Crawford et al. (2018b) assigned a value of  $1.3 \times 10^{-5} \text{ m}^{2/5} \text{ s}^{-1/5} \text{ }^\circ\text{C}^{-1}$  to  $C$ , supplied by Crawford (2018), when quantifying the distribution of freshwater input from ice island melt through the eastern Canadian Arctic. The use of this previous value would cause a very minimal skew in the distribution of freshwater input, slightly overestimating the freshwater input at the higher latitudes in their study region.

The calibration of the bulk heat transfer coefficient ( $C$ ) is specific to the approach used to assign an ice temperature when deriving the driving temperature,  $\Delta T$ . This paper follows the approach of Weeks and Campbell (1973), White et al. (1980), Løset (1993a), Kubat et al. (2005) and Ballicater Consulting (2012), empirically estimating the ice interface temperature (Eq. 2) based on work by Josberger (1977). This contrasts with a line of successive papers (e.g. Bigg et al., 1997; Martin and Adcroft, 2010; FitzMaurice et al., 2016; Wagner and Eisenman, 2017) that assign a constant ice temperature ( $T_{\text{ice}}$ ) of  $-4 \text{ }^\circ\text{C}$  based on Løset (1993b) This study reported -temperature measurements at ~50 cm depth of several iceberg sails in the

Barents Sea and show that this temperature will be reached within a few meters of the ice-ocean interface using a model (Løset 1993b). Yet another paper assigns  $T_{ice} = -15\text{ }^{\circ}\text{C}$  for sidewall melt calculations (FitzMaurice et al., 2017). These discrepancies underscore the need for harmonization on how  $T_{ice}$  is derived or assigned in the context of the forced convection basal ablation model. We recommend that the melting point of ice be assigned to  $T_{ice}$  in future studies that use this model, as it is the actual temperature of the ice-water interface, easy to calculate, and physically meaningful. Our recommendation is also supported by the model comparison and sensitivity study of FitzMaurice and Stern (2018); this work found that the thermodynamic and forced convection models agree when  $T_{ice} = M_p$  and infer that the heat flux into the iceberg interior is small. For the sake of inter-comparison, example calculations using  $L$ ,  $S_A$ ,  $\Theta$ , and  $u$  values representative of conditions at PII-A-1-f, the calibrated value of  $C$  would decrease by one order of magnitude if  $T_{ice} = -4\text{ }^{\circ}\text{C}$ . The calibrated  $C$  value would then be in-line with that derived by Weeks and Campbell (1973). The value of  $C$  would decrease by two orders of magnitude if  $-15\text{ }^{\circ}\text{C}$  was assigned to  $T_{ice}$ .

The in situ dataset of oceanographic conditions in the vicinity of PII-A-1-f and the GreenEdge sea ice camp was of paramount importance for validating and correcting the CMEMS data for bias. However, the calibration could be further improved by obtaining a longer time series of oceanographic data in closer proximity to the ice island. In general, there is a paucity of in situ oceanographic data collected in the vicinity of ice islands. Collection of such data will allow for further, improved drift and deterioration analyses of icebergs and ice islands. Modifying the sIPR to resolve smaller magnitudes of thickness change is also recommended. This would make it possible to relate these higher-quality thinning measurements to corresponding surface ablation and recorded oceanographic conditions.

## 5.2 Ablation rates and contributions to overall deterioration

Field measurements of ice island thinning are extremely sparse. Scambos et al. (2008) installed numerous instruments on two Antarctic tabular icebergs while investigating their deterioration processes with field and remote sensing data. Unfortunately, it was impossible to process thickness measurements collected by a radio-echo sounder installed on one of the icebergs (Scambos et al. 2008). Prior to this-our study, the spot values reported by Halliday et al. (2012) from a  $17\text{ km}^2$  drifting in the Labrador Sea were the only known field observations of ice island thinning. The PII-A-1-f thinning dataset greatly improves on this previous work, as the long-term sIPR time-series and repeat mIPR transects together produce a comprehensive dataset that allows us to assess the spatial and temporal variations in thinning and ablation. It would be highly beneficial to re-deploy the sIPR on a drifting ice island in the future to 1) augment the number of observations of ice island thinning and 2) begin comparing basal ablation occurring to drifting vs. grounded ice islands.

The average  $M_b$  of  $3.4\text{ cm d}^{-1}$  reported by Halliday et al. (2012) was greater than that observed for PII-A-1-f. Since the basal ablation rate for a drifting ice island should be lower than that of a grounded ice island due to decreased  $\Delta u$  and the potential protection of a meltwater plume, this greater melt rate was likely due to the higher  $\Theta$  off the coast of Labrador. Elevated ocean temperatures also contributed to the high,  $13.5\text{ m month}^{-1}$  basal ablation rate that was-Jansen et al (2007) estimated for the grounded Antarctic ice island, “A38-B”. Basal ablation was reported to cause 96 % of the thinning of this



ice island that had a surface extent of approximately 7600 km<sup>2</sup> (Jansen et al., 2007), whereas 73 % of the thinning of PII-A-1-f was a result of this process. -We note that, when present, the densification of firn is a factor in the surface ablation of Antarctic tabular icebergs. Scambos et al. (2008) document firn characteristics on two Antarctic tabular icebergs, and a follow-up field study that combines such observations with repeated thickness measurements would be valuable for assessing the contributions of surface and basal processes to Antarctic iceberg thinning.

While basal ablation was responsible for the majority of the thinning of PII-A-1-f, this process caused the ice island volume to reduce by approximately 5 %. However, basal ablation indirectly influences further deterioration by reducing the relative thickness of the ice island and decreasing fracture resistance (Goodman et al., 1980; Jansen et al., 2005). This will happen more quickly if an ice island is grounded, as basal ablation could increase by a factor of three after an ice island stops ~~freely~~-drifting freely (Jansen et al., 2007). Future research, including finite element modelling (e.g., Sazidy et al. ~~submitted~~forthcoming), is warranted to assess if and how thinning contributed to the September 2017 calving event. This fracture occurred along the paired feature that was substantially thinner than adjacent ice surfaces and which emanated from the root of the sidewall notch. We also recommend that further research be conducted into the relationship between the presence of the paired feature, the propagation of the sidewall notches before and during the time that PII-A-1-f was grounded in southern Baffin Bay, and the ultimate September 2017 fracture. The enlargement of sidewall notches, similarly to cusped deterioration patterns observed by Ballicater Consulting (2012), is a recurring deterioration mechanism that is not considered in deterioration studies or models at this point in time.

## 6 Conclusion

This study focuses on the thinning of PII-A-1-f, an ice island that originated from a calving event at the Petermann Glacier in 2012 and was grounded in western Baffin Bay over a 2-year monitoring time span. A unique field dataset was collected over four visits between October 2015 and September 2016 and was used to report ice island thinning and ablation rates and calibrate the popular forced convection basal ablation model.

The time series of ice island thinning recorded with a customized sIPR showed the ice island thinned by 4.7 m over the 11 months that on-ice data were collected. Basal ablation was responsible for 73 % of the observed thinning. ~~and caused the volume of PII A 1 f to decrease by an estimated 5 % over the duration that the ice island was remotely monitored via RADARSAT 2. This was a relatively small change in comparison to the 67 % volume reduction that resulted from processes that contributed to reducing the areal extent of the ice island. However,~~ Overall, PII-A-1-f was likely more susceptible to fracture than if it had been freely drifting, due to enhanced basal ablation and a correspondingly faster reduction in relative thickness.

It is important to model ice island basal ablation accurately for predicting the impact of meltwater input on the ocean system (Jansen et al., 2007). Additionally, basal ablation will alter the relative thickness of an ice island, which will influence fracture likelihood (Goodman et al., 1980), drift patterns (Barker et al., 2004) and grounding locations (Sackinger



et al., 1991). This is the first study to calibrate the forced convection basal ablation model for ice island or iceberg use with field data of ice island thinning, which removed uncertainty regarding estimated ablation rates from remotely sensed datasets. The calibrated value of the bulk heat transfer coefficient ( $1.2 \times 10^{-5} \text{ m}^{2/5} \text{ s}^{-1/5} \text{ }^{\circ}\text{C}^{-1}$ ) is in-line with the larger of two values assigned in previous iceberg and ice island basal ablation models, and we recommend that this value, specific to our approach of deriving a driving temperature, be used in future modelling endeavours. It is important to conduct such field studies to develop and validate methods for modelling ice island thickness change (i.e., surface and basal ablation), as this will inform future deterioration investigations and improve ice island drift and deterioration forecasting in both of the polar regions (Barker et al., 2004). The calibration of the forced convection basal ablation model might also be used to generally predict when grounded ice islands might thin enough to drift free, assuming certain shoal bathymetry and ice island morphology. This may be especially useful along the eastern coast of Canada where shipping and offshore industry operates, and where ice island grounding is a common occurrence. Overall, the work presented in this study highlights the value of systematic ice island field data collection. This is necessary for deterioration analyses, connecting morphology and deterioration, and developing high-quality models for operational and research purposes.

*Author contribution* AJC, GC and DM were responsible for project design. LM designed and produced the stationary and mobile ice penetrating radars and assisted in data processing and interpretation. AJC carried out field work and data analyses. DD and MB provided oceanographic data associated with the GreenEdge sea ice camp. DM, GC, LD, LM and DD contributed to the preparation of the manuscript, which was led by AJC.

*Competing interests* The authors declare that they have no conflict of interest.

*Data availability* Amundsen CTD data is available from the Polar Data Catalogue (PDC) under the file identifier 12713. Data collected at PII-A-1-f by the authors (i.e., data associated with ablation stakes, the weather station and sonic ranger, sIPR and mIPR) is also available from PDC (CCIN 13091), as is the RADARSAT-2 imagery (geotiffs) and digitized polygons of the areal extent of PII-A-1-f. CMEMS products are available at <http://marine.copernicus.eu/>. Data associated with the GreenEdge project are available, either through download or correspondence with a project coordinator, at <http://www.greenedgeproject.info/>.

*Acknowledgements* We would like to acknowledge the field assistance of Jaypootie Moesesie, Graham Clark, Jill Rajewicz, Jonathan Gagnon, Luke Copland, Abigail Dalton, Lauren Candlish, Hugo Jacques, Tom Desmeules, Natalie Theriault, Alison Cook, Alexis Burt, Leah Braithwaite, Julie Payette, Claire Bernard-Grand'Maison, Eric Brossier and Christian Haas. Adam Garbo and Iain Burnett helped with field preparation. This unique dataset would not be possible to collect without the access provided through the collaboration of ArcticNet and the CCGS *Amundsen*. We thank Canadian Coast Guard (CCG) captains

905 Alain Lacerte and Alain Gariépy as well as Transport Canada pilots Olivier Talbot, Alain Roy and Guillaume Carpentier for providing safe transport to our field site. We also greatly thank the entire CCG crew, who were instrumental in our field success. The hard work of ArcticNet staff, including Keith Lévesque, Phillipe Archambault, Louis Fortier, Jean-Éric Tremblay, Tim Papakyriakou, Leah Braithwaite, Annisa Merzouk and Colleen Gombault deserves great recognition. We thank Doug King, Geography and Environmental Studies, Carleton University and Douglas MacAyeal, University of  
910 Chicago for reviewing an earlier version of the manuscript.

Instrument development and field work were funded by the Northern Transportation Adaptation Initiative of Transport Canada, the Polar Knowledge Canada Safe Passage project (#1516 – 065) and Polar Knowledge Canada’s Northern Scientific Training Program. A.J. Crawford received personal funding from the Garfield Weston Foundation, the Natural Sciences and Engineering Research Council (Canada) and Environment and Climate Change Canada. This study has been  
915 conducted using E.U. Copernicus Marine Service Information. RADARSAT-2 scenes were acquired through a joint partnership agreement between Water and Ice Research Lab and the Canadian Ice Service, Environment and Climate Change Canada. RADARSAT data and products are © MacDonald, Dettwiler and Associates Ltd. (2010-2017), all rights reserved. RADARSAT is an official mark of the Canadian Space Agency. Some of the data presented herein were collected by the Canadian research icebreaker CCGS *Amundsen* and made available by the Amundsen Science program, which was  
920 supported by the Canada Foundation for Innovation and Natural Sciences and Engineering Research Council of Canada. The views expressed in this publication do not necessarily represent the views of Amundsen Science or that of its partners. Some of the data presented in this analysis were collected by the GreenEdge project. The GreenEdge project is funded by the following French and Canadian programs and agencies: ANR (Contract #111112), CNES (project #131425), IPEV (project #11164), CSA, Fondation Total, ArcticNet, LEFE and the French Arctic Initiative (GreenEdge project). The GreenEdge  
925 project would not have been possible without the support of the Hamlet of Qikiqtarjuaq and the members of the community as well as the Inuksuit School and its Principal, Jacqueline Arsenault. The GreenEdge project is conducted under the scientific coordination of the Canada Excellence Research Chair on Remote sensing of Canada's new Arctic frontier and the CNRS & Université Laval Takuvik Joint International laboratory (UMI3376). The GreenEdge field campaign was successful thanks to the contribution of J. Ferland, G. Bécu, C. Marec, J. Lagunas, F. Bruyant, J. Larivière, E. Rehm, S. Lambert-  
930 Girard, C. Aubry, C. Lalande, A. LeBaron, C. Marty, J. Sansoulet, D. Christiansen-Stowe, A. Wells, M. Benoît-Gagné, E. Devred and M.-H. Forget from the Takuvik laboratory, C.J. Mundy and V. Galindo from University of Manitoba as well as F. Pinczon du Sel and E. Brossier from Vagabond. The GreenEdge project also thanks Michel Gosselin, Québec-Océan, the CCGS *Amundsen* and the Polar Continental Shelf Program for their in-kind contribution in polar logistic and scientific equipment.

935

## References

- Abramoff, M.D., Magalhaes, P.J., and Ram, S.J.: Image processing with ImageJ, *Biophotonics Int.*, 11, 36–42, 2004.
- Amundsen Science Data Collection: CTD data collected by the CCGS Amundsen in the Canadian Arctic. Processed data. Version 3, doi:AN/ccin12713. Archived at [www.polardata.ca](http://www.polardata.ca). Accessed on 18 October 2016.
- 940 Ballicater Consulting: Ice Island and Iceberg Studies 2012, Canadian Ice Service, Environment Can., Ottawa, Ont., Contract Report, 73 pp., 2012.
- ~~Bamber, J., van den Broeke, M., Ettema, J., Lenaerts, J. and Rignot, E.: Recent large increases in freshwater fluxes from Greenland into the North Atlantic, *Geophys. Res. Lett.*, 39(19), doi:10.1029/2012GL052552, 2012.~~
- Barker, A., Sayed, M., and Carrieres, T.: Determination of iceberg draft, mass and cross-sectional areas, in: Proceedings of the 14th International Offshore and Polar Engineering Conference, Toulon, France, 23–28 May 2004, 1–6, 2004.
- 945 Bigg, G. R., Wadley, M. R., Stevens, D. P., and Johnson, J. A.: Modelling the dynamics and thermodynamics of icebergs, *Cold Reg. Sci. Tech.*, 26, 113–135, 1997.
- Bouhier, N., Tournadre, J., Rémy, F., and Gourves-Cousin, R.: Melting and fragmentation laws from the evolution of two large Southern Ocean icebergs estimated from satellite data, *Cryosphere*, 12, 2267–2285, doi: 10.5194/tc-12-2267-2018,
- 950 2018.
- Crawford, A.J.: Ice island deterioration in the Canadian Arctic: Rates, patterns and model evaluation, MSc thesis, Department of Geography and Environmental Studies, Carleton University, Ottawa, Ont, Canada. 140 pp., 2013.
- Crawford, A.J.: Ice island deterioration, PhD dissertation, Department of Geography and Environmental Studies, Carleton University, Ottawa, Ont, Canada. 205 pp., 2018.
- 955 Crawford, A., Crocker, G., Mueller, D., Desjardins, L., Saper, R., and Carrieres, T.: The Canadian Ice Island Drift, Deterioration and Detection (CI2D3) Database, *J. Glaciol.*, 1–5, doi:10.1017/jog.2018.36, 2018a.
- Crawford, A., Mueller, D., Desjardins, L., and Meyers, P.: The aftermath of Petermann Glacier calving events (2008–2012): Ice island size distributions and meltwater dispersal, *J. Geophys. Res.*, 123, 8812–8827, doi:10.1029/2018JC014388, 2018b.
- 960 Crawford, A., Mueller, D., and Joyal, G.: Surveying drifting icebergs and ice islands: Deterioration detection and mass estimation with aerial photogrammetry and laser scanning, *Remote Sens.*, 10(4), 575, doi:10.3390/rs10040575, 2018c.
- Dowdeswell, J.A., and Bamber, J.L: Keel depths of modern Antarctic icebergs and implications for sea-floor scouring in the geological record, *Mar. Geol.*, 243, 120–130, doi:10.1016/j.margeo.2007.04.008, 2007.
- ~~Enderlin, E. M., Howat, I. M., Jeong, S., Noh, M. J., van Angelen, J. H., and van den Broeke, M. R.: An improved mass budget for the Greenland ice sheet, *Geophys. Res. Lett.*, 41(3), 866–872, doi:10.1002/2013GL059010, 2014.~~
- 965 ~~FitzMaurice, A., Cenedese, C. and Straneo, F.: Nonlinear response of iceberg side melting to ocean currents, *Geophys. Res. Lett.*, 44(11), 5637–5644, doi:10.1002/2017GL073585, 2017.~~

- FitzMaurice, A. and Stern, A.: Parameterizing the basal melt of tabular icebergs, *Ocean Modelling*, 130, 66–78, doi:10.1016/j.ocemod.2018.08.005, 2018.
- 970 [FitzMaurice, A., Straneo, F., Cenedese, C. and Andres, M.: Effect of a sheared flow on iceberg motion and melting, \*Geophysical Research Letters\*, 43\(24\), 12,520–12,527, doi:10.1002/2016GL071602, 2016.](#)
- Foldvik, A., Gammelsrød, T., and Gjessing, Y.: Measurements of the radiation temperature of Antarctic icebergs and the surrounding surface water, *Annals Glaciol.*, 1, 19–22, doi:10.3189/S026030550001689X, 1980.
- Fraser, A.D., Massom, R.A., Michael, K.J., Galton-Fenzi B.K. and Lieser J.L. East Antarctic landfast sea ice distribution and  
975 variability, 2000–2008. *J. Climate*, 25, 1137–1156. doi:10.1175/JCLI-D-10-05032.1, 2012.
- Fuglem, M. and Jordaan, I.: Risk analysis and hazards of ice islands, in *Arctic Ice Shelves and Ice Islands*, edited by: Copland, L., and Mueller, D., Springer Netherlands, Dordrecht., 395–415, 2017.
- Gladstone, R. M., Bigg, G. R., and Nicholls, K. W.: Iceberg trajectory modeling and meltwater injection in the Southern Ocean, *J. Geophys. Res.*, 106(C9), 19903–19915, 2001.
- 980 Goodman, D. J., Wadhams, P., and Squire, V. A.: The flexural response of a tabular ice island to ocean swell, *Annals Glaciol.*, 1, 23–27, 1980.
- Haas, C., and Druckenmiller, M.: Ice thickness and roughness measurements, in: *Field Techniques for Sea Ice Research*, edited by Eicken, H., Gradinger, R., Salganek, M., Shirasawa, K., Perovich, D., and Leppäranta, M., University of Alaska Press, Fairbanks, Alaska, USA, 49–117, 2009.
- 985 Halliday, E.J., King, T., Bobby, P., Copland, L., and Mueller, D.R.: Petermann Ice Island ‘A’ survey results, offshore Labrador, in: *Proceedings of the Arctic Technology Conference*, Houston, Texas, USA, 3–5 December 2012, OTC 23714, 2012.
- Hellmer, H. H., and Olbers, D. J.: A two-dimensional model for the thermohaline circulation under an ice shelf, *Antarctic Sci.*, 1(04), doi:10.1017/S0954102089000490, 1989.
- 990 [Higgins, A. K.: North Greenland ice islands, \*Polar Rec.\*, 25\(154\), 207–212, 1989.](#)
- Holland, D. M., and Jenkins, A.: Modeling thermodynamic ice-ocean interactions at the base of an ice shelf, *J. Phys. Oceanogr.*, 29(8), 1787–1800, 1999.
- IOC, SCOR, and IAPSO, The international thermodynamic equation of seawater – 2010: Calculation and use of thermodynamic properties, *Manuals and Guides No. 56*, Intergovernmental Oceanographic Commission, UNESCO, Paris, France, 196 pp., 2010.
- 995 Jansen, D., Sandhager, H., and Rack, W.: Model experiments on large tabular iceberg evolution: ablation and strain thinning, *J. Glaciol.*, 51(174), 1–11, 2005.
- Jansen, D., Schodlok, M., and Rack, W.: Basal melting of A-38B: A physical model constrained by satellite observations, *Remote Sens. Environ.*, 111(2–3), 195–203, doi:10.1016/j.rse.2007.03.022, 2007.

- 1000 [Josberger, E.G.:1977, A laboratory and field study of iceberg deterioration, in: In: A.A. Hussein \(Editor\), Proceedings of the First International Conference on Iceberg Utilization for Fresh Water Production, Weather Modification and other Applications, edited by: Hussein, A.A., -IA. Pergamon Press, New York, pp. 245-264, 1977.](#)
- Keghouche, I., Counillon, F. and Bertino, L.: Modeling dynamics and thermodynamics of icebergs in the Barents Sea from 1987 to 2005, J. Geophys. Res., 115(C12), doi:10.1029/2010JC006165, 2010.
- 1005 Kelly, D.: gsw: Gibbs Sea Water Functions (v. 1.0-5), R package. 2017.
- [Kubat, I., Sayed, M., Savage, S. B. and Carrieres, T.: An operational model of iceberg drift, Int. J. Offshore Polar Eng., 15\(2\), 125–131, 2005.](#)
- Kubat, I., Sayed, M., Savage, S.B., Carrieres, T., and Crocker, G.: An operational iceberg deterioration model, in: Proceedings of the 17th International Offshore and Polar Engineering Conference, International Society of Offshore and
- 1010 Polar Engineers, Lisbon, Portugal, 1-6 June 2007, 652–657, 2007.
- Leucci, G., Negri, S., and Carrozzo, M. T.: Ground Penetrating Radar (GPR): an application for evaluating the state of maintenance of the building coating, Annals Geophys., 46(3), 481-489, 2003.
- [Lichey, C., and Hellmer, H. H.: Modeling giant-iceberg drift under the influence of sea ice in the Weddell Sea, Antarctica, J. Glaciol., 47\(158\), 452–460, 2001.](#)
- 1015 Løset, S.: Numerical modelling of the temperature distribution in tabular icebergs, Cold Reg. Sci. Tech., 21(2), 103–115, 1993a.
- [Løset, S.: Thermal energy conservation in icebergs and tracking by temperature, J. Geophys. Res., 98\(C6\), 10001, doi:10.1029/93JC00138, 1993b.](#)
- Macheret, Y. Y., Moskalevsky, M. Y., and Vasilenko, E. V.: Velocity of radio waves in glaciers as an indicator of their
- 1020 hydrothermal state, structure and regime, J. Glaciol., 39(132), 373–384, 1993.
- Martin, T., and Adcroft, A.: Parameterizing the fresh-water flux from land ice to ocean with interactive icebergs in a coupled climate model, Ocean Model., 34(3–4), 111–124, doi:10.1016/j.ocemod.2010.05.001, 2010.
- Massom, R. A., Hill, K.L., Lytle, V.I., Worby, A.P., Paget, M.J., and Allison, I.: Effects of regional fast-ice and iceberg distributions on the behaviour of the Mertz Glacier polynya, East Antarctica. Ann. Glaciol., 33, 391–398,
- 1025 doi:10.3189/172756401781818518, 2001.
- Merino, N., Le Sommer, J., Durand, G., Jourdain, N. C., Madec, G., Mathiot, P., and Tournadre, J.: Antarctic icebergs melt over the Southern Ocean: Climatology and impact on sea ice, Ocean Model., 104, 99–110, doi:10.1016/j.ocemod.2016.05.001, 2016.
- Mingo, L., and Flowers, G.: An integrated lightweight ice-penetrating radar system, J. Glaciol., 56(198), 709–714, 2010.
- 1030 Mingo, L., Flowers, G., Crawford, A.J., ~~and~~ Mueller, D., [Bigelow, D.G.: A stationary iceimpulse-penetrating radar system for autonomous monitoring-deployment in cold and temperature environments of englacial and subglacial properties. Forthcoming. Revised manuscript submitted, 09/2019ForthcomingForthcoming.](#)

Natural Resources Canada: Tools and Applications | Natural Resources Canada. <http://www.nrcan.gc.ca/earth-sciences/geomatics/geodetic-reference-systems/tools-applications/10925#ppp>. last access: 2016.

- 1035 [Oziel, L., Massicotte, P., Randelhoff, A., Ferland, J., Vladioiu, A., Lacour, L., Galindo, V., Lambert-Girard, S., Dumont, D., Cuypers, Y., Bouruet-Aubertot, P., Mundy, C.-J., Ehn, J., Bécu, G., Marec, C., Forget, M.-H., Garcia, N., Coupel, P., Raimbault, P., Houssais, M.-N. and Babin, M.: Environmental factors influencing the seasonal dynamics of spring algal blooms in and beneath sea ice in western Baffin Bay, \*Elem Sci Anth\*, 7\(1\), 34, doi:10.1525/elementa.372, 2019.](#)
- 1040 Sackinger, W.M., Jeffries, M.O., Li, F., and Lu, M.: Ice island creation, drift, recurrences, mechanical properties, and interactions with Arctic offshore oil production structures, Final Report for the U.S. Department of Energy, University of Fairbanks, Fairbanks, Alaska, USA, 38 pp., 1991.
- Sazidy, M., Crocker, G., and Mueller, D.: A 3D numerical model of ice island calving due to buoyancy-driven flexure, in: Proceedings of the 25th International Conference on Port and Ocean Engineering under Arctic Conditions, Delft, The Netherlands, 9-13 June 2019, ~~Submitted~~[Forthcoming](#).
- 1045 [Scambos, T., Ross, R., Bauer, R., Yermolin, Y., Skvarca, P., Long, D., Bohlander, J. and Haran, T.: Calving and ice-shelf break-up processes investigated by proxy: Antarctic tabular iceberg evolution during northward drift, \*Journal of Glaciology\*, 54\(187\), 579–591, 2008.](#)
- 1050 Stern, A. A., Johnson, E., Holland, D. M., Wagner, T. J. W., Wadhams, P., Bates, R., Abrahamsen, E. P., Nicholls, K. W., Crawford, A., Gagnon, J., and Tremblay, J.-E.: Wind-driven upwelling around grounded tabular icebergs, *J. Geophys. Res.: Oceans*, 120(8), 5820–5835, doi:10.1002/2015JC010805, 2015.
- [USGS \(United States Geological Survey\): Earthshots: Satellite Images of Environmental Change; Petermann Glacier, Greenland: <https://earthshots.usgs.gov/earthshots/node/70#ad-image-5-1> last access: 10 November 2019.](#)
- Wagner, T. J. W., and Eisenman, I.: How climate model biases skew the distribution of iceberg meltwater, *Geophys. Res. Lett.*, 44(8), 3691–3699, doi:10.1002/2016GL071645, 2017.
- 1055 Weeks, W. F., and Campbell, W. J.: Icebergs as a fresh-water source: An appraisal, *J. Glaciol.*, 12(65), 207–233, 1973.
- White, F. M., Spaulding, M. L., and Gominho, L.: Theoretical estimates of the various mechanisms involved in iceberg deterioration in the open ocean, Technical Report, Office of Research and Development, United States Coast Guard, Washington, DC., 133 pp., 1980.
- Wilson, N.: Characterization and interpretation of polythermal structure in two subarctic glaciers, MSc thesis, University of British Columbia, Vancouver, BC., 241 pp., 2012.
- 1060 Wilson, N.: Radar Tools. Software program. available at: <http://doi.org/10.5281/zenodo.43972>, 2013.

**Table 1: Ablation and total thinning magnitudes and rates per ablation period.** Air temperature ( $T_a$ ) data for each period is also included. Inconsistencies between the summed ablation and thinning magnitudes or between the ratios of surface : basal ablation magnitudes versus rates are due to rounding.  $M_s$  and  $M_b$  represent daily surface and basal ablation rates, respectively.

Ablation period	Dates	Surface ablation (cm) & $M_s$ (cm d <sup>-1</sup> )	Basal ablation (cm) & $M_b$ (cm d <sup>-1</sup> )	Total thinning (cm) & <i>rate</i> (cm d <sup>-1</sup> )	Mean $T_a$ (°C)	Days $T_a > 0^{\circ}\text{C}$
1	2015-11-16: 201 <del>56</del> -12-04	2 ( <i>0.1</i> )	14 ( <i>0.7</i> )	16 ( <i>0.8</i> )	-13	3
2	201 <del>56</del> -12-05: 2016-07-14	0	200 ( <i>0.9</i> )	200 ( <i>0.9</i> )	-15	33
3	2016-07-15: 2016-09-14	106 ( <i>1.6</i> )	78 ( <i>1.2</i> )	184 ( <i>2.8</i> )	3	65
1-3	2015-11-16: 2016-09-14	108 ( <i>0.4</i> )	295 ( <i>1.0</i> )	403 ( <i>1.3</i> )	-11	101

**Table 2: Calibrated values for  $C_i$  for the individual calibration intervals associated with each measured change in ice thickness.**  
The corresponding mean driving temperature ( $\Delta T$ ) and velocity ( $u$ ) values, found with the CMEMS data, are also provided.


Calibration Interval	Date range	$C_i$ ( $\text{m}^{2/5} \text{s}^{-1/5} \text{ } ^\circ\text{C}^{-1}$ )	$\Delta T$ ( $^\circ\text{C}$ )	$u$ ( $\text{m s}^{-1}$ )
1	2015-11-15 : 2016-01-30	$7.8 \times 10^{-6}$	0.32	0.12
2	2016-01-30 : 2016-05-07	$8.3 \times 10^{-6}$	0.24	0.12
3	2016-05-07 : 2016-07-09	$2.6 \times 10^{-5}$	0.12	0.11
4	2016-07-09 : 2016-08-01	$2.1 \times 10^{-5}$	0.18	0.11
5	2016-08-01 : 2016-08-22	$2.1 \times 10^{-5}$	0.17	0.11
6	2016-08-22 : 2016-09-18	$1.6 \times 10^{-5}$	0.23	0.14




1090

**Table 3: Descriptive statistics and sensitivity analysis results.** The key driving variables in the forced convection basal ablation model were individually perturbed in 8 equal increments based on the variable's range. The sensitivity of the model was assessed by as the mean percent increase in predicted cumulative basal ablation ~~over the each time period that basal ablation was derived with~~following each incremental increase in the value assigned to a given variable. The sensitivity was analysed over the longer time period (2015-11-15 to 2016-09-18) due to the certainty in the bulk basal ablation.


	Range	Increment	Mean-increase in variable (%)	Mean increase in cumulative basal ablation (m)	Mean increase in cumulative basal ablation output (%)
$\Delta u$ (m s <sup>-1</sup> )	0.11 : 0.18	0.01	6.1	0.22	4.8
$\Delta T$ (°C)	0.10 : 0.37	0.03	18.6	0.59	18.6
$C$ (m <sup>2/5</sup> s <sup>-1/5</sup> °C <sup>-1</sup> )	$7.8 \times 10^{-6}$ : $2.6 \times 10^{-5}$	$2.3 \times 10^{-5}$	16.5	0.56	16.5




**Figure 1: Remote monitoring of PII-A-1-f and data collection locations.** (a) Dates and locations of “Petermann Ice Island (PII)-A-1-f” as observed with RADARSAT-2 synthetic aperture radar (SAR) observations while the ice island drifted from the Petermann Glacier (PG), through Nares Strait (NS), Kane Basin (KB) and Baffin Bay (BB) before grounding north of the Cumberland Peninsula (CP). The polygon in the top inset shows the location of the larger-scale map. The bottom inset shows the proximity of the grounding location to the GreenEdge sea ice camp. (b) RADARSAT-2 ScanSAR image (100 m nominal resolution) showing the surface feature and sidewall notches. (c) – (e) RADARSAT-2 Fine-Quad SAR (8 m nominal resolution) scenes acquired on 27 June 2016, 27 September 2016 and 23 September 2017, respectively. Conductivity, temperature, depth (CTD) cast locations are denoted by triangles in (c) and (d) and the star in (c) denotes the location of the stationary ice penetrating radar and weather station; the ~3 km mobile IPR transect conducted in May 2016 is shown as a black line. All RADARSAT-2 images are presented as colour composites (ScanSAR polarizations: red = HH, blue and green = HV. Fine-Quad polarizations: red = HH, green = VV, blue = HV).




**Figure 2: Instruments installed on Petermann Ice Island-A-1-f on 20 October 2015.** (a) Stationary ice penetrating radar (sIPR) components deployed with antennas ‘in-line’ and attached to eight sleeves that slid over stake anchors. (b) Automatic weather station components, including a camera that acquired a weekly image of the sIPR and ablation stakes (not shown). The two systems were installed 30 m apart on two small ridges at the location denoted in Fig. 1c. PV = photovoltaic panel, SR = sonic ranger.



**Figure 3: Ablation and thickness change.** (a) “Petermann ice island (PII)-A-1-f” thickness change as measured by the stationary ice penetrating radar (sIPR) (grey dots) and interpolated between thickness measurements (black line). The first thickness observation was not included in the interpolation due to uncertainty imposed by sIPR measurement settings. (b) Surface and basal ablation, as well as the change in surface height (ht) relative to the start of sIPR data collection. The latter, represented by the thick blue line, includes ice ablation (positive values) as well as snow accumulation (negative values). Basal ablation is calculated with the linearly interpolated thickness values in a) and should be taken as an estimate of daily magnitudes. The vertical lines denote three surface ablation periods that correspond with the periods described in Table 1. The timing of CTD profiling by the CCGS *Amundsen* and the time span associated with oceanographic data collection at the GreenEdge sea ice camp are also denoted.



**Figure 4: Oceanographic data collected or modelled over the time span that PII-A-1-f basal ablation rates were derived.** CMEMS product data for the grid cells in which “Petermann Ice island (PII)-A-1-f” (solid black lines) and the GE sea ice camp (solid green lines) were located: (a) conservative temperature ( $\Theta$ ), (b) absolute salinity ( $S_A$ ), and (c) velocity ( $u$ ). As PII-A-1-f was grounded,  $u$  is equivalent to the differential velocity between the ice island and the ocean current. Field data collected by the CCGS *Amundsen* and the GreenEdge project are shown in the respective panels. The CMEMS data, corrected for bias in  $S_A$  and  $u$ , are shown as dotted lines in (b) and (c), respectively. The vertical lines denote the calibration intervals, numbered in (b), associated with thickness change measurements that were used for the calibration of the forced convection basal melt model.



**Figure 5: Spatial variation in morphology and thinning between May and September 2016.** Upper-case letters are used to denote transect segments referred to in the text. (a) Thickness observations collected with mobile ice penetrating radar. The grey box corresponds with the thin ice section also denoted in (c). (b) Thickness change. The pink boxes indicate thickness change gradients also identified in (c). Thinning data were categorized based on the amount of overlap between the May and September radar footprints. Grey shading in (a) and (b) denotes uncertainty in thickness measurements and thinning amounts, respectively. The larger, single points denote the respective thickness and thinning measured with corresponding data collected by the sIPR at the main site. (c) Thickness change displayed on a Fine-Quad RADARSAT-2 synthetic aperture radar scene acquired on 28 September 2016 (shown in greyscale). A surface feature is apparent as a line of high backscatter (white) pixels starting at the sidewall notch. The geolocation error of the RADARSAT-2 scene was evaluated and minimal error in the position of the scene was observed.