

Author's Responses

January 2, 2020

Dear Dr. Nisancioglu,

Thank you again for facilitating the review process. Below are point by point responses to reviewer comments.

Sincerely,
Jacob Downs

Comment

cf Bahadory and Tarasov, GMD 2018

Response

Coupling an ice sheet model to an earth systems model as in Bahadory and Tarasov (2018) is undoubtedly more physically principled, and would capture feedbacks between the ice sheet and climate system that are neglected in the PDD model. We now note this point in the model limitations section. Given our emphasis on determining precipitation and surface mass balance via an inverse modeling approach, we believe that the computational cost of this approach outweigh the benefits.

Comment

you are assuming continuous frozen base then? If there is a transition between cold and warm based conditions, then your statement is no longer apriori valid.

Response

This is a good question. We do not assume a continuous frozen base. The ice is allowed to slide throughout the entire ice sheet domain, however sliding velocities are negligible in the ice sheet interior.

Comment

Given the attention to uncertainty quantification, there should be some consideration of the uncertainties introduced from the above. If you review there relevant litterature, there are other choices for treating these components. Just stating you chose something because somebody else used it is crappy science that ice sheet modellers have too often relied on.

Response

This is a good point. The lapse rate was included in past sensitivity experiments. However, we found that the sensitivity of the precipitation anomaly to the rate factor was low compared to other tested parameters. Therefore we excluded it in the most recent draft.

Comment

Need to explicitly state so in the text

Response

We mention this in the model limitations section.

Comment

not addressing the stated query

Response

We attempted to clarify. We define a Gaussian prior distribution from which we draw “candidate” retreat histories. The covariance matrix for this prior is selected so that the randomly drawn retreat histories from the prior are plausible in the sense of being correlated on short time scales. Implausible retreat histories are rejected in order to obtain a distribution of retreat histories that match the observational record.

Comment

not addressing point. You are assuming that mean and covariance fully characterize the distribution. This implies you are assuming an underlying multivariate normal distribution.

Response

We have edited the text to emphasize that the unscented transform is intended to estimate moments of a distribution, rather than fully characterize the distribution itself. Some versions of the UT, such as the fifth order method outlined in the manuscript, are not limited to the Gaussian assumption since they estimate higher order moments. However, we largely agree with this point, since we do not explicitly use or report information about higher order moments.

Comment

incorrect. If you do the math with the observed mean global temperature (278K) and the simplification of Clausius-Clapeyron using the ideal gas law, you get a 10% increase per K.

Response

According to Held and Soden (2006) at temperatures typical of the lower troposphere, the saturation vapor pressure increases by around 7% per 1 degree C increase in temperature.

Comment

I find this claim almost vacuous within the context of this study. You are explicitly assuming ice dynamical changes don't matter, ie your inference is focussed on inferring past precip changes with a 1D isothermal flowline model with time independent sliding. How does this relate to inferring future changes in the GrIS in response to future changes in climate when the largest uncertainties are local climate change, ice dynamical (ie surges/destabilization), and submarine melt and grounding line retreat where relevant.

Response

Our results add support to the idea that surface mass balance was the main driver of retreat in land terminating sectors of the GrIS. In these regions the feedback between increasing temperature and increased accumulation may be an especially important one in the future.

Comment

observed terminus positions in an isothermal flowline ice dynamics model

Response

Thanks, this has been changed.

Comment

Scientific writing needs to be precise and accurate. The above statement is not correct if emulators are used.

Response

We would argue that emulators are a separate topic from MCMC. The unscented transform and emulators are both strategies for reducing the cost of inference, and the two strategies are actually related, which we now highlight in the discussion.

Comment

I don't know how you can infer this from the cited studies. I'm assuming you are equating that the sensitivity to relevant tuning parameters is higher for SMB than basal sliding. But that is not the same as your claim. Those models did not assume an isothermal structure and therefore had basal temperature controlled switching of basal sliding.

I don't know how you can infer this from the cited studies. I'm assuming you are equating that the sensitivity to relevant tuning parameters is higher for SMB than basal sliding. But that is not the same as your claim. Those models did not assume an isothermal structure and therefore had basal temperature controlled switching of basal sliding.

Response

Cuzzone et al. (2019) show that in west central Greenland, simulated retreat is not significantly affected by mesh resolution, as surface mass balance is the primary driver of retreat. Lecavalier et al. (2014) state that the modeled margin position is determined almost entirely by surface mass balance rather than the flow law or basal sliding.

Comment

Need to be clear that you are introducing another source of uncertainty given that the GIA output from Caron et al, is unlikely to be self-consistent with your changing ice sheet load.

Response

We address this informally in the discussion. Estimated precipitation anomalies are similar whether we assume a static or dynamic bed using GIA output.

Comment

that is not the Clausius-Clapeyron relation

Response

Thanks, this was a poorly worded. It is an expression derived from the Clausius-Clapeyron relation. This has been fixed.

Comment

I see little point in bothering with a higher-order momentum balance if you are likely introducing larger uncertainties from the use of an isothermal model

Response

We agree. It is likely that the higher-order momentum balance is not terribly important here given the surface mass balance dominated retreat pattern, although we did not necessarily anticipate this result.

Comment

Seriously, you are doing Bayesian inversion with only 3 model runs? In this case, if this inversion is so fast, then the cost of running a full 3D ice sheet model should be irrelevant.

Response

We are doing 45 runs in parallel over three iterations, not 3 model runs total.

Comment

Should cite past applications thereof, eg Tarasov et al, EPSL, 2012, and pollard et al, GMD, 2016

Response

Thanks, we have added these citations in the discussion.

Comment

.. and/or changes ..

Response

Fixed, thank you.

References

Taimaz Bahadory and Lev Tarasov. LCice 1.0-a generalized Ice Sheet System Model coupler for LOVE-CLIM version 1.3: Description, sensitivities, and validation with the Glacial Systems Model (GSM version D2017.aug17). *Geoscientific Model Development*, 2018. ISSN 19919603. doi: 10.5194/gmd-11-3883-2018.

Joshua K. Cuzzone, Nicole Jeanne Schlegel, Mathieu Morlighem, Eric Larour, Jason P. Briner, Helene Seroussi, and Lambert Caron. The impact of model resolution on the simulated Holocene retreat of the southwestern Greenland ice sheet using the Ice Sheet System Model (ISSM). *Cryosphere*, 2019. ISSN 19940424. doi: 10.5194/tc-13-879-2019.

Isaac M. Held and Brian J. Soden. Robust responses of the hydrological cycle to global warming. *Journal of Climate*, 2006. ISSN 08948755. doi: 10.1175/JCLI3990.1.

Benoit S. Lecavalier, Glenn A. Milne, Matthew J.R. Simpson, Leanne Wake, Philippe Huybrechts, Lev Tarasov, Kristian K. Kjeldsen, Svend Funder, Antony J. Long, Sarah Woodroffe, Arthur S. Dyke, and Nicolaj K. Larsen. A model of Greenland ice sheet deglaciation constrained by observations of relative sea level and ice extent. *Quaternary Science Reviews*, 102:54–84, oct 2014. ISSN 02773791. doi: 10.1016/j.quascirev.2014.07.018.

West Greenland ice sheet retreat history reveals elevated precipitation during the Holocene thermal maximum

Jacob Downs¹, Jesse Johnson², Jason Briner³, Nicolás Young⁴, Alia Lesnek³, and Josh Cuzzone⁵

¹Department of Mathematical Sciences, University of Montana, Missoula MT, USA

²Department of Computer Science, University of Montana, Missoula MT, USA

³Department of Geology, University at Buffalo, Buffalo NY, USA

⁴Lamont-Doherty Earth Observatory, Columbia University, Palisades NY, USA

⁵Department of Earth System Science, University of California Irvine, Irvine CA, USA

Correspondence: Jacob Downs (jacob.downs@umontana.edu)

Abstract. We investigate changing precipitation patterns in the Kangerlussuaq region of west central Greenland during the Holocene thermal maximum, using a new chronology of ice sheet terminus position through the Holocene and a novel inverse modeling approach based on the unscented transform (UT). The UT is applied to estimate changes in annual precipitation in order to reduce the misfit between modeled and observed terminus positions. We demonstrate the effectiveness of the UT for time-dependent data assimilation, highlighting its low computational cost and trivial parallel implementation. Our results indicate that Holocene warming coincided with elevated precipitation, without which modeled retreat in the Kangerlussuaq region is more rapid than suggested by observations. Less conclusive is if high temperatures during the HTM were specifically associated with a transient increase in precipitation, as the results depend on the assumed temperature history. Our results highlight the important role that changing precipitation patterns had in controlling ice sheet extent during the Holocene.

10 1 Introduction

During the early Holocene (~11.7 - 8 ka BP), terrestrial and marine climate proxies from the northern hemisphere reveal a warmer than present peak in temperature (Kaufman et al., 2004; Marcott et al., 2013). This period of elevated temperatures, likely initiated by greater than modern insolation, is referred to as the Holocene thermal maximum (HTM). Its onset, duration, and severity were likely spatially variable (Kaufman et al., 2004). Records of HTM warming can be found in Greenland ice core records. For example, temperatures measured in the Dye-3 borehole show a pronounced HTM signal occurring from 15 7 to 4 ka BP and having value of 2.5° above present temperatures (Dahl-Jensen et al., 1998; Miller et al., 2010), whereas at the GISP2 site, the HTM appears to occur slightly earlier, following the 8.2 ka BP cold event (Kobashi et al., 2017) (Figure 1).

While warming during the HTM is well established, less is known about the regional changes in precipitation that accompanied increased temperatures. Ice core records provide long term estimates of accumulation (Alley et al., 1993), but these point 20 measurements near ice divides are not representative of the precipitation across the ice sheet, particularly at lower elevations near the coast. Because the HTM was accompanied by lower Arctic sea ice extent (Polyak et al., 2010), it is possible that additional moisture was available to the GrIS from open Arctic waters. This is supported by proxy evidence showing an increase

in winter precipitation in western Greenland coincident with HTM warming (Thomas et al., 2016). However, temperature is known with greater certainty than precipitation.

Understanding feedbacks between temperature and precipitation during the HTM has implications for the future of the GrIS. Warming and declining sea ice are projected to cause an increase in Arctic precipitation (Bintanja and Selten, 2014; 5 Singarayer et al., 2006). On a global scale, the moisture content of the atmosphere increases by around 7% for every degree of warming, according to the Clausius-Clapeyron relation. On a regional scale, declining arctic sea ice is expected to cause changes in atmospheric circulation, bringing more moisture to the Arctic (Bintanja and Selten, 2014). While there are important differences between HTM and modern climate, the history of retreat in West Greenland may provide insights into how the GrIS will respond to a warmer and possibly wetter future climate.

10 Modeling studies indicate that Holocene retreat in land terminating regions of the GrIS were controlled primarily by surface mass balance rather than ice dynamics (Cuzzone et al., 2019; Lecavalier et al., 2014). Given the primary importance of surface mass balance in controlling modeled retreat, we explore the hypothesis that enhanced winter snowfall during the HTM may have slowed retreat by partially offsetting increased surface melt (Thomas et al., 2016). We investigate changes in precipitation in a land terminating sector of the western central GrIS, near Kangerlussuaq, taking advantage of a new chronology of 15 ice sheet terminus position (Young et al., 2019, In Review) and a novel inverse modeling approach based on the unscented transform (UT) (Julier and Uhlmann, 1997). In particular, we use the UT to estimate changes in annual precipitation during the Holocene by reducing the misfit between modeled and observed terminus positions in an [isothermal](#) flowline ice dynamics model (Brinkerhoff et al., 2017) (section 2.1).

20 The inverse problem is posed as a Bayesian inference problem, and its solution involves estimating a non-Gaussian posterior probability distribution. Markov Chain Monte Carlo (MCMC) methods, such as Metropolis Hastings method (Chib and Greenberg, 1995), provide one means of solving the inference problem by generating random samples from the posterior distribution. Generating samples from the posterior, however, requires repeatedly running the ice dynamics model with different precipitation histories as input, which is intractable even for a relatively computationally inexpensive flowline model.

25 The unscented transform provides a computationally efficient and trivially parallelizable alternative to MCMC methods. The basic idea of the unscented transform is to use a small, fixed number of deterministic sample points in order to estimate the statistical moments (e.g. mean and covariance) of the posterior distribution. Sigma points, each of which represents a different precipitation history input to the ice dynamics model, are generated a priori (section 2.5.3). Consequently, all model runs can be performed simultaneously in parallel resulting in at least a 100 fold speed up compared to MCMC methods.

2 Numerical methods for inference

30 2.1 Ice-sheet model

We use the 1D, [isothermal](#), flowline model with higher-order momentum balance described in Brinkerhoff et al. (2017). The momentum conservation equations are simplified using the Blatter-Pattyn approximation, assuming hydrostatic pressure and

negligible vertical resistive stresses (Blatter, 1995; Pattyn, 2003). Default parameter values used in this work are specified in Table 1.

We adopt a linear sliding law of the form

$$\tau_b = \beta^2 N u_b \quad (1)$$

5 where τ_b is basal shear stress, β^2 is a constant basal traction parameter, N is effective pressure, and u_b is sliding speed. Based on borehole water pressure measurements in Wright et al. (2016), basal water pressure P_w is assumed to be a fixed fraction $P_{frac} = 0.85$ of ice overburden pressure P_0 . Effective pressure is therefore given by

$$N = P_0 - P_{frac} P_0. \quad (2)$$

10 The basal traction parameter β^2 is tuned to minimize the misfit between modeled and observed surface velocities from Moug-
inot et al. (2017) for modern Isunnguata Sermia.

2.2 Flowline Selection and Moraine Age Constraints

To define the path followed by ice, we assume that flow follows the modern surface velocity field inland of the present day margin. In ice free regions, the direction of ice flow is inferred from bedrock topography (Figure 1). Since the direction of ice flow is unknown and time varying, we cannot directly quantify the uncertainty introduced by errors in flowline selection. To 15 account for some of this uncertainty, we perform inversions on two plausible, adjacent paleo-flowlines in the Kangerlussuaq area.

The rate of Holocene retreat on each flowline is estimated using constraints on ice sheet terminus position from (Young et al., 2019, In Review) (Figure 1). Terminus position data in Young et al. (2019, In Review) indicates that in the early Holocene (11.6 ka BP), the ice sheet margin was some tens of kilometers inland of the present day coastline. Although the moraine patterns 20 are spatially complex, generally speaking there was a period of moderate retreat (~ 10 km on the northern flowline and ~ 30 km on the southern flowline) from 11.6 to 10.3 ka BP, followed by rapid retreat (~ 100 km on both flowlines) from 10.3 to 8.1 ka BP. By 8.1 ka BP, the margin position was within 20 km of its present position on both flowlines (Figure 5). The modern terminus position provides one additional constraint. Moraine ages have uncertainties of up to ± 400 years.

For modern bedrock geometry along the flowlines, we use BedMachine v3 (Morlighem et al., 2017). Isostatic uplift and 25 relative sea level changes are accounted for using a Glacial isostatic adjustment model (Caron et al., 2018). This model, combined with the retreat chronology in Young et al. (2019, In Review), indicate that ice remained grounded on both the northern and southern flowlines from 11.6 ka BP onward.

2.3 Positive degree day model

Surface mass balance is estimated using a positive degree day (PDD) model (Johannesson et al., 1995). Annual surface mass 30 balance is constructed in the PDD model using estimates of average monthly precipitation and temperature. Inputs into the PDD model include the unknown ice surface elevation S , modern monthly temperature T_m and precipitation P_m along the

flowlines, as well as the seasonal temperature anomaly ΔT . As in Cuzzone et al. (2019), modern temperature and precipitation are computed as 30 year averages from 1980-2010 using Box (2013).

To assess the sensitivity of modeled retreat history to temperature, we perform experiments using temperature reconstructions from both Buizert et al. (2018) and Dahl-Jensen et al. (1998). For the spatially explicit Buizert et al. (2018) reconstruction, 5 monthly temperature anomalies are computed as averages along the flowlines. In contrast, Dahl-Jensen et al. (1998) reconstruct temperature only at the GRIP and Dye-3 borehole locations (Figure 1). Since a full Holocene reconstruction is unavailable at the Dye-3 borehole site, which is closer to Kangerlussuaq, we use the temperature reconstruction at GRIP. The HTM is roughly 0.3° C warmer and 1,500 years later in Dahl-Jensen et al. (1998) than Buizert et al. (2018) (Figure 6)

A limitation of the Dahl-Jensen et al. (1998) reconstruction is that it does not resolve seasonal temperatures. To address this, 10 we calculate the difference between monthly and mean annual temperatures in Buizert et al. (2018) and apply those offsets to the mean annual temperature at GRIP from Dahl-Jensen et al. (1998).

Surface temperature T is computed monthly as

$$T = T_m + \Delta T + \alpha(S - S_m) \quad (3)$$

where S_m is the modern surface elevation, and $\alpha = 5^\circ \text{ C km}^{-1}$ is the lapse rate (Abe-Ouchi et al., 2007). Following Ritz et al. 15 (2001) and Cuzzone et al. (2019), precipitation P along the flowline is estimated based on the Clausius-Clapeyron relation. In particular precipitation is estimated by

$$P = P_T + \Delta P = P_m \exp(\lambda_p(T - T_m)) + \Delta P. \quad (4)$$

The term P_T accounts for changes in precipitation solely due to changes in temperature. Here $\lambda_p = 0.07$, which results in a 7% increase in precipitation for every 1° C increase in temperature above modern (Abe-Ouchi et al., 2007; Ritz et al., 2001).

20 The term P_T does not capture the effects of many unknown climate factors that may have caused dynamic, regional changes in Holocene precipitation. Therefore, we introduce a precipitation anomaly term ΔP , analogous to the temperature anomaly ΔT . This time-dependent function, which has units of meters water equivalent (m.w.e.) a^{-1} , is used to adjust precipitation uniformly across a flowline in order to reduce mismatch between modeled and observed terminus positions. Unlike ΔT , which can be inferred from ice cores, ΔP will be used as a control variable to be determined using the inverse methods detailed in 25 section 2.5.3. Equations 3 and 4 provide a method of accounting for elevation changes through time and downscaling inputs to match the mesh resolution of the model ($\sim 1 \text{ km}$).

Positive degree days and snowfall are computed month-by-month based on mean monthly temperature and precipitation (Johannesson et al., 1995). Snow is melted first at a rate of $5 \times 10^{-3} \text{ m.w.e. per degree day}$ followed by ice at a rate of $8 \times 10^{-3} \text{ m.w.e. per degree day}$. Snow melt is initially supposed to refreeze in the snowpack as superimposed ice. Runoff begins 30 when the superimposed ice reaches a given fraction (60%) of the snow cover (Reeh, 1991). A listing of ice-flow and PDD model parameters is provided in Table 1, and all data sets used in the model are shown in Table 2.

Table 1. Summary of primary model parameters used in this work. Default values are provided where applicable.

Description	Symbol	Value	Units
PDD Model Parameters			
Std. deviation	σ	5.5	C
Std. deviation, accumulation	σ_a	5 C	
Ablation rate, snow	λ_s	5×10^{-3}	m.w.e. C ⁻¹ d ⁻¹
Ablation rate, ice	λ_i	8×10^{-3}	m.w.e. C ⁻¹ d ⁻¹
Precipitation param.	λ_p	7×10^{-2}	C ⁻¹
Superimposed ice fraction	p_{max}	0.6	-
Ice Flow Parameters			
Rate factor	A	3.5×10^{-25}	s ⁻¹ Pa ⁻³
Basal traction	β^2	1.2×10^{-3}	Pa a m ⁻¹
Water pressure fraction	P_{frac}	0.85	-
Precipitation Prior			
Prior kernel variance	σ_p^2	5×10^{-4}	-
Prior time Scale	τ	8×10^3	-
Measurement Prior			
Prior kernel variance	σ_ℓ^2	1×10^6	-

2.4 Modeling Limitations

A limitation of our modeling approach is that we do not account for potential ice dynamical effects caused by changes in surface runoff or subglacial hydrology. Modeling melt water runoff would be difficult in a flowline model due to flux of melt water in and out of the path of ice flow. Another limitation is that our model is isothermal. Unless ice temperature is treated in a vertically averaged sense, resolving temperature would require a 2D mesh, which would considerably increase the computational cost of the model. We consider the consequences of this simplification in section 3.3, where we test sensitivity to the ice hardness parameter.

The PDD scheme outlined in section 2.3 does not account for changes in surface mass balance due to orographic forcing, or other complex interactions between the ice sheet and climate system that could be captured by coupling the ice sheet model to an Earth system model (Bahadory and Tarasov, 2018). Since our emphasis is on estimating precipitation and surface mass balance using an inverse modeling approach, we believe the computational cost of such an approach outweighs the benefits. Uncertainties related to feedbacks between ice dynamics and climate are assessed via extensive sensitivity testing (section 3.3).

2.5 Data Assimilation Approach

In order to assess the initial mismatch between modeled and observed retreat histories, we perform a reference experiment with $\Delta P = 0$ and ΔT estimates from both Buijzer et al. (2018) and Dahl-Jensen et al. (1998). To improve the fit to observations, we assimilate terminus position data to obtain improved estimates of Holocene precipitation anomalies. Previous modeling

Table 2. Citations for the primary data sets used in this work.

Data	Citation
Terminus position chronology	Young et al. (2019, In Review)
Bedrock elevation	Morlighem et al. (2017)
Modern ice surface velocity	Mouginot et al. (2017)
Modern precipitation	Box (2013)
Temperature reconstructions	Buizert et al. (2018)
	Dahl-Jensen et al. (1998)
Glacial Isostatic Adjustment	Caron et al. (2018)

studies indicate that Holocene retreat in land-terminating sectors of the GrIS were dominated by surface mass balance rather than ice dynamics (Cuzzone et al., 2019; Lecavalier et al., 2014). Uncertainty in Holocene climate, and consequently surface mass balance, is therefore likely the primary cause of discrepancies between modeled and observed terminus positions.

In principle ΔP , ΔT , or both could be tuned to improve the fit between modeled and observed terminus positions. We focus 5 on precipitation because it is more poorly constrained than temperature. In the upcoming sections, we introduce a framework for time-dependent data assimilation based on the unscented transform (UT). Sections 2.5.1 - 2.5.2 outline the basic tenets of the UT. Sections 2.5.3 - 2.5.7 outline how the UT can be applied to estimate precipitation anomalies.

2.5.1 Overview of the Unscented transform

In what follows, the notation $\mathbf{x} \sim \mathcal{N}(\mathbf{x}_0, P_x)$ means that \mathbf{x} is a normally distributed random variable with mean vector \mathbf{x}_0 and 10 covariance matrix P_x . Suppose that $\mathbf{x} \sim \mathcal{N}(\mathbf{x}_0, P_x)$, and $\mathcal{F} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a nonlinear function. We would like to estimate the distribution of the non-Gaussian random variable

$$\mathbf{y} = \mathcal{F}(\mathbf{x}) + \boldsymbol{\epsilon}. \quad (5)$$

where $\boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, R)$ is the measurement noise.

In general, the non-Gaussian probability distribution for \mathbf{y} can be approximated using Markov chain Monte Carlo (MCMC) 15 methods such as the Metropolis-Hastings algorithm (Chib and Greenberg, 1995). However, if the nonlinear function is time-consuming to compute, generating thousands of MCMC samples is often intractable. As a computationally efficient alternative to MCMC methods, Julier and Uhlmann (1997) introduced a method for approximating the mean and covariance of \mathbf{y} called the unscented transform (UT)¹.

The term unscented transform has been applied somewhat broadly to a family of methods that approximate the statistical 20 moments of a non-Gaussian random variable using a small, deterministic set of sample points called sigma points. It is known

¹According to Jeffrey Uhlmann, the creator of the UT, the term “unscented” was inspired by a stick of deodorant and has no technical significance.

primarily in the context of the unscented Kalman filter. However, the UT can be applied more generally as an alternative to traditional MCMC methods. Sigma points and weight sets are designed to accurately estimate moments of a transformed random variable using a minimal number of function evaluations.

A set of vectors, called sigma points, are chosen with the same weighted sample mean and weighted covariance structure as \mathbf{x} . There are many algorithms for generating sigma points sets with different numbers of points and orders of accuracy. A commonly used set of $2n + 1$ sigma points is given by

$$\boldsymbol{\chi}_i = \begin{cases} \mathbf{x}_0 & i = 0 \\ \mathbf{x}_0 - \sqrt{n + \kappa} [\sqrt{P_x}]_i & i = 1, \dots, n \\ \mathbf{x}_0 + \sqrt{n + \kappa} [\sqrt{P_x}]_i & i = n + 1, \dots, 2n \end{cases} \quad (6)$$

with corresponding mean / covariance weights given by

$$w_i^{(m)} = w_i^{(c)} = \begin{cases} \kappa / (n + \kappa) & i = 0 \\ 1/2(n + \kappa) & \text{otherwise} \end{cases} \quad (7)$$

The notation $[\sqrt{P_x}]_i$ refers to the i -th row of a matrix square root (typically computed by Cholesky factorization) of P_x , and κ is a free parameter controlling the scaling of the sigma points around the mean. Julier and Uhlmann (1997) recommend a default value of $\kappa = 3 - n$. However, κ can be fine tuned to reduce prediction errors for a given problem.

The nonlinear function \mathcal{F} is applied to each sigma point to yield a set of transformed points

$$\mathbf{y}_i = \mathcal{F}(\boldsymbol{\chi}_i). \quad (8)$$

The mean $\bar{\mathbf{y}}$ and covariance matrix P_y of \mathbf{y} are then estimated as weighted sums

$$\bar{\mathbf{y}} = \sum_{i=0}^{2n} w_i^{(m)} \mathbf{y}_i \quad (9)$$

$$P_y = \sum_{i=0}^{2n} w_i^{(c)} (\mathbf{y}_i - \bar{\mathbf{y}})(\mathbf{y}_i - \bar{\mathbf{y}})^T + R. \quad (10)$$

A visual example of this algorithm is shown in Figure 2.

2.5.2 Bayesian Inference using the UT

Given a measurement \mathbf{y}_o , we would like to estimate the posterior distribution

$$P(\mathbf{x}|\mathbf{y}_o) \propto P(\mathbf{y}|\mathbf{x})P(\mathbf{x}). \quad (11)$$

Using an approach called statistical linearization (Sarkka, 2013) the joint distribution for $[\mathbf{x}, \mathbf{y}]^T$ can be approximated by

$$\begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix} \sim \mathcal{N} \left(\begin{bmatrix} \mathbf{x} \\ \boldsymbol{\mu} \end{bmatrix}, \begin{bmatrix} P_x & P_{xy} \\ P_{xy}^T & P_y \end{bmatrix} \right) \quad (12)$$

with

$$\boldsymbol{\mu} = \sum_i w_i^{(m)} \mathcal{Y}_i \quad (13)$$

$$P_y = \sum_i w_i^{(c)} (\mathcal{Y}_i - \boldsymbol{\mu})(\mathcal{Y}_i - \boldsymbol{\mu})^T + R \quad (14)$$

5

$$P_{xy} = \sum_i w_i^{(c)} (\mathcal{X}_i - \mathbf{x}_0)(\mathcal{Y}_i - \boldsymbol{\mu})^T. \quad (15)$$

Here, \mathcal{X}_i and $\mathcal{Y}_i = \mathcal{F}(\mathcal{X}_i)$ are sigma points and transformed sigma points respectively. Matrices P_y and P_{xy} are known as the measurement covariance and cross covariance respectively.

Given a measurement \mathbf{y}_o , the joint distribution then easily yields a Gaussian approximation of the posterior distribution.

10 Letting

$$K = P_{xy} P_y^{-1} \quad (16)$$

we have

$$\mathbf{x}' = \mathbf{x}_o + K [\mathbf{y}_o - \boldsymbol{\mu}] \quad (17)$$

$$15 \quad P' = P_x - K P_y K^T \quad (18)$$

where \mathbf{x}' and P' are approximations of the posterior mean and covariance respectively. Readers familiar with Kalman filters might recognize that \mathbf{x}' and P' are computed using a Kalman “update” step given a measurement \mathbf{y}_o and Kalman gain K (Sarkka, 2013).

2.5.3 Assimilating Glacier Length Observations

20 Time-dependent data assimilation using the UT involves running the ice sheet model with a set of different precipitation anomaly histories, each corresponding to a different sigma point. This is followed by a post-processing step, which incorporates the ice sheet terminus chronology data via a correction of the prior mean vector and covariance matrix. Implementation of the unscented transform is straightforward and easily parallelizable since each model run is independent. In the following section, we outline the mathematical details of this process.

25 We seek to find ΔP histories that match the observed retreat history on both flowlines. An optimal solution should reproduce the observed retreat history within uncertainty, while not overfitting the data. We discretize the problem by estimating the precipitation anomaly at times t_1, t_2, \dots, t_n ranging from 11.6 - 0 ka BP. In practice, we use a regular grid of 44 points, spaced

roughly 250 years apart. Precipitation anomaly values at these time points are denoted by $\Delta p_1, \Delta p_2, \dots, \Delta p_n$ respectively and assembled in a vector

$$\Delta \mathbf{p} = [\Delta p_1, \Delta p_2, \dots, \Delta p_n]^T. \quad (19)$$

Given a multivariate Gaussian prior $\Delta \mathbf{p} \sim \mathcal{N}(\Delta \mathbf{p}_0, P_0)$, which encodes assumptions about the structure of the precipitation

5 anomaly (section 2.5.4), we would like to estimate the mean and covariance of the posterior distribution

$$P(\Delta \mathbf{p} | \ell_0) \propto P(\ell_0 | \Delta \mathbf{p}) P(\Delta \mathbf{p}). \quad (20)$$

Here, the measurement vector

$$\ell_0 = [\ell_1, \ell_2, \dots, \ell_m]^T \quad (21)$$

contains measured glacier lengths at discrete points in time. Our procedure for defining the measurement mean \mathbf{y}_0 and covari-

10 ance R are discussed in section 2.5.5.

We can think of the ice sheet model as a function that maps precipitation anomaly inputs to glacier length outputs with some additive observation noise

$$\ell = \mathcal{F}(\Delta \mathbf{p}) + \epsilon. \quad (22)$$

Discrete precipitation anomaly values are linearly interpolated for input into the ice-dynamics model, which has time steps on

15 the order of months. The function \mathcal{F} returns glacier lengths at the same m discrete times as in ℓ_0 .

To predict the posterior distribution, we use the methods outlined in sections 2.5.1 and 2.5.2. First, sigma points are generated based on the prior distribution for $\Delta \mathbf{p}$. To reduce computational costs, we use a minimal set of $n+1 = 45$ sigma points \mathcal{P}_i with corresponding weights $w_i^{(m)} = w_i^{(c)}$ generated using the method presented in Menegaz et al. (2011). Their method includes one free parameter $0 < w_0 < 1$, which can be tuned to reduce prediction errors. While this method has a lower order of accuracy

20 than other methods, we find it often produces comparable results to other larger sigma point sets in practice.

Sigma points are propagated through the model to obtain transformed points $\mathcal{L}_i = \mathcal{F}(\mathcal{P}_i)$. In this context, sigma points \mathcal{P}_i correspond to different time-dependent precipitation anomaly histories, while the transformed points \mathcal{L}_i correspond to the resulting glacier length histories given those precipitation anomalies as input (Figure 3). The structure of the sigma points reflects the mathematical formulation of the Menegaz et al. (2011) sigma points. Hence they are not merely random samples

25 from the prior distribution.

Transformed sigma points are computed simultaneously in parallel, using one core per sigma point. After all transformed sigma points have been computed, the mean and covariance of the posterior distribution are estimated as outlined in Section 2.5.2. In parallel, this procedure takes roughly the same amount of time as a single forward model run. Unlike a standard filtering approach to data assimilation, all measurement data is incorporated simultaneously rather than time step by time step.

30 For that reason, the Kalman update step corrects the entire time-dependent precipitation history at once. Moreover, unlike in

Kalman smoothing, we approximate the full posterior distribution rather than the probability distributions

$$P(\Delta p_i | \ell_0) \\ i = 1, \dots, n \quad (23)$$

Note that the variables Δp_k with $k \neq i$ are marginalized out of the Kalman smoothing distributions. The use of time-dependent sigma points distinguishes our approach from standard Kalman filtering or Kalman smoothing approaches, and does not rely 5 on the assumptions that states (Δp_i 's) and measurements (ℓ_i 's) satisfy the Markov property.

2.5.4 Gaussian Process Prior for Regularization

We adopt a Gaussian process prior (Rasmussen, 2004) to control the temporal smoothness of ΔP . A Gaussian process can be thought of as a distribution over functions. That is, random samples from a Gaussian process are functions rather than individual points or vectors. A collection of random variables $\{f(t) : t \in \mathcal{T}\}$ is said to be drawn from a Gaussian process 10 with mean function $m(\cdot)$ and covariance function $k(\cdot, \cdot)$ if for any finite set of elements $t_1, \dots, t_n \in \mathcal{T}$, the random variables $f(t_1), \dots, f(t_n)$ have the distribution

$$\mathbf{f} \sim \mathcal{N}(\mathbf{m}, K) \quad (24)$$

with

$$\mathbf{f} = [f(t_1), f(t_2), \dots, f(t_n)]^T, \quad (25)$$

15

$$\mathbf{m} = [m(t_1), m(t_2), \dots, m(t_n)]^T, \quad (26)$$

and

$$K = \begin{bmatrix} k(t_1, t_1) & \dots & k(t_1, t_n) \\ \vdots & \ddots & \vdots \\ k(t_n, t_1) & \dots & k(t_n, t_n) \end{bmatrix}. \quad (27)$$

The set \mathcal{T} is called the index set, and specifies the domain of the Gaussian process. Here, the index set represents points in 20 time.

The prior distribution for Δp has mean vector Δp_0 and covariance matrix $P_0 = K$ of the form shown in Equation 27. We use a squared exponential covariance function

$$k(t, t') = \sigma_p^2 \exp\left(\frac{(t - t')^2}{2\tau^2}\right) \quad (28)$$

where σ^2 is a scaling constant, and τ is a characteristic time scale. Variables Δp_i and Δp_k are more highly correlated the closer 25 they are in time. In effect, this acts as a form of temporal regularization, in which smooth precipitation history functions are preferred over less smooth ones. We discuss the choice of the mean Δp_0 in section 3.2.

2.5.5 Measurement Mean and Variance

Observations of terminus position are available roughly every 1000 years between 11.6 and 7.2 ka BP, with a gap from 7.2 ka BP to present. In contrast, model time steps are on the order of months. Due to these disparate time scales, we use the following procedure to estimate the measurement mean ℓ_0 and covariance matrix R on a time scale more appropriate for the ice sheet model.

We define a Gaussian process prior of “candidate” glacier length histories $\ell_p(t)$ as follows. The mean function $\hat{\ell}_p(t)$ is obtained by linearly interpolating between glacier length observations. The Brownian covariance kernel for the Gaussian process is defined by

$$k_p(t, t') = \sigma_\ell^2 \min(t, t'). \quad (29)$$

10 Candidate retreat histories are generated by drawing random samples from the Gaussian process. The Brownian covariance kernel results in random glacier length histories that are somewhat noisy, but correlated over shorter time scales (Figure ??). Candidate length histories are resampled so that the mean moraine formation times and uncertainties match the observations in Young et al. (2019, In Review). Hence, highly implausible candidate histories are rejected. The average length and variance of this culled set of samples is computed at a series of time slices to obtain a plausible measurement mean ℓ_0 and diagonal
15 measurement covariance matrix R (Figure 4).

2.5.6 Iterative Optimization Procedure

Optimizations are conducted in multiple passes. In the first pass, the measurement covariance matrix R is multiplied by a factor of 1/4 so that the measurements are initially weighted more than the prior. This produces a reasonable fit to the data, even given a poor initial estimate of ΔP . The optimal precipitation anomaly from a given iteration is used as the prior mean in
20 the next iteration. We use the same prior covariance matrix P_0 for regularization in each iteration. After two to three iterations, modeled and observed terminus positions match within measurement uncertainty (Section 3). In our experience, the results of iteration are not dependent on the choice of prior mean in the first iteration, but we find that convergence can be improved by choosing a sensible initial guess as in Section 3.2.

2.5.7 Approach to Sensitivity Testing

25 As described, the data assimilation method accounts for measurement but not model uncertainty. It can easily be extended to account for uncertainties in the ice flow and PDD model parameters. We define an augmented state vector

$$\mathbf{u} = [\Delta p, \boldsymbol{\theta}]^T \quad (30)$$

where $\boldsymbol{\theta}$ is a vector of scalar parameters including the natural logarithm of the rate factor for ice, the basal traction parameter, a parameter controlling precipitation scaling with temperature, and the PDD melt rate parameters for ice and snow

$$30 \quad \boldsymbol{\theta} = [\ln(A), \beta^2, \lambda_p, \lambda_i, \lambda_s]^T. \quad (31)$$

The prior distribution for the augmented state vector is given by

$$\mathbf{u} \sim \mathcal{N} \left(\begin{bmatrix} \Delta \mathbf{p}_0 \\ \boldsymbol{\theta}_0 \end{bmatrix}, \begin{bmatrix} P_0 & 0 \\ 0 & \Theta \end{bmatrix} \right) \quad (32)$$

where $\boldsymbol{\theta}_0$ is the parameter mean vector and Θ is a diagonal matrix containing parameter variances.

The unscented transform is applied to the augmented function

$$5 \quad \boldsymbol{\ell} = \hat{\mathcal{F}}(\mathbf{u}) + \boldsymbol{\epsilon} \quad (33)$$

to obtain estimates of the joint distribution for $[\Delta \mathbf{p}, \boldsymbol{\theta}, \boldsymbol{\ell}]^T$ and the conditional distribution for $[\Delta \mathbf{p}, \boldsymbol{\theta}]^T | \boldsymbol{\ell}_0$. Since parameters are included as state variables, sigma points reflect a variety of precipitation histories and parameter sets. A model run for a particular sigma point is initialized from an appropriate steady state using the parameter set for that point.

2.6 Model Initialization

10 Model runs are initialized by tuning the precipitation anomaly to obtain a steady state at 12.6 ka BP, with a margin position 5 km beyond the 11.6 ka BP moraine. We invert for a precipitation anomaly time series that forces a retreat of 5 km over 1000 years to obtain an initial ice sheet configuration with the correct terminus position at 11.6 ka BP (Figure 5). This initialization procedure is intended to ease the ice sheet out of steady state in order to avoid strong transient effects at the beginning of model runs.

15 3 Results

3.1 Reference Experiment

To assess the initial misfit between modeled and observed Holocene retreat, the model is forced with ΔT reconstructions from Buizert et al. (2018) and Dahl-Jensen et al. (1998) and a zero precipitation anomaly. Precipitation is scaled with temperature according to Equation 4, neglecting possible influences from changing Arctic sea ice cover, atmospheric circulation, or other 20 unknown climate factors. Modeled ice retreat is far more rapid than observed on both flowlines (Figure 8). Colder temperatures during the early Holocene (Figure 8 a) lead to a somewhat more plausible retreat history using the Dahl-Jensen et al. (1998) temperature forcing versus the Buizert et al. (2018) forcing. However, by 8 ka BP, ice has retreated inland of the present day margin in both reconstructions.

3.2 Precipitation Anomaly Inversions

25 We estimate precipitation anomalies on the northern and southern flowlines using both the Buizert et al. (2018) and Dahl-Jensen et al. (1998) temperature reconstructions. Given the rapid retreat in the reference experiment, we expect that a positive precipitation anomaly will be required to match observed terminus positions in the early Holocene. Therefore, in the first round

of optimization, we assume a prior mean of the form

$$\Delta P = \frac{1}{2}(1 - \hat{\tau}) \quad (34)$$

where $\hat{\tau}$ is a rescaled time variable that increases from zero at 11.6 ka BP to one at 0 ka BP. The results of the iterative optimization procedure are insensitive to the prior mean selected in the first iteration. Using a sigma point scaling parameter 5 $w_0 = 0.5$ for the Menegaz et al. (2011) sigma point set ensures that a wide region around the mean is explored in each iteration.

Positive precipitation anomalies are predicted throughout most of the Holocene for both temperature reconstructions (Figure 6). While differences between the northern and southern flowlines are relatively minor, there are significant differences in precipitation between the Buizert et al. (2018) and Dahl-Jensen et al. (1998) inversions. In the Buizert et al. (2018) inversion, the largest precipitation anomalies (up to 1 m.w.e. a^{-1}) occur during the early Holocene. Precipitation remains relatively high 10 during the HTM (10 - 6 ka BP), but dips before the 8.2 ka BP cold event. For the Dahl-Jensen et al. (1998) inversion, ΔP is relatively low during the early Holocene, but increases during the HTM (8.5 - 3 ka BP). Unlike the Buizert et al. (2018) inversion, there is an evident trend between HTM warming and increased snowfall (Figure 9).

Forcing the model with mean estimated precipitation anomalies yields plausible retreat histories on both flowlines. Modeled and observed retreat chronologies match within uncertainty (Figure 8). A significant fraction of Holocene precipitation 15 (typically $> 90\%$) falls as snow. Hence, a positive precipitation anomaly can be interpreted directly as additional snowfall / accumulation. Average HTM snowfall is around 35% higher than modern in both temperature reconstructions (Figure 9). However, overall trends in Holocene snowfall differ between reconstructions.

3.3 Sensitivity Testing

We assess the sensitivity of Holocene precipitation anomalies to modeling uncertainties by performing an HTM inversion using 20 the methodology described in Section 2.5.7. To obtain accurate uncertainty estimates, we use a fifth order accurate sigma point set based on Li et al. (2017) (Appendix A), as we find that the second order Menegaz et al. (2011) set likely underestimates covariance. Inversions are performed on the northern flowline using the Buizert et al. (2018) temperature reconstruction. Model runs are initialized from steady states around at 10.5 ka BP, 500 years prior to the Buizert et al. (2018) HTM. Prior and posterior parameter values are reported in Table 3.

25 Mean HTM precipitation anomalies are within 2 cm.w.e. a^{-1} of the inversion presented in section 3.2 (Figure 10). Parameter uncertainties contribute to uncertainty in ΔP . Overall, however, temperature uncertainty is far more significant than uncertainty in model parameters. Differences in model initialization do not significantly impact the results of the inversion. Although sensitivity tests are initialized from steady states at 10.5 ka BP, while inversions in Section 3.2 are initialized from a transient state at 11.6 ka BP, the mean HTM precipitation anomalies are nearly identical. Estimated posterior parameter values are also 30 similar to the assumed prior values (Table 3).

Table 3. Summary of primary model parameters used in this work. Default values are provided where applicable. Prior parameter uncertainties assumed for the sensitivity test are shown in the 2σ column.

Description	Symbol	Prior Mean	2σ Range	Posterior Mean	Units
Ablation rate, snow	λ_s	5×10^{-3}	$3 - 7 \times 10^{-3}$	2.8×10^{-3}	m.w.e. $\text{C}^{-1} \text{d}^{-1}$
Ablation rate, ice	λ_i	8×10^{-3}	$6 - 10 \times 10^{-3}$	8.9×10^{-3}	m.w.e. $\text{C}^{-1} \text{d}^{-1}$
Precipitation param.	λ_p	7×10^{-2}	$5 - 9 \times 10^{-2}$	9.8×10^{-2}	C^{-1}
Rate factor	A	3.5×10^{-25}	$2.1 - 5.7 \times 10^{-25}$	2×10^{-25}	$\text{s}^{-1} \text{Pa}^{-3}$
Basal traction	β^2	1.2×10^{-3}	$1.1 - 1.3 \times 10^{-3}$	1.2×10^{-3}	Pa a m^{-1}

4 Discussion

We infer changes in Holocene precipitation in the Kangerlussuaq region of western Greenland using a new chronology of ice sheet terminus position from Young et al. (2019, In Review) and an inverse modeling procedure based on the unscented transform. We find that scaling precipitation with temperature via the Clausius–Clapeyron equation (Equation 4) results in 5 excessively fast retreat during the early Holocene for both the Buizert et al. (2018) and Dahl-Jensen et al. (1998) temperature reconstructions. Thus, temperature driven changes in precipitation / accumulation alone are not sufficient to reproduce the observed pattern of retreat. Inversions show that adding precipitation throughout the Holocene and specifically during the HTM yields a good fit to observations (Figures 6, 8). Since a large fraction of precipitation falls as snow, a positive precipitation anomaly can be interpreted fairly directly as an increase in accumulation beyond what would be expected from temperature 10 changes alone.

There are considerable differences in predicted precipitation anomalies depending on the assumed temperature history. Inversions using the Buizert et al. (2018) temperature reconstruction show generally decreasing snowfall through the Holocene. Due to high snowfall in the early Holocene (nearly 100% higher than modern) and a dip in HTM precipitation associated with the 8.2 ka BP cold event, average HTM snowfall roughly matches the overall Holocene average, which is about 35% above 15 modern. In contrast, inversions using the Dahl-Jensen et al. (1998) temperature reconstruction show a clear trend between HTM warming and increased snowfall (Figure 9). This trend could be interpreted as a transient increase in snowfall due to reduced Arctic sea ice cover or changes in atmospheric circulation during the HTM.

A large positive ΔP correction during the early Holocene in the Buizert et al. (2018) inversion likely reflects the dependence of precipitation on temperature. Low temperatures during the early Holocene result in low precipitation. Without an additional 20 moisture source to increase snowfall, the ice sheet retreats far more rapidly than observed. In the Dahl-Jensen et al. (1998) reconstruction, warming occurs more gradually in the early Holocene. Accumulation and ablation are more closely balanced, resulting in a smaller precipitation anomaly correction from 11.6 to 10 ka BP.

When interpreting results of precipitation inversions, it is important to consider that the Buizert et al. (2018) and Dahl-Jensen et al. (1998) reconstructions are obtained using different methodologies. The Greenland-wide Buizert et al. (2018) reconstruction is obtained by merging the TraCE-21ka coupled ocean-atmosphere general circulation model of the last deglaciation (Liu 25

et al., 2009; He et al., 2013) with borehole temperature reconstructions at GISP2, NGRIP, and NEEM. Our estimated precipitation history in the Kangerlussuaq region is significantly different from TraCE-21ka, particularly during the early Holocene and parts of the HTM (Figure 7). In contrast to Buizert et al. (2018), the Dahl-Jensen et al. (1998) temperature reconstruction at GRIP offers only a pointwise estimate containing no explicit spatial or seasonal information.

5 While the Buizert et al. (2018) temperature reconstruction is arguably more suitable for our purposes, since it resolves spatial and seasonal patterns in ΔT , there is still considerable uncertainty in temperature, particularly near the ice margins. Despite this, there is some consensus between the Buizert et al. (2018) and Dahl-Jensen et al. (1998) inversions. For example, average HTM snowfall is roughly 35% higher than modern in both cases (Figure 9).

10 Predicted HTM precipitation anomalies are not particularly sensitive to uncertainties in the ice sheet or PDD model parameters, or to the initialization procedure. In the sensitivity test presented in Section 3.3, model runs are initialized using steady states at 10.5 ka BP. Runs in the full Holocene inversions in Section 3.2, which use fixed parameter sets, are initialized from a transient state at 11.6 ka BP. Even considering these differences in model initialization, estimated HTM precipitation anomalies are comparable in all inversions (Figure 10). Additionally, ΔP does not appear to be sensitive to uncertainties in bedrock geometry, as inversions conducted with a static bedrock geometry yielded similar results to inversions accounting for isostatic 15 uplift.

4.1 Conclusions

4.1.1 The Unscented Transform as a Data Assimilation Method

Significant strides have been made in time-dependent data assimilation in glaciology using adjoint based methods. Goldberg and Heimbach (2013) infer the initial thickness and basal conditions for a synthetic ice sheet given snapshots of ice thickness 20 at discrete times. Larour et al. (2014) demonstrate a data assimilation framework within the Ice Sheet System Model (ISSM), capable of obtaining temporal estimates of surface mass balance and basal friction given surface altimetry.

In contrast to adjoint based approaches, the unscented transform (UT) does not require computing the Jacobian or Hessian of an objective function, or special checkpointing code for time-dependent problems. Adjoint based methods are advantageous for extremely high-dimensional problems, as the required number of model runs is independent of the number of parameters. However, the UT provides more accurate uncertainty estimates than linearization (Julier and Uhlmann, 1997). Hessian 25 information can be used to improve uncertainty estimates (Isaac et al., 2015). However, this methodology uses purely local approximations to the nonlinear function around the maximum a posteriori probability (MAP) estimator, which may affect the quality of uncertainty estimates.

The unscented transform has advantages over Markov chain Monte Carlo methods for inference problems with a relatively 30 small number of unknown parameters (< 1000 or so parameters). In this work, we optimize for $n = 44$ parameters representing precipitation anomaly values at discrete points in time. Since function evaluations at each sigma point are independent, the UT is trivially parallelizable. Consequently, on a high end desktop, the iterative optimization process presented in Section 2.5.6 takes roughly the same amount of time as performing three successive forward model runs.

MCMC methods can be parallelized to some extent by utilizing multiple interacting Markov chains running in parallel (e.g. Chowdhury and Jermaine, 2018) or by combining samples from independent chains in a post processing step (e.g. Neiswanger et al., 2013). Nonetheless, computational bottlenecks persist in MCMC methods since individual Markov chains are inherently serial. Consequently, UT approximations to the posterior distribution can be generated hundreds to thousands of times faster
5 for small parameter sets.

A drawback of the UT is that its accuracy is inherently limited by using a predetermined number of sample points. In our case, for example, using a minimal set of $n + 1$ sigma points for n unknown parameters yields accurate mean estimates but appears to underestimate covariance when compared to a higher order cubature method. Without testing multiple sigma point sets and scaling parameters, it can be difficult to assess the accuracy of UT estimates of the posterior. MCMC methods,
10 in contrast, can provide arbitrarily accurate estimates of the posterior given sufficient computation time. Moreover, they can resolve the full posterior distribution rather than computing its moments as in the UT.

An alternative to traditional MCMC methods is to use surrogate models or emulators (Gong and Duan, 2017). Here, a computationally inexpensive surrogate model is trained to approximate the output of a more complex model function. The surrogate model can then be used in place of the full model for the purpose of MCMC sampling, significantly reducing the
15 overall computational cost. A few types of surrogate models have already been applied to glaciological problems including deep neural networks (Tarasov et al., 2012) and Gaussian processes (e.g. Chang et al., 2016; Pollard et al., 2016).

Surrogate models require an initial training phase. Pollard et al. (2016), for example, perform 625 ice sheet model runs using combinations of four unknown simulation parameters. A Gaussian process surrogate model is then fitted to this training data in order to interpolate the model in parameter space for inference. The number and selection of training points significantly
20 affects the performance of the surrogate model.

Interestingly, there is a strong connection between the unscented transform and Gaussian process surrogate models. It can be shown that sigma points for the UT are optimal training points for a Gaussian process surrogate model in the sense of minimizing the variance of the expected value of the posterior distribution (Sarkkä et al., 2016). While the details are technical, one can think of UT sigma points as optimal training points given that (i) the prior distribution is a multivariate Gaussian and
25 (ii) the Gaussian process surrogate model has a polynomial covariance function.

4.1.2 Modeling Conclusions

Our work follows a number of previous observationally constrained paleo ice sheet modeling studies (e.g. Tarasov and Peltier, 2002; Lecavalier et al., 2014; Calov et al., 2015). Perhaps most relevant to this work is Lecavalier et al. (2014), who model the deglaciation of Greenland from the Last Glacial Maximum using a 3-D thermomechanically coupled ice sheet model. Model
30 runs in Lecavalier et al. (2014) are informed by constraints on relative sea level, ice core thinning, and LGM ice sheet extent.

In contrast to previous modeling studies, the computational efficiency of the flowline model outlined in Brinkerhoff et al. (2017) makes time-dependent data-assimilation, sensitivity testing, and robust uncertainty estimation tractable. Sensitivity testing indicates that estimated precipitation is insensitive to parameter uncertainties in the PDD and ice-dynamics models. This conclusion supports earlier findings showing that modeled Holocene retreat in land terminating sectors of the GrIS is

more sensitive to surface mass balance than other factors like the flow law or basal sliding (Cuzzone et al., 2019; Lecavalier et al., 2014).

A drawback of our modeling approach is that we cannot account for inherently map plane effects such as changes in ice flow direction, or convergent / divergent flow in or out of the assumed flow path. These factors likely contribute to small discrepancies in estimated precipitation anomalies between the northern and southern flowlines (Figure 6). **Beyond computational efficiency, there are number of reasons why a flowline model is appropriate for our purposes. The modern flow field near Kangerlussuaq can be characterized by relatively simple east-to-west flow, and there is an absence of strongly convergent flow into outlet glaciers. Given the low bedrock relief in the region, and the surface mass balance dominated retreat pattern (Van Tatenhove et al., 1996; Cuzzone et al., 2019), it is reasonable to assume that the flow regime was similar during the Holocene.**

10 In this work, we do not treat temperature as a random variable with its own covariance structure. However, differences between the Buizert et al. (2018) and Dahl-Jensen et al. (1998) inversions indicate that temperature is the dominant source of uncertainty in ΔP . This result underscores the importance of generating improved, regionally specific, temperature reconstructions constrained by proxy records. More regionally specific estimates of temperature would help to decrease uncertainty in the estimated precipitation history.

15 Despite lingering uncertainties, our modeling results indicate that the Holocene thermal maximum was accompanied by elevated snowfall, which slowed ice retreat in the Kangerlussuaq region of the GrIS. Inversions conducted using both the Buizert et al. (2018) and Dahl-Jensen et al. (1998) temperature reconstructions show average HTM snowfall around 35% higher than modern. More specifically, in the Dahl-Jensen et al. (1998) inversion, there is a clear trend between HTM warming and increased accumulation, which could be interpreted as transient increase in precipitation due to reduced Arctic sea ice 20 cover or changes in atmospheric circulation.

Code availability. The flowline ice sheet model and PDD model used in this work are available at https://github.com/JacobDowns/flow_model.

Appendix A: A Higher Order Method for Estimating Covariance

Estimating the mean and covariance of the posterior distribution requires approximating Gaussian weighted expectation integrals of the form

$$E[\mathcal{F}(\mathbf{x})] = \int_{\mathbb{R}^n} \mathcal{F}(\mathbf{x}) \mathcal{N}(\mathbf{x} | \mathbf{0}, I) d\mathbf{x} \quad (\text{A1})$$

via numerical integration rules, also called cubature rules, which approximate the expectation integral as a weighted sum

$$E[\mathcal{F}(\mathbf{x})] \approx Q[\mathcal{F}(\mathbf{x})] = \sum_{i=1}^N w_i \mathcal{F}(\mathbf{x}_i). \quad (\text{A2})$$

Here, the notation $\mathcal{N}(\mathbf{x} | \mathbf{0}, I)$ refers to a Gaussian probability density function evaluated at the point \mathbf{x} . General Gaussian weight functions $\mathcal{N}(\mathbf{x} | \mathbf{x}_0, P_x)$ are handled by changing variables. Letting $\sqrt{P_x}$ be a matrix square root of the covariance matrix, we have

$$\begin{aligned} & \int_{\mathbb{R}^n} \mathcal{F}(\mathbf{x}) \mathcal{N}(\mathbf{x} | \mathbf{x}_0, P_x) d\mathbf{x} \\ &= \int_{\mathbb{R}^n} \mathcal{F}(\mathbf{x}_0 + \sqrt{P_x} \boldsymbol{\xi}) \mathcal{N}(\boldsymbol{\xi} | \mathbf{0}, I) d\boldsymbol{\xi}. \end{aligned} \quad (\text{A3})$$

5 Cubature rules, including the unscented transform, are constructed to exactly integrate polynomial functions $\mathcal{F}(\mathbf{x})$ up to a certain degree d . Suppose that $\mathbf{x} = [x_1, x_2, \dots, x_n]^T$ is a point in \mathbb{R}^n . A monomial of degree d refers to a function $x_1^{i_1} x_2^{i_2} \cdots x_n^{i_n}$ where the exponents are non-negative integers that sum to d . A polynomial of degree d is a linear combination of monomials with highest degree d .

10 Li et al. (2017) describe a fifth-order cubature rule using fully symmetric sets of sigma points. A set $X = \{\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_N\}$ is fully symmetric if it is closed under the operations of coordinate position and sign permutations. Their cubature rule has the form

$$\begin{aligned} Q[\mathcal{F}(\mathbf{x})] &= w_1 \mathcal{F}(0, 0, \dots, 0) \\ &+ w_2 \sum_{\text{full sym}} \mathcal{F}(\lambda, 0, \dots, 0) \\ &+ w_3 \sum_{\text{full sym}} \mathcal{F}(\lambda, \lambda, 0, \dots, 0). \end{aligned} \quad (\text{A4})$$

The notation $\sum_{\text{full sym}} \mathcal{F}(\cdot)$ refers to a sum of the function \mathcal{F} evaluated at all points in the fully symmetric set generated by the given point.

15 Due to the symmetry of the sigma points and the Gaussian weight function, all moments (that is, integrals of Gaussian weighted monomial functions) containing an odd order exponent are automatically satisfied. Exploiting this fact, and the symmetries of the sigma points, it can be shown that satisfying the remaining moment constraint equations up the fifth order reduces to solving the following system of four equations in four unknowns w_1, w_2, w_3 and λ

$$\begin{aligned} E[1] &= w_1 + 2nw_2 + 2n(n-1)w_3 \\ E[x_i^2] &= 2\lambda^2 w_2 + 4(n-1)\lambda^2 w_3 \\ E[x_i^4] &= 2\lambda^4 w_2 + 4(n-1)\lambda^4 w_3 \\ E[x_i^2 x_j^2] &= 4\lambda^4 w_3. \end{aligned} \quad (\text{A5})$$

20 By slightly modifying this cubature rule

$$\begin{aligned} Q[\mathcal{F}(\mathbf{x})] &= w_1 \mathcal{F}(0, 0, \dots, 0) \\ &+ w_2 \sum_{\text{full sym}} \mathcal{F}(\lambda_1, 0, \dots, 0) \\ &+ w_3 \sum_{\text{full sym}} \mathcal{F}(\lambda_2, \lambda_2, 0, \dots, 0) \end{aligned} \quad (\text{A6})$$

we introduce a new free parameter λ_2 that allows scaling of the sigma points about the mean. The new moment constraint equations become

$$\begin{aligned}
E[1] &= w_1 + 2nw_2 + 2n(n-1)w_3 \\
E[x_i^2] &= 2\lambda_1^2 w_2 + 4(n-1)\lambda_2^2 w_3 \\
E[x_i^4] &= 2\lambda_1^4 w_2 + 4(n-1)\lambda_2^4 w_3 \\
E[x_i^2 x_j^2] &= 4\lambda_2^4 w_3.
\end{aligned} \tag{A7}$$

Using that $E[1] = 1$, $E[x_i^2] = 1$, $E[x_i^4] = 3$, and $E[x_i^2 x_j^2] = 1$ we obtain

$$\begin{aligned}
\lambda_1 &= \frac{\lambda_2 \sqrt{n-4}}{n - \lambda_2^2 - 1} \\
w_2 &= \frac{4-n}{2\lambda_1^4} \\
w_3 &= \frac{1}{4\lambda_2^2}
\end{aligned} \tag{A8}$$

$$w_1 = 1 - 2nw_2 - 2n(n-1)w_3$$

with $n > 4$ and $n - \lambda_2^2 - 1 \neq 0$. A drawback of the original cubature rule, as well our modified version here, is that it requires negative weights, which can lead to numerical instability.

Author contributions. JD was the lead author of this manuscript. JJ and JC assisted JD with ice sheet modeling efforts. JB, NY, and AL provided the moraine age data used in this work. JD and JJ wrote the manuscript with input from all authors.

10 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. JD and JJ were supported by NSF grant 1504457. JD would like to thank Doug Brinkerhoff for providing code for the flowline ice sheet model used in this work, as well as many elucidating discussions. JD would also like to thank John Bardsley for providing many mathematical insights into inverse problems. Our thanks to reviewers Stephen Price, Lev Tarasov, and editor Kerim Hestnes Nisancioglu, who helped to significantly improve the quality of this manuscript.

References

Abe-Ouchi, A., Segawa, T., and Saito, F.: Climatic conditions for modelling the Northern Hemisphere ice sheets throughout the ice age cycle, <https://doi.org/10.5194/cp-3-423-2007>, 2007.

Alley, R. B., Meese, D. A., Shuman, C. A., Gow, A. J., Taylor, K. C., Grootes, P. M., White, J. W., Ram, M., Waddington, E. D., Mayewski, P. A., and Zielinski, G. A.: Abrupt increase in Greenland snow accumulation at the end of the Younger Dryas event, *Nature*, <https://doi.org/10.1038/362527a0>, 1993.

Bahadory, T. and Tarasov, L.: LCice 1.0-a generalized Ice Sheet System Model coupler for LOVECLIM version 1.3: Description, sensitivities, and validation with the Glacial Systems Model (GSM version D2017.aug17), *Geoscientific Model Development*, <https://doi.org/10.5194/gmd-11-3883-2018>, 2018.

10 Bintanja, R. and Selten, F. M.: Future increases in Arctic precipitation linked to local evaporation and sea-ice retreat, *Nature*, <https://doi.org/10.1038/nature13259>, 2014.

Blatter, H.: Velocity and stress fields in grounded glaciers: a simple algorithm for including deviatoric stress gradients, *Journal of Glaciology*, <https://doi.org/10.1017/S002214300001621X>, 1995.

Box, J. E.: Greenland ice sheet mass balance reconstruction. Part II: Surface mass balance (1840-2010), *Journal of Climate*, <https://doi.org/10.1175/JCLI-D-12-00518.1>, 2013.

15 Brinkerhoff, D., Truffer, M., and Aschwanden, A.: Sediment transport drives tidewater glacier periodicity, *Nature Communications*, <https://doi.org/10.1038/s41467-017-00095-5>, 2017.

Buizert, C., Keisling, B. A., Box, J. E., He, F., Carlson, A. E., Sinclair, G., and DeConto, R. M.: Greenland-Wide Seasonal Temperatures During the Last Deglaciation, *Geophysical Research Letters*, <https://doi.org/10.1002/2017GL075601>, 2018.

20 Calov, R., Robinson, A., Perrette, M., and Ganopolski, A.: Simulating the Greenland ice sheet under present-day and palaeo constraints including a new discharge parameterization, *Cryosphere*, <https://doi.org/10.5194/tc-9-179-2015>, 2015.

Caron, L., Ivins, E. R., Larour, E., Adhikari, S., Nilsson, J., and Blewitt, G.: GIA Model Statistics for GRACE Hydrology, *Cryosphere*, and *Ocean Science*, *Geophysical Research Letters*, <https://doi.org/10.1002/2017GL076644>, 2018.

Chang, W., Haran, M., Applegate, P., and Pollard, D.: Calibrating an Ice Sheet Model Using High-Dimensional Binary Spatial Data, *Journal 25 of the American Statistical Association*, <https://doi.org/10.1080/01621459.2015.1108199>, 2016.

Chib, S. and Greenberg, E.: Understanding the metropolis-hastings algorithm, *American Statistician*, <https://doi.org/10.1080/00031305.1995.10476177>, 1995.

Chowdhury, A. and Jermaine, C.: Parallel and Distributed MCMC via Shepherding Distributions, *Proceedings of the Twenty-First International Conference on Artificial Intelligence and Statistics*, 84, 1819–1827, <http://proceedings.mlr.press/v84/chowdhury18a.html>, 2018.

30 Cuzzone, J. K., Schlegel, N. J., Morlighem, M., Larour, E., Briner, J. P., Seroussi, H., and Caron, L.: The impact of model resolution on the simulated Holocene retreat of the southwestern Greenland ice sheet using the Ice Sheet System Model (ISSM), *Cryosphere*, <https://doi.org/10.5194/tc-13-879-2019>, 2019.

Dahl-Jensen, D., Mosegaard, K., Gundestrup, N., Clow, G. D., Johnsen, S. J., Hansen, A. W., and Balling, N.: Past temperatures directly from the Greenland Ice Sheet, *Science*, <https://doi.org/10.1126/science.282.5387.268>, 1998.

35 Goldberg, D. N. and Heimbach, P.: Parameter and state estimation with a time-dependent adjoint marine ice sheet model, *Cryosphere*, <https://doi.org/10.5194/tc-7-1659-2013>, 2013.

Gong, W. and Duan, Q.: An adaptive surrogate modeling-based sampling strategy for parameter optimization and distribution estimation (ASMO-PODE), *Environmental Modelling and Software*, 95, 61–75, <https://doi.org/10.1016/j.envsoft.2017.05.005>, 2017.

He, F., Shakun, J. D., Clark, P. U., Carlson, A. E., Liu, Z., Otto-Bliesner, B. L., and Kutzbach, J. E.: Northern Hemisphere forcing of Southern Hemisphere climate during the last deglaciation, *Nature*, <https://doi.org/10.1038/nature11822>, 2013.

5 Isaac, T., Petra, N., Stadler, G., and Ghattas, O.: Scalable and efficient algorithms for the propagation of uncertainty from data through inference to prediction for large-scale problems, with application to flow of the Antarctic ice sheet, *Journal of Computational Physics*, <https://doi.org/10.1016/j.jcp.2015.04.047>, 2015.

Johannesson, T., Sigurdsson, O., Laumann, T., and Kennett, M.: Degree-day glacier mass-balance modelling with applications to glaciers in Iceland, Norway and Greenland, *Journal of Glaciology*, <https://doi.org/10.1017/S002214300016221>, 1995.

10 Julier, S. J. and Uhlmann, J. K.: New extension of the Kalman filter to nonlinear systems, *Int Symp AerospaceDefense Sensing Simul and Controls*, <https://doi.org/10.1117/12.280797>, 1997.

Kaufman, D. S., Ager, T. A., Anderson, N. J., Anderson, P. M., Andrews, J. T., Bartlein, P. J., Brubaker, L. B., Coats, L. L., Cwynar, L. C., Duvall, M. L., Dyke, A. S., Edwards, M. E., Eisner, W. R., Gajewski, K., Geirsdóttir, A., Hu, F. S., Jennings, A. E., Kaplan, M. R., Kerwin, M. W., Lozhkin, A. V., MacDonald, G. M., Miller, G. H., Mock, C. J., Oswald, W. W., Otto-Bliesner, B. L., Porinchu, D. F., Rühland, K., Smol, J. P., Steig, E. J., and Wolfe, B. B.: Holocene thermal maximum in the western Arctic (0–180°W), *Quaternary Science Reviews*, <https://doi.org/10.1016/j.quascirev.2003.09.007>, 2004.

15 Kobashi, T., Menviel, L., Jeltsch-Thömmes, A., Vinther, B. M., Box, J. E., Muscheler, R., Nakaegawa, T., Pfister, P. L., Döring, M., Leuenberger, M., Wanner, H., and Ohmura, A.: Volcanic influence on centennial to millennial Holocene Greenland temperature change, *Scientific Reports*, <https://doi.org/10.1038/s41598-017-01451-7>, 2017.

20 Larour, E., Utke, J., Csatho, B., Schenk, A., Seroussi, H., Morlighem, M., Rignot, E., Schlegel, N., and Khazendar, A.: Inferred basal friction and surface mass balance of the Northeast Greenland Ice Stream using data assimilation of ICESat (Ice Cloud and land Elevation Satellite) surface altimetry and ISSM (Ice Sheet System Model), *Cryosphere*, <https://doi.org/10.5194/tc-8-2335-2014>, 2014.

Lecavalier, B. S., Milne, G. A., Simpson, M. J., Wake, L., Huybrechts, P., Tarasov, L., Kjeldsen, K. K., Funder, S., Long, A. J., Woodroffe, S., Dyke, A. S., and Larsen, N. K.: A model of Greenland ice sheet deglaciation constrained by observations of relative sea level and ice extent, *Quaternary Science Reviews*, 102, 54–84, <https://doi.org/10.1016/j.quascirev.2014.07.018>, 2014.

25 Li, Z., Yang, W., Ding, D., and Liao, Y.: A Novel Fifth-Degree Cubature Kalman Filter for Real-Time Orbit Determination by Radar, *Mathematical Problems in Engineering*, <https://doi.org/10.1155/2017/8526804>, 2017.

Liu, Z., Otto-Bliesner, B. L., He, F., Brady, E. C., Tomas, R., Clark, P. U., Carlson, A. E., Lynch-Stieglitz, J., Curry, W., Brook, E., Erickson, D., Jacob, R., Kutzbach, J., and Cheng, J.: Transient simulation of last deglaciation with a new mechanism for bolling-allerod warming, *Science*, <https://doi.org/10.1126/science.1171041>, 2009.

30 Marcott, S. A., Shakun, J. D., Clark, P. U., and Mix, A. C.: A reconstruction of regional and global temperature for the past 11,300 years, *Science*, 339, 1198–1201, <https://doi.org/10.1126/science.1228026>, 2013.

Menegaz, H. M., Ishihara, J. Y., and Borges, G. A.: A new smallest sigma set for the Unscented Transform and its applications on SLAM, in: *Proceedings of the IEEE Conference on Decision and Control*, <https://doi.org/10.1109/CDC.2011.6161480>, 2011.

35 Miller, G. H., Brigham-Grette, J., Alley, R. B., Anderson, L., Bauch, H. A., Douglas, M. S., Edwards, M. E., Elias, S. A., Finney, B. P., Fitzpatrick, J. J., Funder, S. V., Herbert, T. D., Hinzman, L. D., Kaufman, D. S., MacDonald, G. M., Polyak, L., Robock, A., Serreze, M. C., Smol, J. P., Spielhagen, R., White, J. W., Wolfe, A. P., and Wolff, E. W.: Temperature and precipitation history of the Arctic, *Quaternary Science Reviews*, <https://doi.org/10.1016/j.quascirev.2010.03.001>, 2010.

Morlighem, M., Williams, C. N., Rignot, E., An, L., Arndt, J. E., Bamber, J. L., Catania, G., Chauché, N., Dowdeswell, J. A., Dorschel, B., Fenty, I., Hogan, K., Howat, I., Hubbard, A., Jakobsson, M., Jordan, T. M., Kjeldsen, K. K., Millan, R., Mayer, L., Mouginot, J., Noël, B. P., O'Cofaigh, C., Palmer, S., Rysgaard, S., Seroussi, H., Siegert, M. J., Slabon, P., Straneo, F., van den Broeke, M. R., Weinrebe, W., Wood, M., and Zinglersen, K. B.: BedMachine v3: Complete Bed Topography and Ocean Bathymetry Mapping of Greenland From Multibeam Echo Sounding Combined With Mass Conservation, *Geophysical Research Letters*, <https://doi.org/10.1002/2017GL074954>, 2017.

Mouginot, J., Rignot, E., Scheuchl, B., and Millan, R.: Comprehensive annual ice sheet velocity mapping using Landsat-8, Sentinel-1, and RADARSAT-2 data, *Remote Sensing*, <https://doi.org/10.3390/rs9040364>, 2017.

Neiswanger, W., Wang, C., and Xing, E.: Asymptotically Exact, Embarrassingly Parallel MCMC, *Proceeding UAI'14 Proceedings of the Thirtieth Conference on Uncertainty in Artificial Intelligence*, pp. 1–16, <http://arxiv.org/abs/1311.4780>, 2013.

Pattyn, F.: A new three-dimensional higher-order thermomechanical ice sheet model: Basic sensitivity, ice stream development, and ice flow across subglacial lakes, *Journal of Geophysical Research*, <https://doi.org/10.1029/2002JB002329>, 2003.

Pollard, D., Chang, W., Haran, M., Applegate, P., and DeConto, R.: Large ensemble modeling of the last deglacial retreat of the West Antarctic Ice Sheet: Comparison of simple and advanced statistical techniques, *Geoscientific Model Development*, <https://doi.org/10.5194/gmd-9-1697-2016>, 2016.

Polyak, L., Alley, R. B., Andrews, J. T., Brigham-Grette, J., Cronin, T. M., Darby, D. A., Dyke, A. S., Fitzpatrick, J. J., Funder, S., Holland, M., Jennings, A. E., Miller, G. H., O'Regan, M., Savelle, J., Serreze, M., St. John, K., White, J. W., and Wolff, E.: History of sea ice in the Arctic, *Quaternary Science Reviews*, <https://doi.org/10.1016/j.quascirev.2010.02.010>, 2010.

Rasmussen, C. E.: Gaussian Processes in machine learning, *Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics)*, 2004.

Reeh, N.: Parameterization of Melt Rate and Surface Temperature on the Greenland Ice Sheet, *Polarforschung*, https://doi.org/10.1007/978-3-7091-0973-1_2, 1991.

Ritz, C., Rommelaere, V., and Dumas, C.: Modeling the evolution of Antarctic ice sheet over the last 420,000 years: Implications for altitude changes in the Vostok region, *Journal of Geophysical Research Atmospheres*, <https://doi.org/10.1029/2001JD900232>, 2001.

Sarkka, S.: *Bayesian Filtering and Smoothing*, Cambridge University Press, <https://doi.org/10.1017/CBO9781139344203>, 2013.

Sarkkä, S., Hartikainen, J., Svensson, L., and Sandblom, F.: On the relation between Gaussian process quadratures and sigma-point methods, in: *Journal of Advances in Information Fusion*, 2016.

Singarayer, J. S., Bamber, J. L., and Valdes, P. J.: Twenty-first-century climate impacts from a declining Arctic sea ice cover, *Journal of Climate*, <https://doi.org/10.1175/JCLI3649.1>, 2006.

Tarasov, L. and Peltier, W. R.: Greenland glacial history and local geodynamic consequences, *Geophysical Journal International*, <https://doi.org/10.1046/j.1365-246X.2002.01702.x>, 2002.

Tarasov, L., Dyke, A. S., Neal, R. M., and Peltier, W. R.: A data-calibrated distribution of deglacial chronologies for the North American ice complex from glaciological modeling, *Earth and Planetary Science Letters*, <https://doi.org/10.1016/j.epsl.2011.09.010>, 2012.

Thomas, E. K., Briner, J. P., Ryan-Henry, J. J., and Huang, Y.: A major increase in winter snowfall during the middle Holocene on western Greenland caused by reduced sea ice in Baffin Bay and the Labrador Sea, *Geophysical Research Letters*, <https://doi.org/10.1002/2016GL068513>, 2016.

Van Tatenhove, F. G., Van Der Meer, J. J., and Koster, E. A.: Implications for deglaciation chronology from new AMS age determinations in central West Greenland, *Quaternary Research*, <https://doi.org/10.1006/qres.1996.0025>, 1996.

Wright, P. J., Harper, J. T., Humphrey, N. F., and Meierbachtol, T. W.: Measured basal water pressure variability of the western Greenland Ice Sheet: Implications for hydraulic potential, *Journal of Geophysical Research: Earth Surface*, <https://doi.org/10.1002/2016JF003819>, 2016.

Young, N. E., Briner, J. P., Miller, G. H., Crump, S. E., Lesnek, A. J., and Thomas, E. K.: Arctic deglaciation interrupted by abrupt coolings, 5 *Science Advances*, 2019, In Review.





Figure 1. K and N mark the locations of Kangerlussuaq and Nuuk respectively. D and G mark the locations of the Dye-3 and GISP2 boreholes respectively. Northern and southern paleo-flowlines are shown as blue and red lines running left to right. The inset shows a detailed view of the modeled region. Modern bedrock elevation is expressed in meters above sea level. Historical moraines dating from 11.6 to 7.2 ka BP are indicated by labeled lines.



$$\mathbf{x}_0 = \sum_i w_i^{(m)} \chi_i$$

$$\mathcal{Y}_i = \mathcal{F}(\chi_i)$$

$$\bar{\mathbf{y}} \approx \sum_i w_i^{(m)} \mathcal{Y}_i$$

$$P_x = \sum_i w_i^{(c)} (\chi_i - \mathbf{x}_0)(\chi_i - \mathbf{x}_0)^T$$

$$P_y \approx \sum_i w_i^{(c)} (\mathcal{Y}_i - \bar{\mathbf{y}})(\mathcal{Y}_i - \bar{\mathbf{y}})^T + R$$

Figure 2. (a) White points are samples from a 2D Gaussian distribution $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, I)$. The red ellipse represents the covariance of the distribution, while red points are sigma points χ_i used for the unscented transform (UT). (b) White points are samples from the transformed, non-Gaussian distribution for $\mathbf{y} = \mathcal{F}(\mathbf{x}) + \epsilon$ with $\epsilon \sim \mathcal{N}(\mathbf{0}, R)$. Red points are transformed sigma points $\mathcal{Y}_i = \mathcal{F}(\chi_i)$. UT approximations of the mean $\bar{\mathbf{y}}$ and covariance P_y of \mathbf{y} are estimated via weighted sums of transformed sigma points.




Figure 3. (a) A subset of 6 out of a total of 45 Menegaz et al. (2011) sigma points \mathcal{P}_i representing different precipitation anomaly histories. (b) Six corresponding glacier length histories \mathcal{L}_i , or transformed sigma points, obtained by inputting the sigma points \mathcal{P}_i into the ice dynamics model \mathcal{F} .




Figure 4. The measurement mean ℓ_0 (solid black line) and 95% confidence bands (gray shaded region) for the northern flowline are estimated by generating random retreat histories with the same mean moraine formation ages and variances as the observations. The green, blue, and orange lines represent four random plausible retreat histories. Red dots denote the estimated mean moraine formation ages, while red lines show 95% confidence intervals.




Figure 5. Initial ice sheet configurations for the northern (a) and southern (b) paleo-flowlines. Solid black and blue lines represent the bedrock elevation and ice surface respectively. Moraine positions are indicated by arrows with associated ages expressed in thousands of years before present.




Figure 6. (a) Buizert et al. (2018) and Dahl-Jensen et al. (1998) mean annual temperature anomalies are shown as blue and red lines respectively. Gray and Black lines show average December, January, February and June, July, August temperature anomalies for the Buizert et al. (2018) reconstruction. Shaded blue / pink regions demarcate the extent of the HTM for the Buizert et al. (2018) and Dahl-Jensen et al. (1998) reconstructions. (b) Estimated precipitation anomaly histories for the Buizert et al. (2018) and Dahl-Jensen et al. (1998) inversions are shown by blue and red lines respectively. Solid lines show estimates for the northern flowline, while dashed lines are for the southern flowline.




Figure 7. Average precipitation (both solid and liquid) on the northern flowline from the Buizert et al. (2018) reconstruction (black line), which is based on the TraCE-21ka coupled ocean-atmosphere general circulation model of the last deglaciation, versus estimated precipitation derived in this work based on reconciling the modeled and observed retreat chronologies (blue line).




Figure 8. Modeled retreat histories for the northern (a) and southern (b) flowlines using a variety of ΔT and ΔP forcings. Blue and red lines show modeled retreat using Buizert et al. (2018) and Dahl-Jensen et al. (1998) temperature forcings respectively. Dashed lines show the results of forcing the model with $\Delta P = 0$, while solid lines show the results of using optimal estimated precipitation anomalies. Black diamonds / bars show observed terminus positions along with 95% confidence intervals. Shaded blue / pink regions demarcate the extent of the HTM for the Buizert et al. (2018) and Dahl-Jensen et al. (1998) reconstructions respectively.




Figure 9. HTM snowfall averaged across the northern and southern flowlines for the Buzert et al. (2018) (solid blue line) and Dahl-Jensen et al. (1998) (solid red line) inversions. Shaded blue / pink regions demarcate the extent of the HTM for the Buzert et al. (2018) and Dahl-Jensen et al. (1998) reconstructions respectively. Dashed lines show pre-HTM, HTM, and post-HTM average snowfall for each inversion.




Figure 10. The blue line / shaded region show the estimated mean ΔP and 95% confidence bands for the sensitivity test, accounting for uncertainty in a number of ice-flow and PDD model parameters. The black dashed line shows the mean estimated ΔP from a previous inversion assuming no uncertainty in model parameters.