

Brief communication: An alternative method for estimating the scavenging efficiency of black carbon by meltwater over sea ice

Tingfeng Dou^{1,2}, Zhiheng Du², Shutong Li², Yulan Zhang², Qi Zhang⁴, Mingju Hao⁵, Chuanjin Li², Biao Tian⁴, Minghu Ding⁴, Cunde Xiao^{3,2,4}

5 ¹College of Resources and Environment, University of Chinese Academy of Sciences, Beijing 100049, China

²State Key Laboratory of Cryospheric Sciences, Chinese Academy of Sciences, Lanzhou 730000, China

³State Key Laboratory of Land Surface Processes and Resource Ecology, Beijing Normal University, Beijing 100875, China

⁴Institute of Tibetan Plateau and Polar Meteorology, Chinese Academy of Meteorological Sciences, Beijing 100081, China

10 ⁵Ministry of Education Key Laboratory for Earth System Modeling, Department of Earth System Science, Tsinghua University, Beijing 100084, China

Correspondence to: Tingfeng Dou (doutf@ucas.ac.cn)

Abstract. The meltwater scavenging coefficient (MSC) of black carbon (BC) is a ~~key-crucial~~ parameter in snow and sea ice model, as it determines the BC enrichment in the surface layer of melting snow over sea ice and therefore modulates the BC-snow-albedo feedbacks. We present a new method for MSC estimation by sampling ~~melt-refreeze ice layer~~ice layer within
15 the snowpack and its overlying snow and measuring their physical characteristics ~~in the snowpits~~ in Elson Lagoon- northeast of Barrow, Alaska during the melting season. The bias of estimated MSC ranges from -5.4% to 7.3%, which is not exactly dependent on the degree of ablation. The average MSC value ($14.5\% \pm 2.6\%$) calculated by this proposed method is slightly lower than that derived from the repeating sampling (RS) method in Elson Lagoon, while still is within its best estimate range. Further estimation demonstrates that the MSC in Canada Basin ($23.6\% \pm 2.1\%$) is close to that in Greenland
20 ($23.0\% \pm 12.5\%$), and larger than that in Chukchi Sea ($17.9\% \pm 5.0\%$) on the northwest of Barrow. Elson Lagoon has the lowest MSC ($14.5\% \pm 2.6\%$) in the study areas. ~~It is concluded that MSC exhibited a regional difference in the western Arctic during the sampling period.~~ The method suggested in this study provides a possible approach for large-scale measurements of MSC over the sea-ice area in the Arctic, ~~of course, this method depends on the presence of a melt-refreeze ice layer in the observation area.~~

25 1 Introduction

BC is among the most efficient particulate species at absorbing visible light, which can reduce the surface albedo and potentially accelerate snow melting (Flanner et al., 2007; Goldenson et al., 2012; Dou et al., 2012; 2017). Previous studies suggested an annual-mean radiative forcing of $0.1\text{--}0.3 \text{ W m}^{-2}$ over the Arctic region from BC deposition (Flanner et al., 2009; Jiao et al., 2014). However, large uncertainties still exist in the sea ice region due to lack of field measurements and poor
30 understanding of BC enrichment by overlying snow melting.

The enrichment of BC in melting snow largely depends on MSC, as it reflects the ratio of BC concentration in the meltwater departing the snow layer to the bulk concentration in the exact layer (Flanner et al., 2007). MSC which leads to enhanced concentrations of BC in surface snow ~~has-is been found to be~~ considerably less than 100% by very few previous studies (e.g., 35 Conway et al., 1996; Xu et al., 2012; Doherty et al., 2013). In present snow and sea ice models (e.g., Flanner et al., 2007; Goldenson et al., 2012; Holland et al., 2012), MSC is valued as a constant of 20% and 3% for hydrophilic BC and hydrophobic BC, respectively, which were derived from the observations conducted at Snowdome (2050 m) of the mid-latitude Blue Glacier (Conway et al., 1996).

40 More recently, the MSC of BC was re-evaluated based on the field measurements in Elson Lagoon (Barrow, Alaska) and at Dye-2 station (Greenland) during the melting season (Doherty et al., 2013). They suggested a ~~general-rough~~ range of 10% to 30% in the study area. The method adopted in previous studies requires continuous sampling for about 2–3 weeks at each site, and thus is laborious to ~~be-used~~ apply for large-scale measurements in the polar area. Here, as an alternative, an experimental approach for calculating MSC was proposed which may provide a new way for MSC measuring, and a further 45 comparison between the regional differences of MSCs is presented as well.

The ~~melt-refreeze~~ ice layer within the snowpack results from the refreezing of meltwater that percolates into ~~the~~ snow, and thus the concentration of BC in the ice layer can represent the BC values in the meltwater departing the snow. The BC concentrations in the ~~melt-refreeze~~ ice layer and ~~in~~ its overlying snow layer were together to determine the MSC ~~associately~~ 50 considering the thickness and density of the two layers. ~~We conducted The-the~~ field measurements and sampling ~~were conducted~~ in Elson Lagoon, ~~the~~ Chukchi Sea and Canada Basin during the melt season (Fig. 1). After constraining the uncertainties of this new method, the estimated MSC ~~was-is~~ compared to ~~those values~~ derived from the RS method in the same area, and further the spatial variability of MSC in the western Arctic will be discussed.

2 Field measurements and sample analysis

55 We collected the snow samples in Elson Lagoon northeast of Barrow (Barrow expedition), in ~~the~~ Chukchi Sea (Barrow expedition) and in the Canada Basin (1st South Korean Arctic Ocean expedition) during the late spring and summer over the past decade (2010 to 2018). The snow physical characteristics (~~including the snow thickness, stratification and density~~) were also measured during the three Barrow sea ice expeditions (~~the~~ year 2015, 2017 and 2018) and the 1st South Korean Arctic Ocean expedition (~~the~~ year 2010). In the 3rd Chinese Arctic expedition (~~the~~ year 2008), only snow physics (~~thickness, stratigraphy and density~~) were observed. 60

The field measurements involve snow thickness, snow density and stratification which are conducted in Elson Lagoon, the

Chukchi Sea and Canada Basin. In Elson Lagoon, we measured the snow depth along a 10km line before melt onset (April 15, 2015), and determined the average value of snow depth. A far-shore site is chosen ~12 km away from the coast with snow depth close to the mean value (31.6 ± 5.4 cm) of this region. The site location is shown in Fig. 1. The snow stratification was firstly recorded, and then snow density was measured at 2.5 cm vertical resolution using SnowFork instrument, and four points was measured per time in each layer. We applied the average value of snow density to characterize the snow layer. The snow depth was recorded at ablation stakes next to the snow pit. In the Chukchi Sea, due to the presence of ice ridge, the spatial variation of snow depth is more significant than the Elson Lagoon. We firstly selected a relatively smooth area of sea ice, and measured the snow depth along a 200m line in the centre region of the flat ice on April 6th, 2017. The observation site was chosen at a location close to the average snow depth and the measurement procedure is the same as that applied in Elson Lagoon. Note that due to the interannual variability in the ice situation over the Chukchi Sea, there was a deviation for the observation sites in 2017 and 2018 (Fig .1). In Canada Basin, we conducted the measurements at a 100m line over floe ice to determine the average snow depth due to smaller ice size and limited operating time. Snow density was measured using a Tel-Tru densitometer (Tel-Tru Manufacturing Co., Inc., Rochester, NY) with an accuracy of 1 g, and a snow shovel of 2.5-cm in depth. The thickness of snow and the position of the melt-refreeze ice layer were measured with a ruler.

The sample collection was performed at three stages in Elson Lagoon and the Chukchi Sea during the expeditions in 2015 and 2017. At the stage before snow-melting onset, we collected snow from 4 cm above the sea ice up to the snow surface. At the early stage of melting, the upper snow layer was firstly collected, and then the underlying ice layer was sampled separately in the same snow pit. The newly fallen snow was also collected ~~during the once new~~ snowfall occurred. In order to study the spatial distribution of BC, we dug up three snow pits to sample parallelly at each site (50 meters apart from each other) and measured the physical characteristics synchronously. Observations show that the differences in BC concentrations of the three snow pits are negligible, as the standard deviation value was one order of magnitude lower than the mean concentration. We took the average BC concentration from all three pits as the BC concentration at that exact site. ~~In~~ At the end of the snow-melting season and when most of the snowpack had melted, we collected the top 4-cm layer of snow to analyze the BC concentration in the melted snow. In 2018, we just collected samples of melting snow in the Chukchi Sea. Table 1 shows More-more details ~~are showed in Table S1~~.

Sampling was performed using a pre-cleaned plastic shovel and single-used vinyl gloves. Samples were stored in polyethylene bags that had been thoroughly washed with abundant deionized ultrapure water in the laboratory ~~prior to~~ before use. In the laboratory, the snow samples were allowed to melt in ambient temperature (18–20 °C) and immediately filtered through quartz-fibre filters (25 mm, Whatman® QM-A). The filters were stored in an insulated cabinet with blue ice and kept in low temperature avoiding any bacteria to produce and transited to the laboratory in the University of Chinese Academy of Sciences for analysis.

We used two analytical methods to measure the concentration of BC. The quartz filters were firstly dried between 60 °C and 70 °C and then measured using an optical transmission analytical method (Model OT-21, Magee Scientific, California USA).
100 | The OT-21 is widely used in the measurement of atmospheric BC aerosol. ~~A~~Thereafter ~~that~~, a 1.0 cm² punch was cut from each filter, and was analyzed for elemental carbon (EC) using the “Thermos-optical NIOSH 5040” method (Sunset Laboratory Inc., Forest Grove, U.S., which has been applied to measure EC in Svalbard snow (Forsström et al., 2013). A comparison between EC and BC in a previous study (Dou et al., 2017) showed that the values obtained from two different methods are highly correlated ($R^2 = 0.97$). For consistency, we adopt BC referring to BC and EC. Five blank ~~filters~~ were
105 | processed following the same analytical procedure as the samples, except that they were filtered with ultrapure water. The measured BC background of the filters ($0.03 \pm 0.02 \text{ ng g}^{-1}$) ~~are~~ is an order of magnitude lower than the concentration of the ice layer. The values in Table ~~S1 and Table S2~~ have been corrected by excluding blank ~~contributions~~.

3 Results and discussion

During two Arctic Ocean Expeditions (the year 2008 and 2010), ice layers developed in almost all snowpacks over sea ice in
110 | the measurement area, and the snow stratigraphy and thickness exhibited highly spatial variabilities. The observed thickness of ice layers ranges from ~0.3 cm to ~2.8 cm. During the field measurements in Elson Lagoon in 2015, we recorded that the ice layer came into being on May 18th and May 22th, the early stage of the sea-ice melting season. The ice layer was observed in the Chukchi Sea on May 25th–28th, 2017, and on May 30th–31th, 2018.

115 | The ice layer results from the refrozen meltwater that percolates into cold snow along with layer-parallel capillary barriers by heat conduction into surrounding subfreezing snow (Pfeffer et al., 1998; Massom et al., 2001; Colbeck et al., 2009). It detains BC particles in the meltwater, leaving the upper snow layer. Except for the formation mechanism mentioned above, ice layers could also generate from the radiation crust or liquid precipitation re-freezing (Massom et al. 2001). However, the BC concentrations in these two types of ice layers are in the same order of magnitude as those of new or recently-fallen snow.
120 | Besides, the radiation crust usually forms on the snow surface (Colbeck et al., 2009; Dou et al., 2013). The ice layer frozen from liquid precipitation is mostly formed during winter season before the snow-melt onset (Sturm et al., 2002; Langlois et al., 2017). These two types of ice layers cannot reflect the BC scavenging with meltwater, and thus were not considered in this study.

125 | By measuring BC in the selected melt-refreeze ice layer and its overlying snow, ~~it can be drawn~~ we observed that the concentration of the ice layer is $0.42 \pm 0.08 \text{ ng g}^{-1}$ in the measurement area, ~~meaning suggesting~~ that ~0.42 ng of BC particles can be carried away from the snow layer by 1 gram water. Before estimating MSC, we compared the BC concentration ~~of in~~ the ice layer ~~is also compared~~ with those of other snow layers in the measurement area at different ablation stages. The BC

130 concentration increased from $1.32 \pm 0.20 \text{ ng g}^{-1}$ in the new snow to $2.42 \pm 0.63 \text{ ng g}^{-1}$ in the generally melting snow (Fig. S42), and the concentration in the surface layer increased up to $15.91 \pm 1.12 \text{ ng g}^{-1}$ ~~in~~ at the end of snow ablation.

The MSC is estimated based on the observations of BC, snow density and thickness. By determining the burden of BC per area (ng BC/cm^2) in the ice layer and the average original BC mass per unit area in the unmelted snowpack, the scavenging efficiency (MSC) is given by

135 ~~According to the definition by Flanner et al. (2007), MSC can be given by:~~

$$MSC = h_i \cdot \rho_i \cdot C_{bi} / h_1 \cdot \rho_1 \cdot C_{b1} \quad (1)$$

where h_i (cm), ρ_i (g cm^{-3}) and C_{bi} (ng g^{-1}) are respectively the thickness, density and BC mass concentration of the ice layer (Fig. S23); h_1 (cm), ρ_1 (g cm^{-3}) and C_{b1} (ng g^{-1}) are the same variables but for the snow layer before the melt event (Fig. S23). Note that determining scavenging efficiency with this method requires measuring the above factors at a given site at least
140 twice, before and after the melt event.

If the snow physics and BC concentration were not measured before the melt event, we would choose another method to calculate MSC. We assumed that as the surface snow melts, BC particles scavenged by meltwater are refrozen in the melt-refreeze ice layer, that is, $h_1 \cdot \rho_1 \cdot C_{b1} = h_i \cdot \rho_i \cdot C_{bi} + h_2 \cdot \rho_2 \cdot C_{b2}$, where h_2 (cm), ρ_2 (g cm^{-3}) and C_{b2} (ng g^{-1}) are
145 respectively the thickness, snow density and BC mass concentration of the melting snow overlying the ice layer (Fig. S23). ~~By determining the loading of BC per unit area (ng cm^{-2}) in the ice layer and in the partially melted snow layer above it, the scavenging efficiency (MSC) is given by:~~ So that

$$MSC = h_i \cdot \rho_i \cdot C_{bi} / (h_i \cdot \rho_i \cdot C_{bi} + h_2 \cdot \rho_2 \cdot C_{b2}) \quad (2)$$

150 ~~In fact,~~ ~~†~~ The assumption behind the proposed new method also implies that all of the melt water generated from the original snow column is conserved in the ice layer and its overlying snow. Thus, $h_1 \cdot \rho_1$ is also equal to $(h_i \cdot \rho_i + h_2 \cdot \rho_2)$ in the assumption.

155 Since the new method largely depends on the conservation of snow mass and BC content before and after the ablation event, we validate the above presumption using the observations that involve snow sampling both before and after the melt event at ~~6~~ six sites during the Barrow expeditions (Table S21). The average of the snow density and BC concentration of the whole layer of snow were used to represent the situation (ρ_1, C_{b1}) of the upper part (h_i) of the snow layer before ablation. Here, deviations from 100% conserved is used to measure the conservation of BC ($(h_i \cdot \rho_i \cdot C_{bi} + h_2 \cdot \rho_2 \cdot C_{b2}) / h_1 \cdot \rho_1 \cdot C_{b1} - 100\%$) and snow ($(h_i \cdot \rho_i + h_2 \cdot \rho_2) / h_1 \cdot \rho_1 - 100\%$), and to evaluate the uncertainty in the derived scavenging efficiencies. The loss
160 of snow mass and BC content after the ablation event are both smaller than 7.0% (Fig. 2a4a), indicating that most of the meltwater and BC within it was re-frozen in the ice layer and the BC content was substantially conserved. The assumption of the proposed new method is valid in the measurement area during the sampling period.

165 According to Eq. (2), we estimated the MSC (MSC_2) in the measurement area and compared it with the MSC_1 calculated based on Eq. (1). The result indicates that there is a slight difference in the MSCs calculated separately by the two methods. The bias of MSC $((\text{MSC}_2 - \text{MSC}_1) / \text{MSC}_1)$ caused by the deviation of snow and BC from 100% conserved before and after melt is small than 7.2% (Fig. 2b4b). Further analysis showed that there is no ~~obvious-apparent~~ correlation between the estimated bias of MSC and the degree of snow melting (Fig. 2b4b).

170 With the new method, we calculated the MSC in Elson Lagoon and compared it with that estimated by Doherty et al. (2013) in the same area. The result indicates that the MSC in Elson Lagoon is $14.5\% \pm 2.6\%$, close to the average estimation ($16.2\% \pm 8.5\%$) by repeated sampling (RS) introduced by Doherty et al. (2013) and is still within its best estimation [14.0% - 20.0%]. Our estimation of the MSC is also broadly consistent with that adopted by Flanner et al. (2007) in their model study. They assumed that the MSC is 3% for the hydrophobic BC and 20% for hydrophilic BC, given that the total BC is a
175 combination of the two types of BC.

MSC shows regional variability due to the differences in hydrophilicity and particle size of BC. BC particles are more effectively removed from the melting snow in the regions with smaller particle size and larger proportion of hydrophilic BC, corresponding to greater MSC and weaker enrichment of BC in the surface snow layer. The MSC exhibits significant spatial variability due to the different particle sizes and hydrophilicity (Flanner et al., 2007). From the observations in this study (Chukchi Sea, Elson Lagoon and Canada Basin) and the results of Doherty et al. (2013) (Elson Lagoon and Dye-2, Greenland), we investigated the spatial differences of MSC in the western Arctic. The average of the MSCs in the Canada Basin ($23.6\% \pm 2.1\%$) is basically the same as that at the Dye-2 site, Greenland ($23.0\% \pm 12.5\%$), while is larger-more significant than that of Chukchi Sea ($17.9\% \pm 5.0\%$); and Elson Lagoon has the lowest MSC ($14.5\% \pm 2.6\%$) (Fig. 35). The
180 average of the MSCs in the western Arctic is $18.0\% \pm 3.8\%$.

This study proposes a new method for large-scale measurements of MSC over the Arctic sea ice. We need to note that there may be uncertainty in the spatial difference of MSC obtained from the observations at limited sites due to random variability in BC properties during the measurement period. The estimation of MSC, in reality, requires the existence of a melt-refreeze ice layer. However, the limited data from our measurements cannot support a more extensive investigation. We reviewed the snow stratigraphy records obtained during the 3rd Chinese Arctic expedition in summer 2008 and the expedition hiking through the North Pole from 88 °N to 90 °N in late spring 1995 (Xiao et al., 1997), separately. The records show that the ice layers were widely developing over high latitudes of the Arctic Ocean, which is also confirmed by the observations in Svalbard in late spring 2007–2009 (Eckerstorfer et al., 2011). The widely distributed melt-refreeze ice layer in the Arctic suggests broader applicability for this new method in estimating the MSC of BC in the Arctic, for example, along the cruise
190 lines where it is not pragmatic to carry out long-term continuously sampling. Nevertheless, we do not rule out that a melt-

season ice layer may not form in regions of intense melt. In that case, we cannot obtain the MSC value in that region using the proposed approach.

200 This technique assumes that the refreezing process does not preferentially exclude BC, that said, the BC concentration in the ice layer will be identical to that in the melt water. In fact, some of BC impurities may be expelled during the freezing process of melt-water, thus, this assumption may result in an underestimation of the BC content in the melt-water, in turn leading to an underestimation of MSC. Besides, this method does not account for influxes of BC from snowfall during the melt season, which may also lead to an underestimation of MSC in the case of snowfall occurring after snow-melt onset.
205 Since the collected ice layer represents the event of melting water refreeze at a specific period, thus, the method provides an estimate of the average seasonal scavenging efficiency but does not capture temporal variations efficiently.

3 Conclusions

The MSC of BC ~~has-is been found to be~~ much less than 100% in ~~few~~ previously ~~few~~ studies, leading to enhanced concentrations of BC in surface snow, lowering albedo and accelerating the rate of snow melting. This study proposes a new
210 experimental approach to determine the MSC by sampling the melt-refreeze ice layer and its overlying snow in the snowpits during the melting season, assuming the complete conservation of snow and BC content before and after the ablation event. The method is different from the established methods which require repeated sampling (RS method) over an extended period. The present observations confirm that the theory adopted in the proposed method is valid in the study area and the estimation bias of the calculated MSCs is not dependent on the melting degree during the ablation.

215 Further estimation with the new method demonstrated that the MSC exhibits regional differences in the western Arctic. In the measurement period, the average MSC in Canada Basin is the largest, which is close to that estimated in Greenland, followed by those in the Chukchi Sea and ~~in~~ Elson Lagoon. The spatial difference is suggested to be considered in the future simulation of BC-in-snow over the sea ice, rather than setting MSC as a constant in the snow and sea ice model. Combined
220 with all available observations, we estimated an average of ~~-~~MSC in the western Arctic of $18.0\% \pm 3.8\%$ ranging from 13.0% to 30.0% .

Data availability. The observations of snow thickness, snow density and BC concentrations applied in this study are available as the Supplement.

225 *Author contribution.* TD designed the experiments and performed the analyses. TD, ZD, SL, YZ, QZ, MH and CL conceived field measurements and snow sampling. All authors participated in the writing of the paper.

Competing interests. The authors declare that they have no conflict of interest.

230

Acknowledgments. This study is funded by the National Key Research and Development Program of China (2018YFC1406103), the National Nature Science Foundation of China (NSFC, [41971084](#), 41425003, ~~41401079~~) and the Key Project of CAMS (KJZD-EW-G03-04). We appreciate the State Key Laboratory of Cryosphere Science of the Chinese Academy of Sciences to supply the accommodation and ice logistics support during the visit in Barrow. We also thank UIC Corporation for providing the logistic support for the field measurements over sea ice.

235

References


- Colbeck, S., Akitaya, E., Armstrong, R., Gubler, H., Lafeuille, J. and Morris, E.: The international classification for seasonal snow on the ground, International Hydrological Programme of the United Nations Educational, Scientific and Cultural Organization (UNESCO-IHP), Paris, in the series of IHP Technical Documents in Hydrology: IACS Contribution, 2009.
- Conway, H., Gades, A. and Raymond, C.F.: Albedo of dirty snow during conditions of melt, *Water Resour. Res.*, 32, 1713-1718, doi:10.1029/96WR00712, 1996.
- Doherty, S. J., Grenfell, T.C., Forsström, S., Hegg, D.L., Brandt, R.E. and Warren, S. G.: Observed vertical redistribution of black carbon and other insoluble light-absorbing particles in melting snow, *J. Geophys. Res. Atmos.*, 118, 5553-5569, doi:10.1002/jgrd.50235, 2013.
- Dou, T., Xiao, C., Shindell, D. T., Liu, J. et al.: The distribution of snow black carbon observed in the Arctic and compared to the GISS-PUCCINI model, *Atmos. Chem. Phys.*, 12, 7995-8007, doi:10.5194/acp-12-7995-2012, 2012.
- Dou, T. and Xiao, C.: Measurements of physical characteristics of summer snow cover on sea ice during the Third Chinese Arctic Expedition, *Sciences in Cold and Arid Regions*, 5, 0309-0315, 2013.
- Dou, T., Xiao, C., Du, Z., Schauer, J. J. et al.: Sources, evolution and impacts of EC and OC in snow on sea ice: a measurement study in Barrow, Alaska, *Sci. Bull.*, 62, 1547-1554, 2017.
- Eckerstorfer, M. and Christiansen, H. H.: The “High Arctic Maritime Snow Climate” in Central Svalbard. *Arct. Antarct. Alp. Res.*, 43, 11-21, 2011.
- Flanner, M.G., Zender, C.S., Randerson, J.T. and Rasch, P.J., Present-day climate forcing and response from black carbon in snow, *J. Geophys. Res.*, 112, D11202, doi:10.1029/2006JD008003, 2007.
- Flanner, M. G., Zender, C. S., Hess, P. G., Mahowald, N. M., Painter, T. H., Ramanathan, V. and Rasch, P. J.: Springtime warming and reduced snow cover from carbonaceous particles, *Atmos. Chem. Phys.*, 9, 2481-2497, doi:10.5194/acp-9-2481-2009, 2009.
- Forsström, S., Isaksson, E., Skeie, R. B. et al.: Elemental carbon measurements in European Arctic snow packs, *J. Geophys. Res. Atmos.*, 118:13614-13627, 2013.

260


- Goldenson, N., Doherty, S.J., Bitz, C.M. et al.: Arctic climate response to forcing from light-absorbing particles in snow and sea ice in CESM, *Atmos. Chem. Phys.*, 12, 7903-7920, doi:10.5194/acp-12-7903-2012, 2012.
- Holland, M., Bailey, D. A., Briegleb, B. P., Light, B. and Hunke, E.: Improved sea ice shortwave radiation physics in CCSM4: The impact of melt ponds and aerosols on Arctic sea ice, *J. Climate*, 25, 1413–1430, doi:10.1175/JCLI-D-11-00078.1, 2012.
- Jiao, C., Flanner, M.G., Balkanski, Y. and Bauer, S.E.: An AeroCom assessment of black carbon in Arctic snow and sea ice, *Atmos. Chem. Phys.*, 14, 2399-2417, doi:10.5194/acp-14-2399-2014, 2014.
- Langlois, A., Johnson, C.A., Montpetit, B., Royer, A. et al.: Detection of rain-on-snow (ROS) events and ice layer formation using passive microwave radiometry: A context for Peary caribou habitat in the Canadian Arctic, *Remote Sens. Environ.* 189, 84-95, 2017.
- Massom, R. A., Eicken, H., Haas, C., Jeffries, M. O. et al.: Snow on Antarctic sea ice, *Rev. Geophys.*, 39, 413-445, 2001.
- Pfeffer, W. T. and Humphrey, N. F.: Formation of ice layers by infiltration and refreezing of meltwater, *Ann. Glaciol.*, 26, 83-91, 1998.
- Sturm, M., Holmgren, J. and Perovich, D. K.: Winter snow cover on the sea ice of the Arctic Ocean at the Surface Heat Budget of the Arctic Ocean (SHEBA): Temporal evolution and spatial variability, *J. Geophys. Res.*, 107(C10), 8047, doi:10.1029/2000JC000400, 2002.
- Warren, S. G., Rigor, I. G., Untersteiner, N., Radionov, V. F., Bryazgin, N. N., Aleksandrov, Ye. I., and Colony, R.: Snow depth on Arctic sea ice, *J. Climate*, 12, 1814-1829, 1999.
- Xiao, C., Qin, D., Ren, J.: The feature of sea ice cover, snow distribution and its densification in the central Arctic, *Sci. Geol. Sinica*, 17, 289-296, 1997.
- Xu, B., Cao, J., Joswiak, D. R., Liu, X., Zhao, H. and He, J.: Postdepositional enrichment of black soot in snow-pack and accelerated melting of Tibetan glaciers, *Environ. Res. Lett.*, 7, doi:10.1088/1748-9326/7/014022, 2012.

Table 1: BC concentrations observed in the melt-refreeze ice layer and its overlying snow. The thickness of snow and ice layer, and snow density observed simultaneously are shown. Note that the observations before and after the ablation events in Elson Lagoon and the Chukchi Sea during Barrow expedition are shown as site1-site6. The sampling locations and dates are also shown. Refer to figure 3 for the description of variable names. ‘BC (surface melting snow)’ denotes BC concentration in the top 4-cm layer of melting snow in the end of melt season when most of the snowpack had melted.

Sampling area	Site	Lat (°N)	Lon (°W)	h_1 (cm)	ρ_1 (g/cm ³)	C_{b1} (ng/g)	Sampling date	h_i (cm)	C_{bi} (ng/g)	h_2 (cm)	ρ_2 (g/cm ³)	C_{b2} (ng/g)	Sampling date	BC(Surface melting snow, ng/g)	Sampling date	Expedition
Elson Lagoon	1	71.32	156.37	5.5	0.32	1.72	April 26, 2015	0.7	0.36	3.0	0.36	1.72	May 18, 2015	14.9	May 31, 2015	Barrow Expedition
Elson Lagoon	2	71.32	156.37	5.4	0.30	1.70	April 30, 2015	0.8	0.31	2.5	0.35	1.70	May 22, 2015	15.3	May 31, 2015	Barrow Expedition
Elson Lagoon	3	71.32	156.38	10.9	0.32	1.11	May 7, 2015	1.7	0.41	5.0	0.35	1.98	May 22, 2015	17.9	May 31, 2015	Barrow Expedition
Chukchi Sea	4	71.37	156.54	11.3	0.31	2.11	April 15, 2017	1.8	0.48	5.0	0.36	2.11	May 25, 2017	16.1	June 5, 2017	Barrow Expedition
Chukchi Sea	5	71.37	156.54	13.2	0.29	1.82	April 16, 2017	2.5	0.34	4.0	0.35	1.82	May 26, 2017	16.1	June 5, 2017	Barrow Expedition
Chukchi Sea	6	71.37	156.54	8.5	0.25	2.91	May 1, 2017	1.0	0.55	3.0	0.36	2.91	May 28, 2017	17	June 5, 2017	Barrow Expedition
Chukchi Sea	7	71.37	156.55	--	--	--	--	1.5	0.5	3.0	0.32	2.43	May 30, 2018	14.2	June 10, 2018	Barrow Expedition
Chukchi Sea	8	71.37	156.55	--	--	--	--	0.9	0.36	2.5	0.29	2.11	May 30, 2018	15.9	June 10, 2018	Barrow Expedition
Chukchi Sea	9	71.37	156.55	--	--	--	--	0.5	0.41	2.0	0.24	2.33	May 30, 2018	14.8	June 10, 2018	Barrow Expedition
Chukchi Sea	10	71.37	156.55	--	--	--	--	1.2	0.43	3.5	0.31	2.52	May 31, 2018	17.3	June 10, 2018	Barrow Expedition
Chukchi Sea	11	71.37	156.55	--	--	--	--	0.4	0.31	1.0	0.32	2.14	May 31, 2018	17.5	June 10, 2018	Barrow Expedition
Canada Basin	12	75.03	159.48	--	--	--	--	2.8	0.39	3.5	0.28	2.93	July 22, 2010	--	--	1 st South Korean Arctic Expedition
Canada Basin	13	77.98	159.64	--	--	--	--	1.7	0.54	2.5	0.31	3.81	July 26, 2010	--	--	1 st South Korean Arctic Expedition
Canada Basin	14	79.51	160.02	--	--	--	--	1.9	0.45	2.5	0.32	3.32	August 1, 2010	--	--	1 st South Korean Arctic Expedition




295 **Figure 1: Locations of snow and ice layer sampling and the measurements of snow thickness and density in this study. Barrow**
Expeditions include the field measurements carried out in the Elson Lagoon in 2015, and in the Chukchi Sea in 2017 and 2018; the
3rd Chinese Arctic Expedition was conducted over the Canada Basin and the center-centre region of Arctic Ocean in 2008; the 1st
South Korean Arctic Expedition was conducted over the Canada Basin in 2010; the North Pole Expedition refers to the first
Chinese expedition hiking through the North Pole from 88 °N to 90 °N in 1995 (Xiao et al., 1997); ~~the Svalbard Expedition was~~
conducted by Eckerstorfer et al. (2011) in the field observations in 2007–2009. The open circle indicates the point at which the ice
layer is observed. Solid triangles and squares mark the locations for both sampling and on-site measurements. ~~Cross marks the~~
location of Dye-2 where the MSC was estimated by Doherty et al. (2013).
 300




305

Figure 2: The BC concentrations in the melt-refreeze ice layer and melting snow, and its concentrations in the new snow and the surface layer of melting snow are also shown as a comparison. New-snow samples were only collected in Elson Lagoon and the Chukchi Sea during the measurement period. The box indicates the mean (upper) and median (bottom) values of the observations, and the whiskers constrain the full extent of the observations.




310

Figure 3: Conceptual sketch of snow overlying sea ice before and after the melt event. Variables relating to the snow and ice layer mentioned in Eq. (1) and Eq. (2) are shown.



315 | Figure 24: Deviations from 100% conserved for snow and BC after ablation (a), snow ablation ($h_1 \cdot \rho_1 - h_2 \cdot \rho_2$) during the melt event and the bias $((MSC_2 - MSC_1) / MSC_1) \cdot 100\%$ of estimated MSC based on Eq. (2) (b). The ticks on the X-axis are matching sites given in Table S21.



320 | Figure 35: MSC of BC in different regions over the western Arctic. Superscript ‘*’ indicates the results of this study (red), and ‘^D’
indicates the results of Doherty et al. (2013). Elson Lagoon best^D and Dye-2 best^D indicate the best-best-estimated range of MSC_z
respectively in Elson Lagoon and Dye-2, Greenland published in Doherty et al. (2013). The values of the western Arctic were
estimated based on the observations in all measurement regions, and the best-best-estimated values in Dye-2 and Elson Lagoon
were employed in the estimation. The box indicates-shows the mean (upper) and median (bottom) values, and the whiskers depict
325 | the extent.