

We thank Dr. Ron Kwok, Dr. Rasmus Tonboe and Dr. Rachel Tilling for their insightful and constructive comments and we enclose our responses below.

A 'track changes' manuscript with our proposed changes is appended to this response. Line numbers used in our responses refer to this manuscript.

For each reviewer we reproduce their comments in blue and our responses in black.

### **Response to Reviewer 1, Dr Ron Kwok:**

Equation (2), in the manuscript, is the proper way to calculate the simple quantity. Equation (1) was first introduced in Section 3b of Kwok and Cunningham [2015] – a transcription error – and corrected in subsequently publications that utilizes path length calculations: see Kwok and Markus [2017] and Kwok and Kacimi [2018]. While it is useful (for the community) to note the impact of using Equation (1), the reviewer (and author of Kwok and Cunningham [2015]) feels and requests that– if this article were to be published – it should be noted that the equations are correctly written in the subsequent publications listed above.

We are happy to include this acknowledgement and have included the additional citations (Kwok and Markus, 2017; Kwok and Kacimi, 2018) on line 63. To better acknowledge this transition, we have rephrased lines 63-64 from “Some authors ... have used this formulation” to “Some works ...”

The authors neglected, however, to note that Kwok and Cunningham [2008] first discussed seasonally varying snow density, and have used a modified seasonally varying snow density model in all of their freeboard and thickness calculations from ICESat [Kwok et al., 2009] and CryoSat-2 [Kwok and Cunningham, 2015] data sets. It is appreciated that the authors note that varying densities, though far from perfect, have been discussed though not in the same manner as that here, and are being used in thickness calculations.

We are again happy to include this discussion with the citations suggested. This has been done in lines 123-126 and line 227.

### **Response to Reviewer 2, Dr Rasmus Tonboe:**

Some of these variables such as the snow depth is affecting the radar scattering horizon and the snow-ice interface in opposite directions so that the correction for one and not the other may lead to even larger errors than doing nothing. Here I see the correction of the range for the propagation speed of microwaves in the snow to be related to the scattering horizon depth variability.

However, the magnitude of the range correction described in this MS is probably overestimated because there is evidence that the scattering horizon is not synonymous with the snow ice interface (Armitage and Ridout, 2015). The scattering horizon is more likely within the snowpack also on first-year ice because the first-year ice snow cover may be saline thus preventing penetration into the bottom snowpack (Nandan et al., 2017). I think that a short discussion of that should be included.

This is a significant point which is worthy of acknowledgement in the paper, and is similar to the first point raised by Reviewer 3 (Dr Rachel Tilling). We have now ammended the manuscript, principally adding Subsection 4.3 (lines 189 – 199). Here we state that

remedying the biases analysed may move sea ice thickness estimates further from their true (and presently unknown) values. We have also ammended line 8 (in the abstract) to remove the implication that fixing these biases will definitely improve the accuracy of estimates.

While it was stated in the paper that this work uses the assumption of full radar wave penetration, the impact of this assumption on the results was not stated and is now summarised in lines 209-210.

Also today's snow depth compared to the modified Warren climatology which is used for estimating the magnitude of the range correction should be included in the discussion.

This discussion has now been included in Subsection 4.4 (lines 201-210). Since snow depths over MYI are likely lower now than in the modified W99, sea ice thickness is likely overestimated in this regard. This introduces a similar issue to that discussed directly above.

### Specific Comments

P1, L3: "This implies that that the scattering horizon is synonymous with the snow-ice interface. However there is evidence that the scattering horizon is above the snow ice interface especially if the snow is saline. This depth (scat. horiz.) is not well known, so how to apply the correction?"

We have ammended line 3 to clarify that the assumption of full snowpack penetration is invoked in publicly available sea ice thickness products. However, we now acknowledge in Subsection (4.3) that this assumption has not held in numerous investiagtions (e.g. Nandan et al., 2007; Willatt, et al., 2009; Willatt, et al., 2011; King et al., 2018).

We believe the work in this manuscript to qualitatively hold for lower snow penetration depths induced by a raised scattering horizon, although the size of the biases would be reduced. We have now pointed this out in lines 209-210. In this case, the correct equation for the propagation correction should still be selected and seasonal densification would still take place and should be accounted for. To address the reviewer's point, we now discuss the effect of an elevated scattering horizon in Section 4.3.

P1,L6: "winter ice" is sometimes synonymous with "first-year ice", move "in winter" to the end of the sentence to avoid confusion.

We accept this suggestion and have rearranged the sentence accordingly.

P1, L18: less snow gives more potential for ice growth, increasing temperatures the opposite. This sentence is contradicting.

In combination with feedback from Reviewer 3 (Dr. Rachel Tilling) that this paragraph is unnecessary long and dense with information, we have removed this sentence entirely.

P6, L135:The NP is normally not covered by satellites and so it is not a good spot for comparison or verification.

We have recalculated a representative densification rate based on a spatial average of the Arctic Basin shown in supplementary figure (S2). This newly calculated rate is slightly

higher than that calculated for the North Pole (6.50 vs 6.45 kgm<sup>-3</sup>/year), and we have repeated our analysis and updated our figures with this slightly higher value.

#### P10, L197: How do you know the depth of penetration?

Currently the depth of penetration is not well known and as such is assumed to be total in publicly available sea ice thickness products. To avoid the implication that radar range estimates can be simply corrected for this issue, we have ammended this sentence to read:

*“We suggest this is done before further work is undertaken to estimate the extent of and incorporate the effects of partial radar wave penetration into the snow cover”*

#### Responses to Reviewer 3, Dr Rachel Tilling:

##### Major Comments

*“The authors’ statements about improving the accuracy of sea ice thickness estimates are simplistic and misleading. In the abstract they state that “Correcting these biases would improve the accuracy of sea ice thickness products” and this is echoed throughout the text. This conclusion doesn’t account for opposing biases that also exist.”*

The reviewer highlights several features of SIT retrieval algorithms that introduce overestimating biases. If SIT is indeed overestimated on the whole, then correcting the underestimating biases highlighted in this manuscript will make SIT estimates larger, further removing them from the true value. We therefore agree that the statement “Correcting these biases would improve the accuracy of sea ice thickness products” (L8) is incorrect if this is the case.

We have rephrased the final sentence of the abstract (L8) from:

*“Correcting these biases would improve the accuracy of sea ice thickness products, which feed a wide variety of model projections...”*

To:

*“Correcting these biases would impact a wide variety of model projections, calibrations, validations and reanalyses.”*

We have also included a new subsection (4.3) to discuss the overestimating biases introduced by incomplete radar wave penetration of the snowpack and to acknowledge that if SIT is currently overestimated then fixing these biases may not make SIT retrievals closer to the true value.

Finally, we have changed the wording in the Summary (Subsection 4.6). Rather than referring to how much sea ice thickness is underestimated, we now refer to the biases that are introduced by the treatments examined in our analysis (lines 236 and 242).

*It should be clearer that the study is only concerned with the impact of evolving snow density on the radar propagation correction, and not the conversion of sea ice freeboard to thickness (for which all groups apply an evolving snow density). This is suitably explicit in the title of section 3 and a couple of places in the text, but not*

throughout.

We have now stated this explicitly in lines 127 and 163.

### Minor Comments

Introduction: Unnecessarily dense with information. The first two paragraphs could be condensed and combined.

We have removed information from the first two paragraphs and combined them. This is visible in lines 17-22 of the track changes document.

P1L6: Rearrange for absolute clarity that 15 cm applies to sea ice thickness, not growth rate.

We have rearranged and added the value for growth rate bias (to distinguish from absolute SIT bias).

P2L35: Reference needed

This was an error and the propagation correction is in fact of comparable magnitude (ratio 0.25 : 0.29 for  $300\text{kgm}^{-3}$  snow). This has been ammended and illustrated in Section 1 of the supplement.

Figure 1 (a) and (b): Larger text for numbers and y-axis labels

This figure has been updated with larger x and y axis labels and ticks.

P8L154: "...\*\*effectively\*\* setting the rate to zero \*\*for the radar range correction\*\* introduces..."

This line has now been ammended.

P8L155-157: Again, make it clear that these calculations do account for seasonal variation in snow density, even though they will still be sensitive to uncertainties in the density assumptions.

Line 165 now includes clarification that the evolving snow density is on a "seasonal" scale with respect to the 'snow loading correction'.

# Brief Communication: Conventional assumptions involving the speed of radar waves in snow introduce systematic underestimates to sea ice thickness and seasonal growth rate estimates

Robbie D.C. Mallett <sup>1</sup>, Isobel R. Lawrence <sup>1</sup>, Julianne C. Stroeve <sup>1,2,3</sup>, Jack C. Landy <sup>4</sup>, and Michel Tsamados <sup>1</sup>

<sup>1</sup>Centre for Polar Observation and Modelling, Earth Sciences, University College London, London, UK

<sup>2</sup>National Snow and Ice Data Center, University of Colorado, Boulder, CO, USA

<sup>3</sup>Centre for Earth Observation Science, University of Manitoba, Winnipeg, Canada

<sup>4</sup>School of Geographical Sciences, University of Bristol, Bristol, UK

**Correspondence:** Robbie Mallett (robbie.mallett.17@ucl.ac.uk)

## Abstract.

Pan-Arctic sea ice thickness has been monitored over recent decades by satellite radar altimeters such as CryoSat-2, which emit Ku-band radar waves that are ~~conventionally assumed~~ assumed in publicly available sea ice thickness products to penetrate overlying snow and scatter from the ice-snow interface. Here we examine two expressions for the time delay caused by slower radar wave propagation through the snow layer and related assumptions concerning the time-evolution of overlying snow density. Two conventional treatments lead to systematic underestimates of ~~winter ice thickness~~ ice thickness of up to 15 cm and thermodynamic growth rate of up to ~~15-10~~ cm over multiyear ice in winter. Correcting these biases would ~~improve the accuracy of sea ice thickness products, which feed~~ impact a wide variety of model projections, calibrations, validations and reanalyses.

## 10 1 Introduction

Sea ice is a key moderator of the global climate system, limiting the exchange of heat, moisture and momentum fluxes between the ocean and the atmosphere. It also plays a crucial role in ocean circulation and Arctic Ocean primary productivity (e.g. Sévellec et al., 2017; Chan et al., 2017). During autumn, open water areas form new ice that can grow thermodynamically by 1.5 to 2.5 m over a winter season. ~~Ridging and rafting can locally increase the ice thickness and, if the ice survives the following melt season, further~~ Further deformation and thermodynamic ice growth can lead to thicknesses in excess of 5 m.

Today the Arctic is undergoing a period of profound transformation, with the area and thickness of the floating sea ice cover in rapid decline (e.g. Stroeve and Notz, 2018; Kwok, 2018). These changes are ~~partially a result of a distinct change in seasonality in the Arctic with the melt season starting earlier and lasting longer, enhancing the ice-albedo feedback (Stroeve and Notz, 2018). Delays in freeze-up have reduced both the amount of snow that accumulates on the sea ice over winter (Stroeve et al., 2019; In~~ Revision), and together with recent increases in driven by a variety of factors including later freeze-ups, earlier melt onsets and

increased winter air temperatures, ~~have reduced the rate of thermodynamic ice growth (Graham et al., 2017; Stroeve et al., 2018)~~  
~~Winter (Graham et al., 2017; Stroeve et al., 2018).~~

As well as being a sensitive indicator of climate change, winter sea ice thickness also functions as a prognostic variable in the polar climate system, affecting the amount and distribution of sea ice that will survive the summer melt season. Accurate  
25 knowledge of sea ice thickness is particularly important where data are assimilated into forecasting systems and other complex models which often exhibit sensitive dependence on initial conditions (Day et al., 2014).

Sea ice thickness has been observed through various methods including submarines, ice mass-balance buoys, electromagnetic induction sounding and satellite laser and radar altimetry (e.g. Schweiger, 2017; Kwok, 2018). The CryoSat-2 mission has played a leading role over the last decade, providing radar ranging observations from which the sea ice thickness may be  
30 derived (Wingham et al., 2006; Laxon et al., 2013; Tilling et al., 2018).

Ku-band radar altimeters such as CryoSat-2 and Sentinel-3 do not directly measure sea ice freeboard, but instead measure 'radar freeboard' through a time-of-flight calculation. The radar freeboard is the difference in radar ranging between the snow-ice interface and the local, instantaneous sea level (assuming perfect radar wave penetration through the snowpack). Since the radar wave speed is reduced in snow, a priori knowledge of the snow depth and density is required to convert the radar  
35 freeboard to the true ice freeboard. Following the freeboard calculation, sea ice thickness can then be estimated through the assumption of hydrostatic equilibrium (e.g. Laxon et al., 2003). This again requires a priori knowledge of snow depth and density to account for freeboard reduction due to the weight of overlying snow. The magnitude of freeboard correction for the weight of overlying snow is typically twelve times that for the effect of slower wave propagation in snow.

An important consideration in the conversion of radar freeboard ( $F_r$ ) to ice freeboard ( $F_i$ ) and in turn ice thickness is therefore the time delay due to slower radar pulse propagation in snow (Kwok, 2014). In this study we highlight two different  
40 approaches to the calculation of this time delay used in published literature. Correct handling of this time delay has a significant impact on the retrieval of sea ice thickness and volume from radar altimetry, as we show here. This is particularly the case as snow settles and densifies over the winter season.

We further investigate the impact of assuming a fixed snow density throughout winter when calculating this time delay. At  
45 present no groups producing publicly available sea ice thickness products from CryoSat-2 factor monthly evolution of snow density into their correction for slower radar wave propagation in snow, despite often including an evolving density in their calculation of the floe's hydrostatic equilibrium. The impact of this assumption is assessed and found to produce underestimates of the rate of winter thermodynamic sea ice growth, with October-April growth currently being underestimated by over 10 cm over multiyear ice.

## 50 2 Different Treatments of the Radar Propagation Correction

The correction to the radar range to account for slower radar wave propagation in snow,  $\delta h = F_i - F_r$ , is often expressed as the product of snow depth,  $Z$ , and some function of wave velocity in snow,  $f(c_s)$  (e.g. Tilling et al., 2018; Kwok, 2014) such that:

$$\delta h = Z \times f(c_s) \quad (1)$$

We now present a short derivation of  $f(c_s)$  and thus  $\delta h$  through consideration of the extra time taken,  $\delta t$ , for a radar wave to travel a distance  $Z$  through a specified snow depth rather than through free space. The time delay induced by the snow layer is expressed:

$$\delta t = t_{snow} - t_{vacuum} \quad (2)$$

$$\delta t = Z/c_s - Z/c \quad (3)$$

$$\delta t = Z(1/c_s - 1/c) \quad (4)$$

60 Where  $c_s$  the wave speed in snow, and  $c$  is the radar wave speed in free space ( $3 \times 10^8 \text{ ms}^{-1}$ ). To convert this time delay ( $\delta t$ ) into a path difference ( $\delta h$ ), one multiplies by the speed of the wave in free space:

$$\delta h = \delta t \times c = Z(c/c_s - 1) \quad (5)$$

~~Tilling et al. (2018) use~~ Some works (Tilling et al., 2018; Kwok and Markus, 2018; Kwok and Kacimi, 2018) have used this formulation to correct the radar range for the slower wave propagation speed through snow. ~~Some authors~~ Other works have used an alternative form of Eq. (5), generated by multiplying  $\delta t$  in Eq. (4) by the wave speed in snow (Kwok, 2014; Kurtz et al., 2014; Kwok and Cunningham, 2015; Ricker et al., 2015; Armitage and Ridout, 2015; Hendricks et al., 2016; Landy et al., 2017; Xia and Xie, 2018):

$$\delta h = Z_r(1 - c_s/c) \quad (6)$$

For Eq. (6) to be true,  $Z_r$  must be regarded as:

$$70 \quad Z_r = Z(c/c_s) \quad (7)$$

However,  $Z_r$  is conventionally interpreted as the real snow depth ( $Z$ ) and  $\delta h$  is therefore erroneously reduced by a factor of  $c_s/c$ . When Eq. (7) is incorporated into Eq. (6),  $\delta h$  is redefined in terms of  $Z$  and becomes Eq. (5).

Conventional interpretation of  $Z_r$  as the real snow depth therefore leads to a bias in the freeboard ( $B_f$ ) where:

$$B_f = Z \times \frac{(c - c_s)^2}{c \times c_s} \quad (8)$$

75 Bias in the freeboard then propagates into estimates of sea ice thickness by a multiplicative factor of  $\rho_w/(\rho_w - \rho_i)$ , where  $\rho_w$  represents the density of seawater and  $\rho_i$  represents the density of sea ice. Because first year ice (FYI) is generally denser than multiyear ice (MYI), a fixed snow thickness will cause a greater bias on the thickness of first year ice. However, typical biases over FYI are generally expected to be lower due to reduced snow accumulation. The bias in sea ice thickness ( $B_{SIT}$ ) due to conventional, erroneous use of Eq. (6) is therefore:

$$80 \quad B_{SIT} = Z \times \frac{(c - c_s)^2}{c \times c_s} \times \frac{\rho_w}{\rho_w - \rho_i} \quad (9)$$

Equation (9) illustrates that the bias grows linearly with snow depth. In addition to this,  $B_{SIT}$  is also dependent on the speed of the radar wave in snow, which is itself a function of snow density. Several empirical relationships have been proposed for the relationship between snow density and radar wave speed, however the most commonly used three (Hallikainen et al., 1982; Tiuri et al., 1984; Ulaby et al., 1986) deviate negligibly from each other in the typical density range for snow observed on  
85 Arctic sea ice (Fig. S1). In this investigation, we use the relationship from Ulaby et al. (1986):

$$c_s = c(1 + 0.51\rho_s)^{-1.5} \quad (10)$$


As snow density increases,  $c_s$  decreases and  $B_{SIT}$  increases. This positive relationship between  $f(c_s)$  and snow density is shown in Fig. 1a. Because both snow depth and snow density generally increase throughout the season as snow accumulates, compacts and settles, any  $\delta h$  generated through incorrect expression of  $f(c_s)$  becomes increasingly underestimated.

90 Furthermore,  $B_{SIT}$  increases even as a fixed snow water equivalent densifies and shrinks in volume. This is because  $B_{SIT}$  scales more rapidly with increasing snow density than it reduces with decreasing snow depth. The increase in bias with snow density for constant SWE is illustrated in Fig. 1b.

Since  $B_{SIT}$  is explicitly a function of snow depth and implicitly a function of snow density via Eq. (10), its spatial mapping requires the use of an Arctic snow distribution. Here we use snow depths and densities from Warren et al. (1999) (henceforth  
95 ‘W99’) to illustrate these underestimates. To be consistent with current data products that rely on W99 for their snow depth distribution, we halve snow depths over first-year ice as per Laxon et al. (2013) and only consider the Central Arctic basin (see Fig. S2) where W99 is considered most reliable (Kwok and Cunningham, 2015). Data on sea ice type and extent were taken from the sea ice type product of the EUMETSAT Ocean and Sea Ice Satellite Application Facility (OSISAF; Aaboe et al., 2016).

100 We find that where sea ice thicknesses are calculated using W99 snow depths and densities in the Central Arctic, thickness underestimates increase throughout the winter to values exceeding 15 cm in April over multi-year ice (Fig. 1c). Over FYI the mean bias increases from 4.2 cm in October to 9.8 cm in April (compared to 6.4 cm and 13.6 cm for MYI). In April, 28% of MYI has a bias exceeding 15 cm, and 7% exceeds 16 cm.

How does this bias impact sea ice thickness products currently available to the science community? Most commonly-used  
105 products do not correct for slower wave speed using the W99 density distributions in time or space, but instead use a reference



**Figure 1.** (a) Difference between conventional use of Eq. (6) and Eq. (5) as a function of snow density. This bias increases with snow density, ultimately exceeding a factor of 0.1 of the snow depth for dense snow. (b) Sea ice thickness bias for a fixed mass of snow increases as it densifies and contracts with time. Solid lines indicate bias for first year ice, dashed lines for multiyear ice assuming fixed densities of  $916.7$  and  $882 \text{ kg m}^{-3}$  respectively. (c) Monthly thickness bias introduced by conventional and erroneous use of Eq. (6) when calculated using W99 density and depth distributions. Pixels are only displayed where sea ice type is known in all years 2010-2018. Black line indicates region where multiyear ice is present in over 50% of years. Monthly averages derived from years 2010-2018.

density to calculate a fixed value for  $f(c_s)$  in Eq. (1). This value is fixed not only across the Arctic basin, but throughout the winter. In the CryoSat-2 sea ice thickness product from the Alfred Wegener Institute (AWI; Hendricks et al., 2016),  $f(c_s)$  is taken as  $(1 - c_s/c)$  as in Eq. (6). Citing the reference spring snow density given by Kwok (2014) of  $350 \text{ kg m}^{-3}$ , they generate a fixed  $\delta h$  of  $0.22Z$ .

110 On the other hand, the Centre for Polar Observation and Modelling (CPOM) takes  $f(c_s)$  to be  $(c/c_s - 1)$  (Eq. (5); Tilling et al., 2018). However, the CPOM product uses a lower reference density of  $300 \text{ kg m}^{-3}$  (taken from (Kwok et al., 2011)), generating a reference  $\delta h$  of  $0.25Z$ . AWI's use of a higher reference density mitigates the difference introduced by their

erroneous interpretation of Eq. (6). Were AWI to use a similar reference density to CPOM's  $300 \text{ kg m}^{-3}$  with Eq. (6), their reference  $\delta h$  would be  $0.19Z$ , contrasting starkly with CPOM's  $0.25Z$ .

115 The decision to use a fixed snow density for the wave-speed propagation correction throughout the winter introduces biases of its own with regard to the rate of thermodynamic growth; this is discussed in the next section.

### 3 Impact of Seasonal Snow Density Evolution on the Radar Wave Propagation Correction

Despite recent developments in pan-Arctic scale snow density modelling (Petty et al., 2018b), the Arctic snow density distribution remains poorly constrained in time and space. Because of this, representative values for pan-Arctic average snow density are often combined with the snow depth distributions from W99 to calculate the radar wave propagation correction (Kurtz et al., 2014; Hendricks et al., 2016; Tilling et al., 2018).

This constant value contrasts with the ubiquitous inclusion of density evolution in the adjustment to an ice floe's hydrostatic equilibrium due to the weight of overlying snow. A density evolution curve was derived from W99 by Kwok and Cunningham (2008) and implemented in sea ice thickness estimates derived from ICESat and CryoSat-2 (Kwok et al., 2009; Kwok and Cunningham, 2015).  
125 It is notable that Kwok and Cunningham (2015) include density evolution in both their calculation of the propagation correction and the adjustment to hydrostatic equilibrium.

To investigate the impact of an evolving snow density on ~~freeboard conversion~~the propagation correction, we calculated the ~~wave-speed~~propagation correction over Arctic sea ice by two methods: The control method used a fixed reference density in the wave speed correction (i.e.  $300 \text{ kg m}^{-3}$ ) as done by CPOM and AWI. The other method incorporated a rate of snow densification obtained from the ~~Central Arctic data published in~~ W99 in the Central Arctic.

The control method used the parameters employed by Tilling et al. (2018) producing a radar wave speed in snow of  $2.4 \times 10^8 \text{ m s}^{-1}$  corresponding to a reference density of  $300 \text{ kg m}^{-3}$  when converted using Eq. (10). As discussed in Sect. 2, estimates of absolute sea ice thickness are sensitive to the choice of reference snow density. However, the estimated rate of thermodynamic growth (the focus of this section) is more responsive to the density's time derivative, which for a fixed value ( $\rho_s = 300 \text{ kg m}^{-3}$ ) is zero. As such, our results with respect to growth rate are applicable to different reference densities such as those used by AWI ( $350 \text{ kg m}^{-3}$ ) and the NASA Goddard Space Flight Center ( $320 \text{ kg m}^{-3}$ ; Kurtz et al., 2014).

For the 'evolving' method, we calculated a representative winter (Oct-Apr) densification rate using the average densification rate of snow ~~at the North Pole over the Arctic Ocean~~ given by W99. This was found to be approximately ~~+6.45~~-6.50  $\text{kg m}^{-3}$  per month. The October starting density was taken as the spatial average of the W99 October ~~North Pole density density field~~  
140 over the same region - this choice served to minimise sea ice thickness bias at the start of the growth season and better enable comparison of growth rate. Snow density in the 'evolving' method can therefore be written as:

$$\rho_s = \underline{6.45} \underline{6.50} t + \underline{275.3} \underline{274.51} \tag{11}$$

Where  $t$  represents the number of months since October.

The ~~North Pole was chosen for two reasons: its density evolution can be trivially read from the published data in W99 and it suffers least from edge effects due to the quadratic fitting.~~ To further justify this choice, the W99 snow density evolution of  
145

five Arctic regions were also examined and found to be similar to the [North Pole basin-wide](#) rate, with the exception of the Laptev Sea which shows only a small (but positive) seasonal densification rate (Fig. S3). As in Sect. 2, we halved the W99 snow depths over FYI and only analysed the Central Arctic basin where W99 is considered most reliable.

When the evolving density shown in Eq. (11) was included in our calculation of the radar wave propagation correction, we found sea ice thickness to grow on average by an extra 10.1 cm between October and April over MYI. This corresponds to an extra 1.7 cm per month when compared to a fixed  $f(c_s)$  of  $0.25Z$ . Density evolution caused FYI to grow an extra 6.4 cm over the same time period, corresponding to an extra 1.1 cm per month.

Given the poor state of knowledge concerning the current distribution of pan-Arctic snow densities and the difficulty in collecting in-situ data, we cannot conclude whether this increased growth should correspond to higher-than-previous thicknesses at the end of winter or lower-than-previous thicknesses at the start of winter. Put another way, in this section we show a systematic bias in the [thermodynamic](#) growth rate rather than absolute [ice](#) thickness values.

Having illustrated the effect of snow densification on the radar wave propagation correction, we now justify its inclusion.


While the absolute values for regional mean densities have conceivably changed since the data was collected for W99, it remains almost certain that snow density still increases over winter for the majority of the Arctic basin as documented in W99. Furthermore, the rate of snow densification shown in W99 is likely now underestimated, with field observations indicating densification rates of  $>20 \text{ kg m}^{-3}$  per month on FYI (Langlois et al., 2007) and FYI now occupying significantly more of the Arctic basin than in the 1954-91 period over which W99 was compiled (Stroeve and Notz, 2018). While significant uncertainty in the true densification rate exists, [effectively](#) setting the rate to zero [for the radar wave propagation correction](#) introduces a systematic bias in sea ice thickness calculations.

Finally, commonly used products (e.g. Tilling et al., 2018; Hendricks et al., 2016) have included [a seasonally evolving](#) snow density in the ‘snow loading correction’ (for change in the hydrostatic equilibrium of the floe due to the weight of snow cover), which features a very similar sensitivity to uncertainty in snow density (Fig. S4).

## 4 Discussion

### 4.1 Different Fixed Densities


To further explore this issue, we calculated the expected difference between sea ice thickness estimates from CPOM and AWI introduced by their usage of  $\delta h = 0.25Z$  and  $\delta h = 0.22Z$  respectively. Since the difference in  $\delta h$  is partially due to different choices of a representative snow density, resulting sea ice thickness differences cannot be seen as bias from a true value until Arctic snow densities are better constrained. This variation is superimposed on the bias introduced by fixed snow densities discussed above. We find that CPOM’s higher value for  $f(c_s)$  produces a higher mean MYI thickness of 5cm in November, growing to 7cm by April. 16% of MYI exhibited a difference of  $> 8$  cm. For FYI, the mean difference is 2.8 cm in November and grows to 4.7cm by April (Fig. S5).



**Figure 2.** Monthly biases in sea ice thickness due to the effect of ignoring snow densification in calculating propagation correction (a) Spatially averaged histograms indicating the area of ice subjected to a given bias. Data separated into pixels that feature MYI for that month in more/less than 50% of years 2010-2018. Pixels that typically feature MYI experience greater bias in all months, largely due to halved W99 snow depths over FYI. (b) Bias maps illustrating sea ice thickness biases. Pixels are only displayed where sea ice type is known in all years 2010-2018, so bias is not displayed in some areas of ambiguous ice type. Black line indicates region where MYI is present in over 50% of years.

#### 4.2 Comparison to Radar Freeboards

To investigate these biases further, we compare them by converting pan-Arctic CryoSat-2 radar freeboard retrievals from late 2010 to early 2018 from Landy et al. (2019, In Review) to estimates of sea ice freeboard using:



**Figure 3.** Percentage bias in sea ice freeboard. The bias induced by two effects was compared to the ice freeboards from Landy et al (2019, In Review). (a) Percentage bias introduced by the use of Eq. (5) vs Eq. (6) when combined with the W99 fits for depth and density. As a fraction of the growing ice freeboard, biases remain relatively constant, indicating they grow at the same rate. (b) Percentage bias introduced by an evolving snow density derived from [the densification rate at the North Pole W99 data](#). This bias increases as a fraction of the ice freeboard from 2.3% to >6%, indicating that thermodynamic growth rates are underestimated.

- 180
- Equation (5) versus Eq. (6) (with conventional, erroneous interpretation) using the depth and density fits from W99
  - A monthly evolving density versus the fixed density used in Hendricks et al. (2016), both with spatially constant density across the Arctic basin

We find that the bias induced by the conventional, erroneous interpretation of Eq. (6) remains relatively constant as a fraction of the sea ice freeboard at around 6% (despite increasing in an absolute sense) (Fig. 3a).

185 We find that the bias induced by the assumption of a non-evolving snow density (in calculation of the propagation [offset correction](#)) grows throughout the season relative to the sea ice freeboard and in an absolute sense. The bias grows from 2.3% to 6% of the ice freeboard (Fig. 3b), indicating that the growth rate is underestimated when a fixed density is assumed.

### 4.3 [Incomplete Radar Wave Penetration of the Snowpack](#)

190 [The biases introduced in this analysis are derived based on the common assumption that Ku-band radar waves penetrate the entire snowpack. However, in-situ studies of Antarctic snow on sea ice indicate that snow with significant morphological features can scatter the radar above the snow-ice interface \(Willatt et al., 2009\). Airborne investigations during the CryoVEx and N-ICE2015 campaigns also revealed elevated dominant scattering horizons \(Willatt et al., 2011; King et al., 2018\). Furthermore, snow salinity has also been shown to elevate the dominant scattering horizon from the snow-ice interface. Nandan et al. \(2017\) found the horizon to be elevated by 7 cm based on FYI data from the Canadian Arctic.](#)

195 [Radar wave scattering from a horizon above the ice-snow interface introduces an overestimating bias on sea ice freeboard and thickness. The size of this bias is potentially larger than those discussed above, and may be dominant in determining the sign of the overall bias. If this is the case and sea ice thickness is overestimated overall, fixing the underestimating biases discussed in this analysis would shift estimates further away from the true value. As such, while improving the realism of the retrieval algorithm, the results may not become more accurate.](#)

### 200 4.4 [Snow Depth Decline Since W99 Collection](#)

[The climatology assembled by Warren et al. \(1999\) was collected from drifting ice stations largely over MYI in the period 1954-91. Since then the average age of MYI has declined and freeze-ups have become increasingly delayed \(Stroeve and Notz, 2018\). This has had the effect of decreasing snow depth over MYI \(Webster et al., 2014\). While W99 has been modified to better apply over FYI using comparatively recent Operation Ice Bridge data \(e.g. Laxon et al., 2013; Webster et al., 2014\), this has not been similarly carried out for MYI snow depths in this analysis or other publicly available products. As such, the snow depths conventionally used for thickness retrievals are likely overestimates over MYI and this introduces an overestimating bias on freeboard and sea ice thickness. This would add to the effect described in Sect. \(4.3\), where fixing underestimating biases may not make the overall estimate closer to the truth.](#)

210 [Furthermore, lower snow depths and/or incomplete radar wave penetration of the snowpack would decrease the magnitude of the biases described here \(as Eq. \(8\) and Eq. \(9\) both scale linearly with snow depth\).](#)

### 4.5 **Broader Implications**

Sea ice thickness is closely tied to sea ice volume, a sensitive indicator of climate change but also a quantity of major interest for the modelling community. The thickness underestimates highlighted in Sect. 2 have some impact on total sea ice volume, although this is well within the currently large uncertainty bounds. Nonetheless, we argue that these uncertainty bounds have been systematically biased low through conventional use of Eq. (6) in some products.

In addition, the fact that these underestimates grow over winter means the seasonal growth rate is also underestimated through conventional use of Eq. (6). While the rate of winter sea ice growth is still uncertain and interannually variable, the use of a fixed, seasonally-constant value for the snow density will bias growth rates low.

220 Accurate characterisation of thermodynamic growth is important to a variety of systems. A higher growth-rate will impact the surface salinity balance as more freshwater than previously estimated is locked up in sea-ice during thermodynamic growth and then ejected to the mixed-layer when ice melts in summer. The rate of sea ice growth is an important variable in the characterization of the negative conductive feedback (thin ice thickens faster: Stroeve et al., 2018; Petty et al., 2018a). Finally, end of winter sea ice thickness moderates subsequent light transmittance through the ice, impacting under-ice ecosystems and related geochemical processes (Nicolaus et al., 2012).

225 Sea ice thickness products featuring the misinterpretations of Eq. (6) have fed several forecast and reanalysis models (e.g. Xia and Xie, 2018). Thickness products featuring a constant-density assumption [built into the propagation correction](#) are near-ubiquitous ([with the exception of Kwok and Cunningham \(2015\)](#)) and have also fed forecast and reanalysis models (e.g. Yaremchuk et al., 2019; Blockley and Peterson, 2018). While these biases may be small compared to the effects of partial radar [wave penetration into the snow cover as a function of snow pack variables \(e.g. salinity, wetness, temperature\)](#) [snowpack](#),  
230 they are ~~remediable and can be simply applied~~ [simply remediable. We suggest this is done](#) before further work is undertaken to ~~correct radar range estimates for partial penetration of the wave~~ [estimate the extent of and incorporate the effects of partial penetration radar wave penetration](#) into the snow cover.

#### 4.6 Summary

We investigated two conventional methods for correcting radar altimetry based sea ice freeboard retrievals for slower radar  
235 wave propagation in snow. We found that a commonly used treatment (conventional use of of Eq. (6)) for this correction ~~leads to~~ [introduces an](#) initial and seasonally-increasing ~~underestimation of~~ [underestimating bias on](#) sea ice thickness from October through to April. While most commonly-used products then transform this bias (where present) by choosing a fixed snow density, we find underestimation of April sea ice thickness to exceed 15 cm over some multiyear ice when this treatment is applied in conjunction with the snow climatology from Warren et al. (1999).

240 We also investigated the impact of assuming a seasonally-fixed snow density on the radar wave propagation correction. While uncertainties in the absolute value of Arctic snow density preclude any conclusion about whether sea ice thickness is being under- or overestimated in this respect, [this treatment is found to introduce an underestimating bias on](#) the thermodynamic growth rate of multiyear ice ~~is found to be underestimated by of~~  $\sim 1.7$  cm per month leading to a  $\sim 10.1$  cm ~~underestimate in~~ [growth bias](#) over the October-April period.

245 While these biases ~~in on~~ sea ice thickness (Sect. 2) and growth rate (Sections 2 & 3) [retrievals](#) are small compared to the total uncertainty, they are systematic and influence the uncertainty bounds. These biases also propagate into derived products and model projections, calibrations and reanalyses.

*Code and data availability.* [https://github.com/robbiemallett/snow\\_density\\_assumptions](https://github.com/robbiemallett/snow_density_assumptions)

*Author contributions.* RDCM carried out the analysis and wrote the manuscript, with continued input from all authors. In addition to  
250 manuscript input, JCS and JCL contributed data to aid analysis and MCT contributed to the processing code.

*Competing interests.* The authors declare no competing interests.

*Acknowledgements.* This work was funded primarily by the London Natural Environmental Research Council Doctoral Training Partnership  
grant (NE/L002485/1). JCL acknowledges support from the European Space Agency Living Planet Fellowship ‘Arctic-Summit’ under grant  
ESA/4000125582/18/I-NS and the Natural Environmental Research Council Project ‘Diatom-ARCTIC’ under Grant NE/R012849/1. MT  
255 acknowledges support from the European Space Agency in part by project ‘Polarice’ under grant ESA/AO/1-9132/17/NL/MP and in part by  
the project ‘CryoSat + Antarctica’ under Grant ESA AO/1-9156/17/I-BG.

## References

- Aaboe, S., Breivik, L.-A., Sørensen, A., Eastwood, S., and Lavergne, T.: Global sea ice edge and type product user's manual, OSI-403-c & EUMETSAT, 2016.
- 260 Armitage, T. W. and Ridout, A. L.: Arctic sea ice freeboard from AltiKa and comparison with CryoSat-2 and Operation IceBridge, *Geophysical Research Letters*, 42, 6724–6731, 2015.
- Blockley, E. W. and Peterson, K. A.: Improving Met Office seasonal predictions of Arctic sea ice using assimilation of CryoSat-2 thickness, *The Cryosphere*, 12, 3419–3438, 2018.
- Chan, P., Halfar, J., Adey, W., Hetzinger, S., Zack, T., Moore, G., Wortmann, U., Williams, B., and Hou, A.: Multicentennial record of Labrador Sea primary productivity and sea-ice variability archived in coralline algal barium, *Nature communications*, 8, 15 543, 2017.
- 265 Day, J., Hawkins, E., and Tietsche, S.: Will Arctic sea ice thickness initialization improve seasonal forecast skill?, *Geophysical Research Letters*, 41, 7566–7575, 2014.
- Graham, R. M., Cohen, L., Petty, A. A., Boisvert, L. N., Rinke, A., Hudson, S. R., Nicolaus, M., and Granskog, M. A.: Increasing frequency and duration of Arctic winter warming events, *Geophysical Research Letters*, 44, 6974–6983, 2017.
- 270 Hallikainen, M., Ulaby, F., and Abdel-Razik, M.: Measurements of the dielectric properties of snow in the 4-18 GHz frequency range, in: 1982 12th European Microwave Conference, pp. 151–156, IEEE, 1982.
- Hendricks, S., Ricker, R., and Helm, V.: User Guide - AWI CryoSat-2 Sea Ice Thickness Data Product (v1.2), 2016.
- King, J., Skourup, H., Hvidegaard, S. M., Rösel, A., Gerland, S., Spreen, G., Polashenski, C., Helm, V., and Liston, G. E.: Comparison of Freeboard Retrieval and Ice Thickness Calculation From ALS, ASIRAS, and CryoSat-2 in the Norwegian Arctic to Field Measurements Made During the N-ICE2015 Expedition, *Journal of Geophysical Research: Oceans*, 123, 1123–1141, 2018.
- 275 Kurtz, N. T., Galin, N., and Studinger, M.: An improved CryoSat-2 sea ice freeboard retrieval algorithm through the use of waveform fitting, *The Cryosphere*, 8, 1217–1237, 2014.
- Kwok, R.: Simulated effects of a snow layer on retrieval of CryoSat-2 sea ice freeboard, *Geophysical Research Letters*, 41, 5014–5020, 2014.
- Kwok, R.: Arctic sea ice thickness, volume, and multiyear ice coverage: losses and coupled variability (1958–2018), *Environmental Research Letters*, 13, 105 005, 2018.
- 280 Kwok, R. and Cunningham, G.: ICESat over Arctic sea ice: Estimation of snow depth and ice thickness, *Journal of Geophysical Research: Oceans*, 113, 2008.
- Kwok, R. and Cunningham, G.: Variability of Arctic sea ice thickness and volume from CryoSat-2, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 373, 20140 157, 2015.
- 285 Kwok, R. and Kacimi, S.: Three years of sea ice freeboard, snow depth, and ice thickness of the Weddell Sea from Operation IceBridge and CryoSat-2, *The Cryosphere*, 12, 2789–2801, 2018.
- Kwok, R. and Markus, T.: Potential basin-scale estimates of Arctic snow depth with sea ice freeboards from CryoSat-2 and ICESat-2: An exploratory analysis, *Advances in Space Research*, 62, 1243–1250, 2018.
- Kwok, R., Cunningham, G., Wensnahan, M., Rigor, I., Zwally, H., and Yi, D.: Thinning and volume loss of the Arctic Ocean sea ice cover: 2003–2008, *Journal of Geophysical Research: Oceans*, 114, 2009.
- 290 Kwok, R., Panzer, B., Leuschen, C., Pang, S., Markus, T., Holt, B., and Gogineni, S.: Airborne surveys of snow depth over Arctic sea ice, *Journal of Geophysical Research: Oceans*, 116, 2011.

- Landy, J. C., Ehn, J. K., Babb, D. G., Thériault, N., and Barber, D. G.: Sea ice thickness in the Eastern Canadian Arctic: Hudson Bay Complex & Baffin Bay, Remote sensing of environment, 200, 281–294, 2017.
- 295 Langlois, A., Mundy, C., and Barber, D. G.: On the winter evolution of snow thermophysical properties over land-fast first-year sea ice, Hydrological Processes: An International Journal, 21, 705–716, 2007.
- Laxon, S., Peacock, N., and Smith, D.: High interannual variability of sea ice thickness in the Arctic region, Nature, 425, 947, 2003.
- Laxon, S. W., Giles, K. A., Ridout, A. L., Wingham, D. J., Willatt, R., Cullen, R., Kwok, R., Schweiger, A., Zhang, J., Haas, C., et al.: CryoSat-2 estimates of Arctic sea ice thickness and volume, Geophysical Research Letters, 40, 732–737, 2013.
- 300 Nandan, V., Geldsetzer, T., Yackel, J., Mahmud, M., Scharien, R., Howell, S., King, J., Ricker, R., and Else, B.: Effect of snow salinity on CryoSat-2 Arctic first-year sea ice freeboard measurements, Geophysical Research Letters, 44, 10–419, 2017.
- Nicolaus, M., Katlein, C., Maslanik, J., and Hendricks, S.: Changes in Arctic sea ice result in increasing light transmittance and absorption, Geophysical Research Letters, 39, 2012.
- Petty, A. A., Holland, M. M., Bailey, D. A., and Kurtz, N. T.: Warm Arctic, Increased Winter Sea Ice Growth?, Geophysical Research Letters, 305 45, 12–922, 2018a.
- Petty, A. A., Webster, M., Boisvert, L., and Markus, T.: The NASA Eulerian Snow on Sea Ice Model (NESOSIM) v1. 0: initial model development and analysis., Geoscientific Model Development, 11, 2018b.
- Ricker, R., Hendricks, S., Perovich, D. K., Helm, V., and Gerdes, R.: Impact of snow accumulation on CryoSat-2 range retrievals over Arctic sea ice: An observational approach with buoy data, Geophysical Research Letters, 42, 4447–4455, 2015.
- 310 Schweiger, A. J.: Unified Sea Ice Thickness Climate Data Record, Polar Science Center, Applied Physics Laboratory, University of Washington, [http://psc.apl.uw.edu/sea\\_ice\\_cdr](http://psc.apl.uw.edu/sea_ice_cdr), accessed on 09/9/2019, 2017.
- Sévellec, F., Fedorov, A. V., and Liu, W.: Arctic sea-ice decline weakens the Atlantic meridional overturning circulation, Nature Climate Change, 7, 604, 2017.
- Stroeve, J. and Notz, D.: Changing state of Arctic sea ice across all seasons, Environmental Research Letters, 13, 103 001, 2018.
- 315 Stroeve, J. C., Schroder, D., Tsamados, M., and Feltham, D.: Warm winter, thin ice?, The Cryosphere, 12, 1791–1809, 2018.
- Tilling, R. L., Ridout, A., and Shepherd, A.: Estimating Arctic sea ice thickness and volume using CryoSat-2 radar altimeter data, Advances in Space Research, 62, 1203–1225, 2018.
- Tiuri, M., Sihvola, A., Nyfors, E., and Hallikaiken, M.: The complex dielectric constant of snow at microwave frequencies, IEEE Journal of oceanic Engineering, 9, 377–382, 1984.
- 320 Ulaby, F. T., Moore, R. K., and Fung, A. K.: Microwave remote sensing: Active and passive. Volume 3-From theory to applications, 1986.
- Warren, S. G., Rigor, I. G., Untersteiner, N., Radionov, V. F., Bryazgin, N. N., Aleksandrov, Y. I., and Colony, R.: Snow depth on Arctic sea ice, Journal of Climate, 12, 1814–1829, 1999.
- Webster, M. A., Rigor, I. G., Nghiem, S. V., Kurtz, N. T., Farrell, S. L., Perovich, D. K., and Sturm, M.: Interdecadal changes in snow depth on Arctic sea ice, Journal of Geophysical Research: Oceans, 119, 5395–5406, 2014.
- 325 Willatt, R., Laxon, S., Giles, K., Cullen, R., Haas, C., and Helm, V.: Ku-band radar penetration into snow cover on Arctic sea ice using airborne data, Annals of Glaciology, 52, 197–205, 2011.
- Willatt, R. C., Giles, K. A., Laxon, S. W., Stone-Drake, L., and Worby, A. P.: Field investigations of Ku-band radar penetration into snow cover on Antarctic sea ice, IEEE Transactions on Geoscience and remote sensing, 48, 365–372, 2009.

- Wingham, D., Francis, C., Baker, S., Bouzinac, C., Brockley, D., Cullen, R., de Chateau-Thierry, P., Laxon, S., Mallow, U., Mavrocordatos, C., et al.: CryoSat: A mission to determine the fluctuations in Earth's land and marine ice fields, *Advances in Space Research*, 37, 841–871, 2006.
- Xia, W. and Xie, H.: Assessing three waveform retracers on sea ice freeboard retrieval from Cryosat-2 using Operation IceBridge Airborne altimetry datasets, *Remote Sensing of Environment*, 204, 456–471, 2018.
- Yaremchuk, M., Townsend, T., Panteleev, G., Hebert, D., and Allard, R.: Advancing Short-Term Forecasts of Ice Conditions in the Beaufort Sea, *Journal of Geophysical Research: Oceans*, 2019.

# Supplementary Information: Common assumptions involving the speed of radar in snow introduce systemic underestimates to sea ice thickness and seasonal growth rate estimates

Robbie D.C. Mallett <sup>1</sup>, Isobel R. Lawrence <sup>1</sup>, Julienne C. Stroeve <sup>1,2,3</sup>, Jack C. Landy <sup>4</sup>, and Michel C. Tsamados <sup>1</sup>

<sup>1</sup>Centre for Polar Observation and Modelling, Earth Sciences, University College London, London, UK

<sup>2</sup>National Snow and Ice Data Center, University of Colorado, Boulder, CO, USA

<sup>3</sup>Centre for Earth Observation Science, University of Manitoba, Winnipeg, Canada

<sup>4</sup>School of Geographical Sciences, University of Bristol, Bristol, UK

**Correspondence:** Robbie Mallett (robbie.mallett.17@ucl.ac.uk)

## 1 The Comparable Impacts on SIT of Snow Loading and Slower Radar Wave Propagation in Snow

$$\text{Ice Freeboard} = \text{Radar Freeboard} + \text{Propagation Correction} \quad (\text{Armitage and Ridout, 2015}) \quad (\text{S1})$$

$$h_i = h_r + h_s(c/c_s - 1) \quad (\text{S2})$$

5 The conversion of a given ice freeboard can be combined with a snow depth to estimate sea ice thickness:

$$SIT = h_i \frac{\rho_w}{\rho_w - \rho_i} + h_s \frac{\rho_s}{\rho_w - \rho_i} \quad (\text{Tilling et al. 2018}) \quad (\text{S3})$$


Substituting Eq. (S2) into Eq. (S3)

$$SIT = h_r \frac{\rho_w}{\rho_w - \rho_i} + h_s \frac{\rho_w}{\rho_w - \rho_i} \left[ \frac{c}{c_s} - 1 \right] + h_s \frac{\rho_s}{\rho_w - \rho_i} \quad (\text{S4})$$


$$10 \text{ Sea Ice Thickness} = \text{Radar Freeboard Component} + \text{Propagation Correction} + \text{Snow Loading} \quad (\text{S5})$$

Comparing the relative impacts of the Propagation Correction term and Snow Loading term is relatively simple given they share a common factor of  $h_s/(\rho_w - \rho_s)$ . The ratio of the two terms is therefore  $c/c_s - 1$  to  $\rho_s/\rho_w$ . For a typical snow density of  $300\text{kgm}^{-3}$ , this ratio is 0.25 to 0.29. (Using  $\rho_w = 1023.9$  and Ulaby et al. (1986) to relate  $\rho_s$  to  $c_s$ )


## 2 Supplementary Figures




**Figure S1.** Three commonly used relationships between radar wave speed and snow density Hallikainen et al. (1982); Tiuri et al. (1984); Ulaby et al. (1986).




**Figure S2.** The region over which snow depths published in Warren et al. (1999) are generally considered reliable (Laxon et al. (2013); Kwok and Cunningham (2015)), and over which freeboards are considered in this study.



**Figure S3.** Winter snow densification rates for five regions and the basin-wide average. We defined the ‘basin-wide area’ as the shaded area in Fig. (S2). We found the basin-wide densification rate to be roughly representative of its constituent regions apart from the Laptev, which exhibited a small but positive densification.



**Figure S4.** While the functional form and magnitude of expressions for the effect of snow weight and slower radar propagation are different, they have a similar error dependence on snow depth. That is to say, the percentage error introduced to the "weight correction" by snow density uncertainty is the same as that for the "propagation correction".



**Figure S5.** Monthly differences in sea ice thickness from the use of  $\delta h = 0.22Z$  and  $\delta h = 0.25Z$  for the propagation by AWI and CPOM respectively.

## 15 **References**

- Armitage, T. W. and Ridout, A. L.: Arctic sea ice freeboard from AltiKa and comparison with CryoSat-2 and Operation IceBridge, *Geophysical Research Letters*, 42, 6724–6731, 2015.
- Hallikainen, M., Ulaby, F., and Abdel-Razik, M.: Measurements of the dielectric properties of snow in the 4-18 GHz frequency range, in: 1982 12th European Microwave Conference, pp. 151–156, IEEE, 1982.
- 20 Kwok, R. and Cunningham, G.: Variability of Arctic sea ice thickness and volume from CryoSat-2, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 373, 20140157, 2015.
- Laxon, S. W., Giles, K. A., Ridout, A. L., Wingham, D. J., Willatt, R., Cullen, R., Kwok, R., Schweiger, A., Zhang, J., Haas, C., et al.: CryoSat-2 estimates of Arctic sea ice thickness and volume, *Geophysical Research Letters*, 40, 732–737, 2013.
- Tilling, R. L., Ridout, A., and Shepherd, A.: Estimating Arctic sea ice thickness and volume using CryoSat-2 radar altimeter data, *Advances*  
25 *in Space Research*, 62, 1203–1225, 2018.
- Tiuri, M., Sihvola, A., Nyfors, E., and Hallikaiken, M.: The complex dielectric constant of snow at microwave frequencies, *IEEE Journal of oceanic Engineering*, 9, 377–382, 1984.
- Ulaby, F. T., Moore, R. K., and Fung, A. K.: *Microwave remote sensing: Active and passive. Volume 3-From theory to applications*, 1986.
- Warren, S. G., Rigor, I. G., Untersteiner, N., Radionov, V. F., Bryazgin, N. N., Aleksandrov, Y. I., and Colony, R.: Snow depth on Arctic sea  
30 ice, *Journal of Climate*, 12, 1814–1829, 1999.