



## Thermokarst lake development in syngenetic ice-wedge polygon terrain in the Eastern Canadian Arctic (Bylot Island, Nunavut)

Frédéric Bouchard<sup>1,2</sup>, Daniel Fortier<sup>2,3</sup>, Michel Paquette<sup>4</sup>, Vincent Boucher<sup>5</sup>, Reinhard Pienitz<sup>2,5</sup>, Isabelle Laurion<sup>2,6</sup>

5 <sup>1</sup> Géosciences Paris Sud (GEOPS), Université Paris Saclay, Orsay, France

<sup>2</sup> Centre d'études nordiques (CEN), Université Laval, Québec, Canada

<sup>3</sup> Département de géographie, Université de Montréal, Montréal, Canada

<sup>4</sup> Department of Geography and Planning, Queen's University, Kingston, Canada

<sup>5</sup> Département de géographie, Université Laval, Québec, Canada

10 <sup>6</sup> Centre Eau Terre Environnement, Institut national de la recherche scientifique (INRS-ETE), Québec, Canada

Correspondence to: Frédéric Bouchard ([frederic.bouchard@u-psud.fr](mailto:frederic.bouchard@u-psud.fr))

**Abstract.** Thermokarst lakes are widespread and diverse across permafrost regions and they are considered significant contributors to global greenhouse gas emissions. Paleoenvironmental reconstructions documenting the inception and development of these ecologically important water bodies are generally limited to Pleistocene-age permafrost deposits (Yedoma) of Siberia, Alaska, and the western Canadian Arctic. Here we present the gradual transition from syngenetic ice-wedge polygon terrains to a thermokarst lake in the Eastern Canadian Arctic. We combine geomorphological surveys with paleolimnological reconstructions from sediment cores in an effort to characterize local landscape evolution from terrestrial to freshwater environment. Located on an ice-rich and organic-rich polygonal terrace, the studied lake is now evolving through active thermokarst, as revealed by subsiding and eroding shores, and was likely created by water pooling within a pre-existing topographic depression. Organic sedimentation in the valley started during the mid-Holocene, as documented by the oldest organic debris found at the base of one sediment core and dated at 4.8 kyr BP. Local sedimentation dynamics were initially controlled by fluctuations in wind activity, local moisture and vegetation growth/accumulation, as shown by alternating loess (silt) and peat layers. Fossil diatom assemblages were likewise influenced by local hydro-climatic conditions and reflect a broad range of substrates available in the past (both terrestrial and aquatic). Such conditions likely prevailed until ~ 2000 BP, when peat accumulation stopped as water ponded the surface of degrading ice-wedge polygons, and the basin progressively developed into a thermokarst lake. Interestingly, this happened in the middle of the Neoglacial cooling period, likely under wetter-than-average conditions. Thereafter, the lake continued to develop as evidenced by the dominance of aquatic (both benthic and planktonic) diatom taxa in organic-rich lacustrine muds. Based on these interpretations, we present a four-stage conceptual model of thermokarst lake development during the late Holocene, including some potential future trajectories. Such a model could be applied to other formerly glaciated syngenetic permafrost landscapes.



## 1 Introduction

Lakes are remarkably abundant across the circumpolar regions, with several millions of waterbodies spread over an estimated total surface area ranging from  $\sim 1.4$  to  $1.8 \times 10^6$  km<sup>2</sup> (Muster et al., 2017; Paltan et al., 2015; Verpoorter et al., 2014). The vast majority of these aquatic systems are located in permafrost environments, especially in lowland regions with moderate to high excess ground-ice content (typically  $> 30\%$  in volume) and a thick sediment cover (Grosse et al., 2013; Smith et al., 2007). Recent high-resolution mapping efforts reported a significant variability in waterbody size distributions across permafrost regions (Muster et al., 2017). Thermokarst (thaw) lakes occur mainly in ice-rich permafrost regions, where ground-ice melting can result in localized ground surface subsidence, water accumulation, and self-maintained lake expansion (van Everdingen, 1998). These lakes vary greatly in morphology, depth ( $< 1$  m to several meters deep in most cases) and area (from a few meters across to several km<sup>2</sup>) depending on ground-ice content and distribution, lake age, hydro-climatic conditions and local topography (e.g., Côté and Burn, 2002; Hopkins, 1949; Pienitz et al., 2008). Lakes located in unglaciated ice-rich (Yedoma) terrains of Siberia, Alaska and western Canada started to develop since the late Pleistocene and form a separate lake category, up to several tens of meters deep (e.g., Lenz et al., 2016; Morgenstern et al., 2011; Schirrmeister et al., 2011). Thermokarst lake evolution involves a remarkably diverse suite of hydro-climatic, geomorphological and ecological processes (Bouchard et al., 2017; Grosse et al., 2013). Although modern thermokarst processes and landforms may involve anthropogenic causes, thermokarst development during the Holocene can be associated to three main drivers: climate (increased air and ground temperatures), ground disturbances (through fluvial, thermal or ecological mechanisms, e.g. slumps or fires), and snow accumulation (French, 2017). In the latter case, the insulating capacity of a thick snow cover can result in significantly higher ground temperatures (near 0°C) during the winter and can trigger localized or widespread permafrost thawing. When thaw depth exceeds the maximum thickness of winter ice cover, mean annual lake-bottom temperature remains above 0 °C, resulting in the formation of a talik (thaw bulb) underneath the lake (Burn, 2002). Once initiated, thermokarst lakes in continuous permafrost tend to develop laterally; first by the coalescence of polygonal and/or ice-wedge trough pools overlying melting ice-wedge networks (Czudek and Demek, 1970; Prėskienis et al., in prep), and then by thermal and mechanical shoreline erosional processes, such as wave-induced erosional niche development or mass wasting through thaw slumping and block failures (Kokelj and Jorgenson, 2013). Ultimately, and depending on local landscape conditions (e.g., soil type, vegetation cover, topography), thermokarst lake development ends with one or more of the following: rapid drainage resulting from shoreline breaching, either during higher-than-average lake-level episodes (e.g., Jones and Arp, 2015; Lantz and Turner, 2015; Mackay and Burn, 2002; Turner et al., 2010) or due to ice-wedge melting and thermal erosion gullying (e.g., Fortier et al., 2007; Godin and Fortier, 2012); lake-level drawdown due to factors that lead to increased evaporation (Bouchard et al., 2013a; Riordan et al., 2006); subsurface drainage (groundwater infiltration) through an open talik (Yoshikawa and Hinzman, 2003); or terrestrialization via rapid peat accumulation and lake infilling (Payette et al., 2004; Roach et al., 2011).

Thermokarst lakes also play a key role in the global carbon cycle. Because they form in areas where organic carbon is stored in frozen soils (Hugelius et al., 2014; Schuur et al., 2015), they have been identified as biogeochemical hotspots through their



65 release of substantial amounts of carbon dioxide (CO<sub>2</sub>) and particularly of methane (CH<sub>4</sub>) to the atmosphere (e.g., Abnizova  
 et al., 2012; Laurion et al., 2010; Matveev et al., 2018; Walter et al., 2007). A fundamental yet less often considered aspect is  
 the age (millennium-old vs. modern) of the carbon processed and released by thermokarst ecosystems, which is linked to their  
 potential to generate a positive feedback on climate (Elder et al., 2018; Mann et al., 2015; Vonk et al., 2013). Carbon older  
 than ~ 500 to 1000 years can be considered as ‘in excess’ in the system, thus representing a net atmospheric contribution from  
 70 a formerly stable reservoir (Prėskienis et al., in prep and references therein). Work conducted in the Eastern Canadian Arctic  
 and in Alaska has shown that radiocarbon age can indeed vary by several orders of magnitude over a small area depending on  
 waterbody properties (Bouchard et al., 2015a; Elder et al., 2018). Yet, the majority of studies focusing on the age and sources  
 of CO<sub>2</sub> and CH<sub>4</sub> released by thermokarst lakes come from Yedoma regions, which represent a small fraction (~ 4-6 %) of total  
 permafrost areas (~ 1.0-1.4 x10<sup>6</sup> km<sup>2</sup> out of 23 x10<sup>6</sup> km<sup>2</sup> in total; e.g., Strauss et al., 2017). Lakes formed in formerly glaciated  
 75 terrains are widespread across the Arctic and can contribute significantly to global greenhouse gas emissions (Wik et al., 2016).  
 Here we document the development of a thermokarst lake in a tundra valley of the Eastern Canadian Arctic (Bylot Island,  
 Nunavut, Canada) during the Holocene. We test the hypothesis that this lake developed following local landscape dynamics,  
 and not solely because of climate. The lake is located within an ice-rich and organic-rich syngenetic permafrost environment  
 (Fortier and Allard, 2004). It thus serves as an interesting case-study landscape under-represented in the thermokarst literature.  
 80 We combine high-resolution lake mapping, geomorphological observations and paleolimnological reconstructions (both litho-  
 and biostratigraphy) in an effort to 1) document the inception and evolution of a thermokarst lake in a syngenetic ice-wedge  
 polygon terrain, 2) characterize the transition from terrestrial to aquatic conditions in a tundra valley geosystem, and 3) present  
 a conceptual model of thermokarst development during the Holocene.

## 2 Study site

85 Bylot Island (Nunavut) is located in the Eastern Canadian Arctic, within the continuous permafrost zone (Fig. 1a). Most of the  
 island is mountainous, and several glaciers spread from its center to peripheral lowland areas (Fig. 1b). These glacial valleys  
 were shaped during the successive Pleistocene glaciations (Klassen, 1993), and since the Holocene they developed highly  
 dynamic biogeosystems rich in vegetation, ground ice, peat, and aquatic environments (Allard, 1996; Fortier and Allard, 2004).  
 The prevailing climate is polar with a slight marine influence. Based on the 1980-2010 climate normals from the nearby village  
 90 of Pond Inlet (Mittimatalik) on Baffin Island (72° 41’ N; 77° 58’ W), the mean annual air temperature is -14.6 °C, with average  
 daily temperatures ranging from -33.4 °C in January to 6.6 °C in July, and total precipitations of 189 mm, of which 91 mm fall  
 as rain between June and September (Environment Canada, 2019). Thawing and freezing degree-days are around 475 and  
 5735, respectively. Winter (defined as when continuous daily mean air temperature < 0° C) lasts from early September to mid-  
 June, for an average total of 283 days per year. A station operated since 2004 by the Center for Northern Studies (CEN) at the  
 95 study site provides similar climate data (CEN, 2018).



The study site (73° 09' N; 79° 58' W) is located in the valley locally named Qarlikturvik, which has a NE-SW orientation and a surface area of ~ 65 km<sup>2</sup> (~ 15 km-long x 4-5 km-wide) (Fig. 1c). A terminal moraine, located roughly halfway between the actual glacier (C-79) front and the seashore and sitting on marine clay, was <sup>14</sup>C-dated to ~ 9.8 kyr BP (Allard, 1996). Holocene glacial retreat, accompanied by a marine transgression phase, ended around 6 kyr BP. Deposited marine clays were subsequently covered by glacio-fluvial sand and gravel (Fortier and Allard, 2004). Today, a braided river flows through a glacio-fluvial outwash plain, carrying sediments towards a delta aggrading in Navy Board Inlet. This outwash plain is bordered on both sides by a 3- to 5-m thick terrace, crisscrossed by networks of tundra polygons associated with the formation of syngenetic ice wedges. Along the southern bank of the river, the upper portion of this terrace is composed of alternating mineral (wind-blown sand and silt) and organic (peat) material, which started to accumulate over glacio-fluvial sands and gravels before 3700 years ago (Fortier and Allard, 2004). These peaty loess deposits in which permafrost aggrades syngenetically are typically ice-rich, and their organic matter content can reach over 50 %. The active layer depth in such deposits generally ranges between 40 to 80 cm (down to 1 m in sandy/gravelly material), and the maximum depth of permafrost on Bylot Island has been estimated to be over 400 m (Allard et al., 2016; Smith and Burgess, 2000).

The sampled lake, informally named Gull Lake, is located within the lake- and pond-rich polygonal terrace, near the terminal moraine (Fig. 1c). Limnological observations conducted during the ice-free season indicate relatively low concentrations of dissolved organic carbon (DOC), nutrients and ions in Gull Lake compared to the surrounding ice-wedge troughs and coalescent polygonal ponds, as well as a thermally homogenous and well-oxygenated water column. However, dissolved oxygen concentrations decrease rapidly under the winter ice cover and at the bottom of the lake, near the water-sediment interface (Bouchard et al., 2015a; Prëskienis et al., in prep). Greenhouse gas (GHG) sampling and dating showed that this lake is a relatively small but spatially variable source of dissolved and ebullition GHG, with millennium-age methane released in its center (up to 3.5 kyr BP) and peripheral shallow zones (up to 2.8 kyr BP). The age of methane emitted off-shore is almost corresponding to the maximum age of syngenetic organic sedimentation in the valley (3.7 kyr BP) (Bouchard et al., 2015a; Fortier and Allard, 2004; Prëskienis et al., in prep).

### 3 Materials and methods

#### 3.1 Lake watershed and geomorphology

A portable sonar system, equipped with an internal GPS antenna (Humminbird model 859XD) and mounted on a small zodiac, was used to map the lake bottom in July 2014 (Bouchard et al., 2015b). Lake-depth signals were continuously recorded along regularly spaced (20-25 m apart) navigation lines, mainly of SW-NE and SE-NW orientations. Depth data were interpolated between navigation lines using the compatible software (AutoChart) to produce a geo-referenced 3D bathymetric map. The acquired data were also used to calibrate and extrapolate lake bottom depths inferred from GPR mapping conducted the following year.



Ground penetrating radar (GPR) surveying on lake ice cover allows accurate description of lake bottom topography (Moorman, 2001; Paquette et al., 2015; You et al., 2017). Three GPR survey lines crossing the lake were conducted in May 2015 (Fig. 2) using a sleigh-dragged Sensors and Software PulseEkko GPR and 50 MHz antennas. GPR line processing was performed using Ekko project software and included Dewow, Lowpass temporal filter to diminish background noise and a background average subtraction to remove the overwhelming ice/water boundary signal. The base of the ice cover and lake bottom depth were manually identified and corresponded well to the dielectric properties of ice and water. Signal travel velocities of  $0.06 \text{ m ns}^{-1}$  and  $0.13 \text{ m ns}^{-1}$  were used respectively for unfrozen and frozen ground.

### 3.2 Sediment core sampling and logging

Sediment cores were collected during two consecutive years: a short core (54 cm) from a boat during the ice-free season of 2014 (July), and a longer core (109 cm) from the ice cover in spring 2015 (June). The coring site location ( $> 4 \text{ m}$  deep) was located in 2014 and 2015 using the bathymetric data from the sonar and GPR surveys (Bouchard et al., 2015b). Each core was retrieved using a handheld percussion corer equipped with a 7-cm diameter clear polycarbonate tube (Aquatic Research Instruments). The 2014 core was subsampled in the field immediately after retrieval at 1-cm intervals, and the subsamples were transferred into polyethylene bags and brought back to the laboratory, where they were kept in the dark at  $4 \text{ }^{\circ}\text{C}$ . For the 2015 core, water from above the sediment surface was removed immediately after retrieval in order to prevent the mixing of the water-sediment interface (Bouchard et al., 2011). The core was then stored vertically at non-freezing conditions for at least 48 h, allowing the upper sediments to slowly consolidate by dewatering. It was finally sealed with foam blocks to minimize potential disturbances during subsequent transport, brought back to the laboratory and stored in the dark at  $4 \text{ }^{\circ}\text{C}$  for further analyses.

The 2014 core was visually examined in the field before and during subsampling to identify general stratigraphic units (e.g., gyttja, peat, silt and sand), whereas the 2015 core was described with more detail in the laboratory. First, a non-destructive computed tomographic scan was performed to visualize internal sedimentary structures and infer sediment density (Supplement S1) (Calmels and Allard, 2004). This core was then cut along its longitudinal axis with a rotating saw and split in two halves. One half was covered with a plastic film to minimize surface oxidation and desiccation and kept aside for archives (in the dark at  $4 \text{ }^{\circ}\text{C}$ ), and stratigraphic descriptions were done on the other half. Subsampling was then performed on the working half at 1-cm intervals, and the subsamples were freeze-dried (at least 48 h, depending on water content) and transferred into polyethylene bags for further analyses.

### 3.3 Lithological and chronological analyses

Sediment subsamples from the 2014 and 2015 cores were used to perform physical and chronological analyses. About  $0.5 \text{ g}$  of dry sediment was extracted to perform loss-on-ignition (LOI) measurements by a subsequent combustion at  $550 \text{ }^{\circ}\text{C}$  for 4 h (Heiri et al., 2001). Organic content (LOI) and wet/dry sediment mass measurements were used to determine the sediment dry bulk density and to correlate both cores, in addition to visual descriptions. Supplementary subsamples were used to perform



grain-size distribution analysis by sieving material coarser than 62.5  $\mu\text{m}$  (i.e. sand and gravel) (ASTM, 2004) and using  
 160 hydrometry for fine sediments (i.e. silt and clay) (ASTM, 2017).

Bulk sediment samples and, when present, fossil organic/wood fragments were carefully extracted and dried in glass bottles  
 at 105 °C (Björck and Wohlfarth, 2001). Samples were pre-treated (HCl-NaOH-HCl) and combusted to  $\text{CO}_2$  at the  
 Radiocarbon Dating Laboratory (Université Laval, Québec QC, Canada) and  $^{14}\text{C}$  dated by accelerator mass spectrometry  
 (AMS) at Keck Carbon Cycle AMS Facility (University of California, Irvine CA, USA). Radiocarbon dates were reported  
 165 using Libby's half-life (5568 yr), corrected for natural fractionation ( $\delta^{13}\text{C} = -25\text{‰}$  PDB), and calibrated with the CALIB 7.1  
 online program (Stuiver et al., 2019) using the IntCal13 calibration data set (Reimer et al., 2013).

### 3.4 Diatom analysis

Fossil diatom analysis was conducted at the Aquatic Paleoecology Laboratory of CEN on 60 subsamples from the 2015 core  
 (each cm for the top 12 cm, then each 2 cm towards core bottom) following Bouchard et al. (2013b). Diatom valves were  
 170 extracted from ~ 50-mg samples using acid ( $\text{H}_2\text{SO}_4\text{-HNO}_3$ ) digestion techniques and mounted on microscope slides using  
*Naphrax*, a highly refractive resin (Battarbee et al., 2001). For each subsample, an average of 400 diatom valves were counted  
 along transects using a Leica DMRX light microscope. Identification was carried out to the lowest taxonomic level possible  
 (i.e., species or variety/morphotype) at a 1000  $\times$  magnification. Taxonomic identification mainly followed Antoniadis et al.  
 (2008; 2009), Fallu et al. (2000), Krammer (2000, 2002), Krammer and Lange-Bertalot (1986; 1988; 1991a; 1991b), Lavoie  
 175 et al. (2008), and Zimmermann et al. (2010). Diatom taxa representing at least 1% (relative abundance) in at least one sample  
 were plotted on diagrams using the C2 software (Juggins, 2014). Photos of most of these taxa were taken with a Leica DFC490  
 camera (mounted on the microscope) and were used to prepare plates of the representative taxa (Supplement S2). In the  
 following sections, taxa names are given as they appeared originally in consulted floras.

## 4 Results

### 180 4.1 Lake basin morphology

Gull Lake has an irregular shape and bathymetry. It is mostly SW-NE elongated, parallel to the main valley axis, with  
 maximum length and width of ~ 500 and 250 m, respectively, for a total surface of  $116 \times 10^3 \text{ m}^2$ . There is no apparent inlet;  
 however, the lake receives influx of snowmelt water during the spring. A small outlet, draining towards a nearby lake and the  
 proglacial river to the North, is observable along the northern shore (Figs. 1 and 2). Based on the bathymetric map and satellite  
 185 imagery, the lake can be split into two zones: a shallow platform and a deeper basin. The shallow platform (< 2 m deep)  
 occupies the periphery of the lake while gently dipping towards lake center. Submerged ice-wedge polygons can be seen on  
 this platform, as well as degraded furrows observed during the ice-free period with a submersible camera (Bouchard et al.,  
 2015b; Video Supplement VS1). This morphology confirms that the lake is currently evolving through lateral thermokarst  
 encroachment. The deeper (~ 2-4 m deep) basin occupies the center of the lake. This section is relatively bumpy, with shallower



190 areas that can be distinguished from a boat or on the satellite image (Fig. 2). This central basin appears asymmetrical, with maximum depths ( $> 4$  m deep) concentrated within the SW portion of the lake.

GPR surveys conducted on top of lake ice during spring (early June) 2015 provide further information about Gull Lake morphology and winter conditions (Fig. 3) (Fortier et al., 2019). Lake ice thickness averages  $2.1 \text{ m} \pm 0.1 \text{ m}$  in the central basin, i.e. in areas where ice is not grounded. Lake depth is typically deeper than  $3.2 \text{ m}$  on average within the central basin. In contrast, mean depth is  $< 1 \text{ m}$  within the peripheral platform, where the ice cover extends to the bottom of the lake. Apart from the  $\sim 30 \text{ m}$  long transition between the shallow platform and the deeper central basin, local slopes are generally gentle (rarely exceeding  $4^\circ$ ). Finally, many strong electromagnetic reflectors are visible into the ground on all GPR lines (Fig. 3). A first series of these reflectors are located underneath lake bottom, in both shallow and deeper zones, at an average depth of  $0.49 \text{ m}$  under the sediment surface (range  $0\text{--}1.53 \text{ m}$ ). Another group of deeper reflectors are visible only under the shallow peripheral platform, at a depth of  $2.6 \pm 1.4 \text{ m}$ . The signal velocity ( $> 0.13 \text{ m ns}^{-1}$ ) based on the shape of some hyperboles suggests that they occur in frozen material. All these reflectors are located from  $\sim 5$  to  $40 \text{ m}$  apart (apparent distance along GPR lines).

## 4.2 Lake sediment stratigraphy

### 4.2.1 Lithostratigraphy

Based on the description of the 2015 core, sedimentary units or zones appear as follows (Fig. 4) (Fortier and Bouchard, 2019a; 2019b), from bottom to top (equivalent in the 2014 core, when observed, is mentioned at the end of each paragraph):

- **Lithozone 1** (109–80 cm). This unit is composed of mostly sand and gravel ( $> 50 \%$ ) with scattered peat and organic debris. Subzones 1a (109–103 cm) and 1c (86–80 cm) contain only sand and gravel, whereas subzone 1b (103–86 cm) contains organic debris, mostly in the form of cm-scale pieces of agglomerated peat. Compared to other units, lithozone 1 has a relatively high mean density ( $\sim 2 \text{ g cm}^{-3}$ ), typical of dominantly mineral material. Water content (20–40 %) and LOI ( $< 10 \%$ ) are relatively low, except for the above-mentioned peat and organic debris, as shown for example by a peak at 93–94 cm with 60 % water content and 25 % LOI. Based on  $^{14}\text{C}$  dating of one subsample (107–108 cm), this unit contains organic matter older than 5500 cal. yr BP (4805  $^{14}\text{C}$  BP) (Table 1). No equivalent was found in the shorter 2014 core.
- **Lithozone 2** (80–10 cm). This unit is composed of medium to dark brown porous peat, moderately decomposed, interbedded with mm- to cm-thick silt and sand laminations. These silt/sand laminations are generally thicker ( $> 1 \text{ cm}$ ) and more present at the base of the unit compared to the top. The average proportion of silt vs. sand in the mineral fraction is around 70 % vs. 30 %, respectively. An intermediate subzone 2b (55–35 cm), richer in sand ( $\sim 50 \%$ ) and marked by convoluted horizons, separates subzones 2a (80–55 cm) and 2c (35–10 cm), notably dominated by peat. From the bottom to the top, there is a generally decreasing trend in density, from  $\sim 2 \text{ g cm}^{-3}$  (mostly mineral) to  $\sim 1 \text{ g cm}^{-3}$  (mostly organic material), with the exception of the above-mentioned subzone 2b. Meanwhile, there is an



upward increase in water content (from 20 to > 60 %) and LOI (from < 10 to > 20 %), again with the exception of subzone 2b. This unit was observed in the 2014 core, at depths between 54 and 10 cm.

- **Lithozone 3** (10-0 cm). This unit is composed of laminated dark organic lacustrine mud (gyttja) overlying an organic-poor silt layer (10 cm deep). The relative proportion of silt vs. sand in the mineral fraction is higher (> 80 % vs. < 10 %) compared to the underlying unit. With the exception of this silty mineral layer, the density is relatively low (~ 1.25 g cm<sup>-3</sup>), typical of organic material. It has a high water content (60 to 80 %), similar to subzone 2c, but a medium LOI (~ 15 %), except for the basal silt layer (< 10 %). The basal silty layer was <sup>14</sup>C-dated in both 2014 and 2015 cores (Table 1), giving an age of around 2000 cal. yr BP (~ 2100 <sup>14</sup>C BP). Similar sediments were observed in the 2014 core at the same depth (10-0 cm).

#### 230 4.2.2 Biostratigraphy

A total of 230 diatom taxa belonging to 52 genera were identified within the 60 thin sections prepared from the 2015 core (Pienitz et al., 2019). The average number of taxa for a given level was 43, ranging from a minimum of 5 (108-109 cm) to a maximum of 60 taxa (56-58 cm and 18-20 cm). Among these, 15 so-called “major taxa” (species or species groups) representing more than 5 % in relative abundance in at least one level were selected to show major ecological changes that occurred in the past (Fig. 5). These changes were used to delimit diatom zones (or “biozones”), which are similar to the sedimentary units (or “lithozones”) described above, although exact upper and lower limits are slightly different. These major biozones are as follows, from bottom to top:

- **Biozone 1** (109-74 cm). Compared to the entire core, this unit is characterized by a poor diversity in major taxa (n<10) and in total counted taxa (n<30 in average per level). The diversity is especially low in subzone 1a (109-102 cm), with only 5 major taxa and an average of 13 counted taxa per level. Notably, the *Diploneis-Geissleria* group, practically not observed anywhere else along the core, is overwhelmingly dominant within this subzone (> 20% of relative abundance). These species are generally associated with cold, oligotrophic, organic-poor, low conductivity and mostly alkaline (pH = ~ 8) waters, typical of Arctic streams and wetland headwaters (Antoniades et al., 2008; Zimmermann et al., 2010). The overlying subzones 1b (102-86 cm) and 1c (86-74 cm) are notably more diverse in identified taxa (average total counted taxa of 33 and 34, respectively) and dominated by aerophilous/moss-associated species (e.g., *Chameapinnularia soehrensii*, *Pinnularia sinistra*, *Diatomella balfouriana*), generally living in circumneutral to slightly acidic waters, typical of high-latitude peatlands (D. Antoniades, pers. comm.; Zimmermann et al., 2010). Abundant organic debris, in the form of cm-scale pieces of peat, were indeed observed in this unit (Fig. 4).
- **Biozone 2** (74-12 cm). This unit marks an increase in the abundance of major taxa (n>10), with the appearance of mostly small, benthic diatom genera (e.g., *Cavinula*, *Achnanthisidium*, *Fragilaria*, *Staurosirella*) typical of shallow tundra ponds in ice-wedge polygon terrains, with cold waters and long-lasting ice cover (Antoniades et al., 2008; Ellis et al., 2008; Pienitz et al., 1995; Zimmermann et al., 2010). Total counted taxa are also much higher in this unit (nearly





50 in average per level), except in subzone 2b (48-34 cm) where the number of identified species per level ranges around 35. This intermediate subzone corresponds to the convoluted silt/sand horizons of lithozone 2 (subzone 2b; Fig. 4) and is mostly dominated by epiphytic, moss-associated genera (*Encyonema*, *Eunotia*, *Caloneis*) (Antoniades et al., 2008; Ellis et al., 2008; Zimmermann et al., 2010).

- **Biozone 3** (12-0 cm). This unit is similar to the underlying biozone 2, with a slightly higher number of major taxa (n=13) although a slightly lower total number of counted taxa (n=47 on average per level). Thin sections were more concentrated in diatom valves within this zone, especially in the upper part (7-0 cm). Moreover, several taxa with a generally wide geographic distribution in lakes and wetlands and preferring high-nutrient waters (e.g., *Cavinula cocconeiformis*, *Cymbopleura naviculiformis*, *Eunotia bilunaris*) (Guiry and Guiry, 2019), not observed in other zones, were counted within this unit (Pienitz et al., 2019). This biozone is the equivalent of lithozone 1 (laminated lacustrine mud; Fig. 4).

## 5 Discussion

Combining geomorphological and paleolimnological observations of Gull Lake basin and bottom sediments, we can reconstruct landscape dynamics in Qarlikturvik valley and lake development during the second half of the Holocene. This evolution was however strongly controlled by the early Holocene deglaciation of the valley, for which data from earlier studies are available. Hence, we first adopt a chronological approach in this section, covering the whole Holocene, in order to better 'set the stage' for Gull Lake's inception. We then present a four-stage conceptual model for thermokarst lake development in syngenetic permafrost of formerly glaciated terrains. Finally, we discuss the implications of some of our results on carbon dynamics in the Arctic.

### 5.1 Holocene history of the Qarlikturvik valley and ground-ice development

At the beginning of the Holocene, the Qarlikturvik valley was likely occupied by an inland-based glacier with a front advancing into shallow marine waters (Allard, 1996). This interpretation is based on the presence of marine shells (*Mya truncata* species) <sup>14</sup>C-dated at 9860 yr BP, and found within ice-contact deposits (sands, gravels and pebbles) with lithological properties corresponding to the surrounding Precambrian and Cretaceous-Tertiary rocks. Lacelle et al. (2018) and Coulombe et al. (2019) later proposed that Laurentide ice and Bylot ice were converging in the valley. Glacial retreat was then accompanied by a marine transgression phase, associated with the deposition of silts and clays, and which lasted until about 6000 yr BP (<sup>14</sup>C ages in shells at different altitudes ranging from 9860 to 6100 yr BP). The fossiliferous marine sediments were observed within pingo cores along the southern shore of the proglacial river, as upheaved and slightly deformed strata (Allard, 1996).

The second half of the Holocene was marked by the deposition of glacio-fluvial, eolian and organic sediments over the valley floor. First, following marine regression after ~ 6 kyr BP, a glacial outwash plain probably occupied the entire valley, overlying the marine silts and clays and depositing glacio-fluvial sands and gravels (Allard, 1996, Fortier and Allard, 2004). Small



streams with cold, alkaline, low-DOC and nutrient-poor waters were likely widespread within the plain, as inferred from diatoms observed at the base of the core collected in Gull Lake (Fig. 5; Antoniadou et al., 2008; Zimmermann et al., 2010). The  $^{14}\text{C}$  date of 4.8 yr BP (5.5 cal. yr BP) at the base of the core is interpreted as reworked pieces of peat transported by glacio-fluvial waters that sedimented in channels of the outwash (Table 1). This glacio-fluvial period lasted about two millennia, until eolian (fine sand and silt) and organic (peat) sediments started to accumulate in the valley around 3.7 kyr BP, as based on  $^{14}\text{C}$  dating of *Salix* twigs and peat macrofossils (Fortier and Allard, 2004). A greater initial accumulation rate of  $> 2 \text{ mm yr}^{-1}$  occurred in this period, followed by a reduction to  $< 1 \text{ mm yr}^{-1}$  after 2.2 kyr BP (Allard, 1996; Fortier et al., 2006). Effective organic (peat) sedimentation during this period is further supported by the presence, in the Gull Lake core, of abundant benthic and epiphytic diatom species generally preferring moss substrates typical of more acidic peatland/wetland environments (Antoniades et al., 2008; Pienitz, 2001; Zimmermann et al., 2010). These eolian and organic layers (stratified silt and peat) were gradually incorporated (syngenetically) into the permafrost as they accumulated and froze (see below), forming the polygonal terrace within which numerous ponds and lakes later formed and are still visible today (Fig. 1). Syngenetic permafrost and associated ground-ice development followed sediment deposition in the Qarlikturvik valley. At the late Holocene (roughly 3500 years ago), the valley floor had completely emerged from the sea and the proglacial river running through the valley had started to cut into its own alluvial deposits. At the same time, cooler regional temperatures (the Neoglacial) resulted in slower melting of upstream glaciers, thus lower flow of the river, which enhanced the above-mentioned colonization of the outwash plain by eolian and organic sediments (Fortier and Allard, 2004). Neoglacial cooling was reported at numerous sites across the Eastern Canadian Arctic, based on diverse paleoenvironmental indicators (summarized in Fortier et al., 2006). Repeated thermal frost cracking during severe winters had likely started as soon as the downstream portion of the valley was exposed (i.e. around 6000 yr BP) and later affected the whole glacio-fluvial outwash plain (and the upper section of the underlying marine clay unit), resulting in the formation of a first generation of ice wedges and related polygon networks. The development of the silty peat terrace, starting at 3000-3500 yr BP, brought a change in thermal contraction properties of the ground, triggering the formation of a second generation of ice-wedge polygons (Fortier and Allard, 2004). As a result, ice wedges of several meters wide and 6-8 m deep are extending today through the whole sedimentary sequence (likely down to the marine clays). These ice wedges define a complex patchwork of high-center and low-center polygons with a diameter ranging from 5 to 40 m, with the top of ice-wedges located a few centimeters below the base of the active layer in the transient layer (Allard, 1996; Shur et al. 2005). About two thousand years ago, Gull Lake inception site was located in such a typical tundra landscape.

## 5.2 Thermokarst lake evolution: a conceptual model for syngenetic permafrost in formerly glaciated terrains

Based on the geomorphology and paleolimnology of Gull Lake's basin, we developed a conceptual model of its formation and evolution during the late Holocene, including some potential future trajectories. This four-stage model is summarized in Fig. 6.



The initial conditions, before Gull Lake was formed (Stage 0), were summarized above for the first half of the Holocene, including landscape evolution and ground-ice development. At the beginning of the late Holocene, the site was characterized by a network of syngenetic ice wedges extending through several meters of frozen peat, eolian silt, glacio-fluvial sands, and likely marine clays (Fig. 6a). Based on present-day lake morphology (Fig. 2) and a volumetric ice content exceeding 50 % in the silty peat unit (Fortier and Allard, 2004), which contributed to subsequent subsidence of the ground surface upon thaw, a pre-existing depression 1-2 m deep must have existed at that time to explain the current thickness of this specific unit found at lake bottom (i.e., ~ 0.7 m; Fig. 4). The ice wedge network in Qarlikturvik valley still possesses such topographic lows today, and Gull Lake was probably initiated in such a depression, which collected snow and snowmelt waters, especially during years of higher precipitation and weaker winds. Local topography and its interaction with climate, especially dominant winds during winter (i.e. windward vs. leeward slopes), strongly impacts snow cover depth, density, and duration (French, 2014). There can thus be a complex local heterogeneity in thermokarst development, regardless of regional climate variations. Notably, lake inception (i.e. transition from terrestrial to aquatic sedimentation) was <sup>14</sup>C dated at around 2100 yr BP (Table 1 and Fig. 4), which corresponds to a slightly warmer episode of the Neoglacial cooling period with wetter-than-average local conditions (Fortier et al., 2006).

Such conditions resulted in active layer deepening and thermokarst initiation with ice wedge melting and development of small and shallow ponds, either over the ice wedges or at their junctions (Grosse et al., 2013) (stage 1; Fig. 6b). Diatom communities changed from moss-associated or aerophilous (terrestrial) species to dominantly benthic or planktonic (aquatic) taxa typical of tundra ponds (Fig. 5) (Ellis et al., 2008). This mixed tundra landscape, combining terrestrial and freshwater environments, characterizes high-centered polygon networks.

Although being completely frozen during the winter, these small water bodies were a significant source of latent heat to extract in autumn, delaying the freezing front propagation in the underlying ground (Kokelj and Jorgenson, 2013) (stage 2; Fig. 6c). Ponds then started to coalesce over and at the edge of ice-wedge polygons, extending the aquatic surface area over the terrestrial one. At this stage, however, the aquatic conditions did not last year-round for these water bodies, as they were too shallow to maintain liquid water below the ice cover (> 2 m) in winter. Shallow coalescent ponds are still a common feature in the valley in present days (Prėskienis et al., in prep).

Eventually, lateral expansion by both thermal and mechanical erosion, as well as thaw consolidation and subsidence beneath waterbodies, led to the formation of a lake *sensu stricto*, that is with bottom waters remaining liquid throughout the winter (stage 3; Fig. 6d). Year-round aquatic conditions started to prevail, leading to the gradual accumulation of organic-rich lacustrine mud (gyttja). This is illustrated by the presence of abundant benthic and planktonic diatom species in the upper part of the analyzed core (0-10 cm; Figs. 4 and 5), typical of lacustrine ecosystems across several regions (Guiry and Guiry, 2019). However, the specific morphology of the lake, with a central deeper basin presenting relatively steep slopes surrounded by a gently sloping shallow platform, suggests that this evolution did not follow a strictly linear trend. The central basin, covered by at least 10 cm of lacustrine sediments at the coring site, appears notably older than the peripheral platform, where no or negligible gyttja was observed at 1-m depth (Video Supplement VS1). Assuming a relatively low sedimentation rate of 0.1 to



0.2 mm per year, typical of thermokarst lakes (Bouchard et al. 2011; Coulombe et al., 2016), the central basin was likely formed several centuries before the surrounding shallow platform.

Gull Lake's bathymetry clearly indicates that lake bottom in the central basin is generally deeper than the maximum ice cover thickness. Hence, thermal conditions are met for the development of a talik underneath the lake. A temperature sensor installed within surface sediments of the deepest portion of the central basin in 2014-2015 showed that it never froze, although it stayed close to 1-2 °C for 9 months throughout the year, i.e. from mid-October to mid-June (Supplement S3). However, an unfrozen zone could not be clearly detected along the GPR lines (Fig. 3). Moreover, the bathymetry in the central basin is notably heterogeneous, with shallows (~ 2 m deep) in contact with winter ice cover (GPR line12; Fig. 3), and a deeper portion limited to the SW section of the lake (Fig. 2). The inferred talik is therefore not typically bowl-shaped and underlying the whole lake central basin, as generally reported from other thermokarst lake basins (Morgenstern et al., 2011 and references therein). A potential explanation for this heterogeneity in the deep basin bathymetry is likely related to the differential surface subsidence above ice wedges *vs.* above polygon centers, the latter being tempered by the presence of fibrous peat. Based on GPR lines, the lake bottom bumpy topography can indeed be interpreted as the former surface of ice-wedge polygon ridges and troughs.

Notably, numerous reflectors were identified along GPR lines underneath the lake bottom (at > 35 cm depth on average), and these reflectors were located 5 to 40 m apart laterally, similar to what is today observable between ice wedges within the valley (Fig. 3) (Fortier et al., 2019). The other (deeper) series of reflectors, found only underneath the peripheral platform (average depth of 2.6 m), thus in frozen material, likely represent the top of the glacio-fluvial unit (sand and gravel). This unit was indeed observed in the sediment core, which was collected in the deep basin so in a non-frozen state (Fig. 4). Fortier and Allard (2004) reported a similar depth for the glacio-fluvial unit, based on permafrost coring and GPR surveys on the polygonal terrace a few tens of meters from Gull Lake's northern shore.

Gull Lake is currently slowly expanding laterally by thermokarst in the syngenetic frozen silt-peat terrace, and the deepest parts of the bottom have now almost reached the underlying glacio-fluvial sand (Fig. 4). The future evolution of the lake to its final disappearance might thus include one or both of the following scenarios: 1) a gradual terrestrialization *via* gyttja accumulation and lake infilling (stage 4a; Fig. 6e), 2) a rapid lateral drainage *via* shoreline breaching resulting from fluvial erosion (e.g., thermo-erosion gullying) (stage 4b; Fig. 6f).

Lake or pond infilling causing terrestrialization has been reported from Arctic coastal Alaska (Jorgenson and Shur, 2007), subarctic eastern Canada (Payette et al., 2004), and boreal interior Alaska (Kanevskiy et al., 2014; Roach et al., 2011). The presence of lacustrine sediments in the upper section (10 cm) of the collected cores suggests that such a process has begun, although quite slowly. The absence of a visible inlet, as well as the relatively low concentrations of organic matter and nutrients measured at different times of the year (Prèskienis et al., in prep), might explain the slow sedimentation rate in Gull Lake. Added to current observations that 1) lake shores migrate laterally by both thermal and mechanical erosion, and 2) lake peripheral platform progressively deepens by thaw subsidence, it is likely that the complete lake infilling resulting in full-scale terrestrialization might not happen in a foreseeable future. More realistically, partial infilling might have time to occur before natural landscape evolution results in partial lake drainage.



Partial or complete drainage or shrinkage of thermokarst lakes can be related to their long-term water balances in relation to regional climate (e.g., precipitation vs. evaporation) (Bouchard et al., 2013a; Riordan et al., 2006). However, thermokarst lake drainage can also be a rapid, catastrophic event that has been reported from several permafrost regions across the Arctic (summarized for instance in Grosse et al., 2013 or Kokelj and Jorgenson, 2013). The drivers for an abrupt drainage are mostly related to local geomorphology and natural landscape evolution, that is, factors that are external to the lakes themselves. These include ice wedge melting in the surrounding basin creating a drainage network, retrogressive development of thermo-erosion gullies towards a given lake, coastal erosion or tapping by another lake or a river. In the case of the Qarlikturvik valley in general, and Gull Lake specifically, it is likely that future evolution will involve lake drainage to a certain extent. First, networks of rapidly evolving thermo-erosion gullies, developed along melting ice-wedge networks, have been reported elsewhere in the valley (Fortier et al., 2007). Once initiated, these gully networks developed extremely rapidly during the first year (average erosion rates of several meters per day) and at a much slower but quasi-steady rate afterwards for the following decade (Godin and Fortier, 2012). Obviously, such processes have a strong impact on local hydrology, including snow redistribution and surface/subsurface hydrological connectivity (Godin et al., 2014). Second, field observations suggest that a nearby lake, located immediately downslope of Gull Lake (informally named “Gull Lake 2” or GL-2; Fig. 1c), has partly drained in the past. This is based on the observation of former lake shores, pingo development, and the absence of well-developed ice-wedge polygons in the immediate surroundings of the lake (unpublished data). Such a partial drainage could likely happen to Gull Lake in the future, affecting at least the shallow peripheral platform and leaving a residual smaller lake corresponding to the current deeper basin.

### 5.3 Implications for Arctic carbon dynamics

Several square kilometres of the Qarlikturvik valley are currently underlain by a syngenetic permafrost terrace composed of alternating mineral (silt) and organic (peat) layers 3-5 meters thick and with an average organic matter content of 40 % (ranging from 15 to 65 %) (Fortier and Allard, 2004). Assuming that bulk organic matter contains 58 % of organic carbon (Pribyl, 2010), the terrace presents slightly over 20 % of total organic carbon (TOC). This value is roughly one order of magnitude higher than the TOC values generally reported from the Yedoma domain of Siberia, Alaska and NW Canada, that is 2-3 % (e.g., Schirmer et al., 2011; Strauss et al., 2017). For this, we are assuming volumetric ground-ice content (40-70 %) and bulk density ( $1-1.5 \times 10^3 \text{ kg m}^{-3}$ ) that are comparable as in other circumpolar regions (Fortier et al., 2006). Since the thickness of the organic-rich permafrost terrace on Bylot Island is roughly one order of magnitude lower than the average thickness of Yedoma deposits (30-50 m), specific carbon stocks in both types of landscapes are more or less equivalent (i.e., ranging from ~ 600 to 1500  $\text{kg C m}^{-2}$ ; Schirmer et al., 2011). However, this comparison does not take into account the lability of the organic matter, with much older parent material of a different diagenetic state in Yedoma landscapes (Tank et al., in review and references therein). Moreover, the comparison does not include any spatial considerations about the total carbon stocks: Yedoma landscapes are estimated to cover ~  $1.0-1.4 \times 10^6 \text{ km}^2$  (Strauss et al., 2017), whereas such numbers for syngenetic glaciated terrains are, to our knowledge, currently not available. More generally, there are considerable regional differences in



the biogeochemical response to permafrost thaw, and the key role played for instance by local relief and parent material (geology) on controlling carbon mobilization has been recognized just recently (Tank et al., in review and references therein). Nevertheless, if future climate change was to result in widespread and deep (several meters) thaw of ice-rich permafrost, it is worth noting that the thawing of one meter of organic-rich frozen ground in this valley of Bylot Island could mobilize an order of magnitude more organic carbon ( $200\text{--}300 \text{ kg C m}^{-3}$ ) than an equivalent layer in Yedoma landscapes ( $20\text{--}30 \text{ kg C m}^{-3}$ ; Schirmer et al., 2011). Therefore, the short-term carbon feedback potential caused by GHG emissions from the present study landscape is likely much higher than from Yedoma regions, as to obtain the same amount of organic carbon released from thawing of the upper 3 m permafrost terrace on Bylot Island, almost the entire Yedoma complex (30 m depth) would have to thaw, which is extremely unlikely in the foreseeable future. For the specific case of Gull Lake, since the lake bottom has almost reached the much less organic-rich layer of glacio-fluvial sands, future emissions from the deep basin are likely to slow down, although lateral expansion will likely fuel emissions from the current peripheral platform. To make reasonable estimations from syngenetic glaciated terrains at the global scale, we need to know not only its global coverage, but the thickness and lability of the organic layer in a range of locations.

## 6 Conclusions

Combining high-resolution lake mapping (sonar and GPR), geomorphological observations and paleolimnological reconstructions (litho- and biostratigraphy) from Gull Lake on Bylot Island, we developed a conceptual model of thermokarst lake evolution (past, present and future) in a syngenetic glaciated permafrost landscape of the Eastern Canadian Arctic during the Holocene. Such a model explains multiple steps of local landscape evolution from terrestrial to freshwater environment and is currently underrepresented in the thermokarst literature, which is dominated by Pleistocene-age permafrost deposits (Yedoma).

Based on our results, there is no linear, unidirectional, or cyclic evolution of thermokarst lakes, at least not at our study site. We rather infer natural landscape evolution (e.g., local topography, ground ice development, snow cover distribution and depth, surface hydrology), in conjunction with regional climate, as the main driver. Thermokarst lake development on Bylot Island during the Holocene appears as a self-developing process occurring within a mature landscape: once initiated, it proceeds regardless of climate. Notably, Gull Lake inception likely started during a cooler climate period (Neoglacial) around 2000 years ago, underscoring the importance of precipitation and local snow distribution over temperature only.

The valley surrounding Gull Lake is currently underlain by a syngenetic permafrost terrace which, although much thinner than Yedoma ice complex (3–5 m vs. 30–50 m, respectively), contains an equivalent amount of stored organic carbon per surface unit ( $\sim 600$  to  $1500 \text{ kg C m}^{-2}$ ). If future climate change was to result in the thawing of several meters of ice-rich permafrost across the Arctic, the carbon feedback potential caused by GHG emissions from the present study landscape will thus likely be much higher than from Yedoma regions.



#### 450 **Data availability**

The following datasets have been submitted to the Nordicana D collection at Centre d'études nordiques (CEN – Centre for Northern Studies) (<http://www.cen.ulaval.ca/nordicanad/>) and are currently under review for formatting. The complete citations of each dataset appear in the reference list of this manuscript.

- Fortier et al.: Ground-penetrating radar (GPR) surveys over a thermokarst lake, Bylot Island, Nunavut, Canada. doi: 10.5885/45609CE-E3573955017A4904.
- Fortier and Bouchard: Computed tomography (CT) scanning of a lake sediment core, Bylot Island, Nunavut, Canada. doi: 10.5885/45612CE-AB27C20EB10D4509.
- Fortier and Bouchard: Loss-on-ignition and grain size analysis of a lake sediment core, Bylot Island, Nunavut, Canada. doi: 10.5885/45603CE-21852993EE434926.
- Pienitz et al.: Identification of fossil diatoms present in a lake sediment core, Bylot Island, Nunavut, Canada. doi: 10.5885/45600CE-C0960664FE8F4038.

#### **Video Supplement**

**VS1.** Underwater camera video of submerged degraded ice-wedge polygons at the bottom of the peripheral shallow platform of Gull Lake, Bylot Island, Nunavut, Canada (as in Bouchard et al., 2015b). Water depth is approximately 1 m. Footage was collected in July 2014. doi: 10.5446/43923. Accessible at <https://doi.org/10.5446/43923>.

#### **Supplement Materials**

1. Computed tomography (CT) scanning of a 109-cm long sediment core collected in June 2015 in Gull Lake, Bylot Island, Nunavut, Canada. Methods as in Calmels and Allard (2004). 1 Table (Table S1), 2 figures (Figs. S1-S2).
2. Plates (photographs) of the most abundant fossil diatoms found in a 109-cm long sediment core collected in June 2015 in Gull Lake, Bylot Island, Nunavut, Canada. Methods as in Bouchard et al. (2013b). 1 Figure (Fig. S3).
3. Lake-water temperatures recorded at the bottom of a thermokarst lake (2014-2015), Bylot Island, Nunavut, Canada. 1 Figure (Fig. S4).

#### **Author contribution**

FB, DF, RP and IL designed the research goals and methods. FB, MP and VB conducted a first analysis of the data and produced the figures. DF and IL funded fieldwork sampling and analyses. FB prepared the manuscript with contributions from all co-authors.



### Competing interests

The authors declare that they have no conflict of interest.

### Acknowledgements

480 We are grateful to the team of Gilles Gauthier (Dept. of biology, U. Laval), the Center for Northern Studies (CEN) and the staff of the Sirmilik National Park (Parks Canada) for logistical support and access to Bylot Island. We sincerely thank Jean-Philippe Tremblay and Yoan Lechasseur for fieldwork preparation assistance, Vilmantas Prėskienis, Audrey Veillette and Maxime Tremblay for their help in the field, Arianne Lafontaine and Andréanne Lemay for their help in the laboratory, and Stéphanie Coulombe for inspiring discussions while preparing the first draft of the manuscript. We are also very thankful to  
485 the community and the people of Pond Inlet (Mittimatalik). This research was funded by ArcticNet, the Natural Sciences and Engineering Research Council of Canada (NSERC), the Polar Continental Shelf Program (PCSP) of Natural Resources Canada, the NSERC Discovery Frontiers grant “Arctic Development and Adaptation to Permafrost in Transition” (ADAPT), the EnviroNorth Training Program, and the W. Garfield Weston Foundation.





## 490 References

- Abnizova, A., Siemens, J., Langer, M., and Boike, J.: Small ponds with major impact: The relevance of ponds and lakes in permafrost landscapes to carbon dioxide emissions, *Global Biogeochemical Cycles*, 26, GB2041, doi: 10.1029/2011gb004237, 2012.
- Allard, M.: Geomorphological changes and permafrost dynamics: Key factors in changing arctic ecosystems. An example from Bylot Island, Nunavut, Canada, *Geoscience Canada*, 23, 205-212, 1996.
- Allard, M., Sarrazin, D., and L'Hérault, E.: Borehole and near-surface ground temperatures in northeastern Canada, v. 1.4 (1988-2016). *Nordicana D8*, doi: 10.5885/45291SL-34F28A9491014AFD, 2016.
- Antoniades, D., Hamilton, P. B., Douglas, M. S. V., and Smol, J. P.: Diatoms of North America: The freshwater floras of Prince Patrick, Ellef Ringnes and northern Ellesmere Islands from the Canadian Arctic Archipelago. *Iconographia Diatomologica*, Vol. 17, A.R.G. Gantner Verlag, 649 p., 2008.
- Antoniades, D., Hamilton, P. B., Hinz, F., Douglas, M. S. V., Smol, J. P.: Seven new species of freshwater diatoms (Bacillariophyceae) from the Canadian Arctic Archipelago. *Nova Hedwigia*, 88, 57-80, doi: 10.1127/0029-5035/2009/0088-0057, 2009.
- ASTM: D6913-04 – Standard Test Methods for Particle-Size Distribution (Gradation) of Soils Using Sieve Analysis, ASTM International, West Conshohocken, PA, 2004.
- ASTM: D7928-17 – Standard Test Method for Particle-Size Distribution (Gradation) of Fine-Grained Soils Using the Sedimentation (Hydrometer) Analysis, ASTM International, West Conshohocken, PA, 2017.
- Battarbee, R., Jones, V., Flower, R., Cameron, N., Bennion, H., Carvalho, L., and Juggins, S.: Diatoms, in: *Tracking Environmental Change Using Lake Sediments. Volume 3: Terrestrial, Algal, and Siliceous Indicators*, edited by: Smol, J., Birks, J., Last, W., Bradley, R., and Alverson, K., *Developments in Paleoenvironmental Research*, 3, Springer Netherlands, 155-202, 2001.
- Björck, S., and Wohlfarth, B.: <sup>14</sup>C Chronostratigraphic Techniques in Paleolimnology, in: *Tracking Environmental Change Using Lake Sediments. Volume 1: Basin Analysis, Coring, and Chronological Techniques*, edited by: Last, W., and Smol, J., *Developments in Paleoenvironmental Research*, 1, Springer Netherlands, 205-245, 2001.
- Bouchard, F., Francus, P., Pienitz, R., and Laurion, I.: Sedimentology and geochemistry of thermokarst ponds in discontinuous permafrost, subarctic Quebec, Canada, *Journal of Geophysical Research-Biogeosciences*, 116, G00M04, doi: 10.1029/2011JG001675, 2011.
- Bouchard, F., Turner, K. W., MacDonald, L. A., Deakin, C., White, H., Farquharson, N., Medeiros, A. S., Wolfe, B. B., Hall, R. I., Pienitz, R., and Edwards, T. W. D.: Vulnerability of shallow subarctic lakes to evaporate and desiccate when snowmelt runoff is low, *Geophys. Res. Lett.*, 40, 6112-6117, doi: 10.1002/2013GL058635, 2013a.



- Bouchard, F., Pienitz, R., Ortiz, J. D., Francus, P., and Laurion, I.: Palaeolimnological conditions inferred from fossil diatom assemblages and derivative spectral properties of sediments in thermokarst ponds of subarctic Quebec, Canada, *Boreas*, 42, 575-595, doi: 10.1111/bor.12000, 2013b.
- Bouchard, F., Laurion, I., Prêskienis, V., Fortier, D., Xu, X., and Whiticar, M. J.: Modern to millennium-old greenhouse gases emitted from ponds and lakes of the Eastern Canadian Arctic (Bylot Island, Nunavut), *Biogeosciences*, 12, 7279-7298, doi: 10.5194/bg-12-7279-2015, 2015a.
- Bouchard, F., Fortier, D., Paquette, M., Bégin, P. N., Vincent, W. F., and Laurion, I.: Lake bottom imagery: a simple, fast and inexpensive method for surveying shallow freshwater ecosystems of permafrost regions, *Proceedings of the 7th Canadian Permafrost Conference and the 68th Canadian Geotechnical Conference*, Quebec City, Canada, 2015b.
- Bouchard, F., MacDonald, L. A., Turner, K. W., Thienpont, J. R., Medeiros, A. S., Biskaborn, B. K., Korosi, J., Hall, R. I., Pienitz, R., and Wolfe, B. B.: Paleolimnology of thermokarst lakes: a window into permafrost landscape evolution, *Arctic Science*, 3, 91-117, doi: 10.1139/as-2016-0022, 2017.
- Brown, J., Ferrians, O. J., Heginbottom, J. A., and Melnikov, E. S.: Circum-Arctic map of permafrost and ground-ice conditions, Revised Feb. 2001, National Snow and Ice Data Center/World Data Center for Glaciology, Boulder, Colorado, 1998.
- Burn, C. R.: Tundra lakes and permafrost, Richards Island, western Arctic coast, Canada, *Canadian Journal of Earth Sciences*, 39, 1281-1298, doi: 10.1139/e02-035, 2002.
- Calmels, F., and Allard, M.: Ice segregation and gas distribution in permafrost using tomodesitometric analysis, *Permafrost and Periglacial Processes*, 15, 367-378, doi: 10.1002/ppp.508, 2004.
- CEN: Climate station data from Bylot Island in Nunavut, Canada, v. 1.9 (1992-2018): <http://www.cen.ulaval.ca/nordicanad/dpage.aspx?doi=45039SL-EE76C1BDAADC4890>, access: 2019-06-07, 2018.
- Côté, M. M., and Burn, C. R.: The oriented lakes of Tuktoyaktuk Peninsula, Western Arctic Coast, Canada: a GIS-based analysis, *Permafrost and Periglacial Processes*, 13, 61-70, doi: 10.1002/ppp.407, 2002.
- Coulombe, O., Bouchard, F., and Pienitz, R.: Coupling of sedimentological and limnological dynamics in subarctic thermokarst ponds in Northern Québec (Canada) on an interannual basis, *Sedimentary Geology*, 340, 15-24, doi: 10.1016/j.sedgeo.2016.01.012, 2016.
- Coulombe, S., Fortier, D., Lacelle, D., Kanevskiy, M., and Shur, Y.: Origin, burial and preservation of late Pleistocene-age glacier ice in Arctic permafrost (Bylot Island, NU, Canada), *The Cryosphere*, 13, 97-111, doi: 10.5194/tc-13-97-2019, 2019.
- Czudek, T., and Demek, J.: Thermokarst in Siberia and its influence on the development of lowland relief, *Quaternary Research*, 1, 103-120, doi: 10.1016/0033-5894(70)90013-x, 1970.
- Elder, C. D., Xu, X., Walker, J., Schnell, J. L., Hinkel, K. M., Townsend-Small, A., Arp, C. D., Pohlman, J. W., Gaglioti, B. V., and Czimczik, C. I.: Greenhouse gas emissions from diverse Arctic Alaskan lakes are dominated by young carbon, *Nat. Clim. Chang.*, 8, 166-171, doi: 10.1038/s41558-017-0066-9, 2018.



- Ellis, C. J., and Rochefort, L.: Century-scale development of polygon-patterned tundra wetland, bylot island (73 °N, 80 °W),  
 555 Ecology, 85, 963-978, doi: 10.1890/02-0614, 2004.
- Ellis, C. J., and Rochefort, L.: Long-term sensitivity of a High Arctic wetland to Holocene climate change, Journal of Ecology,  
 94, 441-454, doi: 10.1111/j.1365-2745.2005.01085.x, 2006.
- Ellis, C. J., Rochefort, L., Gauthier, G., and Pienitz, R.: Paleoeological Evidence for Transitions between Contrasting  
 Landforms in a Polygon-Patterned High Arctic Wetland, Arctic, Antarctic, and Alpine Research, 40, 624-637, doi:  
 560 10.1657/1523-0430(07-059)[ellis]2.0.co;2, 2008.
- Environment Canada: 1981-2010 Climate Normals & Averages:  
[http://climate.weather.gc.ca/climate\\_normals/results\\_1981\\_2010\\_e.html?searchType=stnName&txtStationName=Pond+Inlet&searchMethod=contains&txtCentralLatMin=0&txtCentralLatSec=0&txtCentralLongMin=0&txtCentralLongSec=0&stnID=1774&dispBack=1](http://climate.weather.gc.ca/climate_normals/results_1981_2010_e.html?searchType=stnName&txtStationName=Pond+Inlet&searchMethod=contains&txtCentralLatMin=0&txtCentralLatSec=0&txtCentralLongMin=0&txtCentralLongSec=0&stnID=1774&dispBack=1), access: 2019-06-07, 2019.
- 565 Fallu, M. A., Allaire, N., and Pienitz, R.: Freshwater diatoms from northern Québec and Labrador (Canada): species-  
 environment relationships in lakes of boreal forest, forest-tundra and tundra regions, Bibliotheca Diatomologica, J. Cramer,  
 Berlin/Stuttgart, 200 pp., 2000.
- Fortier, D., and Allard, M.: Late Holocene syngenetic ice-wedge polygons development, Bylot Island, Canadian Arctic  
 Archipelago, Canadian Journal of Earth Sciences, 41, 997-1012, doi: 10.1139/e04-031, 2004.
- 570 Fortier, D., Allard, M., and Pivot, F.: A late-Holocene record of loess deposition in ice-wedge polygons reflecting wind activity  
 and ground moisture conditions, Bylot Island, eastern Canadian Arctic, The Holocene, 16, 635-646, doi:  
 10.1191/0959683606hl960rp, 2006.
- Fortier, D., Allard, M., and Shur, Y.: Observation of rapid drainage system development by thermal erosion of ice wedges on  
 Bylot island, Canadian Arctic Archipelago, Permafrost and Periglacial Processes, 18, 229-243, doi: 10.1002/ppp.595, 2007.
- 575 Fortier, D., and Bouchard, F.: Computed tomography (CT) scanning of a lake sediment core, Bylot Island, Nunavut, Canada,  
 v. 1.0 (2015-2015), Nordicana D54, doi: 10.5885/45612CE-AB27C20EB10D4509, 2019a.
- Fortier, D., and Bouchard, F.: Loss-on-ignition and grain size analysis of a lake sediment core, Bylot Island, Nunavut, Canada,  
 v. 1.0 (2015-2015), Nordicana D52, doi: 10.5885/45603CE-21852993EE434926, 2019b.
- Fortier, D., Paquette, M., and Bouchard, F.: Ground-penetrating radar (GPR) surveys over a thermokarst lake, Bylot Island,  
 580 Nunavut, Canada, v. 1.0 (2015-2015), Nordicana D53, doi: 10.5885/45609CE-E3573955017A4904, 2019.
- French, H. M.: The periglacial environment, 4th ed., John Wiley & Sons, Chichester (UK), 515 pp., 2017.
- Godin, E., and Fortier, D.: Geomorphology of a thermo-erosion gully, Bylot Island, Nunavut, Canada, Canadian Journal of  
 Earth Sciences, 49, 979-986, doi: 10.1139/e2012-015, 2012.
- Godin, E., Fortier, D., and Coulombe, S.: Effects of thermo-erosion gullying on hydrologic flow networks, discharge and soil  
 585 loss, Environmental Research Letters, 9, doi: 105010, 10.1088/1748-9326/9/10/105010, 2014.
- Grosse, G., Jones, B., and Arp, C.: Thermokarst Lakes, Drainage, and Drained Basins, in: Treatise on Geomorphology, edited  
 by: Shroder, J. F., Glacial and Periglacial Geomorphology, 8, Academic Press, San Diego, CA, 325-353, 2013.



- Guiry, M. D., and Guiry, G. M.: AlgaeBase: <http://www.algaebase.org>, access: 2019-06-07, 2019.
- Heiri, O., Lotter, A. F., and Lemcke, G.: Loss on ignition as a method for estimating organic and carbonate content in  
 590 sediments: reproducibility and comparability of results, *Journal of Paleolimnology*, 25, 101-110, doi:  
 10.1023/A:1008119611481, 2001.
- Hopkins, D. M.: Thaw Lakes and Thaw Sinks in the Imuruk Lake Area, Seward Peninsula, Alaska, *The Journal of Geology*,  
 57, 119-131, 1949.
- Hugelius, G., Strauss, J., Zubrzycki, S., Harden, J. W., Schuur, E. A. G., Ping, C. L., Schirmer, L., Grosse, G., Michaelson,  
 595 G. J., Koven, C. D., O'Donnell, J. A., Elberling, B., Mishra, U., Camill, P., Yu, Z., Palmtag, J., and Kuhry, P.: Estimated stocks  
 of circumpolar permafrost carbon with quantified uncertainty ranges and identified data gaps, *Biogeosciences*, 11, 6573-6593,  
 doi: 10.5194/bg-11-6573-2014, 2014.
- Jones, B. M., and Arp, C. D.: Observing a Catastrophic Thermokarst Lake Drainage in Northern Alaska, *Permafrost and  
 Periglacial Processes*, 26, 119-128, doi: 10.1002/ppp.1842, 2015.
- 600 Jorgenson, M. T., and Shur, Y.: Evolution of lakes and basins in northern Alaska and discussion of the thaw lake cycle, *Journal  
 of Geophysical Research-Earth Surface*, 112, F02S17, doi: 10.1029/2006jf000531, 2007.
- Juggins, S.: C2 version 1.7.6. Software for ecological and palaeoecological data analysis and visualisation. Univ. of Newcastle,  
 Newcastle upon Tyne, 2014.
- Kanevskiy, M., Jorgenson, T., Shur, Y., O'Donnell, J. A., Harden, J. W., Zhuang, Q., and Fortier, D.: Cryostratigraphy and  
 605 Permafrost Evolution in the Lacustrine Lowlands of West-Central Alaska, *Permafrost and Periglacial Processes*, 25, 14-34,  
 doi: 10.1002/ppp.1800, 2014.
- Klassen, R. A.: Quaternary Geology and Glacial History of Bylot Island, Northwest Territories, *Memoir 429*, n° 429, edited  
 by: Geological Survey of Canada, Ottawa, 1993.
- Kokelj, S. V., and Jorgenson, M. T.: Advances in Thermokarst Research, *Permafrost and Periglacial Processes*, 24, 108-119,  
 610 doi: 10.1002/ppp.1779, 2013.
- Krammer, K.: The genus *Pinnularia*. In H Lange-Bertalot, Ed. *Diatoms of Europe – Diatoms of the European inland waters  
 and comparable habitats*, vol. 1. A. R. G. Gantner Verlag K. G., Ruggel, 2000.
- Krammer, K.: *Cymbella*. In H. Lange-Bertalot, Ed. *Diatoms of Europe – Diatoms of the European inland waters and  
 comparable habitats*, vol. 3. A. R. G. Gantner Verlag K. G., Ruggel, 2002.
- 615 Krammer, K., and Lange-Bertalot, H.: *Bacillariophyceae 1. Teil: Naviculaceae*. In H Ettl, J Gerloff, H Heynig and D  
 Mollenhauer, Eds. *Süßwasserflora von Mitteleuropa*. Gustav Fischer Verlag, Stuttgart/New York, 1986.
- Krammer, K., and Lange-Bertalot, H.: *Bacillariophyceae 2. Teil: Bacillariaceae, Epithemiaceae, Surirellaceae*. In H Ettl, J  
 Gerloff, H Heynig and D Mollenhauer, Eds. *Süßwasserflora von Mitteleuropa*. Gustav Fischer Verlag, Stuttgart/New York,  
 1988.
- 620 Krammer, K., and Lange-Bertalot, H.: *Bacillariophyceae 3. Teil: Centrales, Fragilariaceae, Eunotiaceae*. In H Ettl, J Gerloff,  
 H Heynig and D Mollenhauer, Eds. *Süßwasserflora von Mitteleuropa*. Gustav Fischer Verlag, Stuttgart/New York, 1991a.



- Krammer, K., and Lange-Bertalot, H.: Bacillariophyceae 4. Teil: Achnanthaceae, Kritische Ergänzungen zu Navicula (Lineolatae) und Gomphonema. In H Ettl, J Gerloff, H Heynig and D Mollenhauer, Eds. Süßwasserflora von Mitteleuropa. Gustav Fischer Verlag, Stuttgart/New York, 1991b.
- 625 Lacelle, D., Fisher, D. A., Coulombe, S., Fortier, D., and Frappier, R.: Buried remnants of the Laurentide Ice Sheet and connections to its surface elevation, *Sci Rep*, 8, 13286-13286, doi: 10.1038/s41598-018-31166-2, 2018.
- Lantz, T. C., and Turner, K. W.: Changes in lake area in response to thermokarst processes and climate in Old Crow Flats, Yukon, *Journal of Geophysical Research-Biogeosciences*, 120, 513-524, doi: 10.1002/2014JG002744, 2015.
- Laurion, I., Vincent, W. F., MacIntyre, S., Retamal, L., Dupont, C., Francus, P., and Pienitz, R.: Variability in greenhouse gas emissions from permafrost thaw ponds, *Limnology and Oceanography*, 55, 115-133, doi: 10.4319/lo.2010.55.1.0115, 2010.
- 630 Lavoie, I., Hamilton, P. B., Campeau, S., Grenier, M., and Dillon, P. J.: Guide d'identification des diatomées des rivières de l'Est du Canada, Presses de l'Université du Québec, Québec, 241 pp., 2008.
- Lenz, J., Wetterich, S., Jones, B. M., Meyer, H., Bobrov, A., and Grosse, G.: Evidence of multiple thermokarst lake generations from an 11 800-year-old permafrost core on the northern Seward Peninsula, Alaska, *Boreas*, 45, 584-603, doi: 10.1111/bor.12186, 2016.
- 635 Mackay, J. R., and Burn, C. R.: The first 20 years (1978-1979 to 1998-1999) of ice-wedge growth at the Illisarvik experimental drained lake site, western Arctic coast, Canada, *Canadian Journal of Earth Sciences*, 39, 95-111, doi: 10.1139/e01-048, 2002.
- Mann, P. J., Eglinton, T. I., McIntyre, C. P., Zimov, N., Davydova, A., Vonk, J. E., Holmes, R. M., and Spencer, R. G. M.: Utilization of ancient permafrost carbon in headwaters of Arctic fluvial networks, *Nat Commun*, 6, 7856, doi: 10.1038/ncomms8856, 2015.
- 640 Matveev, A., Laurion, I., and Vincent, W. F.: Methane and carbon dioxide emissions from thermokarst lakes on mineral soils, *Arctic Science*, 4, 584-604, doi: 10.1139/as-2017-0047, 2018.
- Moorman, B. J.: Ground-Penetrating Radar Applications in Paleolimnology. Volume 1: Basin Analysis, Coring, and Chronological Techniques, in: *Tracking Environmental Change Using Lake Sediments. Volume 1: Basin Analysis, Coring, and Chronological Techniques*, edited by: Smol, J. P., and Last, W. M., Springer Netherlands, 23-47, 2001.
- 645 Morgenstern, A., Grosse, G., Günther, F., Fedorova, I., and Schirrmeister, L.: Spatial analyses of thermokarst lakes and basins in Yedoma landscapes of the Lena Delta, *The Cryosphere*, 5, 849-867, doi: 10.5194/tc-5-849-2011, 2011.
- Muster, S., Roth, K., Langer, M., Lange, S., Cresto Aleina, F., Bartsch, A., Morgenstern, A., Grosse, G., Jones, B., Sannel, A. B. K., Sjöberg, Y., Günther, F., Andresen, C., Veremeeva, A., Lindgren, P. R., Bouchard, F., Lara, M. J., Fortier, D., Charbonneau, S., Virtanen, T. A., Hugelius, G., Palmtag, J., Siewert, M. B., Riley, W. J., Koven, C. D., and Boike, J.: PeRL: a circum-Arctic Permafrost Region Pond and Lake database, *Earth Syst. Sci. Data*, 9, 317-348, doi: 10.5194/essd-9-317-2017, 2017.
- 650 Paltan, H., Dash, J., and Edwards, M.: A refined mapping of Arctic lakes using Landsat imagery, *Int. J. Remote Sens.*, 36, 5970-5982, doi: 10.1080/01431161.2015.1110263, 2015.



- 655 Paquette, M., Fortier, D., Mueller, D. R., Sarrazin, D., and Vincent, W. F.: Rapid disappearance of perennial ice on Canada's  
 most northern lake, *Geophysical Research Letters*, 42, 1433-1440, doi: 10.1002/2014GL062960, 2015.
- Payette, S., Delwaide, A., Caccianiga, M., and Beauchemin, M.: Accelerated thawing of subarctic peatland permafrost over  
 the last 50 years, *Geophysical Research Letters*, 31, L18208, doi: 10.1029/2004GL020358, 2004.
- Pienitz, R.: Analyse des microrestes végétaux: diatomées, in: *Écologie des tourbières du Québec-Labrador*, edited by: Payette,  
 660 S., and Rochefort, L., Les Presses de l'Université Laval, Québec, 311-326, 2001.
- Pienitz, R., Bouchard, F., and Boucher, V.: Identification of fossil diatoms present in a lake sediment core, Bylot Island,  
 Nunavut, Canada, v. 1.0 (2015-2015), *Nordicana D51*, doi: 10.5885/45600CE-C0960664FE8F4038, 2019.
- Pienitz, R., Doran, P. T., and Lamoureux, S. F.: Origin and geomorphology of lakes in the polar regions, in: *Polar Lakes and  
 Rivers: Limnology of Arctic and Antarctic Aquatic Ecosystems*, edited by: Vincent, W., and Laybourn-Parry, J., Oxford  
 665 University Press, Oxford, U.K., 25-41, 2008.
- Pienitz, R., Smol, J. P., and Birks, H. J. B.: Assessment of freshwater diatoms as quantitative indicators of past climatic change  
 in the Yukon and Northwest Territories, Canada, *Journal of Paleolimnology*, 13, 21-49, doi: 10.1007/bf00678109, 1995.
- Prëskienis, V., Laurion, I., Bouchard, F., et al.: Seasonal patterns in greenhouse gas emissions from High Arctic lakes and  
 ponds on a polygonal landscape. To be submitted to *Limnology & Oceanography*, in prep.
- 670 Pribyl, D. W.: A critical review of the conventional SOC to SOM conversion factor, *Geoderma*, 156, 75-83, doi:  
 10.1016/j.geoderma.2010.02.003, 2010.
- Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Ramsey, C. B., Buck, C. E., Cheng, H., Edwards, R. L.,  
 Friedrich, M., Grootes, P. M., Guilderson, T. P., Hafliðason, H., Hajdas, I., Hatté, C., Heaton, T. J., Hoffmann, D. L., Hogg,  
 A. G., Hughen, K. A., Kaiser, K. F., Kromer, B., Manning, S. W., Niu, M., Reimer, R. W., Richards, D. A., Scott, E. M.,  
 675 Southon, J. R., Staff, R. A., Turney, C. S. M., and van der Plicht, J.: IntCal13 and Marine13 Radiocarbon Age Calibration  
 Curves 0–50,000 Years cal BP, *Radiocarbon*, 55, 1869-1887, doi: 10.2458/azu\_js\_rc.55.16947, 2013.
- Riordan, B., Verbyla, D., and McGuire, A. D.: Shrinking ponds in subarctic Alaska based on 1950-2002 remotely sensed  
 images, *Journal of Geophysical Research-Biogeosciences*, 111, G04002, doi: 10.1029/2005jg000150, 2006.
- Roach, J., Griffith, B., Verbyla, D., and Jones, J.: Mechanisms influencing changes in lake area in Alaskan boreal forest, *Global  
 680 Change Biology*, 17, 2567-2583, doi: 10.1111/j.1365-2486.2011.02446.x, 2011.
- Schirmermeister, L., Kunitsky, V., Grosse, G., Wetterich, S., Meyer, H., Schwamborn, G., Babiy, O., Derevyagin, A., and Siegert,  
 C.: Sedimentary characteristics and origin of the Late Pleistocene Ice Complex on north-east Siberian Arctic coastal lowlands  
 and islands – A review, *Quaternary International*, 241, 3-25, doi: 10.1016/j.quaint.2010.04.004, 2011.
- Schuur, E. A. G., McGuire, A. D., Schadel, C., Grosse, G., Harden, J. W., Hayes, D. J., Hugelius, G., Koven, C. D., Kuhry,  
 685 P., Lawrence, D. M., Natali, S. M., Olefeldt, D., Romanovsky, V. E., Schaefer, K., Turetsky, M. R., Treat, C. C., and Vonk, J.  
 E.: Climate change and the permafrost carbon feedback, *Nature*, 520, 171-179, doi: 10.1038/nature14338, 2015.
- Shur, Y., Hinkel, K. M., and Nelson, F. E.: The transient layer: implications for geocryology and climate-change science,  
*Permafrost and Periglacial Processes*, 16, 5-17, doi: 10.1002/ppp.518, 2005.



- Smith, L. C., Sheng, Y. W., and MacDonald, G. M.: A first pan-Arctic assessment of the influence of glaciation, permafrost,  
 690 topography and peatlands on northern hemisphere lake distribution, *Permafrost and Periglacial Processes*, 18, 201-208, doi:  
 10.1002/ppp.581, 2007.
- Smith, S., and Burgess, M. M.: Ground Temperature Database for Northern Canada, Geological Survey of Canada, Ottawa,  
 Open File Report 3954, 28, 2000.
- Strauss, J., Schirrmeister, L., Grosse, G., Fortier, D., Hugelius, G., Knoblauch, C., Romanovsky, V., Schädel, C., Schneider  
 695 von Deimling, T., Schuur, E. A. G., Shmelev, D., Ulrich, M., and Veremeeva, A.: Deep Yedoma permafrost: A synthesis of  
 depositional characteristics and carbon vulnerability, *Earth-Science Reviews*, 172, 75-86, doi:  
 10.1016/j.earscirev.2017.07.007, 2017.
- Stuiver, M., Reimer, P. J., and Reimer, R. W.: CALIB 7.1: <http://calib.org>, access: 2019-07-23, 2019.
- Tank, S. E., Vonk, J. E., Walvoord, M. A., McClelland, J. W., Laurion, I., and Abbott, B. W.: Landscape matters: Predicting  
 700 the biogeochemical effects of permafrost thaw on aquatic networks with a state factor approach. Submitted to *Permafrost and  
 Periglacial Processes*, manuscript # PPP-19-0058, in review.
- Turner, K. W., Wolfe, B. B., and Edwards, T. W. D.: Characterizing the role of hydrological processes on lake water balances  
 in the Old Crow Flats, Yukon Territory, Canada, using water isotope tracers, *Journal of Hydrology*, 386, 103-117, doi:  
 10.1016/j.jhydrol.2010.03.012, 2010.
- 705 van Everdingen, R.: Multi-language glossary of permafrost and related ground-ice terms. Boulder (CO): National Snow and  
 Ice Data Center/World Data Center for Glaciology, 1998 (revised 2005).
- Verpoorter, C., Kutser, T., Seekell, D. A., and Tranvik, L. J.: A global inventory of lakes based on high-resolution satellite  
 imagery, *Geophysical Research Letters*, 41, 6396-6402, doi: 10.1002/2014GL060641, 2014.
- Vonk, J. E., Mann, P. J., Davydov, S., Davydova, A., Spencer, R. G. M., Schade, J., Sobczak, W. V., Zimov, N., Zimov, S.,  
 710 Bulygina, E., Eglinton, T. I., and Holmes, R. M.: High biolability of ancient permafrost carbon upon thaw, *Geophysical  
 Research Letters*, 40, 2689-2693, doi: 10.1002/grl.50348, 2013.
- Walter, K. M., Edwards, M. E., Grosse, G., Zimov, S. A., and Chapin, F. S.: Thermokarst lakes as a source of atmospheric  
 CH<sub>4</sub> during the last deglaciation, *Science*, 318, 633-636, doi: 10.1126/science.1142924, 2007.
- Wik, M., Varner, R. K., Anthony, K. W., MacIntyre, S., and Bastviken, D.: Climate-sensitive northern lakes and ponds are  
 715 critical components of methane release, *Nature Geoscience*, 9, 99, doi: 10.1038/ngeo2578, 2016.
- Yoshikawa, K., and Hinzman, L. D.: Shrinking thermokarst ponds and groundwater dynamics in discontinuous permafrost  
 near council, Alaska, *Permafrost and Periglacial Processes*, 14, 151-160, doi: 10.1002/ppp.451, 2003.
- You, Y., Yu, Q., Pan, X., Wang, X., and Guo, L.: Geophysical Imaging of Permafrost and Talik Configuration Beneath a  
 Thermokarst Lake, *Permafrost and Periglacial Processes*, 28, 470-476, doi: 10.1002/ppp.1938, 2017.
- 720 Zimmermann, C., Poulin, M., and Pienitz, R.: Diatoms of North America: The Pliocene-Pleistocene freshwater flora of Bylot  
 Island, Nunavut, Canadian High Arctic. *Iconographia Diatomologica*, Vol. 21, A.R.G. Gantner Verlag, 2010.



## Tables


725 **Table 1: Radiocarbon ( $^{14}\text{C}$ ) dates obtained in sediment cores collected in Gull Lake. Other  $^{14}\text{C}$  dates obtained near the base of the frozen silt-peat terrace in the surroundings (see text for details) are also summarized (Allard, 1996; Fortier et al., 2006; Ellis and Rochefort, 2004; 2006; Ellis et al., 2008; unpublished data).**

Sample ID	Environment	Dated material	Depth (cm)	$^{14}\text{C}$ age (yr BP)	$\pm$	Calib. age (cal yr BP)	1- $\sigma$ range	Source
ULA-5673	aquatic	lake sed. (bulk)	9.25	<b>2025</b>	20	<b>1972</b>	1949 - 1995	
ULA-5672	aquatic	lake sed. (bulk)	10.25	<b>2100</b>	20	<b>2073</b>	2042 - 2117	
ULA-5959	aquatic	wood/plant	107.5	<b>4805</b>	15	<b>5505</b>	5488 - 5588	Bouchard et al.
ULA-4894	aquatic	lake sed. (bulk)	0.5	<b>1650</b>	20	<b>1552</b>	1533 - 1563	(this study)
ULA-4895	aquatic	lake sed. (bulk)	9.5	<b>2065</b>	20	<b>2032</b>	1993 - 2057	
ULA-4896	aquatic	peat	53.5	<b>2635</b>	20	<b>2756</b>	2748 - 2760	
Beta-143333	terrestrial	peat	297	<b>3100</b>	50	<b>3303</b>	3245 - 3374	Ellis and
Beta-143337	terrestrial	peat	229	<b>2590</b>	50	<b>2725</b>	2541 - 2772	Rochefort (2006)
Beta 143339	terrestrial	peat	209	<b>1660</b>	40	<b>1565</b>	1528 - 1611	Ellis and
Beta 152437	terrestrial	peat	182	<b>1470</b>	40	<b>1358</b>	1316 - 1385	Rochefort (2004)
UL-2356	terrestrial	wood + peat	233	<b>3670</b>	110	<b>4010</b>	3848 - 4151	Fortier et al.
UL-2152	terrestrial	wood + peat	241	<b>3270</b>	100	<b>3506</b>	3387 - 3607	(2006)
UL-1048	terrestrial	peat	135	<b>2210</b>	120	<b>2208</b>	2060 - 2346	
UL-1034	terrestrial	peat	230	<b>2510</b>	90	<b>2575</b>	2489 - 2739	Allard (1996)
UL-1035	terrestrial	peat	250	<b>2600</b>	90	<b>2687</b>	2496 - 2840	
UL-1025	terrestrial	peat	320	<b>2900</b>	90	<b>3045</b>	2894 - 3165	
ULA-6508	terrestrial	peat	82	<b>3045</b>	15	<b>3249</b>	3214 - 3323	Veillette (unpub. data)
UL-2427	terrestrial	peat	301	<b>3040</b>	90	<b>3228</b>	3080 - 3362	
UL-2614	terrestrial	peat	275	<b>3350</b>	90	<b>3593</b>	3475 - 3693	
UL-2418	terrestrial	wood	N/A	<b>3560</b>	90	<b>3855</b>	3720 - 3972	Fortier (unpub. data)
UL-2584	terrestrial	wood	N/A	<b>3300</b>	100	<b>3537</b>	3403 - 3640	
UL-2416	terrestrial	wood + peat	155	<b>3440</b>	100	<b>3706</b>	3586 - 3832	
UL-2264	terrestrial	peat	210	<b>2750</b>	90	<b>2869</b>	2765 - 2943	





## Figures



730 **Figure 1: Study area location and context.** a) Location of Bylot Island (Nunavut), Canada, within the continuous permafrost zone (source: Brown et al., 1998). Pleistocene ice-rich permafrost distribution in non-glaciated regions of Siberia and Alaska (Yedoma) is also shown (source: Strauss et al., 2017). b) Location of the study site, on the southwestern lowlands of Bylot Island (satellite photo: Terra-MODIS, 22 July 2012). b) Location of Gull Lake, in Qarlikturvik valley (glacier C-79 in the background). An early Holocene terminal moraine (TM) and a small outlet, draining towards “Gull Lake 2” (GL-2) and the proglacial river, are also shown.





Figure 2: Bathymetry and GPR survey lines conducted on Gull Lake. Sediment core location is shown (red “x”). GPR line cross-sections (12, 13, 14) are shown in Fig. 3. The central basin is deeper and surrounded by a shallow platform where degraded ice-wedge polygons are visible. The boundary between the central basin and the shallow platform is shown by the dashed line. Satellite image: GeoEye-1, 18 July 2010.



**Figure 3:** Interpreted GPR cross-sections obtained along survey lines (see Fig. 2 for line locations). The upper figure is the raw GPR profile with color arrows indicating distinct reflectors such as the base of the ice cover (light blue), lake bottom (brown), former surface of ice-wedge polygon ridges and troughs (green) and the top of the glacio-fluvial sand and gravel unit (red). Complete data are available at <https://doi.org/10.5885/45609XX-E3573955017A4904>.

745






Figure 4: Lithostratigraphy of a sediment core collected in Gull Lake in June 2015. The displayed CT-scan image, as well as visual descriptions and LOI data, were used to split the sedimentary sequence into 3 distinct units (lithozones). Complete data are available at <https://doi.org/10.5885/45612XX-AB27C20EB10D4509> (CT-scan) and at <https://doi.org/10.5885/45603XX-21852993EE434926> (LOI). CT-scan details are summarized in Supplement S1.

750



755 **Figure 5: Biostratigraphy (fossil diatoms) of a sediment core collected in Gull Lake in June 2015. Data are displayed as relative abundance (%) of dominant taxa, i.e. representing more than 5 % in at least one level. Complete data are available at <https://doi.org/10.5885/45600XX-C0960664FE8F4038>. Plates (photographs) of the most abundant species are shown in Supplement S2.**





760 **Figure 6: Four-stage conceptual model of thermokarst lake inception and evolution through the Holocene. a) Stage 0: initial**  
**conditions with networks of ice wedges developed in frozen silt-peat and glacio-fluvial sand and gravel (and likely reaching**  
**underlying marine silts and clays). A pre-existing topographic depression of 1-2 m was collecting drifting snow and meltwater. b) Stage 1:**  
**thermokarst inception, i.e. deepening of the active layer, melting of the top of ice wedges, development of a hummocky**  
 765 **surface. c) Stage 2: thermokarst pond coalescence, formation of a small lake with a maximum depth still above maximum ice cover**  
**thickness. d) Stage 3: thermokarst lake mature development by lateral expansion (thermal and mechanical erosion) and bottom**  
**deepening (subsidence). Lake maximum depth is now below maximum ice cover thickness, triggering the formation of a talik. e)**  
**Stage 4a: possible future evolution by lake infilling (gyttja accumulation). f) Stage 4b: possible future evolution by lake drainage**  
**(partial or complete) and re-activation of ice-wedge cracking and growth.**