



Spatio-temporal variability and decadal trends of snowmelt processes on Antarctic sea ice observed by satellite scatterometers

Stefanie Arndt¹, Christian Haas¹

¹Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, 27570 Bremerhaven, Germany

5 *Correspondence to:* Stefanie Arndt (stefanie.arndt@awi.de)

Abstract. The timing and intensity of snowmelt processes on sea ice are key drivers determining the seasonal sea-ice energy and mass budgets. In the Arctic, satellite passive microwave and radar observations have revealed a trend towards an earlier snowmelt onset during the last decades, which is an important aspect of Arctic amplification and sea ice decline. Around Antarctica, snowmelt on perennial ice is weak and very different than in the Arctic, with most snow surviving the summer.

10 Here we compile time series of snowmelt-onset dates on seasonal and perennial Antarctic sea ice from 1992 to 2014/15 using active microwave observations from European Remote Sensing Satellite (ERS-1/2), Quick Scatterometer (QSCAT) and Advanced Scatterometer (ASCAT) radar scatterometers. We define two snowmelt transition stages: A weak backscatter rise indicating the initial warming and metamorphism of the snowpack (pre-melt), followed by a rapid backscatter rise indicating the onset of thaw-freeze cycles (snowmelt).

15 Results show large interannual variability with an average pre-melt onset date of 29 November and melt onset of 10 December, respectively, on perennial ice, without any significant trends over the study period, consistent with the small trends of Antarctic sea ice extent. There was a latitudinal gradient from early snowmelt onsets in mid-November in the northern Weddell Sea to late (end-December) or even absent snowmelt conditions in the southern Weddell Sea.

20 We show that QSCAT Ku-band (13.4 GHz signal frequency) derived pre-melt and snowmelt onset dates are earlier by 25 and 11 days, respectively, than ERS and ASCAT C-band (5.6 GHz) derived dates. This offset has been considered when constructing the time series. Snowmelt onset dates from passive microwave observations (37 GHz) are later by 13 and 5 days than those from the scatterometers, respectively.

25 Based on these characteristic differences between melt onset dates observed by different microwave wavelengths, we developed a conceptual model which illustrates how the evolution of seasonal snow temperature profiles affects different microwave bands with different penetration depths. These suggest that future multi-frequency active/passive microwave satellite missions could be used to resolve melt processes throughout the vertical snow column.

1 Introduction

Sea-ice extent in the Southern Ocean has experienced large seasonal and inter-annual variations during the last decades, while ice extent has changed little on decadal time scales (Parkinson and Cavalieri, 2012; Stammerjohn et al., 2012; Turner et al.,



2015). This is in strong contrast to the Arctic, where sea ice extent has decreased strongly in all seasons (e.g. Meier et al., 2014). The strong seasonal variability of Antarctic sea ice impacts processes and interactions between atmosphere, sea ice and ocean, and is a key component driving the polar marine ecosystem (Massom et al., 2001). The presence of snow on the ice dramatically alters these interactions through its impact on ice thermodynamics, mass balance, and light transmission. The role of snow on Antarctic sea ice is exacerbated by the fact that the ice remains snow-covered in most regions throughout the summer (Massom et al., 2001).

However, during spring and summer the entire snow column experiences substantial seasonal changes in its physical properties associated with variations in snow temperature profiles, liquid water content, grain size distribution, and stratification (e.g. Haas et al., 2001; Massom et al., 2001; Nicolaus et al., 2009). Detecting these variations in the snowpack on both Arctic and Antarctic sea ice is highly relevant as they modify the energy and mass budgets of the ice-covered oceans, and strongly influence the retrieval of sea-ice parameters from satellite remote sensing algorithms (e.g. Willmes et al., 2014).

In contrast to sea ice in the Southern Ocean, snow and sea ice properties in the Arctic undergo strong changes during the seasonal cycle (Sturm and Massom, 2017). Due to different radiation and turbulent flux regimes of the Arctic atmosphere (e.g. Andreas and Ackley, 1982; Nicolaus et al., 2006) during the spring-summer transition liquid water forms rapidly within the snow. The subsequent albedo-feedback processes accelerate seasonal snowmelt and the disappearance of snow, and lead to widespread formation of surface melt ponds resulting in substantial alterations of dielectric properties of the ice and snow surface associated with increasing microwave emissivity and decreasing radar backscatter. This distinct seasonal cycle of surface properties and related microwave signatures of Arctic sea ice has been utilized to identify different snow and sea-ice melt stages and the time of melt onset from passive [e.g. Markus et al., 2009] and active satellite microwave remote sensing observations (e.g. Markus et al., 2009). Those satellite retrievals of melt onset in the Arctic have shown that melt onset occurred earlier by 1 to more than 10 days decade⁻¹ since 1979 when routine satellite observation commenced, consistent with warmer air temperatures and accelerated ice retreat in the Arctic during the recent past decades.

However, on Antarctic sea ice, the retrieval of snowmelt onset is more challenging as thawing and melting are subtle (Haas et al., 2001; Nicolaus et al., 2006; Nicolaus et al., 2009), whilst other seasonal snow processes affect the snow microwave properties. For example, during winter and spring, due to the widely thick snow which depresses the relatively thin sea ice below the water level, there is widespread flooding at the snow/ice interface and refreezing of slush creating snow-ice (e.g. Eicken et al., 1994). This process is referred to as flood-freeze cycling (Fritsen et al., 1998) and contributes significantly to the sea-ice mass budget particularly of first-year ice in the Southern Ocean (Eicken et al., 1995; Eicken et al., 1994; Haas et al., 2001; Jeffries et al., 1997; Nicolaus et al., 2009).

In contrast, during spring and summer, diurnal thaw-freeze cycles cause strong snow metamorphism. Under more intensive melting conditions, snow changes from the pendular to the funicular regime (e.g. Denoth, 1980) where the liquid snow melt



water percolates through the snowpack to lower, colder layers or to the ice surface where it refreezes to form superimposed ice (Arndt et al., 2016;Tison et al., 2008;Haas et al., 2008;Haas et al., 2001;Nicolaus et al., 2009;Willmes et al., 2009).

These different predominant snow processes and the absence of melt ponding result in different Antarctic sea ice microwave signatures than in the Arctic. The absence of longer periods with large amounts of liquid water within the snow mean that strong, persistent decreases of backscatter or increases of emissivity typical for Arctic sea ice do not frequently occur in the Antarctic. In contrast, strong snow metamorphism leads to large-grained, polygonal granular textured and salt-free (snow) grains and melt clusters (Colbeck, 1997), increasing radar volume- and surface scattering (Colbeck, 1997;Abdalati and Steffen, 1995;Onstott and Shuchman, 2004), and decreasing microwave emissivity (Willmes et al., 2006).

The presence of frequent diurnal thaw-freeze cycles has been utilized by Willmes et al. (2009) and Arndt et al. (2016) to develop algorithms to observe melt processes and to retrieve snowmelt onset dates from passive microwave radiometers, such as the Special Sensor Microwave/Imager (SSM/I). Their algorithms are based on analyses of the magnitude of diurnal 37 GHz, vertical polarized Brightness Temperature changes observed from ascending and descending morning and afternoon satellite passes approximately 12 hours apart. Both studies found large interannual snowmelt onset variability but no pronounced trends. So far, there have only been few studies of Antarctic sea ice melt onset using active microwave sensors, i.e. radars. Drinkwater and Liu (2000) applied an Arctic-like algorithm searching for rapid drops in radar backscatter. However, such drops are only occasionally observed on seasonal sea ice near the ice edge, or on the Larsen Ice Shelves (Bevan et al., 2018). On sea ice, drops in radar backscatter can also result from flooding events (Lytle and Ackley, 2001), potentially close to the time of complete deterioration of the ice. Drinkwater and Liu (2000) stated that such backscatter drop events are sparse and short lived. Therefore, their algorithm may not be easily applicable to most regions of Antarctic sea ice.


In contrast, Haas (2001) used C-band ERS scatterometer data from 1992 to 1999 to show that increasing radar backscatter from winter to summer is typical of perennial ice, and is consistent with the theoretical considerations of backscatter from metamorphic snow and superimposed ice discussed above. On average, backscatter increased by 5.6 dB during 96 days, commencing on 15 November.

In this study, we update and extend the work by Haas (2001) with new satellite scatterometer data to study if any long-term changes in Antarctic melt onset on sea ice emerged since 1992-99. For this purpose, we compile time series of European Remote Sensing (ERS)-1/2, QuikSCAT (QSCAT), and Advanced Scatterometer (ASCAT) scatterometer data from 1992 to 2015. A similar time series has been compiled and analyzed with QSCAT and ASCAT data for the Arctic (Mortin et al., 2014). We apply the algorithm of Haas (2001) to both perennial and seasonal ice regimes (Figure 1), and revise it to include early melt season effects on backscatter, defined as pre-melt phase. In addition, we use twice daily QSCAT observations between 1999 and 2009 to retrieve the magnitude of diurnal backscatter changes similar to earlier work with passive microwave observations discussed above (Arndt et al. (2016);Willmes et al. (2009); Figures 2 and 3).

There are two short overlap periods of ERS-2 and QSCAT in 1999/2000 and of QSCAT and ASCAT in 2008/9. During those we note clear differences in backscatter behavior and melt onset timing observed by QSCAT and the other scatterometers,



which we attribute to the different radar frequencies employed by ERS and ASCAT (C-band; 5.6 GHz), and QSCAT respectively (Ku-band; 13.4 GHz), and their different penetration depths (e.g. Ulaby et al., 1986). The time differences between melt onset detected by the C-band and Ku-band sensors were corrected to construct the long time series from 1992 to 2015. Finally, we compare results with previously published melt onset dates retrieved from satellite passive microwave sensors using frequencies of 37 GHz (Arndt et al., 2016). The time differences between those retrievals and the radar results are again discussed in the context of temporal snow evolution, penetration depth, and the sensitivity of 37 GHz signals primarily to the uppermost snow layers. The results obtained here demonstrate the potential to observe snow processes at different depths from space, opening new avenues for multi-sensor studies of energy and mass budgets of snow on sea ice in the Southern Ocean.



10

Figure 1. Map of Antarctica showing the 12 study locations in the perennial sea-ice zone (black/squares, locations 1-12, same as used by Haas (2001)) and 6 study locations on seasonal sea ice (blue/circles, regions A-F).



Figure 2. Time series of 6-day mean backscatter coefficients (σ^0) from 1992/1993 to 2014/2015 for the study locations on perennial sea ice (Figure 1, 1-12). Purple and blue lines show the scatterometer-derived snow pre-melt (σ^0_{pre}) and melt (σ^0_{melt}) onset dates when the algorithm criteria were met (Section 2.2.1). Red triangles indicate the snowmelt onset retrieved from diurnal backscatter variations ($d\sigma^0$) of QSCAT data. Green and yellow triangles show the Temporary Snowmelt Onset (TeSMO) and Continuous Snowmelt Onset (SMO) derived from passive microwave observations from Arndt et al. (2016).




Figure 3. As Figure 2, but for study sites on seasonal sea ice (Figure 1, A-F).

2. Snowmelt retrieval from satellite scatterometer data

2.1 Data set

5 All presented scatterometer data sets were obtained from the NASA Scatterometer Climate Record Pathfinder (SCP) project, sponsored by NASA (<http://www.scp.byu.edu/>). For the following analysis, all data are interpolated to a 25x25 km² SSM/I polar stereographic grid, using nearest-neighbor resampling.

10 The scatterometer data from the European Space Agency's (ESA) European Remote Sensing (ERS) 1 and 2 missions are provided as 6-day averages of vertical co-polarized C-band (5.3 GHz, wavelength of 5.7 cm) backscatter. This low temporal resolution was chosen in order to cover the polar oceans and maintain a reasonable stability of successive backscatter maps from this early mission (Ezraty and Cavanié, 1999). Backscatter σ^0 is obtained over a 500 km wide swath and normalized to an incident angle of 40°. ERS-1 was operational from 1991 to 1996, while ERS-2 continued observations until 2001.

15 The NASA QuikSCAT (QSCAT) mission acquired measurements from 1999 to 2009, i.e. overlapping with both the ERS and ASCAT observations by 2 years, respectively (see below). In contrast to the ERS scatterometer, QSCAT operated at Ku-band, i.e. 13.4 GHz, equivalent to shorter wavelengths (2.2 cm). In this study, we used the QSCAT Scatterometer Image Reconstruction (SIR; Long et al. (1993)) backscatter product in both polarizations (vertically and horizontally co-polarized) normalized to a 40° incident angle. The SIR algorithm enhances the spatial resolution of the daily data product to 4.45 km.



Data are available twice daily, with a morning pass between 04:00 and 12:00 and afternoon pass between 12:00 and 20:00 local time. This allows observations of diurnal backscatter variations caused by colder conditions in the morning and warmer conditions in the afternoon.

Finally, the ESA Advanced Scatterometer (ASCAT) was launched in 2006 on the MetOp-A and MetOp-B satellites. The scatterometer is an upgraded successor of the scatterometers onboard the ERS-1/2 platforms mentioned above, operating in C-band at 5.6 GHz in vertical co-polarization (VV). In this study, we also used the SIR product, similar to the QSCAT product, with a spatial resolution of 4.45 km. Due to a lower spatial coverage of ASCAT, the data are provided as 2-day averages.

To avoid biases of sea ice and snow signatures due to signals from larger amounts of open water in each pixel (e.g. Drinkwater and Liu, 2000), sea-ice concentration (SIC) data from Nimbus-7 SMMR and DMSP SSM/I-SSMIS Passive Microwave Data were used (Comiso, 2000). The data are available daily since 1978 on a 25 km SSM/I polar stereographic grid. During our study period, starting in 1992, ice concentration data of three different SSM/I sensors were used (F11 from January 1992 to 1995, F13 from May 1995 to December 2008, and F17 from December 2006 onwards).

2.2 Methods

We restrict our analysis of melt onset dates to 18 individual study locations of single pixels of scatterometer data carefully chosen to represent typical ice conditions in seasonal and perennial ice regimes (Figure 1). This single-pixel-based approach was chosen over more regional analyses to avoid blurring of backscatter signatures due to the large spatial and temporal variability of sea ice properties and drift. Like the studies of Markus et al. (2009), Mortin et al. (2014), Willmes et al. (2009), and Arndt et al. (2016) our approach is also Eulerian, i.e. not taking into account ice drift and advection for which there is little accurate data. However, as most Antarctic sea ice is first-year ice at the beginning of the melt season, Eulerian and Lagrangian approaches show similar temporal behavior of melt signals (Fig. 3 in Haas (2001)).

We applied two melt detection algorithms based on two different scatterometer observables: First we analyze daily mean backscatter σ^0 which is available from all satellites. Second, we analyze the magnitude of diurnal backscatter variations $d\sigma^0$, defined as the absolute difference between morning and evening satellite overpasses only available from QSCAT (Section 2.1).


Analyses for both algorithms were only carried out for locations where sea-ice concentration remained above 70% for at least three weeks into the melting season. This avoided contamination of results by wind-roughened water (Drinkwater and Liu, 2000), and effectively eliminated regions of deteriorating, thin ice where surface flooding and break up into small floes and brash ice may occur, e.g. in the marginal ice zone.

2.2.1 Melt onset retrieval from time series of daily mean backscatter

The seasonal cycle of surface properties of Antarctic perennial sea ice is evident in its backscatter time series with higher backscatter in summer than in winter (Haas, 2001). These are well visible in Figure 2 which show the complete time series



compiled for 12 perennial sea-ice regions: While location 1-7 are chosen from north to south in the Weddell Sea, all other locations are distributed throughout the perennial sea-ice regime around Antarctica (Figure 1). As expected before, all perennial sea-ice regions show a distinct seasonal cycle with a sharp increase during spring and a subsequent slow backscatter decrease towards autumn/winter (Figure 4 a). In contrast, the backscatter time series at 6 locations with seasonal sea-ice distributed around Antarctica (Figure 1) are much less systematic and show a much weaker seasonal cycle than those of perennial sea ice (Figures 3 and 4 b). However, Drinkwater and Liu (2000) defined the Antarctic-wide snowmelt onset by a sudden drop in the radar backscatter signal during spring. Their algorithm indicated widely missing onset dates on perennial sea ice, especially in the southwestern Weddell Sea, since radar backscatter behaves differently there than expected by their algorithm.



10

Figure 4. Typical annual time series of mean daily QSCAT radar backscatter (σ^0 , black) and its diurnal variations ($d\sigma^0$, red) for location 2 on perennial (a) and location D on seasonal sea ice (b) in 2003/2004. Colored vertical lines illustrate snow pre-melt (σ^0_{pre} , magenta), and melt (σ^0_{melt} , blue) onset derived from backscatter time series only, and snowmelt onset derived from QSCAT diurnal backscatter variations ($d\sigma^0$, red). Grey shaded areas indicate the time period with sea-ice concentration less than 70%, when the algorithm developed here cannot be applied. Bold lines denote the time period for the spring/summer transition retrieval analysis. Histograms in c and d show frequency of occurrence of the magnitude of diurnal backscatter variations $d\sigma^0$.

15



In the following extensive analysis of the seasonal cycles in the backscatter time series, we distinguish between two major snowmelt stages. Firstly, sporadic, temporal backscatter rises are defined as pre-melt, preceding the larger backscatter rises at melt onset. We interpret these subtle radar backscatter rises to be caused by changing physical properties associated with the spring warming of upper ice layers and with increases in brine volume (e.g. Nandan et al., 2017), or the appearance of liquid water in the pendular regime in the middle or lower snowpack. Secondly, the most distinct, largest rise in backscatter during spring or summer is defined as melt onset. At this time, the high intensity of thawing of the snowpack transitions the snow to the funicular regime when liquid water percolates to cause strong snow metamorphism and superimposed ice formation (Haas et al., 2001). Thaw-refreeze cycles may occur diurnally or with periods of several days. However, on Antarctic sea ice their intensity is usually insufficient to initiate albedo feedbacks with accelerated, irreversible snow melt and melt pond formation. Thawing can also be interrupted by periods of cooler conditions for periods of hours, days or weeks.

In order to retrieve every summer's pre-melt and melt onset, we split the time series into annual sections extending from the beginning of July of one year to end of June in the following year, with the spring and summer season approximately centered (Figure 4 a, b). Here, the time prior to pre-melt onset is referred to as winter, between pre-melt and snowmelt onset as pre-melt stage, after snowmelt onset as snowmelt stage, and after 31 January as autumn/winter again, as noted in Figure 4 a. Following the approach by Haas (2001), and to be consistent with their analysis of ERS data which are only available as 6-day averages (approximately one week), we first down-sampled the QSCAT and ASCAT data by averaging over 6-day intervals. Then, we analyze local backscatter maxima and their preceding local minima from three-point running means of each σ^0 -time series. We define the first instance after 1 October when the difference between a local maximum and the preceding minimum is larger than 2 dB as pre-melt onset (σ^0_{pre} , Figure 4 a, purple line). From there, we search for the instance when the difference between a local maximum and the preceding local minimum is larger than 3 dB. We define that instance as snowmelt onset (σ^0_{melt} , Figure 4 a, blue line). Note that the algorithm only responds to substantial, long-lived changes of snow microwave properties due to the 3-point smoothing of the 6-day averaged data. The thresholds are based on the average rise of the backscatter values during the spring/summer transition.

We apply this pre-melt and melt onset detection algorithm to both, perennial and seasonal sea ice. If no backscatter rises of the required magnitude are found during summer, no pre-melt or melt dates are assigned. This may occur in cooler years, on perennial ice located far south where melt can be small or absent, on seasonal ice when snowmelt is superimposed on other processes like flooding, or where the ice completely disintegrates shortly after snow melt commenced. For example, no melt onset was found at location D in 2003/2004 as shown in Figure 4 b.

2.2.2 Melt onset retrieval from diurnal backscatter variations observed by QSCAT

Willmes et al. (2009) described the onset of snow surface melt on Antarctic sea ice as the appearance of dominant diurnal thaw-freeze cycles in the surface layer. Analyzing the time series of the absolute difference between two daily brightness



temperature values from ascending and descending satellite passes (dT_B , 37GHz, vertically polarized), they found an increase in dT_B once temporary thawing commences in summer.

Since QSCAT data are also available twice a day from ascending and descending passes, the backscatter time series can be utilized to derive diurnal variations at the chosen locations. Therefore, we use the regionally adaptive approach by Arndt et al. (2016) to identify the snowmelt onset during austral spring: After applying a 5-day running mean to each $d\sigma^0$ time series between 1 October and 31 January, a histogram of $d\sigma^0$ values with a bin width of 0.5 dB is computed for each location. All histograms only contain data from 1 October until 31 January or until the sea-ice concentration drops below 70% (see Section 2.2). For example, Figures 4 c and d show typical histograms for the respective perennial and seasonal ice study locations. Generally, we found two kinds of histograms: unimodal and bimodal $d\sigma^0$ distributions. A mode is defined as a local maximum bounded by at least one lower bin on each side. Multimodal distributions of $d\sigma^0$ (Figure 4 c) indicate the presence of distinctly different diurnal backscatter values, which we interpret as different melt stages including differences in the strength of diurnal thaw-freeze cycles. Normally, at least two modes of the $d\sigma^0$ distributions can be observed in those cases. A histogram is considered to be unimodal if a single mode exceeds a fraction of more than 90% of all included data (Figure 4 d). Locations with unimodal distributions are not further considered in the following analysis, as they do not reveal the characteristic diurnal snow backscatter variations. For example, only few occurrences of strong diurnal backscatter variations were found at the seasonal ice locations, and therefore in most years a reliable melt onset from diurnal variations could not be observed there (Figure 4 b). In contrast, bimodal $d\sigma^0$ -distributions are widely detected on perennial sea ice (Figure 4 c). In a next step, an iterative threshold selection algorithm (Ridler and Calvard, 1978) is applied to the $d\sigma^0$ time series of each location with a multimodal distribution to derive individual $d\sigma^0$ -thresholds delineating winter from summer conditions. Finally, snowmelt onset is defined as the first time when $d\sigma^0$ exceeds the respective local threshold for at least 3 consecutive days (Figure 4 a).

3. Results

3.1 Perennial sea ice

25

30



Table 1: Mean snowmelt onset dates indicated in Figure 2 for the 12 study locations on perennial sea ice shown in Figure 1 (mean \pm 1 standard deviation).

Region	From scatterometer data			From passive microwave data	
	Pre-melt Onset	Snowmelt Onset	Diurnal thawing-refreezing Onset (2000/01-2008/09 only)	Temporary Snowmelt Onset (TeSMO)	Continuous Snowmelt Onset (SMO)
Northwestern Weddell Sea (region 1-4)	24 November \pm 16 days	06 December \pm 16 days	09 December \pm 9 days	13 December \pm 11 days	18 December \pm 13 days
Southeastern Weddell Sea (region 5-7)	27 November \pm 25 days	16 December \pm 19 days	19 December \pm 13 days	21 December \pm 11 days	09 January \pm 9 days
Bellingshausen Sea (region 8)	01 December \pm 29 days	04 December \pm 27 days	19 October \pm 20 days	19 October \pm 28 days	11 December \pm 19 days
Amundsen Sea (region 9-10)	24 November \pm 23 days	06 December \pm 18 days	02 December \pm 10 days	05 December \pm 16 days	21 December \pm 17 days
Ross Sea (region 11-12)	11 December \pm 18 days	15 December \pm 17 days	13 December \pm 8 days	16 December \pm 10 days	27 December \pm 7 days
All regions	29 November \pm 10 days	10 December \pm 12 days	09 December \pm 5 days	12 December \pm 8 days	21 December \pm 9 days




Figure 5. Summary of mean radar backscatter σ^0 and range of seasonal variability for all study locations during the complete 23-year study period (cf. Figures 2 and 3). Boxes are the first and third quartiles. Whiskers display the 20- and 80-percentiles. Circles indicate mean, horizontal lines median values. Abbreviations according to Figure 1: WS: Weddell Sea, BS/AS: Bellingshausen and Amundsen Seas, RS: Ross Sea.

As expected (Section 1), all backscatter time series show strongly increasing backscatter during the summer, although with some interannual variability (Figure 2). Mean backscatter and the range of seasonal variations are summarized in Figure 5. Mean perennial ice backscatter ranges between -12.0 and -16.2 dB, i.e. is variable between regions and consistently higher than on seasonal ice. Superimposed on mean backscatter are seasonal variations ranging between 1.6 dB at location 8 in the Bellingshausen Sea up to 4.7 dB in the northwestern Weddell Sea and western Ross Sea (locations 1-3 and 12; range between 20 and 80 percentiles in Figure 5). Separating for the different sensor classes indicates that, averaging all 12 study locations on perennial sea ice, the mean amplitude between seasonal minimum and maximum is 13.35dB for the QSCAT time period, while it is only 7.66 and 9.35dB for ERS and ASCAT, respectively (Figure 2).

Figure 2 also includes the pre-melt and melt onset dates between 1992/93 and 2014/15 retrieved with the algorithms described. Average pre-melt and melt onset dates in the different regions are summarized in Table 1. Note that in some years melt events could not be observed because the temporal behavior of backscatter changes did not possess the characteristic rises expected by our algorithms. On average, the initial pre-melt onset is on 29 November, 11 days prior to the actual melt onset on 10 December retrieved from rapidly rising daily backscatter and its diurnal variations. In the Weddell Sea, there are strong latitudinal differences with earlier snowmelt onset in the northwest and increasingly later snowmelt onset in the southeast: On average, the earliest pre-melt onsets are observed in mid-November in the northwestern Weddell Sea (locations 2 und 6), and in the Amundsen Sea (location 9), while the latest pre-melt onsets are found in mid-December in the Ross (locations 11 and 12) and southeastern Weddell Seas (location 7). Regions in the northwestern Weddell Sea (locations 2 and 3) and



Bellingshausen and Amundsen Seas (locations 8 and 9) had the earliest actual snowmelt dates, at the end of November, while again the southeasternmost region in the Weddell Sea (location 7) had the latest onset dates, in the end of December. Regions in the southeastern Weddell Sea (locations 5-7) and western Amundsen Sea (location 10) frequently do not reveal any distinct pre-melt or melt phases (Figure 2).

5 Snowmelt onset dates based on diurnal variations in the radar backscatter can only be derived with QSCAT data from 2000/01 to 2008/09. Here, on average, snowmelt onset occurred on 9 December, in an interval from 27 November (northwestern Weddell Sea, location 1) to 21 December (central/northwestern Weddell Sea and Ross Sea, locations 4, 5 and 12).

For comparison, Figure 2 and Table 1 also include Temporary SnowMelt Onset (TeSMO) and subsequent continuous SnowMelt Onset (SMO) retrieved from 37 GHz passive microwave brightness temperature observations [Arndt et al., 2016].

10 On average, TeSMO occurs on 12 December, i.e. 3 days later than snow melt observed by scatterometers. The average date of SMO occurs another 12 days later on 24 December. On average, the earliest TeSMO starts in mid-November in the Bellingshausen Sea (location 8), while the latest temporal snowmelt onsets are observed in the end of December in the southeastern Weddell Sea (location 6). Again, regions in the northwestern Weddell Sea (locations 1 and 2) showed the earliest SMO in the beginning of December, while the southernmost locations 5 and 6 show the latest SMO in mid-January.

15 Both, scatterometer and passive microwave observed melt onset parameters consistently show strong gradients towards later snowmelt from north to south as well as the decreasing occurrence of melt events towards the south, in particular in the Weddell Sea. However, we note strong differences in the actual timing of melt events observed by the different sensors which are further discussed in Section 4.4.


3.2 Seasonal sea ice

20 In addition to the perennial sea ice locations, 6 study locations on seasonal sea ice were chosen, distributed around the Antarctic continent (Figures 1 and 3). Mean backscatter ranges between -16.7 and -18.4 dB, i.e. much lower than on perennial ice (Figure 5). In agreement with previous studies discussed in the introduction, none of the seasonal sea-ice regions show any consistent seasonal backscatter cycle. While the Weddell and Ross Seas (locations A and D) show slight increases in backscatter during the transition from winter to summer, backscatter decreases at locations B and C in the Indian Ocean. Locations E and F in the
25 Bellingshausen and Amundsen Seas reveal very little backscatter variation throughout the spring and summer time. However, most regions show a sharp decrease in the backscatter signal with increasing sea-ice melt, consistent with (Drinkwater and Liu, 2000). As expected from the strongly different seasonal backscatter changes over seasonal sea ice compared to perennial ice, snowmelt onset on seasonal ice is only sporadically detected by our algorithm (Figure 3), demonstrating the spurious nature of backscatter signals on seasonal ice. At location A in the Weddell Sea pre-melt is frequently observed, with an average
30 date of 31 October. In contrast, hardly any pre-melt has been observed at locations B, C, and F in the Indian and Bellingshausen Seas. Overall, subsequent actual melt onset is rarely observed either, but most frequently occurring in the Weddell Sea. During the 23-year study period it was observed 8 times with an average melt onset date of 16 November. Snowmelt onset over seasonal ice could hardly be derived from passive microwave observations either (Arndt et al., 2016). However, the years and



regions when and where it was possible were similar to the ones when also the scatterometer data yielded results. On average, melt onset was observed on 12 November with the passive microwave data (Figure 3).

3.3 Compilation of snowmelt onset time series and analysis of long-term trends



5 **Figure 6.** Averaged time differences between pre-melt and snowmelt onset dates retrieved from QSCAT and ERS/ASCAT at the different study locations on perennial sea ice for the overlap periods 1999/2000 and 2008/2009, respectively. Negative (positive) differences indicate an earlier (later) transition detected from Ku-band. Boxes are the first and third quartiles. Whiskers display the 20- and 80-percentiles. Circles indicate mean, lines median values.

10 Similar to the work of Markus et al. (2009);Mortin et al. (2014);and Stroeve et al. (2014) in the Arctic, here we used the backscatter time series of ERS-1/2, QSCAT and ASCAT to study the inter-annual variability and trends of the previously described scatterometer-derived pre-melt and snowmelt onset dates. However, as ERS/ASCAT and QSCAT use different radar bands with different penetrations depths, and have been shown to retrieve different average melt onset dates, we first quantified the differences between retrieved melt onset dates during the overlap periods of ERS-2 and QSCAT in 1999/2000 and of
 15 QSCAT and ASCAT in 2008/2009. Figure 6 shows the average time differences between Ku-band and C-band retrievals for the 12 study locations on perennial sea ice (Figure 1), for both overlap periods, respectively. On average, Ku-band QSCAT data detected pre-melt and melt onset earlier by 25 and 11 days, respectively, than the C-band ERS-2 data. Similarly, QSCAT detected earlier pre-melt and melt onset dates than ASCAT. Pre-melt was observed 21 days earlier, while melt onset was observed 42 days earlier. However, the latter difference is only based on one sample from the Weddell Sea (location 1), while
 20 at all other perennial ice sites melt onset was not retrievable concurrently from both QSCAT and ASCAT data in 2008/09.




Figure 7. Time series of snowmelt onset dates in the perennial sea-ice zone for individual regions 1-12 (Figure 1), and all regions averaged (bottom right, upper panel). Grey circles (and grey line in bottom right) show the uncorrected QSCAT melt onset dates, while black symbols show dates corrected 11 days, the difference between Ku- and C-band melt onset dates, respectively.




Figure 8. Time series of snowmelt onset dates averaged for all regions derived from backscatter time series (σ_{melt}^0 , see Figure 7), diurnal backscatter variations ($d\sigma^0$), as well as the Temporary Snowmelt Onset (TeSMO) and Continuous Snowmelt Onset (SMO) derived from passive microwave observations.

5

Based on the results above, we corrected the QSCAT pre-melt and melt onset dates by 25 and 11 days, respectively, to derive a consistent time series of scatterometer-derived pre-melt and melt onset dates from 1992/93 to 2014/15 and to retrieve any potential trends. Figure 7 shows the resulting time series of the snowmelt onset for each study location on perennial sea ice. In addition, Figure 8 summarizes these results and also compares them with all other previously described sea ice snowmelt onset dates from scatterometer and passive microwave observations in Antarctica. Neither single locations nor the regional averages reveal any significant temporal trend in the retrieved snowmelt onset dates. Instead, the respective snowmelt onset dates show strong interannual variations with the tendency towards later onset dates.

10

Figure 8 and Table 1 also show that passive microwave observed TeSMO and SMO dates occur generally later than those observed by C-band scatterometer. More detail is provided in Figure 9 which summarizes the time differences between all observed pre-melt and melt onset dates for all locations. Results show, on average, that 67 (57) % of the pre-melt (snowmelt) onset dates retrieved from scatterometer observations occur earlier than from passive microwave observations. Particularly large negative differences are observed in the Weddell Sea, where on average about half (a quarter) of the pre-melt (snowmelt) onset differences are larger than 20 days. Overall, the mean pre-melt (snowmelt) onset difference between passive microwave and scatterometer observations is 13 (5) days, while only 46 (33) % of the differences are smaller than 10 days, and 36 (24) % are smaller than 20 days. The smallest differences are most notably in the western Amundsen Sea (location 10), while biggest differences are detected in the southernmost Weddell Sea regions (locations 4 and 6). Again, in the Weddell Sea there is a pronounced gradient of larger differences from northwest to southeast.

20

Similarly, melt onset from diurnal backscatter variations $d\sigma^0$ during the QSCAT era occurred earlier than passive microwave observed melt onset (Figure 9). However, these differences are much smaller than those from the actual time series above, with a mean (mode) or 4 (2) days.

25




Figure 9. Mean time differences between retrieved snowmelt onset dates from scatterometer observations (σ_{pre}^0 , σ_{melt}^0 , $d\sigma^0$) and from passive microwave observations (Temporary SnowMelt Onset, according to Arndt et al. (2016)) for all study locations. Snowmelt onset dates from passive microwave observations and backscatter time series are retrieved from 1992 to 2014/15, while the retrieval of diurnal backscatter variations $d\sigma^0$ is only performed with QSCAT from 2000 to 2008. Positive (negative) differences indicate a later (earlier) transition observed by scatterometers. Boxes are the first and third quartiles. Whiskers display the 20- and 80-percentiles. Circles indicate the mean, dashes the median. Abbreviations according to Figure 1: WS: Weddell Sea, BS/AS: Bellingshausen and Amundsen Seas, RS: Ross Sea.

10 4. Discussion

4.1 Regional differences of seasonal backscatter variations

The previously described seasonal cycle of snow processes and the related backscatter signal (Section 1) are typically found on perennial sea ice (Figures 2 and 4 a). The strong summer backscatter rise may be interrupted by few temporary backscatter drops, which can be due to local and temporary flooding or brief, strong melt events with high snow liquid water content associated with the transition towards a funicular snow regime (Figures 2 and 4 a). In the Weddell Sea section, both the magnitude of the seasonal cycle and the actual backscatter values decrease from northwest to southeast (Figure 5), with fewer melt onset dates detected in the South. This is consistent with generally colder and dryer climatic conditions further south, away from the marginal ice zone and close to the Antarctic ice sheet, where also warm air advection from the North is hampered by the Antarctic Circumpolar Trough [e.g. Simmonds and Keay, 2000; Turner et al., 2015]. Consequently, both seasonal and diurnal variations in the lower atmosphere and snowpack are weak leading to less snow metamorphism and melt in the southern part of the Antarctic sea-ice regime. The absence of distinct snowmelt processes in the southern Weddell Sea were also observed and described in previous studies on snowmelt detection from passive microwave observations [Arndt et al., 2016].



In contrast, most seasonal sea ice is not showing similarly strong seasonal backscatter cycles. Instead, the younger and thinner seasonal ice is warmer and more salty than perennial ice with larger brine volume at the snow/ice interface causing low backscatter through winter [Nicolaus et al., 2009; Yackel et al., 2007]. Also, repeated flood-freeze cycles during winter as well as flooding events during spring and summer cause low backscatter in winter and backscatter drops during spring and summer [Drinkwater and Liu, 2000]. In addition, the early break-up of seasonal sea ice associated with the formation of leads and thin ice between ice floes contributes to the declining backscatter coefficients. Moreover, due to the a comparatively large ocean heat flux of the Southern Ocean, Antarctic sea ice might strongly melt from below and even retreat completely before the pendular-funicular transition in the snowpack have started or even dominant snow melt can develop at its surface in summer. These numerous processes coincide and compete with actual surface snowmelt processes, meltwater percolation, and superimposed ice formation, making it therefore difficult to consistently retrieve snowmelt processes and dates on seasonal sea ice by scatterometer (Section 3.2) and passive microwave algorithms [Arndt et al., 2016]. We therefore consider the found snowmelt onset dates on Antarctic seasonal sea ice highly uncertain and potentially not meaningful.

4.2 Inter-annual variations in the retrieved snowmelt onset dates

A main achievement of our work is the compilation of a long time series of derived snowmelt onset dates on perennial sea ice from 1992/1993 to 2014/2015, and that no significant trend in the retrieved snowmelt onset dates was found, neither from scatterometer nor from passive microwave observations. Instead we found strong inter-annual variability of all 5 snowmelt onset parameters (Table 1). However, previous studies have shown that during the same time period the duration of the summer open water season, when the ice concentration is below 15% for at least 5 days, shows significant trends in some regions. Stammerjohn et al. (2008) showed that the period of open water was longer by 85 ± 20 days (total change from 1979 to 2004) in the Bellingshausen and Amundsen Seas (and somewhat less in the southern Weddell Sea), whereas in the perennial sea ice regime of the western Ross Sea it decreased by 60 ± 10 days. In general, that study revealed a significant correlation between ice-season duration and sea-ice advance in most regions of the Southern Ocean, while weak correlation was found with the respective sea-ice retreat dates. The absence of similar trends for melt indicates that the timing of snowmelt and snowmelt processes are not important processes for Antarctic sea ice extent variability. Instead, we suggest that seasonal and interannual ice concentration and extent variations rather depend on local oceanic conditions governing ocean heat flux and ice bottom melt, and on large-scale atmospheric circulation patterns, which affect the speed and direction of sea-ice drift and intensity of ice deformation and redistribution (Maksym et al., 2012; Stammerjohn et al., 2008; Turner et al., 2014; Turner et al., 2016). Our melt onset retrieval algorithms utilize seasonal changes of microwave properties caused by snowpack transitions from the pendular to the funicular snow regime associated with snow metamorphism, ultimately leading to the formation of superimposed ice (Haas, 2001). Changes of these properties also affect other retrieval algorithms, of, e.g., sea-ice concentration (Willmes et al., 2014) and of ice thickness from radar altimetry (e.g. Ricker et al., 2014). For example, increasing snow metamorphism may lead to reduced radar penetration and raising elevations of radar scattering horizons, and therefore to



retrievals of apparently increasing ice thicknesses. Combination of altimetric sea-ice thickness data and backscatter changes and melt processes observed by scattermeters may therefore help to improve our understanding of sea-ice surface processes and seasonal mass balance of Antarctic sea ice in the future.

4.3 Sensitivity of different microwave wavelengths to evolving snow temperature and moisture profiles during the melt season

5




Figure 10. Conceptual model of the evolution of vertical air, snow and sea ice temperature profiles as well as snow metamorphism processes in the Antarctic sea ice regime during spring and summer in four characteristic stages (a-d). Note that snow temperatures close to 0°C lead to the appearance of liquid water and strong snow metamorphism. Diurnal thaw-freeze cycles during the during stage b to d cause significant differences in the snowpack properties and air temperatures between day- and nighttime. Microwaves with different wavelengths are sensitive to changing snow properties at different depths (left panel), and therefore indicate different melt-onset dates in the same region.

10

An important result of this study is that the retrieved perennial ice snowmelt onset dates from Ku-band (QSCAT), C-band (ERS and ASCAT), and 37 GHz passive microwaves (PMW) were different. On average, melt was detected first with Ku-band, followed by C-band (11 days later on average), and then PMW (13 days later on average), see Figures 6, 8, 9, and Table 1). This behavior could be related to the different wavelengths of those three observations with their different penetrations depths, and different sensitivity to increasing Rayleigh scattering due to metamorphic processes with cycles of variable liquid water content, growing snow grains and ice layer formation (e.g. Onstott, 1992; Ulaby et al., 1986). However, the observed

15



behavior indicates that medium wavelength Ku-band signals with their specific wavelength sense these processes first, followed by longer wavelength C-band signals with their deeper penetration, and that short-wavelength PMW signals with the least penetration are affected last.

We therefore propose a conceptional model for the temporal evolution of snow column temperature, moisture, and metamorphism profiles which can qualitatively explain the observed microwave behavior and which suggests that depth-dependent changes of snow properties can be retrieved by multi-frequency satellite microwave observations. The model is illustrated in Figure 10.

During the spring and pre-melt phase, subject to strong diurnal variations, the snow and ice warm from above due to atmospheric turbulent and shortwave radiative heating, with the highest temperatures in the upper, interior snow layers while lower layers remain colder due to the cold, underlying sea ice with its large heat capacity (Cheng et al., 2003; Haas et al., 2001). However, at the very snow surface, temperatures can also remain lower than in the interior due to longwave radiative cooling of the surface, which results in a sub-surface temperature maximum (Figure 10 a), and can eventually lead to subsurface melt, as observed in, e.g., blue ice regions on the Antarctic ice sheet (e.g. Brandt and Warren, 1993; Cheng et al., 2003; Liston and Winther, 2005). While the magnitude of the difference between the snow surface temperature and the sub-surface temperature is debated (Brandt and Warren, 1993), the subsurface temperature maximum depends on the snow's extinction coefficient and is larger with denser snow, which is typical for perennial Antarctic sea ice (Nicolaus et al., 2009).

As the snow warming continues and successively affects lower and lower snow layers, first the interior and then lower layers reach near-melting or sporadic melting temperatures (Figure 10 b and c) with increases of liquid water content and diurnal or multi-day transitions into the funicular snow regime, when they are subject to strong, irreversible snow metamorphism and ice lenses or superimposed ice formation (Cheng et al., 2003; Haas et al., 2001). Only at a later stage, during instances of large atmospheric heat fluxes, the very snow surface becomes warm enough to be subjected to strong snow metamorphism as well (Figure 10 d).

The different stages in Figure 10 a-d are frequently interrupted or ending by colder periods which cause the prominent thaw/refreeze cycles. While the appearance of liquid water will generally reduce radar backscatter and increase microwave emissivity, it is the refreezing of the respective snow layers and irreversible snow metamorphism that causes the backscatter increases utilized in this paper.

With their intermediate penetration, it is therefore plausible that Ku-band signals sense initial snow property changes in the interior snow column first (Figure 10 b), while C-band and PMW signals receive their strongest contributions from the unchanged lower and topmost layers and therefore show no response at that time. C-band signals respond next, when the warming and sporadic wetting has reached the lower layers near the snow/ice interface (Figure 10 c). Finally, PMW signals, receiving their main power from the topmost snow layer, are only affected once the very surface is affected by thaw-refreeze cycles as well.

It should be noted that the penetration depth of both Ku- and C band into dry snow is at least more than a meter (Ulaby et al., 1986). However, increased backscatter along the propagation path through the snow at any depth will result in the observed



overall backscatter increases. The longer wavelengths of C-band signals may not be sensitive to the initial metamorphism in the interior snow layers which is already sensed by the shorter Ku-band signals (Figure 10 b). Overall, at the 12 study locations on perennial ice the mean amplitude between the seasonal minimum and maximum is 13.35dB for the QSCAT time period, but only 7.66 and 9.35dB for ERS and ASCAT, respectively (Figure 2). This also indicates that Ku-band backscatter is more strongly affected by a certain degree of snow metamorphism than C-band backscatter.

The mean time differences of 11 days between melt onset detected by Ku-band and C-band, and 13 days between C-band and PMW provides a reference for the time scales of initial warming and snow metamorphism in the Antarctic. Interestingly, in the Arctic Mortin et al. (2014) found hardly any temporal difference between melt onset dates observed by the same sensors. In agreement with our conceptual model above we interpret that behavior by the fact that snow on Arctic sea ice warms very rapidly throughout the entire snow column during melt-onset when it is already close to the melting temperature, and therefore all wavelength respond at approximately the same time. This is consistent with the generally very rapid melt and disappearance of snow on sea ice in the Arctic discussed in the introduction.

5. Summary and conclusions

In this study, we compiled a time series of snowmelt onset dates on Antarctic sea ice from 1992 onwards using different radar scatterometer observations (ERS-1/2, ASCAT: 5.3 GHz; QSCAT: 13.4 GHz), in extension of previous work by Haas (2001). Doing so, we defined two major snowmelt stages: Pre-melt onset associated with the initial warming and increasing appearance of liquid water in the snowpack, followed by snowmelt onset related to diurnal thawing and refreezing of the snowpack.

Results show that the magnitude of seasonal and diurnal backscatter variations is highly dependent on latitude, related to earlier and more frequent snowmelt in the north (mid of November in the northern Weddell Sea). In contrast, regions farther south and closer to the continent reveal weaker seasonal and diurnal backscatter variations, and therefore less snowmelt attributed to the prevalent atmospheric conditions of cold and dry air and the absence of warm air advection events from the North.

For practical reasons we were unable to update this time series to include the 2015/16 to 2018/19 summer seasons, when Antarctic sea ice extent in September had reached record minima (e.g. Schlosser et al., 2018). It would be interesting to see in a follow up study if significantly earlier melt onset dates would be observed during those recent years. However, we believe that they will not, based on the facts that previous changes of ice extent did not strongly affect melt onset dates as mentioned above, that those recent changes probably have a strong contribution from oceanic processes, and that most potential surface changes may occur closer to the marginal ice zone where our algorithm is not applicable.

Based on the observed successive timing of melt events retrieved from different sensors, i.e. C- and Ku-band radars and 37 GHz passive microwave radiometers, we developed a conceptual model of the temporal evolution of snow temperature and metamorphism and their effect on, and detectability by, different microwave sensors during the spring/summer transition: The model explains qualitatively how snow metamorphism occurs first in interior snow layers and mostly affects Ku-band signals. Once warming has reached the lower snow pack the resulting metamorphism there can be detected by C-band signals. The



topmost snow layer remains coldest the longest by radiative cooling. Only when it warms and is affected by thaw/refreeze cycles at last do short wavelength passive microwave signals respond as well.

The results obtained here demonstrate the potential to observe snow processes at different depths from space, opening new avenues for studies of energy and mass budgets of snow on sea ice in the Southern Ocean. In addition, improved observations of snow metamorphism will contribute to better understanding and correction of uncertainties and spatial variability of space-borne retrievals of sea-ice concentration, snow depth and sea-ice thickness. However, improved microwave modeling informed by new in-situ observations of microwave properties and snow melt processes on Antarctic sea ice are required first to inform such activities.

Acknowledgement

Satellite data of radar backscatter were kindly provided by the Scatterometer Climate Record Pathfinder (SCP) project, sponsored by NASA (<http://www.scp.byu.edu/>), and sea-ice concentration data from the NASA National Snow and Ice Data Center Distributed Active Archive Center, Boulder, Colorado, USA. We acknowledge discussions about microwave emission and scattering behavior with Wolfgang Dierking, Sascha Willmes, and Marcus Huntemann. The work was funded by the Helmholtz Alliance “Remote Sensing and Earth System Dynamics” (HA-310) and the Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, Bremerhaven, Germany.

References

- Abdalati, W., and Steffen, K.: Passive Microwave-Derived Snow Melt Regions on the Greenland Ice-Sheet, *Geophysical Research Letters*, 22, 787-790, Doi 10.1029/95gl00433, 1995.
- Andreas, E. L., and Ackley, S. F.: On the differences in ablation seasons of Arctic and Antarctic sea ice, *Journal of the Atmospheric Sciences*, 39, 440-447, 10.1175/1520-0469, 1982.
- Arndt, S., Willmes, S., Dierking, W., and Nicolaus, M.: Timing and regional patterns of snowmelt on Antarctic sea ice from passive microwave satellite observations, *Journal of Geophysical Research - Oceans*, 121, 5916-5930, 10.1002/2015JC011504, 2016.
- Bevan, S. L., Luckman, A. J., Kuipers Munneke, P., Hubbard, B., Kulessa, B., Ashmore, D. W. J. E., and Science, S.: Decline in Surface Melt Duration on Larsen C Ice Shelf Revealed by The Advanced Scatterometer (ASCAT), 5, 578-591, 2018.
- Brandt, R. E., and Warren, S. G.: Solar-heating rates and temperature profiles in Antarctic snow and ice, *Journal of Glaciology*, 39, 99-110, 1993.
- Cheng, B., Vihma, T., and Launiainen, J.: Modelling of superimposed ice formation and subsurface melting in the Baltic Sea, *Geophysica*, 39, 31-50, 2003.
- Colbeck, S. C.: A Review of Sintering in Seasonal Snow, CRREL Report, DTIC Document, 1997.
- Denoth, A.: The pendular-funicular liquid transition in snow, *Journal of Glaciology*, 25, 93-98, 1980.



- Drinkwater, M. R., and Liu, X.: Seasonal to interannual variability in Antarctic sea-ice surface melt, *Ieee Transactions on Geoscience and Remote Sensing*, 38, 1827-1842, Doi 10.1109/36.851767, 2000.
- Eicken, H., Lange, M. A., Hubberten, H. W., and Wadhams, P.: Characteristics and distribution patterns of snow and meteoric ice in the Weddell Sea and their contribution to the mass balance of sea ice, *Annales Geophysicae-Atmospheres Hydrospheres and Space Sciences*, 12, 80-93, 10.1007/s00585-994-0080-x, 1994.
- Eicken, H., Fischer, H., and Lemke, P.: Effects of the snow cover on Antarctic sea ice and potential modulation of its response to climate change, *Annals of Glaciology*, 21, 369-376, 10.3189/S0260305500016086, 1995.
- Ezraty, R., and Cavané, A.: Intercomparison of backscatter maps over Arctic sea ice from NSCAT and the ERS scatterometer, *Journal of Geophysical Research: Oceans*, 104, 11471-11483, 1999.
- 10 Fritsen, C. H., Ackley, S. F., Kremer, J. N., and Sullivan, C. W.: Flood-Freeze Cycles and Microalgal Dynamics in Antarctic Pack Ice, Antarctic sea ice: biological processes, interactions and variability, 1-21, 10.1029/AR073p0001, 1998.
- Haas, C.: The seasonal cycle of ERS scatterometer signatures over perennial Antarctic sea ice and associated surface ice properties and processes, *Ann Glaciol*, 33, 69-73, 10.3189/172756401781818301, 2001.
- 15 Haas, C., Thomas, D. N., and Bareiss, J.: Surface properties and processes of perennial Antarctic sea ice in summer, *Journal of Glaciology*, 47, 613-625, 10.3189/172756501781831864, 2001.
- Haas, C., Nicolaus, M., Willmes, S., Worby, A., and Flinspach, D.: Sea ice and snow thickness and physical properties of an ice floe in the western Weddell Sea and their changes during spring warming, *Deep-Sea Res Pt II*, 55, 963-974, 10.1016/J.Dsr2.2007.12.020, 2008.
- Jeffries, M. O., Worby, A. P., Morris, K., and Weeks, W. F.: Seasonal variations in the properties and structural composition of sea ice and snow cover in the Bellingshausen and Amundsen Seas, Antarctica, *Journal of Glaciology*, 43, 138-151, 10.3189/S0022143000002902, 1997.
- 20 Liston, G. E., and Winther, J.-G.: Antarctic surface and subsurface snow and ice melt fluxes, *Journal of Climate*, 18, 1469-1481, 2005.
- Long, D. G., Hardin, P. J., and Whiting, P. T.: Resolution Enhancement of Spaceborne Scatterometer Data, *Ieee Transactions on Geoscience and Remote Sensing*, 31, 700-715, Doi 10.1109/36.225536, 1993.
- Lytle, V. I., and Ackley, S. F.: Snow-ice growth: a fresh-water flux inhibiting deep convection in the Weddell Sea, Antarctica, *Ann Glaciol*, 33, 45-50, Doi 10.3189/172756401781818752, 2001.
- 25 Maksym, T., Stammerjohn, S. E., Ackley, S., and Massom, R.: Antarctic Sea Ice-A Polar Opposite?, *Oceanography*, 25, 140-151, 10.5670/oceanog.2012.88, 2012.
- Markus, T., Stroeve, J. C., and Miller, J. A.: Recent changes in Arctic sea ice melt onset, freezeup, and melt season length, *Journal of Geophysical Research*, 114, C12024, 10.1029/2009jc005436, 2009.
- 30 Massom, R. A., Eicken, H., Haas, C., Jeffries, M. O., Drinkwater, M. R., Sturm, M., Worby, A. P., Wu, X. R., Lytle, V. I., Ushio, S., Morris, K., Reid, P. A., Warren, S. G., and Allison, I.: Snow on Antarctic Sea ice, *Rev Geophys*, 39, 413-445, 10.1029/2000rg000085, 2001.
- Meier, W. N., Hovelsrud, G. K., Oort, B. E., Key, J. R., Kovacs, K. M., Michel, C., Haas, C., Granskog, M. A., Gerland, S., and Perovich, D. K.: Arctic sea ice in transformation: A review of recent observed changes and impacts on biology and human activity, *Rev Geophys*, 52, 185-217, 2014.
- 35 Mortin, J., Howell, S. E. L., Wang, L. B., Derksen, C., Svensson, G., Graverson, R. G., and Schroder, T. M.: Extending the QuikSCAT record of seasonal melt-freeze transitions over Arctic sea ice using ASCAT, *Remote Sensing of Environment*, 141, 214-230, Doi 10.1016/J.Rse.2013.11.004, 2014.



- Nandan, V., Geldsetzer, T., Mahmud, M., Yackel, J., and Ramjan, S.: Ku-, X- and C-Band Microwave Backscatter Indices from Saline Snow Covers on Arctic First-Year Sea Ice, *Remote Sensing*, 9, 757, 2017.
- Nicolaus, M., Haas, C., Bareiss, J., and Willmes, S.: A model study of differences of snow thinning on Arctic and Antarctic first-year sea ice during spring and summer, *Annals of Glaciology*, 44, 147-153, 2006.
- 5 Nicolaus, M., Haas, C., and Willmes, S.: Evolution of first-year and second-year snow properties on sea ice in the Weddell Sea during spring-summer transition, *Journal of Geophysical Research*, 114, 10.1029/2008JD011227, 2009.
- Onstott, R., and Shuchman, R.: SAR measurements of sea ice, Chapter, 3, 81-115, 2004.
- Onstott, R. G.: SAR and scatterometer signatures of sea ice, *Microwave remote sensing of sea ice*, 73-104, 1992.
- 10 Parkinson, C. L., and Cavalieri, D. J.: Antarctic sea ice variability and trends, 1979-2010, *Cryosphere*, 6, 871-880, 10.5194/tc-6-871-2012, 2012.
- Ricker, R., Hendricks, S., Helm, V., Skourup, H., and Davidson, M.: Sensitivity of CryoSat-2 Arctic sea-ice freeboard and thickness on radar-waveform interpretation, *The Cryosphere*, 8, 1607-1622, 10.5194/tc-8-1607-2014, 2014.
- Ridler, T., and Calvard, S.: Picture thresholding using an iterative selection method, *IEEE trans syst Man Cybern*, 8, 630-632, 1978.
- 15 Schlosser, E., Haumann, F. A., and Raphael, M. N.: Atmospheric influences on the anomalous 2016 Antarctic sea ice decay, *Cryosphere*, 12, 1103-1119, 10.5194/tc-12-1103-2018, 2018.
- Stammerjohn, S., Martinson, D. G., Smith, R. C., Yuan, X. J., and Rind, D.: Trends in Antarctic annual sea ice retreat and advance and their relation to El Niño -Southern Oscillation and Southern Annular Mode variability, *Journal of Geophysical Research: Oceans*, 113, 2008.
- Stammerjohn, S., Massom, R., Rind, D., and Martinson, D.: Regions of rapid sea ice change: An inter-hemispheric seasonal comparison, *Geophysical Research Letters*, 39, 10.1029/2012gl050874, 2012.
- 20 Stroeve, J. C., Markus, T., Boisvert, L., Miller, J., and Barrett, A.: Changes in Arctic melt season and implications for sea ice loss, *Geophysical Research Letters*, 41, 1216-1225, Doi 10.1002/2013gl058951, 2014.
- Sturm, M., and Massom, R. A.: *Snow & Sea Ice*, in: *Sea Ice*. 3rd Edition, edited by: Thomas, D., Wiley-Blackwell, New York (UAS) & Oxford (UK), 65-109, 2017.
- 25 Tison, J.-L., Worby, A., Delille, B., Brabant, F., Papadimitriou, S., Thomas, D., De Jong, J., Lannuzel, D., and Haas, C. J. D. S. R. P. I. T. S. i. O.: Temporal evolution of decaying summer first-year sea ice in the Western Weddell Sea, *Antarctica*, 55, 975-987, 2008.
- Turner, J., Barrand, N. E., Bracegirdle, T. J., Convey, P., Hodgson, D. A., Jarvis, M., Jenkins, A., Marshall, G., Meredith, M. P., Roscoe, H., Shanklin, J., French, J., Goosse, H., Guglielmin, M., Gutt, J., Jacobs, S., Kennicutt, M. C., Masson-Delmotte, V., Mayewski, P., Navarro, F., Robinson, S., Scambos, T., Sparrow, M., Summerhayes, C., Speer, K., and Klepikov, A.: Antarctic climate change and the environment: an update, *Polar Rec*, 50, 237-259, 10.1017/S0032247413000296, 2014.
- 30 Turner, J., Hosking, J. S., Bracegirdle, T. J., Marshall, G. J., and Phillips, T.: Recent changes in Antarctic sea ice, *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 373, 20140163, 2015.
- Turner, J., Hosking, J. S., Marshall, G. J., Phillips, T., and Bracegirdle, T. J.: Antarctic sea ice increase consistent with intrinsic variability of the Amundsen Sea Low, *Climate Dynamics*, 46, 2391-2402, 10.1007/s00382-015-2708-9, 2016.
- 35 Ulaby, F. T., Moore, R. K., and Fung, A. K.: *Microwave Remote Sensing. Active and Passive. Vol. 3. From Theory to Applications.*, Addison Wesley Pub., London, U.K., 1065-2162 pp., 1986.



Willmes, S., Bareiss, J., Haas, C., and Nicolaus, M.: The importance of diurnal processes for the seasonal cycle of sea-ice microwave brightness temperatures during early summer in the Weddell Sea, Antarctica, *Annals of Glaciology*, 44, 297-302, 2006.

Willmes, S., Haas, C., Nicolaus, M., and Bareiss, J.: Satellite microwave observations of the interannual variability of snowmelt on sea ice in the Southern Ocean, *Journal of Geophysical Research-Oceans*, 114, 10.1029/2008jc004919, 2009.

- 5 Willmes, S., Nicolaus, M., and Haas, C.: The microwave emissivity variability of snow covered first-year sea ice from late winter to early summer: a model study, *The Cryosphere*, 8, 891-904, 2014.