

## Response to „Review by Editor from 28 June 2020“

Dear Jean-Louis,

We highly appreciate the great and intense work you put into revising our manuscript as your critical questions and remarks throughout the review process considerably improved our manuscript.

Regarding the given comments in the manuscript, we adjusted the following in the manuscript:

First of all, we implemented all suggested language/grammatical changes.

Comment 1: We agree that the term “snow redistribution” promises more than the manuscript is providing. We therefore followed the suggestion to change that towards “snow distribution” throughout the manuscript.

Comment 2: We added the reference Dieckmann et al. (1986).

Comment 3: We agree that this sentence was misleading and therefore rephrased it to [Atka Bay is seasonally sea-ice covered, and the water depth ranges between 80 m and 275 m with maximum depth in the central bay \(Kipfstuhl, 1991\).](#)

Comment 4: We deleted the duplicate reference.

Comment 5: We agree that the described missing melting is related to surface melting only. The sentence is changed towards: [In the following summer, the ice does not disappear by surface melting in-situ, but breaks up and drifts out of the bay once the conditions are sufficiently unstable.](#)

Comment 6/7: The given time period of 2010/11 to 2018/19 represents the time span of the continuous measurements in Atka Bay. This is why we focus our study on that period and highlight during that time the exceptional years 2012 and 2014.

Comment 8: We added the meant direction and changed therefore the sentence towards: [At each sampling site, up to five measurements are taken in an undisturbed area, one as the center measurement and four more at a distance of approx. 5 meters in each direction \(north, east, south, and west\), in order to account for the spatial variability of sea-ice and snow properties.](#)

Comment 9: We agree that for these calculations, cases of dry and flooded snow need to be distinguished. We therefore add a second equation, taking also the latter case into account: [As soon as F gets negative, a flooding of the snow/ice interface is assumed, and with that the formation of snow ice. As the latter is assumed to have the same density as sea ice \(Knight, 1988\), freeboard is calculated for the flooded case as](#)

$$-F = S - \frac{I + S - P \left( \frac{\rho_P - \rho_W}{\rho_W - \rho_I} \right)}{1 + \frac{\rho_S}{\rho_W - \rho_I}}$$

Comment 10: Thanks for that comment; we actually calculated the density of the platelet ice layer as suggested by you but phrased it in the manuscript in a “misunderstanding” way. We therefore rephrased it to:

[The platelet-layer density  \$\rho\_P\$  is calculated by means of the platelet-layer ice volume fraction  \$\beta\$  as  \$\rho\_P = \beta \cdot \rho\_I + \(1 - \beta\) \cdot \rho\_W\$ .](#)

Comment 11: We rephrased the sentence towards: [Colors indicate the relative frequency of each shown wind direction to wind speed pair.](#)

Comment 12: Due to the adjusted calculation, the average value between measured and calculated freeboard values is “negative” zero (-0,0045 m). The new statistics add up to: [According to Equation 1.1, 66% of the calculated freeboard values are smaller than the measured values. The difference between measured and calculated freeboard values ranges from -0.54 to 1.11 m with an average of  \$0.00 \pm 0.19\$  m. Neglecting the underlying buoyant platelet layer in the calculation reduces the freeboard by  \$0.07 \pm 0.15\$  m, whereas neglecting the snow layer on top of the sea ice increases the freeboard by  \$0.19 \pm 0.29\$  m \(Figure 5\).](#)

Comment 13: We agree and changed the gray scale towards the same as in Figure 4. However, we stick to the setting of the figure itself for consistency with the previous figure. Also, there is platelet ice visible, e.g., at ATKA11 in 2012.

Comment 14: The decreasing snow depth towards the ice shelf might be related to both, the easterly winds and the edge effect as shown in the Discussions.

Comment 15: We used consistently R throughout the manuscript.

Comment 16: We do not have an extensive data set on snow quality/properties in Atka Bay and can therefore not provide a proper comparison between the pack ice and fast ice area, respectively.

Comment 17: We agree and changed the sentence towards:

It is particularly remarkable that the average annual maximum plateau-layer thickness of 4 m (Table 1) is consistent with an earlier investigation at Atka Bay [...]

Comment 18: We agree and added the ice rumples and ice rise also in our Figure 1c and do reference it instead of Hoppmann et al. (2015).

Comment 19/20: We agree that the way, the sentence is written, is misleading. We therefore rephrased it to:  
A south-north survey across the bay at the beginning of austral summer 2018, however, revealed a trend of decreasing snow depth towards the northern fast-ice edge, with a stronger gradient approx. 5 km from the ice edge (Figure 7), which is in line with the northern boundary of the ice-shelf edge (Figure 1) and can therefore be explained by associated decreasing effects of the prevailing offshore winds. Consequently, also the measured snow depth is less in that part of the bay.

Comment 21: We agree that “linear” is kind of misleading in the given context. We therefore rephrased it to:  
In contrast, the evolution of the second-year fast-ice thickness shows a different pattern.

Comment 22: We do consider Atka Bay as “key region” of the Southern Ocean as here the interactions and feedback mechanisms can not only be studied between atmosphere, sea ice and ocean, but also includes governing interactions to the neighboring ice shelves. Furthermore, the Weddell Sea is one of the key regions of the global deep-water formation and contains at the same time relatively warm subsurface waters which are predicted to eventually push under the Filchner Ronne Iceshelf in the near future as stated in Vernet et al. (2019).

Comment 23: We agree (as mentioned in Comment 1) that Snow redistribution might be too ambiguous in the given context. We therefore rephrased it also here towards snow distribution only.

Comment 24: We feel super sorry that we missed to add you in the acknowledgements, Jean-Louis. Of course, we changed that immediately (and owe you a beer at the next opportunity!) ☺

# 1 Seasonal and interannual variability of landfast sea ice in Atka Bay, 2 Weddell Sea, Antarctica

3 Stefanie Arndt<sup>1</sup>, Mario Hoppmann<sup>1</sup>, Holger Schmithüsen<sup>1</sup>, Alexander D. Fraser<sup>2,3</sup>, Marcel Nicolaus<sup>1</sup>

4 <sup>1</sup>Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und Meeresforschung, 27570 Bremerhaven, Germany

5 <sup>2</sup>Institute for Marine and Antarctic Studies, University of Tasmania, Hobart 7001, Tasmania, Australia

6 <sup>3</sup>Antarctic Climate & Ecosystems Cooperative Research Centre, University of Tasmania, Hobart 7001, Tasmania, Australia

7 *Correspondence to:* Stefanie Arndt (stefanie.arndt@awi.de)

8 **Abstract.** Landfast sea ice (fast ice) attached to Antarctic (near-)coastal elements is a critical [element-component](#) of the local  
9 physical and ecological systems. Through its direct coupling with the atmosphere and ocean, fast ice properties are also a  
10 potential indicator of processes related to a changing climate. However, in-situ fast-ice observations in Antarctica are  
11 extremely sparse because of logistical challenges and harsh environmental conditions. Since 2010, a monitoring program  
12 observing the seasonal evolution of fast ice in Atka Bay has been conducted as part of the Antarctic Fast Ice Network (AFIN).  
13 The bay is located on the north-eastern edge of Ekström Ice Shelf in the eastern Weddell Sea, close to the German wintering  
14 station Neumayer III. A number of sampling sites have been regularly revisited [each year](#) between annual ice formation and  
15 breakup ~~each year~~ to obtain a continuous record of sea-ice and sub-ice platelet-layer thickness, as well as snow depth and  
16 freeboard across the bay.

17 Here, we present the time series of these measurements over the last nine years. Combining them with observations from the  
18 nearby Neumayer III meteorological observatory as well as auxiliary satellite images enables us to relate the seasonal and  
19 interannual fast-ice cycle to the factors that influence ~~its~~ [their](#) evolution.

20 On average, the annual consolidated fast-ice thickness at the end of the growth season is about two meters, with a loose platelet  
21 layer of four meter thickness beneath, and 0.70 meter thick snow on top. Results highlight the predominately seasonal character  
22 of the fast-ice regime in Atka Bay without a significant interannual trend in any of the observed variables over the nine-year  
23 observation period. Also, no changes are evident when comparing with sporadic measurements in the 1980s and 90s.  
24 ~~However,~~ [It is shown that](#) strong easterly winds in the area govern the year-round snow ~~re~~ distribution and also trigger the  
25 breakup of fast ice in the bay during summer months.

26 Due to the substantial snow accumulation on the fast ice, a characteristic feature is frequent negative freeboard, associated  
27 flooding of the snow/ice interface, and a likely subsequent snow ice formation. The buoyant platelet layer beneath negates the  
28 snow weight to some extent, but snow thermodynamics is identified as the main driver of the energy and mass budgets for the  
29 fast-ice cover in Atka Bay.

30 The new knowledge of the seasonal and interannual variability of fast-ice properties ~~in~~ from the present study helps to improve  
31 our understanding of interactions between atmosphere, fast ice, ocean and ice shelves in one of the key regions of Antarctica,  
32 and ~~lays the foundation~~ calls for ~~more~~ intensified multi-disciplinary studies in this region.

## 33 **1 Introduction**

34 The highly dynamic pack ice of the open polar oceans is continuously in motion under the influence of winds and ocean  
35 currents (Kwok et al., 2017). In contrast, landfast sea ice (short: fast ice) is attached to the coast or associated geographical  
36 features, such as for example a shallow seafloor (especially in Arctic regions) or grounded icebergs, and is therefore immobile  
37 (JCOMM Expert Team on Sea Ice, 2015). Fast ice is a predominant and characteristic feature of the Arctic (Dammann et al.,  
38 2019; Yu et al., 2014) and Antarctic coasts (Fraser et al., 2012), especially in winter. Its seaward edge may vary between just  
39 a few meters and several hundred kilometers from where it is attached to, mostly depending on the local topography and  
40 coastline morphology. The main processes for fast-ice formation are either in-situ thermodynamic growth, or dynamic  
41 thickening and subsequent attachment of ice floes of any age to the shore (Mahoney et al., 2007b).

42 In the Arctic, coastal regions that are characterized by an extensive fast-ice cover in winter are for example found in the  
43 Chukchi Sea and Beaufort Sea (Druckenmiller et al., 2009; Mahoney et al., 2014; Mahoney et al., 2007a), the Canadian Arctic  
44 (Galley et al., 2012), the East Siberian and Laptev Seas (e.g. Selyuzhenok et al., 2017), and the Kara Sea (Olason, 2016).

45 While the fast-ice cover in these regions comes with its own particular impacts on the respective coastal systems, what is a  
46 common to them feature is that they have undergone substantial changes in recent decades (Yu et al., 2014). These include a  
47 reduction of fast-ice area (Divine et al., 2003), later formation and earlier disappearance (Selyuzhenok et al., 2015) and a  
48 reduction of thickness (Polyakov et al., 2003).

49 Along the Antarctic coastline, the fast-ice belt extends even further from the coast (Fraser et al., 2012; Giles et al., 2008) due  
50 to the presence of grounded icebergs in much deeper waters of up to several hundred meters (Massom et al., 2001a).  
51 Embayments and grounded icebergs provide additional protection against storms and currents, and are often favorable for the  
52 formation of a recurrent and persisting fast-ice cover (Giles et al., 2008). Fast-ice around Antarctica is still usually seasonal  
53 rather than perennial, and reaches thicknesses of around 2 meters (Jeffries et al., 1998; Leonard et al., 2006), although it may  
54 attain greater ages and thicknesses in some regions (Massom et al., 2010). It mostly forms and breaks up annually as a response  
55 to various environmental conditions, such as heavy storms (Fraser et al., 2012; Heil, 2006). Its immediate response to both  
56 local atmospheric conditions and lower latitude variability of atmospheric and oceanic circulation patterns via the respective  
57 teleconnections (Aoki, 2017; Heil, 2006; Mahoney et al., 2007b) make fast ice a sensitive indicator of climate variability and  
58 even climate change (Mahoney et al., 2007a; Murphy et al., 1995). Based on the complexity and significance of fast ice in the  
59 Antarctic climate system, there is an urgent need for prognostic Antarctic fast ice in regional models, and later in global climate  
60 models, to capture its potential major impacts on the global ocean circulation, as developed recently for the Arctic (Lemieux  
61 et al., 2016).

62 Although fast ice only represents a rather small fraction of the overall sea-ice area in Antarctica (Fraser et al., 2012), it may  
63 contribute significantly to the overall volume of Antarctic sea ice, especially in spring (Giles et al., 2008). The presence and  
64 evolution of Antarctic fast ice is often associated with the formation and persistence of coastal polynyas, regions of particularly  
65 high sea-ice production (Fraser et al., 2019; Massom et al., 2001a; Tamura et al., 2016; Tamura et al., 2012) and Antarctic  
66 Bottom Water formation (Tamura et al., 2012; Williams et al., 2008). Also, it forms an important boundary between the  
67 Antarctic ice sheet and the pack ice/ocean, for example prolonging the residence times of icebergs (Massom et al., 2003),  
68 mechanically stabilizing floating glacier tongues and ice shelves, and delaying their calving (Massom et al., 2010; Massom et  
69 al., 2018). Therefore, one particularly interesting aspect of Antarctic fast ice is its interaction with nearby ice shelves, floating  
70 seaward extensions of the continental ice sheet that are present along nearly half of Antarctica's coastline. Under specific  
71 oceanographic conditions, supercooled Ice Shelf Water favors the formation of floating ice crystals within the water column  
72 (Foldvik, 1977; Dieckmann et al., 1986), as opposed to the regular process of sea-ice formation by heat transport from the  
73 ocean towards the colder atmosphere. These crystals may be advected out of an ice-shelf cavity and rise to the surface  
74 (Hoppmann et al., 2015b; Mahoney et al., 2011; Hughes et al., 2014). They are eventually trapped under a nearby fast-ice  
75 cover and may accumulate in a layer reaching several meters in thickness (Gough et al., 2012; Price et al., 2014; Brett et al.,  
76 2020). This sub-ice platelet layer has profound consequences for the local sea-ice system, and forms an entirely unique habitat.  
77 Thermodynamic growth of the overlying solid fast ice into this layer (by heat conduction from the ocean into the atmosphere)  
78 leads to subsequent consolidation, and the resulting incorporated platelet ice may contribute significantly to the local fast-ice  
79 mass and energy budgets. This phenomenon has been documented at various locations around Antarctica (Langhorne et al.,  
80 2015 and references therein), and where present, is a defining feature of the local coastal system. Refer to (Hoppmann, in  
81 review) for a comprehensive review of platelet ice.

82 The effects of fast ice on the exchange processes between ocean and atmosphere are further amplified by the accumulation of  
83 snow, as it forms a thick layer over large portions of the Antarctic sea ice (Massom et al., 2001b). However, the snow cover  
84 has opposing effects on the energy and mass budgets of sea ice in the region. On the one hand, due to its low thermal  
85 conductivity, snow acts as a barrier to heat transfer from sea ice to the atmosphere and effectively reduces ice growth at the  
86 bottom (Eicken et al., 1995). On the other hand, snow contributes significantly to sea-ice thickening at the surface through two  
87 distinct seasonal processes: snow-ice and superimposed ice formation. In winter/spring, the heavy snow load leads to the  
88 depression of the sea-ice surface below water level, causing flooding of the snow/ice interface. The subsequent refreezing of  
89 the snow/water mixture forms a salty layer of so-called snow-ice (e.g. Eicken et al., 1994; Jeffries et al., 1998; 2001). In  
90 contrast, in [spring and](#) summer, [surface and](#) internal snowmelt leads to melt water [that can percolating to the snow/ice interface](#)  
91 [where it refreezes](#) and forms fresh superimposed ice, [as it percolates through snow and eventually to the snow-ice interface](#)  
92 (Haas, 2001; Haas et al., 2001; Kawamura et al., 2004). Both processes contribute significantly to sea-ice growth from the top,  
93 and thus to the overall sea-ice mass budget in the Southern Ocean.



94 Beyond its contribution to the general sea-ice mass and energy budget in the Southern Ocean, fast ice also plays an important  
95 role for the ice-associated ecosystem, as it provides a stable habitat for microorganisms (e.g. Günther and Dieckmann, 1999)  
96 and serves as a breeding ground for, e.g., Weddell seals and Emperor penguins (Massom et al., 2009).  
97 Fast ice and its properties as described above have been studied around Antarctica for a long time, especially related to  
98 logistical work at the summer and overwintering bases close to the coast of the continent. In order to commonly coordinate  
99 and facilitate this research, and thus establish an international network of fast-ice monitoring stations around the Antarctic  
00 coastline, the international Antarctic Fast Ice Network (AFIN) was initiated during the International Polar Year (IPY)  
01 2007/2008 (Heil et al., 2011). Active international partners are, e.g., Australia and China working at Davis Station and  
02 Zhongshan Station on the eastern rim of Prydz Bay in East Antarctica (Heil, 2006; Lei et al., 2010), New Zealand working out  
03 of Scott Base in McMurdo Sound in the Ross Sea (Langhorne et al., 2015, and references therein), Norway at the fast ice in  
04 front of Fimbul Ice Shelf at Troll Station (Heil et al., 2011) and Germany in Atka Bay at Neumayer III (Hoppmann et al.,  
05 2013), both in the vicinity of Dronning Maud Land. The regular, AFIN-related monitoring program at Neumayer III started in  
06 2010 in order to fill the observational gap in the Weddell Sea sector.  
07 Here we present a decade of annual in-situ fast-ice observations in Atka Bay, which is one of the longest and most continuous  
08 time series within AFIN so far. The main dataset is a record of fast-ice thickness, snow depth, freeboard, and sub-ice platelet-  
09 layer thickness that was collected by a number of overwintering teams between 2010 and 2018. In addition to determining the  
10 spatio-temporal variability of the fast-ice cover, we co-analyze this data with meteorological observations and satellite imagery  
11 in order to determine how snow and platelet ice influence the local fast-ice mass budget. In doing so, we aim to improve our  
12 understanding of the interaction between the atmosphere, fast ice, ocean and ice shelves in one of the key regions in Antarctica.

## 13 **2 Study site and measurements**


### 14 **2.1 Study site: Atka Bay**

15 The main study area of this paper is Atka Bay, an 18 km-by-25 km embayment in front of the Ekström Ice Shelf located on  
16 the coast of Dronning Maud Land in the eastern Weddell Sea, Antarctica, at 70°35'S/ 7°35'W (Figure 1). Atka Bay is flanked  
17 towards the east, south and west by the edges of the ice shelf which rise as high as 20 meters above sea level. The cavity  
18 geometry of the Ekström Ice Shelf is one of the best known in Antarctica (Smith et al., 2020). Atka Bay is seasonally sea-ice  
19 covered, and the water depth ranges between ~~100 and 500 m, with a steep canyon of 275 m depth~~80 m and 275 m with  
20 maximum depth in the central bay (Kipfstuhl, 1991). Since the 1980s, when the first German research station Georg-von-  
21 Neumayer Station was established in the region, a variety of measurements has been carried out in the bay. Today's German  
22 research station Neumayer III is located at a distance of about 8 km from the bay, where drifting snow regularly forms natural  
23 ramps between the sea ice and the ice-shelf surface. Prior investigations of the interactions between ice shelf, sea ice and ocean  
24 in the bay and its surroundings have been carried out by Kipfstuhl (1991) and Nicolaus and Grosfeld (2004), as well as more

25 recently by Hoppmann et al. (2015a) and Hoppmann et al. (2015b). Ecosystem studies from the 1990's have been published  
26 by Günther and Dieckmann (1999) and Günther and Dieckmann (2001).  
27







30

31 **Figure 1.** Overview of the study site and its surroundings. (a) Atka Bay (red marker) is located at the edge of the northeastern  
 32 Weddell Sea. Coastline data taken from SCAR Antarctic Digital Database. (b) Close-up of map (a) to focus on the study site  
 33 of Atka Bay. The sampling sites of the standard transect (ATKA) are marked with red circles. (c) Enlargement of (b) showing  
 34 in addition to the standard transect (red circles) the parallel transect in the south (ATKA p; light red circles) from ATKA03p

35 to ATKA21p as well the eastern and western perpendicular transects ATKA S-N (east) from ATKA00SNe to ATKA18SNe  
36 and ATKA S-N (west) from ATKA00SNw to ATKA16SNw, each with a distance of 2 kilometers between adjacent sampling  
37 sites (light blue circles). The southern, eastern and western transects were sampled during a field campaign between November  
38 and December 2018. [The dotted black curves indicate the locations of ice rises and rumples in Atka Bay.](#) Background:  
39 Copernicus Sentinel data 01 December 2018, processed by ESA.


## 40 **2.2 Sea-ice conditions**

41 Atka Bay is seasonally covered with sea ice that is attached to the ice shelf to form immobile fast ice. Following the method  
42 of fast-ice time series retrieval detailed in Fraser et al. (2019), we obtained year-round estimates of fast-ice extent in Atka Bay  
43 from MODIS visible and thermal infrared satellite imagery. Hence, the fast-ice extent time series presented here [in Figure 2](#) is  
44 a) produced at a 1 km spatial and 15 day temporal resolution, from 15-day MODIS cloud-free composite images (following  
45 Fraser et al., 2010) and edge-detected non-cloud-filtered composite images; b) spans the time period from March 2000 to  
46 March 2018; and c) is semi-automated in the sense that the fast-ice edge is automatically delineated during times of high  
47 contrast to offshore pack ice/open water, and manually delineated at other times.

48 Accordingly, the initial ice formation in the bay has started in March in recent years (Figure 2), with persistent easterly winds  
49 forcing increased dynamic sea-ice growth towards the western ice-shelf edge of the bay. Once the bay is completely covered  
50 by fast ice usually at the end of April (Figure 2), further in-situ ice growth takes place. In the following summer, the ice does  
51 not [disappear by surface melting in-situ](#), but breaks up and drifts out of the bay once the conditions are sufficiently unstable.  
52 Stabilization and breakup of the ice-covered bay depend on the presence/absence of pack ice offshore of Atka Bay associated  
53 with changing ocean currents and winds, as well as stationary and passing icebergs. Thus, fast-ice breakup in the bay starts  
54 usually in December/January after the pack ice in front of the fast ice has retreated (Figure 2).

55 During our study period from 2010/2011 to 2018/19, there were two exceptions to this “typical” annual cycle: In September  
56 2012, a large iceberg (generally referred to as “B15G”) grounded in front of Atka Bay, sheltering the fast ice and consequently  
57 preventing sea-ice breakup in the following summer (Hoppmann et al., 2015b), resulting in second-year fast ice in the bay in  
58 2013. A year later, in August 2013, the iceberg dislodged itself, drifting westwards following the Antarctic Coastal Current.  
59 Fragments of the iceberg remained grounded in the northern part of the bay, causing it to be blocked again two years later, and  
60 therefore preventing sea-ice breakup in austral summer 2014/2015 for a second time within the study period. The iceberg  
61 fragments became mobile during the course of the following year, resulting in the bay to become ice-free again in the following  
62 summer.

63



64

65

**Figure 2:** Time series of fast-ice extent in Atka Bay between 8° 12'W and 7° 24'W derived from MODIS data between early 2000 to early 2018 (black line). The blue line shows the annual mean extent repeated [each year](#) over the same time period. The average fast-ice extent over the entire time series is  $319.2 \pm 167.8 \text{ km}^2$ , with an uncertainty of  $86.6 \text{ km}^2$ .

66

67

68

69

### 2.3 Sea-ice measurements across Atka Bay

70

71

72

73

74

75

76

77

78

79

80

81

82

83

84

85

Since 2010, the AFIN monitoring protocol has been implemented to study the seasonal evolution of fast ice along a 24-km long west-east transect in Atka Bay (“standard transect”, red circles in Figure 1). Here, six sampling sites have been regularly revisited between annual sea-ice formation and breakup each year to obtain a continuous record of snow depth, freeboard, sea ice- and sub-ice platelet-layer thickness across the bay (Arndt et al., 2019). Sampling sites on the standard transect are referred to in this paper as ATKAx<sub>x</sub>, where xx represents the distance in kilometers to the ice-shelf edge in the west.

Generally, measurements along that standard transect are carried out once a month by the wintering team usually between June and January, when safe access to the sea ice is possible. At each sampling site, up to five measurements are taken in an undisturbed area, one as the center measurement and four more at a distance of approx. 5 meters in each direction ([north, east, south, and west](#)), in order to account for the spatial variability of sea-ice and snow properties. In years of prevailing second-year ice in the bay (2012/2013, 2014/2015), the number of observations per sampling site was reduced to one (the center measurement) due to exceptionally thick snow and ice. Throughout this manuscript, we mainly present the mean values from those up to five single measurements per sampling site. While all measurements along the standard transect from 2010/2011 to 2018/2019 are included in this study, the sea-ice monitoring activities will be continued beyond this work.

In November and December 2018, additional measurements in both, parallel and perpendicular transect lines to the standard transect, have been performed (Figure 1c). Sampling sites on parallel transects are referred to in this paper as ATKAx<sub>p</sub>, where xx represents the distance in kilometers to the ice shelf edge in the west [and “p” refers to “parallel”](#). Along the perpendicular

86 western (w) and eastern (e) transects from south to north, sampling sites are referred to in this paper as ATK AyySNw and  
87 ATK AyySNe, where yy represents the distance in kilometers to the ice-shelf edge in the south.

88 Sea-ice and platelet-layer thickness as well as freeboard are measured with a (modified) thickness tape. In order to enable the  
89 penetration of the usually semi-consolidated platelet layer, the regular metal plates at the bottom of the thickness tape were  
90 replaced by a metal bar of ~2kg. The underside of the platelet layer is determined by gently pulling up the tape and attempting  
91 to feel the first resistance to the pulling. Sea-ice thickness was measured either by pulling this modified tape through the entire  
92 platelet layer until the solid sea-ice bottom is reached (with a high risk of it getting stuck), or using a regular ice thickness tape.  
93 The modified tape is retrieved by pulling a small rope attached to one side of the metal bar. Snow depth was measured using  
94 ruler sticks. Freeboard is defined as the distance between the snow/ice interface and the sea-water level, while the snow/ice  
95 interface above (below) sea-water level is referred to as positive (negative) freeboard.

96 In order to determine the influence of snow and the underlying platelet layer to the observed freeboard (F), we also calculated  
97 ~~this the following~~ parameter, assuming a hydrostatic equilibrium for floating snow-covered sea ice with an additional buoyancy  
98 (the platelet layer below), using Archimedes' principle:

$$99 F = - \frac{I \cdot (\rho_I - \rho_W) + S \cdot \rho_S + P \cdot (\rho_P - \rho_W)}{\rho_W} \quad \text{(Eq. 1.1)}$$

100 ~~As soon as F gets negative, a flooding of the snow/ice interface is assumed, and with that the formation of snow ice. As the~~  
101 ~~latter is assumed to have the same density as sea ice (Knight, 1988), freeboard is calculated for the flooded case as~~

$$102 -F = S - \frac{I + S - P \left( \frac{\rho_P - \rho_W}{\rho_W - \rho_I} \right)}{1 + \frac{\rho_S}{\rho_W - \rho_I}} \quad \text{(Eq. 1.2)}$$

103  
104 ~~where~~ In Equation 1.1 and 1.2, I refers to sea-ice thickness, S to snow depth, P to platelet-layer thickness, the indices I refers  
105 to sea ice, S to snow, P to the platelet layer, and W to water. Constant typical densities of  $\rho_W = 1032.3 \text{ kg m}^{-3}$ ,  $\rho_S = 330 \text{ kg m}^{-3}$   
106 and  $\rho_I = 925 \text{ kg m}^{-3}$  are assumed in this study. The platelet-layer density  $\rho_P$  is calculated ~~by means of the as the product of~~  
107 ~~sea ice density and~~ platelet-layer ice volume fraction  $\beta$  ~~as~~  $\rho_P = \beta \cdot \rho_I + (1 - \beta) \cdot \rho_W$ . In this study, we used a constant ice-  
108 volume fraction of  $\beta = 0.25$ , as suggested by Hoppmann et al. (2015b).


109 The described bore-hole measurements are occasionally complemented by additional total (sea-ice plus snow) thickness  
110 measurements with a ground-based electromagnetic induction instrument (e.g. Hunkeler et al., 2016) as well as autonomous  
111 ice tethered systems, such as Ice Mass balance or Snow Buoys (Grosfeld et al., 2015; Hoppmann et al., 2015a). However, this  
112 paper focusses on the regular bore-hole measurements only, as the additional observations address scientific questions beyond  
113 the scope of this paper.

114

## 15 **2.4 Meteorological conditions and observations at Neumayer III**

16 At the meteorological observatory of the nearby wintering base Neumayer III, atmospheric conditions have been recorded  
17 since 1981 (König-Langlo and Loose, 2007), including the study period from 2010/2011 to 2018/19, and continuing beyond  
18 it (Schmithüsen et al., 2019). Occasionally, automatic weather stations (AWS) were temporarily installed on the sea ice to  
19 record the meteorological conditions directly on the sea ice (Hoppmann et al., 2015a). Since the 2m air temperature and the  
20 wind velocity at the meteorological observatory and the AWS on the ice showed a fairly good agreement in prior studies  
21 (Hoppmann et al., 2015a; Hoppmann et al., 2013), we ~~utilize~~use in this paper the more continuous records of the  
22 meteorological observatory in order to investigate the links between sea-ice conditions and atmospheric conditions. The  
23 Neumayer III data is recorded as minutely averages of typically 10 values per averaging interval. The instrumentation is  
24 checked on a daily basis, any erroneous values, e.g. caused by riming or instrument failure, are removed from the record.  
25 Therefore, the data quality can be considered high, even though there might be gaps in the records due to the validation  
26 routines. Nevertheless, data availability is 99.4% for wind direction, 99.0% for wind speed and 99.7% for air temperature.  
27 Uncertainties are essentially those classified by the manufacturers. Instrument details are given in the metadata of the datasets  
28 since February 2017 in Schmithüsen et al. (2019), earlier data is documented in König-Langlo and Loose (2007).  
29 Generally, in the vicinity of Neumayer III the weather is strongly influenced by cyclonic activities which are dominated by  
30 easterly moving cyclones north of the station. This leads to prevailing persistent and strong easterly winds which exhibit a  
31 seasonal cycle with strongest winds during winter time (Figure 3). The second strongest mode in the wind direction distribution  
32 at 270° (westward) is associated with super geostrophic flows resulting from a high-pressure ridge north of Neumayer III  
33 (König-Langlo and Loose, 2007). These strong winds lead to frequent drifting and blowing snow. Here, we expect snow  
34 transport for 10-m wind velocities exceeding 7.7 m/s for dry snow and exceeding 9.9 m/s for wet snow (Li and Pomeroy,  
35 1997).

36



37

38 **Figure 3:** Distribution of wind speed related to wind directions separated for austral fall (March, April, May; MAM), winter  
 39 (June, July, August; JJA), spring (September, October, November; SON), and summer (December, January, February; DJF)  
 40 for the study period from 2010 to 2018. Colors indicate the relative frequency of each shown [value-pair/wind direction to wind](#)  
 41 [speed pair](#). Dashed vertical lines denote thresholds for 10-m wind speeds for snow transport of dry (7.7 m/s) and wet snow  
 42 (9.9 m/s) (Li and Pomeroy, 1997).

43

### 3 Results

44

#### 3.1 Nine-year record of sea-ice and plateau-layer thickness, snow depth and freeboard along a 24-km W-E transect


45

Figure 4 summarizes all conducted measurements of snow depth, sea-ice and plateau-layer thickness on the standard transect  
 46 from bore-hole measurements for each ATKA sampling site in the study period from 2010 to 2018. In the seven months when  
 47 sea-ice conditions allowed safe access (usually from May/June to December), about eight sets of measurements were taken  
 48 along the standard transect crossing Atka Bay, i.e. once every three to four weeks.

49

Analyzing the average annual maximum values of the investigated parameters (Table 1) for years of seasonal fast ice only  
 50 (excluding 2013 and 2015) and neglecting local iceberg disturbances (ATKA07 in 2017), the highest annual snow  
 51 accumulation of  $0.89 \pm 0.36$  m was measured at ATKA07, while the smallest by far was measured at ATKA24 at the  
 52 easternmost sampling site, with only  $0.28 \pm 0.19$  m. Averaged over the entire bay, the lowest snow accumulation of  $0.51 \pm$   
 53  $0.30$  m was observed in 2016. In contrast, 2011 was the year with the most snow and an average snow depth of  $0.85 \pm 0.20$  m

54 across the bay. The average seasonal fast-ice thickness based on the measurements during the observation period varied  
 55 between  $1.74 \pm 0.31$  m (ATKA21) and  $2.58 \pm 1.28$  m (ATKA07) with a mean value of  $1.99 \pm 0.63$  m. The underlying seasonal  
 56 platelet layer reached an average annual thickness of 3.91 m, which, however, shows a strong gradient in the average annual  
 57 maximum values (Table 1) from  $4.62 \pm 0.67$  m at ATKA07 in the west of the bay to  $2.82 \pm 1.20$  m at ATKA24 in the east.  
 58 In 2013 and 2015, the fast ice in Atka -Bay became second-year ice due to grounded icebergs in front of the bay. Within the  
 59 respective second year, snow depth increased further by an additional  $0.88 \pm 0.43$  in 2013 and by  $0.74 \pm 0.27$  m in 2015. In  
 60 2013, the average fast-ice thickness across the bay increased by an additional  $1.21 \pm 0.42$  m, while in 2015, it increased by an  
 61 additional  $2.79$  m  $\pm 1.48$  m. In the years of prevalent second-year ice in the bay, the thickness of the platelet layer increased  
 62 on average by  $5.13$  m  $\pm 1.43$  m in 2013 (compared to the end of 2012), and  $4.11$  m  $\pm 1.86$  m in 2015 (compared to the end of  
 63 2014). During these periods, ATKA11 experienced the highest annual platelet-layer thickness increase of  $6.82$  m and  $6.44$  m,  
 64 respectively.  
 65



66


67 **Figure 4:** Time series of snow depth, fast-ice and platelet-layer thickness from bore-hole measurements along the standard  
68 transect for each ATKA sampling site (Figure 1) for the time period from 2010 to 2018. Note: In 2010 and 2011, the platelet-  
69 layer thickness was not measured (\*). In 2012/2013 and 2014/2015 Atka Bay was blocked by icebergs, so the fast ice did not  
70 break up and turned into second-year ice instead (–). In 2017, a small iceberg in the vicinity of ATKA07 strongly influenced  
71 the snow measurements ( $\Delta$ ). Reference depth of 0 meters is the snow/ice interface.

72


73 Figure 5 depicts the evolution of the water level with respect to the snow/ice interface (which is the freeboard with an opposite  
74 sign) along the standard transect in the study period from 2010 to 2018. Taking all conducted freeboard measurements from  
75 seasonal fast ice into account, 38% reveal negative data, i.e. flooding can be assumed, with an average negative freeboard of -  
76  $0.10 \pm 0.08$  m. In contrast, considering freeboard measurements from second-year ice only, 55% of the data indicate a negative  
77 freeboard, with an average of  $-0.22 \pm 0.15$  m. Analyzing the average annual maximum of the negative freeboard values (Table  
78 1) for years of seasonal fast ice only, and neglecting local iceberg disturbances (ATKA07 in 2017), there is no distinct gradient  
79 across Atka Bay, but higher average negative freeboard values ( $-0.07$  to  $-0.08$  m) are recorded both in the far west (ATKA03)  
80 and in the east (ATKA16 and ATKA21), whereas the lowest average negative freeboard of  $-0.01 \pm 0.08$  m was measured at  
81 ATKA07. According to Equation 1.1, ~~8966%~~ of the calculated freeboard values are smaller than the measured values. The  
82 difference between measured and calculated freeboard values ranges from  ~~$-0.4954$  to  $0.5411$~~  m with an average of  $0.008 \pm$   
83  $0.190$  m ~~(towards higher measured value)~~. Neglecting the underlying buoyant platelet layer in the calculation reduces the  
84 freeboard by  $0.079 \pm 0.1506$  m, whereas neglecting the snow layer on top of the sea ice increases the freeboard by  $0.1920 \pm$   
85  $0.1729$  m (Figure 5).

86





87



88

89 **Figure 5:** Close-up of Figure 4 which highlights the location of the water level with respect to the snow/ice interface (which  
 90 has the opposite sign of the freeboard) as measured in the field (black solid line) and as calculated according to Equation 1.1  
 91 and 1.2 including snow and platelet-layer thickness (blue dashed line), neglecting the snow cover (dashed yellow line) and  
 92 platelet layer (dashed green line), respectively. Please note that, for the purpose of better illustration, we depict here the actual  
 93 location of the water level rather than the freeboard (the only difference being the opposite sign). This means that, if the water  
 94 level is above the snow/ice interface, this is depicted in the figure accordingly, while the actual freeboard carries a negative  
 95 sign, and vice versa. The reference depth of 0 represents the snow/ice interface.

96

97 **Table 1:** Average annual maximum of snow depth, sea-ice and platelet-layer thickness, as well as freeboard (negative equals  
 98 potential flooding) on the standard transect from bore-hole measurements for each ATKA sampling site (Figure 1) for the time  
 99 period from 2010 to 2018, excluding years of second-year ice due to blocking of the bay (i.e. 2013 and 2015). <sup>1</sup>At ATKA11  
 00 all measurements of the year 2012 are also neglected as the ice has temporarily broken up again. <sup>2</sup>At ATKA07 the snow  
 01 measurements of the year 2017 are also neglected as a small iceberg has strongly influenced the accumulation rates. Standard  
 02 deviations are given in [bracketsparentheses](#).

	ATKA03	ATKA07	ATKA11	ATKA16	ATKA21	ATKA24
Snow depth in m	0.81 (0.35)	0.89 (0.36) <sup>2</sup>	0.74 (0.23) <sup>1</sup>	0.79 (0.37)	0.77 (0.24)	0.28 (0.19)
Ice thickness in m	2.04 (0.31)	2.58 (1.28)	1.97 (0.25) <sup>1</sup>	1.81 (0.36)	1.74 (0.31)	1.83 (0.35)
Platelet-layer thickness in m	3.88 (1.31)	4.62 (0.47)	4.59 (0.83) <sup>1</sup>	3.99 (0.94)	4.21 (0.54)	2.82 (1.20)
Freeboard in m	-0.08 (0.14)	-0.01(0.08) <sup>2</sup>	-0.05(0.08) <sup>1</sup>	-0.07 (0.09)	-0.08 (0.10)	-0.05 (0.09)


03  
04

### 05 3.2 Seasonal snow, sea-ice and platelet-layer accumulation/growth and melt rates

06 Figure 6 summarizes the seasonal snow depth, sea-ice and platelet-layer thickness evolution separated for austral fall (March,  
07 April, May; MAM), winter (June, July, August; JJA), spring (September, October, November; SON), and summer (December,  
08 January, February; DJF) averaged for each ATKA sampling point over the duration of the whole ~~for the~~ study period from  
09 2010 to 2018.

10 Considering the average monthly snow accumulation rates, a slight increase from fall (from 0.04 to 0.09 m per month, across  
11 stations) to spring (0.09 to 0.11 m per month) becomes apparent, if excluding the eastern sampling sites at ATKA21 and  
12 ATKA24. Latter sampling sites show the highest monthly averaged accumulation rates during austral fall (0.11 and 0.08 m  
13 per month), which subsequently decrease to 0.10 and 0.02 m per month, respectively. In contrast, a clear snow loss with a  
14 maximum monthly average of up to  $0.03 \pm 0.12$  m at ATKA11 and a maximum snow loss rate of 0.21 m per month at ATKA07  
15 (80<sup>th</sup> percentile), can be seen mostly during summer months. ~~Also, t~~The seasonal evolution of the platelet layer shows a similar  
16 pattern: between austral autumn and spring, an average monthly thickness increase of up to  $0.82 \pm 0.30$  m at ATKA11 is  
17 observed. Excluding ATKA07, afterwards an average monthly platelet-layer thickness decrease of 0.85 m is calculated for  
18 summer. The maximum decrease of 6.25 m per month occurred at ATKA11 in 2013 (80<sup>th</sup> percentile). However, it is highly  
19 likely that this is a measurement error. In contrast, ATKA07 also reveals an increase in platelet-layer thickness during the  
20 summer months with a monthly average of  $0.67 \pm 1.20$  m. With regard to the growth rates of fast ice in Atka Bay, a contrasting  
21 but expected seasonal development is observed: The highest average monthly fast-ice growth rates of up to approx. 1 m per  
22 month (80<sup>th</sup> percentile) are measured in autumn, and decrease in the following month until spring. These exceptionally high  
23 growth rates result from rapid growth of the solid fast ice into the (unconsolidated) sub-ice platelet layer, i.e. from the  
24 subsequent freezing of the interstitial water between the platelets in the top part of the platelet layer. In other words, some of  
25 the heat within the newly growing ice was already extracted earlier by the ocean during the process of platelet crystal formation  
26 in the supercooled Ice Shelf Water plume. In the subsequent summer months, average monthly sea-ice growth rates increased

27 again to values between 0.07 m (ATKA07) and 0.24 m (ATKA21), except for ATKA24, where sea-ice melt dominates with  
 28 an average monthly melt rate of  $-0.05 \pm 0.22$  m and a maximum monthly sea-ice melt rate of  $-0.58$  m.  
 29




30

31 **Figure 6:** Seasonal snow, sea-ice and platelet-layer accumulation/growth and melt rates separated for austral fall (March,  
32 April, May; MAM), winter (June, July, August; JJA), spring (September, October, November; SON), and summer (December,  
33 January, February; DJF) for each ATKA sampling point for the study period from 2010 to 2018. Boxes are the first and third  
34 quartiles; whiskers the 20<sup>th</sup> and 80<sup>th</sup> percentile. Circles indicate the mean, ~~vertical~~horizontal lines in the boxes the median.  
35 Numbers below the whiskers indicate the respective sampling size, i.e. the number of included years, with a maximum of nine.  
36

### 37 **3.3 Spatial variability of snow depth, sea-ice and platelet-layer thickness**

38 In order to describe the spatial variability of snow depth, sea-ice and platelet-layer thickness in west-to-east as well as in south-  
39 to-north direction across Atka Bay, additional parallel and perpendicular transects to the standard transect have been sampled  
40 in November/December 2018 (Figure 7). Considering the solid sea ice only, the complementary transect data show that sea-  
41 ice thickness over the bay in south-north and west-east direction is rather constant with an average of  $1.68 \pm 0.21$  m. In contrast,  
42 neglecting the measurements in iceberg-affected areas, snow depth data show higher values in the south and in the center of  
43 the bay of up to  $1.00 \pm 0.04$  m, while decreasing significantly towards the eastern ice-shelf edge and northern fast-ice edge to  
44  $0.08 \pm 0.01$  m and  $0.28 \pm 0.09$  m, respectively. The platelet-layer thickness beneath the fast ice shows a large spatial variability.  
45 While all measurements on the standard transect reveal the lowest platelet-layer thickness in the east of the bay at ATKA24  
46 (see Section 3.1), on the parallel transect in the south a maximum platelet-layer thickness of  $7.18 \pm 0.26$  m at the easternmost  
47 sampling point (ATKA21p) is observed. For the perpendicular transects in south-to-north direction, a significantly decreasing  
48 gradient in platelet-layer thickness from the ice-shelf edge towards the northern fast-ice edge is evident. On the western south-  
49 to-north transect, a decrease from  $6.62 \pm 0.25$  m to  $2.33 \pm 0.08$  m was observed, whereas for the eastern transect this strong  
50 gradient is even more apparent with a decrease from  $9.17 \pm 0.11$  m to  $1.88 \pm 0.20$  m.

51



52

53 **Figure 7:** Overview of measurements on the standard transect from west to east (upper left), the  
 54 western perpendicular transect from south to north (upper right) and the respective parallel one to the east (lower right) showing  
 55 the water level, snow depth, fast-ice and platelet-layer thickness across Atka Bay. All measurements were conducted between  
 56 November 25, 2018 and December 07, 2018. For the parallel west-east transect (December 01, 2018), the platelet-layer  
 57 thickness evolution is influenced by a nearby iceberg (see Figure 1c). Also, for the western north-south transect (December  
 58 03, 2018), snow measurements are influenced by several small icebergs in the vicinity between kilometers 4 and 8 (see Figure  
 59 1 c).

## 60 4 Discussion

### 61 4.1 Seasonal and interannual variability of snow depth, sea-ice and platelet-layer thickness

62 The fast-ice regime in Atka Bay is primarily seasonal and the sea-ice cover usually only remains in the bay if a breakup is  
 63 prevented by grounded icebergs in front of it. For example, while in 2013 a 17km-by-10km iceberg (B15G) blocked the entire  
 64 bay, in 2015, only small iceberg fragments of B15G in front of the bay were sufficient to ensure that the sea ice in the bay did  
 65 not break up, but became perennial. It may also occasionally happen that small areas of fast ice remain attached to the ice-  
 66 shelf edge, or that individual ice floes remain in the bay and are incorporated into the newly growing ice in the following  
 67 winter. Not only does the presence and size of the icebergs play a role in the fast-ice seasonality, but also the location and  
 68 associated influence of atmospheric circulation patterns and ocean processes.

69 Considering first of all the seasonal sea ice only, the presented measurements along the standard transect across Atka Bay  
 70 indicate a clear seasonal cycle in all investigated variables, i.e. snow depth, sea-ice and platelet-layer thickness: The initial


71 sea-ice formation in Atka Bay starts in March and proceeds towards a completely fast-ice-covered bay at the end of April. The  
72 continuous sea-ice growth (i.e. ocean-atmosphere heat flux) proceeds with decreasing growth rate through fall and winter until  
73 the thickening snow cover more and more reduces the heat flux between the upper ocean and the atmosphere, preventing  
74 further thermodynamic sea-ice growth. However, the fast-ice thickness still increases in spring and even during austral summer  
75 months (albeit very slowly). This can be explained by the measurement uncertainty with respect to the large spatial variability  
76 of sea-ice thickness even on very small (centimeter) scales, but the consistency in the data suggests that it could also be caused  
77 by consolidation processes within the platelet layer below, i.e. in-situ sea-ice growth by heat transport into a supercooled  
78 plume residing right beneath the solid fast ice similar to observations in McMurdo Sound (Smith et al., 2012; Leonard et al.,  
79 2011; Dempsey et al., 2010; Robinson et al., 2014). So far, in Atka Bay there is only evidence that platelets grow quite large  
80 already while still suspended in the water column (Hoppmann et al., 2015b). To what degree an in-situ growth of platelet  
81 crystals and consolidation processes that go beyond regular freeze-in of the topmost part of the platelet layer by heat conduction  
82 to the atmosphere play a role at Atka Bay still needs to be investigated. In any case the platelet layer is an efficient buffer  
83 between the fast ice and the incoming warmer water in summer (Eicken and Lange, 1989), so the lack of noticeable fast-ice  
84 bottom melt is generally expected. Oceanographic (winter) data is sparse, and the monitoring at Atka Bay has recently been  
85 extended to also include regular CTD casts, whenever the (challenging) conditions and time constraints allow. An analysis of  
86 available CTD data in Atka Bay is currently ongoing, and will be shown in a future dedicated study to close the above  
87 observational gaps with respect to the ocean.

88 Destabilization of the fast ice and the platelet layer below in summer is to a large extent driven by the presence/absence of  
89 pack ice offshore Atka Bay. Thus, the initial breakup and subsequent retreat of the pack ice in front of the bay allow for locally  
90 increasing ocean currents beneath the fast ice and the inflow of warm Antarctic Surface Water from the east (Hoppmann et al.,  
91 2015a; Hattermann et al., 2012) causing both washing out of the platelet layer as well as mixing warm water into the water  
92 column associated with a thinning rate of the platelet layer of approx. one meter per month from December onwards. The  
93 retreating fast ice also initializes the sea-ice breakup in the bay starting usually in December/January (Figure 2). The  
94 diminishing fast-ice zone potentially goes along with an additional thinning of the platelet layer (Figure 8 b) by, e.g., further  
95 washing out mechanisms. Even though the correlation between the change of fast-ice extent and the mean (maximum) platelet-  
96 layer thickness between two consecutive surveys with an an r-coefficient of 0.38 (0.35) is relatively low, Figure 8b shows-indeed  
97 suggests that decreasing platelet-layer thicknesses are generally associated with retreating fast ice in Atka Bay. Also, Massom  
98 et al. (2018) have also shown that pack ice has a stabilizing effect as a buffer against ocean swells. The deployment of complex  
99 oceanographic moorings, either ice- or seafloor-based, would greatly help immensely-to further investigate ocean properties  
00 and currents and their effect on sea ice and the ice shelf, but their deployment, and especially their recovery, is extremely  
01 difficult and risky in the dynamic and harsh conditions of Atka Bay. While such deployments are logistically not feasible at  
02 the moment, it is planned to include suitable instrumentation in the monitoring within the next years.

03 In contrast to the decrease in platelet-layer thickness beneath the fast ice in summer over the entire bay, the snow cover on top  
04 does not show a clear seasonal pattern, but indicates a decrease in snow depth with increasing air temperature. However, even

05 in summer, no consistent snow melt with associated strong mass loss is observed over the entire bay. Rather, a strong variability  
06 in snow depth over all sampling sites and all sampling years with a weak snow loss during summer months (Figure 6) is  
07 observed (Figure 6). ~~This-The latter~~ pattern is a result of both, temporary temperatures above freezing which favor surface  
08 melting (Figure 8a), and comparatively low wind speeds (Figure 3) preventing the accumulation of additional snow blown  
09 over from the surrounding ice shelf. These results match well with results from studies on the seasonal cycle of snow properties  
10 in the inner pack ice zone of the Weddell Sea as for example performed by Arndt et al. (2016), who also showed missing  
11 persistent summer melt as highlighted above.

12 Overall, the 9-year time series for in-situ snow depth, sea-ice and platelet-layer thickness in Atka Bay do not show any trend  
13 over the analyzed study period, whereas their inter-annual variability is dominated by local or temporary effects such as the  
14 presence of icebergs, which may for example lead to small-scale strong snow accumulations (Figure 4) or occasionally even  
15 to a perennial fast-ice regime. It is particularly remarkable that the average annual maximum platelet-layer thickness ~~increase~~  
16 of 4 m (Table 1) is consistent with an earlier investigation at Atka Bay performed in 1982 by Kipfstuhl (1991) between the  
17 western ice-shelf edge and ATKA03, and, at the same time, much higher than in results from another study ~~from in~~ 1995,  
18 where a maximum platelet-layer thickness of 1.5 m was measured in a similar location (Günther and Dieckmann, 1999).  
19 Considering a) the fact that these two studies only sampled one ~~site~~location, and b) the generally large spatial and temporal  
20 variabilities of the platelet-layer ~~properties~~thickness, and the results from our much in the present more detailed study, we infer  
21 that there seems to be no clear trend over the past decades. Our results are also in line with a recent study of Brett et al. (2020),  
22 who also found spatially highly variable platelet layers of 4+m under fast ice in McMurdo Sound. Thereby, our results suggest  
23 that it is likely that the relevant (sub ice-shelf) processes in this region have not changed much either. However, as already  
24 stated above, it is crucial to further look into all the available oceanographic data that are~~is~~ available from the Atka Bay region  
25 in order to support (~~or~~ disprove) these indications provided by the fast-ice monitoring. Another benefit of such comparison  
26 would be to strengthen (~~or~~ weaken) the hypothesis that fast-ice properties can serve as an indicator for the status of an ice-  
27 shelf, as suggested by Langhorne et al. (2015).



28

29 **Figure 8:** Scatter plot comparing (a) the average 2-meter air temperature (see Section 2.4) and the snow accumulation between  
 30 two consecutive surveys, and (b) increasing (positive values) and decreasing (negative values) fast-ice extent and platelet-  
 31 layer thickness between two consecutive surveys. The analysis includes all measurements at all sampling sites throughout the  
 32 study period from 2010 to 2018. Blue circles and red crossed denote the respective mean and maximum values within the time  
 33 frame between the consecutive measurements. Colored solid lines in Figure (b) show the linear regression between both  
 34 parameters with the respective correlation coefficients  $R$ .

35

#### 36 4.2 Spatial variability of fast-ice properties related to the distance to the ice-shelf edges around the bay

37 When neglecting local disturbances, such as icebergs, our results clearly indicate differences in the evolution of platelet-layer  
 38 thickness and snow depth with respect to the distance to the adjacent ice-shelf edges around Atka Bay. In contrast, the fast ice  
 39 itself does not exhibit any large spatial variability, [but measures with](#) at the end of the season a nearly uniform thickness of 2  
 40 meters across the bay, both in west-east and south-north direction (Figure 7).

41 Analysis of the spatial distribution of platelet-layer thickness under the fast ice along the standard transect over the entire bay  
 42 reveals that ATKA24 shows a significantly thinner platelet layer than all other sampling sites. In contrast, the parallel transect  
 43 towards the south reveals a significantly higher platelet-layer thickness at the sampling point closest to the eastern shelf ice  
 44 edge (ATKA21p). Perpendicular sampling transects from close to the southern ice-shelf edge towards the fast-ice edge in the  
 45 north show a strong increase of platelet-layer thickness near the ice-shelf edge, followed by a moderate decrease in platelet-  
 46 layer thickness towards the north, which rapidly decreases about 5 kilometers off the fast-ice edge. This thickness gradient is



47 much more pronounced on the south-north transect in the central area of the bay compared to the western one. Moreover,  
48 considering the entire time series of the west-east transect (Figure 4), the highest platelet-layer thickness is observed in the  
49 central area of the bay (ATKA07 and ATKA11). Summarizing all these observations, we hypothesize that, on the one hand,  
50 the strongest outflow of supercooled water from the ice-shelf cavity (along with associated suspended platelet crystals)- leads  
51 from the south centrally into the bay. On the other hand, local under-water topographic features of the ice shelf (i.e. ice rises)  
52 at the eastern boundary of Atka Bay (Figure 1c; Hoppmann et al., 2015b) might lead to a blocking of ocean currents and thus  
53 the advection of suspended platelet crystals, causing the high platelet-layer thickness at ATKA21p and the consistently low  
54 observed thickness north of this location at ATKA24 (Figure 4). The strongly decreasing gradient in the platelet-layer thickness  
55 towards the northern sea-ice edge is likely due to increasing distance from the source of suspended platelet crystals being  
56 advected from under the ice shelf and related washout effects. This is especially likely since at the time of the corresponding  
57 measurement, the pack ice in front of the bay was already broken open, allowing for wind-induced currents and locally solar-  
58 heated water production (the so-called mode 3 incursions, Jacobs et al. (1992), their (Figure 1). Also, the fact that the  
59 northernmost sampling points are located close the edge of the bay or even already outside of it, raises the probability that the  
60 predominant coastal ocean current transports warm Antarctic Surface Water towards the fast-ice area, which would further  
61 intensify this effect (Hattermann et al., 2012; Hoppmann et al., 2015b). From this, it can be generalized that a smaller amount  
62 of platelet crystals can accumulate under narrow fast-ice areas, since these are exposed to stronger oceanic currents and  
63 associated washout effects, as well as warm water incursions. ~~Comparative analyses to other study regions are not possible at~~  
64 ~~this time, since, to our knowledge, no comparable transects were carried out so far in other Antarctic fast-ice regions with~~  
65 ~~platelet layers beneath.~~ ~~Regarding the properties of the ocean in this region with respect to its interaction with the ice shelf and~~  
66 ~~sea ice.~~ ~~In that respect,~~ Hoppmann et al.; (2015b) used a subset of oceanographic data collected by the nearby PALAOA  
67 hydrographic observatory (Boebel et al., 2006) to link fast ice observations to ocean properties. A more recent study by Smith  
68 et al. (2020) helped to constrain the boundary conditions for Ice Shelf Water outflow by mapping in great detail the cavity  
69 geometry of the Ekström Ice Shelf. This study also shows data from repeated CTD casts through a borehole in the ice shelf,  
70 revealing the buoyant outflow of Ice Shelf Water in a relatively shallow surface layer. While these efforts help to better  
71 understand the complex system of ice shelf-ocean-sea ice interaction in this region, we ~~conclude-suggest~~ that a more  
72 comprehensive, year-round oceanographic study ~~that~~ also implementings a dedicated survey program is still urgently needed  
73 as a complement to the ongoing sea ice monitoring. ~~in order~~ This would allow us to investigate in more detail the outflow of  
74 Ice Shelf Water and the complex processes involved in the redistribution of platelet crystals that emerge from the ice shelf  
75 cavity. ~~Comparative analyses to other study regions are not possible at this time, since, to our knowledge, no comparable~~  
76 ~~transects were carried out so far in other Antarctic fast-ice regions with platelet layers beneath.~~  
77 Examining the spatial distribution of snow over the bay, the considerably lower snow depth at ATKA24 compared to all other  
78 sampling sites is striking (Figure 4), and most likely related to the proximity to the ice-shelf edge in approximately 1 km  
79 distance. Due to the prevailing easterly winds in the bay (Figure 3), an east-west gradient in snow depth could have been  
80 expected over the rest of the bay. However, this gradient cannot be determined on average over the entire time series. This is

81 mainly due to temporary local disturbance factors in the bay, such as icebergs and pressure ridges, which locally dominate the  
82 snow distribution and thus lead to a comparatively homogeneous distribution of snow depth over the central part of Atka Bay.  
83 A south-north survey across the bay at the beginning of austral summer 2018, however, revealed a trend of decreasing snow  
84 depth towards the northern fast-ice edge, with a stronger gradient approx. 5 km from the ice edge (Figure 7), which is in line  
85 with the northern boundary of the ice-shelf edge (Figure 1) and can therefore be explained by associated decreasing [effects of](#)  
86 [the prevailing](#) offshore winds. ~~and e~~Consequently, [also the measured snow depth is less in that part of the bay.](#) ~~less snow~~  
87 [redistribution.](#)

88  
89 Due to the generally thick snow cover on Antarctic sea ice (Kern and Ozsoy-Çiçek, 2016; Markus and Cavalieri, 1998; Massom  
90 et al., 2001b), flooding of the snow/ice interface and the resulting formation of snow-ice is a widespread phenomenon in the  
91 Southern Ocean and contributes significantly to the sea-ice mass budget in the area (Eicken et al., 1995; Jeffries et al., 2001).  
92 While Günther and Dieckmann (1999) observed no flooding in Atka Bay during their study, Kipfstuhl (1991) reported flooding  
93 in relation to snow loads greater than 1 meter, an observation that we can largely confirm with our data. Exceptions are  
94 measurements on comparatively thin ice that already showed a sufficient snow layer, e.g. in the austral winter 2010 and 2011  
95 at ATKA21, leading to a negative freeboard and potential flooding already early in the season (Figure 5). Consequently, taking  
96 all conducted freeboard measurements on the seasonal fast ice into account, 55 % of the data indicate a negative freeboard, i.e.  
97 potential flooding and associated snow-ice formation can be assumed. While the snow cover reduces the buoyancy of the sea  
98 ice and accelerates flooding, the underlying platelet layer counteracts this by adding additional buoyancy. However, neglecting  
99 the platelet layer reduces the freeboard by  $0.09 \pm 0.06$  m, but still a negative freeboard is derived in half of the calculations.  
00 Thus, the spatial distribution of the sign of the freeboard, and therefore also the flooding of the snow/ice interface, is essentially  
01 controlled by the snow layer on top of the fast ice in Atka Bay. The thickness of the underlying platelet layer below, in turn,  
02 contributes to the resulting thickness of the flooded layer and consequently to the thickness of the [expected-potential](#) snow-ice  
03 layer.

#### 04 **4.3 Impact of local disturbances on bay-wide properties and processes**

05 As already stated above, the largest overall effect on the fast-ice properties, the underlying platelet layer and the snow on top  
06 is due to the presence of icebergs in front of Atka Bay, which might entirely prevent a fast-ice breakup. At the same time,  
07 those grounded icebergs that are enclosed by sea ice within the bay add strong local effects. Thus, the large iceberg B15G  
08 grounded in front of Atka Bay sheltering the fast ice in the bay and consequently preventing sea-ice breakup in the following  
09 summer (Hoppmann et al., 2015b) led to second-year fast ice in the bay in 2013. Our measurements have shown that this  
10 hardly had any effect on the monthly snow accumulation rate and platelet-layer growth rate, but rather that these were within  
11 the same range as in the years of seasonal sea-ice cover. Accordingly, for [perennial-second-year](#) sea ice, the total annual snow  
12 and platelet-layer thicknesses are approximately twice as thick as in the other years and average to  $1.30 \pm 0.60$  m and  $7.84 \pm$   
13  $1.33$  m, respectively. Higher snow loads do also increase the probability and extent of surface flooding. This is not only

14 observed for years of ~~perennial-second-year~~ sea ice, but also for local disturbances as a result of the presence of small icebergs  
15 inside of the bay. In contrast, the ~~evolution of the perennial-second-year~~ fast-ice thickness ~~is not as linear~~ shows a different  
16 ~~pattern~~: Considering the large sea-ice thickness of around 2 m, as well as the insulating effect of the thick snow cover on top,  
17 the contribution of congelation growth is very limited. Instead, it is highly likely that dynamical growth as well as growth  
18 related to the consolidation of the platelet layer dominates the thickening of the perennial fast ice, adding up to an average  
19 thickness of  $4.19 \pm 1.90$  m, which is even more than double the thickness of seasonal sea ice in the bay.


#### 21 4.4 Sea-ice growth history

22 A detailed study of sea ice crystal fabric by means of visual inspection of thick/thin sections or with the help of an automated  
23 fabric analyzer can help ~~immensely-greatly~~ to determine the dominant growth processes in a given area of interest. At the same  
24 time, the growth history of fast ice is to a large degree governed by the timing of the formation of a persistent ice cover, and  
25 can only be interpreted accurately by the help of as much auxiliary information as possible, most importantly from regular  
26 satellite imagery such as MODIS, Sentinel-1 or Radarsat.

27 It has been planned since the start of the AFIN monitoring at Atka Bay in 2010 to regularly obtain sea ice cores for crystal  
28 fabric analysis. A set of cores from the six main sampling sites (Figure 1) has been obtained in 2011, and again in 2012. Only  
29 4 out of these 12 cores have been processed so far (all from 2012), which is obviously only a very small sample size compared  
30 to the decade of measurements shown above. While the limited ice core data thereby is insufficient to make general statements  
31 about sea ice growth processes at Atka Bay, we provide this data here to highlight a few major aspects, some of which have  
32 already been discussed earlier.

33 From the (limited) data we have from the four 2012 cores (Figure 9), it is evident that 1. there is no columnar texture at all; 2.  
34 there is a small fraction of granular ice in the top parts of three cores; 3. there is a small fraction of snow ice in one core and  
35 4. all cores are dominated by incorporated platelet ice. The core from the western part of Atka Bay (ATKA03) exhibits a  
36 comparably high fraction of granular ice: a 0.5m long section at the top, and 2 smaller sections a little bit deeper, with some  
37 incorporated platelet ice in between. This crystal fabric is a manifestation of the dynamic conditions under which the initial  
38 growth takes place, and supports the other datasets shown above. The strong easterly winds (Figure 3) keep pushing the initially  
39 forming thin ice towards the western ice shelf edge, which leads to a grinding of the fragile frazil crystals, and subsequently  
40 to a rafting of the newly formed ice. This process seems to be still relevant even after the ice has thickened to  $>0.5$  m, probably  
41 by very strong winds. In this way, the thickening rate of the sea ice is greatly accelerated initially (Figure 4). The absence of  
42 exclusively columnar ice is evidence that there are already platelet crystals emerging from the cavity very early in the season.  
43 While it has been suggested in an earlier study that such crystals would be present in the bay from June onwards (Hoppmann  
44 et al., 2015b), there is a possibility that they might arrive even earlier, at least in parts of the bay close to the outflow of ISW.  
45 While the ice core taken at ATKA11 is not representative at all for sea ice in the bay due to an early breakup event and  
46 subsequent late refreezing, the presence of snow ice is an evidence for a process that we would argue plays an underestimated

47 role in this region. However, we currently do not have any more direct evidence for the wide presence of snow ice at Atka Bay  
 48 (due to the lack of ice core data) other than the observations of negative freeboard in our main dataset (Figure 5), and several  
 49 observations of extensive surface flooding from summer campaigns ([although the latter do not necessarily result in snow-ice  
 50 formation due to the usually warm isothermal temperature profile in the summer ice cover](#)). In order to fill this knowledge  
 51 gap, a dedicated program of obtaining much more core sections from the top of the sea ice at different locations would have  
 52 to be implemented, with a subsequent crystal fabric and/or oxygen isotope analysis. As indicated above, this is currently not  
 53 feasible. The other ice cores taken at ATKA21 and ATKA24 are close to the “typical” sea ice thickness at Atka Bay of 2 m,  
 54 and exhibit the expected granular ice at the top from wind and waves, and incorporated platelet ice throughout the rest of the  
 55 core. [Again, ATKA21 does not show snow-ice formation at the top which might be expected from the snow depth \(Figure 4\)  
 56 and freeboard/water levels \(Figure 5\) measurements](#). No evidence from dynamic growth processes is found in these cores.  
 57 This is in line with our knowledge so far, especially since the sea ice in that area of the bay typically forms later in the year  
 58 and is less influenced by strong winds.  
 59




60  
 61 **Figure 9:** Sea-ice crystal fabric from ice cores obtained at four different fast ice sampling sites in December 2012, derived  
 62 from vertical and horizontal thin sections (0.1m spacing) along the full core length (see also Hoppmann et al., 2015a;  
 63 Hoppmann et al., 2015b; Hoppmann, 2015).

64

65 **4.5 Implications for multi-disciplinary research**

66 Such a multi-layered, thick sea-ice cover not only very efficiently separates the atmosphere from the ocean with respect to ice  
 67 growth, but it also influences the exchange of any fluxes between the two climate system components. Thereby, it also strongly  
 68 impacts the ice-associated ecosystem, which is particularly unique in sub-ice platelet layers (Arrigo, 2014). Günther and  
 69 Dieckmann (1999) concluded from their study that about 99% of the total fast-ice biomass in Atka Bay originates from algae  
 70 initially growing in the sub-ice platelet layer. The maximum Chl-a concentration in their study was around  $490 \text{ mg m}^{-3}$  in the  
 71 bottom of the fast ice, and  $240 \text{ mg m}^{-3}$  in the platelet layer in summer, at a site that had up to 0.35 m of snow cover. The authors  
 72 argued that their total observed fast ice biomass was significantly lower compared to the mostly snow-free fast ice of the Ross  
 73 Sea. However, it was still in the very upper range of biomass usually found in Antarctic fast ice (Meiners et al., 2018). At the  
 74 same time, more recent results from 2012 at Atka Bay reveal that Chl-a concentrations can reach up to  $900 \text{ mg m}^{-3}$  when there  
 75 is much less snow present (Figure- 109).

76



77

78 **Figure 10:** Sea-ice physical and biological properties from cores obtained at different fast ice sampling sites in Nov/Dec 2012  
 79 (after Hoppmann et al., 2013).

80

81 While a few studies exist that investigate shade-adaptation in algae and link algal growth to snow depth on McMurdo Sound  
82 fast ice (e.g. Sullivan et al., 1985; McGrath Grossi et al., 1987; Robinson et al., 1995), so far still comparably little is known  
83 about the adaptation of the ecosystem in the upper ocean to perennial fast-ice conditions and sub-ice platelet layers. These and  
84 similar knowledge gaps that exist with respect to ice-shelf influenced fast-ice regimes can only be addressed by integrated,  
85 multi-disciplinary research in comparably easy to access locations in coastal Antarctica, one of which was introduced in this  
86 physical study.

## 87 **5 Conclusions**

88 This study presents a unique, 9-year long record (2010 to 2018) of snow depth, freeboard, sea-ice and sub-ice platelet-layer  
89 thickness observed at Atka Bay, a coastal Antarctic fast-ice regime in the southeastern Weddell Sea and key region in the  
90 Southern Ocean. As one of the longest time series within the Antarctic Fast Ice Network, and complementary to similar records  
91 in the Ross Sea (Brett et al., 2020; Langhorne et al., 2015 and references therein), this dataset is expected to serve as an  
92 important baseline in the context of climate change and future sea-ice evolution in this region, and will contribute to an  
93 enhanced understanding of the complex interactions between the atmosphere, sea ice, ocean and ice shelves in the Southern  
94 Ocean.

95 For the period of the study presented, and considering individual observations from the 1980s and 1990s, a predominantly  
96 seasonal character of the fast-ice regime in Atka Bay is evident without a noticeable trend for any of the analyzed variables.  
97 The absence of any trend and the seasonality of surface characteristics associated with the year-round snow cover and  
98 negligible surface melting coincides with the prevailing conditions in the Antarctic pack-ice zone. Hence, the described  
99 observations in Atka Bay over the last nine years not only allow to document a baseline of the observed parameters, but also  
00 to capture processes and properties prior to expected future changes of pack ice in, e.g., the Weddell Sea, due to a changing  
01 climate.

02 Atka Bay is dominated by strong cyclonic events leading to easterly winds which determine not only the freeze-up of the bay  
03 in autumn and breakup during summer months, but also govern the year-round snow redistribution on the ice. The consequent  
04 substantial annual snow accumulation determines both, the magnitude and duration of congelation sea-ice growth, as well as  
05 the magnitude and spatial distribution of the frequent negative freeboard and related flooding of the snow/ice interface, and  
06 thus subsequent potential snow-ice formation. In contrast, platelet ice contributes significantly to the total sea-ice mass balance  
07 in this region, both, in its unconsolidated form as an underlying (buoyant) platelet layer, as well as through its incorporation  
08 into the solid sea ice (see also Hoppmann et al., 2015a; Hoppmann et al., 2015b). However, our results indicate that, although  
09 the platelet layer partly offsets the negative freeboard, it is not buoyant enough to lift the snow/ice interface above sea level  
10 against the prevalent weight of the prevalent snow.

11 With regard to the platelet layer and its formation process, we conclude that, although the annual platelet-layer thickness  
12 increase of four meters seems to be independent of the age of the fast ice in the bay, the seasonal and inter-annual variability

13 of this layer and thus the associated ocean properties and processes cannot be understood sufficiently by just considering the  
14 fast-ice properties alone. We therefore recommend to follow the approach of the New Zealand research program at Scott Base  
15 to generally include an oceanographic component into any fast-ice monitoring, especially in regions where ice shelves are  
16 present. This combination would allow for quantifying the seasonal interactions between sea ice, ocean and shelf ice even  
17 more precisely and thus to better understand current patterns and accumulation rates of platelet ice and associated biomass  
18 under the ice as a function of the distance to the shelf-ice and sea-ice edge. These results would provide a solid basis to be  
19 applied to all fast-ice areas around Antarctica, and thus make a fundamental contribution to the understanding of the Antarctic  
20 climate system.

## 21 **Data availability**

22 All presented meteorological data are archived in PANGAEA at <https://doi.pangaea.de/10.1594/PANGAEA.908826>. All fast-  
23 ice data are archived in PANGAEA at <https://doi.pangaea.de/10.1594/PANGAEA.908860>.

## 24 **Author contribution**

25 SA conducted most of the analyses for this paper and did the main writing with input from all co-authors. MH contributed the  
26 sea ice core data and performed the revision with input from all other authors. MN is the principal investigator of the AFIN  
27 work at Neumayer III. MH, MN and SA supervised the sea-ice measurements of the overwintering teams during the study  
28 period. MH and SA participated in field campaigns to collect parts of the presented data. HS contributed the meteorological  
29 datasets and the related analysis. AF contributed the fast-ice extent dataset and the related analysis.

## 30 **Competing interests**

31 The authors declare that they have no conflict of interest.

## 32 **Acknowledgements**

33 We are most grateful to the overwintering teams at Neumayer III from 2010 to 2018 for their conducted measurements on the  
34 fast ice in Atka Bay. Special thanks are due to the respective meteorologists of the teams who led the sea ice work on site.  
35 Also, our work and research at Neumayer III would not have been possible without the extensive support of the AWI logistics.  
36 We also acknowledge the scientific support of Christian Haas, the logistical support of Anja Nicolaus, and the technical support  
37 of Jan Rohde, all from the Sea Ice Physics section at AWI. This work was supported by the German Research Council (DFG)  
38 in the framework of the priority programme “Antarctic Research with comparative investigations in Arctic ice areas” by  
39 grants to SPP1158, HE2740/12, NI1092/2 and AR1236/1, and the Alfred-Wegener-Institut Helmholtz-Zentrum für Polar- und

40 Meeresforschung. This research was also supported under Australian Research Council's Special Research Initiative for  
41 Antarctic Gateway Partnership (Project ID SR140300001). We are grateful to two anonymous reviewers [and the editor Jean-](#)  
42 [Louis Tison](#) for their valuable input, which significantly improved the quality of the presented science.

## 43 **References**

- 44 Aoki, S.: Breakup of land-fast sea ice in Lützow-Holm Bay, East Antarctica, and its teleconnection to tropical Pacific sea  
45 surface temperatures, *Geophysical research letters*, 44, 3219-3227, 2017.
- 46 Arndt, S., Willmes, S., Dierking, W., and Nicolaus, M.: Timing and regional patterns of snowmelt on Antarctic sea ice from  
47 passive microwave satellite observations, *Journal of Geophysical Research - Oceans*, 121, 5916-5930,  
48 10.1002/2015JC011504, 2016.
- 49 Arndt, S., Asseng, J., Behrens, L. K., Hoppmann, M., Hunkeler, P. A., Ludewig, E., Müller, H., Paul, S., Rau, A., Schmidt,  
50 T., Schmithüsen, H., Schulz, H., Stautz bach, E., and Nicolaus, M.: Thickness and properties of sea ice and snow of land-fast  
51 sea ice in Atka Bay in 2010-2018, reference list of 9 datasets, Alfred Wegener Institute, Helmholtz Centre for Polar and Marine  
52 Research, Bremerhaven, PANGAEA, <https://doi.pangaea.de/10.1594/PANGAEA.908860>, 2019.
- 53 Arrigo, K. R.: Sea ice ecosystems, *Ann Rev Mar Sci*, 6, 439-467, 10.1146/annurev-marine-010213-135103, 2014.
- 54 Boebel, O., Kindermann, L., Klinck, H., Bornemann, H., Plötz, J., Steinhage, D., Riedel, S., and Burkhardt, E.: Real-time  
55 underwater sounds from the Southern Ocean, *Eos transactions*, 87, 361,366, 2006.
- 56 Brett, G. M., Irvin, A., Rack, W., Haas, C., Langhorne, P. J., and Leonard, G. H.: Variability in the Distribution of Fast Ice  
57 and the Sub-ice Platelet Layer Near McMurdo Ice Shelf, *Journal of Geophysical Research: Oceans*, 125, e2019JC015678,  
58 10.1029/2019jc015678, 2020.
- 59 Dammann, D. O., Eriksson, L. E. B., Mahoney, A. R., Eicken, H., and Meyer, F. J.: Mapping pan-Arctic landfast sea ice  
60 stability using Sentinel-1 interferometry, *The Cryosphere*, 13, 557-577, 10.5194/tc-13-557-2019, 2019.
- 61 Dempsey, D. E., Langhorne, P. J., Robinson, N. J., Williams, M. J. M., Haskell, T. G., and Frew, R. D.: Observation and  
62 modeling of platelet ice fabric in McMurdo Sound, Antarctica, *Journal of Geophysical Research-Oceans*, 115, Artn C01007  
63 Doi 10.1029/2008jc005264, 2010.
- 64 Dieckmann, G., Rohardt, G., Hellmer, H., and Kipfstuhl, J.: The occurrence of ice platelets at 250 m depth near the Filchner  
65 Ice Shelf and its significance for sea ice biology, *Deep Sea Research Part A. Oceanographic Research Papers*, 33, 141-148,  
66 1986.
- 67 Divine, D., Korsnes, R., and Makshtas, A.: Variability and climate sensitivity of fast ice extent in the north-eastern Kara Sea,  
68 *Polar Research*, 22, 27-34, 10.1111/j.1751-8369.2003.tb00092.x, 2003.
- 69 Druckenmiller, M. L., Eicken, H., Johnson, M. A., Pringle, D. J., and Williams, C. C.: Toward an integrated coastal sea-ice  
70 observatory: System components and a case study at Barrow, Alaska, *Cold Regions Science and Technology*, 56, 61-72,  
71 10.1016/j.coldregions.2008.12.003, 2009.
- 72 Eicken, H., and Lange, M. A.: Development and properties of sea ice in the coastal regime of the southeastern Weddell Sea,  
73 *Journal of Geophysical Research: Oceans*, 94, 8193-8206, 10.1029/JC094iC06p08193, 1989.



- 74 Eicken, H., Lange, M. A., Hubberten, H. W., and Wadhams, P.: Characteristics and distribution patterns of snow and meteoric  
75 ice in the Weddell Sea and their contribution to the mass balance of sea ice, *Annales Geophysicae-Atmospheres Hydrospheres*  
76 *and Space Sciences*, 12, 80-93, 10.1007/s00585-994-0080-x, 1994.
- 77 Eicken, H., Fischer, H., and Lemke, P.: Effects of the snow cover on Antarctic sea ice and potential modulation of its response  
78 to climate change, *Annals of Glaciology*, 21, 369-376, 10.3189/S0260305500016086, 1995.
- 79 Foldvik, A. a. K., T. : Thermohaline convection in the vicinity of an ice shelf, in: *Polar oceans, Proceedings of the Polar Oceans*  
80 *Conference held at McGill University, Montreal, May, 1974*, edited by: Dunbar, M. J., Arctic Institute of North America,  
81 Calgary, Alberta, 247-255, 1977.
- 82 Fraser, A. D., Massom, R. A., Michael, K. J., Galton-Fenzi, B. K., and Lieser, J. L.: East Antarctic landfast sea ice distribution  
83 and variability, 2000–08, *Journal of Climate*, 25, 1137-1156, 2012.
- 84 Fraser, A. D., Ohshima, K. I., Nihashi, S., Massom, R. A., Tamura, T., Nakata, K., Williams, G. D., Carpentier, S., and  
85 Willmes, S.: Landfast ice controls on sea-ice production in the Cape Darnley Polynya: A case study, *Remote Sensing of*  
86 *Environment*, 233, 111315, 2019.
- 87 Galley, R. J., Else, B. G. T., Howell, S. E. L., Lukovich, J. V., and Barber, D. G.: Landfast Sea Ice Conditions in the Canadian  
88 Arctic: 1983-2009, *Arctic*, 65, 133-144, 2012.
- 89 Giles, A. B., Massom, R. A., and Lytle, V. I.: Fast-ice distribution in East Antarctica during 1997 and 1999 determined using  
90 RADARSAT data, *Journal of Geophysical Research: Oceans*, 113, 2008.
- 91 Gough, A. J., Mahoney, A. R., Langhorne, P. J., Williams, M. J. M., Robinson, N. J., and Haskell, T. G.: Signatures of  
92 supercooling: McMurdo Sound platelet ice, *Journal of Glaciology*, 58, 38-50, Doi 10.3189/2012jog10j218, 2012.
- 93 Grosfeld, K., Treffeisen, R., Asseng, J., Bartsch, A., Bräuer, B., Fritzsich, B., Gerdes, R., Hendricks, S., Hiller, W., and  
94 Heygster, G.: Online sea-ice knowledge and data platform < [www. meereisportal. de](http://www.meereisportal.de) >, *Polarforschung*, 85, 143-155, 2015.
- 95 Günther, S., and Dieckmann, G. S.: Seasonal development of algal biomass in snow-covered fast ice and the underlying platelet  
96 layer in the Weddell Sea, Antarctica, *Antarct. Sci.*, 11, 305-315, 1999.
- 97 Günther, S., and Dieckmann, G. S.: Vertical zonation and community transition of sea-ice diatoms in fast ice and platelet layer,  
98 Weddell Sea, Antarctica, in: *Ann Glaciol*, edited by: Jeffries, M. O., and Eicken, H., *Annals of Glaciology, Int Glaciological*  
99 *Soc, Cambridge*, 287-296, 2001.
- 00 Haas, C.: The seasonal cycle of ERS scatterometer signatures over perennial Antarctic sea ice and associated surface ice  
01 properties and processes, *Ann Glaciol*, 33, 69-73, 10.3189/172756401781818301, 2001.
- 02 Haas, C., Thomas, D. N., and Bareiss, J.: Surface properties and processes of perennial Antarctic sea ice in summer, *Journal*  
03 *of Glaciology*, 47, 613-625, 10.3189/172756501781831864, 2001.
- 04 Hattermann, T., Nøst, O. A., Lilly, J. M., and Smedsrud, L. H.: Two years of oceanic observations below the Fimbul Ice Shelf,  
05 Antarctica, *Geophysical Research Letters*, 39, 2012.
- 06 Heil, P.: Atmospheric conditions and fast ice at Davis, East Antarctica: A case study, *Journal of Geophysical Research: Oceans*,  
07 111, 2006.

- 08 Heil, P., Gerland, S., and Granskog, M.: An Antarctic monitoring initiative for fast ice and comparison with the Arctic, *The*  
09 *Cryosphere Discussions*, 5, 2437-2463, 2011.
- 10 Hoppmann, M.: *Sea-Ice Mass Balance Influenced by Ice Shelves*, Jacobs University Bremen, 2015.
- 11 Hoppmann, M., Nicolaus, M., Hunkeler, P. A., Heil, P., Behrens, L. K., König-Langlo, G., and Gerdes, R.: Seasonal evolution  
12 of an ice-shelf influenced fast-ice regime, derived from an autonomous thermistor chain, *Journal of Geophysical Research-*  
13 *Oceans*, 120, 1703-1724, 10.1002/2014jc010327, 2015a.
- 14 Hoppmann, M., Nicolaus, M., Paul, S., Hunkeler, P. A., Heinemann, G., Willmes, S., Timmermann, R., Boebel, O., Schmidt,  
15 T., Kuhnel, M., König-Langlo, G., and Gerdes, R.: Ice platelets below Weddell Sea landfast sea ice, *Annals of Glaciology*, 56,  
16 175-190, 10.3189/2015AoG69A678, 2015b.
- 17 Hoppmann, M. R., M. E. ; Smith, I. J.; Jendersie, S.; Langhorne, P. J.; Thomas, D. N.; Dieckmann, G. S.: Platelet ice, the  
18 Southern Ocean's hidden ice: a review, *Annals of Glaciology*, 61, in review.
- 19 Hughes, K. G., Langhorne, P. J., Leonard, G. H., and Stevens, C. L.: Extension of an Ice Shelf Water plume model beneath  
20 sea ice with application in McMurdo Sound, Antarctica, *Journal of Geophysical Research: Oceans*, 119, 8662-8687,  
21 10.1002/2013jc009411, 2014.
- 22 Hunkeler, P. A., Hoppmann, M., Hendricks, S., Kalscheuer, T., and Gerdes, R.: A glimpse beneath Antarctic sea ice: Platelet  
23 layer volume from multifrequency electromagnetic induction sounding, *Geophysical Research Letters*, 43, 222-231, 2016.
- 24 Jacobs, S., Helmer, H., Doake, C., Jenkins, A., and Frolich, R.: Melting of ice shelves and the mass balance of Antarctica,  
25 *Journal of Glaciology*, 38, 375-387, 1992.
- 26 JCOMM Expert Team on Sea Ice: *WMO Sea-Ice Nomenclature I-III*, 2015.
- 27 Jeffries, M., Li, S., Jana, R., Krouse, H., and Hurst-Cushing, B.: Late winter first-year ice floe thickness variability, seawater  
28 flooding and snow ice formation in the Amundsen and Ross Seas, *Antarctic Sea Ice: Physical processes, interactions and*  
29 *variability*, 74, 69-87, 1998.
- 30 Jeffries, M. O., Krouse, H. R., Hurst-Cushing, B., and Maksym, T.: Snow-ice accretion and snow-cover depletion on Antarctic  
31 first-year sea-ice floes, *Annals of Glaciology*, 33, 51-60, 2001.
- 32 Kawamura, T., Jeffries, M. O., Tison, J.-L., and Krouse, H. R.: Superimposed-ice formation in summer on Ross Sea pack-ice  
33 floes, *Annals of glaciology*, 39, 563-568, 2004.
- 34 Kern, S., and Ozsoy-Çiçek, B.: Satellite Remote Sensing of Snow Depth on Antarctic Sea Ice: An Inter-Comparison of Two  
35 Empirical Approaches, *Remote Sensing*, 8, 450, 10.3390/rs8060450, 2016.
- 36 Kipfstuhl, J.: Zur Entstehung von Unterwassereis und das Wachstum und die Energiebilanz des Meereises in der Atka Bucht,  
37 Antarktis= On the formation of underwater ice and the growth and energy budget of the sea ice in Atka Bay, Antarctica,  
38 *Berichte zur Polarforschung (Reports on Polar Research)*, 85, 1991.
- 39 Knight, C. A.: Formation of slush on floating ice, *Cold Regions Science and Technology*, 15, 33-38, 1988.
- 40 König-Langlo, G., and Loose, B.: *The Meteorological Observatory at Neumayer Stations (GvN and NM-II) Antarctica,*  
41 *Berichte zur Polar-und Meeresforschung (Reports on Polar and Marine Research)*, 76, 25-38, 2007.

- 42 Kwok, R., Pang, S. S., and Kacimi, S.: Sea ice drift in the Southern Ocean: Regional patterns, variability, and trends, *Elem Sci*  
43 *Anth*, 5, 2017.
- 44 Langhorne, P. J., Hughes, K. G., Gough, A. J., Smith, I. J., Williams, M. J. M., Robinson, N. J., Stevens, C. L., Rack, W.,  
45 Price, D., Leonard, G. H., Mahoney, A. R., Haas, C., and Haskell, T. G.: Observed platelet ice distributions in Antarctic sea  
46 ice: An index for ocean-ice shelf heat flux, *Geophysical Research Letters*, 42, 5442-5451, 10.1002/2015gl064508, 2015.
- 47 Lei, R. B., Li, Z. J., Cheng, B., Zhang, Z. H., and Heil, P.: Annual cycle of landfast sea ice in Prydz Bay, east Antarctica,  
48 *Journal of Geophysical Research-Oceans*, 115, C02006, Artn C02006  
49 Doi 10.1029/2008jc005223, 2010.
- 50 Lemieux, J. F., Dupont, F., Blain, P., Roy, F., Smith, G. C., and Flato, G. M.: Improving the simulation of landfast ice by  
51 combining tensile strength and a parameterization for grounded ridges, *Journal of Geophysical Research: Oceans*, 121, 7354-  
52 7368, 2016.
- 53 Leonard, G. H., Purdie, C. R., Langhorne, P. J., Haskell, T. G., Williams, M. J. M., and Frew, R. D.: Observations of platelet  
54 ice growth and oceanographic conditions during the winter of 2003 in McMurdo Sound, Antarctica, *Journal of Geophysical*  
55 *Research-Oceans*, 111, Artn C04012  
56 Doi 10.1029/2005jc002952, 2006.
- 57 Leonard, G. H., Langhorne, P. J., Williams, M. J. M., Vennell, R., Purdie, C. R., Dempsey, D. E., Haskell, T. G., and Frew,  
58 R. D.: Evolution of supercooling under coastal Antarctic sea ice during winter, *Antarct. Sci.*, 23, 399-409, Doi  
59 10.1017/S0954102011000265, 2011.
- 60 Li, L., and Pomeroy, J. W.: Estimates of threshold wind speeds for snow transport using meteorological data, *J Appl Meteorol*,  
61 36, 205-213, 10.1175/1520-0450, 1997.
- 62 Mahoney, A., Eicken, H., Gaylord, A. G., and Shapiro, L.: Alaska landfast sea ice: Links with bathymetry and atmospheric  
63 circulation, *Journal of Geophysical Research: Oceans*, 112, 2007a.
- 64 Mahoney, A., Eicken, H., and Shapiro, L.: How fast is landfast sea ice? A study of the attachment and detachment of nearshore  
65 ice at Barrow, Alaska, *Cold Regions Science and Technology*, 47, 233-255, 2007b.
- 66 Mahoney, A. R., Gough, A. J., Langhorne, P. J., Robinson, N. J., Stevens, C. L., Williams, M. M. J., and Haskell, T. G.: The  
67 seasonal appearance of ice shelf water in coastal Antarctica and its effect on sea ice growth, *Journal of Geophysical Research-*  
68 *Oceans*, 116, Artn C11032  
69 Doi 10.1029/2011jc007060, 2011.
- 70 Mahoney, A. R., Eicken, H., Gaylord, A. G., and Gens, R.: Landfast sea ice extent in the Chukchi and Beaufort Seas: The  
71 annual cycle and decadal variability, *Cold Regions Science and Technology*, 103, 41-56,  
72 <https://doi.org/10.1016/j.coldregions.2014.03.003>, 2014.
- 73 Markus, T., and Cavalieri, D. J.: Snow depth distribution over sea ice in the Southern Ocean from satellite passive microwave  
74 data, *Antarctic sea ice: physical processes, interactions and variability*, 19-39, 1998.
- 75 Massom, R., Hill, K., Lytle, V., Worby, A., Paget, M., and Allison, I.: Effects of regional fast-ice and iceberg distributions on  
76 the behaviour of the Mertz Glacier polynya, East Antarctica, *Annals of Glaciology*, 33, 391-398, 2001a.

- 77 Massom, R., Jacka, K., Pook, M., Fowler, C., Adams, N., and Bindoff, N.: An anomalous late-season change in the regional  
78 sea ice regime in the vicinity of the Mertz Glacier Polynya, East Antarctica, *Journal of Geophysical Research: Oceans*, 108,  
79 2003.
- 80 Massom, R. A., Eicken, H., Haas, C., Jeffries, M. O., Drinkwater, M. R., Sturm, M., Worby, A. P., Wu, X. R., Lytle, V. I.,  
81 Ushio, S., Morris, K., Reid, P. A., Warren, S. G., and Allison, I.: Snow on Antarctic Sea ice, *Rev Geophys*, 39, 413-445,  
82 10.1029/2000rg000085, 2001b.
- 83 Massom, R. A., Hill, K., Barbraud, C., Adams, N., Ancel, A., Emmerson, L., and Pook, M. J.: Fast ice distribution in Adélie  
84 Land, East Antarctica: interannual variability and implications for emperor penguins *Aptenodytes forsteri*, *Marine Ecology*  
85 *Progress Series*, 374, 243-257, 2009.
- 86 Massom, R. A., Giles, A. B., Fricker, H. A., Warner, R. C., Legrésy, B., Hyland, G., Young, N., and Fraser, A. D.: Examining  
87 the interaction between multi-year landfast sea ice and the Mertz Glacier Tongue, East Antarctica: Another factor in ice sheet  
88 stability?, *Journal of Geophysical Research: Oceans*, 115, 2010.
- 89 Massom, R. A., Scambos, T. A., Bennetts, L. G., Reid, P., Squire, V. A., and Stammerjohn, S. E.: Antarctic ice shelf  
90 disintegration triggered by sea ice loss and ocean swell, *Nature*, 558, 383-389, 10.1038/s41586-018-0212-1, 2018.
- 91 McGrath Grossi, S., Kottmeier, S. T., Moe, R. L., Taylor, G. T., and Sullivan, C. W.: Sea ice microbial communities – VI –  
92 Growth and primary production in bottom ice under graded snow cover, *Marine Ecology - Progress Series*, 35, 153-164, 1987.
- 93 Meiners, K. M., Vancoppenolle, M., Carnat, G., Castellani, G., Delille, B., Delille, D., Dieckmann, G. S., Flores, H., Fripiat,  
94 F., Grotti, M., Lange, B. A., Lannuzel, D., Martin, A., McMinn, A., Nomura, D., Peeken, I., Rivaro, P., Ryan, K. G., Stefels,  
95 J., Swadling, K. M., Thomas, D. N., Tison, J. L., van der Merwe, P., van Leeuwe, M. A., Weldrick, C., and Yang, E. J.:  
96 Chlorophyll-a in Antarctic Landfast Sea Ice: A First Synthesis of Historical Ice Core Data, *Journal of Geophysical Research:*  
97 *Oceans*, 123, 8444-8459, 10.1029/2018JC014245, 2018.
- 98 Murphy, E. J., Clarke, A., Symon, C., and Priddle, J.: Temporal variation in Antarctic sea-ice: analysis of a long term fast-ice  
99 record from the South Orkney Islands, *Deep Sea Research Part I: Oceanographic Research Papers*, 42, 1045-1062, 1995.
- 00 Nicolaus, M., and Grosfeld, K.: Ice-Ocean Interactions underneath the Antarctic Ice Shelf Ekströmsen, *Polarforschung*, 72,  
01 17-29, 2004.
- 02 Olason, E.: A dynamical model of Kara Sea land-fast ice, *Journal of Geophysical Research-Oceans*, 121, 3141-3158,  
03 10.1002/2016JC011638, 2016.
- 04 Polyakov, I. V., Alekseev, G. V., Bekryaev, R. V., Bhatt, U. S., Colony, R., Johnson, M. A., Karklin, V. P., Walsh, D., and  
05 Yulin, A. V.: Long-Term Ice Variability in Arctic Marginal Seas, *Journal of Climate*, 16, 2078-2085, 10.1175/1520-  
06 0442(2003)016<2078:LIVIAM>2.0.CO;2, 2003.
- 07 Price, D., Rack, W., Langhorne, P. J., Haas, C., Leonard, G., and Barnsdale, K.: The sub-ice platelet layer and its influence on  
08 freeboard to thickness conversion of Antarctic sea ice, *Cryosphere*, 8, 1031-1039, 10.5194/tc-8-1031-2014, 2014.
- 09 Robinson, D. H., Arrigo, K. R., Iturriaga, R., and Sullivan, C. W.: Microalgal Light-Harvesting in Extreme Low-Light  
10 Environments in Mcmurdo Sound, Antarctica, *J. Phycol.*, 31, 508-520, 10.1111/j.1529-8817.1995.tb02544.x, 1995.
- 11 Robinson, N. J., Williams, M. J. M., Stevens, C. L., Langhorne, P. J., and Haskell, T. G.: Evolution of a supercooled Ice Shelf  
12 Water plume with an actively growing subice platelet matrix, *Journal of Geophysical Research-Oceans*, 119, 3425-3446, Doi  
13 10.1002/2013jc009399, 2014.

- 14 Schmithüsen, H., König-Langlo, G., Müller, H., and Schulz, H.: Continuous meteorological observations at Neumayer station  
15 (2010-2018), reference list of 108 datasets, Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research,  
16 Bremerhaven, PANGAEA, <https://doi.pangaea.de/10.1594/PANGAEA.908826>, 2019.
- 17 Selyuzhenok, V., Krumpfen, T., Mahoney, A., Janout, M., and Gerdes, R.: Seasonal and interannual variability of fast ice extent  
18 in the southeastern Laptev Sea between 1999 and 2013, *Journal of Geophysical Research: Oceans*, n/a-n/a,  
19 10.1002/2015JC011135, 2015.
- 20 Selyuzhenok, V., Mahoney, A., Krumpfen, T., Castellani, G., and Gerdes, R.: Mechanisms of fast-ice development in the south-  
21 eastern Laptev Sea: a case study for winter of 2007/08 and 2009/10, *Polar Research*, 36, Artn 1411140  
22 10.1080/17518369.2017.1411140, 2017.
- 23 Smith, E. C., Hattermann, T., Kuhn, G., Gaedicke, C., Berger, S., Drews, R., Ehlers, T. A., Franke, D., Gromig, R., Hofstede,  
24 C., Lambrecht, A., Läufer, A., Mayer, C., Tiedemann, R., Wilhelms, F., and Eisen, O.: Detailed Seismic Bathymetry Beneath  
25 Ekström Ice Shelf, Antarctica: Implications for Glacial History and Ice-Ocean Interaction, *Geophysical Research Letters*, 47,  
26 e2019GL086187, 10.1029/2019gl086187, 2020.
- 27 Smith, I. J., Langhorne, P. J., Frew, R. D., Vennell, R., and Haskell, T. G.: Sea ice growth rates near ice shelves, *Cold Regions  
28 Science and Technology*, 83-84, 57-70, 10.1016/j.coldregions.2012.06.005, 2012.
- 29 Sullivan, C. W., Palmisano, A. C., Kottmeier, S., Grossi, S. M., and Moe, R.: The Influence of Light on Growth and  
30 Development of the Sea-Ice Microbial Community of McMurdo Sound, in: *Antarctic Nutrient Cycles and Food Webs*, edited  
31 by: Siegfried, W., Condy, P., and Laws, R., Springer Berlin Heidelberg, 78-83, 1985.
- 32 Tamura, T., Williams, G., Fraser, A., and Ohshima, K.: Potential regime shift in decreased sea ice production after the Mertz  
33 Glacier calving, *Nature communications*, 3, 826, 2012.
- 34 Tamura, T., Ohshima, K. I., Fraser, A. D., and Williams, G. D.: Sea ice production variability in Antarctic coastal polynyas,  
35 *Journal of Geophysical Research: Oceans*, 121, 2967-2979, 2016.
- 36 Williams, G., Bindoff, N., Marsland, S., and Rintoul, S.: Formation and export of dense shelf water from the Adélie  
37 Depression, East Antarctica, *Journal of Geophysical Research: Oceans*, 113, 2008.
- 38 Yu, Y., Stern, H., Fowler, C., Fetterer, F., and Maslanik, J.: Interannual Variability of Arctic Landfast Ice between 1976 and  
39 2007, *Journal of Climate*, 27, 227-243, 10.1175/jcli-d-13-00178.1, 2014.
- 40