

5 **Dear Professor Flanner,**

Thank you very much for your handling of our revised manuscript (tc-2019-34). We have addressed, one by one, all comments made by the Reviewers. Below we provide our responses (*in italics*).

Response to the reviews:

10

Anonymous Referee #1

Thank you for the thoughtful and constructive comments. We agree with and have followed all comments from the reviewer 1 especially that the manuscript needed structural revision.

The abstract and the introduction were revised, some parts were removed for the sake of clarity and ease of comprehension.

15 The results part was strongly shortened and also revised. The raw data were mostly removed from the text.

General comments

20 This is an interesting paper, the theme that is discussed is original and relatively little investigated by the glaciology community. I don't have any concern about the methodology and about the presented data and results. It is clear that the authors have expertise in the field of environmental radioactivity and inorganic pollution. Despite these points I cannot support the publication of this paper in its current form. Now the paper would be suitable for publication in a journal specialized on radioactivity, not in a journal whose audience comes from many and diverse fields of science, as is the case with Cryosphere.

25 *We feel that results of our paper are interesting for the broader audience of The Cryosphere because the artificial radioactivity that is associated with cryoconite appears to be a potential tracer of material transfers on and within glaciers and from glaciers to their forefronts. Radioactivity surveys on glaciers and their forefronts might become a useful tool in the assessments of glacier responses to climate change. At the same time, the cryospheric community is mostly unaware of the phenomenon of fallout radioactivity (and other pollution) accumulation in cryoconite. Our results also indicate that the*
30 *cryosphere is an important, and active, element of the global cycling of airborne pollution. Our research is also in line with the recent recognition of the role of biological processes on glacier surfaces in the cryospheric and global biogeochemical cycling.*

35 The manuscript needs a deep language and structural revision. I tried to improve the first part of the paper from this perspective, but I realized that to really fix this side of the manuscript a big effort is needed. This is a job for the authors. Given the importance of this point, I feel that a true and accurate evaluation of the paper will be possible only after this first shape revision.

The structure of the manuscript was revised. Some of the subsections were merged. The language was improved by one of the co-authors (dr Andrew Milton).

40

The readability of the paper is not good, there are several parts where it is difficult to follow the flow and the many data seem not well connected, also because the interpretation of the results is a little bit poor and the paper is unbalanced: the presentation of raw data is very long and detailed (even too much in my opinion), while their interpretation and discussion are poor. There is a sort of gap between what someone is expecting to find in the paper given its title and the actual content.

45 Authors should avoid presenting and discussing extensively on numerical values and ratios in the main text, there are figures to this aim that are more useful and informative, in addition the authors could consider adding or changing some figures with new ones, where also literature data are shown, so as to strengthen the hypothesis presented in this paper. For example, the issue that Pu composition is compatible with an influence from satellite re-entry, is something of new or relatively rare, or is it common to see such results? This is not clear because the data are only compared to global fallout and very few other

cryoconite data, but the findings are not sufficiently highlighted. Would it be possible to extend the comparison (both graphically and in the main text), so as to evidence if what was found by the authors was a local signal or if it is more common? Discussion and interpretation in this sense should be expanded and improved, so as to allow a full comprehension of the paper also to people not well concerned about environmental radioactivity.

5 *Most of raw data was removed from the text. Fig. 5 presents additional data from literature (Łokas et al., 2018; Łokas et al., 2016 and our unpublished data from Arctic glaciers). The issue of ^{238}Pu source from satellite re-entry is presented in the following added sentences: “A third potential source of Pu is pure ^{238}Pu from a satellite re-entry after injection from the stratosphere into the troposphere. The ^{238}Pu enrichment was observed also in other cryoconite from Arctic glaciers (unpublished data) (Fig. 5C) and in air filters in Finland (Salminen-Paatero et al., 2012) or in air filters from NW Poland (Kierepko et al., 2016). “*

Another suggestion is to shorten the manuscript, now it is very long and the impression during the reading is that the same could have been said using less and less words. For example, I wouldn't always treat as separate, soils, cryoconite, radionuclides and heavy metals. Discussing about them together is more difficult for the authors, but it is for sure good from the perspective of the reader, who would better appreciate the importance and the novelty of this paper.

15 *The manuscript was shortened, also by removing most of raw data. The presentation of results for soils and cryoconite was merged. The results section is now divided into two subsections: “3.1. Radionuclide contents and their activity ratios in tundra soil profiles and in cryoconite samples” and “3.2. Heavy metals analyses in tundra soil profiles and in cryoconite samples”. The discussion was also merged and divided only for 2 subsections: “4.1. Radionuclides and heavy metals contamination” and “4.2. The source of radionuclides and heavy metals contamination “.*

20 The abstract should be adjusted, in its current shape it is not very informative and the many points that are touched in it sound a little bit as disconnected. Given the importance of this section it would be desirable to rewrite it. Now it is a brief summary of the entire paper, but the reader misses the main conceptual points of the paper, that in my opinion are: cryoconite is a better absorber than soil, local glacier morphology plays an important role in determining the accumulation of pollutants (this is one of the most interesting points from the glaciological perspective), probably cryoconite age also influences the process, in addition cryoconite seems capable of recording both global and more local events.

25 *The abstract was rewritten following reviewer's suggestions.*

30 Something similar is for the introduction. Also this part is not easy to read, the authors should deeply revise these sections, improving the language and the general structure of the text, maybe cutting some sections that sounds too technical for people who is not in the field of environmental radioactivity. One of the most difficult sections to follow is the results one. I suggest to the authors to merge the results and the discussion ones in one single paragraph. Now the results part consists in a presentation of number and concentrations that is not very informative. The same information is found in the supplementary tables. I see two options: 1-results are removed and its content is added to the following section, creating a single results and discussion paragraph. 2-the results part is strongly shortened and supported by figures that present the data in a graphical way (current figures are almost impossible to read).

35 *The introduction was shortened and rewritten. The language was improved. The results and discussion sections were also corrected. In the result section we left only maximum value for radionuclides and only average values for isotopic ratios. For heavy metals we left ranges because Figure 3 presents only two: Pb and Cd. See replies to the above comments.*

40 Figures must be modified, in many cases font size and details are too little. Some figures are also too dark. To make them well readable, modifications are needed. I suggest to the authors to improve figures, possibly adding some comparison between the samples considered here and ones discussed in previous publications, in particular dealing with isotopic ratios. It would be nice to understand if what is found in cryoconite is a common signal (even if amplified) or it is something of peculiar. This paper could be potentially published in TC, but several additional efforts must be pushed by the authors to this aim, therefore major revisions is my final response.

45

All figures were modified. Fig 1 was lightened and we removed the Fig 1B with the recession of Waldemarbreen. Now the Fig 1 is better readable.

The Figures 2 and 3 were merged (for Fig 2 corrected) and we selected only airborne radionuclides for cryoconite and soil profiles, we removed graphs with natural radionuclides like U and Th isotopes because these data are presented in the supplementary material and also we found higher differences in radioactivity between cryoconite and soils for airborne radionuclides than for natural radionuclides.

We also merged Figures 3 and 4 (now Fig 3). This new figure shows only two, most interesting heavy metals Cd and Pb in cryoconite and soil profiles. We removed other heavy metals from the graph because these data are presented in supplementary material.

We also present additional data for cryoconite (some of that are not published yet) in the corrected Fig 5 (now Fig 6).

The presentation of raw data is very long and detailed, while their interpretation and discussion are poor.

The authors could consider adding or changing some figures with new ones, where also literature data are shown

The result was divided for two subsection: "Radionuclide contents and their activity ratios in tundra soil profiles and in cryoconite samples" and "Heavy metals analyses in tundra soil profiles and in cryoconite samples". Generally, this section was shortened, figures were merged and the selected airborne radionuclides in soils and cryoconite and selected heavy metals in soils and cryoconite and presented on Fig 2 and 3, respectively. Fig 1 was modified, it was lightened and larger font size was used. In Fig 5 we added some new literature data (Łokas et al., 2018; Łokas et al., 2016 and our unpublished data from Arctic glaciers).

Another suggestion is to shorten the manuscript, now is very long and the impression during the reading is that the same could have been said using less and less words eg. I wouldn't always threat as separate, soils, cryoconite, radionuclides and heavy metals....

The manuscript was shortened, the result for soils and cryoconite were merged. The results section was divided into two subsections: "Radionuclide contents and their activity ratios in tundra soil profiles and in cryoconite samples" and "Heavy metals analyses in tundra soil profiles and in cryoconite samples"

Abstract should be adjusted

Abstract was adjusted as suggested by the reviewer.

Introduction should be revised

The introduction was revised, some parts were removed for the sake of clarity

Results should be strongly shortened...

Corrected

Figures must be modified, in many cases font size and details are too little. Some figures are also dark.

Corrected. See above.

Specific comments

Page 1. Line 20: "... changed weathered with local"

Corrected to: "...their activity concentrations are controlled only by mixing of local material derived from different bedrock."

Line 23: I would change to "from additional and more specific sources might be..."

Corrected.

Line 24: change "is visible" with "was detected" and "are" with "is"

Corrected to: "We assumed that the main source of Pu, which was detected only in cryoconite samples, is derived from nuclear tests and non-exploded weapons-grade material."

5 Line 25-28: change "Approximately one third of the total observed Pu activity concentration is related to ²³⁸Pu and can be explained considering re-entry of the SNAP9A satellite, which was powered by a Pu thermoelectric generator."

Changed to: "Approximately one third of the total observed Pu activity concentration is ²³⁸Pu originating from a SNAP9A satellite re-entry, which was powered by a Pu thermoelectric generator."

Line 25-28: this passage must be rephrased

10 *Replaced as: "Local glacial morphology plays an important role for determining the accumulation of airborne pollutants."*

Page 2. Line 2: replace "elements" with "species"

Replaced

15 Line 3: ...adjust this passage

It was adjusted: "Many of the artificial radionuclides were released into the environment due to various human activities mainly in the second half of the 20th century, but the present work is focused only on the ¹³⁷Cs, ^{238,239,240}Pu, ²⁴¹Am."

Line 7: add: "... powered by nuclear thermos-electric generators"

20 *Following part was added: "...disintegration of satellites (SNAP9A-1964, Cosmos 958-1978) powered by nuclear thermos-electric generators."*

Line 11: what is the non-proliferation of nuclear material?

25 *Replaced with: "...radioactivity is closely connected with potential threats to national security and controlling of the spread of nuclear material."*

Line 15: Please reformulate, spreading the message that metals and radioactivity are only two of the several pollutant groups that affect cryosphere

30 *Corrected to: "Additionally, the presence of these artificial radionuclides and heavy metals are only two of the several pollutant groups that affect the cryosphere."*

Line 27: please specify where cryoconite is found also. It's important to say that cryoconite is only found on the ablation surface of glaciers

35 *It was reformulated: "...such holes are form on the ablation zone of glaciers and constitute nutrient-rich pools with diverse biota within the supraglacial zone"*

Line 29-30: this is an important passage, please add a reference for it

40 *It was added two references: "They produce extracellular polymeric substances whose adhesive properties enhance the accumulation of mineral dust and microorganisms (Gadd 2004; Francis 2007)."*

Line 31: change "their" with "is"

Corrected

Line 34: "Cryoconite is usually found...."

45 *Corrected*

Page 3. Line 1: "Such holes are usually found..."

Corrected to: "Such holes are usually found on the ablation zone of glaciers..."

Line 3-4: This information is not correct.

It was reformulated: “Glaciers in Canadian, Russian Arctic, Alaska, Antarctica and Greenland are projected to lose about 40 % of their volume by the end of 2100 but in Central Europe, low-latitude South America, Caucasus, North Asia, Western Canada and US are expected to lose about 80 % of their volume or disappear within decades at current climatic conditions (Radic et al., 2014).”

5

Line 5: “contain”

Corrected

Line 8-9: “provide evidence that cryoconite-derived contaminants were dispersed in the proglacial area by meltwater channels and accumulated in depressions on the glacier forefield”

10

Corrected

Line 10: which properties? May be it will be better to use features, but anyway it is not clear what authors are talking about

This sentence was removed.

15

Line 12-14: it is clear what authors are saying but it would be nice to reformulate this passage

We agree with this suggestion that this topic are quite sensitive and we refused the countries.

Page 4. Line 25: the figure is too dark and the used font is too little

20

This Fig 1 was modified.

Page 5. Line 26: introduce the coefficient r, what is it?

The citation was added: “....(eg. Mukaka, 2012)”

25

Page 6. Line 2: change “understood” with “defined”

It was corrected

Line 10: change with “this is the reason why the activity...”

30

It was corrected

Page 7. Line 1: the authors should modify the figure, increasing the font size and the boxes dedicated to the vertical profiles, now you can't read anything

35

It was modified and corrected: Figures 2 and 3 were merged and the author selected airborne radionuclides and heavy metals and presented on new Fig 2 and 3, respectively.

Page 10. Line 18-20: I guess that this is only a hypothesis, even if very likely. The authors should be more open to doubt and add a reference

40

It was rephrased and references were added: “Probably, they can retain and concentrate airborne radionuclides due to metal-binding properties of extracellular substances that are excreted by microorganisms (Gadd 2004; Francis 2007).”

Page 11. Line 1-2: please add the correlation coefficient to better appreciate the correlation degree between these variables

45

It was added: “A significant correlation was observed between the amount of organic matter present and the concentration of all airborne radionuclides ($r=0.79-0.83$). We noticed significant correlation between ^{210}Pb , ^{137}Cs , Pu isotopes and ^{241}Am with the altitude ($r=0.64-0.91$) and the area of the cryoconite holes ($r=0.58-0.63$), except ^{210}Pb ($r=0.38$). The depth of these cryoconite holes also correlates with ^{210}Pb and Pu isotopes ($r=0.60-0.68$)”

Line 5: “which were located close to the moraine at lower altitude”

50

Corrected

Line 8: please explain why here the collapse of cryoconite holes is more likely
The part of this sentence was removed.

Page 15. Line 8: what is mineral soil

5 *It was corrected: "...sandy fraction of soils.."*

Page 16. Line 19-30: this part sounds like introduction, it should be removed (or drastically shortened) or moved to the introduction

10 *It was shortened and modified. "Both radionuclides and heavy metals are deposited from the atmosphere on land/ocean and glacier surfaces by wet and dry precipitation. Since there are no new sources of radioactive pollution into the atmosphere, the patterns of atmospheric circulation do not influence radionuclide distribution within different glaciers on Svalbard. The amount of precipitation received by the glacier can definitely influence the glacier pollution, since coastal areas (eg. Kaffiøyra) receive twice the amount of precipitation compared to the inland areas (Førland et al., 2011). Post-depositional processes, mainly those connected to cryoconite hole formation and eventual material transport within the glacier, appear to*
15 *have more significant influence on the levels of radionuclide pollution within the glacier. The different scenario concerns the distribution of heavy metals. Since they are still emitted to the atmosphere from different sources, the direction of atmospheric circulation is an important factor shaping the metal pollution distribution within the glacier. Main atmospheric metal discharge origins from Europe and Russia (Isaksen et al., 2016), in coastal regions local winds can enhance the introduction of contaminants due to transport of contaminated sea salt aerosols, however (Lüdke et al., 2005 Kozak et al.,*
20 *2015). "*

Anonymous Referee #2:

General comments:

25 From an analytical perspective, this article represents a great amount of work with regard to sample collection and processing, chemical and radiochemical separations, instrumental measurements, and interpretation of results. A systematic organization of the measurement results is found in the extensive tables of analytical data; these will enable valuable comparisons with past and future data obtained by other researchers in that region.

Perhaps referring to the tables, figures and plots with very brief descriptions should comprise the "Results" section. I'm not sure that it is necessary to mention the ranges of all the analytes in the text when this information can be seen in the tables.

30 *We agree with the reviewer that the results section should be modified. The results for soils and cryoconite were linked. The results section was divided for two subsections: "Radionuclide contents and their activity ratios in tundra soil profiles and in cryoconite samples" and "Heavy metals analyses in tundra soil profiles and in cryoconite samples". Generally, this section was shortened, figures were merged and the author selected airborne radionuclides in soils and cryoconites and only two*
35 *heavy metals in soils and cryoconite and presented on Fig 2 and 3, respectively. We will present additional data on radionuclides in cryoconites (some of that are not published yet) in the corrected Fig 5 (now Fig 6).*

In the results section we left only maximum value for radionuclides and only average values for isotopic ratios. For heavy metals we left ranges because Figure 3 presents only concentrations of two metals: Pb and Cd. We removed other heavy metals from the graph because these data are presented in supplementary material.

40 The "Discussion" section is partitioned into three subsections. Subsection 4.2 "The source of radionuclides and heavy metals contamination" is essentially entirely devoted to radionuclides; perhaps it would be better just to drop the "heavy metals" part from this title since it is discussed in depth in subsection 4.3. Some of the actual discussion of the results, especially for stable heavy metals and correlations among them (subsections 3.3 and 3.4), occurs in the "Results" section. It is
45 recommended to integrate these remarks into the "Discussion" section along with the existing material.

The discussion was modified, it was merged and divided only for 2 subsections: "4.1. Radionuclides and heavy metals contamination" and "4.2. The source of radionuclides and heavy metals contamination ". Most of the raw data of radionuclides and heavy metals in the discussion were removed.

Uncertainties are presented with the tables S2 through S5 giving radionuclide measurement results; however, I cannot find any mention in the text of whether these uncertainties are at the 68% (k=1) or 95% (k=2) confidence level. A general statement about the confidence level of uncertainties should be given early in the text (e.g., at the very beginning of the “Results” section). In addition, the main contributions to the radioanalytical uncertainties should be mentioned (e.g., counting statistics, calibrations, tracer concentrations, etc.).

The information about uncertainties and confidence level was added to the Methods section: “The uncertainties of activity concentrations include the measurement uncertainties (counting statistics and efficiency) and are reported as 1 σ counting statistics.”

Table S6 lists all of the stable element measurement results, but there are no uncertainties attached to these values. A statement of uncertainty “ranges” associated with each element concentration measurement should be given for the reader, perhaps at the column headings or as a footnote.

The information about uncertainties was added to the “Methods” section: “The quality control and assurance for this method is presented in Zaborska et al. (2017).”

My recommendation is that this article should be published with minor revisions as outlined above.

Specific comments to revised article:

1. Although I believe Svalbard and Spitsbergen refer to the same archipelago, the authors should choose one or the other to be used consistently in the article.

The authors choose Spitsbergen.

2. In the abstract, it would be useful to mention that Kaffioyra region is located in Spitsbergen
“Spitsbergen” *was added.*

3. In subsection 2.3.2 “Heavy metal analyses” it is not mentioned if separate subsamples (from those used for radionuclide analyses) were used for stable element determinations. If separate samples were used, how much material and what dissolution methods were used to prepare them for the flame atomic absorption and mass spectrometer measurements?

Following text was added: “Separate sub-samples (0.5 g) were dedicated for measurements of selected heavy metals that are believed to be important anthropogenic contaminants (Pb, Cu, Zn, Cd). Contaminating metals and other metals (Fe, Al, Ni, Cr, Mn) were measured with a flame atomic absorption spectrometer (AAS Shimadzu 6800) using deuterium background correction.”

4. How was sample organic content assessed and related to LOI (which needs to be defined in the text)?

It was defined in the “Methods” section: “Organic matter contents were determined by loss on ignition (LOI) at 600° C for 6 hours.”

5. The plots of depth distribution in fig 2 are important but very difficult to read; a significant improvement in quality is needed.

Figures 2 and 3 were merged and the authors selected airborne radionuclides and selected heavy metals and presented on new Fig 2 and 3.

6. The reference “Łokas et al., 2010” in line 10 on page 5 is missing.....
It was added to the “References” section.
- 5 7. In line 6 on page 17, “ average mass ratios” should be “average atom ratios” to be consistent with all of the previous text.
Corrected.
- 10 8. In line 8 on page 17, “.....initial soils from other west cost of Spistbergen” is confusing.....
It was corrected as “.....initial soils from other west coast of Spitsbergen”.
- 15 9. In line 8-9 on page 17, perhaps to write “We observed an excess of ^{238}Pu in cryoconite samples similar to that which we observed in other initial soils in Spitsbergen, and this enrichment of ^{238}Pu is derived from sources other than ^{239}Pu . This is due to pure ^{238}Pu from a satellite re-entry after injection from the stratosphere into the troposphere.”
Corrected.

Airborne radionuclides and heavy metals in High Arctic terrestrial environment as the indicators of sources and transfers of contamination

20 Edyta Łokas¹, Agata Zaborska², Ireneusz Sobota³, Paweł Gaca⁴, Andrew Milton⁴, Paweł Kocurek⁵, Anna Cwanek¹

¹Department of Nuclear Physical Chemistry, Institute of Nuclear Physics Polish Academy of Sciences, Kraków, 31-342, Poland

25 ²Marine Chemistry and Biochemistry Department, Institute of Oceanology Polish Academy of Sciences, Sopot, 81-712, Poland

³Department of Hydrology and Water Management, Polar Research Centre, Nicholas Copernicus University, Toruń, 87-100, Poland

⁴Ocean and Earth Science, University of Southampton, National Oceanography Centre, European Way, Southampton, SO14 3ZH United Kingdom

30 ⁵Research and Development Laboratory for Aerospace Materials, Rzeszow University of Technology, Rzeszow, 35-959, Poland

Correspondence to: Edyta Łokas (Edyta.Lokas@ifj.edu.pl)

Abstract. A survey of airborne radioactive isotopes (^{137}Cs , ^{238}Pu , $^{239+240}\text{Pu}$, ^{241}Am and ^{210}Pb) and trace metals (Pb, Cu, Zn, Cd, Fe, Al) in tundra soils and cryoconite hole material sampled from several locations in the Kaffiøyra [of Spitsbergen](#) revealed significant variability in spatial concentration. Lithogenic radionuclides (^{230}Th , ^{232}Th , ^{234}U , ^{238}U) show less

35

~~variability than the airborne radionuclides because their activity concentrations are controlled by mixing of local material derived from different types of bedrock.~~

~~region revealed high spatial concentration variability of the analysed samples. Lithogenic radionuclides (^{230}Th , ^{232}Th , ^{234}U , ^{238}U) show less variability than the airborne radionuclides because their activity concentrations are controlled only by mixing of weathered material derived from different bedrock.~~

Activity ratios of the artificial radionuclides differ in most cryoconite samples from global fallout signatures. The contribution of radionuclides from ~~other~~ additional and more specific sources might be enhanced by non-continuous exposure of cryoconite to atmospheric deposition. We assumed that the main source of Pu, which ~~is visible~~ was detected only in cryoconite samples, ~~are is~~ derived from nuclear tests and non-exploded weapons-grade material. Approximately one third of the total observed Pu activity concentration is ^{238}Pu originating from a SNAP9A satellite re-entry, ~~try~~ which was powered by a Pu thermoelectric generator ~~try and subsequent injection of nuclear debris from stratosphere into troposphere~~. In samples from Waldemarbreen we ~~could appreciate the influence of observed the effect of~~ glacial local morphology on the capability of cryoconite effective of trapping and accumulating/storing of airborne radionuclides. Local glacial morphology plays an important role for determining the accumulation of airborne pollutants. ~~The differences in the concentrations of radionuclides between sampling points and differences in the elevation gradient from terminus towards icefall may reflect the homogenous topography of the glacier tongue.~~ The trace metal concentrations in soils were typical or slightly higher than concentrations characteristic for natural background concentrations and the $^{206}\text{Pb}/^{207}\text{Pb}$ ratio also was close to the natural ratio for parent rocks. Conversely, trace metal concentrations in cryoconite samples (Pb and Cd) were higher than in soil samples and definitely exceeded natural values. As a consequence of this research the cryoconite can be considered as a better absorber for airborne contaminants than soil.

1 Introduction

The radioactive nuclides in the environment originate from both natural and anthropogenic sources. The present work is focused on the ^{137}Cs , $^{238,239,240}\text{Pu}$, and ^{241}Am which have been produced through nuclear fission and neutron activation processes and have been released to the environment mainly in the second half of the 20th century. Main sources of anthropogenic radionuclides in Northern Hemisphere are: (i) nuclear weapon tests, atmospheric explosions in Novaya Zemlya, Semipalatinsk, Nevada etc.; (ii) nuclear accidents (Kyshtym-1957, Lake Karchay-1968; Tomsk-1993; Chernobyl-1986; Fukushima-2011); (iii) disintegration of satellites (SNAP9A-1964, Cosmos 958-1978) powered by nuclear thermoelectric generators. Relatively low but constant releases of artificial radioisotopes are also associated with reprocessing of spent nuclear fuel and energy generation in nuclear power plants. Heavy metals are natural elements of the Earth's crust. Since the beginning of the industrial era (from ~1850) loading of metals to the environment due to human activities has increased nearly 10 times, however (Nriagu, 1996). The emission of heavy metals increased substantially after World War II and is still increasing in some countries (Asian sector). The main anthropogenic heavy metal sources include industry,

5 mining, agriculture, fuel burning, waste disposal and transportation (Pacyna and Pacyna, 2001). Artificial radionuclides and heavy metals are transported over the whole world mainly by the atmosphere, oceanic currents, and rivers. These contaminants can be found also in remote Earth locations eg. polar cryosphere. Elevated radionuclide and heavy metal levels have been reported in Asian and European glaciers (Barbante et al., 2004; Aizen et al., 2009; Eyrikh et al., 2017; Baccollo et al., 2017) and even in remote Arctic and Antarctic areas (Hur et al., 2007; Singh et al., 2013).

10 The highest amounts of pollutants on glaciers are most likely stored in cryoconite granules and micro-fauna (Segawa et al., 2013; Łokas et al., 2016; Baccollo et al., 2017; Zawierucha et al., 2018). Cryoconite granules are aggregates of mineral and organic components composed of archaea, algae, cyanobacteria, fungi and heterotrophic bacteria consortia (Cook et al., 2015; Simon et al., 2002; Takeuchi et al., 2001a,b; Hodson et al., 2008). Cyanobacteria play a crucial role in formation of such aggregates as they produce extracellular polymeric substances (EPS) whose adhesive properties enhance the accumulation of mineral dust and microorganisms (Gadd 2004; Francis 2007). Cryoconite holes are formed on the ablation zone of glaciers and constitute nutrient-rich pools with diverse biota within the supraglacial zone (Wharton et al., 1985; Cook et al., 2016). Dark-colored cryoconites significantly increase the absorption of solar light by glaciers, enhancing and speeding up melting processes (Simon et al., 2002; Takeuchi et al., 2001a,b; Hodson et al., 2008). Glaciers in Canadian, Russian Arctic, Alaska, Antarctica and Greenland are projected to lose about 40 % of their volume by the end of 2100 but in Central Europe, low-latitude South America, Caucasus, North Asia, Western Canada and US are expected to lose about 80 % of their volume or disappear within decades at current climatic conditions (Radic et al., 2014). Cryoconite granules contain high concentrations of contaminants and can persist on glacier surfaces for several decades (Łokas et al., 2014), nevertheless, as the glacier ice disappears, their contents is deposited on the freshly exposed areas. The discovery of hot spots of radioactivity in the proglacial zones of Spitsbergen glaciers (Łokas et al., 2014; Łokas et al., 2017b) provided evidence that cryoconite-derived contaminants were dispersed in the proglacial area by meltwater channels and accumulated in depressions of the glacier forefield.

25 The monitoring of anthropogenic radionuclides and heavy metals environmental distribution and fate is important as both pose a potential threats to the environment. Nowadays, during the intense cryosphere melting contaminants accumulated on the glaciers surface over the last decades are released and can pollute local vegetation, surface waters and even drinking water sources (Dong et al., 2017). Moreover, determination of Pu isotopes, ^{241}Am and ^{137}Cs in cryoconites is of great interest to monitor environmental impact of military nuclear technologies development and the rising number (and risk) of nuclear events in civilian nuclear installations as well as the growing risk of nuclear terrorism and undeclared nuclear activities.

30 The main objective of this study is to evaluate the concentrations of artificial (^{137}Cs , $^{238,239+240}\text{Pu}$, ^{241}Am) and natural (^{210}Pb , ^{230}Th , ^{232}Th , ^{234}U , ^{238}U) radionuclides and trace metals (Pb, Cu, Zn, Cd, Fe, Al) in cryoconite holes and tundra soils sampled in several locations in the Kaffiøyra region (Spitsbergen) and to identify and constrain the different sources of contamination based on the isotopic ratios of measured radionuclides and metals ($^{238}\text{Pu}/^{239+240}\text{Pu}$, $^{241}\text{Am}/^{239+240}\text{Pu}$, $^{239+240}\text{Pu}/^{137}\text{Cs}$, $^{240}\text{Pu}/^{239}\text{Pu}$, $^{206}\text{Pb}/^{207}\text{Pb}$ and $^{208}\text{Pb}/^{206}\text{Pb}$). The occurrence of radioactive elements in the environment is related

to both natural and anthropogenic factors. The artificial radionuclides (^{137}Cs , $^{238,239,240}\text{Pu}$, ^{241}Am) were released into the environment due to various human activities mainly in the second half of the 20th century. They were produced through nuclear fission and neutron activation processes. Main sources of anthropogenic radionuclides in Northern Hemisphere are: (i) nuclear weapon tests, atmospheric explosions in Novaya Zemlya, Semipalatinsk, Nevada etc.; (ii) nuclear accidents (Kyshtym 1957, Lake Karchay 1968; Tomsk 1993; Chernobyl 1986; Fukushima 2011); (iii) disintegration of satellites (SNAP9A 1964, Cosmos 958 1978). Relatively small but constant releases of artificial radioisotopes are also associated with reprocessing of spent nuclear fuel and energy generation in nuclear power stations. Studies of the presence of anthropogenic radionuclides in the environment have entered a new era with the renaissance of nuclear energy and associated fuel reprocessing. The renewed interest in environmental radioactivity is closely connected with potential threats to national security and non proliferation of nuclear material. Additionally, the presence of these artificial radionuclides Heavy metals are natural elements of the Earth's crust. Since the beginning of the industrial era (from 1850) loading of metals to the environment due to human activities has increased nearly 10 times (Nriagu, 1996). The emission of heavy metals increased substantially after World War II and is still increasing in some countries (Asian sector). The main anthropogenic heavy metal sources include industry, mining, agriculture, fuel burning, waste disposal and transportation (Pacyna and Pacyna, 2001). Heavy metals pollution usually occurs locally but can be also transported globally by air mass circulation, rivers, oceanic currents, etc. Heavy metals have been found in Asian and European glaciers (Barbante et al., 2004; Aizen et al., 2009; Eyrikh et al., 2017; Baccollo et al., 2017) and even in remote Arctic and Antarctic areas (Hur et al., 2007; Singh et al., 2013). As they melt, glaciers release metals to streams that may pollute both vegetation and drinking water sources (Dong et al., 2017).

The highest concentrations of pollutants on glaciers are most likely stored in cryoconite granules and micro fauna (Segawa et al., 2013; Lokas et al., 2016; Baccollo et al., 2017; Zawierucha et al., 2018). Cryoconite granules are aggregates of mineral and organic components where it is possible to identify biological consortia composed of archaea, algae, cyanobacteria, fungi and heterotrophic bacteria (Cook et al., 2015; Simon et al., 2002; Takeuchi et al., 2001a,b; Hodson et al., 2008). Cyanobacteria play a crucial role in formation of such aggregates. They produce extracellular polymeric substances whose adhesive properties enhance the accumulation of mineral dust and microorganisms. Among the many effects played by cryoconite in glacial environments, their influence on ice albedo is extremely relevant. Given their dark color, cryoconite is able to significantly increase the absorption of solar light by glaciers, enhancing and speeding up melting processes (Simon et al., 2002; Takeuchi et al., 2001a,b; Hodson et al., 2008). Cryoconites are usually found in spatially confined holes, produced by locally increased ice melting caused by the dark color of cryoconite granules themselves. They usually form on the ablation zone of glaciers and constitute nutrient rich pools with diverse biota within the supraglacial zone (Wharton et al., 1985; Cook et al., 2016). Glaciers all over the world are projected to lose 80 % of their volume by the end of 2100 and some of them are expected to disappear within decades at current climatic conditions (Radic et al., 2014). Cryoconite granules containing high concentrations of contaminants and can persist on glacier surfaces for decades (Lokas et al., 2014), nevertheless, as the glacier ice disappears, they deposit their contents on the freshly exposed areas. The discovery

of hot spots of radioactivity in the proglacial zones of Spitsbergen glaciers (Lokas et al., 2014; Lokas et al., 2017b) provides evidence that cryoconite derived contaminants may have accumulated in depressions of the glacier forefield, together with fines redistributed by slope wash.

A critical feature of a cryoconite is its accumulation capability. Given these properties, cryoconite samples are among the best candidates to monitor the spreading of artificial radionuclides and heavy metals into their surrounding environments. Determination of Pu isotopes, ^{241}Am and ^{137}Cs in cryoconite samples is of great interest for several reasons; as a radioecological monitoring tool because of the development of nuclear technologies for military purposes (India, Pakistan and potentially Iran) and the rising number (and risk) of nuclear events in civilian nuclear installations as well as the growing risk of nuclear terrorism and undeclared nuclear activities.

The main objective of this study is to evaluate the concentrations of artificial (^{137}Cs , $^{238,239+240}\text{Pu}$, ^{241}Am) and natural (^{210}Pb , ^{230}Th , ^{232}Th , ^{234}U , ^{238}U) radionuclides and trace metals (Pb, Cu, Zn, Cd, Fe, Al) of cryoconite holes and tundra soils sampled in several locations in the Kaffiøyra region, Svalbard (Fig. 1A) and to identify and constrain the different sources of contamination based on the isotopic ratios of radionuclides (^{238}Pu , $^{239+240}\text{Pu}$, ^{241}Am , $^{239+240}\text{Pu}$, $^{239+240}\text{Pu}$, ^{137}Cs , ^{240}Pu , ^{239}Pu , ^{206}Pb , ^{207}Pb and ^{208}Pb , ^{206}Pb).

15 2 Data and methods

2.1 Study area

The Kaffiøyra region, located in the north-western Spitsbergen, the adjoining Aavatsmarkbreen (75 km²) and the Dahlbreen (132 km²) together comprise an area of about 310 km² (Sobota et al., 2016). It is a coastal lowland at Forlandsundet (Forland Sound). It accounts for 12 % of the area of Oscar II Land. Along with Kaffiøyra, which is 14 km long and 4 km wide, seven land glaciers are located in this region. The main pedogenetic processes in the Kaffiøyra region consist of the accumulation of organic material, gleying and decarbonation. Cryogenic processes, which overlap the pedogenetic processes, also play an important role in the genesis of the soils there. The substantial moisture content of the surface layers and stagnant waters result in a shallow permafrost (Plichta, 2005). In the years 1996–2015, the greatest average active layer thickness of permafrost (215 cm) was recorded at a location in the moraine. A much lower mean value was recorded on the beach (126 cm). An active layer in the tundra reached on average a depth of 150 cm (Sobota et al. 2018). Climate research regarding the Kaffiøyra region has been conducted since 1975. The increase in air temperature noted during the study confirms the general trend observed in Svalbard and in many areas of the Arctic (Nørdli et al., 2014). In the years 1997–2016 an average air temperature during the summer season in this region was 5.4°C.

Glaciers of this region are polythermal (Sobota, 2009, 2011). From the time of their maximum extent in the late 19th and early 20th centuries to 2015, the total area of this region's valley glaciers has decreased on average by about 43 % (Sobota et al. 2016). The Waldemarbreen is an alpine-type glacier running down the valley towards the Kaffiøyra. It contains one accumulation zone and a glacier tongue in the valley. The area of the glacier is 2.40 km² (Sobota, 2017). It borders the

Prins Heinrichfjella ridge in the north and east (500–770 m above sea level), and Gråfjellet (300–350 m) in the south. The glacier consists of two distinct parts separated by a medial moraine. The mean annual mass balance of the Waldemarbreen in 1996–2015 was -0.72 m w.e. (Sobota et al. 2016) and in 2014 ablation of glacier also was very significant. Starting from 1909, the glacier has been receding by 8 m a^{-1} . Between years 1995–2009 the recession rate accelerated to 10 m a^{-1} , and between 2000–2009 reached 11 m a^{-1} (Sobota and Lankauf, 2010). From the time of the maximum advance to 2015 the Waldemarbreen area decreased by 31.4 % and the annual average recession rate was 8 m a^{-1} (Sobota et al., 2016) (Fig. 1B).

2.2 Field campaign

~~Sample locations are shown. The locations of sampling sites are presented~~ in Fig. 1A. Soil samples were collected in 2009 (coded: S01-S06). Sampling sites represent accumulations of typical tundra samples. Generally, this area is predominantly covered with dry moss and lichen tundra with communities of *Saxifraga oppositifolia*, *Salix polaris* – *Sanionia uncinata*, *Luzula arcuata* ssp. *confusa* – *Cetrariella delisei* – *Scorpidium revolvens* (Sobota et al. 2018). Soil profiles S01-S06 were collected at increasing distances from the front of Waldemarbreen (soil profile S06 was located on the flat outwash fan at the foreland of the moraine ridge of Aavatsmarkbreen). The profile depths varied from 12 to 17 cm. All these profiles were covered by a layer of organic material. All soil cores were divided into 1–2 cm sub-samples and dried at 105° C to a constant weight, passed through a 2 mm sieve and then prepared for radionuclide analysis. Additionally, twelve cryoconite samples (coded: 2-13) were collected in the Waldemarbreen in August-September 2014.

~~The samples were taken from cryoconite holes at different altitudes, depths and surface areas (see Tab. S1). Material was collected with disposable plastic Pasteur pipettes and transferred to 15 cm³ plastic test tubes. with disposable plastic Pasteur pipettes from the bottom of randomly selected cryoconite holes and transferred to 15 cm³ plastic test tubes. Cryoconite samples in this study were taken from different altitudes, from cryoconite holes characterised by different depths and surface areas (Tab. S1).~~





Figure 1. Study area location. Profile S01-S06 were collected at increasing distances from the front of Waldemarbreen and cryoconite samples (2-13) from different altitudes of this glacier.

~~Figure 1. A. Location of the study area. Profile S01-S06 were collected at increasing distances from the front of Waldemarbreen and cryoconite samples (2-13) from different altitudes of this glacier. B. Map of recession of the Waldemarbreen in 1909-2017.~~

2.3 Methods

2.3.1 Radionuclide analyses

Gamma emitting radionuclides (^{137}Cs and ^{210}Pb) were measured by planar HPGe (high-purity germanium) detector. The activities of ^{137}Cs were determined using its emission peak at 662 keV and for ^{210}Pb the 46.6 keV. Corrections were made for the effect of self-absorption of low-energy g-rays (46.6 keV) within the sample even though these corrections were insignificant because of very low sample masses (from 2 to 12 g dry masses). The activities of ^{238}Pu , $^{239+240}\text{Pu}$, ^{241}Am , $^{234,238}\text{U}$ and $^{230,232}\text{Th}$ were determined for 0.5–1.15 g cryoconite samples and 5 g dried soil samples. Organic matter contents ~~were determined by loss on ignition (LOI) was decomposed by a muffle oven ignition~~ at 600° C for 6 hours. The samples were then dissolved using concentrated acids (HF, HNO₃, HCl)₃ ~~and a small addition of H₃BO₃~~. Details of the sequential radiochemical procedure used for determination of ^{238}Pu , $^{239+240}\text{Pu}$, ^{241}Am , $^{234,238}\text{U}$ and $^{230,232}\text{Th}$ are described in previous

publications (Łokas et al., 2010; Łokas et al., 2018). The $^{240}\text{Pu}/^{239}\text{Pu}$ ratio was determined using a Thermo Fisher Scientific Neptune MC-ICP-MS (analytical method based on Taylor et al. 2001; Łokas et al., 2018). The combined measurement uncertainties of activity concentrations, which include counting statistics and counting efficiency uncertainties, are reported as 1σ . The Environmental Radioactivity Laboratory of the Henryk Niewodniczanski Institute of Nuclear Physics holds an ISO17025 accreditation for gamma-spectrometric measurements and Pu analyses by alpha-particles spectrometry.

~~The Environmental Radioactivity Laboratory of the Henryk Niewodniczanski Institute of Nuclear Physics holds an ISO17025 accreditation for gamma-spectrometric measurements and Pu analyses by alpha-particles spectrometry.~~

2.3.2 Heavy metal analyses

Sub-samples (0.5 g) were taken for measurement of selected heavy metals considered to be important anthropogenic contaminants (Pb, Cu, Zn, Cd). These metals and others (Fe, Al, Ni, Cr, Mn) were measured by flame atomic absorption spectrometry (AAS Shimadzu 6800) using deuterium background correction. Cd and Pb stable isotope ratios were measured by ICP-MS (Perkin-Elmer Sciex ELAN 9000). Measurement procedures are described in detail in Zaborska et al. (2017). Metal enrichment factors (EF) were calculated according to the formula detailed in Zaborska et al. (2017). Mean background metal concentrations were taken from literature for the Hornsund region. The fraction of anthropogenic Pb (Pb_{ant}) was calculated according to the formula detailed in Zaborska et al. (2017). The excess $^{206}\text{Pb}/^{207}\text{Pb}$ ($^{206}\text{Pb}/^{207}\text{Pb}_{\text{excess}}$) ratio calculation was used to source Pb pollution (Farmer et al., 1996; Bindler et al., 2001). ~~Selected heavy metals that are believed to be important anthropogenic contaminants (Pb, Cu, Zn, Fe, Al) were measured with a flame atomic absorption spectrometer (AAS Shimadzu 6800) using deuterium background correction. Cd and the Pb stable isotopic ratios were measured on a Perkin Elmer Sciex ELAN 9000 ICP MS. The exact measurement procedure and quality assurance is described in detail in Zaborska et al. (2017). The metal enrichment factors (EF) were calculated according to the formula presented in Zaborska et al. (2017). The mean background metal concentrations were taken from literature for the region of Hornsund. The fraction of anthropogenic Pb (Pb_{ant}) was calculated according to formula presented in Zaborska et al. (2017). The excess $^{206}\text{Pb}/^{207}\text{Pb}$ ($^{206}\text{Pb}/^{207}\text{Pb}_{\text{excess}}$) ratio calculation was used to find a source of Pb pollution (Farmer et al., 1996; Bindler et al., 2001).~~

25 *Statistical analyses*

The regression analysis was performed to find correlations between particular radionuclide and metal concentrations. The correlations characterized by $r > 0.5$ are assumed to show moderate correlation, the $r > 0.7$ is according to statistics indicating strong correlation (eg. Mukaka, 2012). ~~The correlations characterized by $r > 0.5$ are assumed to show moderate correlation, the $r > 0.7$ is assumed to show strong correlation.~~ Nonparametric Kruskal-Wallis test was used to find a difference between radionuclide and metal concentrations in soil and cryoconite samples. All statistical calculations were performed using STATISTICA 10.0 licensed program (StatSoft, Inc. Team. 2011).

3 Results

3.1 Radionuclide contents and their activity ratios in tundra soil profiles and in cryoconite samples

Activity concentrations and inventories of anthropogenic radionuclides (^{137}Cs , ^{238}Pu , $^{239+240}\text{Pu}$, ^{241}Am) for tundra soil profiles and for cryoconite are presented in Tab. S2, ~~and in~~ Fig. 2A and Tab. S4, Fig. 3. Radionuclide inventory is ~~understood~~ defined here as the activity of a given radionuclide contained in the soil column at unit surface area. Most of the profiles have concordant depth distributions of all artificial radionuclide activity with distinct peaks occurring in the surface or the first sub-surface layers. The highest activity concentrations of anthropogenic radionuclides are observed in profile S04 at 1 cm depth where significant amounts of organic matter were observed (21 % LOI). Profile S02 shows no detectable ^{137}Cs with a minimal detectable activity (MDC) range of 0.1 to 3 Bq kg⁻¹. Activities of other radionuclides in this profile were not determined. The determined activity concentrations of ^{137}Cs , ^{238}Pu , $^{239+240}\text{Pu}$ and ^{241}Am varied between 7±2 to 65±7 Bq/kg, 0.03±0.01 to 0.09±0.01 Bq/kg, 0.08±0.01 to 2.13±0.16 Bq/kg and 0.09±0.01 to 0.90±0.06 Bq/kg, respectively. In most of the samples the activity concentrations of ^{238}Pu are close to the detection limit, ~~which this is the reason is~~ why the activity ratios of $^{238}\text{Pu}/^{239+240}\text{Pu}$ were calculated for only 3 samples. Activity concentrations for cryoconite range from: 13±3 to 2000±300 Bq kg⁻¹ for ^{137}Cs , 0.08±0.02 to 2.1±0.2 Bq kg⁻¹ for ^{238}Pu , 0.09±0.02 to 43±3 Bq kg⁻¹ for $^{239+240}\text{Pu}$ and 0.25±0.06 to 25±2 Bq kg⁻¹ for ^{241}Am . All of these values are significantly higher than those observed in soil samples collected near this glacier.

~~The activity ratios of $^{238}\text{Pu}/^{239+240}\text{Pu}$, $^{239+240}\text{Pu}/^{137}\text{Cs}$ and $^{241}\text{Am}/^{239+240}\text{Pu}$, as well as atomic ratios of $^{240}\text{Pu}/^{239}\text{Pu}$, were calculated to potentially distinguish the sources of these radionuclides in tundra soils and cryoconite samples. These The mean $^{238}\text{Pu}/^{239+240}\text{Pu}$ activity ratios for tundra varied between 0.029±0.010 and 0.05±0.02 with the mean value of is 0.039±0.010 and it is lower than for cryoconite 0.050±0.009. The analysis of atomic ratios of $^{240}\text{Pu}/^{239}\text{Pu}$ provides important information which allows a more precise identification of the origin of Pu isotopes in environmental samples. The average $^{240}\text{Pu}/^{239}\text{Pu}$ atomic ratios found in tundra profiles (0.179) and in cryoconite (0.147) varied from 0.168±0.001 to 0.191±0.005 with an average of 0.179±0.006 (Lokas et al., 2017a) are presented in Fig. 6. The activity ratios of $^{238}\text{Pu}/^{239+240}\text{Pu}$, $^{239+240}\text{Pu}/^{137}\text{Cs}$ and $^{241}\text{Am}/^{239+240}\text{Pu}$, as well as atomic ratios of $^{240}\text{Pu}/^{239}\text{Pu}$, were calculated to potentially distinguish the sources of these radionuclides in cryoconite samples and are presented in Table S4. The $^{238}\text{Pu}/^{239+240}\text{Pu}$ activity ratios varied between 0.031±0.006 (sample 11) and 0.062±0.008 (samples 4 and 12). The mean value for activity ratios of $^{238}\text{Pu}/^{239+240}\text{Pu}$ is 0.050±0.009. The analysis of atomic ratios of $^{240}\text{Pu}/^{239}\text{Pu}$ provides important information which allows a more precise identification of the origin of Pu isotopes in environmental samples. $^{240}\text{Pu}/^{239}\text{Pu}$ atomic ratios found in cryoconite samples varied from 0.121±0.039 (sample 9) to 0.196±0.019 (sample 5) with an average of 0.147±0.009. The mean $^{239+240}\text{Pu}/^{137}\text{Cs}$ activity ratios in tundra (0.042±0.010) is higher than in the cryoconite samples range between 0.007±0.003 to 0.038±0.008 with the mean value of (0.020±0.004). But The mean $^{241}\text{Am}/^{239+240}\text{Pu}$ activity ratios for tundra and cryoconite are comparable (0.46±0.07 and 0.47±0.07) range between 0.24±0.05 and 0.65±0.12 with the mean value of 0.47±0.07.~~




~~The $^{239+240}\text{Pu}/^{137}\text{Cs}$ activity ratios range between 0.025±0.003 to 0.059±0.009 with the mean value of 0.042±0.010. The $^{241}\text{Am}/^{239+240}\text{Pu}$ activity ratios range between 0.28±0.04 and 0.55±0.11 with the mean value of 0.46±0.07. Artificial~~

radionuclide inventories evaluated for soils vary between 150 ± 40 to 670 ± 100 Bq m⁻² for ¹³⁷Cs, from 0.44 ± 0.15 to 0.96 ± 0.02 Bq m⁻² for ²³⁸Pu, from 6.8 ± 0.6 to 28 ± 2 Bq m⁻² for ²³⁹⁺²⁴⁰Pu and from 1.9 ± 0.2 to 14 ± 3 Bq m⁻² for ²⁴¹Am. The average values of the radionuclide inventories calculated for all of these profiles are: 0.66 ± 0.14 Bq m⁻² for ²³⁸Pu, 20 ± 2 Bq m⁻² for ²³⁹⁺²⁴⁰Pu, 9 ± 1 Bq m⁻² for ²⁴¹Am and 450 ± 70 Bq m⁻² for ¹³⁷Cs.

5 The measurement results for natural radioisotopes (²¹⁰Pb, ^{234,238}U, ^{230,232}Th) for tundra soil profiles and for cryoconite are presented in Figure 2A, ~~and B, Tab. S3 and Tab. S5.~~ Activity concentrations range from 15 ± 1 to 173 ± 4 Bq kg⁻¹ for total ²¹⁰Pb, 13 ± 1 to 28 ± 2 Bq kg⁻¹ for ²³⁴U, 13 ± 1 to 30 ± 2 Bq kg⁻¹ for ²³⁸U, 9 ± 1 to 22 ± 2 Bq kg⁻¹ for ²³⁰Th and 8 ± 1 to 42 ± 3 Bq kg⁻¹ for ²³²Th. The highest activity concentration for ²¹⁰Pb was found in the surface layer of profile S04 and this value is much lower than for cryoconite (12300 ± 600 Bq kg⁻¹). Activity ratios of ²³⁴U/²³⁸U for soils and cryoconite are very comparable, varied between 0.9 ± 0.1 to 1.2 ± 0.2 (Tab. S3). ~~Data on natural radioisotopes (²¹⁰Pb, ^{234,238}U, ^{230,232}Th) for the cryoconite samples are presented in Table S3 (Supplementary material). Activity concentrations range from 490 ± 60 to 12300 ± 600 Bq kg⁻¹ for total ²¹⁰Pb, 12 ± 1 to 43 ± 4 Bq kg⁻¹ for ²³⁴U, 10 ± 1 to 37 ± 4 Bq kg⁻¹ for ²³⁸U, 16 ± 1 to 54 ± 4 Bq kg⁻¹ for ²³⁰Th and 16 ± 1 to 58 ± 4 Bq kg⁻¹ for ²³²Th. The highest activity concentration for ²¹⁰Pb was observed in sample 4. Activity ratio of ²³⁴U/²³⁸U varied between 0.9 ± 0.1 to 1.4 ± 0.3 (Tab. S5).~~

10

15






Figure 2. Schematic of Waldemarbreen with activity concentrations of atmospheric radionuclides (^{210}Pb , ^{137}Cs , $^{239,240}\text{Pu}$, ^{241}Am) in cryoconite and depth distribution in tundra soils.


Figure 2. The model of Waldemarbreen and depth distribution of atmospheric radionuclides (^{210}Pb , ^{137}Cs , $^{239,240}\text{Pu}$, ^{241}Am)

5 ~~(A), depth distribution of natural radionuclides ($^{234,238}\text{U}$, $^{230,232}\text{Th}$) (B) and depth distribution of selected heavy metals (Cu, Pb, Cd, Zn) (C) in tundra soils.~~

3.2 Radionuclide contents and their activity ratios in cryoconite samples

The results of activity concentrations of anthropogenic radionuclides analysis (^{137}Cs , ^{238}Pu , $^{239+240}\text{Pu}$ and ^{241}Am) for all the cryoconite samples are presented in Table S4 and Fig. 3. Activity concentrations range from: 13 ± 3 to $2000\pm 300\text{ Bq kg}^{-1}$ for ^{137}Cs , 0.08 ± 0.02 to $2.1\pm 0.2\text{ Bq kg}^{-1}$ for ^{238}Pu , 0.09 ± 0.02 to $43\pm 3\text{ Bq kg}^{-1}$ for $^{239+240}\text{Pu}$ and 0.25 ± 0.06 to $25\pm 2\text{ Bq kg}^{-1}$ for ^{241}Am . All of these values are significantly higher than those observed in soil samples collected near this glacier.

5






Figure 3. Concentrations of selected heavy metals (Pb, Cd) in cryoconite samples and tundra soil profiles.

Figure 3. Activity concentrations of anthropogenic (^{137}Cs , $^{239,240}\text{Pu}$, ^{241}Am) and natural ($^{234,238}\text{U}$, $^{230,232}\text{Th}$, ^{210}Pb) radionuclides in cryoconite samples from Waldemarbreen.

The activity ratios of $^{238}\text{Pu}/^{239+240}\text{Pu}$, $^{239+240}\text{Pu}/^{137}\text{Cs}$ and $^{241}\text{Am}/^{239+240}\text{Pu}$, as well as atomic ratios of $^{240}\text{Pu}/^{239}\text{Pu}$, were calculated to potentially distinguish the sources of these radionuclides in cryoconite samples and are presented in Table S4. The $^{238}\text{Pu}/^{239+240}\text{Pu}$ activity ratios varied between 0.031 ± 0.006 (sample 11) and 0.062 ± 0.008 (samples 4 and 12). The mean value for activity ratios of $^{238}\text{Pu}/^{239+240}\text{Pu}$ is 0.050 ± 0.009 . The analysis of atomic ratios of $^{240}\text{Pu}/^{239}\text{Pu}$ provides important information which allows a more precise identification of the origin of Pu isotopes in environmental samples. $^{240}\text{Pu}/^{239}\text{Pu}$ atomic ratios found in cryoconite samples varied from 0.121 ± 0.039 (sample 9) to 0.196 ± 0.019 (sample 5) with an average of 0.147 ± 0.009 . The $^{239+240}\text{Pu}/^{137}\text{Cs}$ activity ratios in the cryoconite samples range between 0.007 ± 0.003 to 0.038 ± 0.008 with the mean value of 0.020 ± 0.004 . The $^{241}\text{Am}/^{239+240}\text{Pu}$ activity ratios range between 0.24 ± 0.05 and 0.65 ± 0.12 with the mean value of 0.47 ± 0.07 . Data on natural radioisotopes (^{210}Pb , $^{234,238}\text{U}$, $^{230,232}\text{Th}$) for the cryoconite samples are presented in Table S3 (Supplementary material). Activity concentrations range from 490 ± 60 to 12300 ± 600 Bq kg^{-1} for total ^{210}Pb , 12 ± 1 to 43 ± 4 Bq kg^{-1} for ^{234}U , 10 ± 1 to 37 ± 4 Bq kg^{-1} for ^{238}U , 16 ± 1 to 54 ± 4 Bq kg^{-1} for ^{230}Th and 16 ± 1 to 58 ± 4 Bq kg^{-1} for ^{232}Th . The highest activity concentration for ^{210}Pb was observed in sample 4. Activity ratio of $^{234}\text{U}/^{238}\text{U}$ varied between 0.9 ± 0.1 to 1.4 ± 0.3 (Tab. S5).

3.3.2 Heavy metals analyses in tundra soil profiles and in cryoconite samples

The concentrations of all measured metals and ratios of stable Pb isotopes ($^{206}\text{Pb}/^{207}\text{Pb}$, $^{208}\text{Pb}/^{206}\text{Pb}$) in soil and cryoconite samples are presented in Tab. S6. and S8, respectively. Cd, Pb, Cu and Zn concentrations in soils and cryoconite are shown in Figure 2C, and 4. Pb concentrations varied from 9.6 $\text{mg}\cdot\text{kg}^{-1}$ ~~(sample S05-3)~~ to 25.9 $\text{mg}\cdot\text{kg}^{-1}$ ~~(sample S04-1)~~. Cd concentrations range from 0.05 $\text{mg}\cdot\text{kg}^{-1}$ ~~(sample S01-2)~~ to 0.6 $\text{mg}\cdot\text{kg}^{-1}$ ~~(sample S01-4)~~. The concentrations of Zn ranged from 45.1 $\text{mg}\cdot\text{kg}^{-1}$ ~~(sample S03-1)~~ to 76.9 $\text{mg}\cdot\text{kg}^{-1}$ ~~(sample S04-1)~~. Cu concentrations varied from 11.8 $\text{mg}\cdot\text{kg}^{-1}$ ~~(sample S03-1)~~ to 26.6 $\text{mg}\cdot\text{kg}^{-1}$ ~~(sample S01-2)~~. The results for cryoconite were significantly higher than for soils. Pb concentrations ranged from 19.9 $\text{mg}\cdot\text{kg}^{-1}$ (sample 13) to 97.7 $\text{mg}\cdot\text{kg}^{-1}$ (sample 12). Cd concentrations varied from 0.2 $\text{mg}\cdot\text{kg}^{-1}$ (sample 10) to 0.6 $\text{mg}\cdot\text{kg}^{-1}$ (sample 11). Zn concentrations ranged from 59.6 $\text{mg}\cdot\text{kg}^{-1}$ (sample 7) to 97.5 $\text{mg}\cdot\text{kg}^{-1}$ (sample 11). Cu concentrations varied from 21.5 $\text{mg}\cdot\text{kg}^{-1}$ (sample 13) to 40.1 $\text{mg}\cdot\text{kg}^{-1}$ (sample 9). The stable isotope ratio of $^{206}\text{Pb}/^{207}\text{Pb}$ for soils ranged from 1.190 to 1.217 while $^{208}\text{Pb}/^{206}\text{Pb}$ ranged from 2.014 to 2.057 and for cryoconite are very similar, ranged from 1.169 to 1.199 while $^{208}\text{Pb}/^{206}\text{Pb}$ ranged from 2.025 to 2.061. There was a significant correlation between Pb and Cu and Zn ($r=0.82$, $r=0.71$) in soils (Tab. S7) and cryoconite ($r=0.78$) (Tab. S9). Pb also correlated with naturally derived metals Ni, ~~Cr~~ and Co ($r=0.48-0.59$) in soils and in cryoconite Ni ($r=0.67$). Cu correlated with Zn ($r=0.77$) only in soils. Zn and Cu also correlated with Ni, Cr and Co. Interestingly, Cd did not correlate with any of these metals both in soils and cryoconite. The correlations between contaminating heavy metals and soil composition markers Fe and Al were checked to test the influence of natural variability of soil composition to metal concentration. Generally, there was no significant correlation of contaminating metals with Fe and Al, while naturally derived Cr, Ni and Co correlated with Fe and Al, both in soils and cryoconite. The correlations between contaminating heavy metals and Fe and Al were checked to avoid the influence of natural variability of crust composition to metal concentration. Pb, Zn and Cu correlated with both metals, particularly strong was the correlation of Cu and Zn (r ranging from 0.73 to 0.84). The correlation of Pb with Fe and Al was much lower (r ranging from 0.53 to 0.59). $^{206}\text{Pb}/^{207}\text{Pb}$ inversely correlates with Pb ($r=0.78$), lower but significant correlations were measured for Cu, Zn and Cr. $^{208}\text{Pb}/^{206}\text{Pb}$ did not correlate with any metal concentrations. The metal enrichment factors (normalized to Al) are generally comparable for soils and cryoconite samples and ranged from 1.1 to 1.8 for Pb, from 0.9 to 1.6 for Zn, from 0.8 to 2.1 for Cu and from 0.4 to 7.7 for Cd (Tab. S6) and from 1.5 to 6.1 for Pb, from 1.0 to 1.5 for Zn, from 1.3 to 2.3 for Cu and from 0.5 to 4.5 for Cd (Tab. S8), respectively. For most metals there were significant differences in concentration observed in the soil versus the cryoconite samples (Tab. S10). The only exception is Cd which was similar (and elevated) in both environments. There was no difference in $^{208}\text{Pb}/^{206}\text{Pb}$ ratio between soil and cryoconite samples but this is not very surprising since this ratio is less specific than $^{206}\text{Pb}/^{207}\text{Pb}$ (the $^{208}\text{Pb}/^{206}\text{Pb}$ coal burning signature has a similar ratio to a natural rock).

3.4 Heavy metals analyses in cryoconite samples

~~Table S8 presents concentrations of all measured metals and ratios of stable Pb isotopes ($^{206}\text{Pb}/^{207}\text{Pb}$, $^{208}\text{Pb}/^{206}\text{Pb}$) in cryoconite samples. Metals that are believed to be the most frequent contaminants (Cd, Pb, Zn, Cu) are also shown in Fig. 4.~~

~~esa~~ Pb concentrations ranged from 19.9 mg kg⁻¹ (sample 13) to 97.7 mg kg⁻¹ (sample 12). Cd concentrations varied from 0.2 mg kg⁻¹ (sample 10) to 0.6 mg kg⁻¹ (sample 11). Zn concentrations ranged from 59.6 mg kg⁻¹ (sample 7) to 97.5 mg kg⁻¹ (sample 11). Cu concentrations varied from 21.5 mg kg⁻¹ (sample 13) to 40.1 mg kg⁻¹ (sample 9). The ratio of ²⁰⁶Pb/²⁰⁷Pb ranged from 1.169 to 1.199 while ²⁰⁸Pb/²⁰⁶Pb ranged from 2.025 to 2.061. There was a significant correlation between Pb and Cu (r=0.78) (Tab. S9). Pb also correlated with naturally derived metals Ni and Mn. Zn and Cu correlated with Cr and Ni. Interestingly, like in the soil samples, Cd did not correlate with any of the metals. The correlations between contaminating heavy metals and soil composition markers Fe and Al were checked to test the influence of natural variability of soil composition to metal concentration. There was no significant correlation of contaminating metals with Fe and Al, while naturally derived Cr, Ni and Co correlated with Fe and Al. ²⁰⁶Pb/²⁰⁷Pb inversely correlated with Pb (r=0.86), lower but significant correlations were measured for Cu, Ni and Mn. ²⁰⁸Pb/²⁰⁶Pb correlated with only Pb and Cu (r= 0.58 and r= 0.60 respectively). The metal enrichment factors (normalized to Fe) for cryoconite samples ranged from 1.5 to 6.1 for Pb, from 1.0 to 1.5 for Zn, from 1.3 to 2.3 for Cu and from 0.5 to 4.5 for Cd (Tab. S8).




Figure 4. Concentrations of all measured metals (Cu, Pb, Cd, Zn) in cryoconite samples.

For most metals there were significant differences in concentration observed in the soil versus the cryoconite samples (Tab. S10). The only exception is Cd which was similar (and elevated) in both environments. There was no difference in ²⁰⁸Pb/²⁰⁶Pb ratio between soil and cryoconite samples but this is not very surprising since this ratio is less specific than ²⁰⁶Pb/²⁰⁷Pb (the ²⁰⁸Pb/²⁰⁶Pb coal burning signature has a similar ratio to a natural rock).

4 Discussion

4.1 Radionuclides 4.1. Radionuclides and heavy metals contamination

All soil profiles show typical activity concentrations and inventories as do other profiles found on the raised marine terraces covered by dry tundra in the western part of Spitsbergen (Łokas et al., 2014; 2017a,b; AMAP, 2015; Dowdall et al., 2003; Gwynn et al., 2004; Hardy et al., 1973). Retention of airborne radionuclides in tundra soils is facilitated by the presence of the relatively well-developed organic horizon. ~~The average values of the radionuclide inventories fall within the ranges of the above-mentioned regional evaluations of fallout deposition (Hardy et al., 1973; Holm et al., 1983; AMAP, 2015; Łokas et al., 2017a,b).~~

Cryoconite materials are highly enriched with radionuclides of atmospheric origin compared with tundra soils. Radionuclide enrichment exceeding the soil levels can be explained to some extent by the focusing of fine-grained material, washed down into very small depressions where the cryoconite samples were collected, and also by the properties of cryoconite material.

Cryoconite granules are the mixtures of mineral particles, organic matter and microorganisms. They can retain and concentrate airborne radionuclides due to metal-binding properties of extracellular substances that are excreted by microorganisms (Gadd 2004; Francis 2007). Differences in activity concentrations of anthropogenic radionuclides (^{137}Cs , Pu isotopes and ^{241}Am) found among the cryoconite samples reflect the different locations of sampling sites, morphological features and contents of organic material. A significant correlation was observed between the amount of organic matter

present and the concentration of all airborne radionuclides ($r=0.79-0.83$). We noticed significant correlation between ^{210}Pb , ^{137}Cs , Pu isotopes and ^{241}Am with the altitude ($r=0.64-0.91$) and the area of the cryoconite holes ($r=0.58-0.63$), except ^{210}Pb ($r=0.38$). The depth of these cryoconite holes also correlates with ^{210}Pb and Pu isotopes ($r=0.60-0.68$) (Tab. S8). Cryoconite samples collected at the highest location (4, 12, 13) with the highest surface area of cryoconite holes (Tab. S1) were characterised by the highest observed activity concentrations. These three points were located very close to each other and the deposited material could have originated from the same source and migrated with supraglacial streams. In contrast, sampling points 5, 6, 7 which were located close to the moraine at lower altitude, were characterised by the lowest activity concentrations of anthropogenic radionuclides. These materials could have been washed out, or mixed with mineral particles from the moraine, or collapsed cryoconite holes (that may be destroyed more easily here than in the middle of the glacier).

The highest activity concentrations of ^{210}Pb was observed in sample 4 (more than 12000 Bq kg^{-1}). More uniform and comparable values (about 5000 Bq kg^{-1}) were found for cryoconite samples 3, 8, 11, 12, 13. This ^{210}Pb distribution is very similar to the distribution of anthropogenic radionuclides. Cryoconite granules can be locally redistributed on the surface of ice, dependent on the glacier morphology and altitude. It was noticed that activity concentrations increase in relation to the altitude above sea level. Material deposited in the highest locations can migrate with supraglacial streams and can be transported along with the glacier. Samples 4 and 13 were collected from typical cryoconite holes with the highest surface area (476 and 1200 cm^2) (Tab. S1). Regardless of the lifespan of individual cryoconite holes, their collapse does not imply removal of cryoconite from glacier surface as the dispersed cryoconite granules initiate formation of new holes (Takeuchi et al., 2001).

The constant delivery of ^{210}Pb from the atmosphere suggests that the concentration of this radionuclide in cryoconite material should be proportional to the exposure time, while high concentrations of the artificial radionuclides indicate significant

contribution of material that was exposed to the stratospheric or tropospheric fallout. The results show that ^{210}Pb is being deposited at a more constant rate and lower values for samples 3, 8, 11, 12, 13 may suggest that the majority of material in these samples was already removed with melt waters or that material in these samples is younger than in sample 4 (shorter contact time with the atmosphere).

5 The effect of glacial morphology on effective trapping and storing of radionuclides was also observed in Georgia glacier (Łokas et al., 2018). The differences in the concentrations of radionuclides between sampling points and the lack of clear differences in the elevation gradient from terminus towards icefall may reflect the heterogeneous topography of the glacier tongue. In the Waldemarbreen we observed differences in the elevation gradient from the top of ice.

Activity concentrations of $^{234,238}\text{U}$ and $^{230,232}\text{Th}$ show little variability within the profiles and do not differ from values reported for soils globally (UNSCEAR, 1993), in the tundra soils in northern Spitsbergen (Dowdall et al., 2003) and in south-western Spitsbergen (Łokas et al., 2017b). Soil activity concentrations of these lithogenic radionuclides are related to their contents in the source minerals (Megumi et al., 1988) and are not influenced by the action of cryoconite. The presence of uranium and thorium are the main elements contributing to natural terrestrial radioactivity. Uranium isotopes (^{234}U and ^{238}U) in terrestrial samples (rocks, soils and sediments) are usually present in radioactive equilibrium. The main source of uranium in the natural environment is the atmospheric precipitation of terrigenous material, soil resuspension and rock weathering.

Our data for uranium and thorium analysis in cryoconite samples are comparable to other results for these materials. Our previous paper (Łokas et al., 2018) showed similar results for the concentration of ^{234}U and ^{238}U (mean value 33 ± 3 and 34 ± 3 Bq kg $^{-1}$, respectively) uranium and thorium in cryoconite granules from the Georgia glacier, while on the Swiss glacier (Baccolo et al., 2017), ^{238}U they and ^{230}Th reach around 118 Bq kg $^{-1}$ (are higher with the mean value for ^{238}U 65 ± 5 Bq kg $^{-1}$). The value of the $^{234}\text{U}/^{238}\text{U}$ activity ratio in analysed cryoconite granules suggests a state of radioactive equilibrium. The activity concentrations of ^{232}Th are also in agreement with cryoconite samples from the Georgia glacier (mean value 42 ± 4 Bq kg $^{-1}$) but higher mean values (65 ± 3 Bq kg $^{-1}$) were observed in Swiss glaciers. The mean value for ^{230}Th in analysed cryoconite samples is lower than for the Georgia glacier (51 ± 4 Bq kg $^{-1}$). The activity ratio between thorium isotopes ($^{230}\text{Th}/^{232}\text{Th}$) in cryoconite does not show a constant value – it varies from 0.4 to 0.7. In only a few samples the activity ratio of $^{230}\text{Th}/^{232}\text{Th}$ is about 1 (samples 2, 3, 4, 12, 13). Isotopes derived from uranium (^{230}Th) and thorium series (^{232}Th) are not correlated in cryoconite samples ($R^2=0.1336$) but in tundra soil profiles are better correlated ($R^2=0.6280$). This may suggest that thorium from the uranium series is removed with melt water from the cryoconite samples located at the top of the glacier. We also observed differences between activity concentrations of uranium and thorium isotopes in analysed tundra soils and cryoconite samples. In tundra soils the values are lower than in cryoconite.

The metal concentrations similarly as radionuclides in soils were typical or slightly higher than concentrations characteristic for natural background concentrations. This had been confirmed by generally low metal enrichment ratios (EF) with the exception of Cd. The metal distributions measured in this study are very similar to metal concentrations characteristic for parent rocks (natural crust) on Spitsbergen. For instance, parent rocks in Hornsund contain very low

concentrations of heavy metals (Pb~11 mg·kg⁻¹, Cd~0.04 mg·kg⁻¹, Zn~58 mg·kg⁻¹, Cu~20 mg·kg⁻¹; Samecka-Cymerman et al., 2011). Visible enrichment in Cd and Pb was found in surface and subsurface soil profile layers (Fig. 5). Similar enrichment in metal concentrations (Pb up to 25.3 mg·kg⁻¹, Cd up to 0.4 mg·kg⁻¹) in soil profiles of Kaffiøyra was found in the 1980's by Plichta (1993). Generally low metal concentrations have been mapped by Ottesen et al. (2010) in alluvial soils in Kaffiøyra region. Alluvial sediments are believed to represent natural metal levels, although recently deposited layers can be polluted by anthropogenic activity as Cd concentrations (ranged from 0.01 to 0.13 mg·kg⁻¹), Cu (13.4 to 43.4 mg·kg⁻¹), Pb (8.9 to 24.3 mg·kg⁻¹) and Zn (58.9 to 79.2 mg·kg⁻¹). Metal concentrations were studied in peat cores collected from nearby Kongsfjorden. The Cu ranged from 7 to 25 mg·kg⁻¹, Pb from 10 to 35 mg·kg⁻¹ while Zn ranged from 20 to 75 mg·kg⁻¹ (Headley, 1996). The region of Kongsfjorden may be have been impacted by former mining activity, however (Hao et al., 2013). More recently, the metal distribution in soils of other Spitsbergen regions (Kongsfjorden and Adventfjorden) was studied by Halbach et al. (2017). Like our study, they found elevated concentrations of some metals eg. Cd (0.13-1.00 mg·kg⁻¹) in surface soils (covered by moss), but low concentrations of metals (Cd=0.04-0.5, Cu=3.3-28.4 mg·kg⁻¹, Pb=2.9-22.7 mg·kg⁻¹ and Zn=25-106 mg·kg⁻¹) in sandy fraction of soils (sampled at 20cm depth). Slightly higher levels of heavy metals (Cu: 17-92 mg·kg⁻¹ and Cd: 0.05-1.20 mg·kg⁻¹) were measured in biologically rich soils in Hornsund region with the highest values in areas of rich vegetation (Wojtuń et al., 2013). Ziólek et al. (2017) reports much higher metal concentrations in peat soils in Bellsund. In their study Pb reached 100 mg·kg⁻¹, Zn reached 140 mg·kg⁻¹, Cu 57 reached mg·kg⁻¹ while Cd reached 8 mg·kg⁻¹. The high contamination of peat soils was in their case caused by the richness of organic matter due to a bird colony. In conclusion, the effect of anthropogenic influence on soils in Kaffiøyra region was very limited.

While soils can receive metals from atmosphere and from the surface run-off, cryoconites are fed only by atmospheric particles. Thus, interestingly, metal concentrations in cryoconite were higher than in soil samples (Tab. S8) and definitely exceeded natural values. Particularly elevated was the concentration of Pb reaching 97.5 mg·kg⁻¹ (EF = 6.1) and Cd reaching 0.6 mg·kg⁻¹ (EF=4.5). The enrichment in other metals was lower (Zn up to 97.5 mg·kg⁻¹ and Cu up to 40.1 mg·kg⁻¹). The maximum concentration of Pb in cryoconite samples from the Hansbreen glacier in Hornsund was lower (82.7 mg·kg⁻¹), the concentrations of Zn (108 mg·kg⁻¹) and Cd (1.53 mg·kg⁻¹) were higher, however (Łokas et al., 2016). Singh et al. (2013) found a similar range of metal concentrations in cryoconite samples from a glacier located also in Kongsfjorden area (Pb up to 85.1 mg·kg⁻¹, Cu up to 44 mg·kg⁻¹, Cd up to 0.14 mg·kg⁻¹, Zn up to 150 mg·kg⁻¹). Chmiel et al. (2009) did not find elevated metal levels in glacier waters in the Scott Glacier in Bellsund, they found enrichment in some metals (Pb = 190 mg·kg⁻¹) in cryogenic sediments, however.

4.2 The source of radionuclides and heavy metals contamination

Activity ratios of ²³⁸Pu/²³⁹⁺²⁴⁰Pu, ²³⁹⁺²⁴⁰Pu/¹³⁷Cs, ²⁴¹Am/²³⁹⁺²⁴⁰Pu and ²⁴⁰Pu/²³⁹Pu atom ratio are commonly used to identify and distinguish between global (stratospheric) and regional (tropospheric) sources of these radionuclides (Oughton et al., 2004; Hirose and Povinec, 2015; Łokas et al., 2013, 2014, 2016 and 2017b).

The $^{238}\text{Pu}/^{239+240}\text{Pu}$ activity ratios in cryoconite samples are consistently higher than for tundra soils, suggesting the presence of contributing sources other than the global Pu fallout in this region. The percentage of global fallout versus local sources was estimated and the model assumes that the $^{238}\text{Pu}/^{239+240}\text{Pu}$ activity ratios for global fallout and Chernobyl plutonium are about 0.025 and 0.24, respectively. Following formula described by Mietelski and Wąs (1997) was used:

$$f_G = (A_R - A_M)/(A_R - A_G) \quad (4)$$

where f_G represents the fraction of plutonium isotopes from global fallout sources (%), A_R represents the $^{238}\text{Pu}/^{239+240}\text{Pu}$ ratio of Chernobyl sources (0.24), A_M represents the measured $^{238}\text{Pu}/^{239+240}\text{Pu}$ activity ratio in cryoconite samples and A_G represents the $^{238}\text{Pu}/^{239+240}\text{Pu}$ activity ratio of global fallout (0.025).

Calculations were performed for each of the studied cryoconite samples, and the fraction of global fallout varied from 82 ± 1 % to 97 ± 1 % (Fig. 5). Samples 4, 8 and 12, with the highest activity ratios of $^{238}\text{Pu}/^{239+240}\text{Pu}$ (0.062), contained also the highest contribution of Pu from other sources (18 % and 17 %, respectively). The highest activity ratios of $^{238}\text{Pu}/^{239+240}\text{Pu}$ (0.075 to 0.09) were observed in the initial soils from the vicinity of the Werenskiöld glacier terminus in south-west part of Spitsbergen. These values fall within the range found also for cryoconite from the Hans glacier (between 0.037 and 0.118 with a mean 0.064) and other Arctic glaciers; (Łokas et al., 2016; preliminary our data). The $^{240}\text{Pu}/^{239}\text{Pu}$ atomic ratios in analysed tundra soils (average of 0.179) (Fig. 6A) are comparable to global fallout ratios (0.183 ± 0.009) reported by Efurð et al. (2005) at 70° N latitude and 0.180 reported by others (Muramatsu et al., 1999; Warneke et al., 2002). Similar results was also reported for tundra soils from Spitsbergen (80 samples) with an average value of 0.179 (Łokas et al., 2017a). The tundra sites exposed to the atmosphere over the whole period of anthropogenic radionuclide release acquired their radionuclide contents primarily from direct atmospheric deposition.





Figure 45. The contribution of global fallout Pu calculated using plutonium isotope ratios (left) contribution of anthropogenic origin Pb calculated using Pb isotopes and end-member method.

In tundra soils, $^{240}\text{Pu}/^{239}\text{Pu}$ atom ratios and the $^{238}\text{Pu}/^{239+240}\text{Pu}$ activity ratios are similar to the global fallout signature (0.180 and 0.025, respectively), however, in the cryoconite samples measured in this study (and other unpublished data from Arctic cryoconite) they are lower (0.147 and 0.050, respectively) (see Fig. 5A, B). Due to the slow accumulation of tundra soils and their continuous exposure to the atmosphere, the vertical profiles of the fallout radionuclides represent their time-averaged deposition on soil surface. In contrast, different cryoconite samples from the same glacier could be exposed to fallout for various periods resulting in preferential uptake of radionuclides from single deposition events and deviations from the global fallout signatures. In a previous article (Lokas et al., 2017a) we assumed that the main source of Pu in the proglacial soils (and now also in cryoconite samples) were nuclear tests and a non-exploded weapons-grade material. The lack of correlation between excess ^{238}Pu and $^{240}\text{Pu}/^{239}\text{Pu}$ atom ratios (Figure 5C) suggests that the excessive ^{238}Pu is derived from sources other than ^{239}Pu , which is pure ^{238}Pu from SNAP9A satellite re-entry (Corcho Alvarado et al., 2014). A similar ^{238}Pu enrichment was observed in other cryoconite samples from Arctic glaciers (unpublished data) (Fig. 5C) and in air filters in Finland (Salminen-Paatero et al., 2012) and NW Poland (Kierepko et al., 2016).

~~In tundra soils, $^{240}\text{Pu}/^{239}\text{Pu}$ atomic ratios and the $^{238}\text{Pu}/^{239+240}\text{Pu}$ activity ratios are similar to global fallout signature (0.180 and 0.025, respectively) but in cryoconite samples they are different (Fig. 6B). Tundra soils develop very slowly; therefore, the $^{240}\text{Pu}/^{239}\text{Pu}$ atomic ratios are more uniform and homogeneous than in cryoconite samples. Soil profile have yearly stratification and averaged Pu isotopic ratios, resulting in small variation of the measured values. In contrast, in cryoconite samples, each such “snapshot” contains Pu, which was previously in a certain site in glacial and came from fallout after a specific deposition event. Similar mechanism was observed in initial soils from the proglacial zones of glacier from Spitsbergen (Lokas et al., 2017a). In the analysed cryoconite samples an average $^{240}\text{Pu}/^{239}\text{Pu}$ atomic ratio is 0.147 (Fig. 6A) and this value is lower than for initial soils and also lower than in cryoconite from the Georgia glacier (0.163) (Lokas et al., 2018). The result for analysed cryoconite is very similar to initial soils and we checked the correlation between excess ^{238}Pu and $^{240}\text{Pu}/^{239}\text{Pu}$ atomic ratios (Figure 6C) and the lack of this indicates that enrichment of ^{238}Pu is derived from sources other than ^{239}Pu . We assumed in a previous article that the main source of Pu in the proglacial soils and now also in cryoconite samples are nuclear tests and a non-exploded weapons grade material. A third potential source of Pu is pure ^{238}Pu from a satellite re-entry after injection from the stratosphere into the troposphere.~~






Figure 5. The relationship between $^{240}\text{Pu}/^{239}\text{Pu}$ atom ratios and $^{238}\text{Pu}/^{239+240}\text{Pu}$ activity ratios in cryoconite samples from this study and other Arctic cryoconite (unpublished data) (A); the relationship between $^{240}\text{Pu}/^{239}\text{Pu}$ atom ratio and sample number in tundra soils, cryoconite samples (this study) and other Arctic cryoconite (unpublished data) (B); the correlation between excess ^{238}Pu and $^{240}\text{Pu}/^{239}\text{Pu}$ atom ratios in cryoconite samples (this study) and other Arctic cryoconite (unpublished data) (C).

Figure 6. The relation between the $^{240}\text{Pu}/^{239}\text{Pu}$ atomic ratio and sample number in tundra soils and cryoconite samples (A); the relation between the $^{240}\text{Pu}/^{239}\text{Pu}$ atomic ratios and the $^{238}\text{Pu}/^{239+240}\text{Pu}$ activity ratios in tundra soils and cryoconite samples (B); the correlation between excess ^{238}Pu and $^{240}\text{Pu}/^{239}\text{Pu}$ atom ratios in cryoconite samples (C).

The average $^{239+240}\text{Pu}/^{137}\text{Cs}$ ratios in tundra soils (0.042) is similar to the global fallout value (0.039) expected for the year 2013 (UNSCEAR, 1982); in cryoconite samples (0.020) it is lower. The average $^{239+240}\text{Pu}/^{137}\text{Cs}$ ratios in tundra soils (0.042) is similar to value of 0.039 in the global fallout expected for the year 2013 (UNSCEAR, 1982) but in cryoconite samples (0.020) it is lower. The observed average value of about 0.021 ± 0.03 in the proglacial zone of the glacier from west Spitsbergen is remarkable as there were no other such low values as at this site. Similar or slightly higher values (0.021 – 0.035) were observed on Svalbard-Spitsbergen by Gwynn et al. (2004) and Łokas et al. (2014). The enrichment of soils in ^{137}Cs relative to plutonium might be due to the contributions from sources other than global fallout or these ratios may actually reflect the deposition ratios at this latitude. The mean $^{241}\text{Am}/^{239+240}\text{Pu}$ ratios for tundra soils (0.47) and cryoconite samples (0.46) fall within the ranges reported for soils from western Spitsbergen (Łokas et al., 2014; Łokas et al., 2013) and other Arctic sites (Holm et al., 1983; Smith et al., 1997; Gwynn et al., 2005).

4.3 Heavy metals

The metal concentrations in soils were typical or slightly higher than concentrations characteristic for natural background concentrations. This had been confirmed by generally low metal enrichment ratios (EF ranging from 0.8 to 2.0). The only exception was Cd reaching up to $0.6 \text{ mg} \cdot \text{kg}^{-1}$ (EF=7.7). The metal distributions measured in this study are very similar to metal concentrations characteristic for parent rocks (natural crust) on Svalbard. For instance, parent rocks in Hornsund

contain very low concentrations of heavy metals (Pb 11 mg·kg⁻¹, Cd 0.04 mg·kg⁻¹, Zn 58 mg·kg⁻¹, Cu 20 mg·kg⁻¹; Samecka-Cymerman et al., 2011). Visible enrichment in Cd and Pb was found in surface and subsurface soil profile layers (Fig. 6). Similar enrichment in metal concentrations (Pb up to 25.3 mg·kg⁻¹, Cd up to 0.4 mg·kg⁻¹) in soil profiles of Kaffiøyra was found in the 1980's by Plichta (1993). Generally low metal concentrations have been mapped by Ottesen et al. (2010) in alluvial soils in Kaffiøyra region. Alluvial sediments are believed to represent natural metal levels, although recently deposited layers can be polluted by anthropogenic activity. Cd concentrations ranged from 0.01 to 0.13 mg·kg⁻¹ (our results show values 0.05–0.6 mg·kg⁻¹), Cu ranged from 13.4 to 43.4 mg·kg⁻¹ (our values 11.8–26.6 mg·kg⁻¹), Pb ranged from 8.9 to 24.3 mg·kg⁻¹ (our values 9.6–25.9 mg·kg⁻¹) and Zn 58.9 to 79.2 mg·kg⁻¹ (our values 45.1–76.9 mg·kg⁻¹). Metal concentrations were studied in peat cores collected from nearby Kongsfjorden. The Cu ranged from 7 to 25 mg·kg⁻¹, Pb from 10 to 35 mg·kg⁻¹ while Zn ranged from 20 to 75 mg·kg⁻¹ (Headley, 1996). The region of Kongsfjorden may have been impacted by former mining activity, however (Hao et al., 2013). More recently, the metal distribution in soils of other Svalbard regions (Kongsfjorden and Adventfjorden) was studied by Halbach et al. (2017). Like our study, they found elevated concentrations of some metals eg. Cd (0.13–1.00 mg·kg⁻¹) in surface soils (covered by moss), but low concentrations of metals (Cd=0.04–0.5, Cu=3.3–28.4 mg·kg⁻¹, Pb=2.9–22.7 mg·kg⁻¹ and Zn=25–106 mg·kg⁻¹) in mineral soils (sampled at 20cm depth). Slightly higher levels of heavy metals (Cu: 17–92 mg·kg⁻¹ and Cd: 0.05–1.20 mg·kg⁻¹) were measured in biologically rich soils in Hornsund region with the highest values in areas of rich vegetation (Wojtuń et al., 2013). Ziółek et al. (2017) reports much higher metal concentrations in peat soils in Bellsund. In their study Pb reached 100 mg·kg⁻¹, Zn reached 140 mg·kg⁻¹, Cu 57 mg·kg⁻¹ while Cd reached 8 mg·kg⁻¹. The high contamination of peat soils was in their case caused by the richness of organic matter due to a bird colony. In conclusion, the effect of anthropogenic influence on soils in Kaffiøyra region was very limited. This is also confirmed by relatively high ²⁰⁶Pb/²⁰⁷Pb ratio (1.190–1.217) measured in Kaffiøyra soils that is close to the natural ratio for parent rocks (>1.215) or/and represents the mixture of natural and anthropogenic Pb sources (Zaborska et al., 2017).

~~While soils can receive metals from atmosphere and from the surface run off, cryoconites are fed only by atmospheric particles. Thus, interestingly, metal concentrations in cryoconite were higher than in soil samples (Tab. S8) and definitely exceeded natural values. Particularly elevated was the concentration of Pb reaching 97.5 mg·kg⁻¹ (EF = 6.1) and Cd reaching 0.6 mg·kg⁻¹ (EF=4.5). The enrichment in other metals was lower (Zn up to 97.5 mg·kg⁻¹ and Cu up to 40.1 mg·kg⁻¹). The maximum concentration of Pb in cryoconite samples from the Hansbreen glacier in Hornsund was lower (82.7 mg·kg⁻¹), the concentrations of Zn (108 mg·kg⁻¹) and Cd (1.53 mg·kg⁻¹) were higher, however (Lokas et al., 2016). Singh et al. (2013) found a similar range of metal concentrations in cryoconite samples from a glacier located also in Kongsfjorden area (Pb up to 85.1 mg·kg⁻¹, Cu up to 44 mg·kg⁻¹, Cd up to 0.14 mg·kg⁻¹, Zn up to 150 mg·kg⁻¹). Chmiel et al. (2009) did not find elevated metal levels in glacier waters in the Scott Glacier in Bellsund, they found enrichment in some metals (Pb = 190 mg·kg⁻¹~~

The relatively high ²⁰⁶Pb/²⁰⁷Pb ratio (1.190 - 1.217) measured in Kaffiøyra soils that is close to the natural ratio for parent rocks (>1.215) or/and represents the mixture of natural and anthropogenic Pb sources (Zaborska et al., 2017). The

5 metallic trace elements, which are transported in the atmosphere attached to airborne particulate matter, tend to be retained and concentrated in the cryoconite material. This finding is confirmed by relatively lower $^{206}\text{Pb}/^{207}\text{Pb}$ ratios in cryoconite samples. These ratios suggested that Pb in cryoconite is a mixture of natural and anthropogenic origin and the Pb of anthropogenic origin prevails. Using the end-member method and based on stable isotopic Pb ratios, the contribution of anthropogenically derived Pb was estimated in cryoconite. It was found that Pb with an anthropogenic origin constituted 10-100 % of Pb accumulated in cryoconite samples (Fig. 5). In the case of the Horsund site (Hornbreen) the anthropogenic lead fraction in cryoconite was lower and ranged from 29 % to 95 % (Łokas et al., 2016). It was also calculated that the excess Pb in cryoconite material ($\text{Pb}_{\text{measured}} - \text{Pb}_{\text{natural}}$), was characterized by $^{206}\text{Pb}/^{207}\text{Pb}$ of 1.156-1.189 (Fig. 5). Side stations located closer to rocky glacier sides were characterized by higher (more natural) computed anthropogenic Pb isotopic ratios (1.17-1.19) as compared to stations located in the central transect of the glacier (1.16-1.17). Since glaciers are fed by airborne particles cryoconite $^{206}\text{Pb}/^{207}\text{Pb}$ ratios can be compared to aerosol $^{206}\text{Pb}/^{207}\text{Pb}$ ratios. The study of aerosols in Kongsfjorden region recognized the dominance of two sources of Pb to that region. In the spring, isotopic signatures of $^{206}\text{Pb}/^{207}\text{Pb} \sim 1.158$ suggests an Eastern Eurasian source, while in the summer a signature of $^{206}\text{Pb}/^{207}\text{Pb} \sim 1.167$ suggests a mixture of North American and Eurasian influence (Bazzano et al., 2015). Thus, the cryoconite $^{206}\text{Pb}/^{207}\text{Pb}$ ratios compare well with aerosol $^{206}\text{Pb}/^{207}\text{Pb}$ ratios particularly in the center part of the glacier. This implies that it is mostly anthropogenic and not natural Pb that accumulates in the cryoconite samples and it can be said that an anthropogenic environmental impact is clearly visible in the Spitsbergen cryosphere.

20 Both radionuclides and heavy metals are deposited from the atmosphere on land/ocean and glacier surfaces by wet and dry precipitation. Since there are no new sources of radioactive pollution into the atmosphere, the patterns of atmospheric circulation do not influence radionuclide distribution within different glaciers on Svalbard. The amount of precipitation received by the glacier can definitely influence the glacier pollution, since coastal areas (eg. Kaffiøyra) receive twice the amount of precipitation compared to the inland areas (Førland et al., 2011). Post-depositional processes, mainly those connected to cryoconite hole formation and eventual material transport within the glacier, appear to have more significant influence on the levels of radionuclide pollution within the glacier. The different scenario concerns the distribution of heavy metals. Since they are still emitted to the atmosphere from different sources, the direction of atmospheric circulation is an important factor shaping the metal pollution distribution within the glacier. Main atmospheric metal discharge origins from Europe and Russia (Isaksen et al., 2016), in coastal regions local winds can enhance the introduction of contaminants due to transport of contaminated sea salt aerosols, however (Lüdke et al., 2005 Kozak et al., 2015).

30 ~~in cryogenic sediments, however. These reports suggest that the metallic trace elements, which are transported in the atmosphere attached to airborne particulate matter, tend to be retained and concentrated in the cryoconite material. This finding is confirmed by relatively lower $^{206}\text{Pb}/^{207}\text{Pb}$ ratios in cryoconite samples (1.169 to 1.199). These ratios suggested that Pb in cryoconite is a mixture of natural and anthropogenic origin and the Pb of anthropogenic origin prevails. Using the end-member method and based on stable isotopic Pb ratios, the contribution of anthropogenically derived Pb was estimated in~~

eryoconite. It was found that Pb with an anthropogenic origin constituted 10-100 % of Pb accumulated in cryoconite samples (Fig. 5). In the case of the Horsund site (Hornbreen) the anthropogenic lead fraction in cryoconite was lower and ranged from 29 % to 95 % (Lokas et al., 2016). It was also calculated that the excess Pb in cryoconite material ($Pb_{\text{measured}} - Pb_{\text{natural}}$), that ranged from $7.9 \text{ mg} \cdot \text{kg}^{-1}$ to $85.7 \text{ mg} \cdot \text{kg}^{-1}$, was characterized by $^{206}\text{Pb}/^{207}\text{Pb}$ of 1.156-1.189 (Fig. 5). Side stations located closer to rocky glacier sides were characterized by higher (more natural) computed anthropogenic Pb isotopic ratios (1.17-1.19) as compared to stations located in the central transect of the glacier (1.16-1.17). Since glaciers are fed by airborne particles cryoconite $^{206}\text{Pb}/^{207}\text{Pb}$ ratios can be compared to aerosol $^{206}\text{Pb}/^{207}\text{Pb}$ ratios. The study of aerosols in Kongsfjorden region recognized the dominance of two sources of Pb to that region. In the spring, isotopic signatures of $^{206}\text{Pb}/^{207}\text{Pb}$ 1.158 suggests an Eastern Eurasian source, while in the summer a signature of $^{206}\text{Pb}/^{207}\text{Pb}$ 1.167 suggests a mixture of North American and Eurasian influence (Bazzano et al., 2015). Thus, the cryoconites $^{206}\text{Pb}/^{207}\text{Pb}$ ratios compare well with aerosol $^{206}\text{Pb}/^{207}\text{Pb}$ ratios particularly in the center part of the glacier. This implies that it is mostly anthropogenic and not natural Pb that accumulates in the cryoconite samples and it can be said that an anthropogenic environmental impact is clearly visible in the Svalbard cryosphere. The recent intensification of wind activity may play an important role as a factor enhancing pollution of cryoconite granules. Unfortunately, due to lack of data in the previous year, the importance of atmospheric circulation variability could not be properly assessed.

~~Both radionuclides and heavy metals are deposited from the atmosphere on land/ocean and glacier surfaces by wet and dry precipitation. Since there are no new sources of radioactive pollution into the atmosphere, the patterns of atmospheric circulation do not influence radionuclide distribution within different glaciers on Svalbard. The amount of precipitation received by the glacier can definitely influence the glacier pollution, since coastal areas (eg. Kaffiøyra) receive twice the amount of precipitation compared to the inland areas (Førland et al., 2011). Post depositional processes, mainly those connected to cryoconite hole formation and eventual material transport within the glacier, appear to be a more significant influence on the levels of radionuclide pollution within the glacier. The different scenario concerns the distribution of heavy metals. Since they are still emitted to the atmosphere from different sources, the direction of atmospheric circulation is an important factor shaping the metal pollution distribution within the glacier. The atmospheric circulation over the Arctic has been intensively studied for decades. It is believed that Svalbard receives air masses mainly from Europe and Russia (Reimann and de Caritat, 2005; Isaksen et al., 2016). Air mass transport direction varies seasonally and is influenced by the Arctic Oscillation (AO), however. The largest transport of contaminants to the Arctic occurs in winter and early spring (so called Arctic haze). In the winters characterized by a positive AO index airborne contaminants originating from Europe and eastern parts of North America are dominant; conversely, during winters characterized by a negative AO index contaminants of Eurasian origin prevail. In the summer, contaminants associated with air masses from North America and Europe dominate as the main transport direction (Macdonald et al., 2005). Local winds can also enhance the introduction of contaminants to coastal areas due to transport of contaminated sea salt aerosol (Lüdke et al., 2005; Kozak et al., 2015). The Kaffiøyra land area is shielded from winds from the seaward direction by the mountainous Prins Karl~~

Forland island from the west. The dominating wind directions are north-west and south-east. Moreover, it was recently found that the winds became stronger (when compared to the period of 1975-2014) due to the intensification of cyclonic activity (Przybylak et al., 2016). The recent intensification of wind activity may play an important role as a factor enhancing pollution of cryoconite granules. Unfortunately, due to lack of data in the previous year, the importance of atmospheric circulation variability could not be properly assessed.

5 Conclusions

- 10 The extent of variations of activity concentrations of airborne radionuclides and some trace metals (Pb and Cd) in cryoconite samples from the Waldemarbreen exceeds ranges observed in soils surrounding this glacier. The mechanisms responsible for accumulation of atmospheric contaminants are not very clear but we observed the effect of glacial morphology on effective trapping and storing of airborne radionuclides. The average mass-atom ratios of $^{240}\text{Pu}/^{239}\text{Pu}$ for tundra soils are comparable to the characteristic ratio for global fallout but for cryoconite samples (0.147) are much lower than for surrounding soils and
- 15 also for initial-other initial soils from other west coast of Spitsbergen. We observed an excess of ^{238}Pu in cryoconite samples similar to that which as we observed in other initial soils in Spitsbergen, and this enrichment of ^{238}Pu is derived from other sources other than ^{239}Pu . This is due to pure ^{238}Pu from a satellite re-entry after injection from the stratosphere into the troposphere. Monitoring of cryoconite samples allows identification of Pu release events in the form of a fallout associated with each separate event. Therefore, cryoconite is an excellent candidate for atmospheric contamination monitoring.
- 20 The lithogenic radionuclide contents vary in much narrower ranges and their variability reflects changing proportions of weathering products derived from diverse parent rocks.

Authors Contribution. E.L. designed the idea of this study and performed gamma and alpha analyses. A.C. prepared the calibration for gamma analyses. P.K. performed uranium analyses. I.S. collected the soil and cryoconite samples. P.G. carried out the purification of Nd(Pu)F₃ alpha-spectrometric sources in preparation for MC-ICP-MS measurements and language improvement. J. A. Milton, carried out MC-ICP-MS analysis. E.L., A.Z. wrote the manuscript. E.L., A.Z., P.G. and I.S. supervised the research and contributed to the interpretation of the data. A.M. make-made some language corrections.

Acknowledgment. The research was founded by grant 2016/21/B/ST10/02327 from the Polish National Science Center (NCN) (PI: Edyta Łokas) and IFJ PAS statutory research resources.

Competing Interests. The authors declare that they have no competing interests (both financial or non-financial).

Supplementary information. Full data about environmental radioactivity are available in the Supplementary Material.

References

- AMAP Assessment Report: Heavy Metals in the Arctic. Oslo, Norway, 2002.
- AMAP Assessment Report: Radioactivity in the Arctic. Oslo, Norway, 2015.
- Aizen V.B., Mayewski P.A., Aizen E.M., Joswiak D.R., Surazakov A.B., Kaspari S., Grigholm B., Krachler M., Handley M., Finaev A.: Stable-isotope and trace element time series from Fedchenkoglacier (Pamirs) snow/firn cores, *J. Glaciol.*, 55, 275-291, <http://dx.doi.org/10.3189/002214309788608787>, 2009.
- Baccolo, G., Di Mauro, B., Massabò, D., Clemenza, M., Nastasi, M., Delmonte, B., Prata, M., Prati, P., Previtali, E., and Maggi, V.: Cryoconite as a temporary sink for anthropogenic species stored in glaciers. *Sci. rep.* 7, 9623, doi: 10.1038/s41598-017-10220-5, 2017.
- 10 Barbante, C., Schwikowski, M., Döring, T., Gäggeler, H.W., Schotterer, U., Tobler, L., van de Velde, K., Ferrari, C., Cozzi, G., Turetta, A., Rosman, K., Bolshov, M., Capodaglio, G., Cescon, P., Boutron, C.: Historical record of European emissions of heavy metals to the atmosphere since the 1650s from alpine snow/ice cores drilled near Monte Rosa. *Env. Sci. Tech.*, 38(15), 4085–4090, www.ncbi.nlm.nih.gov/pubmed/15352445, 2004.
- Bazzano, A., Ardini, F., Becagli, S., Traversi, R., Udistri, R., Cappelletti, D., Grotti, M.: Source assessment of atmospheric lead measured at Ny-Ålesund, Svalbard. *Atmos. Environ.* 113, 20-26. doi:10.1016/j.atmosenv.2015.04.053, 2015.
- 15 Bindler, R., Renberg, I., Anderson, N.J., Appleby, P.G., Emteryd, O., Boyle, J.: Pb isotope ratios of lake sediments in West Greenland: interferences on pollution sources. *Atmos. Environ.* 35, 4675-4685. 10.1016/S1352-2310(01)00115-7, 2001.
- Boyle, J.F., Rose, N.L., Appleby, P.G., Birks, H.J.B.: Recent environmental change and human impact on Svalbard: the Lake-Sediment geochemical record. *J. Paleolimnol.* 31 (4), 515-530. doi: 10.1023/b:jopl.0000022549.07298.6, 2004.
- 20 Chmiel, S., Reszka, M., Rysiak, A.: Heavy metals and radioactivity in environmental samples of the Scott Glacier region. *Quaestiones Geographicae* 28A,1, 2009.
- Cook, J.M., Edwards, A., Takeuchi, N. and Irvine-Fynn, T.: Cryoconite. The dark biological secret of the cryosphere. *Prog. Phys. Geog.* 40, 1–46, <https://doi.org/10.1177/0309133315616574>, 2015.
- Cook, J.M., Edwards, A., Bulling, M., Mur, L.A., Cook, S., Gokul, J.K., Cameron, K.A., Sweet, M., Irvine-Fynn, T.D.: 25 Metabolome-mediated biocryomorphic evolution promotes carbon fixation in Greenlandic cryoconite holes. *Environ. Microbiol.* 18, 4674–4686, doi: 10.1111/1462-2920.13349, 2016.
- [Corcho Alvarado, J.A., Steinmann, P., Estier, S., Bochud, F., Haldimann, M., Froidevaux, P.: Anthropogenic radionuclides in atmospheric air over Switzerland during the last few decades. *Nature Comm.* 5, 3030. doi:10.1038/ncomms4030, 2014.](#)
- 30 Dong, Z., Kang, S., Qin, D., Qin, X., Yan, F., Du, W., Wei, T.: Temporal and diurnal analysis of trace elements in the Cryospheric water at remote Laohugou basin in northeast Tibetan Plateau. *Chemosphere*, 171, 386-398, doi: 10.1016/j.chemosphere.2016.12.088, 2017.

- Eyrikh, S., Eichler, A., Tobler, L., Malygina, N., Papina, T., Schwikowski, M.: A 320 Year Ice-Core Record of Atmospheric Hg Pollution in the Altai, Central Asia. *Environ. Sci. Technol.*, 51 (20), 11597-11606, doi: 10.1021/acs.est.7b03140, 2017.
- Førland, E.J.I., Haugen, J.E., Skaugen, T.E.: Temperature and precipitation development at Svalbard 1900-2100. *Adv. Meteo.* 2011. doi:10.1155/2011/893790, 2011.
- Francis A.J.: Microbial mobilization and immobilization of plutonium. *J. Alloy. Compd.* 444, 500–505, doi: 10.1016/j.jallcom.2007.01.132, 2007.
- Gadd, G.M.: Microbial influence on metal mobility and application for bioremediation. *Geoderma* 122 (2–4), 109–119, doi: 10.1016/j.geoderma.2004.01.002 2004.
- 10 Halbach, K., Mikkelsen, Ø., Berg, T., Steinnes, E.: The presence of mercury and other trace metals in surface soils in the Norwegian Arctic. *Chemosphere* 188, December 2017, Pages 567-574 10.1016/j.chemosphere.2017.09.012, 2017.
- Hao L.Z., Wang, F. Yang, H.Z.: Baseline values for heavy metals in soils on Ny-Alesund, Spitsbergen Island, Arctic: the extent of anthropogenic pollution. *Adv. Mater. Res.*, 779–78, 1260-1265. doi: 10.4028/www.scientific.net/AMR.779-780.1260, 2013.
- 15 Headley, A.D.: Heavy metal concentrations in peat profiles from the high Arctic. *Sci. of The Tot. Env.* 177(1-3): 105-111. doi:10.1016/0048-9697(95)04867-7, 1996.
- Hodson, A. Anesio, A.M., Tranter, M., Fountain, A., Osborn, M., Priscu, J., Laybourn-Parry, J., Sattler, B.: Glacial ecosystems. *Ecol. Monogr.*, 78, 41–67, doi.org/10.1890/07-0187.1, 2008.
- Hur S.D., Cunde X., Hong S., Barbante C., Gabrielli P., Lee K., Boutrone C.F., Ming Y.: Seasonal patterns of heavy metal deposition to the snow on Lambert Glacier basin. *East Antarctica Atmospheric Environment* 41 (38), 8567-8578, doi 10.1016/j.atmosenv.2007.07.012 , 2007.
- 20 Isaksen, K., Nordli, Førland, Ø.E.J., Łupikasza, E., Eastwood, S., Niedźwiedz, T.: Recent warming on Spitsbergen and influence of atmospheric circulation and seaice cover. *J. Geophys. Res. Atmos.* 121 (20), 11913-11931. <http://dx.doi.org/10.1002/2016JD025606>, 2016.
- 25 [Kierepko, R., Mietelski, J.W., Ustrnul, Z., Anczkiewicz, R., Wershofen, H., Holgye, Z., Kapała, J. and Isajenko, K.: Plutonium isotopes in the atmosphere of Central Europe: isotopic composition and time evolution vs. circulation factors. *Sci. Total Environ.*, 569, 937-947, doi: 10.1016/j.scitotenv.2016.05.222, 2016.](#)
- Komárek, M., Ettler, V., Chrastný, V., Mihaljevic, M.: Lead isotopes in environmental sciences: a review. *Environ Int.* 34(4), 562-577. doi:10.1016/j.envint.2007.10.005, 2008.
- 30 Kozak. K., Kozioł. K., Luks. B., Chmiel. S., Ruman. M., Marć. M., Namieśnik. J., Polkowska. Ż.: The role of atmospheric precipitation in introducing contaminants to the surface waters of the Fuglebekken catchment, Spitsbergen. *Polar Res.* 34, 24207. doi: 10.3402/polar.v34.24207, 2015.
- Liu X., Jianga S., Zhang P., Xu L.: Effect of recent climate change on Arctic Pb pollution: A comparative study of historical records in lake and peat sediments. *Env. Poll.* 160 (2012) 161-168. doi: 10.1016/j.envpol.2011.09.019, 2012.

- Lüdke, Ch., Skole, J., Taubner, K., Krievs, M.: Trace metal analysis in arctic aerosols by an inductively coupled plasma-time of flight-mass spectrometer combined with an inductively heated vaporizer. *Spectrochim. Acta Part B* 60,1412-1422. doi: 10.1016/j.sab.2005.08.014, 2005.
- 5 [Łokas E., Mietelski J.W., Kleszcz K., Tomankiewicz E.: A sequential procedure for determining ²³⁸Pu, ²³⁹⁺²⁴⁰Pu, ²⁴¹Am, ⁹⁰Sr, U and Th activities in soils and peats from Spitsbergen. *Nukleonika* 55 \(2\) \(2010\) 195-199.](#)
- Łokas, E., Bartmiński, P., Wachniew, P., Mietelski, J.W., Kawiak, T., Środoń, J.: Sources and pathways of artificial radionuclides to soils at a High Arctic site. *Environ. Sci. Poll. Res.* 21, 12479-12493, doi: 10.1007/s11356-014-3163-6, 2014.
- Łokas, E., Zaborska, A., Koliczka, M., Różycki M. and Zawierucha, K.: Accumulation of atmospheric radionuclides and heavy metals in cryoconite holes on an Arctic glacier. *Chemosphere*, 160, 162–172, doi: 10.1016/j.chemosphere.2016.06.051, 2016.
- 10 Łokas, E., Mietelski, J.W., Anczkiewicz, R., Kierepko, R.: Variations in Pu isotopic composition in soils from the Spitsbergen (Norway): Three potential pollution sources of the Arctic region. *Chemosphere*, 178, 231-238, doi: 10.1016/j.chemosphere.2017.03.054, 2017a.
- 15 Łokas, E., Wachniew, P., Jodłowski, P., Gąsiorek, M.: Airborne radionuclides in the proglacial environment as indicators of sources and transfers of soil material. *J. Environ. Radioact.*, 178-179: 193-202, doi: 10.1016/j.jenvrad.2017.08.018, 2017b.
- Łokas, E., Zawierucha, K., Cwanek, A., Szufa, K., Gaca, P., Mietelski, J.W. and Tomankiewicz, E.: The sources of high airborne radioactivity in cryoconite holes from the Caucasus (Georgia). *Scientific reports*, 8(1), p.10802, doi.org/10.1038/s41598-018-29076-4, 2018.
- 20 Nørdli, Ø., Przybylak, R., Ogilvie, A.E.J., Isaksen, K.: Long-term temperature trends and variability on Spitsbergen: the extended Svalbard Airport temperature series, 1898–2012. *Pollut. Res.* 33, 21–49, <http://dx.doi.org/10.3402/polar.v33.21349>, 2014.
- Nriagu, J.O.: A history of global metal pollution. *Science*, 272 (5259), 223-224, doi: 10.1126/science.272.5259.223, 1996.
- 25 Ottesen, R.T., Bogen, J., Finne, T.E.: *Geochemical atlas of Norway. Part 2: Geochemical atlas of Spitsbergen*. NGU, ISBN: 978-82-7385-143-7, 2010.
- Orheim, A., Bieg, G., Brekke, T., Stenvold, J.: Petrography and geochemical affinities of Spitsbergen Paleocene coals. *Nor. Int. J. Coal Geol.* 70 (1), 116e136. doi:10.1016/j.coal.2006.04.008, 2007.
- Pacyna J.M. and Pacyna E.G.: An assessment of global and regional emissions of trace metals to the atmosphere from anthropogenic sources worldwide. *Environ. Rev.*, 9, 269–298, doi.org/10.1139/a01-012, 2001.
- 30 Plichta, W., Kuczyńska I.: Metal contents in soils of Kaffiøyra, Spitsbergen, Pol. *Polar Res.*, 12, 2, 183-193, 1991.
- Plichta, W.: Gleby Kaffiøyry. In: Grześ, M., Sobota, I. (Eds.), *Kaffiøyra. Zarys środowiska geograficznego Kaffiøyry*. 31–34 Oficyna Wydawnicza TURPRESS, Toruń, NW Spitsbergen., 2005.

- Przybylak, R.: The Climate of the Arctic. Ser. Atmospheric and Oceanographic Sciences Library, vol. 52, Springer International Publishing, 287 pp, 2016.
- Radic, V., Bliss, A., Beedlow, A.C., Hock, R., Miles, E., Graham Cogley, J.G.: Regional and global projections of twenty-first century glacier mass changes in response to climate scenarios from global climate models. *Clim. Dynam.* 42, 37–58, doi: 10.1007/s00382-013-1719-7, 2014.
- Reimann, C., de Caritat, P.: Distinguishing between natural and anthropogenic sources for elements in the environment: regional geochemical surveys versus enrichment factors. *Sci. Total Environ.* 337, 91–107. doi:10.1016/j.scitotenv.2004.06.011, 2005.
- Salminen-Paatero, S., Nygren, U. and Paatero, J.: ²⁴⁰Pu/²³⁹Pu mass ratio in environmental samples in Finland. *J. Environ. Radioact.* 113, 163-170, doi: 10.1016/j.jenvrad.2012.06.005, 2012.
- Samecka-Cymerman, A., Wojtuń, B., Kolon, K., Kempers, A.J. 2011. *Sanionia uncinata* (Hedw.) Loeske as bioindicator of metal pollution in polar regions. *Pol. Biol.* 34, 381-388. doi: 10.1007/s00300-010-0893-x
- Segawa, T. Takeuchi, N., Rivera, A., Yamada, A., Yoshimura, Y., Barcaza, G., Shinbori, K., Motoyama, H., Kohshima, S., Ushida, K.: Distribution of antibiotic resistance genes in glacier environments. *Environ. Microbiol. Rep.*, 5, 127–134, doi: 10.1111/1758-2229.12011, 2013.
- Simon, S.L. and Bouville, A.: Radiation doses to local populations near nuclear weapons test sites worldwide. *Health Phys.* 82, 706–725, www.ncbi.nlm.nih.gov/pubmed/12003020, 2002.
- Singh, S.M., Sharma, J., Gawas-Sakhalkar, P., Upadhyay, A.K., Naik, S., Pedneker, S.M., Ravindra, R.: Atmospheric deposition studies of heavy metals in Arctic by comparative analysis of lichens and cryoconite. *Environ. Monit. Assess.* 185, 1367-1376, doi: 10.1007/s10661-012-2638-5, 2013.
- Sobota, I.: The near-surface ice thermal structure of the Waldemarbreen, Svalbard. *Pol. Polar Res.* 30 (4), 317–338, doi: 10.4202/ppres.2009.17, 2009.
- Sobota, I., Lankauf, K.R.: Recession of Kaffiøyra Region glaciers, Oscar II Land. Svalbard. *Bull. Geogr.* 3, 27–45, 2010.
- Sobota, I.: Snow accumulation, melt, mass loss, and the near-surface ice temperature structure of Irenebreen, Svalbard. *Polar Sci.* 5 (3), 321–336, doi.org/10.1016/j.polar.2011.06.003, 2011.
- Sobota, I., Nowak, M., Weckwerth, P.: Long-term changes of glaciers in northwestern Spitsbergen. *Glob. Planet. Chang.* 144, 182–197, doi: 10.1016/j.gloplacha.2016.07.006, 2016.
- Sobota, I.: Selected problems of snow accumulation on glaciers during long-term studies in north-western Spitsbergen, Svalbard. *Geografiska Annaler: Series A, Phys. Geog.* 1-16, doi.org/10.1080/04353676.2017.1297679, 2017.
- Sobota I., Weckwerth P., Grajewski T., Dziembowski M., Greń K., Nowak M.: Short-term changes in thickness and temperature of the active layer in summer in the Kaffiøyra region, NW Spitsbergen, Svalbard. *Catena*, 160: 141-153., doi: 10.1016/j.catena.2017.09.014, 2018.
- StatSoft, Inc. Team. 2011. STATISTICA (data analysis software system), version 10. Tulsa, Oklahoma, StatSoft, Inc.

- Véron, A.J., Church, T.M., Rivera-Duarte, I., Flegal, A.R., 1999. Stable lead isotopic ratios trace thermohaline circulation in the subarctic North Atlantic. *Deep Sea Res. Part II* 46, 919-935. doi:10.1016/S0967-0645(99)00009-0
- Takeuchi, N., Kohshima, S., Shiraiwa, T. & Kubota, K. Characteristics of cryoconite (surface dust on glaciers) and surface albedo of a Patagonian glacier, Tyndall Glacier, Southern Patagonia Icefield. *Bull. Glaciol. Res.* 18, 65–69, 2001.
- 5 Takeuchi, N., Kohshima, S. & Seko, K.: Structure, formation, and darkening process of albedo-reducing material (cryoconite) on a Himalayan glacier: a granular algal mat growing on the glacier. *Arct. Antarct. Alp. Res.* 33, 115–122, doi.org/10.1080/15230430.2001.12003413, 2001.
- Taylor, R. N., Warneke, T., Milton, J. A., Croudace, I. W., Warwick, P. E., and Nesbitt, R. W.: Plutonium isotope ratio analysis at femtogram to nanogram levels by multicollector ICP-MS. *J Anal. Atom. Spectrom.* 16, 279-284, doi: 10.1039/B009078F, 2001.
- 10 UNSCEAR: Ionizing radiation: sources and biological effects. United Nations Scientific Committee on the effects of Atomic. Radiation, New York, 1982.
- Wharton, R.A., McKay, C.P., Simmons, G.M. & Parker B.C. Cryoconite holes on glaciers. *Bioscience* 35, 499–503, www.ncbi.nlm.nih.gov/pubmed/11539025, 1985.
- 15 Wojtuń, B., Samecka-Cymerman, A., Kolon, K., Kempers, A.J., Skrzypek, G.: Metals Metals in some dominant vascular plants, mosses, lichens, algae, and the biological soil crust in various types of terrestrial tundra, SW Spitsbergen, Norway. *Pol. Biol.* 36 (12), 1799-1809. doi.10.1007/s00300-013-1399-0, 2013.
- Zaborska, A., Beszczyńska-Möller, A., Włodarska-Kowalczyk M.: History of heavy metal accumulation in the Svalbard area: Distribution, origin and transport pathways. *Env. Poll.* 231(1):437-450. doi: 10.1016/j.envpol.2017.08.042, 2017.
- 20 Zawierucha, K. Buda, J., Pietryka, M., Richter, D., Łokas, E., Lehmann-Konera, S., Makowska, N., Bogdziewicz, M.: Snapshot of micro-animals and associated biotic and abiotic environmental variables on the edge of the south-west Greenland ice sheet. *Limnol.* 19, 141–150, doi: 10.1007/s10201-017-0528-9, 2018.
- Ziółek, M., Bartmiński, P., Stach, A.: The Influence of Seabirds on the Concentration of Selected Heavy Metals in Organic Soil on the Bellsund Coast, Western Spitsbergen. *Arctic, Antarctic, and Alpine Res.*, 49 (4), 507-25 520. doi: 10.1657/AAAR0016-024, 2017.



Figure 1: The logo of Copernicus Publications.