

Distribution and seasonal evolution of supraglacial lakes on Shackleton Ice Shelf, East Antarctica

Jennifer F. Arthur ¹, Chris R. Stokes ¹, Stewart S.R. Jamieson ¹, J. Rachel Carr ², Amber A. Leeson ³

5 ¹Department of Geography, Durham University, Durham, DH1 3LE, UK

²School of Geography, Politics and Sociology, Newcastle University, Newcastle-upon-Tyne, NE1 7RU, UK

³Lancaster Environment Centre/Data Science Institute, Lancaster University, Bailrigg, Lancaster, LA1 4YW, UK

Correspondence to: Jennifer F. Arthur (jennifer.arthur@durham.ac.uk)

Abstract. Supraglacial lakes (SGLs) enhance surface melting and can flex and fracture ice shelves when they grow and subsequently drain, potentially leading to ice shelf disintegration. However, the seasonal evolution of SGLs and their influence on ice shelf stability in East Antarctica remains poorly understood, despite some potentially vulnerable ice shelves having high densities of SGLs. Using optical satellite imagery, air temperature data from climate reanalysis products and surface melt predicted by a regional climate model, we present the first long-term record (2000-2020) of seasonal SGL evolution on Shackleton Ice Shelf, which is Antarctica's northernmost remaining ice shelf and buttresses Denman Glacier, a major outlet of the East Antarctic Ice Sheet. In a typical melt season, we find hundreds of SGLs with a mean area of 0.02 km², a mean depth of 0.96 m, and a mean total meltwater volume of 7.45 x10⁶ m³. At their most extensive, SGLs cover a cumulative area of 50.7 km² and are clustered near to the grounding line, where densities approach 0.27 km² per km². Here, SGL development is linked to an albedo-lowering feedback associated with katabatic winds, together with the presence of blue ice and exposed rock. Although below average seasonal (December-January-February, DJF) temperatures are associated with below average peaks in total SGL area and volume, warmer seasonal temperatures do not necessarily result in higher SGL areas and volumes. Rather, peaks in total SGL area and volume show a much closer correspondence with short-lived high magnitude snowmelt events. We therefore suggest seasonal lake evolution on this ice shelf is instead more sensitive to snowmelt intensity associated with katabatic wind-driven melting. Our analysis provides important constraints on the boundary conditions of supraglacial hydrology models and numerical simulations of ice shelf stability.

25 **1 Introduction**

Supraglacial lakes (SGLs) are widespread around the margins of Antarctica and can influence ice shelf dynamics both directly and indirectly (Arthur et al., 2020; Kingslake et al., 2017; Stokes, et al., 2019). Ice shelves flex when SGLs pond and drain on their surfaces, which can weaken them and make them more prone to break-up (Banwell, et al., 2013; Banwell et al., 2019; Banwell and MacAyeal 2015). An often cited example of this was the appearance and abrupt drainage of >3,000 SGLs on the surface of the Larsen B Ice Shelf in 2002 in the days preceding its collapse (Glasser and Scambos 2008; Leeson et al., 2020), triggering a chain reaction of SGL hydrofracture and drainage events (Banwell et al., 2013; Robel and Banwell 2019). SGLs

can indirectly influence ice shelf dynamics by lowering surface albedo, which can intensify surface melt and induce a warming effect on the adjacent ice column (Lüthje et al., 2006; Tedesco et al., 2012; Hubbard et al., 2016). SGLs can also act as reservoirs by storing meltwater for crevasse penetration and hydrofracture (Scambos et al., 2000; 2003; 2009). Thus, the evolution and interaction of SGLs on ice shelves has important consequences for the stability of ice shelves, many of which exert a buttressing effect on inland ice flow (Fürst et al., 2016).

Recent research has shown SGLs form part of active surface meltwater systems on the surface of numerous Antarctic outlet glaciers and ice shelves (Banwell et al., 2019; Bell et al., 2017; Langley et al., 2016; Lenaerts et al., 2017; Moussavi et al., 2020; Kingslake et al., 2017; Stokes et al., 2019) and are particularly widespread on the Antarctic Peninsula and around the margins of the East Antarctic Ice Sheet (Arthur et al., 2020; Kingslake et al., 2017; Stokes et al., 2019). However, few studies have conducted detailed analyses of the seasonal behaviour and spatial distribution of SGLs on ice shelves in East Antarctica. As such, direct observations of lake interactions and drainage are currently constrained to a limited number of ice shelves (Banwell et al., 2019; Bell et al., 2017; Langley et al., 2016; Liang et al., 2019; Moussavi et al., 2020; Kingslake et al., 2015), and our understanding of the importance for SGL evolution on ice-shelf-instability remains limited.

Here, we focus on Shackleton Ice Shelf, which is East Antarctica's sixth largest and most northerly remaining ice shelf in Antarctica (Rignot et al., 2013; Zheng and Zhou, 2019). Over 70% of the ice shelf area exerts a buttressing effect on the major outlet glaciers flowing into it and a number of ice rises and islands act as pinning points for the ice shelf (Fürst et al., 2015; 2016; Fig. 1). The remaining 30% of the ice shelf has been identified as passive shelf ice, i.e. that could be removed without acceleration of inland ice (Fürst et al., 2016). The ice shelf is fed by Denman Glacier, whose catchment holds ice equivalent to ~1.5 m of sea-level and is ~1.9 times larger than the catchment of Pine Island Glacier (Morlighem et al., 2019; Stephenson and Zwally, 1988; Willis et al., 2016). Portions of the ice shelf experience high mean surface melt rates (>200 mm w.e. yr⁻¹) compared to other parts of coastal East Antarctica, such as the Fimbul (> 80 mm w.e. yr⁻¹), Riiser-Larsen (> 70 mm w.e. yr⁻¹), Nivlisen (> 80 mm w.e. yr⁻¹), and Roi Baudouin (> 150 mm w.e. yr⁻¹) ice shelves (Trusel et al., 2013). High densities of SGLs in the ice shelf grounding zone suggest that it contains ice-saturated firn (Alley et al., 2018; Lenaerts et al., 2017). The combination of its topographic setting, intense surface melting and low firn air content (FAC) make Shackleton Ice Shelf potentially vulnerable to hydrofracturing (Alley et al., 2018; Stokes et al., 2019).

Our overall aim is to investigate the long-term distribution and evolution of SGLs on Shackleton Ice Shelf. Our first objective is to analyse the intra-seasonal and inter-seasonal evolution in the spatial distribution, extent and volume of SGLs over multiple melt seasons from 2000 to 2020 using remote sensing. Our second objective is to explore potential glaciological and climatic controls on SGL formation and distribution on this ice shelf in order to improve our understanding of its potential vulnerability.

2 Study site

65 Shackleton Ice Shelf, in Queen Mary Land, East Antarctica (65 °S; 100 °E) covers an area of 33,820 km² (Zheng & Zhou, 2019, Fig. 1). Denman Glacier flows into the ice shelf at speeds over 1800 m a⁻¹ at its grounding line (Rignot et al. 2019) and occupies a trough deeper than 3500 m b.s.l which connects to the Aurora subglacial basin (Morlighem et al., 2019; Roberts et al., 2011). Denman Glacier has accelerated 16% since the 1970s (Rignot et al., 2019) and is thinning inland (Flament and Rémy 2012) and undergoing grounding line retreat (Konrad et al., 2018). The ice shelf surface flow has accelerated by 43%
70 between 1957 and 2016 (Rignot et al., 2019). Several smaller outlet glaciers drain into the ice shelf, including the Scott, Apfel, Remenchus and Northcliffe Glaciers (Fig. 1). The Bunger Hills (a large nunatak) and an extensive blue ice area lie adjacent to the ice shelf grounding zone (Fig. 1). Ice thickness varies across the ice shelf from 231 m (at the calving front) to < 10 m around Mill Island (Fretwell et al., 2013; Wearing et al., 2015). Thinning of Shackleton Ice Shelf at a rate of 0.12 m yr⁻¹ between 2003 and 2008 has been attributed to enhanced basal melting (Pritchard et al., 2012). SGLs on Shackleton Ice Shelf
75 have been previously mapped for the year 2015 from Landsat 8 imagery, to assist interpretation of snowmelt detection from microwave radiometer and scatterometer data (Zheng and Zhou 2019), but no studies of seasonal and/or interannual SGL evolution currently exist for the ice shelf.

3 Data and methods

80 3.1 Satellite imagery acquisition and processing

For the years between 1974 and 2020 we acquired all Landsat 1, 4 and 5 MSS (Multispectral Scanner), TM (Thematic Mapper), Landsat 7 ETM+ (Enhanced Thematic Mapper), Landsat 8 OLI (Operational Land Imager) and Sentinel-2A/B MSI (Multispectral Instrument) imagery over Shackleton Ice Shelf during the austral summer months (November to February) with ≤ 20% cloud cover from the USGS EarthExplorer (<https://earthexplorer.usgs.gov>) and the European Space Agency Copernicus
85 Open Access Hub (<https://scihub.copernicus.eu/dhus>; Supplementary Table 1). For scenes that included cloud cover below the ≤ 20% threshold, it was mostly restricted to the outermost parts of the ice shelf or further inland beyond the grounding line, where cloud free scenes indicated that no lakes form. Eight scenes with cloud cover higher than 20% were used, where cloud cover did not obscure most of the ice shelf and SGLs were visible. These were used to analyse the spatial distribution of lakes across the ice shelf but not in our analysis of lake area and volume (Sect. 3.3). The multispectral bands of the Landsat-8 images
90 were pan-sharpened to 15 m using the panchromatic Band 8 and a Brovey Transform (Gillespie et al., 1987) in QGIS. For more recent years (2017 to 2019), Sentinel-2A/B imagery was used in preference to Landsat 8 or another sensor, such as the Moderate Resolution Imaging Spectroradiometer (MODIS) or Advanced Spaceborne Thermal Emission and Reflectance Radiometer (ASTER), owing to its high spatial resolution (10 m) and revisit period of 5 days, , though Landsat 8 imagery was used where no Sentinel-2 imagery was available with suitable (≤ 20%) cloud cover.

95

It is important to remove the influence of different satellite imagery acquisition conditions before extracting SGL extents and depths. To do this, Digital Numbers were converted to Top of Atmosphere (TOA) reflectance for red and blue bands using the DOS1 (Dark Object Subtraction) atmospheric correction (Moran et al., 1992) in QGIS and the scene-specific metadata file containing reflectance multiplicative and additive scaling factors and solar elevation angle. This process removes the cosine effect at different solar zenith angles (due to different scene time acquisitions) and compensates for differences in solar irradiance.

3.2 Automated mapping of SGL extent

We used the Normalised Difference Water Index adapted for ice (NDWI; Yang and Smith 2013) to classify pixels as ‘water’ or ‘non-water’ (i.e. ice, snow, exposed bedrock or ocean) following Eq. (1):

$$105 \quad NDWI_{ice} = \frac{(Blue-Red)}{(Blue+Red)} \quad (1)$$

This version of the NDWI has been widely applied to mapping surface meltwater in Antarctica (Arthur et al., 2020; Banwell et al., 2019; Bell et al., 2017) and Greenland (Doyle et al., 2013; Williamson et al., 2017; Yang and Smith 2013). We found $NDWI_{ice}$ produced fewer false SGL classifications over blue ice/slush areas than with other versions of the NDWI, such as those using the near-infrared band (Supplementary Fig. 1).

110

The first step in our procedure for mapping SGLs was to apply a bedrock mask derived from Landsat 8 data (Burton-Johnson et al., 2016). Exposed bedrock appears spectrally similar to water in the red band, so masking bedrock avoids mis-classification of rock as water. This was particularly important in our study given the large area of nunataks (Bunger Hills) adjacent to the ice shelf. After rock masking was performed, we applied an NDWI threshold value of 0.25, following previous studies (Banwell et al., 2019; Bell et al., 2017; Dell et al., in review; Doyle et al., 2013; Williamson et al., 2017; Yang and Smith 2013), meaning pixels with $NDWI_{ice} > 0.25$ were assumed to be water-covered. This threshold was found to be the most appropriate for detecting lakes, apart from eleven scenes, where the threshold value was manually tuned in 0.01 increments to improve SGL identification by including pixels which were visibly water-covered in true-colour composites of the area (Supplementary Table 1).

120

Once an NDWI threshold value was applied, each image was reclassified into a binary image and vectorised (converted to polygon shapefiles). Clouds were masked by manually by inspecting each RGB image and deleting any cloud. Some manual editing of SGL extents was necessary where floating ice in the centre of SGLs had resulted in pixels being mis-classified as water. The transition from melting ice to saturated firn, slush, shallow ponds and to deeper SGLs makes it challenging to adopt a clear binary definition of ‘lake’ versus ‘non-lake’ (Dell et al., in review). To account for this uncertainty we used a minimum size threshold of two pixels, in order to remove very small SGLs likely comprised solely of mixed pixels (i.e. pixels containing

125

a combination of water, slush and/or snow or ice; 200 m² for Sentinel-2, 450 m² for Landsat 7/8, 1800 m² for Landsat 4/5, 7200 m² for Landsat 1), following previous studies (Moussavi et al., 2020; Pope et al., 2016, Stokes et al., 2019). For the purposes of our study, we chose not to classify SGLs separately to supraglacial meltwater channels, given the ambiguity in distinguishing between channels and thin, elongate SGLs.

Finally, NDWI classifications were checked against RGB composites and false positives, including small bedrock outcrops not included in the bedrock mask. Dark crevasses were also manually removed. To further reduce noise and improve the visual clarity of the dataset, we then applied the dissolve function in ArcMap to clean up overlapping pixels and the aggregate tool to combine pixels within a distance of two pixels. Apart from potentially reducing large numbers of very small SGLs, these changes were largely cosmetic and are insignificant in terms of the uncertainties associated with identifying SGLs and calculating total SGL area, described in Sect. 3.4.

3.3 Extracting lake depths and volumes

We applied a physically-based radiative transfer model to calculate the water depth of all pixels classified as lake (Pope et al., 2016; Sneed and Hamilton 2007). This method calculates lake water depth (z) using the rate of light attenuation in water, lake-bottom albedo, and optically-deep water reflectance (Philpot, 1989) following Eq. (2):

$$z = \frac{[\ln(A_d - R_\infty) - \ln(R_z - R_\infty)]}{g} \quad (2)$$

where A_d is lake bed reflectance, R_∞ is the reflectance of optically deep water, R_z is the reflectance value of a water-coloured pixel and g is the attenuation co-efficient rate.

The radiative transfer model was applied to a subset area of the ice shelf (48 x 48 km², Fig. 1) to allow lake depth comparisons between years with different satellite imagery coverage. This subset area was selected as the region in which most SGLs were concentrated during each melt season with consistent coverage of cloud-free imagery, enabling us to extract changes in lake area and depth on both the floating ice shelf and the grounded ice upstream of its grounding line. Lake area and depth statistics presented in Sect. 4 are all extracted from this subset area. We excluded nine image scenes from our analysis which did not fully cover this subset area, to maintain full comparability between individual dates. Issues with cloud cover resulted in only one useable scene during some melt seasons, and no suitable imagery in others (2002, 2004-2006; Supplementary Table 1).

A_d was calculated image-by-image by extracting the mean reflectance of a two-pixel buffer around each lake polygon. This is an improvement on previous approaches which used static A_d values across a region (e.g. Sneed and Hamilton, 2007). The reflectance value immediately beyond identified lake areas is assumed to be comparable to the lake bed if it were at the surface

(Pope et al., 2016). R_{∞} is typically approximated as pixel reflectance from open ocean water. Most scenes used in this study did not contain optically-deep water, and we found the average difference between using an R_{∞} of zero and using R_{∞} values averaged from fifteen nearby scenes to be <10% (Supplementary Information). Therefore, we used an R_{∞} of zero for all scenes, following MacDonald et al. (2018) and Banwell et al. (2019).

Red light attenuates more strongly in water than blue light, meaning that there are larger measurable changes in red reflectance over water than for other wavelengths (Box and Ski 2007; Pope et al., 2016). For Sentinel-2 scenes, we used red (Band 4) reflectances for R_z and a g value of 0.8304, following Williamson et al. (2017). For Landsat 8 scenes, we followed the recommendation of Pope et al. (2016) and took an average of the depths calculated using the red (Band 4) and panchromatic (Band 8) band TOA reflectances, using g values of 0.7507 and 0.3817 respectively. For Landsat 7 ETM+ scenes, we used red (Band 3) reflectances for R_z and a g value of 0.8049 following Pope et al. (2016). We are unable to calculate lake depths before the year 2000 (i.e. from Landsat 1 MSS, Landsat 4 TM and Landsat 5 TM scenes) using this physically-based depth method because lab-based absorption coefficients within the full red band wavelength ranges (Pope and Fry 1997) are unavailable. Lake depths within the subset area were extracted using zonal statistics in ArcMap. Lake volumes were calculated by multiplying each pixel area by its calculated water depth within the lake boundaries.

3.4 Errors and uncertainties in extracting lake areas and depths

We quantified the uncertainties in our mapping technique by quantitatively comparing SGL areas derived from NDWI with those derived from manual digitisation in four sample areas (Supplementary Fig. 2a-d). The smaller SGLs (<0.01 km²) generated the largest percentage differences, and we found a mean absolute error of 0.007 km² and a root mean square error of 0.029 km² between manually-digitised SGL areas and those derived from NDWI. We attribute this to manual digitisation being less conservative at 'diffuse', less well-defined SGL edges. Despite these differences, we found very close agreement between the two methods ($R^2 = 0.979$, $p = 0.006$). We also quantified the impact of sensor resolution on lake detection and lake extents, and found there is generally a very good agreement between lake areas derived from Landsat 8 and Sentinel-2 (Supplementary Fig. 3). We therefore assign a conservative uncertainty of 1% to our total SGL area for each time step, following Stokes et al. (2019).

The depth-reflectance algorithm used to extract lake depths assumes that lake-bottom albedos are homogenous, that lake water contains minimal dissolved matter and that there is no scattering of light from the lake surface associated with wind-driven roughness (Sneed and Hamilton 2007). We acknowledge that the last of these assumptions may not always hold if there are wind-driven surface waves. We also acknowledge that wind-blown sediment from the adjacent Bunger Hills rock outcrops could cause variability in lake-bottom albedo. This is known to be an important process on McMurdo Ice Shelf, which is heavily debris-covered due to sediment redistribution from medial moraine and rock outcrops (Glasser et al., 2014; MacAyeal

190 et al., 2019; Banwell et al., 2017; 2019; MacDonald et al., 2020). Where lakes were partially ice-covered, we could not calculate the depth of the central, probably deepest, regions of these lakes. In such cases, we calculated the depth of the outer ring of lake-ice free water, following Banwell et al. (2014). This is likely to underestimate the volumes of deeper lakes, but we record maximum lake volumes that are similar (i.e. not substantially lower) to those reported elsewhere in Antarctica (Bell et al., 2017; Banwell et al., 2014; Langley et al., 2016).

195

3.5 Extraction of lake elevations, slopes, densities and ice surface velocities

Lakes were classified as forming on grounded or floating ice using the Making Earth System Data Records for Use in Research Environments (MEaSUREs) grounding line which is derived from an average of differential interferometric synthetic-aperture radar (inSAR) data from 1992 to 2014 (Rignot et al., 2016). Although a more recent inSAR grounding line exists for parts of Denman and Apfel Glaciers, derived from ERS-1/2 and Sentinel-1 data acquired in 1996, 2015 and 2017 (DLR, 2018), this is incomplete on Shackleton Ice Shelf, and the MEaSUREs grounding line was therefore used as the most recent continental-wide grounding line. Surface elevation and slope values were extracted from the Reference Elevation Model of Antarctica (REMA), which has an 8 m spatial resolution. Ice velocity values were extracted from the MEaSUREs InSAR-based ice velocity mosaic which was derived by averaging data covering the period 1996-2016 (Rignot et al., 2017). Lake area densities were calculated using lake polygon centroids as inputs to the Point Density tool in Arcmap. This tool calculates the cumulative area of SGLs within 1 km² cells by dividing the total number of points that fall within a defined neighbourhood around each raster cell by the area of the neighbourhood. Stokes et al. (2019) apply this method across the East Antarctic Ice Sheet using a 50 km search radius. Here, a 20 km search radius was chosen as an appropriate size given the area of the ice shelf.

210 3.6 Extracting near-surface temperature, surface snowmelt and firn air content

To compare lake areas with surface climatological conditions, we used near-surface air temperature and modelled surface snowmelt. In the absence of any automatic weather stations within 80 km of Shackleton Ice Shelf (and hence any long-term meteorological observations within our 2000-2020 study period), we used monthly averaged near-surface temperature (2 m temperature) from the European Centre for Medium-Range Weather Forecasts ERA5 reanalysis from 1979 to 2019 as the best available option. We extracted near-surface temperature over a ~262 x 370 km box covering the whole ice shelf (Fig. 1). ERA5 reanalysis is provided at 0.25 degree (~31 km) resolution. To investigate whether timings of modelled snowmelt events and extensive SGL ponding were comparable, we extracted mean snowmelt over the ice shelf from daily surface melt flux outputs for the period 2000 to 2019 generated by the Regional Atmospheric Climate Model (RACMO) version 2.3p1, which has a horizontal resolution of 27 km. Details of the model can be found in Van Wessem et al. (2018). To investigate the relationship between lake areas and simulated firn air content (FAC), we extracted mean and minimum FAC (total integrated pore space,

220

i.e. the thickness of the equivalent air column contained in the firn in metres) over the ice shelf from daily FAC for the period 2000 to 2019 generated by RACMO version 2.4p1.

3.7 Hydrological routing analysis

225 To estimate potential meltwater routing across the ice shelf surface (Leeson et al., 2012; Bell et al., 2017) and compare this to mapped distributions of SGLs, a D-infinity algorithm (Tarboton, 1997) was applied to a 100-metre resolution version of the REMA DEM (clipped to Shackleton Ice Shelf). We calculated flow direction and accumulated flow into each cell after filling closed basins (sinks) in the DEM. The output raster was then vectorised to show the potential pathways of surface meltwater routing across the ice shelf once surface depressions have been filled.

230 4 Results

4. Spatial distribution of SGLs

We find that SGLs tend to cluster around the ice shelf grounding zone, near exposed blue ice areas and the Bunger Hills nunataks, whereas they are almost entirely absent towards the ice shelf calving front, where surface melt rates are highest (Trusel et al., 2015; Fig. 2a). Lakes form in particularly high densities on Scott, Apfel and Remenchus Glaciers. Indeed, lake area densities reach their maximum (0.27 km² per km²) on this portion of the ice shelf (Fig. 2b). No lakes form on the heavily crevassed tongue of Denman Glacier. On the ice shelf itself, most lakes are only recorded in a given position on a single date in a particular year, because they advected downstream with ice flow (Fig. 2b-d). The largest recorded feature is an area of coalesced lakes covering up to 11.96 km² which formed in January 2020 on Scott Glacier, alongside more linear, elongate lakes which occupy a ~10 m deep longitudinal depression on the glacier surface (Fig. 2d). Further west, smaller lakes occupy crevasses in the suture zone between Denman and Scott Glaciers (Fig. 2c). Though the spatial distribution of lakes on these parts of the ice shelf remains similar during each melt season, lakes gradually become advected downstream with ice flow each year (Fig. 2f).

240 Upstream of the grounding line, lakes have a high tendency to reoccur annually in the same locations (Fig. 2d). Specifically, seven lakes upstream of Apfel Glacier were present in 62% of scenes mapped (Fig. 2d). Just upstream of this cluster of lakes represents the upper elevation limit (527 m a.s.l) of recorded SGLs, 12.8 km inland from the grounding line (~90 km from the coastline). Lakes only formed at this maximum elevation during the six melt years when total lake volume was highest (2010, 2014, 2015, 2017, 2018 and 2020; see Sect. 4.2).

250 The concentration of SGLs around the ice shelf grounding zone is reflected in their topographic distribution, which is strongly skewed towards low (≤ 10 m) elevations and low ($< 1^\circ$) ice surface slopes close to and just above the grounding line (Fig. 3a-b). SGLs preferentially form on slower-moving ice (< 50 m yr⁻¹), typically occupying smaller tributary glaciers and suture zones between glaciers rather than on fastest-flowing portions of the ice shelf like Denman Glacier (Fig. 3c). Lakes form in very close proximity to exposed bedrock (< 10 m) and are within or downstream of blue ice areas around the ice shelf grounding
255 line (Fig. 3d).

4.2 Seasonal evolution of SGLs

4.2.1 Intra-seasonal and inter-seasonal lake evolution

To investigate the seasonal behaviour of SGL formation and evolution on Shackleton Ice Shelf, we present time series of total lake area, depth and volume over the 2000-2020 melt seasons (Fig. 4). In a typical melt season, we observed hundreds (~100-
260 1000) of SGLs that were, on average, 0.02 km² in area, 0.96 m deep, and held an average total meltwater volume of 7.45×10^6 m³. The earliest that meltwater starts being stored on the ice shelf surface within SGLs is in late November (Fig. 4, Table 1). This is earlier than has been reported in other regions of East Antarctica, where SGLs have been observed forming from mid-December, but typically begin forming in early January (Langley et al., 2016; Leppäranta et al., 2013; Moussavi et al., 2020). We suggest the earlier onset of lake formation reflects an earlier start to the melt season due to the northerly location of this
265 ice shelf.

Total lake area on the ice shelf increases through December/January and peaks in early- to late-January (Fig. 4a), and then decreases through the remainder of the melt season until late February. Mean and maximum lake depths follow a similar seasonal pattern to lake area, and the deepest lakes are generally recorded between early- to mid-January (Fig. 4b). However,
270 mean and maximum lake depths do not always peak simultaneously with lake area (Fig. 4b, Table 1). For example, in 2020, lake depth peaks 30 days before total lake area and volume peak on 1st January. SGLs on grounded ice upstream of the ice shelf have a higher mean depth of 0.94 m (st dev: 0.68, $p = 0.01$) compared to a mean depth of 0.91 m (st dev: 0.54) on the ice shelf (Supplementary Fig. 4). Total meltwater volumes remain low ($< 5 \times 10^6$ m³) during the months of November, December and February and peaks between early- to late-January (Fig. 4c). In 2014, 2015 and 2020, lake volume was more than two
275 standard deviations from the 2000-2020 mean. In 2020, total lake area also deviated by almost three standard deviations from the 2000-2020 mean (Fig. 5b). Total lake area was up to 3.9 times higher and total lake volume up to 4.5 times higher than the long-term mean in the five years 2014, 2015, 2017, 2018 and 2020. At the time of peak total lake volume during these years, lakes covered 0.05%, 0.06%, 0.03%, 0.04% and 0.09% of the total ice shelf area, respectively.

280 We document lakes forming during three years prior to 2000 where suitable imagery was available (1974, 1989 and 1991), though we were unable to include these in our depth and volume analyses (see Sect. 3.3). During these years, lakes occupy

many of the same locations as in later melt seasons, including on Scott and Apfel glaciers, and a smaller blue ice area near the northern part of the ice shelf grounding line. SGLs are less extensive during these years than in 2000-2020 (total area 20.3 km², 5.6 km² and 8.8 km² respectively), though we note imagery was limited to the very start (early November) and end (late February) of the melt season.

4.2.2 Lake drainage events

We observe lakes disappearing in three ways: supraglacial drainage via channels over the ice surface, englacial drainage into the firn, or surface refreezing and/or snow burial (Fig. 6). Lakes commonly refreeze at the end of a melt season, indicated by relict frozen lake scars which are similar to refrozen lakes in previous studies (e.g. Langley et al., 2016; Leeson et al., 2020; Tuckett et al., 2019; Fig. 6a-c). This is the dominant mechanism of lake disappearance and is commonly preceded by ice ‘lids’ that grow outwards from lake centres (Fig. 6a-c).

Earlier in the melt season, some lakes drained supraglacially into others via surface channels (Fig. 6d-f). During the peak of the melt season in some years, large portions of the ice shelf firn layer become saturated, causing meltwater to accumulate and coalesce in large lakes on portions of the ice shelf, such as on the tongue of Scott Glacier. This occurred when total lake area exceeded 10 km² (in February 2004, January 2014, 2015, 2017, 2018 and 2020). During these seasons of more extensive surface melt, supraglacial rivers develop by mid-January, fed by lakes upstream of the grounding line (e.g. Fig. 6g). The largest of these rivers reaches ~14 km on Apfel Glacier, though these feed other lakes rather than exporting meltwater off the ice shelf edge, as has been observed on Nansen Ice Shelf in Antarctica (Bell et al., 2017).

We also recorded several instances of lakes apparently draining englacially into the firn over a ≤ 7 -day period during January 2018 and 2020 (Fig. 6h-l). For example, one chain of lakes (Fig. 9k-l) lost ~36% of their 2.2 km² total area and ~24% of their 3.5 x 10⁶ m³ total volume in ≤ 7 days. If this chain of lakes had refrozen, we would have expected to record a reduction in area of the surrounding lakes. However, the surrounding lakes did not change in area over this period, suggesting they also did not refreeze. Neither did downstream lakes grow in size, and nor was this chain of lakes associated with any major streams/rivers which could have exported the meltwater supraglacially. Therefore, we interpret these lakes to have drained englacially into the firn. In other years, this same chain of lakes remains undrained and freezes over (as in Fig. 6a-c).

4.2.3 Lake evolution and relationship with climate

We compare total lake area during the 2000-2020 melt seasons with mean near-surface temperature extracted from ERA5 reanalyses to investigate the relationship between seasonal lake coverage and temporal fluctuations in air temperature on Shackleton Ice Shelf (Fig. 7). We find that warmer melt seasons do not necessarily correlate with more extensive lake coverage.

For example, although 2019/2020 records the highest total lake area (28.55 km²), it has a lower mean December-January-February (DJF) near-surface temperature (-9.3 °C; Fig. 7c). Similarly, in January 2015 and 2014, the next most extensive lake years, mean DJF temperature was also low (-8.4 °C and -8.6 °C respectively), although lakes started to form earliest in the melt season (19th November 2015 and 2nd December 2014; Fig. 7b). However, in the melt seasons with the five lowest mean DJF temperatures (1999-2000, 2007-2008, 2011-2012, 2012-2013 and 2015-2016), total lake area did not exceed 5 km² (Table 2, Fig. 7b). Likewise, we recorded a relatively high maximum lake extent in 2016-17 (10.29 km²), which was the melt season with the highest mean DJF temperature (-6.4 °C; Fig. 7c). Overall, from 2000-2020, total lake area is only weakly positively correlated with mean surface air temperature in the preceding part of the melt season, i.e. from 1st November to the date of lake observation ($r^2= 0.15$, $p= 0.04$; Fig. 8). Therefore, peak years in lake coverage exhibit only a weak relationship with near-surface temperature, suggesting that other factors such as localised albedo interactions may act as more important controls.

To investigate whether there was a correspondence between peak lake area and volume with surface melt rates (which include meltwater available to refreeze in the firn), we compared our lake dataset to modelled surface melt from RACMO 2.3p1. Multiple large (>4 mm w.e. day⁻¹) but generally short lived (≤ 1 week) spikes in melt are modelled, separated by periods of little-to-no surface melt (Fig. 9). A correspondence between peak lake area, volume and surface melt is particularly apparent in 2014, 2015, 2017 and 2018 (Fig. 9). For example, in 2014-2015, total lake area (18.3 km²) and volume (31.5 x10⁶ m³) peak five days after a large surface melt event (9 mm w.e day⁻¹; Fig. 9b). Similarly, in 2017-2018, total lake area (12.4 km²) and volume (15.2 x10⁶ m³) peak eight days after a spike in surface melt (11 mm w.e day⁻¹; Fig. 9d). Though total lake area and volume may have peaked more rapidly following these modelled surface melt events, temporal gaps in cloud-free imagery prevent us from constraining these timings further. Unfortunately, we have insufficient observations to detect whether a correspondence exists between peak lake area, volume and surface melt in other melt seasons (Supplementary Fig. 5). We also note that at the time of submission, RACMO2.3p1 data for the 2019-2020 melt season are unavailable which prevents us from comparing surface melt with total lake area and volume over this period. According to simulations by RACMO 2.3p1 over the whole ice shelf, total lake area is not significantly correlated to minimum firn air content (FAC; r^2 0.01, $p = 0.56$) and mean FAC (r^2 0.009, $p = 0.63$) during the years for which we had lake observations (Fig. 10a-b). However, there is a qualitative correspondence between years of particularly low mean and minimum FAC and years of high melt (2014, 2015 and 2017; Fig. 10c). During these years, mean FAC ranged from 14.5 to 14.59 m and minimum FAC did not exceed 0.31 m.

5 Discussion

5.1 Spatial distribution of SGLs

The clustering of SGLs in the grounding zone of Shackleton Ice Shelf can be explained by localised katabatic wind-driven feedbacks. Extensive blue ice areas upstream of the grounding line and adjacent to the Bunger Hills nunataks (Fig. 1) are

345 indicative of persistent snow surface scouring by easterly katabatic winds. These winds descend from the polar plateau,
generated by large surface slopes ($>20^\circ$) and reach 19 m s^{-1} in summer (Doran et al., 1996). Surface meltwater production is
enhanced as the winds warm adiabatically as they descend to the grounding zone, which lowers humidity and raises near-
surface air temperatures (Doran et al., 1996; Lenaerts et al., 2017). The lower albedo of blue ice areas (~ 0.5) compared to
350 refrozen snow (~ 0.7) exerts a positive feedback on surface melting by enhancing solar radiation absorption (Kingslake et al.,
2017; Lenaerts et al., 2017; Stokes et al., 2019). SGLs have been associated with strong katabatic wind-induced scouring of
blue ice areas on several other East Antarctic ice shelves (Bell et al., 2017; Kingslake et al., 2017; Lenaerts et al., 2017; Stokes
et al., 2019). This albedo-lowering feedback is locally enhanced by the presence of the Bunger Hills and surrounding nunataks
(Fig. 1), whose lower albedo further enhances surface melting and ponding on these portions of the ice shelf. This is a typically
localised effect (i.e. within 10 km; Kingslake et al., 2017; Leppäranta et al., 2013; Winther et al., 1996; Stokes et al., 2019).

355

It is also clear that firn air content (FAC) is an important control on the spatial distribution of SGLs on Shackleton Ice Shelf.
Lake clustering near the grounding line means repeated meltwater percolation and refreezing into the snow and firn layers is
likely to have formed refrozen ice lenses or superimposed ice zones (Hubbard et al., 2016). The large coalesced lakes we
observed in 2014, 2015 and 2020 (e.g. Fig. 2d) are suggestive of the firn layer becoming saturated in this portion of the shelf.
360 Simulations from RACMO 2.3p1 indicate that the FAC may be as low as $\leq 1 \text{ m}$ on parts of the ice shelf (Fig. 10a, 10c). SGLs
could therefore become more extensive in the ice shelf grounding zone in future, because saturated firn promotes surface
ponding by limiting meltwater percolation and its lateral flow within the firn layer (Alley et al., 2018; Hubbard et al., 2016;
Lenaerts et al., 2017). The absence of SGLs further down-ice from the grounding zone can be explained by increased
accumulation (snowfall) rates towards the ice shelf calving front which result in a corresponding higher FAC, meaning surface
365 meltwater can be retained in a deeper snowpack and firn before it can pond into SGLs (Bell et al., 2017; Stokes et al., 2019).
Thus, the clustering of SGLs around the grounding line is suggestive of air-depleted firn, which acts as a primary control on
where meltwater can pond on the ice shelf surface.

At a more local scale, lakes on grounded ice upstream of the ice shelf grounding line have a high tendency to reform annually
370 in the same locations (Fig. 2), which we attribute to the glaciological processes controlling SGL formation. Ice surface
depressions on grounded ice are translations of subglacial topographic undulations, and are therefore fixed in space rather than
migrating with ice flow (Echelmeyer et al., 1991; Lampkin and VanderBerg 2011; Langley et al., 2016). Conversely, lakes on
the ice shelf itself are generally shorter-lived because they form in surface depressions that migrate annually with ice flow and
are progressively advected towards the ice shelf front, as on other Antarctic ice shelves (Banwell et al., 2014; Bell et al., 2017;
375 Glasser and Gudmundsson 2012; Luckman et al., 2014; Reynolds and Smith 1981). We speculate that surface undulations
perpendicular to the grounding line (i.e. parallel to ice flow) may form as ice flow converges when it crosses the break in slope
and is deflected around the Bunger Hills (Fig. 2c), which could flex and fracture the ice (Banwell et al., 2014; Glasser and
Gudmundsson, 2012).

380 Though lakes on the ice shelf are commonly more transient and appear in different locations each year, some larger lakes on
the eastern portion of the ice shelf and on Apfel Glacier have a high tendency to reform annually in the same locations (Fig.
2b and 2c). This characteristic has not previously been reported, and we suggest this occurs because these lakes are located on
very slow-moving ice near the grounding line ($<50 \text{ m yr}^{-1}$) where annual ice advection is minimal. Between 2000-2020, SGLs
on the ice shelf are only slightly shallower than those upstream of the grounding line, as has been recorded on Petermann
385 Glacier in Greenland (Supplementary Fig. 4a-b; MacDonald et al., 2018). Furthermore, in years with higher meltwater
volumes, lakes grew deeper on the ice shelf than on grounded ice (Supplementary Fig. 4c-d). This is perhaps surprising as
slow ice flow produces low-amplitude ice surface topography and shallower depressions, so we would expect grounded lakes
to be deeper (Pope et al., 2016). We suggest greater numbers of deeper structural depressions form in the ice shelf grounding
zone where there is an extensional flow regime and very low slopes conducive to meltwater ponding (Fig. 3b). Inter-annual
390 lake deepening amplified by lake bottom ablation could also mean lakes on the ice shelf evolve to be deeper than those on ice
upstream of the grounding line (Banwell and MacAyeal, 2015). We do not observe inter-annual deepening of individual lakes
in our study.

The spatial distribution of SGLs mapped in this study closely follows predicted meltwater routing pathways derived from a
395 surface DEM, which assumes widespread firm saturation across the ice shelf (Supplementary Fig. 6). Hydrological routing
analysis shows meltwater coalesces from steeper upstream parts of the ice shelf and converges in structurally-determined
topographic lows, such as flow stripes and crevasses on faster-flowing ice, downstream of ice rises, and in shear margins
(Supplementary Fig. 6). This could have future implications for ice shelf stability, though we note that our routing analysis
does not allow for water storage in the firm (Sect. 5.3).

400 **5.2 Seasonal evolution of SGLs**

SGLs on Shackleton Ice Shelf exhibit strong intra-seasonal and inter-seasonal variations in area, depth and volume, as do
studies on Langhovde Glacier and the Nansen and Amery ice shelves (Bell et al., 2017; Langley et al., 2016; Moussavi et al.,
2020). In melt seasons when lakes store larger total volumes of surface meltwater on Shackleton Ice Shelf, lakes are more
extensive and some are fed by supraglacial rivers, which are not present in lower-melt years (Fig. 4, Fig. 6g-j). Notably, we
405 find a correspondence between short-lived, high magnitude modelled snowmelt events and peak total lake area and volume
during years of high surface meltwater storage (i.e. the melt seasons with maximum lake area, depth and volume: 2013-2014,
2014-2015, 2016-2017 and 2017-18) (Fig. 9). These spikes in modelled snowmelt could represent localised katabatic wind-
induced melt events, similar to föhn wind-induced melt events on the Antarctic Peninsula modelled by regional climate models
(Kuipers Munneke et al., 2018; Tuckett et al., 2019). Föhn wind events have been shown to increase surface melt intensity and
410 control the large interannual temporal and spatial variability in surface melt on the Antarctic Peninsula (Cape et al., 2015;

Turton et al., in review). Although 2019-2020 modelled surface melt data is currently unavailable for us to compare with total lake area and volume, we note that this was an exceptional melt season (Fettweis, 2020) in which we record the highest total lake area and volume over the 2000-2020 period (Fig. 9e). Previous work based on meteorological observations from Ekström Ice Shelf, East Antarctica, has suggested that snowmelt-albedo feedbacks exert a key control on melt intensity and duration (Jakobs et al., 2019). Therefore, variations in the strength of this snowmelt-albedo feedback could exert an important control on the amount of melt seasonally available to pond into SGLs on Shackleton Ice Shelf.

Interestingly, the years when total lake area, depth and volume were highest do not always correspond with warmer summer near-surface temperatures (Fig. 7). In particular, mean DJF temperatures during years with the highest surface meltwater storage were actually lower than some years with much less extensive and shallower lakes (Table 2). Furthermore, the weak positive relationship we observe between total SGL area and mean surface air temperature in the preceding part of the melt season (Fig. 8) contrasts with previous studies which show SGLs are highly sensitive to small fluctuations in surface temperature (e.g. Langley et al., 2016). Rather, we suggest that seasonal fluctuations in lake area and volume on this ice shelf are more sensitive to snowmelt intensity associated with katabatic wind-driven melting, rather than longer-term patterns of mean near-surface summer temperature. Therefore, intense short-lived melt events appear to determine seasonal variability in SGL extent and volume. This is important because short-lived katabatic events and albedo feedbacks are difficult to resolve in regional climate models such as RACMO, owing to its 27-km resolution (Van Wessem et al., 2018). Although RACMO2.3p1 output is available at 5.5 km resolution in some regions of Antarctica, it does not yet extend to our study region. This finding underscores the need for finer resolution regional climate model simulations of surface melt when considering future lake distributions (Arthur et al., 2020; Stokes et al., 2019). The weak relationship that we observe between SGL extents and near-surface temperature is perhaps unsurprising given the resolution of ERA5 is insufficient to resolve localised mixing and albedo-lowering processes around the ice shelf grounding zone, but our preliminary comparison between SGL development and climate offers some clear hints that short-lived melt events are more important than longer-term seasonal conditions.

Regarding individual lake development, the lakes that we observed to drain englacially within a ≤ 7 -day period did so more rapidly than lakes on the Amery Ice Shelf where lakes have drained within 10 days (Moussavi et al., 2020). This is still slower than on the Greenland Ice Sheet, where lakes drain on hourly-daily timescales (e.g. Das et al., 2008; Doyle et al., 2013; Selmes et al., 2013; Stevens et al., 2015; Williamson et al., 2017). We note that the lake drainages we observed could have drained on sub-daily timescales comparable to Greenland, but this is not detectable from the temporal resolution of the satellite imagery used in this study. Ice shelf thicknesses in the region where lakes drained englacially (Fig. 6k-l) reach 150 - 200 m, which is comparable to the ~ 200 m thickness of Larsen B through which lakes are thought to have drained prior to its collapse (Banwell et al., 2013; MacAyeal and Sergienko 2013; Scambos et al., 2003). Therefore, lakes could potentially have drained through the full ice shelf thickness via lake bed crevasses. However, because no lakes drained fully, we suggest they are more likely to have drained into the firn and subsequently refrozen.

Most lakes do not drain and instead refreeze at the end of the melt season, which is consistent with the decline in total lake area and volume from late-January to late-February (Fig. 7) and is common elsewhere in East Antarctica (e.g. Langley et al., 2016; Leppäranta et al., 2013). Once meltwater penetrates the snowpack and reaches a sub-freezing firn temperature, it refreezes and raises the surrounding firn temperature by releasing latent heat (Buzzard et al., 2018; Bevan et al., 2018; Jakobs et al., 2019). Meltwater refreezing within the firn produces larger snow grains with a lower albedo, which are more likely to absorb incoming solar radiation and further enhance surface melting (Gardner and Sharp 2010). Thus, our observations of lakes repeatedly refreezing at the end of each melt season are likely to indicate meltwater storage and refreezing in the ice shelf firn layer near the grounding line.

5.3 Implications for ice shelf stability

Future warming is expected to enhance surface melting and increase surface meltwater production in Antarctica (Bell et al., 2018; Kingslake et al., 2017; Rintoul et al., 2018; Trusel et al., 2015). Increased lake coverage and meltwater storage in future years could cause the ice shelf to flex (Banwell and MacAyeal 2015; Banwell et al., 2019), which could lead to fracture. Previous work has suggested that FAC can be used to infer shelf stability and likelihood of hydrofracture (Alley et al., 2018; Holland et al., 2011; Lenaerts et al., 2017). We find simulated minimum FAC is close to zero in the grounding zone of Shackleton Ice Shelf, consistent with simulated FAC on the Roi Baudouin (Lenaerts et al., 2017) and pre-collapse Larsen B (Leeson et al., 200) ice shelves. Therefore, despite limited (<1 %) lake coverage on the ice shelf at present, its susceptibility to widespread ponding and hydrofracture is likely to gradually increase as FAC is depleted (Alley et al., 2018; Leeson et al., 2020; Lenaerts et al., 2017). Our meltwater routing analysis suggests that the ice shelf could potentially support and route water along surface troughs to the ice shelf edge, similar to active shear-margin rivers present on Nansen Ice Shelf (Supplementary Fig. 6; Bell et al., 2017), which could partially mitigate the effect of surface meltwater loading (Bell et al., 2017).

6 Conclusions

This study presents the first quantitative, multiyear study of SGL evolution on East Antarctica's sixth largest ice shelf. Between the melt seasons of 1999-2000 and 2019-2020, melt seasons were characterised by hundreds of SGLs that were, on average, 0.02 km² in area, 0.96 m deep, and held a mean total meltwater volume of 7.45 x10⁶ m³.

We also find that:

- Lakes cluster around the grounding line, controlled by localised albedo-lowering feedbacks caused by interactions between katabatic-driven melting, blue ice and exposed bedrock (Kingslake et al., 2017; Lenaerts et al., 2017; Stokes et al., 2019).

- 475
- Lakes on the ice shelf are more transient than those on grounded ice, which reform annually in the same locations. Some lakes on slow-moving parts of the ice shelf near the grounding line also frequently occupy the same ice surface depressions, which likely results from minimal ice advection in these areas.
 - Lakes begin to form from mid-November and their total volume peaks in early-late January each year, which reached a maximum of $28.4 \times 10^6 \text{ m}^3$ on 31st January 2020.
- 480
- SGLs were most extensive, deepest, and formed at the highest recorded elevation (527 m) in 2014, 2015, 2017, 2018 and 2020. At the peak of these melt seasons, parts of the ice shelf grounding zone became saturated with meltwater which coalesced into large lakes, and lakes were fed by supraglacial rivers.
 - Short-lived high magnitude snowmelt events correspond with peak total lake area and volume during these years of high surface meltwater storage. Seasonal variability in lake extents is more sensitive to the intensity of individual
- 485
- melt events rather than mean near-surface summer temperatures, which we suggest is driven by localised katabatic wind-induced melting.

Our findings may be used to inform the boundary conditions and validation of supraglacial hydrology models, for example to validate estimates of lake distributions and to constrain volume-dependent lake drainage thresholds. In particular, our data

490

quantify the meltwater volumes being stored on the ice shelf surface, highlighting the importance of increased meltwater storage within SGLs during melt seasons experiencing intense, short-lived melt events. An improved understanding of East Antarctica's present surface hydrology has significance for constraining future dynamic change of the ice sheet and its contribution to sea-level rise.

495








Figure 1. Overview map of Shackleton Ice Shelf with LIMA mosaic (Bindschadler et al. 2008) in the background. The grounding line (thick black line) is taken from Rignot *et al.* (2016). The subset area used for detailed analysis of lake area and depth extractions is outlined in green and the black dashed box outlines the area used to extract ERA5 reanalysis near-surface air temperature. The light blue areas visible on the ice sheet are blue-ice areas.



535 **Figure 2. Recurrence frequency of lakes mapped between 1974 and 2020 on Shackleton Ice Shelf out of a total of 58 individual satellite scenes. Surface melt flux derived from QuickScat scatterometer (2000 - 2009 average from Trusel *et al.* (2013)) on the portion of the ice shelf where lakes predominantly form (a). Lakes are concentrated in the grounding zone downstream of blue ice and adjacent to exposed rock (b and c). Lakes on the ice shelf itself are short-lived in a given location (d) and migrate down-shelf with ice flow over multiple melt seasons (f). Lakes on grounded ice upstream of the grounding line typically occupy fixed ice surface depressions and therefore frequently reform in the same locations during multiple melt seasons (e). Note the scale bar frequency corresponds to number of times a lake was recorded in the available cloud-free satellite imagery, rather than the number of years over this time period.**
540

545

550




555

Figure 3. Frequency distributions of supraglacial lakes on Shackleton Ice Shelf by topographic variables: (a) individual lake mean elevations; (b) ice surface slope of individual lakes; (c) ice flow speed; (d) distance of each lake to nearest exposed bedrock. The four panels show a subset of lakes (in blue) on the ice shelf and their spatial association with low elevations, low surface slopes low ice flow speeds and exposed rock/ blue ice. The grounding line is represented by the dashed black line.


560

565




570

575



580

585




590

595


Figure 4. Evolution of total lake area, depth and volume across Shackleton Ice Shelf over the 2000-2020 melt seasons. The stem plots present time series of meltwater area (a), depth (b) and volumes (c) of lakes inside the subset area (green box in Figure 1), where each point and stem represent a single date of lake observations. Grey bars indicate the percentage of cloud cover within this subset area. Full lake area, depth and volume data for individual years is included in Table 1.

600

605



615 **Figure 5.** Total lake volume anomaly relative to long-term (2000-2020) mean (a) and total lake area anomaly relative to long-term (2000-2020) mean (b). Dashed black lines represent one, two and three standard deviations from the long-term mean.



620 **Figure 6. Examples of lake drainage and refreeze-burial events on Shackleton Ice Shelf: lakes refreezing and becoming buried by snow (a-c); a lake draining via a supraglacial stream (d-f); supraglacial streams/rivers feeding lakes (g); a lake before and after through-ice drainage (h-j); and a chain of lakes partially draining in ≤ 7 days (k-l). Background images are Landsat 8 (a-c) and Sentinel-2A (d-l).**

625

630

635

640

645

650







Figure 7. Total supraglacial lake area and mean near-surface (2 m) temperature for 1999 – 2004 (a), 2010 – 2015 (b) and 2015 – 2020 (c). Mean near-surface temperature is the mean extracted over the whole Shackleton Ice Shelf (shown as a 5-day moving average).






655

660

665




670 Figure 8. Scatter plot of total lake area inside the subset area highlighted in Figure 1 (green box) and mean surface air temperature from 1st November to the date of each observation.



675

680

Figure 9. Seasonal variations in total lake area, volume and modelled snowmelt on Shackleton Ice Shelf during the melt seasons 2013/14 (a), 2014/15 (b), 2016/17 (c), 2017/18 (d) and 2019/20 (e) within the subset area (green box, Figure 1). Peak total lake area and volume is associated with a spike in snowmelt. Snowmelt rates are mean values over the entire ice shelf.



690 Figure 10. (a) Mean firn air content (total integrated pore space, i.e. the thickness of the equivalent air column contained in the firn in metres) over Shackleton Ice Shelf from 2000-2019. (b) Scatterplot comparing minimum firn air content and total lake area. (c) Time series comparing mean firn air content, minimum firn air content and total lake area.

Table 1. Total lake area and volume within the 48 x 48 km² subset area on Shackleton Ice Shelf.

Date	Lake depth (m)			Lake area (km²)			Lake volume (x10⁶ m³)	
	Median	Mean	Maximum	Mean	Median	Maximum	Total	Total
6th Feb 2000	0.69	0.69	1.52	0.014	0.004	0.203	1.19	0.85
8th Feb 2004	0.5	0.55	2.17	0.028	0.004	0.986	13.47	8.64
27th Jan 2008	0.78	0.78	3.27	0.022	0.008	0.459	4.09	3.48
17th Feb 2010	0.59	0.68	2.51	0.028	0.007	0.364	3.71	2.34
24th Feb 2010	0.48	0.49	0.95	0.026	0.007	0.376	5.88	3.20
4th Feb 2011	0.71	0.77	2.47	0.031	0.009	0.359	6.26	4.95
23rd Feb 2012	0.57	0.57	2.21	0.015	0.004	0.139	1.33	0.72
2nd Dec 2013	0.96	0.98	3.58	0.004	0.001	0.178	3.04	2.82
3rd Jan 2014	1.80	1.78	6.56	0.026	0.002	1.867	16.45	33.94
19th Nov 2014	0.70	0.69	1.00	0.031	0.007	0.171	0.56	0.41
6th Jan 2015	1.72	1.78	6.46	0.045	0.005	6.717	18.37	31.50
7th Feb 2015	1.57	1.51	3.13	0.017	0.002	0.949	3.38	4.67
23rd Feb 2015	0.89	0.98	2.09	0.044	0.016	0.196	1.19	1.07
9th Jan 2016	1.77	1.87	5.72	0.006	0.001	0.182	1.57	2.93
25th Jan 2016	1.84	1.91	5.61	0.009	0.003	0.223	2.19	4.38
26th Feb 2016	1.16	1.25	3.74	0.012	0.002	0.277	3.13	3.62
27th Jan 2017	1.35	1.44	3.99	0.019	0.002	1.464	10.29	16.35
31st Jan 2017	0.63	0.71	3.09	0.020	0.002	1.573	8.60	7.19
28th Feb 2017	0.46	0.58	2.23	0.041	0.003	0.477	1.71	0.89
29th Dec 2017	1.52	1.66	3.41	0.002	0.001	0.022	0.16	0.26
19th Jan 2018	0.97	0.61	1.05	0.007	0.001	0.684	12.48	15.25
26th Jan 2018	0.86	0.90	3.89	0.009	0.001	0.729	9.66	9.41
8th Feb 2018	0.46	0.52	3.02	0.008	0.001	0.876	4.34	2.26
15th Feb 2018	1.03	1.29	4.47	0.017	0.003	0.081	0.20	0.43
29th Jan 2019	1.17	1.21	2.69	0.019	0.003	0.656	5.24	6.36
18th Feb 2019	0.53	0.52	1.58	0.031	0.001	1.50	7.54	4.48
28th Feb 2019	0.52	0.53	1.87	0.047	0.008	1.494	6.18	3.45

19th Dec 2019	0.59	0.88	2.15	0.023	0.011	0.522	4.63	2.89
1st Jan 2020	0.91	0.95	5.23	0.012	0.001	0.954	12.12	14.22
11th Jan 2020	0.71	0.73	3.47	0.019	0.001	1.627	17.25	16.48
31st Jan 2020	0.69	0.76	2.94	0.045	0.001	11.961	28.55	28.44
28th Feb 2020	0.41	0.41	0.60	0.082	0.008	3.204	14.32	0.67

695

Table 2. Average December-January-February (DJF) near-surface temperatures from ERA5 on Shackleton Ice Shelf from 2000-2020.

Melt season	DJF Temperature (°C)	Melt season	DJF Temperature (°C)
1999-2000	-10.0	2009-2010	-8.9
2000-2001	-7.9	2010-2011	-9.4
2001-2002	-7.5	2011-2012	-10.3
2002-2003	-8.1	2012-2013	-9.6
2003-2004	-7.2	2013-2014	-8.6
2004-2005	-7.1	2014-2015	-8.4
2005-2006	-8.1	2015-2016	-10.2
2006-2007	-7.9	2016-2017	-6.4
2007-2008	-10.6	2017-2018	-8.6
2008-2009	-8.0	2018-2019	-8.3
		2019-2020	-9.3

700

705

Data Availability

The mapped supraglacial lake polygons are available as digital GIS shapefiles (.shp) in Supplementary Data. Sentinel imagery is available from the Copernicus Open Access Hub (<https://scihub.copernicus.eu>) and Landsat scenes are available from the United States Geological Survey EarthExplorer (<https://earthexplorer.usgs.gov/>). ERA5 reanalysis surface temperature is available from the Copernicus Climate Change Service Climate Data Store (<https://cds.climate.copernicus.eu/>). We acknowledge the Norwegian Polar Institute's Quantarctica package.

Author contribution

JFA, CRS and SSJ designed the initial study. JFA undertook the data collection and conducted the analysis, with guidance from all authors. JFA led the manuscript writing, with input from all authors.

Competing Interests

The authors declare that they have no conflict of interest.

Acknowledgements

JFA was funded by the IAPETUS Natural Environment Research Council (NERC) Doctoral Training Partnership (grant number NE/L002590/1). CRS and SSRJ were supported by NERC grant NE/R000824/1. We acknowledge RACMO data from M.R van den Broeke, the Norwegian Polar Institute's Quantarctica package, and Copernicus Climate Change Service Information (ERA5 reanalysis). JFA acknowledges BWJ Miles for helpful discussions during this work. We would like to thank Jan Lenaerts and an anonymous reviewer for providing constructive comments which led to improvement of this manuscript, and the editor (Stef Lhermitte) for handling the manuscript.

References

- Alley, K.E., Scambos, T.A., Anderson, R.S., Rajaram, H.: Continent-wide estimates of Antarctic strain rates from Landsat 8-derived velocity grids, *J. Glaciology*, 64, 1–12, doi:10.1017/jog.2018.23, 2018.
- Alley, K.E., Scambos, T.A., Miller, J.Z., Long, D.G., MacFerrin, M.: Quantifying vulnerability of Antarctic ice shelves to hydrofracture using microwave scattering properties. *Remote. Sens. Environ.* 210, doi:10.1016/j.rse.2018.03.025, 2018.
- Arthur, J.F., Stokes, C.R. Jamieson, S.S.R., Carr, J.R., Leeson, A.A.: Recent understanding of Antarctic supraglacial lakes using satellite remote sensing, *Prog. Phys. Geogr.*, 1-32, doi:tbc, 2020.

- Banwell, A.F., Caballero, M., Arnold, N.S., Glasser, N.F., Cathles, L.M., MacAyeal, D.R.: Supraglacial lakes on the Larsen B ice shelf, Antarctica, and at Paakitsoq, West Greenland: A comparative study, *Ann. Glaciol*, 55, doi:10.3189/2014AoG66A049, 2014.
- Banwell, A.F. and MacAyeal, D.R. (2015) Ice-shelf fracture due to viscoelastic flexure stress induced by fill/drain cycles of supraglacial lakes, *Antarct. Sci*, 27, doi:10.1017/S0954102015000292, 2015.
- Banwell, A.F., MacAyeal, D.R. and Sergienko, O. V.: Breakup of the Larsen B Ice Shelf triggered by chain reaction drainage of supraglacial lakes, *Geophys. Res. Lett.*, 40, doi:10.1002/2013GL057694, 2015.
- Banwell, A.F., Willis, I.C., Macdonald, G.J., Goodsell, B., MacAyeal, D.R.: Direct measurements of ice-shelf flexure caused by surface meltwater ponding and drainage, *Nat. Commun.*, 10, doi:10.1038/s41467-019-08522-5, 2019.
- Bell, R.E., Banwell, A.F., Trusel, L.D., Kingslake, J.: Antarctic surface hydrology and impacts on ice-sheet mass balance, *Nat. Clim. Change*, 8, doi:10.1038/s41558-018-0326-3, 2018.
- Bell, R.E., Chu, W., Kingslake, J., Das, I., Tedesco, M., Tinto, K.J., Zappa, C.J., Frezzotti, M., Boghosian, A., Sang Lee, W.: Antarctic ice shelf potentially stabilized by export of meltwater in surface river, *Nature*, 544, doi:10.1038/nature22048, 2017.
- Bevan, S.L., Luckman, A.J., Kuipers Munneke, P., Hubbard, B., Kulessa, B., Ashmore, D.W.: Decline in Surface Melt Duration on Larsen C Ice Shelf Revealed by The Advanced Scatterometer (ASCAT), *Earth Space Sci*, 5, doi:10.1029/2018EA000421, 2018.
- Box, J.E. and Ski, K.: Remote sounding of Greenland supraglacial melt lakes: Implications for subglacial hydraulics, *J. Glaciology*, 53, doi:10.3189/172756507782202883, 2007.
- Burton-Johnson, A., Black, M., Peter, T.F., Kaluza-Gilbert, J.: An automated methodology for differentiating rock from snow, clouds and sea in Antarctica from Landsat 8 imagery: A new rock outcrop map and area estimation for the entire Antarctic continent, *Cryosphere*, 10, doi:10.5194/tc-10-1665-2016, 2016.
- Buzzard, S.C., Feltham, D.L. and Flocco, D.: A Mathematical Model of Melt Lake Development on an Ice Shelf, *J. Adv. Model. Earth Sys*, 10, doi:10.1002/2017MS001155, 2018.
- Das, S.B., Joughin, I., Behn, M.D., Howat, I.M., King, M.A., Lizarralde, D., Bhatia, M.P.: Fracture propagation to the base of the Greenland ice sheet during supraglacial lake drainage, *Science*, 320, doi:10.1126/science.1153360, 2008.
- Dell, R., Arnold, N., Willis, I., Banwell, A., Williamson, A., Pritchard, H.: Lateral meltwater transfer across an Antarctic Ice Shelf, *Cryosphere*, doi: 10.5194/tc-2019-316, in review.
- Doran, P.T., McKay, C.P., Meyer, M.A., Andersen, D.T., Wharton, R.A., Hastings, J.T.: Climatology and implications for perennial lake ice occurrence at Bunger Hills Oasis, East Antarctica, *Antarct. Sci*. 8, doi:10.1017/s0954102096000429, 1996.
- Doyle, S.H., Hubbard, A.L., Dow, C.F., Jones, G.A., Fitzpatrick, A., Gusmeroli, A., Kulessa, B., Lindback, K., Pettersson, R., Box, J.E.: Ice tectonic deformation during the rapid in situ drainage of a supraglacial lake on the Greenland Ice Sheet, *Cryosphere*, 7, doi:10.5194/tc-7-129-2013, 2013.
- Dell, R., Arnold, N., Willis, I., Banwell, A., Williamson, A., Pritchard, H.: Lateral meltwater transfer across an Antarctic ice shelf, *Cryosphere*, doi: 10.5194/tc-2019-316, in review.

- Echelmeyer, K., Clarke, T.S. and Harrison, W.D.: Surficial glaciology of Jakobshavns Isbræ, West Greenland: Part I. Surface morphology, *J. Glaciol.*, 37, doi:10.3189/S0022143000005803, 1991.
- 770 DLR.: Grounding line for Denman Glacier, derived from ERS-1/2 and Copernicus Sentinel-1 data acquired in 1996, 2015 and 2017, Antarctic Ice Sheet CCI [release date 2018-07-02, v1.0], 2018.
- Fettweis, X.: 5-day forecast of the 2019-2020 Antarctic ice sheet SMB simulated by MARv3.11 forced by GFS: https://climato.uliege.be/cms/c_5652669/fr/climato-antarctica, last access: 17 April 2020.
- Flament, T. and Rémy, F.: Dynamic thinning of Antarctic glaciers from along-track repeat radar altimetry, *J. Glaciol.*, 58, doi:10.3189/2012JoG11J118, 2012.
- 775 Fretwell, P., Pritchard, H.D., Vaughan, D.G., et al.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica P. Fretwell et al.: Improved ice bed, surface and thickness datasets for Antarctica, *Cryosphere*, doi:10.5194/tc-7-375-2013, 2013.
- Fürst, J.J., Durand, G., Gillet-Chaulet, F., Tavard, L., Rankl, M., Braun, M., Gagliardini, O.: The safety band of Antarctic ice shelves, *Nat. Clim. Change*, 6, doi:10.1038/nclimate2912, 2016.
- 780 Gardner, A.S. and Sharp, M.J.: A review of snow and ice albedo and the development of a new physically based broadband albedo parameterization, *J. Geophys. Res. Earth Surf.*, 115, doi:10.1029/2009JF001444, 2010.
- Gillespie, A.R., Kahle, A.B. and Walker, R.E.: Color enhancement of highly correlated images. II. Channel ratio and 'chromaticity' transformation techniques, *Remote Sens. Environ.*, 22, doi:10.1016/0034-4257(87)90088-5, 1987.
- 785 Glasser, N.F. and Gudmundsson, G.H.: Longitudinal surface structures (flowstripes) on Antarctic glaciers, *Cryosphere*, 6: doi:10.5194/tc-6-383-2012, 2012.
- Glasser, N. and Scambos, T.: A structural glaciological analysis of the 2002 Larsen B ice-shelf collapse, *J. Glaciol.*, 54, doi:10.1136/archdischild-2015-309290, 2008.
- Holland, P.R., Corr, H.F.J., Pritchard, H.D., Vaughan, D.G., Arthern, R.J., Jenkins, A., Tedesco, M.: The air content of Larsen Ice Shelf. *Geophys. Res. Lett.*, 38, doi:10.1029/2011GL047245, 2011.
- 790 Hubbard, B., Luckman, A., Ashmore, D., Bevan, S., Kulesa, B., Kuipers, Munneke, P., Philippe, M., Jansen, D., Booth, A., Sevestre, H., Tison, J., O'Leary, M., Rutt, I.: Massive subsurface ice formed by refreezing of ice-shelf melt ponds, *Nature Comms.*, 7, doi:10.1038/ncomms11897, 2016.
- Jakobs, C.L., Reijmer, C.H., Kuipers Munneke, P., König-Langlo, G., van den Broeke, M.R.: Quantifying the snowmelt-albedo feedback at Neumayer Station, East Antarctica, *Cryosphere*, 13, doi:10.5194/tc-13-1473-2019, 2019.
- 795 Kingslake, J., Ely, J.C., Das, I., Bell, R.E.: Widespread movement of meltwater onto and across Antarctic ice shelves, *Nature*, 544, doi:10.1038/nature22049, 2017.
- Kingslake, J., Ng, F. and Sole, A.: Modelling channelized surface drainage of supraglacial lakes, *J. Glaciol.*, 61, doi:10.3189/2015JoG14J158, 2015.
- 800 Konrad, H., Shepherd, A., Gilbert, L., Hogg, A.E., McMillan, M., Muir, A., Slater, T.: Net retreat of Antarctic glacier grounding lines, *Nat. Geosci.*, 11, doi:10.1038/s41561-018-0082-z, 2018.

- Kuipers Munneke, P., Luckman, A.J., Bevan, S.L., Smeets, C.J.P.P., Gilbert, E., van den Broeke, M.R., Wang, W., Zender, C., Hubbard, B., Ashmore, D., Orr, A., King, J.C., Kulesa, B.: Intense Winter surface melting on an Antarctic Ice Shelf, *Geophys. Res. Lett.*, 45, doi: 10.1029/2018GL077899, 2018.
- 805 Lampkin, D.J. and VanderBerg, J.: A preliminary investigation of the influence of basal and surface topography on supraglacial lake distribution near Jakobshavn Isbrae, western Greenland, *Hydrol. Process.*, 25, doi:10.1002/hyp.8170, 2011.
- Langley, E.S., Leeson, A.A., Stokes, C.R., Jamieson, S.S.R.: Seasonal evolution of supraglacial lakes on an East Antarctic outlet glacier, *Geophys. Res. Lett.*, 43, doi:10.1002/2016GL069511, 2016.
- Leeson, A.A., Forster, E., Rice, A., Gourmelen, N., van Wessem, J.M.: Evolution of supraglacial lakes on the Larsen B ice
810 shelf in the decades before it collapsed, *Geophys. Res. Lett.*, 47, doi:10.1029/2019GL085591, 2020.
- Leeson, A.A., Shepherd, A., Palmer, S., Sundal, A., Fettweis, X.: Simulating the growth of supraglacial lakes at the western margin of the Greenland ice sheet, *Cryosphere*, 6, doi:10.5194/tc-6-1077-2012, 2012.
- Lenaerts, J.T.M., Lhermitte, S., Drews, R., Lightenberg, S.R.M., Berger, S., Helm, V., Smeets, C.J.P.P., van den Broeke, M.R., van de Ber, W.J., van Meijgaard, E., Eijkelboom, M., Eisen, O., Pattyn, F.: Meltwater produced by wind-albedo interaction
815 stored in an East Antarctic ice shelf, *Nat. Clim. Change*, 7, doi:10.1038/nclimate3180, 2017.
- Leppäranta, M., Järvinen, O. and Mattila, O.P.: Structure and life cycle of supraglacial lakes in Dronning Maud Land, *Antarct. Sci.*, 25, doi:10.1017/S0954102012001009, 2013.
- Liang, Q.I., Zhou, C., Howat, I.M., Jeong, S., Liu, R., Chen, Y.: Ice flow variations at Polar Record Glacier, East Antarctica, *J. Glaciol.*, 65, doi:10.1017/jog.2019.6, 2019.
- 820 Luckman, A., Elvidge, A., Jansen, D., Kulesa, B., Kuipers Munneke, P., King, J., Barrand, N.E.: Surface melt and ponding on Larsen C Ice Shelf and the impact of föhn winds. *Antarct. Sci.*, 26, doi:10.1017/S0954102014000339, 2014.
- Macayeal, D.R. and Sergienko, O. V.: The flexural dynamics of melting ice shelves, *Ann. Glaciol.*, 54, doi:10.3189/2013AoG63A256, 2013.
- MacDonald, G.J., Banwell, A.F. and MacAyeal, D.R.: Seasonal evolution of supraglacial lakes on a floating ice tongue,
825 Petermann Glacier, Greenland, *Ann. Glaciol.*, 59, doi: doi:10.1017/aog.2018.9, 2018.
- Moran, M.S., Jackson, R.D., Slater, P.N., Teillet, P.M.: (1992) Evaluation of simplified procedures for retrieval of land surface reflectance factors from satellite sensor output, *Remote Sens. Environ.*, 41, doi:10.1016/0034-4257(92)90076-V, 1992.
- Morlighem, M., Rignot, E., Binder, T., et al.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nat. Geosci.*, 13, doi:10.1038/s41561-019-0510-8, 2019.
- 830 Moussavi, M.S., Pope, A., Halberstadt, A.R.W., Trusel, L.D., Cioffi, L., Abdalati, W.: Antarctic supraglacial lake detection using Landsat 8 and Sentinel-2 imagery: towards continental generation of lake volumes, *Remote Sens.*, 12, doi: 10.3390/rs12010134, 2020.
- Philpot, W.D.: Bathymetric mapping with passive multispectral imagery, *Applied Optics*, 28, doi:10.1364/ao.28.001569, 1989.
- Pope, A.: Reproducibly estimating and evaluating supraglacial lake depth with Landsat 8 and other multispectral sensors, *Earth
835 Space Sci.*, 3, doi:10.1002/2015EA000125, 2016.

- Pope, A., Scambos, T.A., Moussavi, M., Tedesco, M., Willis, M., Shean, D., Grigsby, S.: Estimating supraglacial lake depth in West Greenland using Landsat 8 and comparison with other multispectral methods, *Cryosphere*, 10, doi:10.5194/tc-10-15-2016, 2016.
- 840 Pope, R.M. and Fry, E.S.: Absorption spectrum (380–700 nm) of pure water. II. Integrating cavity measurements, *Appl. Opt.*, 36, doi: 10.1364/AO.36.008710, 1987.
- Pritchard, H.D., Ligtenberg, S.R.M., Fricker, H.A., Vaughan, D.G., van den Broeke, M.R., Padman, L.: Antarctic ice-sheet loss driven by basal melting of ice shelves, *Nature*, 484, doi:10.1038/nature10968, 2012.
- Reynolds, R.W. and Smith, T.M.: Lakes on George VI Ice Shelf, *Polar Rec.*, 20, doi: 10.1017/S0032247400003636, 1981.
- 845 Rignot, E., Jacobs, S., Mouginot, J., Scheuchl, B.: Ice-shelf melting around Antarctica, *Science*, 341, doi:10.1126/science.1235798, 2013.
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M.J., Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *PNAS*, 116, doi:10.1073/pnas.1812883116, 2019.
- Rintoul, S.R., Chown, S.L., DeConto, R.M., England, M.H., Fricker, H.A., Masson-Delmotte, V., Naish, T.R., Siegert, M.J., Xavier, J.C.: Choosing the future of Antarctica, *Nature*, 558, doi:10.1038/s41586-018-0173-4, 2018.
- 850 Robel, A. and Banwell, A.F.: A speed limit on ice shelf collapse through hydrofracture, *Geophys. Res. Lett.*, 46, doi:10.1029/2019GL084397, 2019.
- Roberts, J.L., Warner, R.C., Young, D., Wright, A., van Ommen, T.D., Blankenship, D.D., Siegert, M., Young, N.W., Tarbacco, I.E., Forieri, A., Passerini, A., Zirizzotti, A., Frezzotti, M.: Refined broad-scale sub-glacial morphology of Aurora Subglacial Basin, East Antarctica derived by an ice-dynamics-based interpolation scheme, *Cryosphere Discuss*, 5, doi: 855 10.5194/tc-5-551-2011, 2011.
- Scambos, T., Hulbe, C. and Fahnestock, M.: Climate-induced Ice Shelf Disintegration in the Antarctic Peninsula, *Ant. Pen. Clim. Var.: Hist Paleo. Persp.*, 79, doi:10.1029/AR079p0079, 2003.
- Selmes, N., Murray, T. and James, T.D.: Characterizing supraglacial lake drainage and freezing on the Greenland Ice Sheet, *Cryosphere Discuss.*, 7, doi:10.5194/tcd-7-475-2013, 2013.
- 860 Sneed, W.A. and Hamilton, G.S.: Evolution of melt pond volume on the surface of the Greenland Ice Sheet. *Geophys. Res. Lett.*, 34, doi: 10.1029/2006GL028697, 2007.
- Stevens, L.A., Behn, M.D., McGuire, J.J., Das, S.B., Joughin, I., Herring, T., Shean, D.E., King, M.A.: Greenland supraglacial lake drainages triggered by hydrologically induced basal slip, *Nature*, 522, doi:10.1038/nature14480, 2015.
- Stephenson, S.N., and Zwally, H.J.: Ice-shelf topography and structure determined using satellite-radar altimetry and Landsat 865 imagery, *Ann. Glaciol.*, 12, 1988.
- Stokes, C.R., Sanderson, J.E., Miles, B.W., Jamieson, S.S.R., Leeson, A.A.: Widespread development of supraglacial lakes around the margin of the East Antarctic Ice Sheet, *Sci. Rep.*, (9), doi: 10.1038/s41598-019-50343-5, 2019.
- Tarboton, D.G.: A New Method for Determination of Flow Directions and Upslope in Grid Elevation Models, *Water Resour. Res.*, 33, doi:10.1029/96WR03137, 1997.

- 870 Trusel, L.D., Frey, K.E., Das, S.B., Karnauskas, K.B., Kuipers Munneke, P.K., van Meijgaard, E., van den Broeke, M.R.: Divergent trajectories of Antarctic surface melt under two twenty-first-century climate scenarios, *Nat. Geosci.*, 8, doi:10.1038/ngeo2563, 2015.
- Trusel, L.D., Frey, K.E., Das, S.B., Kuipers Munneke, P., van den Broeke, M.R.: Satellite-based estimates of Antarctic surface meltwater fluxes, *Geophys. Res. Lett.*, 40, doi: 10.1002/2013GL058138, 2013.
- 875 Tuckett, P.A., Ely, J.C., Sole, A.J., Livingstone, S.J., Davison, B.J., van Wessem, J.M., Howard, J.: Rapid accelerations of Antarctic Peninsula outlet glaciers driven by surface melt, *Nat. Commun.*, 10, 2019.
- Wearing, M.G., Hindmarsh, R.C.A. and Worster, M.G.: Assessment of ice flow dynamics in the zone close to the calving front of Antarctic ice shelves, *J Glaciol.*, 61, doi: 10.3189/2015JoG15J116, 2015.
- Van Wessem, J., Van De Berg, W., Noël, B.P.Y., et al.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2 - Part 2: Antarctica (1979-2016), *Cryosphere*, 12, doi: 10.5194/tc-12-1479-2018, 2018.
- 880 Williamson, A.G., Arnold, N.S., Banwell, A.F., Willis, I.C.: A Fully Automated Supraglacial lake area and volume Tracking (“FAST”) algorithm: Development and application using MODIS imagery of West Greenland, *Remote Sens. Environ.*, 196, doi: 10.1016/j.rse.2017.04.032, 2017.
- Willis, I.C., Pope, E.L., Leysinger Vieli, G.J.M.C., Arnold, N.S., Long, S.: Drainage networks, lakes and water fluxes beneath the Antarctic ice sheet, *Ann. Glaciol.*, 57, doi: 10.1017/aog.2016.15, 2016.
- 885 Winther, J., Elvehøy, H., Bøggild, C.E., Sand, K., Liston, G.: Melting, runoff and the formation of frozen lakes in a mixed snow and blue-ice field in Dronning Maud Land, Antarctica, *J. Glaciol.*, 42, doi: 10.3189/S0022143000004135, 1996.
- Yang, K. and Smith, L.C.: Supraglacial streams on the Greenland ice sheet delineated from combined spectral-shape information in high-resolution satellite imagery, *IEEE Geosci. Rem. Sens. Lett.*, 10, doi: 10.1109/LGRS.2012.2224316, 2013.
- 890 Zheng, L. and Zhou, C.: Comparisons of snowmelt detected by microwave sensors on the Shackleton Ice Shelf, East Antarctica, *Int. J. Remote Sens.*, 41, doi:10.1080/01431161.2019.1666316, 2019.