

Dear Editor,

Thank you for the information about “publishing subject to minor revisions”. This response letter includes three answer-sections to the main comments of the editor, the reviewer 1 and the reviewer 2. Editorial comments regarding grammatical and language issues are not mentioned here, but are carefully inserted in the manuscript. Here, we address the main comments and how we answered to them.

Attached at this letter to the editor you will find the manuscript with tracked changes.

Answer to Editors comments:

Nature as a continuum couldn't be classified by clear boundaries, especially at our study-scale. So, we changed the main aim from the study to give a prediction of the lower permafrost (PF) limit to an assessment of the lower occurrence of probable permafrost. This change can be tracked through the manuscript by different sections. We put the assessment of the lower occurrence of probable PF in a better context of existing data (e.g. MAAT of the weather station of Zhadang). Two additional paragraphs regarding the atmospheric and topographic influence on PF-occurrence discuss the results of the PF-assessment in the discussion section with corresponding literature. Rockglaciers as permafrost indication can be used but carefully as the editor stated out. Therefore, we discussed corresponding literature and stated out that the rockglaciers in our study area are suitable for a zonal permafrost assessment. The reason for this lies in the low elevational range of these landforms. The distance between the rooting zone and the terminus is short both in length and elevation in comparison to other rockglaciers. So, the rockglaciers do not creep out the periglacial environment.

The uncertainty for displacement measurements were given by a sensitivity coefficient included by the co-author E. Reinosch based on his publication (<https://doi.org/10.5194/tc-14-1633-2020>).

Instead of a zoom in the permafrost map(s) for the investigated catchment in the Fig. 1, we add in the Fig. description: Note the proposed Permafrost occurrence of more than 90 % in study area Qugaqie Basin (black feature).

Answer to Reviewer 1 comments:

We agree that the term “probable permafrost” is not well defined, and that the lack of a precise definition might cause confusion. Therefore, we change the term to “probable occurrence of permafrost”. The use of the term “probable” is motivated by the fact that we do not have borehole-data (MAGST) and a modelled permafrost distribution and we can only assess the occurrence indirectly. The spatial heterogeneity of our data (mapping, InSar and ERT) and of natural variations in permafrost occurrence also prevents us from providing precise limits. This issue goes strongly in the direction to the editor's comment (see above).

We answer the reviewer's remaining question in the quantitative interpretation of the elevation (or elevation ranges) as inferred from each method in the final manuscript as follows. Each result section of the corresponding methods received a concluding sentence indicating the inferred elevation of probable permafrost occurrence.

One aspect of our study is that it provides localized ground-truth data in an area that is otherwise only accessible by simulation or theoretical considerations within large-scale models. Therefore, in order to compare the result with existing literature, we find it useful to infer one single value. In reality, the occurrence of permafrost naturally varies depending on topo-climatic factors (see editors comments). To take this into account, the single value is transformed into a range, which also includes the two lowest, active protalus ramparts. We state out as a general outcome of the publication “the probable occurrence permafrost is higher than 5300 - 5450 m. a.s.l”

The added values of the ERT-results, are that ERT supplements InSar-findings and vice versa. Surface creep without ice could happen without permafrost conditions and subsurface ice without creep over 2 consecutive years is not related to permafrost. With the help of ERT we were able to provide evidence for the existence of ground ice at specific test sites. The localized site surveys are critical for arriving at the 5300 – 5450 m estimate. Profile A (Fig. 8) ranges from 5090 to 5230m and no subsurface ice content was detected in the lateral moraine, which could be supposed.

Due to the reviewers aspect of a marginally supplement of InSAR-data, we believe that the surface movement, derived by InSAR, is one key parameter of periglacial landscapes (Eckerstorfer et al., 2018) and is used to infer permafrost conditions (Kneisel and Käb, 2007; Strozzi et al., 2004). Therefore, InSAR-data plays a major role in this study, because this data validates the two-years rule of frozen ground (=permafrost) through the creeping behavior (compare Fig. 10A).

A differentiation of seasonal and perennial frozen ground of the study area is given by Reinosch et al. (2020). Based on InSAR time series analyses of the typical periglacial surface changes the authors developed a model to distinguish annual surface changes (seasonal frozen ground) from perennial, more or less constant creeping surface changes (perennial frozen ground) for the study area. We focus directly on periglacial landforms according to perennial creep caused by down moving ice content.

Regarding moving the Table 2 and the corresponding text to section 3.2. is probably debatable but since the compiled resistivity values are results and are needed for the explanation of the ERT results, we would prefer to leave the table in section 4.2. We hope that the proximity of the table to the ERT discussion will help the reader to follow our arguments more easily. So, we will change the headline of Table 2: Resistivity values for different materials derived by field measurement. The used terms of the interpreted material followed Hauck and Kneisel (2008) and Mewes et al. (2017).

More details about the mapping procedure were mentioned in the method section.

Answer to Reviewer 2 comments:

As the reviewer 2 mentioned different geometries (i.e.ascending and descending) are used in the description of the Synthetic Aperture Radar Interferometry. We explained the use of different geometry in the corresponding method section. Both ascending (satellite travelling south to north) and descending (satellite travelling north to south) acquisitions are therefore sensitive to vertical surface displacement and towards the East or West respectively but very insensitive to displacement towards the North or

South. We always select the geometry with the highest sensitivity towards the expected displacement direction to calculate our displacement and velocity results. To that end we calculate a sensitivity coefficient for each pixel which is explained in the following. If both ascending and descending velocity LOS data is available for the same pixel, then we use only the geometry with the higher sensitivity coefficient, i.e. better sensitivity, to calculate the downslope velocity and ignore the other geometry to keep the precision of the projection as high as possible.

All observation periods of the shown surface velocity refer to the entire observation period of 2015 - 2018 but described different parts of the landforms. We made changes to the tables and text to describe the same parts of the landforms in both. We also added the observation period to a number of figure captions and paragraphs in the text to make it easier for the reader to follow.

We changed all units to mm/y for an easier comparability.

Now, the discussion chapter describes now a section about the uncertainty of the mapping procedure of the periglacial landforms which contains the IPA mapping guidelines and the important literature by Brardinoni et al. (2019).

Literature:

Brardinoni, F., Scotti, R., Sailer, R. and Mair, V.: Evaluating sources of uncertainty and variability in rock glacier inventories, *Earth Surf. Process. Landforms*, 44(12), 2450–2466, doi:10.1002/esp.4674, 2019.

Eckerstorfer, M., Eriksen, H. Ø., Rouyet, L., Christiansen, H. H., Lauknes, T. R. and Blikra, L. H.: Comparison of geomorphological field mapping and 2D-InSAR mapping of periglacial landscape activity at Nordnesfjellet, northern Norway, *Earth Surf. Process. Landforms*, 43(10), 2147–2156, doi:10.1002/esp.4380, 2018.

Kneisel, C. and Kääh, A.: Mountain permafrost dynamics within a recently exposed glacier forefield inferred by a combined geomorphological, geophysical and photogrammetrical approach, *Earth Surf. Process. Landforms*, 32(12), 1797–1810, doi:10.1002/esp.1488, 2007.

Mewes, B., Hilbich, C., Delaloye, R. and Hauck, C.: Resolution capacity of geophysical monitoring regarding permafrost degradation induced by hydrological processes, *Cryosphere*, 11(6), 2957–2974, doi:10.5194/tc-11-2957-2017, 2017.

Reinosch, E., Buckel, J., Dong, J., Gerke, M., Baade, J. and Riedel, B.: InSAR time series analysis of seasonal surface displacement dynamics on the Tibetan Plateau, *Cryosph.*, 14, 1633–1650, doi:10.5194/tc-2019-262, 2020.

Insights in a remote cryosphere: A multi method approach to assess permafrost occurrence at the Qugaie basin, western Nyainqêntanglha Range, Tibetan Plateau

5 Johannes Buckel¹, Eike Reinosch², Andreas Hördt¹, Fan Zhang³, Björn Riedel², Markus Gerke², Antje Schwalb⁴, Roland Mäusbacher⁵

¹Institute for Geophysics and extraterrestrial Physics, Technische Universität Braunschweig, Braunschweig, 38106, Germany

10 ²Institute for Geodesy and Photogrammetry, Technische Universität Braunschweig, Braunschweig, 38106, Germany

³Key Laboratory of Tibetan Environment Changes and Land Surface Processes, Institute of Tibetan Plateau Research, Chinese Academy of Sciences, Beijing, 100101, China.

⁴Institute of Geosystems and Bioindication, Technische Universität Braunschweig, Braunschweig, 38106, Germany

⁵Geographical Institute, Friedrich Schiller University of Jena, Jena, 07743, Germany

15

Correspondence to: Johannes Buckel (j.buckel@tu-braunschweig.de)

Abstract. Permafrost as a climate-sensitive parameter, its occurrence and distribution plays an important role in the observation of global warming. However, field-based permafrost distribution data and information on the subsurface ice content ~~at-in~~ the large area of the southern mountainous Tibetan Plateau (TP) is very sparse. Existing models based on
20 boreholes and remote sensing approaches suggest permafrost probabilities for most of the Tibetan mountain ranges. Field data to validate permafrost models are generally lacking because access of the mountain regions in extreme altitudes is limited. ~~Our~~The study provides geomorphological and geophysical field data from a north-orientated high-altitude catchment in the western Nyainqêntanglha Range. Our multi-method-approach combines (A) geomorphological mapping ~~-data~~, (B) subsurface ice ~~-content~~occurrence derived from electrical resistivity tomography (ERT) data, and (C) multi-annual
25 ~~displacement-creeping~~ rates from Interferometric Synthetic Aperture Radar (InSAR) analysis ~~allows an improved prediction~~to assess the lower occurrence of ~~catchment-wide~~the probable permafrost ~~occurrence~~. Our analysis indicates a lower boundary of permafrost occurrence ~~in a range of 5350 and 5500~~at around ~~5500-5400~~m a.s.l. in the Qugaie basin. Periglacial landforms such as rockglaciers and protalus ramparts are located in the periglacial zone ~~between~~from 5300 ~~and~~
5600 m a.s.l. ~~Therefore, our analysis indicates a lower boundary of permafrost occurrence at around 5500 m a.s.l.~~The
30 ~~altitudinal periglacial landform distribution is~~ supported by ERT data detecting ice-rich permafrost in a rockglacier at 5500 m a.s.l. ~~and ice lenses around the rockglacier (5450 m a.s.l.)~~. The highest, multiannual ~~displacement-creeping~~ rates up to 150 mm/y are observed typically on these rockglaciers ~~-indicating permafrost occurrence by displacemnet.~~ However, seasonality of rockglacier ~~creeping rates~~surface displacement like in other high mountain areas is missing. This study closes the gap of unknown ~~state of periglacial features and potential~~ permafrost occurrence in ~~a high-elevated basin at the western~~

35 | [Nyainqêntanglha Range](#) (Tibetan ~~mountain areas~~ [Plateau](#)) and ~~showssuppose~~ -compared to other high mountain regions- a higher-~~elevated~~ permafrost occurrence ~~in elevation~~.

Copyright statement (will be included by Copernicus)

1 Introduction

Information on permafrost (defined by perennially cryotic ground, Ballantyne 2018) distribution is of great importance in
40 | times of global warming, especially in high mountain areas (Hock et al., 2019), because these areas are climatically sensitive
(Barsch, 1996). The International Panel on Climate Change (IPCC) reported for 2019 the strongest observed increase of
permafrost temperature (globally averaged across polar and high-mountain regions) since 2007 (Hock et al., 2019). Due to
its thermal definition, it is difficult to evaluate permafrost conditions without ice-occurrence. Periglacial landforms of high
mountain environments are particularly well suited to detect and to study changes of permafrost temperature and the related
45 | ice content (Kneisel and Kääb, 2007, Kääb, 2013, Knight et al., 2019). These changes have an increasing impact on people
and their livelihood (Gruber et al., 2017), e.g., due to the importance of long-term ground ice as water resource (Jones et al.,
2019) in arid/semiarid regions (Halla et al., submitted). The frozen water storages [have a strong impact on](#) ~~impact highly~~
water budgets by permafrost degradation and glacier melt (Bibi et al., 2018; Song et al., 2020), especially at the so-called
Asian water tower, which provides water for more than 1.4 billion people (Immerzeel et al., 2010). The occurrence of
50 | natural hazards increases due to thawing permafrost (Zhang and Wu, 2012, Yu et al., 2016), for example by destabilizing
mountain slopes and rock walls (Deline et al., 2015). The scientific and social importance leads to a stronger focus on
permafrost areas, especially on the Tibetan Plateau (TP) where permafrost conditions react fast to climate warming (Cheng
and Wu, 2007).

Permafrost research in engineering has a 60-year-long tradition on the TP (Chen et al., 2016; Yang et al., 2010). The
55 | continuous use and life span of infrastructure depends on stable surface conditions which are strongly decreased ~~d~~ by
permafrost degradation. The engineering corridors for infrastructure projects like the Qinghai-Tibetan Highway/Railway and
pipelines (Yang et al., 2010; Yu et al., 2016) were accompanied by monitoring permafrost sites based on borehole
temperature (Hu et al., 2020; Li et al., 2009a), ground temperature data (Cheng and Wu, 2007; Ma et al., 2006) and
geophysics: Small scaled ground ice distribution was investigated by ground penetration radar (Wang et al., 2020; Wu et al.,
60 | 2005; You et al., 2017) and by [electrical resistivity tomography \(ERT\)](#) (You et al., 2013, 2017) close to the important
Highways/Railways. Compared to the central and eastern parts of the TP permafrost surveys ~~in other parts in western and~~
~~southern-of-the~~ TP are very scarce (Chen et al., 2016). Additional permafrost studies outside the engineering corridors are
limited to modelling results and large scale permafrost distribution maps (Ran et al., 2012, ~~Cao et al., 2019~~) [Cao et al., 2019](#),
[Obu et al., 2019](#)). Implications [offer](#) a temperature warming followed by permafrost degradation for the entire TP ~~are~~ [is](#) hard

65 to deduce due to inadequate distribution and small number of stations recording air temperature (Yang et al., 2010).
Therefore, modelling approaches are gaining increasing importance in order to estimate the consequences of the current
temperature rise ~~at-on~~ the TP. This warming temperature trend is reconstructed by $\delta^{18}\text{O}$ records in four spatially well-
distributed ice cores back to the beginning of the last century (Yao et al., 2006). Sun et al. (2020) confirm the relationship
between the temperature increase and permafrost degradation on the TP by a slow adaption until the year 2100 based on a
70 numerical heat conduction permafrost model. New statistical and machine learning approaches ~~determine-suggest~~ that the
permafrost extent on the entire TP is 45.9% (2003-2010) and they predict future permafrost degradation of 25.9% by the
2040s and 43.9% by the 2090s (Wang et al., 2019). Cheng and Wu (2007) also conclude that more than “half of the
permafrost may become relict and/or even disappear by 2100”. ~~Large scale modelling of PF conditions on the TP (Sun et al.,
2020) show that some areas (e.g. the study area in Figure 1, B and C) is located at the interface between continuous
permafrost and seasonally frozen ground, which makes it a suitable environment to validate such large scale models and to
precise the interface with ground truthed data. The validation is important, because the final conclusion would be that some
higher region on the TP is not completely underlying permafrost conditions, unlike expected and modelled at other places at
the TP (Cao et al., 2019; Ran et al., 2012).~~

80 ~~This study aims to supplement the previously summarized studies with an assessment of probable occurrence of permafrost
in remote high mountain regions away from the Tibetan engineering corridors and to provide a ground truthing for existing
permafrost studies and maps on the TP. The use of the term “probable” is motivated by the fact that we do not have
borehole-data (Mean annual ground surface temperature). Furthermore, and no small-scaled modelled permafrost
distribution is available, and therefore, So, we assess its the occurrence indirectly. The spatial heterogeneity of our data
(mapping, InSar and ERT) and of topographic variations in permafrost occurrence also prevents us from providing precise
85 elevational limits, sothus we provide an assessment of probable occurrence of permafrost in a range according to the findings
of the three methods. This study aims to complement the summarized, previous results with the occurrence of permafrost in
remote high mountain regions away from the engineering corridors and to provide an important supplement and ground truth
for existing permafrost studies at the TP~~

90 ~~Our study area Large scale modelling of PF conditions on the TP (Sun et al., 2020) show that some areas (e.g. the study area
in (Figure 1, B and C) is located at the interface between continuous permafrost and seasonally frozen ground according
large-scale modelling results of PF-conditions on the TP (Sun et al., 2020). , whichThe location makes it a suitable
environment to validate such large-scale models and to precise the interface with ground-truthed data. The validation is
important, because the final conclusion would be that some higher region on the TP is not completely underlying permafrost
conditions, unlike expected and modelled at other places at the TP (Cao et al., 2019; Ran et al., 2012).~~

95 ~~The identification of periglacial landforms, subsurface ice and surface creeping rates on these landforms leads to an
assessment of the probable occurrence of permafrost. We use periglacial landforms, subsurface ice and surface creeping rates~~

on these landforms to assess the lower boundary occurrence of probable permafrost occurrence. Periglacial landforms such as active (creeping) rockglaciers and protalus ramparts can contain ice (Barsch, 1996; Schrott, 1996; Scapozza, 2015) and are considered indicators of permafrost occurrence (Frauenfelder et al., 1998; Kneisel and Kääb, 2007; López-Martínez et al., 2012). ~~Especially on the TP only sparse less literature is found that describes periglacial landforms and permafrost occurrence (except Fort and van Vliet-Lanoe, 2007 and Wang and French, 1995). However, these is periglacial indicators are essential for creating large-scale permafrost distribution maps making them useful for the creation of large-scale permafrost distribution maps (e.g. Schmid et al., 2015).~~

We present a multi-method approach to provide a reliable prediction of subsurface ice and permafrost occurrence to answer the following research questions:

- How are periglacial landforms distributed?
- Do periglacial landforms like rockglaciers and protalus ramparts contain ice?
- Are they active and ~~which~~ what creeping rates can we observe?

We created (A) an inventory of periglacial landforms indicating potential subsurface ice-occurrence, we (B) acquired Electrical Resistivity Tomography (ERT) data to validate the ice occurrence of selected landforms, and we (C) ~~then~~ used multi-annual surface ~~creeping displacement~~ rates from InSAR time series analysis to corroborate the hypothesis of long-term ice occurrence due to permafrost conditions above a special elevation. ~~We make use of the fact that tThe deformation of debris supersaturated with ice causes surface displacement by downwards permafrost creep (Barsch, 1996; Delaloye et al., 2010). [1] interpret~~ Therefore, the remote sensing approach allows a transfer and an extrapolation of our findings about ice occurrence from geophysical measurements to other periglacial landforms in our study area. As a result, the study provides a lower limit occurrence of probable permafrost occurrence by combining these three methods for a catchment in the high-altitude mountains of the TP.


As a result, the study provides probable occurrence of permafrost by combining these three methods for a catchment in an high-altitude mountain range of the TP.

2 Study area

The Nyainqêntanglha Range (~~Figure 1~~) (~~figure 1~~) was formed during the Himalayan-Tibetan orogenesis as part of the central Lhasa block (Kapp et al., 2005; Keil et al., 2010). From Tertiary to Quaternary, the Nyainqêntanglha area was controlled and compressed by a fracture belt which folded and rose violently, forming the Nyainqêntanglha Mountains, with the highest peak of 7162 m a.s.l. (Kidd et al., 1988; Keil et al., 2010). Our study area, the Qugaqie catchment is characterized by ~~cretaceous~~ Cretaceous red beds and sandstone in the northern part and by early tertiary granodiorites in the center. The

bedrock of the southern part consists of biotite adamellites and glaciers in the highest zone (Kapp et al., 2005; Yu et al., 2019). The atmospheric circulation pattern and the topographic characteristics are responsible for a similar glacier distribution pattern in all North-oriented catchments of the Nyainqêntanglha range, including the Qugaqie Basin (Kang et al., 2009; Bolch et al., 2010). ~~In~~On the Lee side of the main Nyainqêntanglha crest and therefore in the Lee site of the moisture of the Indian Summer Monsoon (ISM) the glaciers are smaller in area and length (Bolch et al., 2010) (Figure 1, B). Bolch et al. (2010) also investigated the glacier shrinkage based on satellite data. They observed a glacier retreat of about $-9.9 \pm 3.1\%$ between 1976 and 2009. The Zhadang glacier located in the Qugaqie head lost an area of almost 0.4 km^2 in the same time span and covered an area of 2.36 km^2 in 2009. The corresponding retreat rate is 14 %, slightly larger than the regional average, which could indicate a slightly faster deglaciation of the smaller, north-orientated glaciers in the Nyainqêntanglha range.

The Qugaqie catchment is a sub-catchment of the Nam Co catchment, which is influenced by a strong climate seasonality driven by different wind systems throughout the year (Yao et al., 2013): Westerlies dominate in the winter months and provide cold, dry continental air from east to northeast (Figure 1, A, blue arrows), with temperature minima below -20°C . The dry season ends with the onset of the ~~Indian Summer Monsoon (ISM)~~ (Figure 1, A, red arrows), which provides moisture from May to September (Mügler et al., 2010). 80% of annual precipitation ($295\text{--}550 \text{ mm/y}$) occurs during the monsoon dominated summer months (Wei et al., 2012). The influence of the East Asian Monsoon on our study area is minor but it is an important source of moisture for the eastern TP (Figure 1, A, black arrows). Consequently, the study area of the Qugaqie Basin, situated in the Western Nyainqêntanglha Range (Figure 1, B), is characterized by semiarid climate and a large amount of solar radiation due to the high elevation and reduced cloud cover (Li et al., 2009). With an area of almost 60 km^2 the basin drains into the dimictic lake Nam Co (Figure 1, B) and the relief extends from 4722 m a.s.l. to an elevation up to 6119 m a.s.l.






Figure 1: A: Location of the **overall** study area **within the Tibetan plateau (TP)** (Background on SRTM DEM v4; Jarvis et al., 2008). Different wind systems influencing climate of the TP are shown by blue (Westerlies), red (Indian Summer Monsoon) and black (East Asian Monsoon) arrows based on Yao et al. (2012). B: Overview map of the Nam Co catchment with **the** altitude colors and the study area of the Qugaqie catchment (thick black lines). Note the greater glacier extents on the south-oriented mountain range. Bathymetric data originated from Wang et al. (2009). **Glacier extents originated from GLIMS database (Bajracharya et al., 2014; Liu and Guo, 2014) Glacier extents originated from the GLIMS database (Guo et al., 2015, Cogley et al., 2015) Liu and Guo, 2014, Cogley et al., 2015).** C: Permafrost distribution in the Nyainqêntanglha range based on Zou et al. (2017). **Note the proposed permafrost occurrence coverage of more than 90 % in the study area, marked by the black edging.**

Detailed information about permafrost occurrence and distribution in the study area is very scarce. Tian et al. (2006) determined a lower limit of permafrost based on soil probes at an elevation of around 5400 m a.s.l. along the northern slopes of Mt. Nyainqêntanglha **Feng-peak** (Figure 1, B). This is generally higher than in other regions (>4500 m a.s.l.) of the TP (Ran et al., 2012). Schütt et al. (2010) sampled lacustrine sediments from a permafrost lense in an outcrop at the Gangyasang Qu's entry into the North-western end of Lake Nam Co at 4722 m a.s.l.. Zou et al. (2017) distinguish between seasonally frozen ground and permafrost on their distribution map over the TP (Figure 1, C). According to their map permafrost is


165 ~~available-existent~~ at elevation higher than 5000 m a.s.l. in the Qugaqie basin. A coarse overview including a distinction between glacial and periglacial processual states around the lake Nam Co is given by Keil et al. (2010). [A two-years temperature-dataset on the Zhadang glacier, recorded at 5680 m a.s.l. by an automatic weather station \(2009-2011\) in 2 m height, shows a mean annual air temperature \(MAAT\) of -6.8°C \(Zhang et al., 2013\) and suggests permafrost conditions for the surrounding periglacial landscape.](#)

3 Data and Methods

170 We have used three different methods (A-C) to gain insights into permafrost-indicating periglacial landforms and to ~~derive assess~~ the lower ~~limitoccurrence~~ of ~~the~~-probable permafrost-~~occurrence~~ in the Qugaqie catchment-~~(Fehler! Verweisquelle konnte nicht gefunden werden.Figure 2)~~. The following methods [\(Figure 2\)](#) indicate information about permafrost conditions:

- 175 (A) Geomorphological Mapping: A map visualizes the distribution and characteristics of periglacial landforms and geomorphometric features.
- (B) Geophysical methods: Electrical resistivity tomography (ERT) identifies ice content and reveals the subsurface structure of periglacial landforms.
- (C) Microwave remote sensing: Interferometric Synthetic Aperture Radar (InSAR) time-series analysis of ESA’s Sentinel-1 satellite data detects perennial, constant creeping rates of active periglacial landforms.

180






Figure 2: Schematic workflow of applied methods to ~~derive~~assess lower ~~boundary~~occurrence of probable permafrost ~~occurrence~~.

(A) A geomorphological map visualizes the distribution and characteristics of landforms and geomorphometric features with the focus on periglacial landforms on a catchment-wide/regional scale. Periglacial landforms like rockglaciers (Barsch, 1996) and protalus ramparts (Scapozza, 2015) can potentially preserve ice over a long period of time (Ballantyne, 2018) and their activity and perennial creeping is an indicator for permafrost occurrence (Delaloye et al., 2010; Eckerstorfer et al., 2018; Esper Angillieri, 2017). This circumstance is validated (B) by ERT to detect subsurface ice on a local scale. (C) InSAR time series analysis detects perennial creeping ~~rates~~ which ~~are~~is typical of active, periglacial landforms. Permafrost occurrence is indicated by the activity of landforms and the corresponding surface structures like bulges, furrows, ridges or lobes. If the creep of landforms can be traced back over more than two years (according to the definition of permafrost) and show typical surface characteristics, like bulges, furrows, ridges or lobes, permafrost occurrence is corroborated. We make use of the fact that the deformation of debris supersaturated with ice causes surface displacement by downwards permafrost creep (Barsch, 1996; Delaloye et al., 2010). Therefore we concretize surface displacement (rates) as permafrost creep (creeping rates) in this study. Although the continuous movement of periglacial landforms and the presence of ice can be implied from InSAR data alone, ground truth at selected locations by ERT is essential to exclude other possible interpretations.

We assess the lower occurrence of probable permafrost by the mean altitudinal distribution of periglacial landforms, by the subsurface ice occurrence which has been validated with geophysics, and by the active status which is indicated by perennial surface creeping rates. ~~We derived the lower limit of the probable permafrost occurrence from the mean altitudinal distribution of periglacial landforms, the subsurface ice occurrence and activity of which has been validated with geophysics and perennial surface displacement rates (Figure 3). .~~ An occurrence of sporadic Permafrost is not excluded in lower elevation, but cannot be validated by the used methods and due to scale issues.




Figure 3: Schematic, hypsometric distribution of mapped landforms. Red features show active, multiannual creeping structures (furrows, lobes, bulges, ridges) of periglacial landforms indicating the lower boundary occurrence of probable permafrost occurrence. Modified from Barsch (1996) after Höllermann (1983).

3.1 Inventory of ~~glacial and periglacial~~ cryospheric landforms mesoscale landforms

The mapping procedure consists of the elementary mapping steps, described by Knight et al. (2011) and Otto and Smith (2013). Pre-Mapping includes analyses of digital elevation models (DEM) and mapping of landforms on ~~aerial-optical~~ images in a scale of 1:10 000 (named here as Mesoscale following to Höllermann (1983)). The DEM used in this study originates from TanDEM-X data (2015) with a resolution of 12 m (©DLR). The ~~aerial-optical~~ images are based on Digital globe, BING maps (2013) and Google Earth data (2007-2012). Geomorphological symbols were used after Kneisel et al. (1998) ~~in the field~~ for field mapping and after Otto (2008) for the digitized visualization in ArcGIS. During the field campaign, the main focus was on the mapping of periglacial landforms at the Mesoscale (Höllermann, 1983). These landforms are components of the periglacial zone which is defined by seasonally-frozen and perennially-frozen ground ~~excluding glacially-conditioned environments~~ (French, 2017). A differentiation between seasonally-frozen and perennially-

frozen movement behavior is given by the data and a derived model by Reinosch et al. (2020). This data was used for the preparation of the cryospheric landform identification. Next to optical and InSAR data, the periglacial landforms were identified in the field by an inspection of the form, the substrat, the catchment and the potential process which formed the landform. The results section describes the inventory statistically and includes morphological field observations which could not be included in the map due to scale issues. For example, small-scaled death ice holes were not included in the mesocale geomorphological map. During post-mapping we integrated the manually field-mapped information into ArcGIS. Additional features like a stream network, lakes, ridges, glacier extents and moraines were delineated with the help of the mentioned DEM, a hillshade map (azimuth 315°, altitude 45°) and the mentioned aerial-optical images. Glacier extents were digitized based on aerial-optical images of the year 2013 (BING maps). Rockglaciers were mapped and identified following the comprehensive description by Barsch (1996). We followed the geomorphological mapping approach based on the baseline concepts (V 4.0) of the IPA Action Group “Rock glacier inventories and kinematics” and mapped the extended geomorphological footprint of the rockglaciers. Additional Mapping criteria of rockglaciers in the field were visible creeping structures on the surface (ridges, furrows, and lobes as those shown in Figure 4, A). In the Qugaqie Basin, moisture originates from glaciers and ice patches located in cirques.




Figure 4: (A) Panorama view on a rockglacier (No. 1) with marked creeping structures (lobe in black, ridges in yellow, and furrows in dashed red) in the Qugaqie Basin in hanging valley 3. (B) Example of a protalus rampart in hanging valley 3 of the Qugaqie Basin. A bulge (in black) formed through creeping of rockfall deposits. The length of the bulge is approximately 500 m. The location of the photos ~~at~~ can be found in Figure 9 B. (photos: J. Buckel)

Protalus ramparts (Figure 4, B) were mapped as periglacial features or permafrost-related landforms as suggested by Scapoza (2015). A straight headwall for the sediment source is required, as the sediment originated by rockfalls and is accumulated at the foot of the rockwall. Infiltrating moisture originating from precipitation and snow melt freezes the sediment deposit and creates a bulge parallel to the rockwall. These ice-permeated rockfall deposits creep downwards. Scapoza (2015) also noted the challenge to differentiate protalus ramparts from initial talus rockglaciers in the sense of Barsch (1996). Protalus ramparts mapped in the present study show no ridges, furrows, or lobes at the surface, but the mapped rockglaciers do. It is pertinent to point out that our mapping procedure both in the field and during post-mapping consistently differentiates between rockglaciers and protalus ramparts based on the above-mentioned criteria. An incorrect determination as pronival ramparts can be minimised by the absence of longer existing snow fields due to arid climate conditions during the winter and the strong solar radiation and less cloud cover due to the extreme altitude (compare Hedding, 2016).

3.2 Ice detection by ERT

Electrical resistivity tomography (ERT) is a widely-used method in geomorphology (Schrott and Sass, 2008). The application works especially well for subsurface ice detection due to strong differences between frozen (high resistivity values) and unfrozen ground (low resistivity values) (Hauck and Vonder Mühll, 2003; Hauck and Kneisel, 2008). Since the end of the 1990ies the method has been established for ~~Permafrost-permafrost~~ detection in solid rock (Krautblatter et al., 2010; Hartmeyer et al., 2012) and in debris-ice mixtures, like rockglaciers (Von der Mühll et al., 2002; Kneisel et al., 2008; Rosset et al., 2013; Emmert and Kneisel, 2017; Mewes et al., 2017).

For the usual four-point measurement of the ground electrical resistivity, two electrodes feed current into the ground, which establishes an electric field in the subsurface. Another pair of electrodes is used to measure the voltage drop between two other locations on the surface. In order to obtain information on the two-dimensional distribution of electrical resistivity in the subsurface, a linear arrangement of the four electrodes is used to measure at different positions along the profile and with varying distances between the electrodes (Wenner array). The apparent resistivity (in Ωm) of each measurement can be calculated from the injected current, the applied voltage, and a factor, which takes the geometry of the arrangement into account. Subsequently, inverse modelling techniques are used to reconstruct the resistivity structure of the subsurface from the measured apparent resistivity data (Loke and Barker, 1995).

We performed ERT measurements during a field campaign in July 2018. We worked with multi-electrode (50) equipment “GeoTom-MK” (GEOLOG2000, Augsburg, Germany), a maximum spacing of 2 meters, allowing a maximum profile length

of 98 m with a single measurement. To obtain longer sections, we used the roll-along procedure illustrated in Figure 5. For this procedure, two cables were available (denoted A and B), each equipped with 25 channels. First, both are connected with the control unit to obtain pseudosection number 1 (Figure 5). Next, cable B (and all connected electrodes) remains at the same location, whereas cable A is moved to the right of cable B to measure the Pseudosection number 2, and so on. The

location of the ERT profiles was partly constrained by logistical conditions. Due to the high altitude, the crew had to stay at one level for three days to get adapted to altitude. The measurement locations were not accessible by vehicles, and a few hours were needed every day to reach the sites, resulting in limited productivity. Therefore, we tried to locate the profiles efficiently to obtain a representative data set of the valley. We covered different landform features (moraine, valley bottom, Rockglacier) where permafrost conditions were assumed. Blocky surfaces constitute a challenge for ERT measurements due to instability and a lack of fine material necessary to provide sufficient contact for the electrodes. In cases where no soil material could be found that closed the gaps between the boulders, we inserted the end of each electrode into a sponge saturated with salt water to improve connectivity to the fine material. The saturated sponge kept the fine material wet and diminished desiccation through high solar radiation. The ERT-data was processed with the Res2Dinv-Software (©Geotomo Software).




Figure 5: Measurement setup for the Roll-along procedure (adapted from [N El Sayed et al., 2018](#)).

3.3 ~~Surface displacement~~Creeping rates by InSAR-analyses

InSAR time series analysis is an active microwave remote sensing technique, which can exploit the phase change of the backscattered microwaves to determine relative surface displacement in the order of millimetres to centimetres (Osmanoğlu et al., 2016). Both the amplitude and the phase of the microwave backscatters are used for InSAR. After precisely co-registering all acquisitions, it is possible to calculate the average phase change of each resolution cell over time, which

contains a number of different signals, including whether a resolution cell moved closer to the receiver, i.e. the satellite, or further away from it. These images of phase change are called interferograms. The accuracy of the derived motion is dependent on a number of different factors, including the frequency of the emitted wave, the atmospheric delay, the accuracy of its modelling, the topographic data used to correct the images, the choice of reference points, the surface characteristics of the observed structure and the frequency of the data acquisitions (Hu et al., 2014).

The reliability of an interferogram is often described by its so-called coherence. Coherence is a measure of phase stability with a value near zero representing poor reliability and values near one representing high reliability (Crosetto et al., 2016). If the backscatter characteristics of the observed surface changes too much between two acquisitions, e.g. due to snow cover, vegetation or events occurring between the acquisitions like rock falls, the coherence is poor and no phase change can be determined reliably. Coherence also decreases with increasing displacement and displacements larger than half the SAR wavelength (~2.8 cm for Sentinel-1) cannot be determined accurately. For this study we chose a coherence threshold of 0.3 and discarded areas with coherence values below 0.3. This threshold is similar to the one chosen by (Sowter et al., (2013) and provides good spatial data coverage while also excluding unreliable data. The issue of low coherence or decorrelation is exacerbated for interferograms with a long temporal baseline i.e. a long time period between data acquisitions.- No Sentinel-1 data is available for a period of 48 to 96 days during the summers of 2016 and 2017. These longer temporal baselines cause decorrelation during the summer months on some of the faster landforms. Freezing and thawing of the ground leads to reduced coherence values in autumn and spring. In our study area we observe decorrelation most heavily in spring and summer, when the active layer thaws and is inundated by precipitation. The coherence over periglacial landforms in the Quagaie Basin is relatively good, due to the lack of high vegetation on actively moving landforms and the relatively sparse snow cover in winter visible on optical Sentinel-2 acquisitions.

Exploiting the phase change with InSAR provides only relative surface motion towards the satellite or away from it. The Line-Of-Sight (LOS) of the satellite is therefore very important, as motion with a very different direction compared to this LOS is severely underestimated (Hu et al., 2014). The severity of this underestimation depends on the angle between the LOS and the direction of the surface displacement. An angle close to 0° will cause only minor underestimation, while displacement with a direction near 90° to the LOS will be severely underestimated or even completely overlooked. The Sentinel-1 satellites follow a circumpolar orbit and observe the earth obliquely with an incidence angle of 33°-43° (Yague-Martinez et al., 2016). Both ascending and descending acquisitions are therefore sensitive to vertical surface displacement and towards the East or West but very insensitive to displacement towards the North or South. Both ascending (satellite travelling south to north) and descending (satellite travelling north to south) acquisitions are therefore sensitive to vertical surface displacement and towards the East or West respectively, but very insensitive to displacement towards the North or South. We always select the geometry with the highest sensitivity towards the expected displacement direction to calculate our displacement and velocity results. To that end we calculate a sensitivity coefficient for each pixel which is explained in the following.

320 The surface displacement data presented in this study represents a spatial subset of a surface displacement model originally based on [Reinosch et al. \(2020\)](#). ~~Mean velocities were calculated by dividing the cumulative displacement observed during the observation period by the length of the observation period (2015-2018). All surface velocity data of periglacial landforms have been projected along the direction of the steepest slope under the assumption that the motion of the described landforms is mainly gravity-driven by an ice-debris mixture. Hereafter we will refer to the mean surface velocity of periglacial landforms projected along the steepest slope as “creeping rates” to reflect this assumption. All surface displacement rates of the periglacial landforms have been projected along the direction of the steepest slope under the assumption that the motion of the described landforms is mainly gravity driven. To produce more representative motion information for all slopes, we projected both ascending and descending LOS velocities into the direction of the steepest slope. We assume that all motion on slopes is directed downslope and parallel to the surface. For this approach, we~~ We calculate a sensitivity coefficient to compensate for the underestimation of the displacement signal caused by the disparity between the LOS and the assumed displacement direction. We followed an approach developed for the study of landslides (Notti et al., 2014), as the displacement of landslides is gravity-driven, which we also assume to be true for the periglacial landforms investigated in this study. ~~For this approach, we calculate a~~ The sensitivity coefficient is based on the difference between the angle of the LOS vector of the satellite and the expected downslope ~~displacement~~ creeping direction. The strength of this coefficient increases with the difference between the angles of the vectors. ~~The coefficient can vary between 0 for areas where the satellite’s sensitivity is low to 1 where the sensitivity is very high. Values below 0.2 are not used to avoid excessive amplification of displacements and associated errors. The coefficient can vary between 0.2 for areas where the satellite’s sensitivity is low to 1 where the sensitivity is very high.~~ The LOS velocity can therefore be increased by a factor of up to 5 by this downslope projection. ~~If both ascending and descending velocity LOS data is available for the same pixel, then we use only the geometry with the higher sensitivity coefficient, i.e. better sensitivity, to calculate the downslope velocity and ignore the other geometry to keep the precision of the projection as high as possible.~~

For our analysis of the Quagaie Basin, we processed 278 interferograms from 74 ascending acquisitions (June 2015 to December 2018) and 257 interferograms from 63 descending acquisitions (November 2015 to December 2018) (Table 1).
345 The temporal baselines, i.e. the time period between two data acquisitions, of individual interferograms is mostly 12 to 36 days with a maximum of 72 and 96 days for ascending and descending orbits, respectively. All data acquisitions originate from ESA’s Sentinel-1 a/b satellite constellation. Both ascending and descending datasets were processed using Small Baseline Subset (SBAS) time series analysis (Berardino et al., 2002), with a coherence threshold of 0.3. We observe low coherence during the thawing period in spring and the freezing period in autumn. We therefore employed the intermittent
350 SBAS approach (Sowter et al., 2013) for the entire data set, allowing us to improve our spatial coverage by interpolating time series results for data points with intermittently low coherence. Data points need to exceed the coherence threshold of 0.3 in at least 75 % of all interferograms to qualify for interpolation, otherwise they were discarded. We used the TanDEM-X 12x12 m DEM to remove the topographic phase from our interferograms (DLR, 2017).

Table 1: Summary of ISBAS processing parameters

<u>Geometry</u>	<u>Observation period</u>	<u>Acquisitions</u>	<u>Interferograms</u>	<u>Temporal baseline</u>	<u>Coherence threshold</u>
<u>ascending</u>	<u>2015-06-05 to 2018-12-22</u>	<u>74</u>	<u>278</u>	<u>12 to 72 days</u>	<u>0.3</u>
<u>descending</u>	<u>2015-11-15 to 2018-12-29</u>	<u>63</u>	<u>257</u>	<u>12 to 96 days</u>	<u>0.3</u>


~~Surface displacement~~Creeping rates presented in this study were not verified by independent measurements (GPS measurements, laser scans, optical remote sensing etc.), as no such data sets exist for our study area. Reference points are located on bedrock whenever possible and on ridges or stable, vegetated moraines with good coherence if no coherent bedrock was available~~We took great care in the selection of our reference areas (choosing bedrock whenever possible) but the lack of data sets challenges a validation of surface displacements for the assessment of their reliability (compare Figure 9, A).~~ Areas which are likely unmoving on a multiannual scale, such as the old moraines at the entrance of the Qugaqie Basin, display LOS velocities of ± 2.4 mm/yr during our observation period. This does not provide information regarding the accuracy of the seasonal variations of our surface displacement results but it indicates that the multiannual LOS velocity results are reliable. We use this variation of ± 2.4 mm/yr over likely stable areas as the precision of the mean LOS velocity during our observation period. The precision of the downslope velocity was determined by dividing the precision of the LOS velocity by the sensitivity coefficient. It therefore varies between 2.4 mm/yr and 12.0 mm/yr for areas with a sensitivity coefficient of 1 and 0.2 respectively (Reinosch et al., 2020).

4 Results and Interpretation

4.1 The cryosphere of the Qugaqie Basin

The geomorphological map in Figure 6 shows features of the mesoscale cryosphere in the Qugaqie Basin: glaciers, moraines, protalus ramparts and rockglaciers. The moraine distribution suggests that former glaciers extended to the present shoreline of the Nam Co at their largest size during Marine Isotope Stage (MIS) 3 (Dong et al., 2014). Multiple smaller moraines are displayed in closer proximity to today's glaciers (Figure 6). Glacial landforms like valley glaciers, cirque and wall glaciers increase in number and size towards the south due to a higher elevation and shorter distance to the main ridge (Figure 6). Only the Genpu (1.56 km²) and the Zhadang (1.41 km²) glaciers are considered valley glaciers, most of the other glaciers are located in the head of the hanging valleys as cirque glaciers. The northward orientation of all glaciers is a result of the Lee-effect towards incoming moisture from the southern direction. The topographic barrier of the Nyainqêntanglha Range detains precipitation and causes an asymmetric, ~~and uneven~~ north-south distribution of glacier extents expressed by smaller extents in the northern catchments draining in the Nam Co like Qugaqie (compare Bolch et al., 2010). The glacial zone with a

380 | cumulative glacier area of 4.07 km² ([bing maps 2013](#)) extends from 5500 m a.s.l. to the highest elevation (6086 m a.s.l.) with a mean elevation of 5770 m a.s.l..




385

Figure 6: Geomorphological map of the Qugaqie basin. The locations of the ERT profiles are shown with purple lines. Periglacial landforms are greenish (rockglaciers and protalus ramparts). The black rectangle represents the boundary of the map shown in Figure 9.

390

The altitudinal (mean) landform distribution illustrates the statistical analyses and displays a typical high-mountain pattern (Figure 7). Debris and talus cones can be found in lower altitudes. The periglacial landforms (i.e. protalus ramparts and rockglaciers) are located between elevations of 5300 m and 5600 m a.s.l. and the average number of periglacial landforms is situated around 5500 m a.s.l. ~~However, the highest number (29) of these landforms occurs from a minimum elevation of 5400 m a.s.l.~~ We conclude from this altitudinal distribution a probable occurrence of permafrost higher than 5300 m.a.s.l., which has to be supported by validating ice occurrence and the status of activity of these landforms.~~that the lower boundary occurrence of probable permafrost occurrence of around 5500 5400 m a.s.l., which has to be supported by ice occurrence and the status of activity.~~



395

Figure 7: Altitudinal (mean) landform distribution of the Qugaqie basin derived from the landform inventory.

Most rockglaciers are located in cirques and three are supplied by glacial melt water resulting in greater extents compared to rockglaciers without a glacier in their catchment (Figure 6, No. 1, 2 and 3). Additionally, moraine deposits, talus slopes and protalus ramparts ~~moraine deposits~~ provide the sediment accumulation at the base required for the formation of a rockglacier

besides water availability (Knight et al., 2019). The altitudinal distribution of the rockglaciers extends from 5363 m to 5789 m a.s.l. with a mean elevation around 5500 m a.s.l. (Figure 7, [Table 2](#)). Rockglacier surfaces display clear creep structures and rockglacier-typical bulges, furrows and lobes (Figure 4, A). There is no pronounced lichen growth, and the uppermost material is extremely unstable. These field observations allow the conclusion of an active status of the rockglaciers, which indicates ice occurrence and, thus, permafrost conditions (according to Barsch, 1996). The altitudinal distribution of protalus ramparts has a narrower range of min-max values, but they are located at a similar mean elevation. The mean area of the individual protalus ramparts is only half of the mean area of the individual rockglaciers, i.e., protalus ramparts are generally smaller than rockglaciers ([Table 2](#), Figure 6), but there are twice as many. Protalus ramparts are situated in front of rocky slopes and are characterized in contrast to rockglaciers by a shorter dimension in down slope (Figure 4 and Figure 6).

The mesoscale periglacial landforms (mean elevation) are situated between 5300 and 5600 m a.s.l. This altitudinal distribution serves as one component of the three methods for assessing the probable occurrence of permafrost in the catchment.

Table 2: Statistical description of cryotic landforms based on DEM-analyses.

	No.	cumulative area [m²]	area (mean)	elevation (min, max)	elevation (mean)
Protalus rampart	22	1018014	46273	5292, 5685	5530
Rockglacier	10	830185	83019	5363, 5789	5523
Glacier	11	4075580	370507	5504, 6086	5771

4.2 ERT-based ice detection

ERT is a common method to detect ground-ice in the subsurface, inferring permafrost conditions (Lewkowicz et al., 2011), if ground ice is present for two consecutive years. With the help of ERT we were able to provide evidence for the existence of ground ice at specific test sites. The localized site surveys are critical for descibing the 5300 – 5450 m estimate because no

high resistivity values representing subsurface ice content were measured. Profile A (Fig. 8) ranges from 5090 to 5230m and represents subsurface conditions in the lower altitudinal areas of the catchment, for example in a lateral moraine.


Figure 6 displays the locations and indicates an altitudinal increase of the four ERT-profiles (A to D). The measured resistivity values were compared with tables by Hauck and Kneisel (2008) and Mewes et al. (2017). These studies also address ice detection in high altitude periglacial environments. Table 3 sums up our measured resistivity values and classifies the values ~~in terms their interpretation~~ in terms of material characteristics. Different studies show resistivity values of till in a range from 1 to 10 kΩm (Reynolds, 2011), from 5 to 10 kΩm (Thompson et al., 2017) and from 50 – 100 kΩm (Vanhala et al., 2009). ~~This addresses one key problem of drawing a conclusion only from resistivity values without additional methods.~~ The diversity of resistivity ranges and the resulting non-uniqueness can be overcome by using additional methods to support the final conclusions.

Table 332: Resistivity values for different materials derived by field measurement. The used terminology of the interpreted material followed Hauck and Kneisel (2008) and Mewes et al. (2017).
~~Resistivity values for different materials derived by field measurement, compiled after Hauck and Kneisel (2008) and Mewes et al. (2017)~~

Material	Resistivity [kΩm]
Sandstone (moist – dry)	0.5 – 5
Till	20– 80
Unfrozen sediment (moist – dry)	1 – 20
Ice-poor permafrost (ice lenses, ice-interspersed till)	50 – 150
Ice-rich permafrost (massive ice body)	150 – 4 000

Profile A (Figure 8) ranges from 5090 to 5230m and represents subsurface conditions in the lower altitudinal areas of the catchment, for example in a lateral moraine. At the surface the profile has a length of 348 m, but the length information in the following text refers to the x-axis which corresponds to planar 2D-view (the topographic effect is not displayed). From

435 ~120 m on, we observe a slope-parallel highly resistive layer (highlighted by the black line in Figure 8, A) with resistivity values ranging between 5 and 100 k Ω m and an average thickness of 10 m. We interpret this layer as compressed till without ice content, based on the resistivity range, the compressed glacial sediment accumulation and the absence of creeping structures indicating ice. According to Yu et al. (2019) the underlying bedrock consists of sandstone, which explains the low resistivity values below the resistive moraine deposits. Between 0 and 20 m along the profile, the electrodes were directly
440 attached to the outcropping, weathered sandstone. The resistivity values around 5 k Ω m correspond to dry sandstone bedrock, which is exposed to strong solar radiation. The hydraulically impermeable till cover is not present between 20 and 120 m, and moisture infiltrates as slope water saturating the sandstone bedrock underneath the moraine and decreasing electrical resistivity.



445 **Figure 8:** Electrical resistivity sections along the four ERT profiles recorded in July 2018 with a standard spacing of 2 m. Profile C and D are located on rockglacier No. 1). Note the increasing elevation between profiles A and D.

Profile B (Figure 8; B) is located in hanging valley 3 on top of an old, a terminal moraine crossing the stream, which drains the hanging valley- (Figure 6). Surrounding dead ice holes indicate former subsurface ice occurrence behind the former moraine terminus. Complete vegetation cover of compresia pygmea interspersed with individual rockstones suggests an old and stable surface. From the high resistivity anomalies of up to 150 kΩm, we conclude that ice-poor permafrost in contrast to ice-rich permafrost in profile C and D is present as an ice lens at 5450 m a.s.l.

Profile C and D (possibly the highest-elevated ERT-measurements worldwide) show the typical two-layer structure of rockglacier No. 1 with equally high resistivity values (Figure 8; C, D). The first layer is characterized by lower resistivity values (1 – 20 kΩm), indicating the unfrozen active layer during the summer months. The active layer thickness varies between two and five meters. The second layer shows high resistivity values of up to 3500 kΩm and covers the complete section from below the active layer to the maximum depth of investigation. No internal heterogeneities are visible due to the lack of current flow within this highly resistive unit.~~No internal heterogeneities are visible within this highly resistive unit,~~ which we interpret as a mixture of ice and sediment. According to Table 3, we interpret the second layer to be ice-rich permafrost. Similar resistivity values of ice-rich rockglacier material, reaching maximum values of 1000 kΩm have been reported in several studies from Häberli and Vonder Mühll (1996), Vanhala et al., (2009) and Mewes et al., (2017). Profile C and D confirm the presence of subsurface ice at an elevation around 5500 m a.s.l., which we use as evidence for the lower limit occurrence of probable permafrost-~~occurrence~~.

The relatively large altitudinal steps between our four ERT profiles do not allow excluding the occurrence of subsurface ice in other, lower parts of the valley. Therefore, we use the following perennial creeping rates to exclude this case. The detection of subsurface ice is the second component of the three methods for estimating the probable occurrence of permafrost. Inferred by ERT-data subsurface ice can be expected at selected locations from an altitude of 5450 m and higher.

4.3 ~~Surface displacement~~Creeping rates of periglacial landforms


The creeping rates for rockglaciers and protalus ramparts, including statistical information, are shown in Table 4. ~~Rockglaciers display the highest mean velocities (26.8 mm/yr), followed by protalus ramparts (12.7 mm/yr). Table 3 summarizes the InSAR surface displacement results for rockglaciers and protalus ramparts.~~ The fastest moving areas of landforms display lower coherence values and spatial data gaps. The low coherence values in those areas are likely connected-related to the long temporal baselines of interferograms in summer of 2016 of up to 72 days and 96 days for ascending and descending data respectively. Long temporal baselines on relatively fast moving landforms may lead to aliasing effects if the displacement exceeds a quarter of the wavelength of the satellite (Crosetto et al., 2016). This would correspond to a LOS displacement of ~~~1.4 mm~~ for Sentinel-1, which emits a wavelength of ~~5.6 mm~~. ~~1.4 mm~~ in 72 days or 96 days corresponds to a LOS velocity of approximately ~~7.1 mm/yr~~ for ascending and ~~5.3 mm/yr~~ for descending data respectively. Displacement in areas with higher LOS velocities than these thresholds is likely to be underestimated with

the InSAR technique and display poor coherence values near or below the coherence threshold of 0.3. Coherence values do not drop significantly in winter, which is likely due to the semi-arid climate and therefore relatively thin snow cover.

Table 4: Summary of InSAR- derived creeping rates for the periglacial landforms. The values represent the mean of all data points over the entire observation period (2015-2018) on the respective landform ~~and, T~~ the distribution is represented by the standard deviation. Creeping rate precision is calculated by dividing the LOS precision of 2.4 mm/yr by the sensitivity coefficient. The percentage of interpolated time periods describes how many interferograms are incoherent and therefore require interpolation with the ISBAS algorithm.

Landform	De <u>Creeping</u> <u>rate</u> wnslope velocity [mm/yr]	Summer acceleration [%]	Downslope velocity <u>Creeping rate</u> precision [mm/yr]	Coherence	Interpolated time periods [%]	Data points
Protalus ramparts	12.7±9.2	13±60	6.5±3.5	0.74±0.08	4±5	7984
Rock- glaciers	26.8±20.4	-9±36	7.0±4.0	0.65±0.11	9±6	5402

~~The active status, the mean altitudinal distribution and validated ice occurrence by ERT of the periglacial landforms corroborates display the lower boundary occurrence of probable permafrost occurrence around 55005400 m a.s.l. (Figure 9, blue line) transparent yellow feature) and the corresponding area of higher elevation. “Around” means that ice lenses at lower altitude are included, as well as lower active creeping periglacial landforms of the altitudinal range. Therefore, an occurrence of permafrost is not excluded in lower elevation, but can’t be validated by the used methods due to scale issues.~~






Figure 9: Surface-displacement rates from periglacial landforms. The elevation higher >5400 m a.s.l. (transparent yellow) displays the area of probable permafrost occurrence. Black rectangles display the two fastest rockglaciers in Figure 11. The mean distribution of active periglacial landforms corroborates the occurrence of permafrost at altitudes above the 5500 m a.s.l. isoline

~~highlighted in blue. Figure 10. Camera position shows the Photographs in Figure 4. A: Landsat 8 satellite image, recorded 30-01-2018. Triangles indicate stable reference points. Dashed lines indicate the outlines of the periglacial landforms. B: Creeping rates from periglacial landforms move in slope direction over the observation period 2015 - 2018. The Black rectangles mark the location of The Black rectangles mark the location of the two fastest rockglaciers in Figure 10. The Camera positions correspond to The Camera positions correspond to the Photographs in Figure 4.~~

Protalus ramparts in the Qugaqie Basin display lower average surface velocities than rockglaciers. The creeping rate of protalus ramparts (12.7mm/yr) is lower and shows more pronounced seasonal variations than that of rockglaciers (26.8 mm/yr). ~~The surface velocity creeping rate of protalus ramparts generally ranges from 10 mm/yr to 80 mm/yr and shows more pronounced seasonal variations than rockglaciers.~~ Rockglacier No. 1 of hanging valley 3 which we also studied with ERT measurement displays ~~slope velocities~~ creeping rates of up to 70 mm/yr in most areas, with the fastest moving part reaching 153 mm/yr (Figure 10, B), similar to the rockglacier No. 2 (Figure 10, C). A Velocity_ time series of creeping rates of the rockglacier No. 1 is shown in Figure 10 A (black line) and of the rockglacier No. 2 in Figure 10 A (grey line). The spatial distribution of the ~~surface velocity~~ creeping rates is relatively uniform in areas with good InSAR sensitivity, i.e. slopes with an East or West aspect, but displays significantly higher noise level in areas with poor InSAR sensitivity, i.e. slopes with a North or South aspect.

We do not observe a clear correlation between variations in ~~surface velocity~~ creeping rates and possible seasonal forcing mechanisms such as temperature or precipitation. Neither protalus ramparts nor rockglaciers display clear acceleration of ~~surface velocity~~ creeping in summer compared to winter (Figure 10 ~~Fehler! Verweisquelle konnte nicht gefunden werden.,~~ A).




Figure 10: Time series (A) of the ~~velocities~~creeping rates of the two largest rockglaciers No. 1 (black line) and No. 2 (grey line). Summer months with air temperatures $>0^{\circ}\text{C}$ (according to Zhang et al., 2013) are shown in red. Time series represent the moving average of the 10 nearest values in time based on the median of time series points, located in B (black dots) and C (grey dots). Dashed lines in (A) indicate the ~~mean downslope velocity~~creeping rate over during the entire observation period between 2015 and 2018. Black time series (rockglacier No. 1 in B) ~~are is~~ based on ascending data and grey time series (rockglacier No. 2 in C) on descending data.

The third component for assessing the occurrence of permafrost is based on the movement rates of periglacial landforms. Based on the assumption that a measurable movement rate is determined by perennial ice in the subsurface, the observed active status of the periglacial landforms allows the conclusion of permafrost occurrence in the corresponding landform.

4.4 Assessment of the lower permafrost limit of the Qugaqie valley

Integrating the findings of the applied methods allows to estimate a ~~allows to estimate~~ lower permafrost limit. The active status, the altitudinal distribution of the periglacial landforms and validated ice-occurrence by ERT suggest a lower limit of probable ~~lower permafrost limit. The active status, the altitudinal distribution of the periglacial landforms and validated ice-occurrence by ERT suggest a lower limit of probable~~ permafrost between 5300 – 5450 m a.s.l. This range includes ice lenses detected by ERT-data as well as all creeping landforms indicating an active status and therefore an existence of ice.

5 Discussion

One critical issue for the estimation of the lower ~~permafrost boundary is the question~~ occurrence of probable permafrost by the used approach is the focus on periglacial landforms. These landforms are characterized by blocky material and a special thermal regimes that lowers the internal temperature by 5 – 7°C due to the Balch and the chimney effects (Harris and Pedersen, 1998). The latter originated by findings of von Wakonigg (1996) in the eastern alps and requires a significant snow cover during winter. Due to the semi-arid climate ~~resulting in thin snow cover~~ in the study area and the extreme cold mean annual air temperature of -6.8°C ~~at~~ at 5680 m a.s.l this effect would play a minor role. The Balch effect assumes ~~that a thin snow cover favours permafrost conditions because of denser and therefore heavier air in the blocky material. This effect amongst other leads to our interpretation of ice-rich permafrost in the rockglaciers. Using the ERT-measurements we found ice-poor permafrost in ice-lenses in mineral soils next to the rockglacier that corroborates the idea of permafrost conditions outside of blocky material.~~

The next critical issue for the estimation of the lower occurrence of probable permafrost is the question ~~on~~ whether the huge resistivities observed on profile A (Figure 8, A, black lines) indicates ice or not. In general, subsurface material determination without additional cross-validating techniques by other geophysical methods or borehole data remains uncertain (Hauck and Kneisel, 2008; Guglielmin et al., 2018). Therefore, the geomorphological knowledge of the study area is essential for an interpretation of the subsurface: In this case, the measured resistivity values of Profile A (Figure 8) of up to 100 kΩm are consistent with both till or ice-poor permafrost (Schrott and Sass, 2008). From the resistivity values it is therefore not possible to determine whether the till contains ice or not. However, field observations allow the conclusion that no ice was measured because clear creep structures would have to be recognizable due to a significant slope. Furthermore, InSAR analysis of this location shows no clear perennial ~~surface-displacement~~ creep behaviour (Reinosch et al., 2020),

making the presence of subsurface ice unlikely. In order to uniquely identify ice, it would have been desirable to apply additional geophysical methods, like ground penetrating radar, refraction seismic tomography, or capacitively coupled resistivity (Mudler et al., 2019). In particular, the combination of electrical and seismic methods allows to derive a petrophysical four-phase model (Hauck et al., 2008; Mewes et al., 2017; Halla et al., 2020) and to estimate the sediment-to-ice ratios from electrical resistivity and seismic velocities. However, due to the extremely difficult logistical constraints in this remote location, these methods could not be applied, and we thus rely on combining evidence from field observations with geophysical results.

The approach by Kneisel and Käab (2007) uses a similar combination of methods as used in this study to describe periglacial morphodynamics of a glacier forefield including a rockglacier. ERT profiles show the same range of layer thickness of 2-5 meters as in our profiles in the summer months. They recommend the joint application of geoelectrical and surface-movement data to investigate periglacial landforms and to assess the permafrost distribution, because the combination of both tools allows a more comprehensive characterization of permafrost characteristics like ice-rich or ice-poor. Also, in our case, we believe the ground-based geophysical surveys are useful, as predicting subsurface ice content and deriving permafrost distribution maps only by modelling and/or using remote sensing includes various sources of error:

- Low resolution (1 km gridded) of the permafrost-distribution models over the entire TP (Zou et al., 2017; Figure 1, C) prevents detailed analyses of permafrost occurrence at a meso-scale, especially in high mountain relief.
- Surface displacement patterns originate from different surface processes and take place in different time-intervals, such as freeze-thaw cycles, seasonal creeping or constant, multiannual ~~displacements—creep~~ (Reinosch et al., 2020) (Reinosch et al., 2020).
- Remote Sensing approaches can only guess the geomorphological process behind the surface displacement. Surrounding landscape features, underlying material and sediment source areas are essential factors that need to be considered during the interpretation of remote sensing images.
- Without ground based validation (e.g. ERT-data) large-scaled permafrost distribution maps cannot accurately be used to predict permafrost occurrence in the remote, high mountain areas.

Geomorphological field evidence allows a small-scaled interpretation and, in combination with remote sensing data, an extrapolation to larger scales. The periglacial landforms in this study show lower ~~creeping rates~~ rates of surface displacement than similar landforms of other regions. Other studies employing InSAR techniques observe ~~creeping~~ displacement rates from centimetres to several meters per year for rockglaciers in the Western Swiss Alps (Strozzi et al., 2020), in western Greenland (Strozzi et al., 2020) and in the Argentinian Andes (Villarroel et al., 2018; Strozzi et al., 2020). Furthermore, all of them clearly indicate seasonal variations of the rockglacier movement, with faster rates in summer and ~~decreasing velocity~~ reduced creeping rates in winter months (Delaloye et al., 2008, 2010). In our study area neither rockglaciers nor protalus ramparts display significantly accelerated ~~surface displacement~~ creep in summer (Figure 10, A). The lack of

585 | seasonality and ~~small surface displacement~~the lower creeping rates compared to rockglaciers in the alps and the semi-arid
Andes (Strozzi et al., 2020) might be related to the semi-arid climate conditions (lack of moisture) and the short time-span of
three months with positive air temperatures in the Qugaquie basin (Zhang et al., 2013). Strozzi et al. (2020) figured out that
their highest rockglacier “Dos Lenguas” (4300 m a.s.l.) in the Andes is characterized by “less amplitude variations of the
annual cycle than observed for the Swiss Alps”. Hence, we hypothesize that the seasonality of rock glacier ~~surface~~
590 | ~~displacement~~creeping behaviour is less pronounced the lower the mean annual air temperatures and the shorter the timespans
of positive air temperature are. It seems that the magnitude of seasonal variations of the ~~surface displacement~~creeping rates
also decreases with a lower availability of moisture, because strongest seasonality is observed in moist regions such as the
Alps. Additionally, catchments in the Qugaquie basin are quite small for sediment release, so the extent of our rockglaciers is
limited by a small debris input. Probably for similar reasons, protalus ramparts investigated in this study creep with an
595 | average velocity of 1-3 mm/y, while comparable ~~surface displacement~~creeping rates for protalus ramparts range from 40
cm/y up to 100 cm/y in the Swiss alps (e.g., Scapozza, 2015).

The optical image-based process of rockglacier mapping and outlining is subject to several uncertainties, like the quality of
optical imagery and the rather subjective mapping style (Brardinoni et al., 2019). However, rockglacier inventories become
increasingly important due to their function as indicators of stored water resources and as indicators for climate change
600 | (Bolch and Gorbunov, 2014; Robson et al., 2020). A An IPA-working group was installed to reduce the uncertainties of such
inventories and to standardize mapping procedures. This year (2020) standardized guidelines were published on:
(<https://www3.unifr.ch/geo/geomorphology/en/research/ipa-action-group-rock-glacier/>)—which we followed in our mapping
procedure. Additionally with the opportunity to perform a field-based mapping a decrease of these uncertainties is likely
which we followed in our mapping procedure. Additionally with the opportunity to perform a field-based mapping a
605 | decrease of these uncertainties is likely.

Using rockglaciers and their long-term ice content as indicators for permafrost occurrence must be critically evaluated
because rockglaciers can overcome long distances and the terminus is far away from the routing zone (Bolch and Gorbunov,
2014; Halla et al., 2020). In this case rockglaciers are not suited for permafrost distribution assessment, because the ice-
debris mass creeps out of the continuous permafrost zone, as the rockglacier distribution in combination with modelled
610 | permafrost occurrence demonstrates in the northern Tien Shan (Bolch and Gorbunov, 2014). In our study, periglacial
landforms are characterized by a small extent and a low altitudinal range in extreme elevation. The rockglacier terminus is
close to the rooting zone, and they do not ~~span~~span a significant elevation range. Temperature data (MAAT of -6.8°),
~~elevated~~acquired at Zhadang glacier (Zhang et al., 2013), and ~~various different~~various large-scaled permafrost distribution maps
(Zou et al., 2017, Obu et al., 2019) suggest a high permafrost probability higher than 5400 m a.s.l. in the study area.
615 | Nevertheless a detailed, small scaled model of permafrost distribution would help to make a prognosis of permafrost
occurrence by localizing probabilities especially in lower areas of the catchment. “Permakart” considers topographic

parameters and different slope characteristics by using a topo-climatic key to handle the heterogeneity of high mountain areas (Schrott et al., 2012).

Often, rockglaciers can overcome long distances and the terminus is far away of the routing zone (Boleh and Gorbunov, 2014; Halla et al., 2020). In this case rockglaciers are not suited for permafrost distribution assessment, because the ice-debris mass creeps out of the continuous permafrost zone, like the rockglacier distribution in combination with modelled permafrost occurrence demonstrates in the northern Tien Shan (Boleh and Gorbunov, 2014). In our study periglacial landforms are characterized by a small extent and a low altitudinal range in extreme elevation. Rockglacier terminus is close to the rooting zone, so they do not exceed a significant elevation. Temperature data (MAAT of -6.8°), elevated at Zhadang glacier, and different, large scaled permafrost distribution maps (Zou et al., 2017; Obu et al., 2019) suggest a high permafrost probabilities higher than 5400 m a.s.l. in the study area. Nevertheless a detailed, small scaled model of permafrost distribution would help to make a detailed prognosis of permafrost occurrence by localizing probabilities especially in lower areas of the catchment. “Permakart” considers topographic parameters and different slope characteristics by using a topo-climatic key to handle the heterogeneity of high mountain areas (Schrott et al., 2012).

6 Conclusion and future work

In spite of the adverse logistical conditions in the study area, we were able to give insights to the cryosphere and to ~~constrain~~ assess a lower permafrost occurrence in the Qugaqie Basin on the TP using a multi-method approach. Thus, we add an important piece of information to the literature in a region where, due to its high altitude, ground truth data is usually difficult to obtain. Geomorphological mapping identifies the altitudinal distribution of periglacial landforms. ERT-measurements validate ice occurrence of one periglacial landform, a rockglacier. The activity of the periglacial landforms is derived from surface displacement ~~rates-analysis~~ of high resolution InSAR-data over three years. By combining the three findings we ~~conclude that~~ assess the lower boundary occurrence of probable permafrost ~~occurrence is located at an elevation around 5500 m a.s.l.~~ The main outcome is summarized as follows:

- The altitudinal distribution of periglacial landforms ranges between 5300 m a.s.l. and 5600 m a.s.l. and averages around 5500 m a.s.l. Protalus ramparts are more frequent ~~of~~ periglacial landforms, while rockglaciers have a larger extent and move faster.
- ERT measurements outside of blocky material of the periglacial landforms indicate ice-poor permafrost as ice-lenses (70 – 150 kΩm) at 5450 m a.s.l.
- ERT measurements on a rockglacier confirm perennial ice occurrence around 5500 m a.s.l. Resistivity values of more than 200 kΩm indicate ice-rich permafrost.

- Surface displacement ~~rates-analysis~~ extrapolates the status of active creeping to other permafrost related landforms. Especially rockglaciers show ~~creeping surface displacement~~ rates up to a maximum of 150 mm/y (average 27 mm/y). Protalus ramparts have much lower surface displacement rates (average 13 mm/y).
- ~~Seasonality of rockglacier creep surface displacement~~ is lacking probably due to low average temperatures and semi-arid climate conditions.
- ~~The lower limit of probable occurrence of permafrost is higher than 5300 – 5450 m. a.s.l.~~
- ~~The lower boundary occurrence of probable permafrost occurrence is supposed at an elevation around 5500/5400 m a.s.l. in the Qugaqie basin.~~

Our results illustrate the benefit of combining field-based and remote sensing techniques and recommend interdisciplinary approaches to geomorphological and geocryological issues. Nevertheless, the current results should be compared with a permafrost model of the study area in order to make a prognosis and zonation of the permafrost distribution. We also follow the suggestion by Strozzi et al. (2020) to include rock glaciers and the monitoring of rock glacier velocities as an essential climate variable in the Global Climate Observing System (GCOS) of the World Meteorological Organization due to the essential contribution of the results as climate sensitive parameters. As a next step, we plan to provide a rock glacier inventory for the Nyainqêntanglha Range based on InSAR-data as a status quo to understand the sensitivity and the vulnerability of high mountain cryosphere referred to climate warming.

Data availability. The data sets can be obtained on request to the authors.

Team list and Author contributions. JB designed the study, conducted fieldwork, collected and processed geomorphological and geophysical data, wrote the manuscript and conceptualized figures. ER was in charge of InSAR analyses. AH and BR conducted field logistics and data acquisition. AS, FZ, MG, RM and AH was responsible for funding acquisition. All authors contributed to the revision of the text.

The authors declare that they have no conflict of interest.

Acknowledgements

We thank all colleagues who contributed to this study, especially Bernd Wünnemann for fruitful discussions and valuable comments during field work, Jussi Baade for providing the TanDEM-X data (©DLR), Zhengliang Yu for the support in the field and Guoshuai Zhang and his team for logistical support at the Nam Co Station for Multisphere Observation and Research, Chinese Academy of Sciences. We thank Matthias Bucker for his insightful and highly valuable suggestions during this manuscript's revision phase. We thank the two anonymous reviewers for their comprehensive work on comments and suggestions to improve the manuscript.

This research is a contribution to the International Research Training Group (GRK 2309/1) “Geo-ecosystems in transition on the Tibetan Plateau (TransTiP)” funded by Deutsche Forschungsgemeinschaft (DFG).

References

- 680 | ~~racharya, S. (submitter), Shrestha, F., Bajracharya, S., Maharjan, S. and Guo, W.: GLIMS Glacier Database, ,~~
~~doi:10.7265/N5V98602, 2014.~~
- Ballantyne, C. K.: Periglacial geomorphology, first., edited by J. Wiley and & Sons, Wiley-Blackwell, Oxford., 2018.
- Barsch, D.: Rockglaciers: Indicators for the Present and Former Geoecology in High Mountain Environments, Springer
Berlin Heidelberg, Berlin, Heidelberg., 1996.
- 685 Berardino, P., Fornaro, G., Lanari, R. and Sansosti, E.: A new algorithm for surface deformation monitoring based on small
baseline differential SAR interferograms, IEEE Trans. Geosci. Remote Sens., 40(11), 2375–2383,
doi:10.1109/TGRS.2002.803792, 2002.
- Bibi, S., Wang, L., Li, X., Zhou, J., Chen, D. and Yao, T.: Climatic and associated cryospheric, biospheric, and hydrological
changes on the Tibetan Plateau: a review, Int. J. Climatol., 38, e1–e17, doi:10.1002/joc.5411, 2018.
- 690 Bolch, T. and Gorbunov, A. P.: Characteristics and Origin of Rock Glaciers in Northern Tien Shan
(Kazakhstan/Kyrgyzstan), Permafr. Periglac. Process., 25(4), 320–332, doi:10.1002/ppp.1825, 2014.
- Bolch, T., Yao, T., Kang, S., Buchroithner, M. F., Scherer, D., Maussion, F., Huintjes, E. and Schneider, C.: A glacier
inventory for the western Nyainqentanglha range and the Nam Co Basin, Tibet, and glacier changes 1976–2009,
Cryosphere, 4(3), 419–433, doi:10.5194/tc-4-419-2010, 2010.
- 695 Cao, B., Zhang, T., Wu, Q., Sheng, Y., Zhao, L. and Zou, D.: Brief communication : Evaluation and inter-comparisons of
Qinghai-Tibet Plateau permafrost maps based on a new inventory of field evidence, Cryosphere, 13(February), 511–519,
doi:10.5194/tc-13-511-2019, 2019.
- Chen, J., Zhao, L., Sheng, Y., Li, J., Wu, X., Du, E., Liu, G. and Pang, Q.: Some Characteristics of Permafrost and Its
Distribution in the Gaize Area on the Qinghai—Tibet Plateau, China, Arctic, Antarct. Alp. Res., 48(2), 395–409,
700 doi:10.1657/AAAR0014-023, 2016.
- Cheng, G. and Wu, T.: Responses of permafrost to climate change and their environmental significance, Qinghai-Tibet
Plateau, J. Geophys. Res. Earth Surf., 112(2), 1–10, doi:10.1029/2006JF000631, 2007.
- Cogley, G. (submitter), Moelg, N., Frey, H., Guo, W., Raup, B. H., Sakai, A., Liu, S., Nuimura, T., Paul, F. and Bolch, T.:
GLIMS Glacier Database, , doi:10.7265/N5V98602, 2015.
- 705 Crosetto, M., Monserrat, O., Cuevas-González, M., Devanthéry, N. and Crippa, B.: Persistent Scatterer Interferometry: A
review, ISPRS J. Photogramm. Remote Sens., 115, 78–89, doi:10.1016/j.isprsjprs.2015.10.011, 2016.

- Delaloye, R., Perruchoud, E., Bodin, X., Kääb, A., Kellerer-pirklbauer, A., Krainer, K., Lambiel, C., Roer, I. and Thibert, E.: Recent interannual variations of rock glacier creep in the European Alps, Proc. 9th Int. Conf. Permafrost, Fairbanks, Alaska, 2008.
- 710 Delaloye, R., Lambiel, C. and Gärtner-Roer, I.: Overview of rock glacier kinematics research in the Swiss Alps, *Geogr. Helv.*, 65(2), 135–145, doi:10.5194/gh-65-135-2010, 2010.
- Deline, P., Gruber, S., Delaloye, R., Fischer, L., Geertsema, M., Giardino, M., Hasler, A., Kirkbride, M., Krautblatter, M., Magnin, F., McColl, S., Ravel, L. and Schoeneich, P.: Ice Loss and Slope Stability in High-Mountain Regions, in *Snow and Ice-Related Hazards, Risks and Disasters*, edited by J. F. Shroder, W. Haeberli, and C. Whiteman, pp. 521–
- 715 561, Academic Press, Boston., 2015.
- Dong, G., Yi, C. and Caffee, M.: Be dating of boulders on moraines from the last glacial period in the Nyainqentanglha mountains , Tibet, *Sci. China Earth Sci.*, 57(2), 221–231, doi:10.1007/s11430-013-4794-z, 2014.
- Eckerstorfer, M., Eriksen, H. Ø., Rouyet, L., Christiansen, H. H., Lauknes, T. R. and Blikra, L. H.: Comparison of geomorphological field mapping and 2D-InSAR mapping of periglacial landscape activity at Nordnesfjellet, northern
- 720 Norway, *Earth Surf. Process. Landforms*, 43(10), 2147–2156, doi:10.1002/esp.4380, 2018.
- Emmert, A. and Kneisel, C.: Internal structure of two alpine rock glaciers investigated by quasi-3-D electrical resistivity imaging, *Cryosphere*, 11(2), 841–855, doi:10.5194/tc-11-841-2017, 2017.
- Esper Angillieri, M. Y.: Permafrost distribution map of San Juan Dry Andes (Argentina) based on rock glacier sites, *J. South Am. Earth Sci.*, 73, 42–49, doi:10.1016/j.jsames.2016.12.002, 2017.
- 725 Fort, M. and van Vliet-Lanoe, B.: Permafrost and periglacial environment of Western Tibet, *Landf. Anal.*, 5, 25–29, 2007.
- Frauenfelder, R., Allgöwer, B., Haeberli, W. and Hoelzle, M.: Permafrost Investigations With GIS - A Case Study in the Fletschhorn Area , Wallis , Swiss Alps, in *Seventh International Conference on Permafrost*, pp. 291–295., 1998.
- French, H. M.: *The Periglacial Environment*, fourth., John Wiley & Sons, Ltd, Oxford., 2017.
- Gruber, S., Fleiner, R., Guegan, E., Panday, P., Schmid, M. O., Stumm, D., Wester, P., Zhang, Y. and Zhao, L.: Review
- 730 article: Inferring permafrost and permafrost thaw in the mountains of the Hindu Kush Himalaya region, *Cryosphere*, 11(1), 81–99, doi:10.5194/tc-11-81-2017, 2017.
- Guglielmin, M., Ponti, S. and Forte, E.: The origins of Antarctic rock glaciers: periglacial or glacial features?, *Earth Surf. Process. Landforms*, 43(7), 1390–1402, doi:10.1002/esp.4320, 2018.
- Guo, W., Liu, S., Xu, J., Wu, L., Shangguan, D., Yao, X., Wei, J., Bao, W., Yu, P., Liu, Q. and Jiang, Z.: The second
- 735 Chinese glacier inventory: Data, methods and results, *J. Glaciol.*, 61(226), 357–372, doi:10.3189/2015JoG14J209, 2015.
- Häberli, W. and Vonder Mühll, D.: On the characteristics and possible origins of ice in rock glacier permafrost, *Zeitschrift für Geomorphol. Suppl.*, 104, 43–57, 1996.
- Halla, C., Blöthe, J. H., Baldi, C. T., Trombotto, D., Hilbich, C., Hauck, C. and Schrott, L.: Ice content and interannual water storage changes of an active rock glacier in the dry Andes of Argentina, *Cryosphere*, (March), 1–35, 2020.
- 740 Harris, S. A. and Pedersen, D. E.: Thermal regimes beneath coarse blocky materials, *Permafr. Periglac. Process.*, 9(2), 107–

- 120, doi:10.1002/(SICI)1099-1530(199804/06)9:2<107::AID-PPP277>3.0.CO;2-G, 1998.
- Hartmeyer, I., Keuschnig, M. and Schrott, L.: A scale-oriented approach for the long-term monitoring of ground thermal conditions in permafrost-affected rock faces , Kitzsteinhorn , Hohe Tauern Range , Austria, *Austrian J. Earth Sci.*, 105(May 2014), 2012.
- 745 Hauck, C. and Kneisel, C.: *Applied Geophysics in Periglacial Environments*, Cambridge University Press, Cambridge., 2008.
- Hauck, C. and Vonder Mühl, D.: Evaluation of geophysical techniques for application in mountain permafrost studies, *Zeitschrift für Geomorphol. N.F.*, 132, 161–190, 2003.
- Hauck, C., Bach, M. and Hilbich, C.: A four-phase model to quantify subsurface ice and water content in permafrost regions
 750 based on geophysical data sets, *Proc. 9th Int. Conf. Permafr.*, (Archie 1942), 675–680, 2008.
- Hedding, D. W.: Pronival ramparts: A review, *Prog. Phys. Geogr.*, 40(6), 835–855, doi:10.1177/0309133316678148, 2016.
- Hock, R., Rasul, G., Adler, C., Cáceres, B., Gruber, S., Hirabayashi, Y., Jackson, M., Kääb, A., Kang, S., Kutuzov, S., Milner, A., Molau, U., Morin, S., Orlove, B. and Steltzer, H.: High Mountain Areas, in *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, edited by H.-O. Pörtner, D. c. Roberts, V. Masson-Delmote, P. Zhai, M.
 755 Tignor, E. Poloczanska, K. Mintenbeck, A. Alegriá, M. Nicolai, A. Okem, J. Petzold, B. Rama, and N. M. Weyer, p. 94., 2019.
- Höllermann, P.: Blockgletscher als Mesoformen der Periglazialstufe, *Bonner Geogr. Abhandlungen*, 67, 1983.
- Hu, G., Zhao, L., Li, R., Wu, X., Wu, T., Xie, C., Zhu, X. and Hao, J.: Estimation of ground temperatures in permafrost regions of the Qinghai-Tibetan Plateau from climatic variables, *Theor. Appl. Climatol.*, (February), doi:10.1007/s00704-
 760 020-03135-1, 2020.
- Hu, J., Li, Z.-W., Li, J., Zhang, L., Ding, X.-L., Zhu, J.-J. and Sun, Q.: 3-D movement mapping of the alpine glacier in Qinghai-Tibetan Plateau by integrating D-InSAR, MAI and Offset-Tracking: Case study of the Dongkemadi Glacier, *Glob. Planet. Change*, 118, 62–68, doi:10.1016/j.gloplacha.2014.04.002, 2014.
- Immerzeel, W. W., Van Beek, L. P. H. and Bierkens, M. F. P.: Climate change will affect the asian water towers, *Science*
 765 (80-.), 328(5984), 1382–1385, doi:10.1126/science.1183188, 2010.
- Jarvis, A., Guevara, E., Reuter, H. I. and Nelson, A. D.: Hole-filled SRTM for the globe: version 4: data grid, 2008.
- Jones, D. B., Harrison, S., Anderson, K. and Whalley, W. B.: Rock glaciers and mountain hydrology: A review, *Earth-Science Rev.*, 193(March), 66–90, doi:10.1016/j.earscirev.2019.04.001, 2019.
- Kääb, A.: PERMAFROST AND PERIGLACIAL FEATURES | Rock Glaciers and Protalus Forms, in *Encyclopedia of Quaternary Science* (Second Edition), edited by S. A. Elias and C. J. Mock, pp. 535–541, Elsevier, Amsterdam., 2013.
 770
- Kang, S., Chen, F., Gao, T., Zhang, Y., Yang, W., Yu, W. and Yao, T.: Early onset of rainy season suppresses glacier melt: A case study on Zhadang glacier, Tibetan Plateau, *J. Glaciol.*, 55(192), 755–758, doi:10.3189/002214309789470978, 2009.
- Kapp, J. L. D. A., Harrison, T. M., Kapp, P., Grove, M., Lovera, O. M. and Lin, D.: Nyainqentanglha Shan: A window into

- 775 the tectonic, thermal, and geochemical evolution of the Lhasa block, southern Tibet, *J. Geophys. Res. Solid Earth*, 110(8), 1–23, doi:10.1029/2004JB003330, 2005.
- Keil, A., Berking, J., Mügler, I., Schütt, B., Schwalb, A. and Steeb, P.: Hydrological and geomorphological basin and catchment characteristics of Lake Nam Co, South-Central Tibet, *Quat. Int.*, 218(1–2), 118–130, doi:10.1016/j.quaint.2009.02.022, 2010.
- 780 Kidd, W. S. F., Yusheng, P., Chengfa, C., Coward, M. P., Dewey, J. F., Gansser, A., Molnar, P., Shackleton, R. M. and Yiyin, S.: Geological Mapping of the 1985 Chinese--British Tibetan (Xizang--Qinghai) Plateau Geotraverse Route, *Philos. Trans. R. Soc. A Math. Phys. Eng. Sci.*, 327(1594), 287–305, doi:10.1098/rsta.1988.0130, 1988.
- Kneisel, C. and Kääb, A.: Mountain permafrost dynamics within a recently exposed glacier forefield inferred by a combined geomorphological, geophysical and photogrammetrical approach, *Earth Surf. Process. Landforms*, 32(12), 1797–1810, doi:10.1002/esp.1488, 2007.
- 785 Kneisel, C., Lehmkuhl, F., Winkler, S., Tressel, E. and Schröder, H.: Legende für geomorphologische Kartierungen in Hochgebirgen (GMK Hochgebirge), *Trierer Geogr. Stud.*, 18, 1998.
- Kneisel, C., Hauck, C., Fortier, R. and Moorman, B.: Advances in Geophysical Methods for Permafrost Investigations, *Permafr. Periglac. Process.*, 178(March), 157–178, doi:10.1002/ppp, 2008.
- 790 Knight, J., Mitchell, W. A. and Rose, J.: Chapter Six - Geomorphological Field Mapping, in *Geomorphological Mapping*, vol. 15, edited by M. J. Smith, P. Paron, and J. S. Griffiths, pp. 151–187, Elsevier., 2011.
- Knight, J., Harrison, S. and Jones, D. B.: Rock glaciers and the geomorphological evolution of deglaciating mountains, *Geomorphology*, 324, 14–24, doi:10.1016/j.geomorph.2018.09.020, 2019.
- Krautblatter, M., Verleysdonk, S., Flores-Orozco, A. and Kemna, A.: Temperature-calibrated imaging of seasonal changes in permafrost rock walls by quantitative electrical resistivity tomography (Zugspitze, German/Austrian Alps), *J. Geophys. Res. Earth Surf.*, 115(2), 1–15, doi:10.1029/2008JF001209, 2010.
- 795 Li, J., Sheng, Y., Wu, J., Chen, J. and Zhang, X.: Probability distribution of permafrost along a transportation corridor in the northeastern Qinghai province of China, *Cold Reg. Sci. Technol.*, 59(1), 12–18, doi:https://doi.org/10.1016/j.coldregions.2009.05.012, 2009a.
- 800 Li, M., Ma, Y., Hu, Z., Ishikawa, H. and Oku, Y.: Snow distribution over the Namco lake area of the Tibetan Plateau, *Hydrol. Earth Syst. Sci.*, 13(11), 2023–2030, doi:10.5194/hess-13-2023-2009, 2009b.
- Liu, S. and Guo, W.: GLIMS Glacier Database, , doi:10.7265/N5V98602, 2014.
- Loke, M. H. and Barker, R. D.: Least-squares deconvolution of apparent resistivity pseudosections, *Geophysics*, 60(6), 1682–1690, doi:10.1190/1.1443900, 1995.
- 805 López-Martínez, J., Serrano, E., Schmid, T., Mink, S. and Linés, C.: Periglacial processes and landforms in the South Shetland Islands (northern Antarctic Peninsula region), *Geomorphology*, 155–156, 62–79, doi:10.1016/j.geomorph.2011.12.018, 2012.
- Ma, W., Shi, C., Wu, Q., Zhang, L. and Wu, Z.: Monitoring study on technology of the cooling roadbed in permafrost region

- of Qinghai–Tibet plateau, *Cold Reg. Sci. Technol.*, 44(1), 1–11, doi:10.1016/j.coldregions.2005.06.002, 2006.
- 810 Mewes, B., Hilbich, C., Delaloye, R. and Hauck, C.: Resolution capacity of geophysical monitoring regarding permafrost degradation induced by hydrological processes, *Cryosphere*, 11(6), 2957–2974, doi:10.5194/tc-11-2957-2017, 2017.
- Mudler, J., Hördt, A., Przyklenk, A., Fiandaca, G., Kumar Maurya, P. and Hauck, C.: Two-dimensional inversion of wideband spectral data from the capacitively coupled resistivity method - First applications in periglacial environments, *Cryosphere*, 13(9), 2439–2456, doi:10.5194/tc-13-2439-2019, 2019.
- 815 Mügler, I., Gleixner, G., Günther, F., Mäusbacher, R., Daut, G., Schütt, B., Berking, J., Schwalb, A., Schwark, L., Xu, B., Yao, T., Zhu, L. and Yi, C.: A multi-proxy approach to reconstruct hydrological changes and Holocene climate development of Nam Co, Central Tibet, *J. Paleolimnol.*, 43(4), 625–648, doi:10.1007/s10933-009-9357-0, 2010.
- Von der Mühll, D., Hauck, C. and Gubler, H.: Mapping of mountain permafrost using geophysical methods, *Prog. Phys. Geogr.*, 26(4), 643–660, 2002.
- 820 N El Sayed, A., SM Barseem, M., M Ezz El Deen, H. and A Ezz El Din, H.: Using of Geo-electrical and Geochemical Techniques to Investigate the Change in Ground Water Quality-South West El Khtatbah City - Cairo-Alexandria Desert Road, Egypt, *Adv. Appl. Sci. Res.*, 8, 2018.
- Notti, D., Herrera, G., Bianchini, S., Meisina, C., García-Davalillo, J. C. and Zucca, F.: A methodology for improving landslide PSI data analysis, *Int. J. Remote Sens.*, 35(6), 2186–2214, doi:10.1080/01431161.2014.889864, 2014.
- 825 Obu, J., Westermann, S., Bartsch, A., Berdnikov, N., Christiansen, H. H., Dashtseren, A., Delaloye, R., Elberling, B., Etzelmüller, B., Kholodov, A., Khomutov, A., Kääb, A., Leibman, M. O., Lewkowicz, A. G., Panda, S. K., Romanovsky, V., Way, R. G., Westergaard-Nielsen, A., Wu, T., Yamkhin, J. and Zou, D.: Northern Hemisphere permafrost map based on TTOP modelling for 2000–2016 at 1 km² scale, *Earth-Science Rev.*, 193(October 2018), 299–316, doi:10.1016/j.earscirev.2019.04.023, 2019.
- 830 Osmanoğlu, B., Sunar, F., Wdowinski, S. and Cabral-Cano, E.: Time series analysis of InSAR data: Methods and trends, *ISPRS J. Photogramm. Remote Sens.*, 115, 90–102, doi:10.1016/j.isprsjprs.2015.10.003, 2016.
- Otto, J. C. and Dikau, R.: Symbols for geomorphologic mapping in high mountains for ArcGIS; Geomorphologic System Analysis of a High Mountain Valley in the Swiss Alps., *Zeitschrift für Geomorphol.* 48 323-341 [online] Available from: <http://geomorphology.sbg.ac.at/research/map-symbols/> (Accessed 15 February 2018), 2008.
- 835 Otto, J. C. and Smith, M. J.: Geomorphological mapping, in *Geomorphological Techniques* (Online Edition), vol. 2.1.6, edited by Cook, S.J., L. E. Clarke, and J. M. Nield, pp. 344–345, British Society for Geomorphology, London., 2013.
- Ran, Y., Li, X., Cheng, G., Zhang, T., Wu, Q., Jin, H. and Jin, R.: Distribution of Permafrost in China: An Overview of Existing Permafrost Maps, *Permafr. Periglac. Process.*, 23(4), 322–333, doi:10.1002/ppp.1756, 2012.
- Reinosch, E., Buckel, J., Dong, J., Gerke, M., Baade, J. and Riedel, B.: InSAR time series analysis of seasonal surface displacement dynamics on the Tibetan Plateau, *Cryosph.*, 14, 1633–1650, doi:10.5194/tc-2019-262, 2020.
- 840 Reynolds, J. M.: *An Introduction to Applied and Environmental Geophysics*, 2nd ed., edited by J. Reynolds, Wiley/Blackwell, Oxford., 2011.

- Rosset, E., Hilbich, C., Schneider, S. and Hauck, C.: Automatic filtering of ERT monitoring data in mountain permafrost, *Near Surf. Geophys.*, 11(4), 423–433, doi:10.3997/1873-0604.2013003, 2013.
- 845 Scapoza, C.: Investigation on protalus ramparts in the Swiss Alps, *Geogr. Helv.*, 70(2), 135–139, doi:10.5194/gh-70-135-2015, 2015.
- Schmid, M. O., Baral, P., Gruber, S., Shahi, S., Shrestha, T., Stumm, D. and Wester, P.: Assessment of permafrost distribution maps in the Hindu Kush Himalayan region using rock glaciers mapped in Google Earth, *Cryosphere*, 9(6), 2089–2099, doi:10.5194/tc-9-2089-2015, 2015.
- 850 Schrott, L.: Some geomorphological-hydrological aspects of rock glaciers in the Andes (San Juan, Argentina), *Zeitschrift für Geomorphol.*, 104, 161–173, 1996.
- Schrott, L. and Sass, O.: Application of field geophysics in geomorphology: Advances and limitations exemplified by case studies, *Geomorphology*, 93(1–2), 55–73, doi:10.1016/j.geomorph.2006.12.024, 2008.
- Schrott, L., Otto, J. C. and Keller, F.: Modelling alpine permafrost distribution in the hohe tauern region, Austria, *Austrian J. Earth Sci.*, 105(2), 169–183, 2012.
- 855 Song, C., Wang, G., Mao, T., Dai, J. and Yang, D.: Linkage between permafrost distribution and river runoff changes across the Arctic and the Tibetan Plateau, *Sci. China Earth Sci.*, 63(2), 292–302, doi:10.1007/s11430-018-9383-6, 2020.
- Sowter, A., Bateson, L., Strange, P., Ambrose, K. and Syafiudin, M. F.: DInSAR estimation of land motion using intermittent coherence with application to the South Derbyshire and Leicestershire coalfields, *Remote Sens. Lett.*, 4(10), 860 979–987, doi:10.1080/2150704X.2013.823673, 2013.
- Strozzi, T., Caduff, R., Jones, N., Barboux, C., Delaloye, R., Bodin, X., Kääb, A., Mätzler, E. and Schrott, L.: Monitoring Rock Glacier Kinematics with Satellite Synthetic Aperture Radar, *Remote Sens.*, 12(3), 559, doi:10.3390/rs12030559, 2020.
- Sun, Z., Zhao, L., Hu, G., Qiao, Y., Du, E., Zou, D. and Xie, C.: Modeling permafrost changes on the Qinghai–Tibetan plateau from 1966 to 2100: A case study from two boreholes along the Qinghai–Tibet engineering corridor, *Permafr. Periglac. Process.*, 31(1), 156–171, doi:10.1002/ppp.2022, 2020.
- 865 Thompson, S. S., Kulessa, B., Benn, D. I. and Mertes, J. R.: Anatomy of terminal moraine segments and implied lake stability on Ngozumpa Glacier, Nepal, from electrical resistivity tomography (ERT), *Sci. Rep.*, 7(January), 1–12, doi:10.1038/srep46766, 2017.
- 870 Tian, K., Liu, J., Kang, S. and Li, C.: A Primary Study of the Environment of Frozen Ground in the Nam Co Basin, Tibet, *Adv. earth Sci.*, 21(12), 1324–1332, 2006.
- Vanhala, H., Lintinen, P. and Ojala, A.: Electrical resistivity study of permafrost on Ridnitšohkka fell in northwest Lapland, Finland, *Geophysica*, 45(1–2), 103–118, 2009.
- Villarreal, C. D., Beliveau, G. T., Forte, A. P., Monserrat, O. and Morvillo, M.: DInSAR for a regional inventory of active rock glaciers in the Dry Andes Mountains of Argentina and Chile with sentinel-1 data, *Remote Sens.*, 10(10), 1–21, doi:10.3390/rs10101588, 2018.

- von Wakonigg, H.: Unterkühlte Schutthalden (Undercooled talus) (in German), Beiträge zur Perma-frostforsch. Österreich. Arb. aus dem Inst. für Geogr. der Karl-Franzens Univ., 33, 209–223, 1996.
- Wang, B. and French, H. M.: Permafrost on the Tibet Plateau, China, Quat. Sci. Rev., 14(3), 255–274, doi:10.1016/0277-880 3791(95)00006-B, 1995.
- Wang, J., Zhu, L., Daut, G., Ju, J., Lin, X., Wang, Y. and Zhen, X.: Investigation of bathymetry and water quality of Lake Nam Co, the largest lake on the central Tibetan Plateau, China, Limnology, 10(2), 149–158, doi:10.1007/s10201-009-0266-8, 2009.
- Wang, T., Yang, D., Fang, B., Yang, W., Qin, Y. and Wang, Y.: Data-driven mapping of the spatial distribution and 885 potential changes of frozen ground over the Tibetan Plateau, Sci. Total Environ., 649, 515–525, doi:10.1016/j.scitotenv.2018.08.369, 2019.
- Wang, Y., Fu, Z., Lu, X., Qin, S., Wang, H. and Wang, X.: Imaging of the internal structure of permafrost in the tibetan plateau using ground penetrating radar, Electron., 9(1), doi:10.3390/electronics9010056, 2020.
- Wei, D., -Ri, X., Wang, Y., Wang, Y., Liu, Y. and Yao, T.: Responses of CO₂, CH₄ and N₂O fluxes to livestock 890 exclosure in an alpine steppe on the Tibetan Plateau, China, Plant Soil, 359(1–2), 45–55, doi:10.1007/s11104-011-1105-3, 2012.
- Wu, T., Li, S., Cheng, G. and Nan, Z.: Using ground-penetrating radar to detect permafrost degradation in the northern limit of permafrost on the Tibetan Plateau, Cold Reg. Sci. Technol., 41(3), 211–219, doi:10.1016/j.coldregions.2004.10.006, 2005.
- Yague-Martinez, N., Rodriguez Gonzalez, F., Brcic, R., Shau, R., Geudtner, D., Eineder, M. and Bamler, R.: Interferometric 895 processing of SLC Sentinel-1 TOPS data, IEEE Trans. Geosci. Remote Sens., 54(4), 2220–2234, doi:10.1109/TGRS.2015.2497902, 2016.
- Yang, M., Nelson, F. E., Shiklomanov, N. I., Guo, D. and Wan, G.: Permafrost degradation and its environmental effects on the Tibetan Plateau: A review of recent research, Earth-Science Rev., 103(1–2), 31–44, 900 doi:10.1016/j.earscirev.2010.07.002, 2010.
- Yao, T., Guo, X., Thompson, L., Duan, K., Wang, N., Pu, J., Xu, B., Yang, X. and Sun, W.: $\delta^{18}\text{O}$ record and temperature change over the past 100 years in ice cores on the Tibetan Plateau, Sci. China, Ser. D Earth Sci., 49(1), 1–9, doi:10.1007/s11430-004-5096-2, 2006.
- Yao, T., Thompson, L., Yang, W., Yu, W., Gao, Y., Guo, X., Yang, X., Duan, K., Zhao, H., Xu, B., Pu, J., Lu, A., Xiang, Y., 905 Kattel, D. B. and Joswiak, D.: Different glacier status with atmospheric circulations in Tibetan Plateau and surroundings, Nat. Clim. Chang., 2(9), 663–667, doi:10.1038/nclimate1580, 2012.
- Yao, T., Masson-Delmotte, V., Gao, J., Yu, W., Yang, X., Risi, C., Sturm, C., Werner, M., Zhao, H., He, Y., Ren, W., Tian, L., Shi, C. and Hou, S.: A review of climatic controls on $\delta^{18}\text{O}$ in precipitation over the Tibetan Plateau: Observations and simulations, Rev. Geophys., 51(4), 525–548, doi:10.1002/rog.20023, 2013.
- 910 You, Y., Yu, Q., Pan, X., Wang, X. and Guo, L.: Application of electrical resistivity tomography in investigating depth of

- permafrost base and permafrost structure in Tibetan Plateau, *Cold Reg. Sci. Technol.*, 87, 19–26, doi:10.1016/j.coldregions.2012.11.004, 2013.
- You, Y., Yu, Q., Pan, X., Wang, X. and Guo, L.: Geophysical Imaging of Permafrost and Talik Configuration Beneath a Thermokarst Lake, *Permafr. Periglac. Process.*, 28(2), 470–476, doi:10.1002/ppp.1938, 2017.
- 915 Yu, W., Han, F., Liu, W. and Harris, S. A.: Geohazards and thermal regime analysis of oil pipeline along the Qinghai–Tibet Plateau Engineering Corridor, *Nat. Hazards*, 83(1), 193–209, doi:10.1007/s11069-016-2308-y, 2016.
- Yu, Z., Wu, G., Keys, L., Li, F., Yan, N., Qu, D. and Liu, X.: Seasonal variation of chemical weathering and its controlling factors in two alpine catchments, Nam Co basin, central Tibetan Plateau, *J. Hydrol.*, 576, 381–395, doi:10.1016/J.JHYDROL.2019.06.042, 2019.
- 920 Zhang, G., Kang, S., Fujita, K., Huintjes, E., Xu, J., Yamazaki, T., Haginoya, S., Wei, Y., Scherer, D., Schneider, C. and Yao, T.: Energy and mass balance of Zhadang glacier surface, central Tibetan Plateau, *J. Glaciol.*, 59(213), 137–148, doi:10.3189/2013JoG12J152, 2013.
- Zhang, Z. and Wu, Q.: Thermal hazards zonation and permafrost change over the Qinghai-Tibet Plateau, *Nat. Hazards*, 61(2), 403–423, doi:10.1007/s11069-011-9923-4, 2012.
- 925 Zou, D., Zhao, L., Sheng, Y., Chen, J., Hu, G., Wu, T., Wu, J., Xie, C., Wu, X., Pang, Q., Wang, W., Du, E., Li, W., Liu, G., Li, J., Qin, Y., Qiao, Y., Wang, Z., Shi, J. and Cheng, G.: A new map of permafrost distribution on the Tibetan Plateau, *Cryosphere*, 11(6), 2527–2542, doi:10.5194/tc-11-2527-2017, 2017.