



1 **Observation-derived ice growth curves show patterns and trends in maximum**  
2 **ice thickness and safe travel duration of Alaskan lakes and rivers**

3 Christopher D. Arp<sup>1</sup>, Jessica E. Cherry<sup>1,2</sup>, Dana R.N. Brown<sup>3</sup>, Allen C. Bondurant<sup>1</sup>, and Karen L.  
4 Endres<sup>4</sup>

5 <sup>1</sup>Water and Environmental Research Center, University of Alaska Fairbanks, Fairbanks, AK  
6 99775, USA

7 <sup>2</sup>Alaska-Pacific River Forecast Center, National Weather Service, Anchorage, AK 99502, USA

8 <sup>3</sup>Institute of Arctic Biology, University of Alaska Fairbanks, Fairbanks, AK 99775, USA

9 <sup>4</sup>Alaska-Pacific River Forecast Center, National Weather Service, Fairbanks, AK 99775, USA  
10

11 *Correspondence to:* Chris Arp ([cdarp@alaska.edu](mailto:cdarp@alaska.edu))

12

13 **Abstract.** The formation, growth, and decay of freshwater ice on lakes and rivers are  
14 fundamental processes of northern regions with wide ranging implications for socio-ecological  
15 systems. Ice thickness at the end of winter is perhaps the best integration of cold-season weather  
16 and climate, while the duration of thick and growing ice cover is a useful indicator for the winter  
17 travel and recreation season. Both maximum ice thickness (MIT) and ice travel duration (ITD)  
18 can be estimated from temperature-driven ice growth curves fit to ice thickness observations. We  
19 simulated and analyzed ice growth curves based on ice thickness data collected from a range of  
20 observation programs throughout Alaska spanning the past 20 – 60 years to understand patterns  
21 and trends in lake and river ice. Results suggest reductions in MIT (thinning) in several northern,  
22 interior, and coastal regions of Alaska and overall greater interannual variability in rivers  
23 compared to lakes. Interior regions generally showed less variability in MIT and even slightly  
24 increasing trends in at least one river site. Average ITD ranged from 214 days in the northern-  
25 most lakes to 114 days across southern-most lakes with significant decreases in duration for half  
26 of sites. River ITD showed low regional variability, but high interannual variability,



1 underscoring the challenges with predicting seasonally-consistent river travel. Standardization  
2 and analysis of these ice observation data provide a comprehensive summary for understanding  
3 changes in winter climate and its impact on freshwater ice services.

4

## 5 **1 Introduction**

6 Arctic amplification is an enhanced warming response in high latitudes relative to increasing  
7 global temperature (Serreze and Barry 2011). Though not yet completely understood, sea ice  
8 decline and associated climate feedbacks are considered to be major drivers of this process  
9 (Serreze and Francis 2006). A salient feature of arctic amplification is greater warming during  
10 winter, which has been strikingly apparent in Alaska during recent years (Wendler et al. 2014,  
11 Walsh and Brettschneider 2019). Terrestrial landscape responses to winter climate change are  
12 perhaps most quantifiable in ice formation and growth on lakes and rivers, and readily described  
13 by ice thickening through the winter (Allen 1977, Engram et al. 2018). Freshwater ice thickness  
14 and duration may function as robust integrators of winter climate, as they respond both to  
15 changes in air temperature and snow accumulation. Additionally, ice thickness and its duration  
16 have important implications for winter travel, subsistence, and recreation in Alaska and across  
17 the Arctic (Brown and Duguay 2010, Schneider et al. 2013, Cold et al. 2020).

18       Some of the longest ice thickness records come from the Barrow Peninsula in northern-  
19 most Alaska where lake ice historically grew greater than 2-m thick in some years by winter's  
20 end as recorded in the 1970s (Weeks et al. 1978) and 1990s (Zhang and Jeffries 2000). In recent  
21 years however, MIT did not exceed 1.2-m in snowy winters, when ice was well insulated and air  
22 temperatures were unusually warm (Arp et al. 2018). The impacts of arctic amplification on



1 freshwater ice should be most evident in this northern coastal region where Alexeev et al. (2016)  
2 demonstrated a linkage between sea ice extent and lake ice growth. This long, yet intermittent,  
3 record of lake ice thickness in northern Alaska comes from a variety of observation efforts  
4 including community-based monitoring facilitated by government agencies (Bilello 1980) and  
5 school science programs (Morris and Jeffries 2010). These same community-based monitoring  
6 programs also contributed to shorter, but still highly valuable, ice thickness records for other  
7 lakes and river in Alaska, which have been maintained and extended by the National Weather  
8 Service's Alaska Pacific River Forecast Center (APRFC). APRFC's interest in ice thickness has  
9 primarily been to facilitate river breakup and ice jam forecasting, but is also of value for  
10 informing safe ice travel, as fall through accidents have increased in recent years (Fleischer et al.  
11 2014).

12 In contrast to ice thickness observations, records and analysis of ice phenology (timing of  
13 freeze-up and break-up) are often long and abundant for many northern regions, likely owing to  
14 the ease of observing water-to-ice transition timing from shorelines, aircraft, or satellites (Brown  
15 and Duguay 2010). Rigorous satellite-based observations show distinct trends towards earlier  
16 break-up on both rivers (Cooley and Palvesky 2016) and lakes (Smejkalova et al. 2017) in  
17 Alaska and longer-term observer records from other northern regions show similar patterns (e.g.,  
18 Magnuson et al. 2000, Weyhenmeyer et al. 2011). Detection of trends in freeze-up timing are  
19 often less certain (Brown and Duguay 2010), though recent analyses suggest late freeze-up  
20 contributions to reduced ice cover duration on lakes (Sharma et al. 2019) and rivers (Yang et al.  
21 2019). Brown et al. (2018) tracked both freeze-up and break-up progression in Alaskan rivers,  
22 highlighting the varying stages of ice formation and decay processes relative to access for over  
23 ice travel, and suggesting the need to move beyond ice phenology as an exact *to-the-day* event.



1 Tracking changes in ice thickness through the winter cold season provides the simplest means of  
2 quantifying this continuum, though these data are distinctly more field intensive.

3       Reported analysis of ice thickness datasets often lack detection of thinning trends despite  
4 progressive winter warming, which may be due to high interannual variability in snowfall and  
5 the dominant role of snow insulation on ice growth (Brown and Duguay 2010). Yet the majority  
6 of studies analyzing ice thickness trends that we are aware of came before unprecedented warm  
7 winters during the last decade in Alaska (Wendler et al. 2014, Walsh and Brettschneider 2019).  
8 We also suspect that the inherent nature of ice thickness data collection, in which the timing of  
9 late winter measurement may vary from year to year relative to slight shifts in the ice growth  
10 season, adds additional noise in detecting trends. Additionally, ice thickness can vary  
11 significantly within a small area depending on snowcover, measurement protocols that differ  
12 among programs, and because many ice thickness observations often depend on volunteers with  
13 high turnover and minimal oversight. Large data gaps in records also make it difficult to  
14 ascertain trends at some long-term sites (Cherry 2019). For these reasons, we are motivated to  
15 standardize ice thickness according to ice growth curves informed by field observations and  
16 calculate relevant metrics, Maximum Ice Thickness (MIT) and Ice Travel Duration (ITD), as  
17 well as to merge analyses of both river and lake ice in Alaska.

18       In this study, we organized winter lake and river ice thickness observations from a variety  
19 of sources to provide an updated analysis of patterns and trends of ice growth in Alaska from  
20 1962 to present. Fitting these data to air temperature-driven ice growth curves simulated with the  
21 Stefan equation (Stefan 1891, Jumikis 1977) provides a robust seasonal estimation of changing  
22 ice thickness for multiple sites with proximate climate data. The ice growth curves were used to  
23 estimate MIT and ITD for four lakes and four rivers distributed across Alaska, with records




1 spanning 20 years or greater, to provide a summary of recent changes in ice of climatic and  
2 societal relevance. Several shorter records are also presented for spatial comparison.

3

## 4 **2 Background and Methods**

### 5 **2.1 Study region and waterbodies**

6 The diverse geographic and northern climatic setting of the State of Alaska presents a fascinating  
7 study region to observe freshwater ice (Arp and Jones 2009, Arp et al. 2013). Even though the  
8 state of Alaska is a geopolitical unit, its vast size (1.5 million km<sup>2</sup>), expansive latitudinal extent  
9 (18°), wide longitudinal extent (58°), lengthy coastline (54,500 km), and complex tectonic  
10 setting create a largely contiguous landscape with several large mountain ranges and expansive  
11 river valleys (Figure 1). These geographical attributes of Alaska interact with climate, glacial  
12 history, and soil conditions (particularly permafrost) to create many lakes (> 400,000) and  
13 extensive and varied river networks (>150,000 km) (Arp and Jones 2009). In contrast to  
14 waterbody extents, Alaska road network is relatively short (<25,000 km) and not connected to  
15 the majority of towns and villages, which limit opportunities to maintain long-term observations  
16 of most waterbodies. Thus the majority of sites with long-term ice observation data are  
17 associated with large towns along roadways or villages adjacent to rivers or lakes (Table 1).  
18 Another restriction for this study was proximity to reliable long-term air temperature data from  
19 weather stations, which are typically associated with larger airports.



1

2 **Figure 1.** Map of the State of Alaska with all observation locations by waterbody and nearest  
3 community indicated and average maximum ice thickness (cm) for each period of record (in  
4 parenthesis). Several shorter term records (<20 years) are shown here for additional context  
5 which are not presented in the main analysis (\* indicates that observations are based on multiple  
6 lakes or rivers within a region; 300 m DEM hillshade is USGS data in the U.S. public domain).



1 **Table 1.** Summary of major ice observation stations and corresponding ice thickness records. Ice  
 2 growth curves simulated using the Stefan equation and the average  $\alpha$  coefficient ( $\text{cm}^0 \text{C}^{-1/2} \text{d}^{-1/2}$ )  
 3 and accumulated freezing degree days (AFDD) at MIT ( $^0 \text{C d}$ ) are reported for each station. Air  
 4 temperature data from National Weather Service stations are indicated by station codes.

Water Body	Community	Region	Period	Years Observed	$\alpha$ (ave)	AFDD (ave)	Weather Station
multiple lakes	Utiqagvik	North Slope	1962-2019	26	2.6	-4166	Barrow-PABR
Koyukuk River	Bettles	Interior	1968-2019	48	1.9	-3265	Bettles-PABT
Yukon River	Eagle	Interior	1999-2019	21	2.4	-2594	Eagle-PAEG
Smith Lake	Fairbanks	Interior	1965-2019	53	1.3	-2895	Fairbanks-PAFA
Tanana River	Nenana	Interior	1989-2019	31	2.2	-2516	Nenana-PANN*
Lake Minchumina	Minchumina	Interior	1997-2019	18	2.0	-2542	Nenana-PANN, McGrath-PAMC
Lake Hood	Anchorage	South-Central	1997-2019	22	2.6	-921	Anchorage-PANC
Kuskokwim River	Bethel	Western	1962-2019	48	3.3	-1826	Bethel-PABE

5 \*Missing data derived from relationship to Fairbanks-PAFA.

6

## 7 **2.2 Alaska ice observation programs**

8 Scientific records of ice thickness measurements in Alaska date back to the 1<sup>st</sup> International Polar  
 9 Year in 1882-83 for lakes located near Barrow (Ray 1885). Starting in the early 1960's the U.S.  
 10 Army Corps of Engineer's Cold Regions Research and Engineering Laboratory (CRREL)  
 11 established ice observing stations in coordination with Canadian and U.S. government agencies  
 12 (Bilello 1980). Stations included 26 lakes and rivers in Alaska where ice thickness data were  
 13 collected at weekly intervals until at least 1974. Observations made by local residents (Alaska  
 14 Natives, homesteaders, lodgekeepers, teachers, and clergy) for up to 15 years, in some cases,  
 15 provided valuable data for developing ice growth and decay models as reported in Bilello (1980)  
 16 and other CRREL reports. Perhaps as importantly, these data provide a comprehensive summary



1 of ice thickness and its variability over climatically diverse region of the Arctic and sub-Arctic  
2 before notable climate warming. The majority of Alaska ice thickness data, as well as snow  
3 depth data, from this program are now archived with the Arctic Data Center (ADC) (Bilello  
4 2019). Observation protocols are not always well described in the reports for this program, but  
5 likely relied on narrow gauge hand-augers and tapes similar to current procedures, and the ice  
6 thickness observations per date may have come from single point measurements.

7 A more recent winter observation program, Alaska Snow and Ice Observation Network  
8 (ALISON), pioneered the integration of science education with snow and ice physics in Alaska  
9 schools from 1999 to 2010 (Morris and Jeffries 2010). While learning about snow and ice  
10 physics, students and teachers collected valuable datasets across a wide range geographic and  
11 climatic settings, including at least 17 lakes ranging from the Barrow Peninsula in the north to  
12 the Kenai Peninsula in the south. Several of these sites overlap with CRREL stations, thus  
13 providing an opportunity for extension of records and temporal comparisons. ALISON's focus  
14 on snow depth, density, and heat flux provided additional data of value for understanding  
15 variability and modeling ice growth (Gould et al. 2005, Jeffries et al. 2005). ALISON datasets  
16 are also archived at ADC (Morris and Jeffries 2019). Often up to 20 snow measurements were  
17 recorded per sampling interval for this program, while ice thickness was typically recorded from  
18 a single point using a thermal resistance (heated) wire known as a TWIT (Thermal Wire Ice-  
19 thickness Thingy) (M. Jeffries personal communications). The TWIT was designed to minimize  
20 local snow disturbance that would affect subsequent measurements, but also represents a single  
21 point measurement of ice thickness.

22 An important Alaska ice record focused on breakup timing of a single river reach is the  
23 Nenana Ice Classic (NIC), where a tripod is set on the Tanana River by the community of





1 Nenana each year and cabled to a clock to record the exact data of river breakup (Sagarin and  
2 Micheli 2001). This community-based monitoring program dates back to 1917 and each year  
3 thousands of people submit breakup guesses into a pool with the closest participants taking home  
4 >\$300,000 US dollars in recent years. Regular ice thickness observations, dating back to at least  
5 1989, have been made by community members who run the NIC, and are published to provide  
6 contestants with additional information to aid in guesses.


7 A fourth and also contemporary Alaska-wide lake and river data source is provided by  
8 the NWS Alaska-Pacific River Forecast Center (APRFC). Prior to the establishment of the  
9 APRFC, the National Weather Services weather forecast offices collected or solicited these data  
10 and maintained them at what is now the National Center for Environmental Information. Historic  
11 monthly ice thickness measurements have been compiled from a variety of sources including the  
12 CRREL dataset (Bilello 1980) and contemporary observation come from NWS scientists and  
13 paid and volunteer observers in remote communities throughout the state. Much of this ice  
14 thickness data for both rivers and lakes is used in operational forecasting of river conditions  
15 specific to river break-up and ice jam flooding predictions. APRFC ice thickness data are  
16 available online (<https://www.weather.gov/aprfc/IceThickness>). Current protocol for APRFC are to  
17 collect single-auger hole and tape observations near the start of the month from November to  
18 March near the same location at each waterbody below undisturbed snow.

19 Lastly, a regional lake ice observation program focused on Alaska's North Slope began in  
20 2012 called the CircumArctic Lakes Observation Network (CALON) (Hinkel et al. 2012). This  
21 project supported by the National Science Foundation collected consistent late winter ice  
22 thickness data from sets of six lakes in each of ten study areas arrayed from the Brooks Range  
23 foothills to the Beaufort Sea coastal plain until 2017. Several of these study areas were



1 associated with field camps or other long-term research locations where prior lake ice data  
2 existed or was expected to continue after CALON concluded. Observations were made at the  
3 same location on each lake every year by drilling 3-5 holes at regular spacing and recording  
4 snow data in association with individual ice thickness measurements. ADC has CALON datasets  
5 archived for ice thickness (Arp 2018a) and snow characteristics (Arp 2018b) separately.

6 Addressing the issue of data comparability among programs is relevant to this study. It is  
7 of course expected that higher numbers of samples per site correspond to more accurate ice  
8 thickness measurements such that CALON protocols have higher accuracy than some of the  
9 other ice observation programs described. Analysis conducted in a 36-member sampling protocol  
10 on two lakes and two rivers (one each in the Interior and the North Slope) showed that more  
11 samples reduced error (Figure 2). This analysis suggests that making one observation ( $n=1$ )  
12 results in potential error up to 18, 10, 8, and 4 cm from the mean for Interior rivers, North Slope  
13 rivers, North Slope lakes, and Interior lakes, respectively. Three observations ( $n=3$ ) results in  
14 potential error up to 9, 6, 5, and 3 cm from the mean for Interior rivers, North Slope rivers, North  
15 Slope lakes, and Interior lakes, respectively. While this analysis provides guidance for  
16 comparing the quality of differing ice observation approaches, we also suggest that professional  
17 experience or local knowledge in selecting locations of representative ice thickness may well  
18 overcome low sample size in many cases. Our expectation is that observers reporting to APRFC  
19 have such experience, as do observers from previous programs.



1

2 **Figure 2.** Margin of error analysis for Interior and North Slope lakes and rivers from samples of  
 3 36 evenly spaced measurements during the winter of 2018-19.

4

5 **2.3 Ice growth curve simulation**

6 For all records with late winter ice observations, we estimated MIT using the Stefan ice growth  
 7 model (modified Stefan equation) (1):

$$8 \quad Z_{ice} = \alpha \sqrt{AFDD} \quad (1)$$

9 where  $\alpha$  is the ice growth (thermal insulation) coefficient in  $\text{cm}^0 \text{C}^{-1/2} \text{d}^{-1/2}$  and AFDD is  
 10 accumulated freezing degree days in  $^0 \text{C d}$  are used to estimate ice thickness ( $Z_{ice}$ ) in cm on a  
 11 daily time-step (Stefan 1891, Jumikis 1977). Mean daily air temperature data used to force this



1 model was acquired from the nearest National Weather Service (NWS) station, the majority of  
2 which were within 20 km of the water body where ice was monitored (Table 1). In the cases of  
3 Shageluk and Lake Minchumina (Figure 1), no adjacent NWS or other meteorological stations  
4 air temperature records were available and we averaged the records between two nearest stations  
5 set in opposing directions. Summation of AFDD was started at the date of estimated ice growth  
6 initiation each winter. When early ice thickness observations (i.e.  $Z_{ice} < 10$  cm) were available or  
7 actual observation of the day of ice initiation, these data guided selecting this date. When these  
8 data were not available, the more common case, we selected the date according to three  
9 consecutive days with mean daily temperature  $< 0^{\circ}$  C for lakes, which is based on previous  
10 camera and sensor observations (Arp et al. 2013) and six consecutive days  $< 0^{\circ}$  C for rivers. The  
11 later criteria for rivers is based on information APRFC observers, but is more limited and thus  
12 considered much more uncertain and a potential source of error in ice growth modeling on rivers.  
13 Both air temperature criteria follow guidance in Bilello et al. (1964) and Ashton (1989) on ice  
14 cover initiation and early growth. Complementing the ice growth model (1), is an ice decay  
15 model developed by Bilello (1980) (2):

$$16 \quad Z_{ice} = \alpha' ATDD \quad (2)$$

17 where  $\alpha'$  is the ice decay coefficient in  $\text{cm}^{\circ} \text{C}^{-1} \text{d}^{-1}$  and ATDD is accumulated thawing degree  
18 days in  $^{\circ} \text{C d}$  with a  $0^{\circ} \text{C}$  and also with summation beginning at the day of ice growth initiation.  
19 Equation 2 is calculated concurrently with equation 1 to estimate the ice growth curve (Figure 3)  
20 for the winter season in lakes and rivers. The ice growth curve was then fit to late winter  $Z_{ice}$   
21 observations for winter season primarily by adjusting  $\alpha$ . In some cases when observation came  
22 after the maximum in ATDD,  $\alpha'$  was also adjusted to provide the best fit, otherwise  $\alpha'$  was left  
23 constant (typically set at 1.0) among simulations with no effect on estimations of MIT or ITD.




1 MIT was then extracted from each record as shown in the example ice growth curve in Figure 3.  
2 All subsequent data analysis use the MIT as a standardized estimate of winter ice growth for  
3 each winter season and waterbody. Individual year estimates of MIT and the original ice  
4 thickness observations and corresponding model parameters will be archived and available at the  
5 ADC (Arp and Cherry 2020).

6 A new metric was also derived from each ice growth curve, which we term Ice Travel  
7 Duration (ITD), and is intended to represent the period of time when most modes of common  
8 travel are safe according to ice thickness and continued thickening. Quantitatively this is defined  
9 as the date when ice thickness surpasses 30 cm to the date of MIT, the latter of which typically  
10 corresponds to the maximum AFDD and the start of ice decay (Figure 3). Our rationale is that 30  
11 cm is typically the thickness when safe vehicle travel is recommended for freshwater ice and that  
12 once ice begins to decay, even though its thickness may well exceed 30 cm, its structural  
13 integrity and strength is changing rapidly such that ice thickness is less relevant to its load  
14 bearing capacity (Gold 1971, Leppäranta 2015). In practice it is common for safe foot and  
15 snowmachine travel on thicknesses less than 30 cm and similar travel is common over thick ice  
16 that is degrading. To evaluate how closely this ITD start date tracked ice conditions identified by  
17 local APRFC observers, we compared “Safe for Vehicle” dates on three rivers and one lake  
18 common to our dataset when “snowmachine” was indicated for “Type of Vehicle” in the APRFC  
19 database (<https://www.weather.gov/aprfc/freezeUp>) (Figure 4a). This comparison showed a close  
20 match with an average offset ranging from +4 days on the Yukon River at Eagle to -5 days on  
21 Lake Minchumina. Similar comparison was made with the ITD end date tracked observer data  
22 (<https://www.weather.gov/aprfc/breakupDB>) (Figure 4b). For this comparison in all but two  
23 cases, observer indications of “Safe for Vehicles” was always later than the date of maximum ice




1 thickness, as we would have anticipated because it is common for travelers to use degrading ice  
2 safely for some period before complete breakup. Thus, our estimates of ITD should be  
3 considered conservative and for many modes of travel (and corresponding levels of caution) our  
4 estimates of ITD are shorter in duration than what is practiced locally.



5

6 **Figure 3.** Example ice growth and decay curves from Barrow lakes in a thick (a) and thin (b) ice  
7 year showing curve fits to observed data, the time when maximum ice thickness (MIT) is  
8 reached, and the period representing ice travel duration (ITD).



1

2 **Figure 4.** Comparison of Ice Travel Duration metric start (a) and end (b) dates for four study  
 3 rivers and lakes to ice condition designations reported by Alaska Pacific River Forecast Center  
 4 (APRFC) observers.

5



## 1 **2.5 Data analysis**

2 Patterns and trends in MIT and ITD records for each station were primarily analyzed graphically  
3 in time-series by comparison among lake and river sites. Detection of trends was done using  
4 linear regression for the entire record available for each waterbody. To separate or break MIT  
5 records into distinct periods of potential interest, we used a combination of piece-wise linear  
6 regression (Systat 2013) to identify significant changes in trends and regime shift detection  
7 methods (Rodionov 2004) to identify significant changes in means and variance. Gaps in  
8 observational data also resulted in gaps in MIT estimates, such that many of the breaks or record  
9 separations graphically selected correspond to these missing years of record. We report  
10 significant trends ( $p < 0.05$ ) and the means and standard deviations of MIT and ITD for all periods  
11 within the record and this served as our primary basis for comparison and analysis. Multiple  
12 regression analysis was used to evaluate relationships between MIT and ITD to air temperature  
13 and upland snow depth data from proximate weather stations when available to evaluate general  
14 controls on interannual variability.

15

## 16 **3 Results**

### 17 **3.1 Patterns and trends in lake ice**

18 Latitudinal patterns in ice thickness exist in Alaska, yet location relative to mountain ranges,  
19 river valleys, and coasts, with varying degrees of sea ice influence, likely had a larger influence.  
20 Spatial patterns of MIT averaged over periods greater than two decades range from 67 cm in  
21 Southcentral (Anchorage) and 70 cm in the Interior (Fairbanks) to 167 cm on the Arctic Coastal  
22 Plain of northern Alaska (Utqiagvik) (Table 2). Intermediate average MIT ranged from 98 to 122





1 cm from shorter term records collected within the last two decades in western Alaska, along the  
 2 Alaska Range separating the Southcentral Region from the Interior, and the Brooks Range  
 3 foothills of the North Slope (Figure 1).

4

5 **Table 2.** Summary of maximum ice thickness (MIT) and ice travel duration (ITD) according to  
 6 the mean, minimum, and maximum values for each period of record reported here by nearest  
 7 community to observed lakes (blue) and rivers (green). Years corresponding to minimums and  
 8 maximums are in parentheses and metrics with significant trends are in **bold**.

<i>Metric</i>	Utiqagvik	Bettles	Eagle	Fair-banks	Nenana	Minchumina	Anchorage	Bethel
Ave MIT (cm)	<b>167</b>	109	117	70	105	<b>98</b>	67	117
Min MIT (year)	114 (2018)	69 (2009)	67 (2004)	53 (1998)	69 (2018)	62 (2014)	46 (2003)	69 (2019)
Max MIT (year)	211 (1970)	182 (2013)	157 (2015)	101 (1971)	137 (1994)	123 (1997)	84 (2011)	159 (1971)
Ave MIT Date	29-May	18-Apr	5-Apr	5-Apr	3-Apr	6-Apr	20-Mar	2-Apr
Ave ITD (days)	<b>215</b>	157	138	135	135	<b>144</b>	110	134
Min ITD (year)	192 (2013)	109 (1994)	101 (2019)	105 (1980)	98 (2019)	108 (2019)	31 (2003)	79 (2019)
Max ITD (year)	239 (1971)	191 (2013)	180 (2013)	165 (2013)	155 (2013)	170 (2013)	158 (2002)	180 (2013)

9

10 Overall the thickest ice and the steepest long-term thinning trend came from lakes on the  
 11 Barrow Peninsula (Figure 5a), set between the Chukchi and Beaufort seas (Figure 1). A  
 12 significant decrease in MIT from 1962 to 2019 of 0.9 cm/yr is most prominent in the period of  
 13 continuous record between 2003 and 2019 with 2.9 cm/yr of thinning ( $r^2=0.35$ ,  $p=0.01$ ) (Figure  
 14 5a). Analyzed separately, the earlier period with more intermittent observations showed no trend  
 15 and an average thickness of 177 cm. In comparison, MIT for four of the past six years was less  
 16 than 121 cm—a thickness not reported once during the previous 22 years with observations  
 17 dating back to 1962. MIT on Barrow Peninsula lakes was typically reached between late May



1 and early June and average number of safe travel days was 215 with a significant decline of 0.5  
2 d/yr ( $r^2=0.54$ ,  $p<0.01$ ) (Figure 6). Air temperature during the ice growth season averaged  $-17.9$   
3  $^{\circ}$  C with an increasing trend over the period of analysis and with temperature above  $-15^{\circ}$  C  
4 during the last four of six years. Upland snow depth averaged 22 cm with no trend. Winter  
5 temperature during the ice growth season explained the majority of variation in MIT for this  
6 record ( $r^2=0.61$ ,  $p<0.01$ ) and snow depth was poorly correlated.

7       The next longest and nearly complete record comes from Smith Lake in Fairbanks dating  
8 back to 1965 (Figure 1). Interestingly, no overall trend was observed over this 54-year period,  
9 though the two thickest MIT years were in 1971 and 1977. A thinning trend of 1.1 cm/yr  
10 ( $r^2=0.27$ ,  $p=0.02$ ) was noted during the first 19 years, while the middle period (1986-2003) had  
11 an average MIT of 72 cm and the last period had average MIT of 66 cm with no trend (Figure  
12 5b). Very thin MIT, less than 60 cm, occurred in all three periods, and MIT exceeding 80 cm  
13 occurred as recently as 2017. Smith Lake is a shallow thermokarst lake with wind-protecting  
14 forest around its entire perimeter, typically with relatively deep uniform snowcover.  
15 Contemporary synoptic comparisons made by APRFC on several larger and less protected  
16 interior lakes within a 100 km radius often show Smith Lake having the thinner ice regionally.  
17 Still, Smith Lake may be widely representative of the many small thermokarst lakes and pond  
18 surrounded by forest in interior Alaska. MIT was typically reached by early April and average  
19 ITD was 135 days with no trend over time (Table 2). Air temperature during the ice growth  
20 season averaged  $-16.8^{\circ}$  C with a slight increasing trend and upland snow depth averaged 36 cm  
21 with no trend. Neither upland snow depth nor air temperature during the ice growth season  
22 explained interannual variation in MIT or ITD for this lake.




1           In contrast, the larger and more southerly, but slightly higher elevation Lake Minchumina  
2 (Figure 1) had a MIT of 98 cm averaged over 18 years of mostly continuous observation. MIT of  
3 this lake became notably thinner over this period, decreasing 1.6 cm/yr ( $r^2=0.28$ ,  $p=0.02$ ) (Figure  
4 5c). Yet, nearly the thickest MIT in this record, 121 cm, was recent, 2013, and then the thinnest  
5 MIT of 62 cm was also recent, 2014 (Table 2). MIT was typically reached by early April and  
6 average ITD was 144 days with a relatively steep declining trend of 1.5 d/yr ( $r^2=0.33$ ,  $p<0.02$ )  
7 over this short period (Figure 6). We estimated that the most recent year of record 2019 had the  
8 shortest ITD of 108 days, while another relatively recent year 2013 had the longest ITD of 170  
9 days (Table 2). The long and relatively cold ice growth season of 2013 was also prominent in  
10 other records of ITD in interior sites. Air temperature during the ice-growth season averaged -  
11 14.8 °C (between Nenana and McGrath stations) with no trend detected.

12           The most southerly site, Lake Hood, located near the Anchorage International Airport  
13 and used as a major floatplane runway, had a similar duration record as Lake Minchumina from  
14 1997 to 2019. Observations on this lake began in 1967, but these appeared anomalously thick,  
15 with many exceeding 140 cm and  $\alpha$  coefficients averaging 4.8. We suspect that observers may  
16 have collected measurements along areas where ski-planes or snow grooming compacted the  
17 snow significantly to allow this level of thickening with modest winter temperatures. For the  
18 period when  $\alpha$  coefficients are within a more normal range (1.6-3.7) (Table 1), a non-significant  
19 increase of 0.3 cm/yr was detected with an average MIT of 67 cm (Figure 4d). MIT was typically  
20 reached by late March and average ITD was 110 days with lots of variation, but no trend over  
21 this period (Figure 5). We estimated a very short ITD of 31 days over the winter of 2002-03  
22 (Table 2) when ice growth only surpassed 30 cm by 31-Dec and MIT was reached early, 2-Feb.  
23 That winter, as well as 2007-08, had numerous freeze-thaw periods through the normal ice




- 1 growth season corresponding to short estimates of ITD (Figure 6). Ice-growth season
- 2 temperature averaged  $-6.4^{\circ}\text{C}$  and upland snow depth averaged 29 cm, with no trends detected.
- 3 We suspect snow on this lake would be greatly reduced in some years due to wind-scour along
- 4 with varying alteration from human activities.



1

2 **Figure 5.** Maximum ice thickness patterns and trends for lakes around the Barrow Peninsula (a),  
3 Fairbanks (b), Minchumina (c), and Anchorage (d) for each station's period of record.



1

2 **Figure 6.** Ice travel duration on lakes estimated from ice growth curves for each period of  
3 record.

4

### 5 3.2 Patterns and trends in river ice

6 In contrast to lake ice records, fewer river ice thickness records were analyzed in this study. All  
7 of these observation sites are on larger rivers within the Yukon and Kuskokwim river drainages  
8 (Figure 1), associated with river-side communities. A primary use of these ice thickness data is  
9 in forecasting ice jam floods and safe travel conditions. Average MIT over these differing  
10 periods of observation ranged from 98 cm on the Tanana River at Nenana (40 years) in the  
11 Interior to 117 cm at both the Kuskokwim River at Bethel (57 years) near the Bering Sea and  
12 Yukon River at Eagle (20 years) near the Canadian border (Figure 1).



1 Ice observations on the Kuskokwim River began with the CRREL program in 1962 with  
2 a decade-long gap starting in 1985 and continued again in 1996 with measurements reported to  
3 APRFC. The earlier period (1962-1984) had average MIT of 124 cm with the thickest ice of 159  
4 cm recorded in 1997, no estimates of MIT less than 100 cm, and no trend (Figure 7a). The  
5 second period running to the present also showed no trend, but had somewhat lower average  
6 MIT of 111 cm and higher interannual variation than the first period. The thinnest MIT, 69 cm,  
7 occurred in 2019 and also corresponded to the earliest river breakup on record, 31-March. Also  
8 in contrast to the first period, MIT was less than 100 cm during eight of 24 winters in the recent  
9 period (Figure 7a). However, thick ice (>130 cm) also occurred in 2009, 2010, 2012, and 2017,  
10 underscoring a recent pattern of higher interannual variability. The average date of peak ice  
11 thickness was 1-April, but was estimated to occur as early as February in some years and as late  
12 as May in others with no trend. ITD also varied widely from 79 days in 2019 to 190 days in 2013  
13 and averaged 144 days, also with no significant trend (Figure 8). The Kuskokwim River ice  
14 growth curve during the winter of 2018-19 suggests 30 cm ice-thickness was reached by mid-  
15 December and MIT was reached in late February. In contrast, during the winter of 2012-13, 30  
16 cm ice thickness was reached in early November and peaked in early May. During the ice growth  
17 period for the full record, air temperature at Bethel averaged  $-11.8^{\circ}\text{C}$  and upland snow depth  
18 averaged 14 cm. Neither ice-growth controlling condition showed any trend during this period,  
19 yet both air temperature and upland snow depth together explained significant portions of  
20 variability in MIT ( $r^2=0.24$ ,  $p<0.03$ ).

21 Ice thickness records on the Tanana River were collected by community members  
22 associated with the Nenana Ice Classic (NIC) and were available back to the winter of 1988-89.  
23 In that year, MIT was 112 cm and the average for the entire record is 105 cm (Table 2). The most



1 recent two years had the lowest MIT on record, 69 and 72 cm, respectively—the latter, 2019, of  
2 which was also the earliest breakup in the 102 year record, tipping the tripod on 14-April. Two  
3 distinct ice thickness periods are noted. The earlier period from 1989 to 2007 had average MIT  
4 of 106 cm with high interannual variability and no trend (Figure 7b). The second period showed  
5 a strong thinning trend of 4.5 cm/yr ( $r^2=0.80$ ,  $p<0.01$ ) ending in the two thinnest ice years, as  
6 previously noted. Ice typically thickened until early April with recorded breakup happening one  
7 month later on average. Ice travel duration on this section of the Tanana River averaged 135 days  
8 and ranged from 108 days in 2019 to 165 days in 2013 (Table 2) —which also corresponded to  
9 the earliest and latest breakup dates for the total 102 year record. Ice growth season air  
10 temperature averaged  $-16.4^{\circ}\text{C}$  over this period and upland snow data was not consistently  
11 available at this station over this period.

12 The second longest and mostly complete river ice observation record was made on the  
13 Koyukuk River at Bettles starting in 1968. Three periods were noted in this record of  
14 approximately equal durations (Figure 7c). The first was characterized by increasing thickness of  
15 MIT by 3 cm/yr ( $r^2=0.33$ ,  $p=0.02$ ) with ice as thin as 81 cm in 1968 and as thick as 182 cm in  
16 1978. The middle and the latest period we identified were less distinct in terms of average MIT,  
17 111 and 103 cm, respectively, and lacked trends, but the latter period had much higher  
18 interannual variability with MIT ranging from 69 cm in 2009 to 182 cm in 2013 (Figure 7c). Ice  
19 typically reached its maximum by mid- to late-April and ITD averaged 157 days with less  
20 interannual variability than other rivers in this set (Figure 8). Ice-growth season air temperature  
21 averaged  $-18.3^{\circ}\text{C}$  and upland snow was quite deep, averaging 54 cm and ranging from 22 cm in  
22 1996 to 100 cm at the start of the record in 1968. Here, upland snow depth explained a






1 significant portion of the variation in MIT ( $r^2=0.16$ ,  $p<0.05$ ), while air temperature was  
2 uncorrelated to ice variability over this period.

3 The Yukon River at Eagle showed an increasing trend in MIT of 1.5 cm/yr ( $r^2=0.17$ ,  
4  $p=0.06$ ) from 1999 to 2019, though not quite significant statistically (Figure 7d). Three distinctly  
5 thin ice years occurred in 1999, 2004, and 2019 when the MIT reached 82 cm or less. Recent  
6 thick ice years in 2015 and 2017, when the MIT was greater than 150 cm, contributed to this  
7 weak trend of increasing ice growth (Figure 7d) at this eastern most station near the Canadian  
8 border. MIT was typically reached in early April in most years and ITD ranged from 101 days in  
9 2019 to 180 days in 2013 (Table 2). Air temperature during the ice growth period averaged -  
10  $16.5^{\circ}\text{C}$  and was not correlated with variation in MIT over this period. Upland snow  
11 observations were not consistently recorded at the station in Eagle.


12

13



1

2 **Figure 7.** Maximum ice thickness patterns and trends for rivers near Bethel (a), Nenana (b),  
3 Bettles (c), and Eagle (d) for each station's period of record.



1

2 **Figure 8.** Ice travel duration on rivers estimated from ice growth curves for each period of  
3 record.

4

#### 5 **4 Discussion**

6 The clearest signal of reduced ice growth and shorter duration ice cover come from lakes in  
7 northern Alaska where the impacts of arctic amplification are known to be most pronounced  
8 (Wendler et al. 2014, Walsh and Brettschneider 2019). The majority of this trend from Barrow  
9 Peninsula lakes is driven by recent declining sea ice impacts on early winter temperature and  
10 snowfall (Alexeev et al. 2016), though strong interannual variability in ice thickness is still  
11 evident (Arp et al. 2018). Barrow Peninsula ice observations from 1962 to 1996 are well within  
12 the range of more distant single year observations of 192 cm in 1955 (Brewer 1958) and 188 cm



1 in 1882 (Ray 1885). In comparison, MIT for four of the past six years was less than 121 cm—a  
2 thickness not reported once during the previous 22 years with observations dating back to 1962.  
3 Snow is typically considered the dominant control on interannual variability in ice thickness for  
4 coastal plain lakes (Zhang and Jeffries 2000), yet warmer winter temperature during the ice  
5 growth season appear to have overcome this driver of ice growth in our analysis. Snow depth on  
6 coastal plain lakes is typically about 60% of upland depth on average (Sturm and Liston 2003),  
7 but this can vary greatly from year to year (Arp et al. 2018) and this changing offset between  
8 tundra snow and lake snowpack may explain the lack of correlation to upland snow records we  
9 observed. The role of arctic amplification and early winter warming is also seen in reduced safe  
10 travel days on ice for Barrow Peninsula lakes (Figure 6a) with the majority of this decrease  
11 coming from slower ice thickening rather than earlier arrival of MIT.

12 Despite the impacts of arctic amplification on winter climate change in other parts of  
13 Alaska (Walsh and Brettschneider 2019), the response of river and lake ice growth in several  
14 records spanning over 50 years often appear muted or highly variable. In western coastal Alaska,  
15 recent thin ice conditions and short ice cover duration on the Kuskokwim River were striking,  
16 yet follow a pattern of enhanced variability over recent decades. Thin ice conditions of the 2018-  
17 19 winter were observed in nearly all records we analyzed and provided much of our motivation  
18 to standardize, summarize, and analyze these records. In contrast, the relatively recent winter of  
19 2012-13 had consistently thick ice and very prolonged ice cover duration across western and  
20 interior Alaska. Such dramatic winter conditions and divergent ice responses underscore the need  
21 for enhanced freshwater ice observation programs.

22 The premise that freshwater ice growth integrates changes in climate deserves  
23 consideration (Allen 1977, Engram et al. 2018). We found few consistent relationships between



1 fundamental drivers of ice growth, air temperature and snow depth, suggesting that other more  
2 complex environmental factors play a role in river and lake ice dynamics. One factor is that snow  
3 accumulation on ice is fundamentally different than terrestrial upland snowpacks where snow  
4 depth is recorded—typically ice on rivers and lakes is thinner and depending on the ice column’s  
5 isostatic balance can slow ice growth through insulation or thicken it through formation of snow-  
6 ice and overflow (Sturm and Liston 2003, Ashton 2011). Particularly on rivers, the combined  
7 hydrologic and thermal conditions of flowing waters can also cause divergent responses in ice  
8 thickness—more and warmer water can cause slower growth or degradation, but also generate  
9 overflow that can refreeze and add thickness to ice covers (Prowse and Beltaos 2002, Brown and  
10 Duguay 2010, Jones et al. 2015). Little is known about changes in groundwater in Alaska, which  
11 also impacts ice growth on rivers, though several studies do point to increases in groundwater  
12 input (Brabets and Walvoord 2009, Liljedahl et al. 2017). Higher water temperatures in relation  
13 to enhanced groundwater input present another potentially important driver affecting ice growth  
14 and decay that deserves evaluation (Jones et al. 2015, Cherry 2019). Many of these interactions  
15 are documented process studies of lake and river ice (Ashton 2011) and observations of these  
16 processes also appear sporadically in monitoring program notes (Bilello 1980), but are  
17 challenging to quantify in long time-series analysis such as this one. Thus, ice thickness at its  
18 seasonal maximum (MIT) and duration of the ice growth season (ITD) do integrate important  
19 and complex changes in climate including the hydrologic cycle, but these responses do need to  
20 be carefully interpreted and compared along with other environmental drivers.

21 This analysis of long-term ice observation records in Alaska using standardized metrics  
22 from ice growth curves provides an important baseline to compare with future observations and  
23 support process studies. Past freshwater ice observations programs in Alaska, including CRREL



1 (Bilello 1980) and ALISON (Morris and Jeffries 2010), both collected basic ice thickness data  
2 that supported numerous process studies adding to our ice dynamics predictive capability (e.g.,  
3 Jeffries et al. 2005, Arp et al. 2010, Ashton 2011). These datasets are now archived by the Arctic  
4 Data Center (arcticdata.io/) as Bilello (2019) and Morris and Jeffries (2019) and many of these  
5 records continued and made readily available by APRFC  
6 (<https://www.weather.gov/aprfc/IceThickness>). A new freshwater ice observation program, Fresh  
7 Eyes on Ice (freshiceonice.org), is working to continue monitoring and analysis of river and lake  
8 ice conditions in Alaska in part through engagement with rural communities and public schools  
9 using a combination of approaches including remote sensing and field-based observations.

10 Our analysis of ice growth curves only represents a portion of the ice formation, growth,  
11 and decay process, whereas more abundant ice phenology studies (e.g. Arp et al. 2013, Cooley  
12 and Pavelsky 2016, Smejkalova et al. 2017, Sharma et al. 2019, Yang et al. 2020) identify  
13 patterns and trends at the very start and very end of this cycle of seasonal ice cover. Perhaps the  
14 two most relevant and also challenging periods are freeze-up and break-up, which in terms of  
15 process and informing travel conditions should be viewed as a continuum rather than momentary  
16 *to-the-day* events (Brown et al. 2018). The critical freeze-up period occurs from when ice first  
17 forms and grows on water surfaces to when ice cover is spatially consistent and thick enough to  
18 support reliably safe travel. The critical break-up period, starts after reaching MIT when ice  
19 decay begins and then proceeds at widely varying rates in space and time for lakes and even  
20 more so for rivers. The importance of understanding how these periods of freeze-up to  
21 contiguous thick ice and decay initiation to break-up progress, and how this progression may be  
22 changing over time, are critical in terms of informing safe winter travel, predicting ice jam flood  
23 hazards, and understanding interactions with river and lake ecosystems (Brown et al. 2018).



1 Focusing on these key ice growth and decay periods and how they may be responding in new  
2 ways to climate change and arctic amplification deserve renewed attention in northern regions.

3

4 *Acknowledgements:* Numerous professional and citizen scientists contributed to ice thickness  
5 observations used in this analysis including many dedicated volunteers from CRREL, ALISON,  
6 and APRFC programs. Augering holes in lake and river ice is cold tedious work and their efforts  
7 and additional observations over many years is greatly appreciated. Known ice observers from  
8 Alaska's North Slope contributed hundreds of observations over recent decades including Ben  
9 Jones, Guido Grosse, Matthew Whitman, Richard Beck, Ken Hinkel, Ben Gaglioti, Andrew  
10 Parsekian, Andrea Creighton, and many others. This research was primarily supported by a grant  
11 from the National Science Foundation (#1836523).

12

### 13 **References**

- 14 Alexeev, V. A., Arp, C. D., Jones, B. M., and Cai, L.: Arctic sea ice decline contributes to  
15 thinning lake ice trend in northern Alaska, *Environ Res Lett*, 11, 2016.
- 16
- 17 Allen, W. T. R.: Freeze-up, break-up, and ice thickness in Canada, Fisheries and Environment  
18 Canada, Report CLI-1-77, 185p., 1977.
- 19
- 20 Arp, C.: Lake ice thickness observations for arctic Alaska from 1962 to  
21 2017. urn:node:ARCTIC. doi:10.18739/A2G27V, 2018a.
- 22
- 23 Arp, C.: Arctic Alaska Tundra and Lake Snow Surveys from 2012-2018. Arctic Data  
24 Center. doi:10.18739/A2G15TB05, 2018b.
- 25
- 26 Arp, C. and Cherry, J.: Seasonal maximum ice thickness data for rivers and lakes in Alaska from  
27 1962 to 2019. Arctic Data Center. urn:uuid:64148cbe-1c32-433a-8b49-a88b29bc752a, 2020.
- 28
- 29 Arp, C. D. and Jones, B. M.: Geography of Alaska Lake Districts: Identification, description, and  
30 analysis of lake-rich regions of a diverse and dynamic state, U.S. Geological Survey, 40 pp.,  
31 2009.
- 32
- 33 Arp, C. D., Jones, B. M., and Grosse, G.: Recent lake ice-out phenology within and among lake  
34 districts of Alaska, U.S.A., *Limnology and Oceanography*, 58, 2013-2028, 2013.
- 35
- 36 Arp, C. D., Jones, B. M., Engram, M., Alexeev, V. A., Cai, L., Parsekian, A., Hinkel, K.,  
37 Bondurant, A. C., and Creighton, A.: Contrasting lake ice responses to winter climate indicate  
38 future variability and trends on the Alaskan Arctic Coastal Plain, *Environ Res Lett*, 13, 125001,  
39 2018.
- 40



- 1 Ashton, G. D.: Thin ice growth, *Water Resources Research*, 25(3): 564-566, 1989.  
2  
3 Ashton, G. D.: River and lake ice thickening, thinning, and snow ice formation, *Cold Regions  
4 Research and Technology*, 68, 3-19, 2011.  
5  
6 Bilello, M. A.: Method for predicting river and lake ice formation, *Journal of Applied  
7 Meteorology*, 3: 38-44, 1964.  
8  
9 Bilello, M. A.: Maximum thickness and subsequent decay of lake, river, and fast sea ice in  
10 Canada and Alaska, U.S. Army, 160 pp., 1980.  
11  
12 Bilello, M.: River and lake ice thickness and snow depth at near maximum ice thickness and  
13 during ice decay in Alaska, 1961-1974. Arctic Data Center. doi:10.18739/A2FF3M027, 2019.  
14  
15 Brabets T. P. and M. A. Walvoord: Trends in streamflow in the Yukon River Basin from 1944 to  
16 2005 and the influence of the Pacific Decadal Oscillation. *Journal of Hydrology* 371:108-119  
17 doi:10.1016/j.jhydrol.2009.03.018, 2009.  
18  
19 Brewer, M. C.: The thermal regime of an arctic lake, *Transactions of the American Geophysical  
20 Union*, 39, 278-284, 1958.  
21  
22 Brown, D. R. N., Brinkman, T. J., Verbyla, D. L., Brown, C. L., Cold, H. S., and Hollingsworth,  
23 T. N.: Changing River Ice Seasonality and Impacts on Interior Alaskan Communities, *Weather,  
24 Climate, and Society*, 10, 625-640, 2018.  
25  
26 Brown, L. C. and Duguay, C. R.: The response and role of ice cover in lake-climate interactions,  
27 *Progress in Physical Geography*, 34, 671-704, 2010.  
28  
29 Cherry, J.E. Alaska Climate Dispatch, Summer 2019. Another Season of Dangerous Ice  
30 Conditions. [https://uaf-accap.org/wp-content/uploads/2019/08/climate-dispatch\\_2019.pdf](https://uaf-accap.org/wp-content/uploads/2019/08/climate-dispatch_2019.pdf)  
31  
32 Cold, H. S., T. J. Brinkman, C. L. Brown, T. N. Hollingsworth, D. R. N. Brown, and K. M.  
33 Heeringa. 2020. Assessing vulnerability of subsistence travel to effects of environmental change  
34 in Interior Alaska. *Ecology and Society* 25(1):20.  
35 <https://doi.org/10.5751/ES-11426-250120>  
36  
37 Cooley, S. W. and Pavelsky, T. M.: Spatial and temporal patterns in Arctic river ice breakup  
38 revealed by automated ice detection from MODIS imagery, *Remote Sensing of Environment*,  
39 175, 310-322, 2016.  
40  
41 Gold, L. W.: Use of Ice Covers for Transportation, *Canadian Geotechnical Journal*, 8(2): 170-  
42 181, <https://doi.org/10.1139/t71-018>, 1971.  
43  
44 Engram, M., Arp, C. D., Jones, B. M., Ajadi, O. A., and Meyer, F. J.: Analyzing floating and  
45 bedfast lake ice regimes across Arctic Alaska using 25 years of space-borne SAR imagery,  
46 *Remote Sensing of Environment*, 209, 660-676, 2018.





- 1 Fleischer, N. L., Melstrom, P., Yard, E., Brubaker, M., and Thomas, T.: The epidemiology of  
2 falling-through-the-ice in Alaska, 1990–2010, *Journal of Public Health*, 36, 235-242, 2014.
- 3
- 4 Gould, M. and Jeffries, M.: Temperature variation in lake ice in central Alaska, USA, *Annals of*  
5 *Glaciology*, 40, 1-6, 2005.
- 6
- 7 Jeffries, M. O., Morris, K., and Duguay, C. R.: Lake ice growth and decay in central Alaska,  
8 USA: observations and computer simulations compared, *Annals of Glaciology*, 40, 1-5, 2005.
- 9
- 10 Jones, C., Kielland, K., and Hinzman, L.: Modeling groundwater upwelling as a control on rivder  
11 ie thickness, *Hydrology Research*, 46.4, 2015.
- 12
- 13 Jumikis A. R.: *Thermal Geotechnics*, Rutgers University Press: New Brunswick, NJ; 375 pp.,  
14 1977.
- 15
- 16 Hinkel, K. M., Lenters, J. D., Sheng, Y. W., Lyons, E. A., Beck, R. A., Eisner, W. R., Maurer, E.  
17 F., Wang, J. D., and Potter, B. L.: Thermokarst Lakes on the Arctic Coastal Plain of Alaska:  
18 Spatial and Temporal Variability in Summer Water Temperature, *Permafrost and Periglacial*  
19 *Processes*, 23, 207-217, 2012.
- 20
- 21 Leppäranta, M.: A growth model for black ice, snow ice and snow thickness in subarctic basins,  
22 *Nordic Hydrology*, 14, 59-70, 1983.
- 23
- 24 Leppäranta, M.: *Freezing of Lakes and the Evolution of their Ice Cover*, Springer Publishing,  
25 301pp., 2015.
- 26
- 27 Liljedahl, A. K., A. Gädeke, S. O'Neel, T. A. Gatesman, T. A. Douglas: Glacierized headwater  
28 streams as aquifer recharge corridors, subarctic Alaska. *Geophysical Research Letters* 44:6876-  
29 6885 doi:10.1002/2017gl073834, 2017.
- 30
- 31
- 32 Magnuson, J. J., Robertson, D. M., Benson, B. J., Wynne, R. H., Livingstone, D. M., Arai, T.,  
33 Assel, R. A., Barry, R. G., Card, V., Kuusisto, E., Granin, N. G., Prowse, T. D., Stewart, K. M.,  
34 and Vuglinski, V. S.: Historical trends in lake and river ice cover in the Northern Hemisphere,  
35 *Science*, 289, 1743-1747, 2000.
- 36
- 37 Morris, K. and Jeffries, M.: Alaska Lake Ice and Snow Observatory Network (ALISON). In:  
38 *Polar Science and Global Climate: An International Resource for Education and Outreach*,  
39 Kaiser, B., Allen, B., and Zicus, S. (Eds.), Pearson Education Limited, Harlow, Essex, UK,  
40 2010.
- 41
- 42 Morris, K. and Jeffries, M.: Alaska Lake Ice and Snow Observatory Network (ALISON) Project  
43 Data, Alaska, 1999-2011. Arctic Data Center. doi:10.18739/A2K35MD3N, 2019.
- 44
- 45 Prowse, T. D. and Beltaos, S.: Climatic control of river-ice hydrology: a review, *Hydrological*  
46 *Processes*, 16, 805-822, 2002.



- 1  
2 Ray, P.H.: Report of the International Polar Expedition to Point Barrow, Alaska, Washington,  
3 D.C., Government Printing Office, 1885.  
4  
5 Rodionov, S. N.: A sequential algorithm for testing climate regime shifts, *Geophysical Research*  
6 *Letters*, 31, 2004.  
7  
8 Sagarin, R. and Micheli, F.: Climate Change in Nontraditional Data Sets, *Science*, 294, 811-811,  
9 2001.  
10  
11 Schneider, W.S., Brewster, K., Kielland, K., and Jones, C.E.: On Dangerous Ice: Changing  
12 Conditions on the Tanana River. University of Alaska Fairbanks, Fairbanks, Alaska. 76 pp.,  
13 2013.  
14  
15 Serreze, M. C. and Barry, R. G.: Processes and impacts of Arctic amplification: A research  
16 synthesis, *Global and Planetary Change*, 77, 85-96, 2011.  
17  
18 Serreze, M. C. and Francis, J. A.: The arctic amplification debate, *Climatic Change*, 76, 241-264,  
19 2006.  
20  
21 Sharma, S., Blagrove, K., Magnuson, J. J., O'Reilly, C. M., Oliver, S., Batt, R. D., Magee, M. R.,  
22 Straile, D., Weyhenmeyer, G. A., Winslow, L., and Woolway, R. I.: Widespread loss of lake ice  
23 around the Northern Hemisphere in a warming world, *Nature Climate Change*, 9, 227-231, 2019.  
24  
25 Smejkalova, T., Edwards, M. E., and Dash, J.: Arctic lakes show strong decadal trend in earlier  
26 spring ice-out, *Scientific Reports*, doi: 10.1038/srep38449, 2017. 1-8, 2017.  
27 Sturm, M. and Liston, G. E.: The snow cover on lakes of the Arctic Coastal Plain of Alaska,  
28 U.S.A., *Journal of Glaciology*, 49, 370-380, 2003.  
29  
30 Stefan J.: Über die Theorie der Eisbildung, insbesondere über die Eisbildung im Polarmee.  
31 *Annals of Physics and Chemistry*, 42: 269–286, 1891.  
32  
33 Walsh, J. E. and Brettschneider, B.: Attribution of recent warming in Alaska, *Polar Sci*, 21, 101-  
34 109, 2019.  
35  
36 Weeks, W. F., Fountain, A. G., Bryan, M. L., and Elachi, C.: Differences in radar return from  
37 ice-covered North Slope lakes, *Journal of Geophysical Research*, 83, 4069-4073, 1978.  
38  
39 Wendler, G., Moore, B., and Galloway, K.: Strong temperature increase and shrinking sea ice in  
40 Arctic Alaska, *The Open Atmospheric Science Journal*, 8, 7-15, 2014.  
41  
42 Weyhenmeyer, G. A., Livingstone, D. M., Meili, M., Jensen, O., Benson, B., and Magnuson, J.  
43 J.: Large geographical differences in the sensitivity of ice-covered lakes and rivers in the  
44 Northern Hemisphere to temperature changes, *Global Change Biology*, 17, 268-275, 2011.  
45



- 1 Yang, X., Pavelsky, T. M., and Allen, G. H.: The past and future of global river ice, *Nature*, 577,
- 2 69-73, 2020.
- 3
- 4 Zhang, T. and Jeffries, M. O.: Modeling interdecadal variations of lake-ice thickness and
- 5 sensitivity to climatic change in northernmost Alaska, *Annals of Glaciology*, 31, 339-347, 2000.
- 6