

Interactive comment on “Systematic bias of Tibetan Plateau snow cover in subseasonal-to-seasonal models” by Shuzhen Hu and Wenkai Li

Anonymous Referee #2

Received and published: 21 July 2020

This paper examines how a set of state-of-the-art subseasonal-to-seasonal (S2S) forecast systems predict the Tibetan Plateau-wide snow cover and surface temperature over the 1999-2010 period. The forecast systems from ECMWF, CMA and NCEP are used in the intercomparison. The evaluation of forecasted snow cover is made against the multi-instrument (IMS) snow cover data. A connection is drawn between the bias in snow cover, which increases systematically with lead time in all models, and a bias in surface temperature. Model experiments with the WRF model support the idea that the latter is induced by the snow excess through land-atmosphere coupling. Few papers have examined snow forecast on the S2S time scale on the Tibetan Plateau (TP), a region with well-known biases in snow and surface temperature. This is an innovative

C1

study that is well-worthy of publication in The Cryosphere. I however recommend a major revision of the paper, before it is in an acceptable form for publication.

MAJOR COMMENTS 1) Some brief description of how the three different models are initialised in terms of snow and land surface is needed, complementing the description of the land-surface models. What sort of snow analysis is used to initialise the different models? Isn't ERA-5, which has strong snow biases over the TP, used to initialise the ECMWF S2S reforecasts? Which observations are used in these snow analyses, both globally and over the TP more specifically? While IMS is not used to initialise the ECMWF forecast model over the TP (see Orsolini Y. et al., 2019), is it used in the other systems? The quality of this initialisation would most certainly influence the snow forecasts at the S2S time scale. The initial snow values may be a key addition to bring in Fig 3. 2) Some more details about how the snow cover fraction conversion is derived in each model is needed. For example, a 100% snow cover may mean very different snow depth or snow water equivalent (the actual prognostic variables) in the different prediction systems. Also, I believe that IMS provides a binary information snow cover being 0 or 1, with the former case meaning that the fractional snow cover is below 50%. When aggregated to a 1-degree grid (L88), isn't there a range or uncertainty in the IMS aggregated value (given that the observed fractional snow cover could actually be 0 or 50%)? Please clarify these points and the implications for forecast verification. 3) The authors consider the snow cover averaged over the entire Tibetan-Plateau, but they do not show any snow cover maps although there is considerable heterogeneity, as shown by the authors in previous publications. I wonder if there could be compensation effects between different geographical sub-regions, that could result in an agreement of the TPSC index between forecasts and IMS. Different prediction systems may have different regional biases. Showing such maps would re-assure the reader that the prediction systems capture the main climatological features of the snow distribution over the TP and its S2S variability (e.g., in subregions as in Li W. et al., IJOC, 2019). Could spatial pattern correlation between IMS and snow forecasts be helpful in this case? 4) The authors could also try to examine the snow-temperature coupling strength (as an

C2

indicator of land-atmosphere interaction) in the respective prediction systems by correlating the local forecasted temperature and forecasted snow, as a function of lead time. See Diro and Lin (2020) or Li et al. (2019).

Diro, G. T., and H. Lin, 2020: Subseasonal Forecast Skill of Snow Water Equivalent and Its Link with Temperature in Selected SubX Models. *Wea. Forecasting*, 35, 273–284. Orsolini, Y., et al. G. (2019), Evaluation of snow depth and snow cover over the Tibetan Plateau in global reanalyses using in situ and satellite remote sensing observations. *The Cryosphere*, Vol. 13.(8) s.2221-2239 Li, F., et al. (2019). Impact of snow initialization in subseasonal to seasonal winter forecasts with the Norwegian Climate Prediction Model. *Journal of Geophysical Research: Atmospheres*, 124. <https://doi.org/10.1029/2019JD030903>

MINOR COMMENTS L92-93: the description here focuses on winter. While winter is the main focus, many plots show year-long results. It would be better to emphasize the whole set used (nb of years, total nb of forecasts, . . .), not only the winters. L252: the snow bias might not only arise from the land surface model (shared between WRF and the NCEP model) but also from the meteorological forcing (e.g., excess precipitation). Please clarify. L244: how is the GDAS snow analysis used in Section 4.3 on numerical modelling compare with the IMS snow analysis, used in the first part of the paper. While it is mentioned that GDAS assimilates IMS, does it assimilate it over the TP? What does it assimilate specifically over the TP (in-situ data)? Would the prediction skill be different if evaluated against GDAS (Fig 1) ? Conclusions: a brief mention of possible, relevant physical processes over the TP leading to snow ablation would be helpful. Could it be the strong surface winds or else the snow sublimation missing in the models? The short length of the period over which the forecasts are evaluated (around 10 years) is a bit of a concern. It appears that the biases are quite strong and systematic, but I wonder if some features in the forecasts would be robust over a longer period: for example, the slowdown in TPSC in early winter (December) seen in ECMWF and NCEP (Fig 3). I realise that if adding another 10 years may entail a lot of

C3

computational work, but it would add to the robustness of the conclusions. At least, a word of caution in the summary is warranted.

Typos / English L43: hydrologic cycle L28: radiative rather than radiant L108: the total variability L113: and the three different L143: the preceding week rather than the last week, seems more appropriate L152: accumulation, leading to a systematic TPSC bias. L168: growth is used for a declining variable. Either decrement, reduction or decline should be clearer. L173: real rate should be observed rate or rate derived from IMS. L173: indices or index L202: growth of SAT, rather than TPSC (it says 1.2 degree). L229: land-atmosphere interactions L256: Hence,

Interactive comment on *The Cryosphere Discuss.*, <https://doi.org/10.5194/tc-2020-131>, 2020.

C4