

Heterogenous CO₂ and CH₄ content of glacial meltwater from the Greenland Ice Sheet and implications for subglacial carbon processes

Andrea J. Pain^{1,2}, Jonathan B. Martin¹, Ellen E. Martin¹, Åsa K. Rennermalm³, Shaily Rahman^{1,4}

¹ University of Florida, Department of Geological Sciences, Gainesville, FL 32611

² Now at: University of Maryland Center for Environmental Science, Horn Point Lab, Cambridge, MD 21613

³ Rutgers, The State University of New Jersey, Department of Geography, Piscataway NJ 08854

⁴ Now at: University of Southern Mississippi, Department of Marine Science, Stennis Space Center, MS 39529

Correspondence to: Andrea J. Pain (ajpain@umces.edu)

10 **Abstract.** Accelerated melting of the Greenland Ice Sheet has increased freshwater delivery to the Arctic Ocean and amplified the need to understand the impact of Greenland Ice Sheet meltwater on Arctic greenhouse gas budgets. We measured carbon dioxide (CO₂) and methane (CH₄) concentrations and δ¹³C values and use geochemical models to evaluate subglacial CH₄ and CO₂ sources and sinks in water discharging from three subglacial outlets of the Greenland Ice Sheet in southwest (a sub-catchment of the Isunnguata Glacier, sub-Isunnguata, and the Russell Glacier) and southern Greenland (Kiattut Sermiat).

15 Meltwater CH₄ concentrations vary by orders of magnitude between sites and are saturated with respect to atmospheric concentrations at Kiattut Sermiat. In contrast, meltwater in southwest sites is supersaturated, even though oxidation reduces CH₄ concentrations by up to 50% during periods of low discharge. CO₂ concentrations range from supersaturated at sub-Isunnguata to undersaturated at Kiattut Sermiat. CO₂ is consumed by mineral weathering throughout the melt season at all sites, however differences in the magnitude of subglacial CO₂ sources result in meltwaters that are either sources or sinks of atmospheric CO₂. At the sub-Isunnguata site, the predominant source of CO₂ is organic matter (OM) remineralization. At Russell and Kiattut Sermiat sites, multiple or heterogeneous subglacial CO₂ sources that maintain atmospheric CO₂ concentrations at Russell but not at Kiattut Sermiat where CO₂ is undersaturated. These results highlight a previously unrecognized degree of heterogeneity in greenhouse gas dynamics under the Greenland Ice Sheet. Future work should constrain the extent and controls of heterogeneity to improve our understanding of the impact of Greenland Ice Sheet melt on

20 Arctic greenhouse gas budgets, as well as the role of continental ice sheets in greenhouse gas variations over glacial-interglacial timescales.

1 Introduction

Glaciers play an important role in global chemical cycles due to the production of fine-grained sediments that participate in carbonate and silicate mineral weathering reactions (Table 1), which are the principal sink of atmospheric CO₂ over geologic timescales (Berner et al., 1983; Walker et al., 1981). Variations in the weathering intensity of comminuted

Deleted: of

Deleted: ³

Deleted: ³

Deleted: ajpain@ufl

Deleted: (GrIS)

Deleted: GrIS

Deleted: (GHG)

Deleted: GrIS

Deleted: Glaciers

Deleted: , but are supersaturated at

Deleted: The

Deleted: at Isunnguata

Deleted: , but

Deleted: have

Deleted: the variability

Deleted: GHG

Deleted: GrIS. Constraining this variability will

Deleted: GrIS

Deleted: GHG

Deleted: GHG

sediments may contribute to glacial-interglacial atmospheric CO₂ variations as sediments are alternately covered by ice and exposed following ice retreat. However, the importance of CO₂ consumption by mineral weathering is poorly understood, including effects from the advance and retreat of continental ice sheets (Ludwig et al., 1999). Recent evaluations of carbon budgets in proglacial environments indicate that mineral weathering results in net sequestration of atmospheric CO₂, suggesting that proglacial systems are underrecognized as Arctic CO₂ sinks (St Pierre et al., 2019), however alternate processes could lead to the production of greenhouse gases in glacial systems. For instance, CH₄ production in anaerobic subglacial environments driven by the remineralization of organic matter (OM) contained in soils and forests covered during glacial margin fluctuations has been suggested as a potential carbon feedback to drive warming (Sharp et al., 1999; Wadham et al., 2008). Because the global warming potential of CH₄ is 25 times greater than CO₂, even limited subglacial methanogenesis has the potential to strongly impact the greenhouse gas composition of glacial meltwater. Combined inorganic and organic subglacial processes may therefore produce glacial meltwater that is a source or sink of greenhouse gas. While the net impact of these processes on modern carbon fluxes is poorly constrained, determining these impacts will improve modern carbon budgets as well as depictions of how fluxes may have evolved during the advance and retreat of continental ice sheets.

Deleted: Recent evaluations of carbon budgets in proglacial environments indicate that mineral weathering results in net sequestration of atmospheric CO₂, suggesting that proglacial systems are underrecognized as Arctic CO₂ sinks (St. Pierre et al., 2019)

Deleted: (GHG)

Deleted: GHG effect

Deleted: GHG.

In subglacial environments where remineralization is limited by low OM availability, the major element solute load of glacial meltwater is typically dominated by products of mineral weathering reactions (Tranter, 2005). The extent of mineral weathering in subglacial environments depends in part on the availability of acids to drive reactions, namely sulfuric and carbonic acids (Table 1). Sulfuric acid is derived from the oxidation of reduced sulfur species, which largely occur as iron-sulfide minerals including pyrite (Tranter, 2005). Sulfide oxidation may occur abiotically, however, the kinetics of microbially mediated sulfide oxidation is several orders of magnitude faster and may lead to local depletion of oxygen given a sufficient supply of sulfide minerals (Sharp et al., 1999). In contrast, carbonic acid may be derived from multiple external or *in situ* sources of CO₂ to the system. The dominant external source is supraglacial meltwater that flows to the subglacial system through moulins following equilibration with atmospheric CO₂ (Fig. 1). Unlike proglacial environments where free exchange between water and the atmosphere may resupply CO₂ consumed by weathering, subglacial environments may be partially or fully isolated from the atmosphere, limiting further atmospheric CO₂ invasion and thus the extent of mineral weathering with carbonic acid. However, additional atmospheric CO₂ may be delivered in open portions of the subglacial environment through exchange in fractures or moulins along subglacial flow paths or in partially air-filled conduits, allowing a much greater magnitude of carbonic acid mineral weathering (Graly et al., 2017). CO₂ may also be derived from *in situ* sources, such as gaseous CO₂ contained in ice bubbles of basal ice, or fluid inclusions in rocks that release volatiles (including CO₂) following mechanical grinding (Macdonald et al., 2018). When OM is available, its remineralization also generates CO₂ (and potentially CH₄) along with nutrients, but low OM availability in many subglacial systems limits remineralization as a CO₂ source (Fig. 1).

Deleted: ,

Deleted: provided

The role of subglacial carbon processes may play an increasingly important role in modern Arctic carbon budgets as disproportionate warming increases glacial meltwater and sediment fluxes to the ocean, particularly from the Greenland Ice

Sheet. The is the last remaining ice sheet in the Northern hemisphere following collapse of other ice sheets since the Last Glacial Maximum (~20 ka). It has been losing mass at increasing rates that averaged 286±20 Gt/yr between 2010-2018, representing a six-fold increase since the 1980s (Mouginot et al., 2019). While mineral weathering significantly modifies the chemical composition of Greenland Ice Sheet subglacial discharge (e.g. Hindshaw et al., 2014; Deuerling et al., 2018; Urra et al., 2019) and should consume CO₂ similar to other glacial and proglacial environments, the recent identification of microbially driven reactions (including methanogenesis) in subglacial environments of the Greenland Ice Sheet indicates that organic processes may also play a role (Dieser et al., 2014; Lamarche-Gagnon et al., 2019; Musilova et al., 2017). The relative importance of subglacial greenhouse gas sinks (CO₂ consumption through mineral weathering) and sources (such as OM remineralization) determine the greenhouse gas composition of subglacial discharge, which may then serve as a source or a sink of atmospheric greenhouse gases. Constraining the relative impacts and variability of these processes underneath the Greenland Ice Sheet will provide important information regarding the current and future impact of Greenland Ice Sheet loss on Arctic carbon budgets, as well as the role of continental ice sheets on carbon cycle feedbacks.

Deleted: (GrIS). The Greenland Ice Sheet (GrIS)

Deleted: all others

Deleted: 25

Deleted: GrIS

Deleted: GrIS

Formatted: Danish

Deleted: GHG

Deleted: GHG

Deleted: GHGs.

Deleted: GrIS

Deleted: GrIS

To evaluate the net impact of carbon processes on the greenhouse gas composition of subglacial discharge of the Greenland Ice Sheet, we compare water chemistry, dissolved CO₂ and CH₄ concentrations, and gas stable isotopic compositions between three subglacial discharge sites draining land-terminating glaciers of the Greenland Ice Sheet over the melt seasons of 2017 and 2018 (Fig. 2). We employ mass balance models utilizing the concentrations of major cations and anions to determine the magnitude of the impact on CO₂ concentrations from mineral weathering reactions (Table 1). These results are combined with measured gas concentrations and $\delta^{13}\text{C}$ to determine the relative importance of mineral weathering compared to OM remineralization on the CH₄ and CO₂ content of subglacial discharge. We also assess the temporal and spatial variability of these processes under the Greenland Ice Sheet to improve our understanding of carbon cycling in Greenland subglacial environments and the implications of Greenland Ice Sheet mass loss on Arctic carbon budgets.

Deleted: GHG

Deleted: GrIS

Deleted: GrIS

Deleted: GrIS

Deleted: GrIS

2 Methods

2.1 Study locations

Our three subglacial discharge locations are located in southwest (Fig. 2a, b) and southern (Fig. 2a, c) Greenland, (pictures given in Supplementary Material). Our sub-Isunnguata watershed (IS; 67°09'27.1" N, 50°03'25.0" W) and Russell Glacier watershed (RU; 67°05'22.1" N, 50°14'18.7" W) drain to the Akuliarusiarsuup Kuua, which is a tributary to the Qinnguata Kuussua. The short stretch of river downstream of the confluence of the Akuliarusiarsuup Kuua and the main stem of the Qinnguata Kuussua near the town of Kangerlussuaq is also known as the Watson River (Fig. 2b). The majority of drainage from the Isunnguata Glacier drains to the northern Isortoq River (Fig. 2), however a sub-catchment of the Isunnguata drains to a stream that directly feeds the Akuliarusiarsuup Kua, which we refer to as the Northern Tributary, while the Akuliarusiarsuup Kuua refers to river that flows between the outlet of the Russell Glacier and the confluence with the Qinnguata Kuussua (Fig. 2b). Watson River discharge is monitored by PROMICE (Programme for Monitoring of the

Deleted: . The

Deleted: Glacier

Deleted: Watson River

Deleted: one

Deleted: the largest proglacial rivers in Greenland.

145 Greenland Ice Sheet; van As et al., 2018) and total discharge was 4.3 and 3.6 km³ of water in 2017 and 2018, respectively (van As et al., 2018). The total catchment size for the Isunnguata is 15,900 km², though the size of the sub-catchment draining to the Northern Tributary is smaller with a drainage area of approximately 40 km² (Lindbäck et al., 2015; Rennermalm et al., 2013), therefore we hereby refer to this site as the sub-Isunnguata watershed. The total drainage area for the Russell glacier is not precisely known, however the catchment draining both the Russell and Leverett glaciers has been estimated at approximately 900 km² (Lindbäck et al., 2015) while the Leverett drainage area alone is estimated at approximately 600 km² (Hawkins et al., 2016), therefore we estimate the Russell drainage area at approximately 300 km². Discharge from the third site in southern Greenland, Kiattut Sermiat (KS; 61°12'13.5" N, 45°19'49.1"W) drains to the Kuusuaq River near the town of Narsarsuaq. While Kuusuaq discharge is not monitored, a previous study using dye tracing techniques estimated approximately 0.22 km³ of discharge in 2013, and its catchment size was estimated at 36 km² (Hawkins et al., 2016).

155 Underlying lithologies differ between sites. Southwest Qinnguata Kuussua sites are located near the boundary between the Archean Craton to the south and the southern Nagssugtoqidian Orogen to the north (Henriksen et al., 2009). The Archean Block is composed of granites and granulite facies orthogneisses that were intruded by mafic dykes during Paleoproterozoic rifting. These rocks were deformed and modified during subsequent continent-to-continent collision in the Paleoproterozoic to create the amphibolite facies gneisses of the southern Nagssugtoqidian Orogen (van Gool et al., 2002).
 160 Kiattut Sermiat lies within the Paleoproterozoic Ketilidian fold belt (Henriksen et al., 2009). Lithologies in this region include the Julianehåb Granite and associated basic intrusions and the sedimentary and volcanic rocks of the Mesoproterozoic Gardar Province that include a suite of alkaline igneous rocks and basaltic dykes with interbedded sandstones (Kalsbeek and Taylor, 1985; Upton et al., 2003).

Previous studies have characterized chemical weathering reactions in subglacial discharge to the Akuliarusiarsuup Kua and Qinnguata Kuussua (Deuerling et al., 2018; Hasholt et al., 2018; Yde et al., 2014), in the Kuusuaq that drains Kiattut Sermiat (Hawkins et al., 2016), and in comparison between these two sites (Urra et al., 2019). There has been extensive work regarding ice sheet dynamics and hydrology in the southeast glaciers draining to the Akuliarusiarsuup Kua and Qinnguata Kuussua (Van As et al., 2017, 2018; Lindbäck et al., 2015) as well as southern Kuusuaq catchments (Warren and Glasser, 1992; Winsor et al., 2014). Subglacial permafrost has been identified near the sub-Isunnguata site (Ruskeeniemi et al., 2018) and mostly likely formed during Holocene fluctuations in the ice sheet margin. While a similar Holocene ice retreat and readvance may have occurred in southern Greenland (Larsen et al., 2016), it is unknown whether this retreat led to the organic deposits below the ice sheet.

2.2 Sample collection

We collected water samples from subglacial discharge sites in spring and fall of 2017, and the summer of 2018 to observe seasonal variations in water chemistry. To minimize atmospheric influence, samples were collected as close as possible to the glacier front where subglacial waters emerge, which was less than 10 m for the sub-Isunnguata site, and approximately 100 m for the Russell glacier site. Subglacial discharge from Kiattut Sermiat site flowed through a glacial meltwater lake prior

Deleted: our water samples were collected from a smaller

Deleted: .

Deleted: 300

Deleted: . This estimate comes from subtracting

Deleted: from the

Deleted: 900

Deleted: catchment that includes the Russell and Leverett drainages (Lindbäck et al., 2015). While discharge

Deleted:),

Deleted: Watson River

Deleted:

Deleted: block

Deleted:

Deleted:

Deleted:

Deleted: Watson River

Field Code Changed

Formatted: Danish

Formatted: Danish

Formatted: Danish

Field Code Changed

Formatted: Danish

Formatted: Danish

Deleted: comparatively

Formatted: Danish

Formatted: Danish

Deleted: Watson River catchment

Deleted: Kiattut Sermiat

Deleted: . Previous studies near these study locations have identified CH₄ and CO₂ supersaturation in subglacial discharge of the Isunnguata site (Christiansen and Jørgensen, 2018; Ryu and Jacobson, 2012), while methanogenic microbial communities have been observed at Russell Glacier (Dieser et al., 2014) and CH₄ supersaturation at the Leverett Glacier (Lamarche-Gagnon et al., 2019), which also flows into the Watson River (Fig. 2c). Subglacial permafrost has been identified near the

Deleted: attributed to

Deleted: formation of

Deleted: composition. Samples

Deleted: discharge site

Deleted: .

Deleted: , and approximately 1.1 km for the

to arriving at the sampling location, which was approximately 1.1 km from the glacial outlet (Fig. 2) and therefore may experience a more interaction with atmospheric gases during transport from the subglacial discharge site to our sampling location. We collected water samples by pumping water through a 0.5-cm flexible PVC tube that was placed in flowing water as far as possible from shore (approximately 1-2 m). A YSI Pro-Plus sensor that was calibrated daily was installed in an overflow cup filled from the bottom to measure specific conductivity (Sp.C), temperature, pH, dissolved oxygen, and oxidation-reduction potential (ORP). These parameters were monitored until stable, between about 10 and 30 minutes, after which samples were collected and preserved in the field according to the solute to be measured after being filtered through a 0.45 μm trace-metal grade Geotech high-capacity disposable canister filter. Samples for cations and anions were collected in HDPE bottles; cation samples were preserved with Optima-grade ultrapure nitric acid ($\text{pH} < 2$) while no preservative was added to anion samples. Dissolved inorganic carbon (DIC) samples were filtered through 0.2 μm filters directly to the bottom of 20 ml Qorpac glass vials and allowed to overflow until sealed tightly with no headspace.

Deleted:).

Deleted:

Deleted: Samples for ammonium (NH_4) were filtered into 15 mL polypropylene containers and frozen until analysis.

Gas samples were collected in duplicate via headspace extractions according to methods outlined in Repo et al. (2007) and Pain et al. (2019). Unfiltered water was pumped into the bottom of 500 mL bottles until they overflowed. Bottles were immediately capped with rubber stoppers fitted with two 3-way inlet valves. 60 mL of water was extracted from one inlet and replaced with 60 mL of atmospheric air (for spring and fall 2017 sampling trips) or ultrapure N_2 gas in a gas bag (summer 2018 sampling trip). Bottles were shaken for 2 minutes to equilibrate headspace gas with water, and headspace gas was extracted and immediately injected into 60 ml glass serum bottles that had been evacuated immediately prior to sample introduction. Samples were stored at room temperature until analysis, which occurred within one week of collection. Measured headspace concentrations were converted to dissolved concentrations using methods outlined in Pain et al. (2019). When atmospheric air was used for headspace extractions, atmosphere samples were collected in tandem and analyzed to correct the calculated dissolved CO_2 and CH_4 concentrations and isotopic compositions for atmospheric CO_2 and CH_4 . This correction altered CH_4 concentrations by up to 22% for one sample from the Russell glacier, though less than 5% for all other samples, and resulted in a correction of $\delta^{13}\text{C}-\text{CH}_4$ of up to 1.3‰. For CO_2 , the correction altered concentrations by up to 15% for one sample collected at Kiattut Sermiat, though less than 10% for all other samples, and resulted in a correction of $\delta^{13}\text{C}-\text{CO}_2$ of up to 0.4‰.

Deleted: each sample for

Deleted: .

In samples collected in fall 2017 and summer 2018, alkalinity was measured in the field laboratory within 3 days of collection by titration with 0.01 N HCl using the Gran method. Because alkalinity measurements were not available for the spring 2017 sampling trip, we estimate alkalinity with PHREEQC modeling and the phreeqc.dat database (Parkhurst, 1997) using major cations and anions, pH, temperature, and DIC concentrations as model inputs.

Deleted: For

Deleted: sampling trips

2.3 Laboratory analysis

Gas samples were analyzed for CO_2 and CH_4 concentrations, and $\delta^{13}\text{C}-\text{CO}_2$ and $\delta^{13}\text{C}-\text{CH}_4$ on a Picarro G2201-i cavity ring-down spectrometer, in the field within a few days of collection. Carbon isotopic compositions are reported in reference to Vienna Pee Dee Belemnite (VPDB). Check standards of known CO_2 and CH_4 concentrations and isotopic compositions were

Deleted: .

measured during each sample run and were accurate within 10%. Anion and cation concentrations were measured on an automated Dionex ICS-2100 and ICS-1600 Ion Chromatograph, respectively. Error on replicate analyses was less than 5%.
255 DIC concentrations were measured on a UIC (Coulometrics) 5011 CO₂ coulometer coupled with an AutoMate Preparation Device. Samples were acidified and the evolved CO₂ was carried through a silver nitrate scrubber to the coulometer where total C was measured. Accuracy was calculated to be ±0.1 mg/L based on measurement of check standards. ▾

Deleted: Dissolved ammonium (NH₄) concentrations were analyzed on a Seal AutoAnalyzer III. Error on check standards was less than 10%.

2.4 Methane modeling

To assess CH₄ sources and sinks, we calculate ε_c, or the carbon isotopic fractionation factor between CO₂ and CH₄ as
260 defined in Whiticar (1999):

$$\epsilon_c = \delta^{13}C_{CO_2} - \delta^{13}C_{CH_4} \quad (5)$$

Formatted: Left, Indent: Left: 0.5"

Values of ε_c reflect methanogenesis pathways (acetoclastic or CO₂ reduction) as well as the extent of oxidation. Values of ε_c between approximately 40 and 55‰ are produced for CH₄ generated via acetoclastic methanogenesis, while CO₂ reduction produces values between approximately 55 and 90‰. Lower values (ε_c between 5 and 30) result when CH₄ oxidation
265 predominates. Modern atmospheric input without additional alteration of CO₂ or CH₄ isotopic systematics results in a ε_c value of approximately 40 (Whiticar, 1999).

Deleted: produced

We calculated CH₄ oxidation using the isotopic method outlined in Mahieu et al. (2008) and Preuss et al. (2013). The fraction of oxidized methane (f_{ox}) in an open system is given by:

$$f_{ox} = \frac{\delta_E - \delta_P}{1000(\alpha_{ox} - \alpha_{trans})} \quad (6)$$

Deleted:

where δ_E is the measured δ¹³C-CH₄ value for each water sample, δ_P is δ¹³C-CH₄ of produced methane, α_{ox} is the oxidation
270 fractionation factor, and α_{trans} is a fractionation factor resulting from diffusive transportation of CH₄. While the exact value of δ_P is unknown, diagenetic alteration of δ¹³C-CH₄ values through oxidation or transport only enrich δ¹³C-CH₄ signatures, therefore the value of δ_P is taken as the most depleted δ¹³C-CH₄ signature assuming it is the least impacted by diagenetic alteration. Literature-reported values for α_{ox} range between 1.003 and 1.049. We calculate the fraction of oxidized methane with the largest fraction factor (α_{ox} = 1.049; Mahieu et al., 2008), which yields the minimum amount of CH₄ oxidation required
275 to explain the observed variations in δ¹³CH₄, and thus is a conservative estimate for CH₄ oxidation, and actual oxidation ratios may be higher. Literature-reported values for α_{trans} vary from 1 for advection-dominated systems to 1.0178 for diffusion-dominated porous media (Visscher et al., 2004; Mahieu et al., 2008; Preuss et al., 2013). We assume that transport is advection dominated and thus assume α_{trans} = 1, however diffusive transport of CH₄ may result in fractionation of CH₄ in the subglacial environment and lead to relatively lower estimates of f_{ox}. Because hydrologic connectivity between subglacial methanogenic
280 meltwater pockets and drainage features is not well described, the relative importance of advective compared to diffusive CH₄

Deleted: .

Deleted: We assume that transport is advection dominated and thus assume α_{trans} = 1

290 [transport within the subglacial drainage system is not well understood; however, it is presumed to be an advection-dominated system in which expanding drainage networks access and drain methanogenic meltwater pockets throughout the melt season.](#)

2.5 Mineral weathering and carbonate modeling

We used major cation and anion concentrations and alkalinity to partition solutes into the four mineral weathering reactions in Table 1 after correcting solute concentrations for marine aerosol deposition using measured chloride concentrations and standard seawater element ratios. The mass balance model followed the methods of Deuerling et al. (2019).
295 After apportioning solutes to mineral weathering reactions, we used the stoichiometries of reactions to calculate the impact of each reaction on dissolved CO₂ concentrations (Table 1). The mineral weathering model apportions solutes to reactions in Table 1 based on the ratios of Ca/Na and Mg/Na in silicate minerals in stream bedload samples, which were taken to be 0.54 and 0.38, respectively, for [sub-Isunnguata](#) and Russell Glacier samples (Deuerling et al., 2019; Hindshaw et al., 2014; Wimpenny et al., 2010, 2011) and 0.39 and 0.27, respectively, for Kiattut Sermiat samples (Da Prat and Martin, 2019). Because
300 mineral weathering reactions may both add and remove CO₂, we discuss both the net impact of mineral weathering on CO₂ concentrations (Net CO_{2-MW}), which may have a positive or negative value:

$$[\text{Net CO}_{2\text{-MW}}] = [\text{CO}_{2\text{-CarbCA}}] + [\text{CO}_{2\text{-CarbSA}}] + [\text{CO}_{2\text{-SilCA}}] \quad (7)$$

as well as the total impact of mineral weathering on CO₂ concentrations (Total CO_{2-MW}),

$$[\text{Total CO}_{2\text{-MW}}] = |[\text{CO}_{2\text{-CarbCA}}]| + |[\text{CO}_{2\text{-CarbSA}}]| + |[\text{CO}_{2\text{-SilCA}}]| \quad (8)$$

305 where changes in the concentrations of CO₂ are defined by their absolute values. To discuss the relative importance of individual reactions, we define proportional contributions of each reaction as follows:

$$\% \text{CO}_{2\text{-CarbCA}} = \frac{|[\text{CO}_{2\text{-CarbCA}}]|}{[\text{Total CO}_{2\text{-MW}}]} * 100 \quad (9a)$$

$$\% \text{CO}_{2\text{-CarbSA}} = \frac{|[\text{CO}_{2\text{-CarbSA}}]|}{[\text{Total CO}_{2\text{-MW}}]} * 100 \quad (9b)$$

$$\% \text{CO}_{2\text{-SilCA}} = \frac{|[\text{CO}_{2\text{-SilCA}}]|}{[\text{Total CO}_{2\text{-MW}}]} * 100 \quad (9c)$$

310 We combine measured CO₂ concentrations with Net CO_{2-MW} in order to determine the magnitude of CO₂ production or consumption in the subglacial environment due to processes besides mineral weathering. This analysis assumes that the concentration of CO₂ measured at the subglacial outlet is equivalent to the net change in CO₂ due to mineral weathering plus the sum of all other subglacial CO₂ sources and sinks. We refer to the sum of all other subglacial CO₂ sources and sinks as CO_{2-total}, which represents the amount of CO₂ that must have been supplied to the subglacial environment to balance the mineral
315 weathering CO₂ sink:

$$\text{CO}_{2\text{-measured}} = \text{Net CO}_{2\text{-MW}} + \text{CO}_{2\text{-total}} \quad (10)$$

Formatted: No widow/orphan control

Formatted: Don't keep lines together

The sources of CO₂ to CO_{2-total} may be evaluated through the use of Keeling plots, which are constructed as the inverse of CO₂ concentrations ([CO₂]⁻¹) versus stable isotopic composition (δ¹³C-CO₂). If variations in the concentration and isotopic composition of CO₂ arise from the mixing of two CO₂ reservoirs with constant isotopic compositions and concentrations (Keeling, 1958), a linear relationship is expected between [CO₂]⁻¹ and δ¹³C-CO₂. The y-intercept of a regression between these variables represents the isotopic composition of the high-CO₂ end member. Because measured CO₂ concentrations include both subglacial CO₂ sources and sinks, which may include considerable consumption through mineral weathering reactions, the magnitude of the total subglacial CO₂ source is taken as CO_{2-total}. We therefore construct Keeling plots between [CO_{2-total}]⁻¹ and measured δ¹³C-CO₂ values because while mineral weathering impacts the concentration of CO₂, its isotopic composition is not appreciably altered (Myrntinen et al., 2012) compared to the range of isotopic compositions of potential CO₂ end members, namely OM remineralization, atmospheric CO₂, and lithogenic CO₂ sources due to mechanical grinding (Fig. 1).

2.6 Discharge relationships

We evaluate the relationship between subglacial CH₄ and CO₂ dynamics and glacial meltwater river discharge records collected downstream of the sub-Isunnguata and upstream of the Russell sampling sites. Proglacial river discharge was collected in the Akuliarusiarsuup Kuua (AK) River at the AK4 site, 2 km downstream of the sub-Isunnguata sampling site (Fig. 2b). The river discharge dataset is an updated and extended version of Rennermalm et al. (2012) using reference and regression models to correct Solinst level logger drift in water stage (Solinst, 2017), and a total of 57 discharge measurements to convert continuous water stage to discharge. The standard uncertainty (i.e. the 68th percent confidence interval or one standard deviation) was determined to be 17% using methods and recommendations provided by (Hersch, 1999; ISO Guide 98-3, 2008; WMO, 2010).

Because diurnal fluctuations in river discharge can be large, and differing water travel times from subglacial outlet sites to the discharge monitoring site induces a lag between maximum daily discharge at subglacial discharge sites and the AK4 site outlet, we compare subglacial CH₄ and CO₂ concentrations to average daily discharge, calculated as the average of hourly discharge estimates over the days on which subglacial discharge water samples were collected. We use the AK4 site discharge records for evaluating concentration-discharge relationships for both sub-Isunnguata and Russell sites. Although this site is upstream of the Russell Glacier, its close proximity to the Russell Glacier suggests it is more likely to reflect local melting patterns similar to those that would be controlling discharge dynamics at the Russell than discharge records collected at the Watson River outlet. While Watson River discharge records are also available through PROMICE (van As et al., 2018), which includes some contributions from the Russell glacier, the Watson River includes discharge from the Akuliarusiarsuup Kuua (draining sub-Isunnguata, Russell, and Leverett catchments) as well as the much larger Qinnguata Kuussua catchment and therefore Watson River discharge records are not likely to be representative of the temporal changes in the magnitude and variability of discharge from the much smaller Russell glacier catchment.

Deleted: → We evaluate the relationship between subglacial CH₄ and CO₂ dynamics and discharge using discharge records provided by PROMICE (Programme for Monitoring of the Greenland Ice Sheet; van As et al., 2018). Discharge records are collected at the outlet of the Watson River, which represents the combined discharge of Isunnguata and Russell glaciers as well as other outlet glaciers including the Leverett Glacier and major tributaries including the Orkendalen River (Fig. 2). Watson River discharge estimates are therefore greater than the true discharge of our individual sampling locations, however we assume that discharge at the Isunnguata and Russell subglacial outlet sites is roughly proportional to Watson River discharge and exhibits similar temporal variability (Rennermalm et al., 2012). Because diurnal fluctuations in river discharge can be large, and differing water travel times from subglacial outlet sites to the Watson River mouth induces a lag of up to 8 hours between maximum daily discharge at subglacial discharge sites and the Watson River outlet, we compare subglacial CH₄ and CO₂ concentrations to average daily discharge, calculated as the average of hourly discharge estimates over the days on which subglacial discharge water samples were collected. Because no discharge information is available for Kiattut Sermiat, we assess discharge relationships at Watson River (Isunnguata and Russell) sites only.

3 Results

3.1 Temporal variability in water chemistry and gas concentrations

Chemical parameters differ spatially between subglacial discharge sites as well as temporally through the 2017-2018 melt seasons. Comparing the means and standard deviations of water samples collected throughout 2017 and 2018, specific conductivity (Sp.C; Fig. 3a) is typically highest at Kiattut Sermiat (26 ± 8 $\mu\text{S}/\text{cm}$), followed by Russell (22 ± 5 $\mu\text{S}/\text{cm}$) and sub-Isunnguata sites (13 ± 9 $\mu\text{S}/\text{cm}$; Fig. 3a). All sites show variability over time, with lowest values occurring in the summer for sub-Isunnguata and Russell, while Sp.C drops continuously with days of the year for Kiattut Sermiat. Sites differ in pH, and average values at Kiattut Sermiat (8.2 ± 0.4) are higher than both Russell (7.2 ± 0.2) and sub-Isunnguata (6.6 ± 0.6 ; Fig. 3b), and while values vary over time, no consistent trend is identified between sites. The saturation of dissolved oxygen (D.O.) with respect to atmospheric concentrations is similar between sites, though sub-Isunnguata ($98\pm 8\%$) values fall below Russell ($115\pm 16\%$) and Kiattut Sermiat ($117\pm 11\%$) during all sampling times and exhibit undersaturation in the mid-summer samples, while Russell and Kiattut Sermiat are consistently supersaturated (Fig. 3c). Alkalinity is similar at Russell (93 ± 31 $\mu\text{eq}/\text{L}$) and Kiattut Sermiat (93 ± 26 $\mu\text{eq}/\text{L}$) which are higher than at sub-Isunnguata (39 ± 25 $\mu\text{eq}/\text{L}$), but all reach minimum values in summer (Fig. 3d). CH_4 concentrations differ by orders of magnitude between sites (Fig. 3e) and are consistently supersaturated with respect to atmospheric concentrations at sub-Isunnguata (648 ± 411 ppm or 1575 ± 997 nM) and Russell (58 ± 33 ppm or 110 ± 78 nM) sites, while close to atmospheric equilibrium at Kiattut Sermiat (4 ± 2 ppm or 9 ± 5 nM). Mean $\delta^{13}\text{C}-\text{CH}_4$ values (Fig. 3f) are similar between sub-Isunnguata ($-54.7\pm 7.5\%$), Russell ($-52\pm 7.3\%$), and Kiattut Sermiat ($-57.6\pm 14.2\%$). Measured CO_2 concentrations (Fig. 3g) are consistently supersaturated with respect to atmospheric concentrations for sub-Isunnguata (685 ± 230 ppm or 58 ± 18 μM), near atmospheric equilibrium for Russell (442 ± 31 ppm or 29 ± 4 μM) and undersaturated for Kiattut Sermiat (263 ± 33 ppm or 19 ± 2 μM). Mean $\delta^{13}\text{C}-\text{CO}_2$ values (Fig. 3h) are lower in spring and fall for sub-Isunnguata ($-16.6\pm 4.0\%$) compared to Russell ($-13.7\pm 2.3\%$) and Kiattut Sermiat ($-16.1\pm 1.6\%$) sites, though similar seasonal variation occurs for all sites with relatively more depleted values in the spring and fall compared to summer.

3.2 Methane oxidation and relationship with discharge

Values of ϵ_c are similar over time for sub-Isunnguata ($38\pm 10\%$) and Russell ($38\pm 9\%$) and are relatively higher in the summer sampling period, while Kiattut Sermiat ϵ_c values are higher on average ($42\pm 13\%$) with lowest values in the summer (Fig. 4a). Estimates of f_{ox} are similar between sub-Isunnguata ($17\pm 15\%$), Russell ($23\pm 15\%$), and Kiattut Sermiat sites ($25\pm 22\%$; Fig. 4b). However, f_{ox} values are higher in the spring and fall sampling times compared to summer for sub-Isunnguata and Russell and approach 50% in the spring, while Kiattut Sermiat values decrease throughout the melt season.

CH_4 concentrations, $\delta^{13}\text{C}-\text{CH}_4$ values, and f_{ox} are negatively correlated to average daily discharge for both sub-Isunnguata and Russell sites (Fig. 5a, b and d), while ϵ_c is positively correlated with discharge for both sub-Isunnguata and Russell (Fig. 5c).

Deleted: the melt season. Specific

Deleted: throughout the melt season

Deleted: throughout

Deleted: melt season

Deleted: KS.

Deleted: throughout the melt season

Deleted:),

Deleted: throughout the melt season

Deleted: %, Fig. 4a)

Deleted: .

Deleted: unrelated

Deleted: Watson River

Moved down [1]: 5c).

Deleted: , but significantly negatively correlated for

Deleted:). The fraction of CH_4 oxidized is moderately negatively correlated with discharge for both Isunnguata and Russell, though the relationship is not significant for either site (Fig. 5b). Discharge

Deleted: ϵ_c for both Isunnguata and Russell (Fig.

Deleted: While the relationship is only significant for Isunnguata, the slopes and intercepts of regression lines for

Moved (insertion) [1]

Deleted: are similar.

3.3 Mineral weathering impacts on CO₂ and relationship with discharge

Mineral weathering leads to net sequestration of CO₂ at all three sites (Fig. 6a). The magnitude of Net ΔCO₂ differs between sites with the lowest average values at sub-Isunnguata (-39±37 μM) followed by Russell (-65±32 μM) and Kiattut Sermiat (-98±17 μM) sites. Individual mineral weathering reactions produce differing contributions between sites and over time, with notable differences between southwest sites (sub-Isunnguata and Russell) and the southern Kiattut Sermiat site (Fig. 6b). For instance, the proportional contribution of Carb_{SA} is similar between sub-Isunnguata (17±11%) and Russell (15±6%), but lower at Kiattut Sermiat (8±1%; Fig. 6b). Kiattut Sermiat has a relatively greater contribution from Carb_{CA} (62±2%) compared to sub-Isunnguata (41±10%) and Russell (38±6%), while Sil_{CA} is lower at Kiattut Sermiat (28±1%) compared to sub-Isunnguata (41±17%), and Russell (47±11%). Kiattut Sermiat additionally exhibits low seasonal variability in the proportional contributions of individual mineral weathering reactions compared to sub-Isunnguata and Russell sites.

CO_{2-total} represents CO₂ concentrations in the subglacial environment prior to addition and/or consumption of CO₂ through mineral weathering (Eq. 10; Fig. 7). Because the Net CO_{2-MW} is always negative (more consumption than production), the value of CO_{2-total} is always greater than measured concentrations (CO_{2-measured}). Regardless of differences in CO_{2-measured} between sites, the average CO_{2-total} values are similar between sites and average 91±47 μM for sub-Isunnguata, 94±33 μM for Russell, and 117±16 μM for Kiattut Sermiat.

For both sub-Isunnguata and Russell sites, average daily discharge is negatively correlated with CO₂ concentrations (Fig. 8a), while positively correlated with δ¹³C-CO₂ (Fig. 8b).

Keeling plots between [CO_{2-total}]⁻¹ and δ¹³C-CO₂ for each site indicate no linear relationship for Russell or Kiattut Sermiat samples, however a strong linear correlation is observed for sub-Isunnguata (r²=0.99; p<0.001) samples with the removal of one outlier, which also had the lowest CO_{2-total} value (Fig 9).

4 Discussion

We observe orders of magnitude variability in dissolved CH₄ and CO₂ concentrations in subglacial discharge of the Greenland Ice Sheet, indicating significant differences in the magnitudes of the sources and sinks of these gases across time and space. Supersaturation of both CO₂ and CH₄ with respect to atmospheric concentrations indicates that sub-Isunnguata discharge is a source of both gases to the atmosphere, neighbouring Russell Glacier discharges water that is a source of CH₄ but near equilibrium with respect to CO₂, while Kiattut Sermiat in southern Greenland is a sink of atmospheric CO₂ but near equilibrium with respect to CH₄ (Fig. 3e, g). Because CH₄ dynamics may be largely microbially driven while CO₂ dynamics include microbial as well as abiotic mineral weathering processes, we first discuss CH₄ dynamics including a comparison of concentrations, isotopic compositions, and extent of oxidation between sites and over the melt season. We then discuss CO₂ concentrations, impacts of mineral weathering reactions (Table 1), and an assessment of subglacial CO₂ sources, including

Deleted: and CO₂ models

Deleted: lowest

Deleted:).

Deleted: the melt season

Deleted: Watson River

Deleted: though is

Deleted: then

Deleted: 8a). Isunnguata samples also show a significant positive correlation between the magnitude of CO_{2-total} and NH₄ concentrations (r²=0.73; p<0.05), unlike Russell or Kiattut Sermiat samples, though several Russell samples fall close to the regression line between CO_{2-total} and NH₄ of Isunnguata samples (Fig. 8b)

Deleted: Because Isunnguata samples exhibit a linear relationship between [CO_{2-total}]⁻¹ and δ¹³C-CO₂ consistent with two endmember mixing, we utilize the δ¹³C-CO₂ of samples and end members defined by the Keeling plot in an isotopic mixing model to calculate the relative contributions of CO₂ sources to CO_{2-total}. We take the y-intercept (-27.5‰; Fig. 8a) of the Keeling plot regression as the δ¹³C-CO₂ for the high-CO₂ end member (assumed to be generated by OM remineralization; CO_{2-OM}) and assume atmospheric CO₂ as the low-CO₂ end member (CO_{2-atm}; δ¹³C-CO₂ = -8‰). The relative (%CO_{2-atm} and %CO_{2-OM}) and absolute (CO_{2-atm} and CO_{2-OM}) contributions of these CO₂ sources to Isunnguata CO_{2-total} vary throughout the melt season, with CO_{2-OM} being the dominant source in the early and late melt season while CO_{2-atm} is the dominant source in the mid-melt season (Fig. 9a). The absolute magnitude of CO_{2-OM} is approximately 5-10 times greater in the early (89 μM) and late (117 μM) melt seasons compared to mid-melt seasons (13-23 μM), while the magnitude of CO_{2-atm} varies relatively little throughout the melt season at around 50 μM (Fig. 9b). Both %CO_{2-atm} and %CO_{2-OM} are significantly (p<0.05) correlated with Watson River discharge, and the correlation is positive for %CO_{2-atm} and negative for %CO_{2-OM} (Fig. 9c). The magnitude of CO_{2-OM} exhibits a power-law relationship with discharge and is highest at lowest discharge, while the magnitude of CO_{2-atm} is invariable as a function of discharge (Fig. 9d).

Deleted: GrIS

Deleted: neighboring

OM remineralization. These assessments will contribute to our understanding of the variability and controls of CH₄ and CO₂ concentrations in subglacial discharge from the Greenland Ice Sheet and may improve predictions of the impact of future ice melt on Arctic carbon budgets.

4.1 Sources and sinks of CH₄

495 Differences in CH₄ concentrations and relationships with discharge between sites imply heterogeneity in both the extent and controls of subglacial methanogenesis under the Greenland Ice Sheet. CH₄ supersaturation occurs at the two subglacial discharge sites that flow to the Akuliarusiarsuup Kuua (sub-Isunnguata and Russell), and concentrations are similar to the ranges reported in discharge of the Leverett Glacier (up to 600 nM; Lamarche-Gagnon et al., 2019), located near the Russell Glacier in this study (Fig. 2b). However, CH₄ concentrations are near atmospheric equilibrium for the Kiattut Sermiat 500 site (Fig. 3e). Because methanogenesis is an anaerobic OM remineralization pathway, it is more likely to occur in subglacial environments isolated from atmospheric O₂ sources. Widespread observations of methanogenesis in glacial meltwater of southwest Greenland from this and other studies (Christiansen and Jørgensen, 2018; Dieser et al., 2014; Lamarche-Gagnon et al., 2019), and limited observations of CH₄ in subglacial discharge in southern Greenland, suggests heterogeneity in subglacial conditions that support methanogenesis. Methanogenesis fuelled by organic material overridden during ice sheet growth has 505 been suggested as a potential climate feedback over glacial interglacial timescales (Wadham et al., 2008), and may contribute to observed variations in CH₄ concentrations.

Subglacial methane concentrations may additionally be controlled by hydrologic factors as the subglacial hydrological network develops throughout the melt season and channelization of meltwater conduits increases subglacial drainage efficiency (Andrews et al., 2015; Cowton et al., 2013). Drainage efficiency impacts both subglacial water residence time as well the transport of aerobic supraglacial meltwater to the ice bed. Both residence time and oxygen delivery may impact subglacial redox status and methanogenesis potential, and favour methanogenesis when oxygen supply rates are low compared to OM remineralization rates. This condition is most likely to be met in distributed subglacial systems that are hydrologically isolated with limited inputs from aerobic supraglacial meltwater. Such a hydrologic control on methanogenesis is supported by the negative correlation between CH₄ concentrations and average daily discharge at both sites (Fig. 5a). This correlation would suggest that either CH₄ production occurs predominantly during periods of low discharge and greater residence time, 515 or higher discharge results in the dilution of a CH₄-laden subglacial water source. While both mechanisms would lead to a similar relationship between discharge and CH₄ concentrations, they carry different implications for subglacial methanogenesis. If limited by residence time, a hydrologic link between glacial hydrology and subglacial biogeochemistry would be established due to the importance of supraglacial discharge for the delivery of terminal electron acceptors to the ice bed for the active production of CH₄. If predominantly controlled by dilution, however, active methanogenesis would not be required; only the existence of meltwater pockets containing CH₄ that may or may not have been recently produced. Further discussion of these mechanisms is outside the scope of this study. While we have limited data to make further inferences about

Deleted: GrIS

Deleted: GrIS.

Deleted: Watson River (

Deleted: adjacent to

Deleted: .

Deleted: 2e

Deleted: This heterogeneity may include variations in the distribution of subglacial organic substrates, such as subglacial permafrost reservoirs identified in previous studies (Ruskeeniemi et al., 2018), which were implicated as an OM source to drive methanogenesis under the Leverett Glacier (Lamarche-Gagnon et al., 2019). Organic deposits under the GrIS may have formed as a consequence of Holocene ice margin fluctuations following the encroachment of ice over previously established soils and terrestrial vegetation (Ruskeeniemi et al., 2018). Methanogenesis fueled

Deleted: between southwest and southern sites in this study

Deleted: methanogenesis

Deleted: .

Deleted: favor

Deleted: at the Russell Glacier

Deleted: significant

Deleted: Watson River

Deleted:), suggesting that

Deleted: . Alternatively,

Deleted: CH₄ concentrations during low

Deleted: could result from

Deleted: relatively small volumes of methanogenic

Deleted: meltwater with increasing volumes of aerobic supraglacial meltwater. The lack of

Deleted: CH₄ concentrations and

Deleted: at the Isunnguata Glacier may indicate a greater influence of

Deleted: outburst events, in which hydrologically isolated methanogenic meltwater pockets are stochastically drained as the

Deleted: drainage network extends throughout the melt season (Fig. 5a). Subglacial outburst events were also implicated by heterogeneous CH₄ concentrations in subglacial

Deleted: of

Deleted: Leverett Glacier, and could contribute to

hydrologic controls of methanogenesis, the presence of several outliers at the sub-Isunnguata site in particular (Fig. 5a) highlights the possibility for additional controls including stochastic drainage events or heterogeneity in subglacial CH₄ concentrations that result in variability in the relationship between concentration and discharge, as was observed in Lamarche-Gagnon et al. (2019).

Deleted: CH₄ concentrations at the Isunnguata Glacier (Lamarche-Gagnon et al., 2019)

While our results suggest heterogeneity in the extent of methanogenesis between outlet glaciers, they suggest homogeneity of the microbial methanogenesis pathway as well as CH₄ oxidation dynamics between sites. Methanogenesis pathways may be evaluated by $\delta^{13}\text{C-CH}_4$ as well as ϵ_c values because they impart distinct $\delta^{13}\text{C}$ signatures to CH₄ and CO₂ (Whiticar and Schoell, 1986). Dieser et al. (2014) measured a microbial $\delta^{13}\text{C-CH}_4$ production signal at the Russell Glacier with values between -63‰ and -64‰, which was interpreted to reflect a possible combination of CH₄ produced through both acetoclastic and CO₂ reduction pathways. The most depleted $\delta^{13}\text{C-CH}_4$ value measured at the sub-Isunnguata in this study was -62.7‰, close to values measured by Dieser et al. (2014) (Fig. 3f), and similar to values reported by Lamarche-Gagnon et al. (2019) for the Leverett Glacier. The similar isotopic ratio between our samples and that measured in active methanogenic communities suggests similar methanogenesis pathways occur across this region, and additionally suggests that the CH₄ we measure in the sub-Isunnguata comes from active methanogenesis rather than old preserved subglacial CH₄ reservoirs, at least in the peak melt season when we observe these depleted $\delta^{13}\text{C-CH}_4$ values.

Deleted: and controls

Deleted: are consistent

Deleted: methanogenesis pathways

Deleted: different isotopic

Deleted: is

Deleted: that of

Deleted: at -62.7‰

Deleted: , suggesting

Deleted: .

While the exact contributions from each methanogenesis pathway cannot be inferred from isotopic information alone, the range of ϵ_c values at outlet glaciers are consistent with predominantly acetoclastic methanogenesis during the peak melt season (Fig. 4a). However, ϵ_c values fall below the expected range from acetoclastic methanogenesis during the early and late melt seasons, and may result from variations in the extent of subglacial CH₄ oxidation. Seasonal variation in CH₄ oxidation is supported by consistency between ϵ_c and f_{ox} values, which both indicate the greatest impact of oxidation (approaching 50%) in the early melt season compared to peak melt season (Fig. 4a, b), with additional evidence of elevated CH₄ oxidation in the late melt season at both sub-Isunnguata and Russell sites. Because our water sampling locations were slightly downstream of glacial discharge outlets, there is also the possibility that outgassing in between the outlet and our sampling location reduced dissolved CH₄ concentrations and led to more enriched isotopic compositions of remaining dissolved CH₄. While it is likely that some outgassing did occur, it is unlikely that the extent of outgassing between the glacial outlet and our sampling location would vary significantly between sampling times, and thus outgassing would not explain temporal differences in concentration, $\delta^{13}\text{C-CH}_4$, f_{ox} , or ϵ_c . While our measured gas concentrations and isotopic compositions likely reflect some modification of CH₄ and CO₂ isotopic compositions due to outgassing, the differences over time are more likely due to changes in subglacial CH₄ dynamics than outgassing.

Deleted: , likely resulting

Deleted: Isunnguata and Russell glaciers

Formatted: Font color: Text 1

The extent of CH₄ oxidation may be controlled by multiple factors including oxygen availability, subglacial residences time, and subglacial hydrology, similar to methanogenesis. A hydrologic control of CH₄ oxidation is supported by relationships between f_{ox} and ϵ_c with average daily discharge (collected at site AK4; Fig. 2b) at both sub-Isunnguata and Russell

Deleted: Watson River

sites: f_{ox} is negatively related with discharge for both sites (Fig. 5b) while ϵ_c is positively correlated with discharge (Fig. 5c). These correlations suggest that CH_4 oxidation is greatest during periods of low flow, which may be associated with longer residence times to allow subglacial CH_4 oxidation, however this relationship could also result from differences in CH_4 sources throughout the melt season as the subglacial drainage network expands. Assuming the former, the delivery of oxygen to the subsurface by supraglacial melting does not appear to be a limiting factor in subglacial CH_4 oxidation, which should increase f_{ox} as more oxygenated supraglacial water is delivered to the subglacial system. Instead, the observed greater CH_4 oxidation during periods of low discharge may reflect mixing of methane-rich subglacial meltwater pockets and aerobic subglacial meltwater leading to CH_4 oxidation. Longer transit times during periods of low flow may allow more subglacial methane oxidation to occur than during peak discharge, when the development of channelized flow paths reduces meltwater residence time in the subglacial environment.

Our results indicate a high degree of heterogeneity in subglacial methanogenesis under the Greenland Ice Sheet, as well as a significant impact of CH_4 oxidation, which serves to reduce atmospheric CH_4 fluxes. Given the observed spatial and temporal heterogeneity of CH_4 concentrations and processes, further investigation of the spatial variability in outlet glacier CH_4 concentrations is needed to determine the impact of Greenland Ice Sheet loss on Arctic and global CH_4 budgets, while a better understanding of the controls of these differences will improve models of how CH_4 fluxes from subglacial discharge will change with continued warming.

4.2 Sources and sinks of CO_2

Dissolved CO_2 concentrations in subglacial discharge are consistently supersaturated with respect to atmospheric concentrations at the sub-Isumnguata site, near atmospheric equilibrium at the Russell Glacier, and undersaturated at Kiattut Sermiat Glacier, indicating that glacial meltwater from the Greenland Ice Sheet can serve as either a source or sink of CO_2 to the atmosphere. Similar to CH_4 differences in dissolved CO_2 dynamics (Fig. 3g) imply variability in carbon processes under the Greenland Ice Sheet. We first discuss potential subglacial CO_2 sources, including OM remineralization, followed by a discussion of CO_2 consumption due to mineral weathering.

4.2.1 Subglacial CO_2 sources

There are many potential sources of CO_2 in the subglacial environment including dissolution of atmospheric gases in air-filled conduits or fractures in ice, CO_2 contained in ice bubbles (Fig. 1; Anklin et al., 1995; Graly et al., 2017) mechanical grinding and volatilization of fluid inclusions in bedrock (Macdonald et al., 2018), and OM remineralization. While previous studies have indicated that additional atmospheric CO_2 input through fractures and air-filled conduits may supply sufficient CO_2 to drive mineral weathering observed in many subglacial environments, including several sites in Greenland (Graly et al., 2017), CO_2 is also a product of OM remineralization, which is believed to account for CH_4 concentrations elevated above atmospheric equilibrium at the two southwest sites in this study. Both CO_2 and CH_4 exhibit negative correlations with average

- Deleted: Glaciers
- Deleted: 5c) although the correlation is only significant at Isumnguata.
- Deleted: greater
- Deleted: . The
- Deleted: CH_4 oxidation following
- Deleted: between draining methanogenic
- Deleted: with
- Deleted: GrIS
- Deleted: in this study
- Deleted: GrIS
- Deleted: Glacier
- Deleted: GrIS
- Deleted: Differences
- Deleted: concentrations also
- Deleted: GrIS (Fig. 3g). Because our approach to assess the magnitude of
- Deleted: (
- Deleted: subglacial
- Deleted:) depends in part on modelling results of
- Deleted: by
- Deleted: (Eq. 10), we first discuss the impacts of mineral weathering reactions, followed by a discussion of subglacial CO_2 sources, including OM remineralization.
- Deleted: sink: mineral weathering reactions
- Deleted: → Although mineral weathering reactions may either increase or decrease dissolved CO_2 concentrations (Table 1), the net impact of mineral weathering at our study sites is to consume CO_2 (Fig. 6a
- Moved down [2]:). Net consumption occurs because the CO_2 source from Carbs_A is ubiquitously low compared to sinks from
- Deleted: 6a), but average values increase from Kiattut Sermiat to 1
- Moved down [3]: readily weatherable basaltic intrusions as has
- Deleted: (Tranter, 2005). At Isumnguata and Russell glaciers, 2
- Moved down [4]: Isumnguata and Russell sites, Carbs_A is notably
- Deleted: Observations of higher CH_4 concentrations as well as 3
- Moved down [5]: may therefore be linked to subglacial
- Deleted: 4.2.2 Subglacial CO_2 sources 4
- Deleted: (Fig. 1; Anklin et al., 1995; Graly et al., 2017). CO_2 m 5
- Deleted: or
- Deleted: the observed extent of
- Deleted: Watson River
- Deleted: reflects OM remineralization that would also contribute 6

daily discharge for both sub-Isunnguata and Russell sites, and could suggest a common OM remineralization source (Fig. 8a).

750 While the magnitude of this source and its relative importance compared to other subglacial CO₂ sources is currently unknown, differing sources of carbonic acid for mineral weathering reactions carry different implications for subglacial CO₂ budgets. For instance, carbonic acid weathering driven by invasion of atmospheric CO₂ would represent a sink of atmospheric CO₂, but carbonic acid weathering driven by OM remineralization would instead serve to consume CO₂ from *in situ* sources and limit the potential for subglacial meltwater to be an atmospheric CO₂ source once discharged from the glacier. Determining the sources of carbonic acid to subglacial weathering reactions is therefore critical to understand the controls of mineral weathering in subglacial environments and its role in atmospheric CO₂ sequestration.

760 Comparisons between measured $\delta^{13}\text{C-CO}_2$ in subglacial discharge samples and likely $\delta^{13}\text{C-CO}_2$ values of CO₂ sources indicate that CO₂ sources differ between sites, with OM remineralization as the most important CO₂ source at the sub-Isunnguata but likely not the predominant or sole source at Russell or Kiattut Sermiat glaciers. Keeling plots of $[\text{CO}_{2\text{-total}}]^{-1}$ versus $\delta^{13}\text{C-CO}_2$ indicate that CO_{2-total} may be represented by a two-end member mixing model for sub-Isunnguata discharge, but not for discharge from the Russell and Kiattut Sermiat glaciers (Fig. 9). Mixing model end members include a ¹³C-enriched, lower concentration CO₂ source and a ¹³C-depleted, higher concentration CO₂ source. The y-intercept of the regression between $[\text{CO}_{2\text{-total}}]^{-1}$ versus $\delta^{13}\text{C-CO}_2$, which represents the isotopic signature of the high-CO₂ endmember, is -27.4‰. This value is close to what would be expected from OM remineralization, as indicated by remineralized OM in Greenlandic heath soils that produced $\delta^{13}\text{C-CO}_2$ of approximately -27 to -25‰ (Ravn et al., 2020), and thawed Alaskan permafrost soils that produced $\delta^{13}\text{C-CO}_2$ of between -20 to -30‰ for (Mauritz et al., 2019), both of which may be similar to subglacial organic matter. The low-CO₂ end member could reflect atmospheric CO₂ input, with an associated $\delta^{13}\text{C-CO}_2$ value of approximately -8‰. While the $\delta^{13}\text{C-CO}_2$ value of the lowest-CO_{2-total} samples in the sub-Isunnguata Keeling plot (e.g. highest $[\text{CO}_{2\text{-total}}]^{-1}$ not including the outlier) is slightly depleted compared to atmospheric values at -12.1‰; however, even the lowest CO₂ concentrations measured at sub-Isunnguata are supersaturated with respect to atmospheric concentrations (Fig. 3g). Thereby suggesting that OM remineralization contributes CO₂ even for low CO₂-concentration samples and isotopically depletes the subglacial CO₂ reservoir.

775 While $\delta^{13}\text{C-CO}_2$ values of Russell and Kiattut Sermiat samples are within the range of sub-Isunnguata samples, suggesting possible contributions of CO_{2-atm} and CO_{2-OM}, scatter in the Keeling plots indicate variability in the CO₂ concentration and/or isotopic composition of end members, or significant contributions of at least one other major subglacial CO₂ source. We address both possibilities here. While atmospheric CO₂ concentrations and $\delta^{13}\text{C}$ values should be relatively invariable, CO_{2-OM} may vary both in concentration and isotopic composition, depending on variability in the quantity and composition of organic deposits, as well as remineralization rates. For instance, if remineralization largely occurs in hydrologically isolated subglacial meltwater pockets, some variability in the concentration and $\delta^{13}\text{C-CO}_2$ of CO_{2-OM} is likely. While no data yet exist to characterize the variability in subglacial OM reservoirs, variability in either concentration or isotopic

Deleted: its

Deleted: as

Deleted: as well as the

Deleted: of that process

Deleted: predominant

Deleted: at Isunnguata discharge

Deleted: , in contrast to

Deleted: 8

Deleted: -

Deleted: -

Deleted: (Fig. 8a).

Deleted: (representing

Deleted:)

Deleted: ‰, which

Deleted: . For instance, CO₂ from

Deleted: ranged between

Deleted: between -20 to -30‰ for

Deleted: An additional correlation is observed between CO_{2-total} and NH₄ concentrations for Isunnguata samples, which would be expected from OM remineralization (Fig. 8b).

Deleted: which should result in a

Deleted: are

Deleted: ‰ (Fig. 8a).

Formatted: Font color: Text 1

Deleted: Supersaturation suggests

Deleted: While both atmospheric CO₂ (CO_{2-atm}) and CO₂ derived from OM remineralization (CO_{2-OM}) provide carbonic acid to drive subglacial mineral weathering as well as CO₂ supersaturation in Isunnguata discharge, their absolute and relative contributions are controlled by different processes. Understanding the controls of CO₂ acquisition may improve understanding of subglacial carbon dynamics as well as the conditions necessary for subglacial environments to become CO₂ sources to the atmosphere. CO_{2-OM} is the dominant source to CO_{2-total} at Isunnguata in the early and late melt seasons (Fig. 9a) when discharge is low (Fig. 9c), while CO_{2-atm} is the dominant CO₂ source in the peak melt season when discharge is high. This switch in dominant CO₂ sources occurs because the magnitude of CO_{2-OM} has a strong negative association with discharge, approaching zero during maximum discharge times (Fig. 9d), while CO_{2-atm} remains relatively constant over the melt season (Fig. 9b) and the range of discharges (Fig. 9d). The high contributions of CO_{2-OM} during low discharge could reflect higher residence times that allow greater biogeochemical modification and accumulation of OM remineralization reaction products, including CO₂. The chemostatic behavior for CO_{2-atm} indicates that invasion of atmospheric CO₂ is independent of the extent of chemical ... [7]

Deleted: non-linear

Deleted: ,

870 composition of CO_{2-OM} could plausibly result in the scatter shown in Figure 9. Additional subglacial CO₂ sources from ice
bubbles, or lithogenic CO₂ liberated by mechanical grinding would be expected to enrich rather than deplete the δ¹³C-CO₂
875 values of the samples relative to modern atmospheric δ¹³C-CO₂ values. Ice bubbles contain gaseous CO₂ at concentrations and
isotopic compositions reflecting atmospheric conditions during ice formation. While heterogeneity may result from gas
bubbles recording changes in atmospheric CO₂, variability in δ¹³C-CO₂ of gas bubble CO₂ should be only a few per mil, which
is small compared to the variation observed in Russell and Kiattut Sermiat samples (e.g., Tipple et al., 2010; Fig. 9). Gas
bubble CO₂ should also be ¹³C-enriched compared to modern atmospheric CO₂ due to fossil fuel contributions, and thus would
be unlikely to cause the observed depletion of ¹³C in the subglacial water.

Deleted: non-linear Keeling plots here.
Deleted: could include atmospheric CO₂ contained in
Deleted: , though both of these sources

Recent work has also highlighted the potential for subglacial mechanical grinding to result in CO₂ production through
the volatilization of bedrock fluid inclusions (Macdonald et al., 2018). While this process was found to produce sufficient CO₂
to drive approximately 20% of mineral weathering in Svalbard subglacial environments, the expected isotopic composition of
880 lithogenic CO₂ is more ¹³C-enriched than our measured δ¹³C-CO₂ values. For instance, estimates of δ¹³C for bulk hydrocarbons
in fluid inclusions in the Ilmaussaq alkaline complex of South Greenland have values of -4.5±1.5‰ (Madsen, 2001), which
is close to the δ¹³C-CO₂ of CO₂ in fluid inclusions in the Bamble granulite sector of South Norway (~ -6‰; Newton et al.,
1980). Because mechanical grinding should not fractionate the δ¹³C-CO₂ values (Lüders et al., 2012), our low δ¹³C-CO₂ values
suggest this source is small relative to other sources.

Deleted: (Tipple et al., 2010; Fig. 8a).
Deleted: variation
Deleted: sample δ¹³C-CO₂ values that are more ¹³C-depleted than
modern atmospheric δ¹³C-CO₂ values (Fig. 8a).

885 One additional source or sink of CO₂ to some of our samples is atmospheric exchange as water flows from the
subglacial outlet site to our sampling sites. However, atmospheric CO₂ exchange after discharge would have the same impact
on Keeling plots as atmospheric CO₂ exchange prior to discharge. Incorporation of an atmospheric source between the outlet
and sample site would be most likely at Kiattut Sermiat where CO₂ concentrations are undersaturated with respect to
atmospheric concentrations, which would promote invasion of atmospheric CO₂. However, the measured δ¹³C-CO₂ values are
890 more ¹³C-depleted than modern atmospheric CO₂ and are not consistent with atmospheric CO₂ as the sole or dominant source
of CO₂ to these samples (Fig. 9).

Moved (insertion) [6]

Deleted: While volatilization of fluid inclusions through
mechanical grinding was found to produce sufficient CO₂ to drive
approximately 20% of mineral weathering in Svalbard subglacial
environments, the expected isotopic composition of lithogenic CO₂
is more ¹³C-enriched than our measured δ¹³C-CO₂ values. Because
mechanical grinding should produce CO₂ with an isotopic
composition reflecting the lithogenic source,

Deleted: its contributions here are likely limited

Moved up [6]: For instance, estimates of δ¹³C for bulk
hydrocarbons in fluid inclusions in the Ilmaussaq alkaline complex
of South Greenland have values of -4.5±1.5‰ (Madsen, 2001),
which is close to the δ¹³C-CO₂ of CO₂ in fluid inclusions in the
Bamble granulite sector of South Norway (~ -6‰; Newton et al.,
1980). There is an additional possibility of atmospheric exchange

Deleted: There is an additional possibility of atmospheric
exchange between the subglacial outlet site and our water sampling
sites that could serve as an additional CO₂ source or sink.

Deleted: Although

Deleted: and

Deleted: ,

Deleted: glacial meltwater

Deleted: 8a).

Deleted: the

895 While more information is needed to determine all possible sources of CO₂ to Russell and Kiattut Sermiat samples,
δ¹³C-CO₂ values of samples from both sites imply mixing between a ¹³C-depleted CO₂ source, such as OM remineralization,
and one or more ¹³C-enriched CO₂ sources, such as atmospheric or lithogenic CO₂. Similar to CH₄, concentrations and isotopic
compositions of gases may be impacted by atmospheric exchange between the glacial outlet and our sampling locations, which
would alter dissolved CO₂ concentrations and δ¹³C-CO₂ compositions to values more similar to atmospheric values. Therefore,
we are unable to distinguish the impacts of atmospheric exchange that occurs prior to discharge from exchange that occurs
between discharge and our sampling locations. However, the impact of this exchange should be relatively constant between
sampling times and sampling locations, therefore outgassing would not account for temporal or spatial variability in CO₂
900 concentrations or isotopic compositions between sites.

4.2.2 Subglacial CO₂ sink: mineral weathering reactions

Mineral weathering leads to net CO₂ consumption in all subglacial discharge samples (Fig. 6), and thus the measured CO₂ concentrations at glacial outlets represent only a fraction of the total CO₂ that would have been present in the absence of mineral weathering reactions (CO_{2-total}; Eq. 10). Net consumption occurs because the CO₂ source from Carbs_A is ubiquitously low compared to sinks from either Carb_{CA} or Sil_{CA} (Fig. 6b). The range in Net CO_{2-MW} is similar between subglacial discharge sites (between 10-150 μM; Fig. 6a), but average values increase from Kiattut Sermiat to Russell to sub-Isunnguata, likely reflecting the relative weatherability of alkaline igneous rocks, granulite facies gneisses, and amphibolite facies gneisses. Kiattut Sermiat is characterized by a relatively high proportion of Carb_{CA} compared to sub-Isunnguata and Russell sites, which may arise from the presence of trace carbonates in abundant and readily weatherable basaltic intrusions as has been implicated in other studies (Urta et al., 2019). The relatively greater influence of carbonate dissolution compared to silicate dissolution on Total CO_{2-MW} at Kiattut Sermiat may also relate to more rapid dissolution kinetics of carbonates, which allow carbonate dissolution to have a large influence on major cation and anion load even when carbonates are only present in trace amounts (Deuerling et al., 2019; Tranter, 2005). At sub-Isunnguata and Russell sites, Sil_{CA} has a greater influence than Carb_{CA} on Total CO_{2-MW}, which could result from either a lower abundance of trace carbonates to participate in weathering reactions, or relatively longer subglacial residence times that would allow a greater accumulation of silicate weathering products.

Despite the impact of Carb_{CA} on Total CO_{2-MW} at Kiattut Sermiat compared to sub-Isunnguata and Russell sites, Carbs_A is notably lower at Kiattut Sermiat than other sites and suggests a limited role for sulfuric acid weathering that may relate to subglacial sulfide oxidation dynamics. Lower abundances of sulfide minerals in the subglacial environment may limit the production of sulfuric acid, and could result from differences in lithology between sites, the depletion of sulfide minerals due to prior weathering (Graly et al., 2014), or weathering occurring in anoxic environments that limit the oxidation of sulfide to sulfuric acid (Deuerling et al., 2019). The kinetics of sulfide oxidation may also significantly differ between sites depending on the relative contributions of abiotic compared to microbially mediated sulfide oxidation, as microbially mediated sulfide oxidation is several orders of magnitude faster than abiotic sulfide oxidation. Rapid microbially mediated sulfide oxidation has been implicated in the development of anaerobic conditions, which could also support subglacial methanogenesis (Sharp et al., 1999). Observations of higher CH₄ concentrations as well as higher contributions of Carbs_A at sub-Isunnguata and Russell compared to Kiattut Sermiat, may therefore be linked to subglacial microbial activity, which is known to vary based on factors such as the presence of organic and fine-grained rock flour to serve as growth substrates, insulation from fluctuations in temperature, and delivery of nutrients and organic matter from supraglacial sources (Sharp et al., 1999). If microbially driven, our results suggest possible linkages between microbial processes and subglacial mineral weathering regimes, with significant impacts to both CH₄ and CO₂ dynamics due to the role of Carbs_A as a CO₂ source (Table 1).

Whether controlled by geochemical, microbial, or mechanical processes, the relationships between CO₂ concentrations and δ¹³C-CO₂ and average daily discharge are similar between sub-Isunnguata and Russell sites (Fig. 8). These similarities suggest that underlying controls in carbonate chemistry may be consistent between sites despite the heterogeneity

Moved down [7]: 5 Conclusions†

Deleted: impact the concentrations of CO₂ and CH₄

Moved (insertion) [2]

Moved (insertion) [3]

Moved (insertion) [4]

Deleted: of GrIS, which act as either sources or sinks of GHG to the atmosphere.

Moved (insertion) [5]

Deleted: of subglacial discharge are likely

Deleted: the availability of subglacial OM to drive methanogenesis as well as the extent of CH₄ oxidation. Regional differences in subglacial OM deposits may account for the occurrence of methanogenesis in southwest outlet glaciers (Isunnguata and Russell) in this and other studies, contrasting with southern Kiattut Sermiat where little CH₄ production occurs. During the early melt season, oxidation consumes nearly 50% of CH₄ produced at southwest sites, and

Deleted: discharge and CH₄ oxidation (ϵ_c and ϵ_{oc}) suggest that CH₄ oxidation depends on longer subglacial residence time during periods of low discharge. While mineral weathering consumes CO₂ throughout the melt season at all three sites, additional CO₂ respired from atmospheric and subglacial sources increases the CO₂ ...

Deleted: of subglacial

Deleted: . The magnitude of additional

985 in measured dissolved CO₂ concentrations. For both sub-Isunnguata and Russell sites, mineral weathering reactions consume CO₂, which implies contributions from *in situ* CO₂ sources (such as atmospheric CO₂ invasion or OM remineralization) to produce measured CO₂ concentrations. The different CO₂ concentrations observed between sites therefore appear to result from the strength of *in situ* CO₂ sources relative to CO_{2-MW}, both of which impart the greatest chemical change at times of low discharge and high subglacial residence time. At Kiattut Sermiat, where measured CO₂ concentrations are lowest, the magnitude of *in situ* sources is insufficient to maintain atmospheric equilibrium, leading subglacial discharge to be a sink of atmospheric CO₂, while CO_{2-total} maintains close to atmospheric equilibrium concentrations at the Russell Glacier. At sub-Isunnguata, however, OM remineralization produces more CO₂ than is consumed by mineral weathering and causes meltwater to be a source of CO₂ to the atmosphere. This finding implies that subglacial mineral weathering serves to partially or fully consume CO₂ produced from *in situ* sources under the Greenland Ice Sheet but does not always serve to directly consume modern atmospheric CO₂.

Deleted: (CO_{2-total})

Deleted: at Kiattut Sermiat

Deleted: GrIS

Deleted: necessarily result in direct consumption of

5 Conclusions

Moved (insertion) [7]

Deleted: The important

990 Subglacial reactions impact the concentrations of CO₂ and CH₄ in subglacial discharge of Greenland Ice Sheet, and differences in the relative magnitudes of microbial and geochemical processes result in a high degree of previously unrecognized heterogeneity between glacial discharge sites of the Greenland Ice Sheet. Our results imply a significant role of OM remineralization in driving this heterogeneity and leading to CO₂ and CH₄ supersaturation at the sub-Isunnguata site, and CH₄ supersaturation at the Russell site. Heterogeneity may result in significant uncertainty in total greenhouse gas flux estimates from subglacial systems of the Greenland Ice Sheet, which will be an increasingly important carbon flux as the Arctic warms in the coming decades. While heterogeneous, the uncertainty in greenhouse gas fluxes from Greenland Ice Sheet meltwater may be reduced by a better understanding of the controls and variability of the weathering reactions and microbial processes driving heterogeneous gas concentrations. Such a process-based understanding could also improve estimates of the impact of greenhouse gas variability associated with the growth and retreat of continental ice sheets on over glacial-interglacial timescales. Subglacial OM remineralization further implies the existence of links between subglacial OM deposits and export of other biogeochemical solutes from the Greenland Ice Sheet, including nutrients as well as redox-sensitive elements. While the export of nutrients from the Greenland Ice Sheet has been the focus of numerous studies (Bhatia et al., 2013; Hawkings et al., 2016; Lawson et al., 2014), little is currently known regarding the role of OM sources in governing these exports. Given the variability in greenhouse gas concentrations observed in this study, constraining the extent of heterogeneity in outlet glaciers of the Greenland Ice Sheet as well as the biogeochemical, hydrologic, and geologic controls of this heterogeneity will be important for upscaling atmospheric fluxes as well as efforts to predict impacts of ice loss on carbon and nutrient budgets due to current and future melting of the Greenland Ice Sheet.

Deleted: subglacial environments of the GrIS determined by

Deleted: and other studies

Deleted: implies

Deleted: GrIS

Deleted: GrIS

Deleted: GHG

Deleted: GrIS

Deleted: GrIS

Author Contribution

Jonathan B. Martin and Ellen E. Martin participated in conceptualization, data collection, data interpretation, reviewing and editing the manuscript, and acquired the funding for this project. [Asa Rennermalm provided the discharge data that was used in analyses of discharge-concentration relationships](#). Shaily Rahman participated in data collection and early interpretation and presentation of results. Andrea J. Pain participated in conceptualization, data collection, analysis, and interpretation, and took the lead on writing the manuscript with contributions from Jonathan B. Martin, [Ellen E. Martin](#), and [Asa Rennermalm](#).

Deleted: and

Acknowledgements

We acknowledge the members of our field teams: Daniel Fischer, Fabio Da Prat, Hailey Hall, Mark Robbins, [Scott Schnur](#), and [Michelle D. Kisbye](#). Additional invaluable support was provided by Steven DiEgidio, Nini Frydkjær Brandt, Inga Gisladottir, and Jacky Simoud. We are grateful for the excellent field support provided by the Kangerlussuaq International Science Station and Polar Field Services (CH2M Hill). This work was funded by the National Science Foundation grant (ANS-1603452). The authors declare that they have no conflict of interest. Data is accessible on the Arctic Data Center (<https://cn.dataone.org/cn/v2/resolve/urn:uuid:c1051a07-cbdf-4061-ae44-c1472f61e3fe>) and major element concentrations used for geochemical modeling (<https://cn.dataone.org/cn/v2/resolve/urn:uuid:65d272f6-d280-4fcc-8aaa-4805f12ca6ae>). We are grateful to [Marek Stibal](#) and an anonymous reviewer who made suggestions that greatly improved the manuscript, as well as [Joseph Graly](#) for assistance improving the accuracy of presented geographical locations.

Deleted: and

Deleted: Discharge data from Watson River were gathered by B. Hasholt and A.B. Mikkelsen on behalf of the University of Copenhagen (2006-2013), and by D. van As and B. Hasholt on behalf of the Geological Survey of Denmark and Greenland (2014-present). ...

Deleted: (<https://cn.dataone.org/cn/v2/resolve/urn:uuid:65d272f6-d280-4fcc-8aaa-4805f12ca6ae>).

References

Andrews, L. C., Catania, G. A., Hoffman, M. J., Gulley, J. D., Lüthi, M. P., Ryser, C., Hawley, R. L. and Neumann, T. A.: Direct observations of evolving subglacial drainage beneath the Greenland Ice Sheet, *Nature*, 514(7520), 80–83, doi:10.1038/nature13796, 2015.

Anklin, M., Barnola, J. M., Schwander, J., Stauffer, B. and Raynaud, D.: Processes affecting the CO₂ concentrations measured in Greenland ice, *Tellus B*, 47(4), 461–470, doi:10.1034/j.1600-0889.47.issue4.6.x, 1995.

Deleted: . M

van As, D., Hasholt, B., Ahlstrøm, A. P., Box, J. E., Cappelen, J., Colgan, W., Fausto, R. S., Mernild, S. H., Bech, A., Noël, B. P. Y., Petersen, D. and van Den Broeke, M. R.: Reconstructing Greenland Ice Sheet meltwater discharge through the Watson River (1949–2017), *Arctic, Antarct. Alp. Res.*, 50(1), doi:10.1080/15230430.2018.1433799, 2018.

[Van](#) As, D., Mikkelsen, A. B., Nielsen, M. H., Box, J. E., Liljedahl, L. C., Lindbäck, K., Pitcher, L. and Hasholt, B.: Hypsometric amplification and routing moderation of Greenland ice sheet meltwater release, *Cryosphere*, 11(3), 1371–1386, doi:10.5194/tc-11-1371-2017, 2017.

Deleted: van

- As, D. Van, Hasholt, B., Ahlström, A. P., Box, J. E., Cappelen, J., Colgan, W., Fausto, R. S., Mernild, S. H., Bech, A., Noël, B. P. Y., Petersen, D., Broeke, M. R. Van Den, As, D. Van, Hasholt, B., Ahlström, A. P., Box, J. E., Colgan, W., Fausto, R. S., Mernild, S. H. and Mikkelsen, A. B.: Reconstructing Greenland Ice Sheet meltwater discharge through the Watson River (1949 – 2017) Reconstructing Greenland Ice Sheet meltwater discharge through the Watson, Arctic, *Antarct. Alp. Res.*, 50(1), doi:10.1080/15230430.2018.1433799, 2018.
- Berner, R. A., Lasaga, A. C. and Garrels, R. M.: The carbonate-silicate geochemical cycle and its effect on atmospheric carbon dioxide over the past 100 million years., *Am. J. Sci.*, 283(7), 641–683, doi:10.2475/ajs.283.7.641, 1983.
- 1075 Bhatia, M. P., Kujawinski, E. B., Das, S. B., Breier, C. F., Henderson, P. B. and Charette, M. A.: Greenland meltwater as a significant and potentially bioavailable source of iron to the ocean, *Nat. Geosci.*, 6(4), 274–278, doi:10.1038/ngeo1746, 2013.
- Christiansen, J. R. and Jørgensen, C. J.: First observation of direct methane emission to the atmosphere from the subglacial domain of the Greenland Ice Sheet, *Sci. Rep.*, 8(1), 2–7, doi:10.1038/s41598-018-35054-7, 2018.
- Cowton, T., Nienow, P., Sole, A., Wadham, J., Lis, G., Bartholomew, I., Mair, D. and Chandler, D.: Evolution of drainage system morphology at a land - terminating Greenlandic outlet glacier, , 118, 29–41, doi:10.1029/2012JF002540, 2013.
- 1080 Deuerling, K. M., Martin, J. B., Martin, E. E. and Scribner, C. A.: Hydrologic exchange and chemical weathering in a proglacial watershed near Kangerlussuaq, west Greenland, *J. Hydrol.*, 556, 220–232, doi:10.1016/j.jhydrol.2017.11.002, 2018.
- Deuerling, K. M., Martin, J. B., Martin, E. E., Abermann, J., Myreng, S. M., Petersen, D. and Rennermalm, A. K.: Chemical weathering across the western foreland of the Greenland Ice Sheet, *Geochim. Cosmochim. Acta*, 245(245), 426–440, doi:10.1016/j.gca.2018.11.025, 2019.
- 1085 Dieser, M., Broemsen, E. L. J. E., Cameron, K. A., King, G. M., Achberger, A., Choquette, K., Hagedorn, B., Sletten, R., Junge, K. and Christner, B. C.: Molecular and biogeochemical evidence for methane cycling beneath the western margin of the Greenland Ice Sheet, *ISME J.*, 8(11), 2305–2316, doi:10.1038/ismej.2014.59, 2014.
- van Gool, J. A., Alsop, G. I., Arting, U. E., Garde, A. A., Knudsen, C., Krawiec, A. W., Mazur, S., Nygaard, J., Piazzolo, S. and Thomas, C. W.: Precambrian geology of the northern Nagssugtoqidian orogen, West Greenland: mapping in the Kangaatsiaq area, *Geol. Greenl. Surv. Bull.*, 191, 13–23, 2002.
- 1090 Graly, J., Harrington, J. and Humphrey, N.: Combined diurnal variations of discharge and hydrochemistry of the Isunnguata Sermia outlet, Greenland Ice Sheet, *Cryosphere*, 11(3), 1131–1140, doi:10.5194/tc-11-1131-2017, 2017a.
- Graly, J. A., Humphrey, N. F., Landowski, C. M. and Harper, J. T.: Chemical weathering under the Greenland Ice Sheet, *Geology*, 551–554, doi:10.1130/G35370.1, 2014.
- 1095 Graly, J. A., Drever, J. I. and Humphrey, N. F.: Calculating the balance between atmospheric CO₂ drawdown and organic carbon oxidation in subglacial hydrochemical systems, *Global Biogeochem. Cycles*, 31(4), 709–727,

Deleted: van

Deleted: .,

1100 doi:10.1002/2016GB005425, 2017b.

Hasholt, B., As, D. Van, Mikkelsen, A. B., Mernild, S. H., Yde, J. C., Hasholt, B., As, D. Van, Mikkelsen, A. B., Mernild, S. H. and Hasholt, B.: Observed sediment and solute transport from the Kangerlussuaq sector of the Greenland Ice Sheet (2006 – 2016), *Arctic, Antarct. Alp. Res.*, 50(1), doi:10.1080/15230430.2018.1433789, 2018.

1105 Hawkins, J., Wadham, J., Tranter, M., Telling, J. and Bagshaw, E.: The Greenland Ice Sheet as a hot spot of phosphorus weathering and export in the Arctic, *Global Biogeochem. Cycles*, 191–210, doi:10.1002/2015GB005237. Received, 2016.

Henriksen, N., Higgins, A. K., Kalsbeek, F. and Pulvertaft, T. C. R.: Greenland from Archaean to Quaternary. Descriptive text to the 1995 Geological map of Greenland, 1:2 500 000. 2nd edition, *GEUS Bull.*, 18(SE-MONOGRAPH), 1–126, doi:10.34194/geusb.v18.4993, 2009.

1110 [Herschy, R.: Hydrometry, 2nd Editio., John Wiley & Sons, Ltd., Chichester, New York, Winheim, Brisbane, Singapore, Toronto., 1999.](#)

Hindshaw, R. S., Rickli, J., Leuthold, J., Wadham, J. and Bourdon, B.: Identifying weathering sources and processes in an outlet glacier of the Greenland Ice Sheet using Ca and Sr isotope ratios, *Geochim. Cosmochim. Acta*, 145, 50–71, doi:10.1016/j.gca.2014.09.016, 2014.

1115 [ISO Guide 98-3 \(2008\). Guide 98-3. Uncertainty of measurement - Part 3: Guide to the expression of uncertainty in measurement \(GUM:1995\), Guid. 98-3, \(148902\), 130, doi:10.1373/clinchem.2003.030528.](#)

Kalsbeek, F. and Taylor, P. N.: Isotopic and chemical variation in granites across a Proterozoic continental margin- the Ketilidian mobile belt of South Greenland, *Earth Planet. Sci. Lett.*, 73(1), 65–80, 1985.

Keeling, C. D.: The concentration and isotopic abundances of atmospheric carbon dioxide in rural areas, *Geochim. Cosmochim. Acta*, 13(4), 322–334, doi:10.1016/0016-7037(58)90033-4, 1958.

1120 Lamarche-Gagnon, G., Wadham, J. ., Sherwood Lollar, B., Arndt, S., Fietzek, P., Beaton, A. D., Tedstone, A. J., Telling, J., Bagshaw, E. A., Hawkins, J. R., Kohler, T. J., Zarsky, J. D., Mowlem, M. C., Anesio, A. M. and Stibal, M.: Greenland melt drives continuous export of methane from the ice-sheet bed, *Nature*, 565(7737), 73–77, doi:http://dx.doi.org/10.1038/s41586-018-0800-0, 2019.

1125 Larsen, N. K., Find, J., Kristensen, A., Bjørk, A. A., Kjeldsen, K. K., Odgaard, B. V., Olsen, J. and Kjær, K. H.: Holocene ice marginal fluctuations of the Qassimiut lobe in South Greenland, *Nat. Publ. Gr.*, 1–11, doi:10.1038/srep22362, 2016.

Lawson, E. C., Wadham, J. L., Tranter, M., Stibal, M., Lis, G. P., Butler, C. E. H., Laybourn-Parry, J., Nienow, P., Chandler, D. and Dewsbury, P.: Greenland ice sheet exports labile organic carbon to the arctic oceans, *Biogeosciences*, 11(14), 4015–4028, doi:10.5194/bg-11-4015-2014, 2014.

Lindbäck, K., Pettersson, R., Hubbard, A. L., Doyle, S. H., As, D., Mikkelsen, A. B. and Fitzpatrick, A. A.: Subglacial water

1130 drainage, storage, and piracy beneath the Greenland ice sheet, , 7606–7614, doi:10.1002/2015GL065393.Received, 2015.

Lüders, V., Plessen, B. and di Primio, R.: Stable carbon isotopic ratios of CH₄-CO₂-bearing fluid inclusions in fracture-fill mineralization from the Lower Saxony Basin (Germany) - A tool for tracing gas sources and maturity, *Mar. Pet. Geol.*, 30(1), 174–183, doi:10.1016/j.marpetgeo.2011.10.006, 2012.

1135 Ludwig, W., Amiotte Suchet, P. and Probst, J.-L.: Enhanced chemical weathering of rocks during the last glacial maximum: a sink for atmospheric CO₂?, *Chem. Geol.*, 159, 147–161 [online] Available from: <http://www.sciencedirect.com/science/article/pii/S0009254199000388%5Cnpapers2://publication/uuid/9F3040A1-7901-480E-B435-BA665074AC32>, 1999.

Deleted: Geol., 159, 147–161

1140 Macdonald, M. L., Wadham, J. L., Telling, J. and Skidmore, M. L.: Glacial Erosion Liberates Lithologic Energy Sources for Microbes and Acidity for Chemical Weathering Beneath Glaciers and Ice Sheets, *Front. Earth Sci.*, 6(November), 1–15, doi:10.3389/feart.2018.00212, 2018.

Madsen, J. K.: A review of the composition and evolution of hydrocarbon gases during solidification of the Ilímaussaq alkaline complex, South Greenland, *Geol. Greenl. Surv. Bull.*, 190, 159–166, 2001.

Mahieu, K., De Visscher, A., Vanrolleghem, P. A. and Van Cleemput, O.: Modelling of stable isotope fractionation by methane oxidation and diffusion in landfill cover soils, *Waste Manag.*, 28(9), 1535–1542, doi:10.1016/j.wasman.2007.06.003, 2008.

1145 Mauritz, M., Celis, G., Ebert, C., Hutchings, J., Ledman, J., Natali, S. M., Pegoraro, E., Salmon, V. G., Schädel, C., Taylor, M. and Schuur, E. A. G.: Using Stable Carbon Isotopes of Seasonal Ecosystem Respiration to Determine Permafrost Carbon Loss, *J. Geophys. Res. Biogeosciences*, 124(1), 46–60, doi:10.1029/2018JG004619, 2019.

1150 Mouginot, J., Rignot, E., Bjørk, A. A., van den Broeke, M., Millan, R., Morlighem, M., Noël, B., Scheuchl, B. and Wood, M.: Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, *Proc. Natl. Acad. Sci.*, 116(19), 201904242, doi:10.1073/pnas.1904242116, 2019.

Musilova, M., Tranter, M., Wadham, J., Telling, J., Tedstone, A. and Anesio, A. M.: Microbially driven export of labile organic carbon from the Greenland ice sheet, *Nat. Geosci.*, 10(5), 360–365, doi:10.1038/ngeo2920, 2017.

1155 Myrntinen, A., Becker, V. and Barth, J. A. C.: A review of methods used for equilibrium isotope fractionation investigations between dissolved inorganic carbon and CO₂, *Earth-Science Rev.*, 115(3), 192–199, doi:10.1016/j.earscirev.2012.08.004, 2012.

Newton, R. C., Smith, J. V. and Windley, B. F.: Carbonic metamorphism, granulites and crustal growth, *Nature*, 288(5786), 45–50, doi:10.1038/288045a0, 1980.

Pain, A. J., Martin, J. B. and Young, C. R.: Sources and sinks of CO₂ and CH₄ in siliciclastic subterranean estuaries, *Limnol. Oceanogr.*, 64(4), 1500–1514, doi:10.1002/lno.11131, 2019.

Parkhurst, L.: Geochemical mole-balance modeling with uncertain data, *Water Resour. Res.*, 33(8), 1957–1970, 1997.

Da Prat, F. A. and Martin, E.: Weathering in the Glacial Foreland of Southern and Western Greenland, *Univ. Florida J. Undergrad. Res.*, 20(2), doi:10.32473/ufjur.v20i2.106168, 2019.

1165 Preuss, I., Knoblauch, C., Gebert, J. and Pfeiffer, E. M.: Improved quantification of microbial CH₄ oxidation efficiency in arctic wetland soils using carbon isotope fractionation, *Biogeosciences*, 10(4), 2539–2552, doi:10.5194/bg-10-2539-2013, 2013.

Ravn, N. R., Elberling, B. and Michelsen, A.: Arctic soil carbon turnover controlled by experimental snow addition, summer warming and shrub removal, *Soil Biol. Biochem.*, 142(December 2019), 107698, doi:10.1016/j.soilbio.2019.107698, 2020.

1170 Rennermalm, A. K., Smith, L. C., Chu, V. W., Forster, R. R., Box, J. E. and Hagedorn, B.: Proglacial river stage, discharge, and temperature datasets from the Akuliarusiarsuup Kuua River northern tributary, Southwest Greenland, 2008–2011, *Earth Syst. Sci. Data Discuss.*, 2100(4), 1–12, doi:10.5194/essd-4-1-2012, 2012.

Rennermalm, A. K., Smith, L. C., Chu, V. W., Box, J. E., Forster, R. R., Broeke, M. R. Van Den and As, D. Van: Evidence of meltwater retention within the Greenland ice sheet, *Cryosph.*, 1433–1445, doi:10.5194/tc-7-1433-2013, 2013.

1175 Repo, M. E., Huttunen, J. T., Naumov, A. V., Chichulin, A. V., Lapshina, E. D., Bleuten, W. and Martikainen, P. J.: Release of CO₂ and CH₄ from small wetland lakes in western Siberia, *Tellus, Ser. B Chem. Phys. Meteorol.*, 59(5), 788–796, doi:10.1111/j.1600-0889.2007.00301.x, 2007.

Ruskeeniemi, T., Engström, J., Lehtimäki, J., Vanhala, H., Korhonen, K., Kontula, A., Claesson Liljedahl, L., Näslund, J. O. and Pettersson, R.: Subglacial permafrost evidencing re-advance of the Greenland Ice Sheet over frozen ground, *Quat. Sci. Rev.*, 199, 174–187, doi:10.1016/j.quascirev.2018.09.002, 2018.

1180 Sharp, M., Parkes, J., Cragg, B., Fairchild, I. J., Lamb, H. and Tranter, M.: Widespread bacterial populations at glacier beds and their relationship to rock weathering and carbon cycling, *Geology*, 27(2), 107–110, doi:10.1130/0091-7613(1999)027<0107:WBPAGB>2.3.CO;2, 1999.

[Solinst: Solinst Technical Bulletin: Understanding pressure sensor drift, 2017.](#)

1185 [St Pierre, K. A., St Louis, V. L., Schiff, S. L., Lehnerr, I., Dainard, P. G., Gardner, A. S., Aukes, P. J. K. and Sharp, M. J.: Proglacial freshwaters are significant and previously unrecognized sinks of atmospheric CO₂, *Proc. Natl. Acad. Sci.*, 116\(36\), 17690–17695, doi:10.1073/pnas.1904241116, 2019.](#)

Tipple, B. J., Meyers, S. R. and Pagani, M.: Carbon isotope ratio of Cenozoic CO₂: A comparative evaluation of available geochemical proxies, *Paleoceanography*, 25(3), 1–11, doi:10.1029/2009pa001851, 2010.

1190 Tranter, M.: Geochemical Weathering in Glacial and Proglacial Environments, in *Surface and Ground Water, Weathering, and Soils: Treatise on Geochemistry*, vol. 5, edited by J. I. Drever., 2005.

Deleted: St. Pierre, K. A., St.

Moved down [8]: Louis, V. L., Schiff, S. L., Lehnerr, I., Dainard, P. G., Gardner, A. S., Aukes, P. J. K. and Sharp, M. J.: Proglacial freshwaters are significant and previously unrecognized sinks of atmospheric CO₂, *Proc. Natl. Acad. Sci.*, 116(36), 17690–17695, doi:10.1073/pnas.1904241116, 2019.

Deleted: Ryu, J. S. and Jacobson, A. D.: CO₂ evasion from the Greenland Ice Sheet: A new carbon-climate feedback, *Chem. Geol.*, 320–321, 80–95, doi:10.1016/j.chemgeo.2012.05.024, 2012.

Moved (insertion) [8]

- 1200 Upton, B. G. J., Emeleus, C. H., Heaman, L. M., Goodenough, K. M. and Finch, A. A.: Magmatism of the mid-Proterozoic Gardar Province, South Greenland: Chronology, petrogenesis and geological setting, *Lithos*, 68(1–2), 43–65, doi:10.1016/S0024-4937(03)00030-6, 2003.
- Urta, A., Wadham, J., Hawkings, J. R., Telling, J., Hatton, J. E., Yde, J. C., Hasholt, B., van As, D., Bhatia, M. P. and Nienow, P.: Weathering Dynamics Under Contrasting Greenland Ice Sheet Catchments, *Front. Earth Sci.*, 7(November), 1205 doi:10.3389/feart.2019.00299, 2019.
- de Visscher, A., de Pourcq, I. and Chanton, J.: Isotope fractionation effects by diffusion and methane oxidation in landfill cover soils, *J. Geophys. Res. Atmos.*, 109(18), 1–8, doi:10.1029/2004JD004857, 2004.
- Wadham, J. L., Tranter, M., Tulaczyk, S. and Sharp, M.: Subglacial methanogenesis : A potential climatic amplifier ?, *Global Biogeochem. Cycles*, 22(December 2007), 1–16, doi:10.1029/2007GB002951, 2008.
- 1210 Walker, J. C. G., Hays, P. B. and Kasting, J. F.: A negative feedback mechanism for the long-term stabilization of earth's surface temperature, *J. Geophys. Res.*, 86(1), 9776–9782, 1981.
- Warren, C. R. and Glasser, N. F.: Contrasting response of south Greenland glaciers to recent climatic change, *Arct. Alp. Res.*, 24(2), 124–132, doi:10.2307/1551532, 1992.
- Whiticar, M. J.: Carbon and hydrogen isotope systematics of bacterial formation and oxidation of methane, *Chem. Geol.*, 161, 1215 291–314, 1999.
- Whiticar, M. J. and Schoell, M.: Biogenic methane formation in marine and freshwater environments: CO₂ reduction vs. acetate fermentation- Isotope evidence, *Geochim. Cosmochim. Acta*, 50, 693–709, 1986.
- Wimpenny, J., James, R. H., Burton, K. W., Gannoun, A., Mokadem, F. and Gislason, S. R.: Glacial effects on weathering processes: New insights from the elemental and lithium isotopic composition of West Greenland rivers, *Earth Planet. Sci. Lett.*, 1220 290(3–4), 427–437, doi:10.1016/j.epsl.2009.12.042, 2010.
- Wimpenny, J., Burton, K. W., James, R. H., Gannoun, A., Mokadem, F. and Gislason, S. R.: The behaviour of magnesium and its isotopes during glacial weathering in an ancient shield terrain in West Greenland, *Earth Planet. Sci. Lett.*, 304(1–2), 260–269, doi:10.1016/j.epsl.2011.02.008, 2011.
- 1225 Winsor, K., Carlson, A. E. and Rood, D. H.: ¹⁰Be dating of the Narsarsuaq moraine in southernmost Greenland: Evidence for a late-Holocene ice advance exceeding the Little Ice Age maximum, *Quat. Sci. Rev.*, 98, 135–143, doi:10.1016/j.quascirev.2014.04.026, 2014.
- [World Meteorological Organization: Manual on Stream Gauging. Volume I: Fieldwork, 2010.](#)
- Yde, J. C., Knudsen, N. T., Hasholt, B. and Mikkelsen, A. B.: Meltwater chemistry and solute export from a Greenland Ice Sheet catchment, Watson River, West Greenland, *J. Hydrol.*, 519(PB), 2165–2179, doi:10.1016/j.jhydrol.2014.10.018, 2014.

Table 1. Mineral weathering reactions and impacts on dissolved CO₂ concentrations.

<i>Eq.</i>	<i>Mineral</i>	<i>Acid</i>	<i>Abbreviation¹</i>	<i>Reaction</i>	<i>Impact on CO₂</i>
(1)	Carbonate	Carbonic	Carb _{CA}	$(\text{Ca, Mg})\text{CO}_3 + \text{H}_2\text{O} + \text{CO}_2 \rightarrow (\text{Ca}^{2+}, \text{Mg}^{2+}) + 2\text{HCO}_3^-$	CO ₂ sink
(2)		Sulfuric	Carb _{SA}	$2(\text{Ca, Mg})\text{CO}_3 + \text{H}_2\text{SO}_4 \rightarrow 2(\text{Ca}^{2+}, \text{Mg}^{2+}) + \text{SO}_4^{2-} + \text{H}_2\text{O} + \text{CO}_2$	CO ₂ source
(3a)	Silicate	Carbonic	Sil _{CA}	$(\text{Ca, Mg})\text{Al}_2\text{Si}_2\text{O}_8 + 2\text{CO}_2 + 3\text{H}_2\text{O} \rightarrow (\text{Ca}^{2+}, \text{Mg}^{2+}) + 2\text{HCO}_3^- + \text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4$	CO ₂ sink
(3b)				$(\text{Na, K})\text{AlSi}_3\text{O}_8 + \text{CO}_2 + 5.5\text{H}_2\text{O} \rightarrow (\text{Na, K}) + \text{HCO}_3^- + 0.5\text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4 + 2\text{H}_4\text{SiO}_4$	CO ₂ sink
(4a)		Sulfuric	Sil _{SA}	$(\text{Ca, Mg})\text{Al}_2\text{Si}_2\text{O}_8 + \text{H}_2\text{SO}_4 + \text{H}_2\text{O} \rightarrow (\text{Ca}^{2+}, \text{Mg}^{2+}) + \text{SO}_4^{2-} + \text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4$	No impact
(4b)				$2(\text{Na, K})\text{AlSi}_3\text{O}_8 + \text{H}_2\text{SO}_4 + 9\text{H}_2\text{O} \rightarrow 2(\text{Na}^+, \text{K}^+) + \text{SO}_4^{2-} + \text{Al}_2\text{Si}_2\text{O}_5(\text{OH})_4$	No impact

¹ Abbreviations are based first on the mineral class (carbonate = carb, silicate = sil) and then on the acid (carbonic acid = CA, sulfuric acid = SA)

Formatted: English (US)






Figure 1: Conceptual diagram of subglacial sources and sinks of CO_2 and CH_4 . Arrows indicate the direction of fluxes. Boxes represent processes, and sources of gases to subglacial meltwaters are indicated by blue text while sinks of gases to subglacial meltwater are indicated by red text. Gas bubbles, mechanical grinding, and OM remineralization are grouped because all are CO_2 and CH_4 sources.

Deleted: Arrow

Deleted: green



Deleted:

Deleted:) subglacial



Deleted: sites flowing to the Watson River in southwest Greenland...

Deleted: and the

Deleted: . Discharge monitoring of the Watson River is collected by PROMICE (van As et al., 2018) at the location represented by the yellow star.

245 Figure 2. Google Earth satellite images of study locations in a) Greenland including b) locations near the town of Kangerlussuaq, including the sub-Isunnguata water sampling location (dark blue circle) and Russell water sampling location (light blue square). The gold star represents the location of AK4, where proglacial river discharge records were collected. (c) Location of Kiattut Sermiat site (orange triangle) near the town of Narsarsuaq in southern Greenland, where water samples were collected.


1250




Deleted:

Deleted: discharge


Figure 3. Chemical parameters at Isunnguata (IS), Russell (RU) and Kiattut Sermiat (KS) subglacial water sampling sites versus day of year for a) specific conductivity (Sp.C), b) pH, c) dissolved oxygen (D.O.) percent saturation, d) alkalinity (Alk), e) measured CH₄ concentrations (left y-axis in ppm and right y-axis in nM), f) $\delta^{13}\text{C}\text{-CH}_4$ values, g) measured CO₂ concentrations (left y-axis in ppm and right y-axis in μM), and h) $\delta^{13}\text{C}\text{-CO}_2$ values. Atmospheric equilibrium concentrations are indicated by dashed lines and taken as 1.9 ppm for CH₄ and 410 ppm for CO₂. Error bars on CH₄ and CO₂ concentrations and stable isotopic compositions represents the standard deviation of replicates and are smaller than symbols for some data points.




1270 Figure 4. CH₄ dynamics over the course of the 2017 and 2018 melt seasons including a) the carbon fractionation factor (ϵ_c) between
 1275 dissolved CO₂ and CH₄ and b) the fraction of CH₄ oxidized (f_{ox}) for sub-Isunnguata (IS), Russell (RU) and Kiattut Sermiat (KS)
 samples. Fields of ϵ_c representing methanogenesis and oxidation values are based on Whiticar (1999). Values of ϵ_c between
 approximately 40 and 55 (gray shaded region in panel a) are produced for methanogenesis via acetate fermentation, while CO₂
 reduction produces values between approximately 50 and 90. Lower values result from a predominant isotopic signature of CH₄
 oxidation. Atmospheric input without additional alteration of CO₂ or CH₄ isotopic systematics results in a ϵ_c value of approximately
 40.



Deleted:



280 **Figure 5. Relationships between average daily discharge and CH₄ dynamics including a) CH₄ concentrations, b) $\delta^{13}\text{C}-\text{CH}_4$, c) f_{ox} , and d) ϵ_c for Isunnguata (IS) and Russell (RU) samples. Regressions are shown by dotted lines for Isunnguata and dashed lines for Russell samples. Horizontal error bars represent the standard deviation of average daily discharge for days samples were collected and are smaller than symbols for some data points.**



Deleted:

Deleted: Discharge relationships with

Deleted: c

Deleted: Average daily discharge estimates are for the outlet of the Watson River and provided by PROMICE (van As et al., 2018). ...





Figure 6. Mineral weathering model results in a) net impact of mineral weathering reactions on CO₂ (Net CO_{2-MW}; Eq. 7) for Isunnguata, Russell, and Kiattut Sermiat subglacial discharge sites (where negative values of Net CO_{2-MW} indicate net sequestration of CO₂ due to mineral weathering), and b) the proportional contribution of each mineral weathering reaction to the total change in CO₂ (% Total CO_{2-MW} Eq. 9a-9c).




Deleted:

Formatted: Font: 9 pt, Bold

Formatted: Normal

Deleted:

Formatted: Line spacing: single








Figure 7. Calculated CO₂-total values for a) sub-Isunnguata, b) Russell, and c) Kiattut Sermiat subglacial discharge against day of the year.



310 **Figure 8.** Relationships between average daily discharge a) CO₂ concentrations, and b) δ¹³C-CO₂. Regressions are shown by dotted lines for sub-Isunnguata and dashed lines for Russell samples. Horizontal error bars represent the standard deviation of average daily discharge for days samples were collected and are smaller than symbols for some data points.



Moved (insertion) [9]

Deleted: Correlations




Figure 9. Keeling plot indicating correlations between the magnitude of $\text{CO}_2\text{-total}$ and $\delta^{13}\text{C-CO}_2$ and for sub-Isunnguata (sub-IS), Russell (RU) and Kiattut Sermiat (KS) samples. Asterisk denotes the outlier not included in the regression between $\text{CO}_2\text{-total}$ and $\delta^{13}\text{C-CO}_2$ for sub-Isunnguata samples. The plotted regression line was constructed using the sub-IS samples only.

Deleted: a)

Deleted: b) NH_4

Deleted: Isunnguata

Deleted: in panel a).




Figure 9. Estimates of contributions of atmospheric CO_2 ($\text{CO}_2\text{-atm}$) compared to that derived from remineralized OM ($\text{CO}_2\text{-OM}$) for Isunnguata subglacial discharge site including a) relative contributions according to the day of the year, b) absolute ... [8]

Moved up [9]: Horizontal error bars represent the standard deviation of average daily discharge for days samples were collected and are smaller than symbols for some data points.

Deleted: Regressions between average daily discharge and proportional and absolute contributions of $\text{CO}_2\text{-atm}$ and $\text{CO}_2\text{-OM}$ sources are shown with dotted lines.

Formatted: English (UK)

Page 13: [1] Deleted	Pain,Andrea J	12/31/20 12:28:00 PM
-----------------------------	----------------------	-----------------------------



Page 13: [2] Deleted	Pain,Andrea J	12/31/20 12:28:00 PM
-----------------------------	----------------------	-----------------------------



Page 13: [3] Deleted	Pain,Andrea J	12/31/20 12:28:00 PM
-----------------------------	----------------------	-----------------------------



Page 13: [4] Deleted	Pain,Andrea J	12/31/20 12:28:00 PM
-----------------------------	----------------------	-----------------------------



Page 13: [5] Deleted	Pain,Andrea J	12/31/20 12:28:00 PM
-----------------------------	----------------------	-----------------------------



Page 13: [6] Deleted	Pain,Andrea J	12/31/20 12:28:00 PM
-----------------------------	----------------------	-----------------------------



Page 14: [7] Deleted	Pain,Andrea J	12/31/20 12:28:00 PM
-----------------------------	----------------------	-----------------------------



Page 33: [8] Deleted	Pain,Andrea J	12/31/20 12:28:00 PM
-----------------------------	----------------------	-----------------------------

