

Environmental Drivers of Circum-Antarctic Glacier and Ice Shelf Front Retreat over the Last Two Decades

Celia A. Baumhoer¹, Andreas. J. Dietz¹, Christof Kneisel², Heiko Paeth² and Claudia Kuenzer^{1,2}

¹German Remote Sensing Data Center (DFD), German Aerospace Center (DLR), Muenchner Strasse 20, D-82234 Wessling, Germany

²Institute of Geography and Geology, University Wuerzburg, Am Hubland, D-97074 Wuerzburg, Germany

Correspondence to: Celia A. Baumhoer (celia.baumhoer@dlr.de)

Abstract. The safety band of Antarctica consisting of floating glacier tongues and ice shelves buttresses ice discharge of the Antarctic Ice Sheet. Recent disintegration events of ice shelves and glacier retreat indicate a weakening of this important safety band. Predicting calving front retreat is a real challenge due to complex ice dynamics in a data-scarce environment ~~being that~~ are unique for each ice shelf and glacier. We explore to what extent easy to access remote sensing and modelling data can help to define environmental conditions leading to calving front retreat. For the first time, we present a circum-Antarctic record of glacier and ice shelf front retreat over the last two decades in combination with environmental variables such as air temperature, sea ice days, snowmelt, sea surface temperature and wind direction. We find that the ~~Antaretic ice sheet~~ Antarctic Ice Sheet area ~~shrank-decreased by -29,618±1,193~~ 29,618±29 km² in extent between 1997-2008 and gained an area of 7,108±~~1,029~~ 144.4 km² between 2009 and 2018. Retreat concentrated along the Antarctic Peninsula and West Antarctica including the biggest ice shelves Ross and Ronne. ~~In several cases G~~ glacier and ice shelf retreat ~~comes along~~ occurred in conjunction with one or several changes in environmental variables. Decreasing sea ice days, intense ~~snow-melt~~ snowmelt, weakening easterlies and relative changes in sea surface temperature were identified as enabling factors for retreat. In contrast, relative increases in ~~mean~~ air temperature did not correlate with calving front retreat. For future studies a more appropriate measure for atmospheric forcing should be considered including above zero-degree days and temperature extreme events. To better understand drivers of glacier and ice shelf retreat it is of high importance to analyse the magnitude of basal melt through the intrusion of warm Circumpolar Deep Water (CDW) driven by strengthening westerlies and to further assess surface hydrology processes such as meltwater ponding, runoff and lake drainage.

1 Introduction

A safety band of floating ice shelves and glacier tongues fringes the ~~Antaretic ice sheet~~ Antarctic Ice Sheet (Fürst et al., 2016). Large glaciers and ice shelves create buttressing effects decreasing ice flow velocities and ice discharge (De Rydt et al., 2015; Gagliardini et al., 2010; Royston and Gudmundsson, 2016). The recent large-scale retreat of ice shelves and glacier fronts along the Antarctic Peninsula (AP) and the West Antarctic Ice Sheet (WAIS) indicates a weakening of this safety band (Cook and Vaughan, 2010; Friedl et al., 2018; Rankl et al., 2017; Rott et al., 2011). The glacier or ice shelf front is defined as the border

between the ice sheet and the ocean whereby ice shelves and glaciers are the floating extensions of the ice sheet (Nicholls et al., 2009). Calving front retreat (including both, glacier and ice shelf front retreat) ~~increases-can increase~~ ice discharge and hence the contribution to global sea level rise (De Angelis and Skvarca, 2003; Seehaus et al., 2015). ~~Increases in ice discharge occur if ice shelf areas with strong buttressing forces are lost (Fürst et al., 2016).~~ The current contribution of the ~~Antarctic ice sheet~~ Antarctic Ice Sheet to global sea level rise is 7.6 ± 3.9 mm (1992-2017) but over ~~the this last decades study period~~ a strong trend of mass loss acceleration was observed for the WAIS after ice shelves and glaciers retreated and thinned (IMBIE, 2018). In contrast, there is no clear trend in the mass balance of the East Antarctic Ice Sheet (EAIS). ~~as~~ Since the 1990s ~~a~~ Altimetry measurements ~~have shown~~ show very a small gain (but with high uncertainties) for the EAIS with 5 ± 46 Gt/yr (1992-2017) (IMBIE, 2018). However, ~~a strong mass loss trend of -47 ± 13 Gt/yr (1989-2017) is calculated using the mass budget method~~ calculated with the mass budget method a strong mass loss trend exists with -47 ± 13 Gt/yr (1989-2017). (Rignot et al., 2019). Glacier terminus positions along the EAIS experienced a phase of retreat between 1974 and 1990 followed by a phase of advance until 2012. The single exception at East Antarctica is Wilkes Land with retreating glacier fronts (Miles et al., 2016). ~~Whether a glacier or ice shelf front advances or retreats depends mainly on four different factors: internal ice dynamics, geometry, external mechanical forcing and external environmental forcing (Alley et al., 2008; Benn et al., 2007b; Cook et al., 2016; Luckman et al., 2015; Walker et al., 2013).~~ The combined influence of those factors make it challenging to create a realistic calving law and making iceberg calving still one of the least understood ice shelf processes (Bassis, 2011). Internal ice dynamics determine the glacier front position by a function of ice velocity, strain rate, ice thickness and water depth (Benn et al., 2007a) creating a critical self-organized system readily changing between frontal advance and retreat (Åström et al., 2014). Frontal retreat starts with the formation of a crevasse originating from a strain rate surpassing the yield stress of ice (Mosbeux et al., 2020). As soon as the surface and/or basal crevasses evolve and fractures propagate, the calving margin creates at the point where the depth of a surface crevasse equals the freeboard above sea level (Benn et al., 2007b, 2007a). For ice shelves, the calving position evolves where crevasses develop into through-cutting fractures (rifts). Those ice shelf rifts can propagate further to the ice front or intersect with other rifts resulting into a tabular iceberg calving event (Benn et al., 2007b; Joughin and MacAyeal, 2005) whereas the extent of the ice shelf and the size of the iceberg is defined by the rift location (Lipovsky, 2020; Mosbeux et al., 2020; Walker et al., 2013). Further boundary conditions such as fjord geometry (Alley et al., 2008; Catania et al., 2018), water depth (Bassis and Walker, 2012; Pollard et al., 2015) and bed topography (Hughes, 1981) influence the stability of the calving margin. The previously described natural cycle of growth and decay of a glacier or ice shelf (Hogg and Gudmundsson, 2017) can be disturbed by external mechanical and environmental forcing. It was found that mechanical forcing arises from tides (Rosier et al., 2014), ocean swell (Massom et al., 2018), iceberg collision (Massom et al., 2015) and tsunamis (Walker et al., 2013) which can destabilize floating ice shelves and glacier tongues initiating iceberg calving (Alley et al., 2008). Glaciers and ice shelves are in direct interaction with the atmosphere and ocean and are hence also sensitive to changes in environmental conditions (Domack et al., 2005; Kim et al., 2001; Vaughan and Doake, 1996; Wouters et al., 2015). Environmental drivers can destabilize floating ice tongues and shelves through various oceanic and atmospheric forced processes causing mechanical weakening and

Formatiert: Schriftart: 10 Pt.

65 fracturing (Pattyn et al., 2018). Instead of sporadic calving by tabular icebergs (a sign of a natural calving event within the calving cycle) more frequent small-scale calving events can occur through environmental forcing (Liu et al., 2015). Destabilization through atmospheric forcing arises from meltwater runoff (Cowton et al., 2018), thinning through surface melt (Howat et al., 2008), hydrofracture (Kopp et al., 2017; Pollard et al., 2015), lake ponding and lake drainage (Banwell et al., 2013; Leeson et al., 2020). Ocean forcing weakens the floating ice by basal melt inducing thinning and grounding line retreat (Konrad et al., 2018; Paolo et al., 2015; Rignot et al., 2013) as well as warmer ocean surface temperatures undercutting the ice cliff at the waterline (Benn et al., 2007b), reducing stabilizing fast ice (Larour, 2004) and enhanced sea ice reduction (Massom et al., 2018; Miles et al., 2016). Environmental forcing on tidewater glaciers has been frequently observed on Greenlandic glaciers (Cowton et al., 2018; Howat et al., 2008; Luckman et al., 2015) but for Antarctica in many regions it is unclear where and at what amount environmental drivers cause calving front retreat (Baumhoer et al., 2018; Pattyn et al., 2018).

75 Current knowledge on environmental forced calving front retreat in Antarctica can be summarized by the following studies.

Glaciers and ice shelves are in direct interaction with the atmosphere and the ocean and are hence sensitive to changes in environmental conditions (Domack et al., 2005; Kim et al., 2001; Vaughan and Doake, 1996; Wouters et al., 2015). Nevertheless, changes in glacier and ice shelf extent can be independent from environmental forcing as glaciers experience a natural cycle of decay and growth (De Rydt et al., 2019; Hogg and Gudmundsson, 2017). Therefore, the identification of driving forces for fluctuations in ice shelf and glacier extent is challenging and has been subject of many discussions in the past. Glacier retreat along the Antarctic Peninsula was first only associated to atmospheric warming (Cook et al., 2016; Mercer, 1978), until more recent studies identified ocean forcing as the main driver (Cook et al., 2016; Wouters et al., 2015). Additionally, the formation of melt ponds on the ice shelf surface has been discussed as an enhancing factor for calving.

85 Meltwater can initiate crevasse propagation resulting in hydrofracture and ice shelf retreat (Scambos et al 2000, Scambos 2017). The poleward shift of the west wind drift causes upwelling Circumpolar Deep Water (CDW). This allows warmer ocean waters to reach increases ocean temperatures at the bottom of ice shelves inducing basal melt and ice shelf thinning which was observed in the Bellingshausen and Amundsen Sea (Dutrieux et al., 2014; Thoma et al., 2008; Wouters et al., 2015). Basal melt, when combined with Especially in combination with a retrograde bed (Hughes, 1981; Scheuchl et al., 2016), has leadled to a retreat of the grounding line the basal melt causes a retreat of the grounding line followed by increased ice discharge (Konrad et al., 2018; Rignot et al., 2013) for example at former Wordie Ice Shelf (Friedl et al., 2018; Walker and Gardner, 2017). In contrast, the retreat of calving fronts along Wilkes Land (EAIS) was associated with a reduction in the duration of the sea ice cover of sea ice days (Miles et al., 2016).

The drivers of ice shelf and glacier retreat and advance can be manifold depending on studied variables, time periods and regions. Therefore, the identification of environmental driving forces for fluctuations in ice shelf and glacier extent is challenging and has been subject of many discussions in the past. So far, there has not been a comprehensive analysis comparing circum-Antarctic glacier terminus change on a continental scale within uniform time scales (Baumhoer et al., 2018). In this paper, we explore Antarctic calving front change over the last two decades by analysing Antarctic coastal change (see

Figure 1) and address the question if environmental drivers forced the observed changes in glacier and ice shelf front fluctuations, assess the connection to changing boundary conditions. We compare changes in -Antarctic calving fronts in two decadal time steps (1997-2008 and 2009-2018) to minimize the effect of short-term glacier front fluctuations. To identify potential links between calving front retreat and recent changes in the Antarctic environmental conditions, we correlated two decades of glacier change with climate data including air and sea surface temperature as well as changes in wind direction, snowmelt and sea ice cover.

105




Figure 1 Coastal change of Antarctica between 1997 and 2018 with enlarged views (counter-clockwise) for Larsen C Ice Shelf, Pine Island Bay, Oates Land, Wilkes Land, Shackleton Ice Shelf, Shirase Bay and Dronning Maud Land. Colours indicate the timing of retreat and advance. The pie chart visualizes losses and gains in the area of the Antarctic Ice Sheet with regard to the indicated year. Background: LIMA Landsat Mosaic.

2 Processed and Analysed Data Sets

2.1 Coastlines

To calculate the decadal retreat and advance of Antarctic glaciers and ice shelves, we use calving front positions of three Antarctic coastline products from 1997, 2009 and 2018. The Antarctic coastline is the border between the ~~Antarctic ice sheet~~Antarctic Ice Sheet and the ocean and hence includes ice shelf and glacier front positions (Baumhoer et al., 2018). Coastal change not only includes changes in floating calving fronts but also of grounded ice walls. Timewise, we refer to first (1997-2008) and second (2009-2018) decade even though the time periods are 11 and 9 years, respectively, limited by the availability of coastline products. The first coastline product was automatically extracted from the high-resolution Radarsat-1 mosaic by adaptive thresholding and finalized by manual correction (Liu and Jezek, 2004). The Radarsat-1 mosaic imagery was acquired between September and October 1997 with a spatial resolution of 25 m. The entire dataset is freely available at the National Snow and Ice Data Center (NSIDC) (1997). Glacier and ice shelf fronts for the year 2009 were manually delineated from the MOA 2009 Surface Morphology Image Map acquired during austral summer 2008/2009 (Nov-Feb) with a spatial resolution of 125 m (Scambos et al., 2007). This dataset is also freely available at NSIDC (2009). The coastline for 2018 was automatically extracted via a fully convolutional network from Sentinel-1 mid resolution (40 m) dual-pol imagery (see methods section).

2.2 ERA 5

We use ERA5-Land monthly averaged atmospheric re-analysis data for air temperature and snowmelt information. It is the most state-of-the-art reanalysis product from the European Centre for Medium-Range Weather Forecasts (ECMWF) replacing the former ERA-Interim product (Hersbach et al., 2020). Data is available from 1981/82 to today at a 9 km spatial resolution. As ground-based meteorological observations are scarce over the Antarctic continent we decided to use the reanalysis data even though modelled data is less accurate compared to in-situ measurements. Studies comparing ~~in situ~~in-situ observations to modelled data prove that ERA5 surface air temperature (2 m temperature) outperforms the former ERA-Interim product. ERA5 temperature data has the ability to capture annual variability and magnitude of temperature change over the Antarctic Ice Sheet (Gossart et al., 2019; Tetzner et al., 2019). The mean absolute error is 2.0° C with higher accuracies in the coastal areas and less accurate results in the interior of the ice sheet (Gossart et al., 2019). Tetzner (2019) report much higher accuracies for the Antarctic Peninsula with a mean absolute error of -0.13° C. Compared to the mean, the variability of temperature is captured with high accuracy at a mean Pearson correlation coefficient of 0.98 (Gossart et al., 2019; Tetzner et al., 2019). We divided temperature measurements into the cooler half (“winter” Apr-Sep) of the year and warmer half (“summer” Oct-Mar) to separate the environmental forcing for different seasons. Snowmelt data should be handled with care as accuracy assessments for the ERA5 Land snowmelt product were not yet performed. Surface mass balance (SMB) data including modelled snowmelt data was found to slightly underestimate the SMB

Feldfunktion geändert

Feldfunktion geändert

(Gossart et al., 2019). Snowmelt is calculated within the summer months December, January and February where most melt occurs. The amount of ~~snow melts~~ snowmelt is calculated in mm water equivalent (mm w. eq) per day.

~~The Antarctic continent is circled by weak and irregular easterly winds driven by high pressure areas over the interior of the Antarctic continent created by cold and dry air. Easterlies weaken in the case of a positive Southern Annular Mode (SAM) and the west wind drift shifts poleward. In order to asses those changes in wind direction, For zonal (West to East) wind speed estimations~~ we use ERA5 monthly averaged zonal (West to East) wind speed estimations data at 10 m above the surface with a resolution of approx. 31 km. ERA-5 captures the spatial variability of near-surface wind speed but underestimates strong winds and coastal winds whereas the wind speed over the interior is captured very accurately ~~especially~~ during summer (Gossart et al., 2019). Over the ocean, ERA5 annual mean zonal wind speed is considerably underestimated compared to ASCAT (Advanced Scatterometer) observations (Belmonte Rivas and Stoffelen, 2019). Overall, the mean absolute error over the continent is 2.8 m/s with high variance in space and time but an accurate representation of the annual variability in zonal wind speed is achieved (Gossart et al., 2019). ERA5 data is freely available at the Copernicus Climate Change Service Climate Data Store (CDS) (C3S, 2017). Zonal Wind was calculated for the summer months (DJF) because ~~especially~~ easterly winds show a weakening trend during summer but not throughout the entire year (Hazel and Stewart, 2019).

2.3 Sea Ice Days

The most recent Global Sea Ice Concentration Climate Data Record (Version 2) (Lavergne et al., 2019) was downloaded from the Ocean and Sea Ice Satellite Application Facility (Osisaf, 2020). We used the daily product (OSI-450, OSI-430-b) during sea ice months April through October in line with previous studies (Massom et al., 2013; Miles et al., 2016). The product covers the time period 1982 to 2018 and is derived from passive microwave data by Nimbus 7 and DMSP satellites. The final sea ice concentration is computed by the passive microwave data in combination with ERA-Interim data. The standard deviation of mismatch between OSI products and ice chart analysis on sea ice concentration is 8 % for ice and open water during winter (JJA). The trend in sea ice extent is very similar between OSI and ice chart products (Brandt-Kreiner et al., 2019) which allows the assumption of a very accurate data set. The product has a resolution of 25 km x 25 km. In order to calculate actual sea ice days we count each pixel per day with a sea ice concentration higher than 15 % as suggested by previous studies (Miles et al 2016, Massom et al 2013). ~~Especially in~~ During the early acquisition years data gaps occur and second-daily acquisitions exist until mid of 1987. In this case, we multiplied the monthly available sea ice days by the proportional amount of missing days. We decided not to use data for the year 1986 as data for entire months were missing and mean sea ice days per year could not be calculated accurately.

2.4 Sea Surface Temperature

Sea surface temperature measured by satellite sensors is also provided by the CDS (C3S, 2017). The most up-to-date product Level 4 (Version 2) consists of multiple satellite observations from NOAA, ERS, Envisat and Sentinel-3 satellites with a 0.05 ° gridded resolution (approx. 5.5 km). We calculated surface temperatures only for months with little to no sea ice cover (Oct-

Mar). This reduces errors over sea ice where the thickness and concertation of sea ice can account for large measurement errors
 175 (Kwok and Comiso, 2002). Uncertainties of Level 4 data vary depending on the year of acquisition and latitude of
 measurement. The median difference compared to in-situ drifter measurements is up to -0.4 K for low latitudes during the
 early acquisition years before 1996. Afterwards the accuracy increases to better than -0.1 K for low latitudes. A stability
 assessment of the Level 4 product calculated a maximum trend of 0.01 K per year (Embury, 2019). A summary of all processed
 climate variables is given in [Table 1](#)~~Table 1~~.

180 **Table 1 Summary of processed climate variable data sets.**

Climate Variable	Time Span	Season	Spatial Resolution	Accuracy	Data	Data Store
Air Temperature	1982-2018	Summer (Oct-Mar) Winter (Apr-Sep)	9 km	0.13 – 2.0 °C	modelled	CDS
Snow Melt	1982-2018	Dec-Feb	9 km	-	modelled	CDS
Zonal Wind	1982-2018	Dec-Feb	31 km	2.8 m/s	modelled	CDS
Sea Ice Days	1982-2018 (not 1986)	Apr-Oct	25 km	8 % std to ice chart	modelled + satellite	Osisaf
Sea Surface Temperature	1982-2018	Oct-Mar	~5.5 km	-0.1 to -0.4°C	multiple satellites	CDS

3 Method

3.1 Extraction of Sentinel-1 Coastline

We use a modified version of the automatic coastline extraction approach published by Baumhoer et al. (2019) to extract the
 185 Antarctic coastline for 2018. To cover the entire Antarctic coastline, 158 dual-pol and 17 single-pol medium resolution
 Sentinel-1 scenes were processed. Dual polarized scenes contain radar backscatter values for the polarizations HH and HV
 whereas single polarized scenes only include the HH polarization. The coastline was split into 18 zones based on ice flow
 divides, defining major ice sheet basins after Rignot et al. (2011). For each zone all available scenes acquired during winter
 months ([June, July, August](#)~~Jun-Aug~~) in 2018 were selected. Depending on the scene availability, each zone was covered at
 190 least by one Sentinel-1 scene and in the best case by three scenes. In case no dual-pol scenes were available single-pol data
 was selected. First, each scene was pre-processed (thermal correction, calibration, terrain correction), masked with a coastline
 buffer of 100 km and tiled into 780x780 pixel tiles. The convolutional neural network (CNN) U-Net was used to segment each
 tile into the class ocean and the class land ice. We trained the U-Net on 15 different Antarctic coastal regions during various
 seasons with 40,036 image tiles from 75 Sentinel-1 scenes. For single-pol scenes we re-trained our network for HH polarized
 195 images only. This decreased the accuracy slightly as only one polarization limits the information on surface backscatter
 characteristics. In the post-processing step the mean of all segmented tiles within one zone was calculated and then thresholded
 by a prediction probability of 50 % for the class land ice. In case of multi-coverage by several satellite scenes more robust
 results were obtained by merging the prediction probabilities. Morphological filtering and the exclusion of higher ice sheet

200 areas by integrating elevation information from the TanDEM-X Polar DEM 90 reduced further errors. Finally, from the binary segmentation results the border between both classes was extracted as the final coastline.

3.2 Derivation of Calving Front Change

205 Calving front change was estimated by calculating the change in area between the coastlines in 1997, 2009 and 2018 over the area of glaciers and ice shelves. As coastline extraction and delineation is a very subjective task and all coastlines originated from different sources and resolutions deviations in many areas occurred. Especially Areas of fast ice, mélange and icebergs trapped in sea ice were error sources. For the automatically extracted coastline 2018, errors existed along the western Antarctic Peninsula and in areas where only single-pol imagery was available. The MODIS derived coastline often included snow covered sea ice which is difficult to distinguish from glacier ice in optical imagery. To minimize errors and mismatches all three coastlines were manually corrected and adjusted. Each coastline product was corrected based on the satellite imagery from which they were originally created. After manual correction any change in area between the coastline products could be attributed to glacier front change. From each coastline a raster (resolution 40 x 40 m) was created with a unique value over the ice covered area. All three raster layers were stacked and summed up, so each raster value pixel is associated with retreat or advance of the specific year. The area change was calculated for each major floating ice shelf and glacier wider than 3 km based on a 40 m resolution raster in Polar Stereographic Projection. To account for inaccuracies in the manual adjustment of all coastlines, we measured the accuracy as the area change over 30 randomly picked stable coastline areas. Any change over those regions can be attributed to errors in manual delineation, imagery resolution differences and errors in orthorectification of the different satellite image mosaics. The error is calculated per km of the coastline and calculated in proportion to the actually-measured front length. On average, the coastlines deviated ± 1.2 pixels per kilometre per year. Broken down for each coastline product, the error between 1997-2009 was 0.4 pixels per km per year, 2.3 pixels between 2009-2018 per km per year and 0.9 pixels between 1997 -2018 per km per year.

220 3.3 Climate Data Correlation

225 In order to assess the influence of climate variables (such as air temperature, sea ice days, sea surface temperature, snowmelt and zonal wind) on glacier and ice shelf retreat, we spatially correlated the percentage of advance and retreat for each minor glacier basin (as defined by Mouginit (2017)) with the decadal mean value of each climate variable. Means were calculated from 37-years of climate data for the time periods 1997-2008, 2009-2018 and 1982-1996. Hence, the inputs of the correlation covered the two time spans 1997-2008 and 2009-2018 for which also the glacier retreat was calculated. To also assess relative changes in the variables to previous times we subtracted the long-termreference mean (1982-1996). Zonal wind, sea ice coverage and sea surface temperature means were calculated within a 100 km seawards buffer along the coastline. Mean air temperature and snowmelt was calculated within a 100 km buffer landwards from the coastline covering the surface of ice shelves. The 100 km landwards buffer was chosen as the best trade-off to calculate averages over the ice shelf area by not including too much of the ice sheet. Due to a circum-Antarctic analysis this is only a rough approximation and especially over

smaller glaciers within steep terrain the average not only includes changes over the glacier tongue but the surrounding area. Again, also relative values were calculated by subtracting the ~~long-term~~reference mean 1982-1996. To remove the effect of different basin sizes and different amounts of ice discharge we took the percentage of advance and retreat within each basin instead of the absolute value for the correlation. Input data for the Pearson correlation were the mean of all assessed climate variables (absolute and relative averages) as well as the percentage of retreat and advance. This created 14 different variables which were correlated with each other based on 188 observations (N=188). The number of observations ~~composes-is derived~~ from 94 assessed glacier basins with variable averages for the two different decades (1997-2008 and 2009-2018).

4 Results

4.1 Advance and Retreat of Antarctic Glaciers and Ice Shelves

Circum-Antarctic glacier and ice shelf front changes were assessed by comparing -Antarctic coastline products during 1997-2008 and 2009-2018. The results are ~~visualized-shown in Figure 2~~Figure 2. Between 1997 and 2008 ice shelf and glacier extents decreased by -29,618±~~29-1,193~~ km² where 69 % of the total changed area retreated and 31 % advanced (see Table 2Table 2 for all change rates). In contrast, during the period 2009 to 2018 a slight area increase of 7,108±~~144.4~~1,029 km² could be observed with 44 % of the total changed area retreating and 56 % advancing. The locations of area change are almost similar for both observation periods. Ice shelves along the Antarctic Peninsula retreated over the entire observation period (1997-2018). The only exception was the advance of the Larsen D Ice Shelf which is located close to the Ronne Ice Shelf in the Weddell Sea (see Figure 2Figure 2). ~~B~~Especially between 1997 and 2008, ~~huge-the large~~ disintegration events of the Larsen B, Wilkins and Wordie Ice Shelf resulted in a 37 % higher calving amount ~~fromat~~ the Antarctic Peninsula, as compared to the amount calved from this region between-during 2009-2018. During the second decade the breakup of iceberg A-68 from the Larsen C Ice Shelf and the further disintegration of the Wilkins Ice Shelf were ~~hotspots-regions~~ of strong area loss. In total, the rate of retreat along the Antarctic Peninsula was 5-6 times higher than glacier advance over the last 20 years.

Table 2 Retreat and advance rates of Antarctic glaciers and ice shelves for each basin between 1997 and 2018 (upper table). Total lost and gained ice shelf/glacier area per decade (lower table). For basin abbreviations see Figure 2.

		1997-2008		2009-2018		1997-2018	
basin		advance (km ² /yr)	retreat (km ² /yr)	advance (km ² /yr)	retreat (km ² /yr)	advance (km ² /yr)	retreat (km ² /yr)
EAIS	A-Ap	249±14.9	164±17.4	295±27.5	112±13	215±11.8	85±6.5
	Ap-B	62±4.8	89±5.6	80±8.9	53±4.2	34±3.8	36±2.1
	B-C	187±1.9	4±2.2	186±3.5	20±1.6	179±1.5	4±0.8
	C-Cp	380±7.7	167±9	363±14.2	153±6.7	323±6.1	111±3.3
	Cp-D	49±4.6	58±5.4	45±8.6	97±4.1	23±3.7	51±2
	D-Dp	154±5	64±5.8	156±9.2	317±4.3	85±3.9	111±2.2
	Dp-E	68±9.5	58±11.1	74±17.5	81±8.3	34±7.5	32±4.1
	E-Ep	6±3	411±3.5	156±5.5	8±2.6	5±2.4	154±1.3
	Jpp-K	163±2.1	2±2.4	252±3.8	2±1.8	203±1.6	1±0.9
	K-A	306±7.8	23±9.1	285±14.3	82±6.8	270±6.2	24±3.4
	EAIS	1626±61.1	1040±71.4	1894±112.9	926±53.4	1370±48.6	607±26.6
AP	Hp-I	20±4.5	328±5.3	25±8.4	305±4	11±3.6	306±2
	I-lpp	87±3.9	938±4.6	59±7.2	592±3.4	12±3.1	715±1.7

WAIS	lpp-J	91±4.5	67±5.3	118±8.4	75±4	71±3.6	38±2
	AP	198±13	1333±15.2	202±24	972±11.4	94±10.3	1060±5.7
	Ep-F	160±7.7	868±8.9	452±14.1	74±6.7	136±6.1	341±3.3
	F-G	103±5.2	101±6.1	73±9.7	106±4.6	59±4.2	73±2.3
	G-H	40±3.6	371±4.2	49±6.7	369±3.2	14±2.9	340±1.6
	H-Hp	26±3.7	35±4.3	6±6.8	97±3.2	3±2.9	51±1.6
	J-Jpp	29±5.8	1127±6.8	643±10.7	26±5.1	13±4.6	317±2.5
	WAIS	358±26	2502±30.4	1222±48	672±22.7	225±20.7	1121±11.3

1997-2008			2009-2018		1997-2018	
TOTAL	advance (km ²)	retreat (km ²)	advance (km ²)	retreat (km ²)	advance (km ²)	retreat (km ²)
WAIS	3942±286	27525±334	11612±456	6382±216	4617±424	22970±232
AP	2178±143	14660±167	1920±228	9232±36	1929±212	21722±116
EAIS	17886±672	11439±785	17990±1072	8801±51	28089±997	12453±545
AIS	24006±1100	53624±1286	31522±1756	24414±303	34636±1632	57146±893
difference	-29618±1193		7108±1029		-22510±1263	

255 The West Antarctic Ice Sheet (WAIS) lost more than three-times as much calving area in the first decade (1997-2008) compared to 2009-2018. ~~75% of~~In West Antarctica, 75% of glacier and ice shelf retreat ~~were caused by~~can be attributed to the calving of Ronne and Ross West Ice Shelf during the first decade. The retreat at WAIS was significantly lower in the second decade whereas the Ross and Ronne ~~Ice Shelf~~ice shelves started to advance again. A different pattern occurs when considering West Antarctic calving front change without the Ronne and Ross ~~Ice Shelf~~ice shelves. Then the glacier and ice shelf retreat during both observation periods is predominant. Pine Island and Thwaites glacier retreated over the entire observation period (1997-2018). The Getz and Abbot ~~Ice Shelf~~ice shelves had stable front positions within the first decade but started to retreat since 2009. Only the Crosson Ice Shelf showed a contradicting pattern with retreat in the first and re-advance in the second decade.

The glaciers and ice shelves of the EAIS showed an overall stable advancing tendency with similar rates for both decades with ~~slightly 2638±418 km²~~more retreat during the first (attributed to calving of Ross East Ice Shelf) and no clear change in advance (slightly 104±872 km²) more advance during the second decade. Strongest advance was observed for the Amery and Filchner ~~Ice Shelf~~ice shelves as well as at the shelves ~~of~~at Dronning Maud Land and Queen Mary Land. During 1997-2008, clear retreat appeared for the Ross East Ice Shelf and along Enderby Land especially ~~in~~at Shirase Bay. Wilkes Land as well as Victoria Land had an almost equal share of retreat and advance. Between 2009 and 2018, the Ross Ice Shelf and Enderby Land entered a phase of advance whereas Wilkes Land, George V and Adélie Land started to retreat predominately.

Formatiert: Hochgestellt

Formatiert: Hochgestellt




Figure 2 Glacier and ice shelf extent changes for major glaciers and ice shelves over the last two decades. Circles indicate the rate of retreat or advance. Major ice sheet basins as defined by Rignot et al. (2011).

4.2 Air Temperature

The mean Antarctic air temperature was equal for both decades during winter but in summer +0.6 °C higher in the second decade (2009-2018). **But However**, the overall mean does not reflect the strong regional differences in air temperature changes as shown in [Figure 3](#). During summer and winter the Antarctic Peninsula and the adjoining coast along Bellingshausen

and Amundsen Sea as well as Queen Mary Land were cooler between 2009-2018 whereas the interior of the ice sheet warmed during this period. Compared to the ~~long-term~~reference mean (1982-1996) summer temperatures between 1997-2008 were cooler except for the Peninsula, Queen Mary Land, and the Ross West and Filchner Ice Sheet ice shelves. In the second decade this changed in sign where the Antarctic Peninsula started to cool down and the interior increased in temperature. For the winter months almost over the entire Antarctica continent a warming tendency was observed compared to the ~~long-term~~reference mean. Only exceptions were Victoria, Oates and George V Land. Additionally, over Getz Ice Shelf cooler temperatures by of 1°C occurred in 2009-2018 compared to the ~~long-term~~reference mean. For Dronning Maud Land a strong change was observed as the ice shelf surfaces warmed up to 1.8°C in the second decade.





Figure 3 Absolute ~~long-term~~air temperature (1982-1996) and relative changes compared to the ~~long-term~~reference mean. Additionally, the difference between the first and second decade is illustrated.

4.3 Sea Ice Days

A decrease or increase in sea ice days refers to the difference in the number of sea ice days per year (during the sea ice season April to OctoberApr-Oct) compared to the ~~long-term~~reference mean. The strongest decrease in sea ice days could be observed at the northern tip of the Antarctic Peninsula with a shorter sea ice cover per year of up to 40 days during the first decade compared to the ~~long-term~~reference mean. During 1997 and 2008 a decrease in sea ice days was observed along the

295 Bellingshausen Sea with up to 20 days less and in the Amundsen Sea with up to 5-10 days less. Compared to the first decade, the mean sea ice cover persisted longer in the second decade along the Antarctic Peninsula and the Bellingshausen Sea whereas in the Amundsen Sea the sea ice days decreased up to 10 days further compared to the first decade. Along the EAIS, sea ice coverage increased 10 days on average except for Dronning Maud Land with a slight decrease of up to 5 days. When comparing the two decades more sea ice days occurred during 2009 to 2018 along the EAIS and the northernmost Antarctic Peninsula
 300 whereas the decrease along West Antarctica continues. Changes in sea ice days were less extreme along East Antarctica with ± 10 days in difference compared to 1982-1996. Between 1997-2008 the duration of sea ice cover was up to 5 days shorter along Dronning Maud and Wilkes Land compared to the ~~long-termreference~~ mean. In the second decade sea ice days along the entire EAIS increased with a highest increase along Wilkes Land with up to 10 days (compared to 1982-1996). The described changes in sea ice days are ~~visualized-shown~~ in ~~Figure 4~~Figure 4.



305 **Figure 4** Difference in the number of sea ice days per year (during the sea ice season ~~April to October~~Apr-Oct) compared to the ~~long-termreference~~ mean (1982-1996) and the difference in sea ice cover duration between both decades.

4.4 Sea Surface Temperature

The changes in sea surface temperature during months with little sea ice cover (~~October to March~~Oct-Mar) relative to the ~~long-termreference~~ mean are shown in ~~Figure 5~~Figure 5. Sea surface temperatures of the ~~southern ocean~~Southern Ocean cooled ($\sim -0.5^\circ$) along the EAIS and warmed along WAIS and the western Antarctic Peninsula ($\sim +0.5^\circ\text{C}$) within the last two decades compared to 1982-1996 which is above the data uncertainty of 0.1 to 0.4 K. The warming was less strong in the first decade and intensified within the second decade especially in the Bellingshausen Sea with maxima at George VI north ($+0.65^\circ\text{C}$) and Pine Island Bay ($+0.35^\circ$) with a slightly weaker increase along the Amundsen Sea ($+0.15^\circ\text{C}$). The cooling along the East
 310 Antarctic was less strong in the second decade. Along East Antarctica for Amery Ice Shelf ($+0.35^\circ\text{C}$), West Ice Shelf ($+0.2^\circ\text{C}$) and Shackleton Ice Shelf ($+0.25^\circ$) warming was observed compared to the first decade.




Figure 5 Mean sea surface temperature changes (October-to-March) compared to 1982-1996 and the difference in sea surface temperatures between both decades.

4.5 Snowmelt

Snowmelt over the Antarctic Peninsula was more extensive during the long-term reference mean compared to the recent two decades. This is the reason why relative snowmelt in 1997-2008 and 2009-2018 was mostly negative with up to -0.3 mm w. eq. per day (see Figure 6). Nevertheless, strong snowmelt occurred also during the recent two decades but more selective over Wilkins and Larsen B Ice Shelf as well as at the northern tip of the Antarctic Peninsula. During the long-term reference mean, snowmelt concentrated on the Antarctic Peninsula, Ronne, Abbot, Shackleton and Amery Ice Shelf as well as Shirase Bay. In the first decade snowmelt expanded to Pine Island Bay, Getz Ice and Ross Ice Shelf as well as along Wilkes, George V and Dronning Maud Land. In all cases, the increase in melt was small with 0.1 mm w. eq. per day. Within the most recent decade melt got more extensive (+0.1 mm w. eq. per day) over George V, Oates and parts of Wilkes Land as well as over Getz and Sulzberger Ice Shelf. Smaller areas of strong surface melt in Pine Island Bay and Dronning Maud Land could only be observed during the first decade.

4.6 Zonal Wind

The Antarctic continent is circled by weak and irregular easterly winds driven by high pressure areas over the interior of the Antarctic continent created by cold and dry air. In case of a positive Southern Annular Mode (SAM) the easterlies weaken and the west wind drift shifts poleward. A positive SAM occurs when air pressure over the Antarctic Ice Sheet lowers but rises over the subtropical ocean. In the first decade, weakening easterlies ($\sim +0.5$ m/s) were observed along the Antarctic Peninsula, Bellingshausen and Amundsen Sea as well as in East Antarctica in the area of Shackleton Ice Shelf. Stronger easterlies occurred only at George V and Dronning Maud Land. Within the second decade the strengthening westerlies along East Antarctica expanded from Amery Ice Shelf to Victoria Land with up to +1 m/s. Also, the westerlies strengthened in the Bellingshausen Sea but weakened in the Amundsen Sea (see Figure 7). Strong dominating easterlies occurred along Dronning Maud Land with up to -0.75 m/s within the second decade.




Figure 6 Mean snowmelt over Antarctica with enlarged views of the Antarctic Peninsula. Snowmelt is given in mm water equivalent per day.




Figure 7 Zonal Wind (West to East) around the Antarctic continent in m/s for 1997-2008 and 2009-2018 compared to the long-term reference mean. Additionally, the difference in wind speed between both decades is illustrated. Positive shifts in zonal wind indicate stronger westerlies with the potential to cause upwelling of warm Circum Polar Deep Water (CDW).

4.7 Southern Annular Mode (SAM)

Changes in zonal wind direction are closely connected to fluctuations in SAM. Positive phases of SAM weaken the easterlies around the Antarctic continent and shift the westerlies poleward closer to the coastline. Phases of a positive SAM occur when air pressure over the Antarctic Ice Sheet lowers but rises over the subtropical ocean. Figure 9 shows the evolution of the SAM Index since 1960. The 5-year moving average of the SAM was positive since 1992 with positive values during the first and second decade. Annual peak events with exceptional high SAM occurred in 1997/98 and 2014/15 shown in blue (annual) and orange (summer only). Note the shift in peaks as the summer value for SAM averages values from December to February (December indicates the year).

Formatiert: Überschrift 2

Formatiert: Englisch (Vereinigte Staaten)

Formatiert: Block, Vom nächsten Absatz trennen




Figure 88 Southern Annular Mode (SAM) since 1960. Values are given for the annual SAM (blue), SAM during summer (orange) and a 5-year moving average for the annual SAM (black). The dashed grey lines indicate the investigated periods 1997/08 and

2009/18 as delimited by the available coastline data. For SAM during summer (Dec-Feb), the beginning of austral summer (hence December) indicates the year of the peak. Data: Marshall et al. (2003).

Formatiert: Standard

4.7.4.8 Correlation between Climate Variables and Calving Front Retreat

Potential drivers of calving front retreat were ~~identified~~diagnosed by correlating the analysed climate variables with the percentage of retreat/advance within each glacier/ice shelf basin. This allows for the assessment of a potential spatial relationship between calving front retreat and changes in environmental variables. The results of the Pearson correlation are displayed in ~~Figure 9~~Figure 8. Dark blue colors indicate a strong positive, dark red color a strong negative correlation. Stars indicate a significant Pearson correlation with stars for $p=0.05$ (*), 0.01 (**) and 0.001 (***). Correlations for retreat and advance are counterparts, hence correlations are of the same magnitude for each climate variable but with reversed sign. Overall, weak to moderate correlations with significance occur for relative summer sea surface temperature, absolute air temperature, snowmelt, relative zonal wind speed and sea ice days. The strongest positive linear relationship ($r=0.44$) exists between calving front retreat and relative sea surface temperature. Slightly weaker is the positive correlation between glacier and ice shelf front retreat and absolute air temperature on the ice shelf surface ($r_{\text{summer}}=0.18$, $r_{\text{winter}}=0.23$). A relatively more positive zonal wind (hence strengthening westerlies) correlates positively with calving front retreat ($r=0.30$) but the absolute strength of zonal winds does not. Decreases in sea ice days correlate positively with calving front retreat ($r_{\text{abs}}=0.33$, $r_{\text{rel}}=0.27$). The mean daily amount of snowmelt correlates weakly but significant with glacier and ice shelf front retreat ($r=0.17$). The correlation of the climate variables among each other reflects that they are closely linked to each other. Higher air ($r_{\text{summer}}=-0.26$, $r_{\text{winter}}=-0.36$) and sea surface temperatures ($r=0.44$) have a negative relationship to an increase in sea ice days. An increase in sea ice days negatively correlated with an increase in zonal winds ($r=0.31$). Stronger snowmelt correlates positively with warmer summer ($r=0.46$) and winter ($r=0.37$) air temperatures. An increase of zonal winds was positively related to decreasing summer air temperatures ($r_{\text{summer}}=-0.31$).




Figure 940 Correlation between glacier and ice shelf change and the analysed climate variables winter/summer air temperature, snowmelt, sea ice days, summer sea surface temperatures and zonal wind speed. Colour and circle size indicate the correlation coefficient. Stars indicate significance levels for $p=0.05$ (*), 0.01 (**) and 0.001 (***). S: summer, W: winter, rel: relative to 1982-1996, DJF: Dec-Feb.

5 Discussion

For the first time, this study presented circum-Antarctic glacier and ice shelf front change over two decades. The diagnosis through correlation indicated a spatial relationship between strengthening westerlies, higher air temperatures, intense snowmelt, a decrease in sea ice cover and rising sea surface temperature and calving front retreat. We want discuss if those external environmental factors were responsible for the observed glacier retreat or if internal glaciological forces were the key driver. To address this question, two major challenges arise. First, a significant correlation between calving front retreat/advance and climate variables cannot alone provide conclusive evidence of a causal link. Second, the retreat forced by environmental drivers has to be entangled from retreat caused through internal glaciological and external mechanical factors. To find solid evidence for drivers of glacier and ice shelf retreat, we will discuss observed glacier/ice shelf retreat on the basis of findings from previous studies in combination with measured changes in analyzed climate variables.

5.1 Antarctic Peninsula

Along the Antarctic Peninsula a decrease in ice shelf extent resulted mainly by the disintegration and retreat of Larsen B, Larsen C, Wilkins and Wordie Ice Shelf. The Wordie Ice Shelf disintegrated in a series of calving events beginning in the 1960s which were controlled by several pinning points (Friedl et al., 2018). The initial destabilization of the ice shelf was likely caused by a shift in flow which caused rift development at the ice rises and a collapse in 1989 (Walker and Gardner, 2017). Additional destabilisation occurred through basal melt induced thinning (Depoorter et al., 2013; Friedl et al., 2018; Rignot et al., 2013; Walker and Gardner, 2017). A later calving event accompanied by thinning and acceleration was associated with upwelling of warm CDW which was strong in the years 2008 to 2011 due to an exceptionally positive SAM (Walker and Gardner, 2017). In addition to the continuous thinning by basal melt, the retrograde bed in a subglacial deep trough destabilized the Wordie Ice Shelf and led to grounding line retreat in 2010/2011 (Friedl et al., 2018). For the major calving event of Wordie in 1989, we observed stronger zonal winds between 1997-2008 compared to the reference period (+0.62 m/s) and the most recent decade (+ 0.28 m/s) (2009-2018). This means that during the break up event the eastern wind direction weakened towards a more westerly direction. This change in wind-direction causes upwelling of warm CDW and enhances basal melt which is in line with the basal melt observed by the studies above. Additionally, we not only observed a peak in SAM during 2008 to 2011 as Walker and Gardner (2017) but also for the loss of the remaining ice tongues in summer 98/99. This strengthens the hypothesis that the retreat of Wordie Ice Shelf was influenced by environmental forcing through enhanced basal melt. Besides, we observed a decreasing sea ice coverage (-14 day compared to the reference mean) at Wordie Bay which has not been discussed in literature but might have further facilitated the destabilization of the ice shelf margins. Despite strong evidence by observed environmental forcing it shall not be neglected that the glaciers flowing into Marguerite Bay are still, at least partly, responding to the collapse initiated in the 1960s.

The Wilkins Ice Shelf also experienced a series of calving events likely caused by environmental and buoyancy forces (Braun and Humbert, 2009; Scambos et al., 2000). Scambos et al. (2000, 2009) linked the calving event in summer 1998 to hydrofracture due to atmospheric surface melt and ponding caused by atmospheric forcing. In contrast, Braun et al. (2009) speculate that basal melt thinned the boundaries and bending stresses from buoyancy forces caused the break-up in 1998. Additionally, the minimal extent of sea ice was discussed as potential factor for retreat (Lucchitta and Rosanova, 1998). Our observations confirm the decrease in sea ice days (-9 days), potential basal melt through enhanced zonal winds (+0.47 m/s), a peak in SAM, increased basal melt (+0.06 mm w. eq. per day), higher sea surface temperatures (+0.13°C) and increased air temperature (+0.58°C) within the first decade compared to the reference time period. This suggests that the calving event in 1998 was probably forced by a combination of several environmental drivers that destabilized and thinned the ice shelf. Whether these factors alone would have caused the retreat or the additional stresses through buoyancy as stated by Braun et al. (2009) were necessary to initiated the final calving event stay unresolved. A second break up event at Wilkins Ice Shelf occurred in three steps during February, May and June 2008 (Scambos et al., 2009) and the remaining ice bridge collapsed in

April 2009 likely due to mechanical external forcing by strong winds (Humbert et al., 2010). For the calving events in 2008, Scambos et al. (2009) found surface melt water as the main pre-condition for calving whereas Braun et al. (2009) excluded melt pond drainage as a driver and attribute the calving to fracture development and rift formation due to buoyancy forces and bending stresses through variable ice thickness. As described above the surface melt did only slightly increase between 1997-2008 compared to the reference period which strengthens the hypothesis by Braun et al. (2009). Therefore, we suggest that a combination of glaciological, environmental forced and mechanically forced processes caused the disintegration of Wilkins Ice Shelf.

The disintegration of Larsen B in 2002 is associated with increased surface melt causing enhanced fracturing (Khazendar et al., 2007; Rack and Rott, 2004) and widespread supraglacial lake drainage (Banwell et al., 2013; Leeson et al., 2020). Enhanced surface melt likely occurred as zonal winds were +0.33 m/s stronger and a positive SAM anomaly occurred. Positive SAM years with stronger westerlies are not associated with upwelling at the Larsen Ice Shelves but with warmer air temperatures in combination with warm winds and surface melt (Rack and Rott, 2004). Recent studies found that increases in foehn days in a series of years are related to positive SAM phases and significantly increase melt (Cape et al., 2015; Leeson et al., 2017). Interestingly, surface melt was high with up to 5 mm w. eq. per day but not more than during the reference period, solely the spatial distribution changed slightly. Mean air temperature between 1997-2008 increased with +0.16°C (within the uncertainties of the ERA5 data) during summer and no change in winter compared to the reference period but in the centre of the ice shelf increases of up to 0.49°C were observed during summer. The observed changes favour supraglacial lake formation as mentioned in literature. But whether the shortly after observed lake drainage is a cause or effect of ice shelf break up remains unclear (Leeson et al., 2020) and internal glaciological factor might also have played a role due to certain zones of weakness in the ice shelf (Khazendar et al., 2007).

In contrast stands the tabular calving event of iceberg A-68 (~5800 km²) from Larsen C Ice Shelf which accounted for almost the entire ice shelf extent decrease for the I-Ipp basin in 2009-2018. The calving event was predictable as a decade earlier a rift developed from a crevasse field at the Gipps ice rise and propagated further until July 2017 when the iceberg calved (Hogg and Gudmundsson, 2017). In the period of the Larsen C calving event, we could not observe any environmental forcing as lower amounts of melt (-0.2 mm w. eq. per day), not remarkable stronger zonal winds of +0.1 m/s and decreased summer air temperatures (-0.26°C) occurred compared to 1982-1996. This strengthens the hypothesis of a natural calving event as proposed by Hogg and Gudmundsson (2017). Nevertheless, slight negative thickness changes were observed by Paolo et al. (2015) between 1994 and 2012, which could indicate a future weakening of the Larsen C Ice Shelf. Larsen D forms the only exception of the strong retreating trend along the Antarctic Peninsula. This ice shelf neither experienced melt nor positive trends in zonal winds. Positive thickness changes indicate so far no potential weakening through basal melt (Paolo et al., 2015). The mass balance of Larsen D-G is, depending on the study, positive (Gardner et al., 2018) to slight negative but still much smaller than the huge negative mass balance of the rest-of-the-remaining Antarctic Peninsula (Rignot et al., 2019). George VI and Stange Ice Shelf were relatively stable during 1997 to 2008 and started to retreat in 2009-2018. Slightly strengthening westerlies (+0.25 m/s) but almost no melt (0.02 mm w. eq. per day) occurred. Summer sea surface temperatures increased by

+0.62°C and +0.38°C for George VI south and Stange ice shelves in the second decade compared to the reference period. Both ice shelves started to double the retreat rate during the second decade compared to the first. George VI is not believed to disintegrate rapidly (Holt et al., 2013). But recent developments might require to reconsider this assumption because calving front retreat, recently detected supraglacial lakes on the ice shelf surface (Dirscherl et al., 2020) and moderate basal melt (Paolo et al., 2015) -occurred.

5.2 WAIS

Glacier and ice shelf front retreat of the WAIS is clearly dominated at Pine Island Bay and at the biggest Antarctic ice shelves Ross and Ronne. Ross Ice Shelf lost an entire area of ~14.000 km² between 1997 and 2008 which can be completely attributed to tabular iceberg calving of B15-B18 in 2000 and B19 in 2002 (in total ~18.000 km²) (Budge and Long, 2018; MacAyeal et al., 2001). Note that the difference in iceberg size versus retreated area arises from the rough estimation of iceberg area (width multiplied by length) as well as the measurement over one decade where re-advance occurred after the calving event. The same applies for Ronne Ice Shelf where the entire area (~12.400 km²) was lost by tabular iceberg calving of A38/39 (1998) and A43/44 (2000) with an area of ~17.500 km² (Budge and Long, 2018; MacAyeal et al., 2001; Wuite et al., 2019). Even though the calved icebergs were of notable size (including the largest ever recorded iceberg B15) those calving events were not exceptional (MacAyeal et al., 2001). The Ross West Ice Shelf already experienced such a calving event before 1962, where the front had about the position of 2004 after the maximum extent around 1997 (Ferrigno et al., 2007). Also the maximum extent of the Ronne Ice Shelf was in 1997 and its minimum extents occurred in 1974 and 2004 (Ferrigno et al., 2005). Measurements of environmental conditions substantiate a natural calving event within the calving cycle of those ice shelves. We could neither observe a reduction of sea ice nor increases in sea surface temperature (0.02 °C) and zonal winds (-0.15 m/s). Only strong melt (0.5 mm w. eq. per day) occurred on the calving area on Ronne Ice Shelf but even higher melt occurred between 1982 and 1996 in the same area. It should be noted that Ronne-Filchner Ice Shelf might be prone to changing environmental conditions in the future as recent studies predict that the Ronne-Filchner Ice Shelf will be affected by increasing basal melt through changing sea ice conditions and wind direction (Darelius et al., 2016). A completely different setting occurs in the Pine Island Bay where Pine Island and Thwaites Glacier showed strong retreat rates with 40±0.15 km²/yr and 288±0.77km²/yr between 1997 and 2018. The instability and unstoppable retreat of both glaciers has been suggested by several studies (Joughin et al., 2014; Milillo et al., 2019; Parizek et al., 2013; Rignot et al., 2014). The deep marine basin in combination with a retrograde slope creates an increased sensitivity to basal melt by ocean forcing. Our climatological analysis indicates potential upwelling of CDW through strengthening westerlies with +0.28 m/s in 1997/2008 and +0.25 m/s in 2009/2018 compared to the reference mean. Additionally, sea surface temperatures raised by 0.28°C at the front of Pine Island but only 0.11°C at Thwaites Glacier from the first to the second decade. Notable changes in sea ice days or snowmelt were not observed. Especially the sea surface temperature is important for the stability of the Thwaites Glacier tongue as it is currently stabilized by fast ice persistency and the acceleration of the ice tongue is associated with changing

500 ocean conditions (Miles et al., 2020). It has to be considered that the initial start of destabilisation in Pine Island Bay occurred
previous to our observation period as mass loss exists since 1979 (Rignot et al., 2019), new rifted areas created in the beginning
of the 1990s (Bindshadler, 2002; Rignot, 2002) and basal melt by ocean forcing exists at least since 1994 (Jacobs et al., 2011).
Once marine ice sheet instability is initiated changes in ocean forcing can no longer directly be linked to ice flow and hence
calving front position (Christianson et al., 2016). Moreover, as soon as the maximum ice thickness at the grounding line is
505 reached through grounding line retreat on a retrograde bed no stable terminus is possible anymore and calving-induced frontal
retreat has to be expected which would no longer be coupled to environmental forced basal melt (Bassis and Walker, 2012).
Consequently, it cannot be ruled out that the measured frontal retreat over the last two decades is actually a response to earlier
destabilisation by ocean forcing. Nevertheless, our observations confirm that Pine Island and Thwaites Glacier are still exposed
to ocean forcing.

510 Besides the fast changing Amundsen Sea Sector, the Bellingshausen Sea at Abbot Ice Shelf is less studied but also vulnerable
to CDW forcing (Christie et al., 2016). Just recently, the velocity at the grounding line increased (+20%, 2007-2014), a dynamic
thinning signal evolved which is related to ice dynamics and lower snowfall (Chuter et al., 2017) and minor grounding line
retreat exists (Christie et al., 2016). The recently dynamic thinning signal observed by Chuter et al. (2017) fits to the glacier
front retreat at Abbot Ice Shelf observed since the second decade where westerlies increased by + 0.24 m/s from the first to
515 the second decade indicating vulnerability to upwelling CDW. Other variables did not change significantly. Along Marie Byrd
Land the Getz Ice Shelf experienced strong grounding line retreat between 2003-2008 due to ocean forcing which slowed
down afterwards (2010-2015) (Christie et al., 2018). This pattern in ocean forcing is visible in our data as zonal winds reduced
by -0.41 m/s from the first to the second decade. Interestingly, this is not apparent in the glacier front movement as glacier
front retreat and advance was balanced between 1997 and 2008 but advance reduced from 103 km²/yr to 73 km²/yr in the
520 second decade. No clear signal in frontal retreat could be explained by several pinning points stabilizing the ice shelf (Christie
et al., 2018) and the low glacier flow velocity (Mouginot et al., 2019). We propose that those stabilized fronts require longer
ocean forcing to respond with frontal retreat. The most stable ice shelves along West Antarctica are Sulzberger and Nickerson
Ice Shelf because a buffer to ocean forcing exists by the bathymetry of the continental shelf (Christie et al., 2018). Additionally,
they belong to the cooler Ross Sea (Rignot et al., 2013). No increase in potential environmental drivers was observed over the
525 last two decades which further strengthens the hypothesis of stable ice shelves not affected by environmental forcing.

5.2.5.3 EAIS

530 The EAIS was long believed to represent the invulnerable part of Antarctica with a stable discharge over the last years (Gardner
et al., 2018). Recent mass balance estimations show that East Antarctica is losing mass since 1979 with an increasing loss
since 1999 (Rignot et al., 2019). This stands in contrast to altimeter measurements recording an almost stable mass balance for
the EAIS since 1992 (IMBIE, 2018). In contrast to the calving front retreat of West Antarctica and the Antarctic Peninsula,
glacier and ice shelf front advance dominated over the last two decades in East Antarctica. The regional differences are

discussed in the following. In Victoria Land, previous studies could not identify a trend in calving front location change and attributed observed changes to glacier size, terminus type and local topographical settings (Fountain et al., 2017; Lovell et al., 2017). Our analysis strengthens the hypothesis that non-climatic drivers regulated calving front change along Victoria Land because no noticeable changes in sea ice, wind direction, snow melt and ocean surface temperatures were measured between 1997-2018 compared to the reference time period. The only noticeable change was the increase of summer (+ 1.1°C) and winter (+0.71°C) air temperatures over Mariner and Tucker Glacier during the second decade compared to 1982-1996 which did not influence the slight advance of Mariner Glacier and stable front of Tucker Glacier.

Oats and George V Land have experienced a negative mass balance since 1999 (Rignot et al., 2019) but did not show a trend in calving front position change. The position of fronts was balanced over the last two decades with switches between advance and retreat. A previous study by Lovell et al. (2017) noticed the potential influence of a decreased sea ice cover on the terminus position but as we did not measure a significant change in sea ice nor wind direction, sea surface temperature and melt the observed frontal changes cannot be connected to environmental forcing. Winter air temperature cooled by up to 1°C and summer air temperature by up to 0.49°C along the coast but frontal advance and retreat occurred. The only noticeable retreat occurred at Mertz Glacier tongue which was caused by mechanical forcing due to an iceberg collision in 2010 (Giles, 2017). The mass loss of East Antarctica is dominated by the loss in Wilkes Land being the only basin in EAIS losing mass since 1979 (Rignot et al., 2019) where the majority of glaciers retreated between 2000 and 2012. This contrasting retreat compared to the rest of EAIS was linked to environmental forcing by a decrease in sea ice days (Miles et al., 2016). For Wilkes Land, we observed a simultaneous retreat in glacier fronts between 2009 and 2018 but could not link it to a decrease in sea ice days. This might be due to the fact that during 1997-2008 sea ice days were only ~ 5 days less vs. ~34 days between 2000 and 2012, as reported by Miles et al. (2016). This suggests that only extreme reductions in sea ice can be related to glacier retreat, implying that sea ice alone cannot be the only explanation for retreat along Wilkes Land. We propose that upwelling CWD weakened the glaciers by basal melt as a very strong positive tendency in zonal wind with up to +0.44 m/s was observed in the second (but not in the first) decade compared to the long-term reference mean. All other measured variables did not change significantly over the observation time period.

A strong retreat within the second decade occurred also at Glenzer Glacier next to Wilkes Land. The retreat is not yet mentioned in literature but satellite imagery reveals that an important pinning point was lost in 2004 and a rift started to propagate and lead to a break up event in April 2020. We suggest that the initial destabilization and loss of the pinning point was caused by an increase in zonal winds (+0.32 m/s) and enhanced surface melt (up to +0.4 mm w. eq. /day) during the first decade. Increased zonal winds persisted during the second decade (+1.23 m/s) indicating further ocean forcing. Nevertheless, the more recent calving event occurred due to a rift that developed over several years. If environmental drivers played a role or the rift developed due to ice flow acceleration initiated by the previous lost pinning point has to be investigated.

The fronts of Shackleton and West Ice Shelf advanced within the last two decades. Both ice shelves are stabilized by several pinning points. Recent studies found that supra glacial lake formation on the Shackleton Ice Shelf is closely connected to short lived high magnitude snow melt events (Arthur et al., 2020). So far, the supra glacial lake coverage on the ice shelf is still

small (< 1%) but it is proposed that widespread ponding and hydrofracture is likely to increase (Arthur et al., 2020). For both ice shelves we observed warmer air temperatures within the first and colder during the second decade (compared to 1982-1996). Snow melt increased only punctually along the margins of both ice shelves (up to 0.3 mm/day). The strongest environmental forcing we observed within the second decade where zonal winds strengthened by +0.97 m/s and +1.2 m/s for the Shackleton and West Ice Shelf, respectively. The only indicator that weakening by basal melt occurred indicates the collapse of iceberg D-15 which calved in 1992 but re-attached to the West Ice Shelf until it split-up in 2015 (Walker et al., 2019). Shackleton shows first signs of instability as ice flow velocities sped up by 43 % between 1957-2016 (Rignot et al., 2019). Further investigations are necessary to prove that the measured enhanced zonal winds caused enhanced basal melt at both ice shelves or if other factors caused acceleration and the break-up of the iceberg.

The front of Amery Ice Shelf gradually advanced between 1997 and 2018. The environmental conditions changed with strengthening westerlies by + 0.26 m/s within the second decade and increased snow melt (+ 0.23 mm w. eq. per day) on the northern part of Amery ("Loose Tooth" region) over the last two decades but a decrease (- 0.1 mm w. eq. per day) in the southern part. The stability of the ice shelf is confirmed by velocity measurements where no speed-up occurred since the 1970s (Rignot et al., 2019) and basal melt rates are low (Paolo et al., 2015) suggesting that the westerlies did not strengthen enough to cause upwelling CDW. In 2019, the tabular iceberg D-28 calved from Amery Ice Shelf. This calving front retreat was predicted by Fricker et al. (2002) based on the observed rift propagation and regular calving cycle of the ice shelf. Still, it remains unclear if the observed increase in surface melt influenced the rift propagation in the Loose Tooth region. Prior to 2006, the rift propagation of Amery Ice Shelf was not influenced by environmental forcing but the authors did not exclude the potential influence of surface melt if the mean air temperature would raise above zero (Bassis et al., 2008).

The area loss of Enderby Land is mostly attributed to the changes in Shirase Bay. Here, very contrasting changes appear where the Prince Harald Ice Shelf retreats and the Shirase Glacier advances and vice versa. This was already mentioned by Jezek (2002). The only change in environmental conditions in this area occurred between 1997 and 2008 with an up to 1 mm w. e. melt per day over Prince Edward Ice Shelf compared to the reference period. For Shirase Glacier no increases in melt were observed. In Shirase Bay, rather the specific glacier morphology and bed topography influence the calving than environmental factors (Nakamura et al., 2007).

Dronning Maud Land is dominated by advance over the entire study period. Historical records reveal that along Dronning Maud Land phases of retreat and advance are typical with bigger calving events between 1963 and the 70s. Retreat continued until the 90s followed by a phase of advance (Kim et al., 2001; Miles et al., 2016). Mass balance estimates also confirm phases between mass loss and mass gain since 1979 with no evidence for long term change (Rignot et al., 2019) and basal melt rate estimates confirm the relatively stable state of the ice shelves (Paolo et al., 2015; Rignot et al., 2013). The environmental conditions along Dronning Maud Land changed only little in regards to sea ice days (-5 days in the first decade) and constant zonal winds between Shirase Bay and Fimbul Ice Shelf and more negative winds (~ 0.35 m/s) between Jelbart and Filchner Ice Shelf (1997-2018 compared to reference mean). The only exceptional change was observed for air temperatures which increased between Lazarev and Filchner Ice Shelf between +0.6-1.6 °C in summer and 0.9-1.8 in winter within the second

decade compared to the reference mean. Despite this remarkable increase in air temperature ice shelf advance persisted which is not exceptional as mean air temperatures were far below the freezing point and surface melt did not increase.

5.4 A Circum-Antarctic Perspective

The above examples illustrated that environmental conditions changed very differently along the Antarctic coastline over the last two decades. Strengthening westerlies occurred along with frontal retreat at the Western Antarctic Peninsula, Larsen B Ice Shelf, Pine Island Bay, Abbot Ice Shelf and Wilkes Land but advance at Shackleton and West Ice Shelf. This can have several reasons. Strengthening zonal winds are a proxy for upwelling CDW (Spence et al., 2014) but further studies are needed to assess the relationship between the magnitude of strengthening westerlies and the amount of caused basal melt. Only a small increase in westerlies could be enough to increase basal melt considerably because basal melt has a nonlinear relationship with ocean temperature and can quadruple by only slight ocean warming (Jenkins et al., 2018). Additionally, the forcing by zonal winds is less strong in areas where CDW upwelling is less efficient due to strong tidal forcing (Hazel and Stewart, 2019; Stewart et al., 2018) and the local topography regulates the amount of CDW accessing the cavities of an ice shelf (Dutrieux et al., 2014). Moreover, the duration of required environmental forcing to cause destabilization of ice shelves solely by basal melt very likely exceeds the time scales addressed in this study. In addition, Shackleton and West Ice Shelf only experienced potential forcing by one driver but it was found that the forcing by several environmental drivers had the greatest impact on calving front retreat in Greenland (Cowton et al., 2018; Howat et al., 2008) which likely also applies for Antarctica. Further studies are necessary to assess the amount of basal melt for individual ice shelves by strengthening westerlies.

Our study found a strong link between increased sea surface temperature and the occurrence of calving front retreat. The measured changes in sea surface temperature were between 0.15 to 0.6°C per decade which is still below the trend uncertainty stability of 0.01 K per year (Embury, 2019). The influences of warmer sea surface temperatures are twofold. Warmer sea surface temperatures lead to the melt of fast ice which destabilizes glacier margins (Larour, 2004) and could result in retreat. Another explanation proposes that warm surface water increases melt at the waterline which creates an overhanging ice cliff that is more likely to collapse (Mosbeux et al., 2020). Our results suggest, that increased sea surface temperatures weakened the margins in the Bellingshausen and Amundsen Sea where highest increases in sea surface temperature were observed. Further destabilization could arise as through the presence of marginal weakening rifts become unstable (Lipovsky, 2020).

Marginal weakening can also arise from a decrease in sea ice days which occurred along the Western Antarctic Peninsula, the Bellingshausen and Amundsen Sea and in Wilkes Land. Decreasing sea ice days occurred with glacier/ice shelf retreat but only if the average reduction per year was on average over 10 days within the decade. This fact might become important in the future. Recently, the turning point of an increasing Antarctic sea ice extent was reached (Ludescher et al., 2019; Parkinson, 2019).

Relative changes in mean air temperature could not be identified as a direct driver for calving front retreat, even though increases of up to 2°C per decade were measured in some coastal areas (e.g. Dronning Maud Land, Victoria Land) which is beyond the uncertainty of the ERA5 air temperature data. This suggests that mean air temperature over a decade is not an

Formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Englisch (Vereinigtes Königreich)

Feldfunktion geändert

Formatiert: Schriftart: (Standard) Times New Roman

Formatiert: Englisch (Vereinigtes Königreich)

635 appropriate way to assess the effect of air temperature on calving front change because the relationship between air temperature
driven surface melt and hydrofracture is known to destabilize ice shelves and can cause glacier front retreat (Arthur et al.,
2020; Banwell et al., 2013; Leeson et al., 2020). More suitable would be the assessment of the amount of positive degree days
and temperature extreme events directly influencing surface melt on the ice shelf with more accurate and higher-resolution
melt data. For example, catchment wide melt and the resulting runoff had a higher impact on the retreat of Greenlandic glaciers
than local air temperatures (Cowton et al., 2018).

640 Howat et al. (2008) found that peak events in air temperature and sea surface temperature can initialize glacier retreat. This
study found that the absolute amount of snowmelt correlated significantly with glacier and ice shelf retreat but not relative
changes in snowmelt. The detailed comparison of snowmelt with calving front retreat revealed strong melt during the decade
of the break up event (e.g. Larsen B, Wilkins Ice Shelf). Nevertheless, snowmelt was often higher during the reference period
than during the actual calving event which might indicate that longer phases of melt weaken the ice shelf over a longer time
645 period (Leeson et al., 2017) and the intensity of surface melt events are of higher importance than just the presence of melt
(Arthur et al., 2020). Moreover, the low resolution of melt data did not always match the actual calving front. We assume that
the use of a not up to date coastline product (e.g. at Larsen B) caused the inaccuracies over the ice shelf margins. Additionally,
no information on the accuracy of the snowmelt product is provided which makes the considered reanalysis data prone to
potentially large biases. We suggest a more detailed study on glacier and ice shelf front retreat in combination with accurate
650 high resolution surface melt data (e.g. satellite-based surface melt estimates as published by Trusel et al. (2013)) to capture all
necessary surface hydrology processes, peak events and an accurate metric for snowmelt. For example, not the amount of melt
but much more the effect of ponding on the ice shelf influences its vulnerability (Joughin and Alley, 2011). Moreover, the
hydrological system of the ice shelf has to be considered as river formation can act as a stabilization by exporting meltwater
to the ocean (Banwell, 2017) and hydrology processes such as lake drainage occurred prior to break up events (Leeson et al.,
655 2020). Those complex surface hydrology processes require much more detailed studies considering the combination of factors
mentioned above and not solely mean surface melt.

For the future assessment on environmental driven calving front change, it has to be considered that the environmental drivers
with the potential to drive calving front retreat are closely connected to positive SAM years. Extreme peaks in positive SAM
can be connected to ice shelf disintegration as has been shown for Wordie Ice Shelf (Walker and Gardner, 2017) and further
660 peaks in positive SAM occurred during the disintegration of Larsen B Ice Shelf and the first break-up event of Wilkins Ice
Shelf. A positive SAM increases sea surface temperatures and air temperatures along AP and WAIS. It also influences the sea
ice cover, snowmelt and foehn effect and enhances westerly winds along the Antarctic coastline (Cape et al., 2015; Kwok and
Comiso, 2002; Marshall, 2007; Tedesco and Monaghan, 2009; Verdy et al., 2006). For the future it should be considered, that
glacier and ice shelf retreat forced by environmental drivers will be even more likely as rising greenhouse gases and ozone
665 depletion will cause more positive phases of SAM (Paeth and Pollinger, 2010; Wang et al., 2014) and further strengthen the
identified environmental forces on calving front retreat.

Formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Englisch (Vereinigtes Königreich)

Formatiert: Englisch (Vereinigtes Königreich)

Feldfunktion geändert

Formatiert: Englisch (Vereinigtes Königreich)

The identification of environmental drivers for calving front retreat presented a major challenge and some questions remain unresolved due to study limitations. The connection between atmospheric forcing and calving front retreat could not be entirely revealed as average surface melt and air temperatures did not allow for an analysis regarding threshold behavior (e.g. above zero-degree days), peak events and surface hydrological processes. Moreover, once ice shelf destabilization is initiated the response in frontal change might no longer linearly be connected to forcing and decouple from external forcings as known for the marine ice sheet instability (Feldmann and Levermann, 2015; Joughin et al., 2014; Robel et al., 2019). As a result, the natural calving cycle and responses to environmental forcing may prevail on longer time scales than the ones addressed in this study due to limited data coverage. Only the detailed analysis of observed environmental changes from this study in combination with information from previous studies on individual ice shelves and glaciers allowed the identification of key drivers for calving front retreat being not solely environmentally driven but a combined result of complex interactions between internal ice dynamics, geometry, external mechanical forcing and environmental drivers.

6 Conclusion

For the first time, we present a circum-Antarctic record of calving front changes over the last two decades. Overall, the extent of the Antarctic Ice Sheet decreased $-29618 \pm 1,19329 \text{ km}^2$ between 1997 and 2008 gained an area of $+7108 \pm 14441,029 \text{ km}^2$ between 2009-2018. Glacier and ice shelf front retreat concentrated along the Antarctic Peninsula and West Antarctica. The only East Antarctic coastal sector experiencing simultaneous calving front retreat of several glaciers was Wilkes Land in 2009-2018. The largest proportion of calving originated from Ross West and Ronne Ice Shelf being responsible for 75 % of the West Antarctic loss during 1997 and 2008 but was found to be in the range of their natural calving cycle. Decreasing sea ice days, strengthening westerlies, intense snowmelt and increasing sea surface temperatures were identified as enabling driving forces for glacier and ice shelf front retreat along the Antarctic Peninsula, West Antarctica and Wilkes Land. In contrast, changes in mean air temperature were not an appropriate measure for atmospheric forcing and rather extreme temperature events, threshold behaviour and surface hydrological processes should be considered for future analysis. ~~relative increases in mean air temperatures could not be linked to calving front retreat.~~ Further studies should assess whether record high air temperatures can also trigger glacier/ice shelf retreat as it has been observed for Greenland. Snowmelt was found to be a strong driver of calving front retreat at the Antarctic Peninsula (up to 5 mm w. eq. per day) but more accurate data on surface melt and surface hydrology is needed to assess the influence of melt in more detail. Increased sea surface temperatures with up to 0.62°C were observed along the Bellingshausen and Amundsen Sea weakening the glacier margins. Between 1997 and 2018, sea ice days decreased most along the Western Antarctic Peninsula with up to -25 days, the Bellingshausen Sea (-10 days) and Amundsen Sea (-5 days). Strengthening westerlies affected ice shelves along the Western Antarctic Peninsula (up to $+0.54 \text{ m/s}$) and West Antarctica ($+0.28\text{-}0.41 \text{ m/s}$) but also East Antarctica ~~at-in~~ Wilkes ($+0.44 \text{ m/s}$) and Queen Mary Land (up to $+1.23 \text{ m/s}$). Despite those changes in environmental conditions the influence of internal ice dynamics, geometry and external mechanical forcings play a crucial role and must not be neglected when assessing drivers of calving front retreat.

by causing upwelling of warm CWD and resulting basal melt. The magnitude of required forcing to immediately trigger calving front retreat was difficult to assess as the regional setting (e.g. bed topography, glacier morphology) as well as the combination and duration of driving forces are likewise important for assessing glacier and ice shelf front retreat. The All identified assessed environmental drivers are closely connected to positive phases of SAM which occurred over the last two decades with local maxima forcing stronger retreat. Rising CO₂-emissions and ozone depletion will further enhance positive phases of SAM putting additional pressure on glaciers and ice shelves. To better assess the vulnerability of glaciers and ice shelves in the future, it is essential to better understand melt, ponding and runoff processes on the ice shelf surface impacting calving front retreat to an unknown extent. Equally important is the understanding of changing wind conditions and their impact on upwelling CWD and basal melt. Another interesting issue related to glacier and ice shelf changes across Antarctica pertains to their specific time scales and response times. A shortcoming of our analysis is the restriction to three snapshots of coastline data that has delimited the design of this study. Although our study has revealed that the Antarctic cryosphere is subject to tremendous changes in recent decades, it is conceivable that our results are at least partly biased by strong inter-annual variability of all considered climate and cryospheric variables. In addition, natural cycles and the effects of man-made global warming may prevail on longer time scales than the ones addressed in this study – the ones that are also characterized by a reasonable data coverage.

Data availability

The Antarctic coastline 2018 is available at the EOC Geoservice of the German Aerospace Center <LINK>. Additional coastlines used for this analysis are available at <https://nsidc.org/data/NSIDC-0103/versions/2> (Radarsat 1997) and <https://nsidc.org/data/NSIDC-0593/versions/1> (MODIS 2009). The supplementary material provides a table on calving front change for each assessed glacier/ice shelf in m/yr and km²/yr. Used climate data is available at the Copernicus Climate Data Store (cds.climate.copernicus.eu) and the Ocean and Sea Ice Satellite Application Facility (www.osi-saf.org/?q=content/sea-ice-products).

Acknowledgements

We thank the Copernicus Climate Data Store for providing the ERA5 climate data and the Ocean and Sea Ice Satellite Application Facility (OSI SAF) for providing the sea ice concentration datasets. In addition, we thank the European Union Copernicus program providing Sentinel-1 data accessible via NASA Alaska Satellite Facility (ASF) and G. J. Marshall for granting open access to the Southern Annular Mode (SAM) Index. Finally, we would like to thank the editor, Eleri Evans, Brad Lipovsky and one anonymous reviewer for their comprehensive comments and suggestions on our manuscript.

Formatiert: Hochgestellt

References

- 730 Alley, R. B., Horgan, H. J., Joughin, I., Cuffey, K. M., Dupont, T. K., Parizek, B. R., Anandakrishnan, S. and Bassis, J.: A simple law for ice-shelf calving, *Science*, 322(5906), 1344–1344, 2008.
- Arthur, J. F., Stokes, C. R., Jamieson, S. S. R., Carr, J. R. and Leeson, A. A.: Distribution and seasonal evolution of supraglacial lakes on Shackleton Ice Shelf, East Antarctica, *The Cryosphere*, 14(11), 4103–4120, <https://doi.org/10.5194/tc-14-4103-2020>, 2020.
- 735 Åström, J. A., Vallot, D., Schäfer, M., Welty, E. Z., O’Neel, S., Bartholomäus, T. C., Liu, Ya., Riikilä, T. I., Zwinger, T., Timonen, J. and others: Termini of calving glaciers as self-organized critical systems, *Nat. Geosci.*, 7(12), 874–878, 2014.
- Banwell, A.: Glaciology: Ice-shelf stability questioned, *Nature*, 544(7650), 306–307, 2017.
- Banwell, A. F., MacAyeal, D. R. and Sergienko, O. V.: Breakup of the Larsen B Ice Shelf triggered by chain reaction drainage of supraglacial lakes, *Geophys. Res. Lett.*, 40(22), 5872–5876, <https://doi.org/10.1002/2013GL057694>, 2013.
- 740 Bassis, J. N.: The statistical physics of iceberg calving and the emergence of universal calving laws, *J. Glaciol.*, 57(201), 3–16, 2011.
- Bassis, J. N. and Walker, C. C.: Upper and lower limits on the stability of calving glaciers from the yield strength envelope of ice, *Proc. R. Soc. -Math. Phys. Eng. Sci.*, 468(2140), 913–931, <https://doi.org/10.1098/rspa.2011.0422>, 2012.
- Bassis, J. N., Fricker, H. A., Coleman, R. and Minster, J.-B.: An investigation into the forces that drive ice-shelf rift propagation on the Amery Ice Shelf, East Antarctica, *J. Glaciol.*, 54(184), 17–27, 2008.
- 745 Baumhoer, Dietz, A., Dech, S. and Kuenzer, C.: Remote Sensing of Antarctic Glacier and Ice-Shelf Front Dynamics—A Review, *Remote Sens.*, 10(9), 1445, <https://doi.org/10.3390/rs10091445>, 2018.
- Baumhoer, C. A., Dietz, A. J., Kneisel, C. and Kuenzer, C.: Automated Extraction of Antarctic Glacier and Ice Shelf Fronts from Sentinel-1 Imagery Using Deep Learning, *Remote Sens.*, 11(21), 2529, <https://doi.org/10.3390/rs11212529>, 2019.
- 750 Belmonte Rivas, M. and Stoffelen, A.: Characterizing ERA-Interim and ERA5 surface wind biases using ASCAT, *Ocean Sci.*, 15(3), 831–852, <https://doi.org/10.5194/os-15-831-2019>, 2019.
- Benn, D. I., Hulton, N. R. J. and Mottram, R. H.: ‘Calving laws’, ‘sliding laws’ and the stability of tidewater glaciers, *Ann. Glaciol.*, 46, 123–130, <https://doi.org/10.3189/172756407782871161>, 2007a.
- Benn, D. I., Warren, C. R. and Mottram, R. H.: Calving processes and the dynamics of calving glaciers, *Earth-Sci. Rev.*, 82(3), 143–179, 2007b.
- 755 Bindschadler, R. A.: History of lower Pine Island Glacier, West Antarctica, from Landsat imagery, *J. Glaciol.*, 48(163), 536–544, <https://doi.org/10.3189/172756502781831052>, 2002.
- Brandt-Kreiner, M., Lavelle, J., Tonboe, R., Howe, E., Laverigne, T., Killie, M. A., Sorensen, A., Eastwood, S. and Neuville, A.: Global Sea Ice ConcentrationClimate Data RecordValidation Report. OSI-450 and OSI-430-b. http://osisaf.met.no/docs/osisaf_cdop3_ss2_valrep_sea-ice-conc-climate-data-record_v1p1.pdf, 2019.
- 760 Braun, M. and Humbert, A.: Recent Retreat of Wilkins Ice Shelf Reveals New Insights in Ice Shelf Breakup Mechanisms, *IEEE Geosci. Remote Sens. Lett.*, 6(2), 263–267, <https://doi.org/10.1109/LGRS.2008.2011925>, 2009.

- Braun, M., Humbert, A. and Moll, A.: Changes of Wilkins Ice Shelf over the past 15 years and inferences on its stability, *The Cryosphere*, 3(1), 41–56, <https://doi.org/10.5194/tc-3-41-2009>, 2009.
- 765 Budge, J. S. and Long, D. G.: A Comprehensive Database for Antarctic Iceberg Tracking Using Scatterometer Data, *IEEE J. Sel. Top. Appl. Earth Obs. Remote Sens.*, 11(2), 434–442, <https://doi.org/10.1109/JSTARS.2017.2784186>, 2018.
- C3S: Copernicus Climate Change Service. ERA5: Fifth generation of ECMWF atmospheric reanalyses of the global climate, Copernic. Clim. Change Serv. Clim. Data Store CDS <https://cds.climate.copernicus.eu/cdsapp#!/home>, last access: 5 February 2020, 2017.
- 770 Cape, M. R., Vernet, M., Skvarca, P., Marinsek, S., Scambos, T. A. and Domack, E.: Foehn winds link climate-driven warming to ice shelf evolution in Antarctica: Foehn winds link climate driven warming, *J. Geophys. Res. Atmospheres*, 120(21), 11,037–11,057, <https://doi.org/10.1002/2015JD023465>, 2015.
- Catania, G. A., Stearns, L. A., Sutherland, D. A., Fried, M. J., Bartholomaeus, T. C., Morlighem, M., Shroyer, E. and Nash, J.: Geometric Controls on Tidewater Glacier Retreat in Central Western Greenland, *J. Geophys. Res. Earth Surf.*, 123(8), 2024–2038, <https://doi.org/10.1029/2017JF004499>, 2018.
- 775 Christianson, K., Bushuk, M., Dutrieux, P., Parizek, B. R., Joughin, I. R., Alley, R. B., Shean, D. E., Abrahamsen, E. P., Anandakrishnan, S., Heywood, K. J., Kim, T.-W., Lee, S. H., Nicholls, K., Stanton, T., Truffer, M., Webber, B. G. M., Jenkins, A., Jacobs, S., Bindschadler, R. A. and Holland, D. M.: Sensitivity of Pine Island Glacier to observed ocean forcing: PIG response to ocean forcing, *Geophys. Res. Lett.*, 43(20), 10,817–10,825, <https://doi.org/10.1002/2016GL070500>, 2016.
- 780 Christie, F. D. W., Bingham, R. G., Gourmelen, N., Tett, S. F. B. and Muto, A.: Four-decade record of pervasive grounding line retreat along the Bellingshausen margin of West Antarctica: West Antarctic Ice Retreat, *Geophys. Res. Lett.*, 43(11), 5741–5749, <https://doi.org/10.1002/2016GL068972>, 2016.
- Christie, F. D. W., Bingham, R. G., Gourmelen, N., Steig, E. J., Bisset, R. R., Pritchard, H. D., Snow, K. and Tett, S. F. B.: Glacier change along West Antarctica's Marie Byrd Land Sector and links to inter-decadal atmosphere–ocean variability, *The Cryosphere*, 12, 2461–2479, [https://doi.org/Christie, Frazer D.W.; Bingham, Robert G.; Gourmelen, Noel; Steig, Eric J.; Bisset, Rosie R.; Pritchard, Hamish D.; Snow, Kate; Tett, Simon F.B.. 2018 Glacier change along West Antarctica's Marie Byrd Land Sector and links to inter-decadal atmosphere–ocean variability. The Cryosphere, 12 \(7\). 2461-2479. https://doi.org/10.5194/tc-12-2461-2018 <https://doi.org/10.5194/tc-12-2461-2018>](https://doi.org/Christie, Frazer D.W.; Bingham, Robert G.; Gourmelen, Noel; Steig, Eric J.; Bisset, Rosie R.; Pritchard, Hamish D.; Snow, Kate; Tett, Simon F.B.. 2018 Glacier change along West Antarctica's Marie Byrd Land Sector and links to inter-decadal atmosphere–ocean variability. The Cryosphere, 12 (7). 2461-2479. https://doi.org/10.5194/tc-12-2461-2018 <https://doi.org/10.5194/tc-12-2461-2018>), 2018.
- 785 Chuter, S. J., Martín-Español, A., Wouters, B. and Bamber, J. L.: Mass balance reassessment of glaciers draining into the Abbot and Getz Ice Shelves of West Antarctica: Getz and Abbot Mass Balance Reassessment, *Geophys. Res. Lett.*, 44(14), 7328–7337, <https://doi.org/10.1002/2017GL073087>, 2017.
- 790 Cook, A. J. and Vaughan, D. G.: Overview of areal changes of the ice shelves on the Antarctic Peninsula over the past 50 years, *The Cryosphere*, 4(1), 77–98, <https://doi.org/10.5194/tc-4-77-2010>, 2010.
- 795 Cook, A. J., Holland, P. R., Meredith, M. P., Murray, T., Luckman, A. and Vaughan, D. G.: Ocean forcing of glacier retreat in the western Antarctic Peninsula, *Science*, 353(6296), 283–286, 2016.
- Cowton, T. R., Sole, A. J., Nienow, P. W., Slater, D. A. and Christoffersen, P.: Linear response of east Greenland's tidewater glaciers to ocean/atmosphere warming, *Proc. Natl. Acad. Sci.*, 115(31), 7907–7912, <https://doi.org/10.1073/pnas.1801769115>, 2018.
- 800 Darelius, E., Fer, I. and Nicholls, K. W.: Observed vulnerability of Filchner-Ronne Ice Shelf to wind-driven inflow of warm deep water, *Nat. Commun.*, 7(1), 12300, <https://doi.org/10.1038/ncomms12300>, 2016.

- De Angelis, H. and Skvarca, P.: Glacier Surge After Ice Shelf Collapse, *Science*, 299(5612), 1560–1562, <https://doi.org/10.1126/science.1077987>, 2003.
- De Rydt, J., Gudmundsson, G. H., Rott, H. and Bamber, J. L.: Modeling the instantaneous response of glaciers after the collapse of the Larsen B Ice Shelf, *Geophys. Res. Lett.*, 42(13), 5355–5363, <https://doi.org/10.1002/2015GL064355>, 2015.
- 805 Depoorter, M. A., Bamber, J. L., Griggs, J. A., Lenaerts, J. T. M., Ligtenberg, S. R. M., Van den Broeke, M. R. and Moholdt, G.: Calving fluxes and basal melt rates of Antarctic ice shelves, *Nature*, 502(7469), 89–89, <https://doi.org/10.1038/nature12567>, 2013.
- Dirscherl, M., Dietz, A. J., Kneisel, C. and Kuenzer, C.: Automated Mapping of Antarctic Supraglacial Lakes Using a Machine Learning Approach, *Remote Sens.*, 12(7), 1203, <https://doi.org/10.3390/rs12071203>, 2020.
- 810 Domack, E., Duran, D., Leventer, A., Ishman, S., Doane, S., McCallum, S., Amblas, D., Ring, J., Gilbert, R. and Prentice, M.: Stability of the Larsen B ice shelf on the Antarctic Peninsula during the Holocene epoch, *Nature*, 436(7051), 681–685, <https://doi.org/10.1038/nature03908>, 2005.
- Dutrieux, P., De Rydt, J., Jenkins, A., Holland, P. R., Ha, H. K., Lee, S. H., Steig, E. J., Ding, Q., Abrahamsen, E. P. and Schroder, M.: Strong Sensitivity of Pine Island Ice-Shelf Melting to Climatic Variability, *Science*, 343(6167), 174–178, <https://doi.org/10.1126/science.1244341>, 2014.
- 815 Embury, O.: Product Quality Assessment Report. Sea Surface Temperature., http://datastore.copernicus-climate.eu/c3s/published-forms/c3sprod/satellite-sea-surface-temperature/documentation/v2.0/D2.SST.2-v2.1_PQAR_of_v2SST_products_v3.1_APPROVED_Ver1.pdf, last access: 28 September 2019, 2019.
- Feldmann, J. and Levermann, A.: Collapse of the West Antarctic Ice Sheet after local destabilization of the Amundsen Basin, *Proc. Natl. Acad. Sci.*, 112(46), 14191–14196, <https://doi.org/10.1073/pnas.1512482112>, 2015.
- Ferrigno, Foley, K. M., Swithinbank, C., Williams Jr, R. S. and Dailide, L.: Coastal-Change And Glaciological Map Of The Ronne Ice Shelf Area, Antarctica: 1974–2002, USGS Coast. Change Maps, I-2600-D, 1–11, 2005.
- Ferrigno, Foley, K. M., Swithinbank, C. and Williams Jr, R. S.: Coastal-Change and Glaciological Map of the Northern Ross Ice Shelf Area, Antarctica: 1962–2004, USGS Coast. Change Maps, 1–11, 2007.
- 825 Fountain, A. G., Glenn, B. and Scambos, T. A.: The changing extent of the glaciers along the western Ross Sea, Antarctica, *Geology*, 45(10), 927–930, 2017.
- Fricker, H. A., Young, N. W., Allison, I. and Coleman, R.: Iceberg calving from the Amery Ice Shelf, East Antarctica, *Ann. Glaciol.*, 34, 241–246–241–246, <https://doi.org/10.3189/172756402781817581>, 2002.
- Friedl, P., Seehaus, T. C., Wendt, A., Braun, M. H. and Höppner, K.: Recent dynamic changes on Fleming Glacier after the disintegration of Wordie Ice Shelf, Antarctic Peninsula, *The Cryosphere*, 12(4), 1347–1365, <https://doi.org/10.5194/tc-12-1347-2018>, 2018.
- 830 Fürst, J. J., Durand, G., Gillet-Chaulet, F., Tavard, L., Rankl, M., Braun, M. and Gagliardini, O.: The safety band of Antarctic ice shelves, *Nat. Clim. Change*, 6(5), 479–482, 2016.
- Gagliardini, O., Durand, G., Zwinger, T., Hindmarsh, R. C. A. and Le Meur, E.: Coupling of ice-shelf melting and buttressing is a key process in ice-sheets dynamics, *Geophys. Res. Lett.*, 37(14), <https://doi.org/10.1029/2010gl043334>, 2010.

- Gardner, A. S., Moholdt, G., Scambos, T. A., Fahnestock, M., Ligtenberg, S., van den Broeke, M. and Nilsson, J.: Increased West Antarctic and unchanged East Antarctic ice discharge over the last 7 years, *The Cryosphere*, 12(2), 521–547, <https://doi.org/10.5194/tc-12-521-2018>, 2018.
- 840 Giles, A. B.: The Mertz Glacier Tongue, East Antarctica. Changes in the past 100 years and its cyclic nature - Past, present and future, *Remote Sens. Environ.*, 191, 30–37, <https://doi.org/10.1016/j.rse.2017.01.003>, 2017.
- Gossart, A., Helsen, S., Lenaerts, J. T. M., Broucke, S. V., van Lipzig, N. P. M. and Souverijns, N.: An Evaluation of Surface Climatology in State-of-the-Art Reanalyses over the Antarctic Ice Sheet, *J. Clim.*, 32(20), 6899–6915, <https://doi.org/10.1175/JCLI-D-19-0030.1>, 2019.
- 845 Hazel, J. E. and Stewart, A. L.: Are the Near-Antarctic Easterly Winds Weakening in Response to Enhancement of the Southern Annular Mode?, *J. Clim.*, 32(6), 1895–1918, <https://doi.org/10.1175/JCLI-D-18-0402.1>, 2019.
- Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., Nicolas, J., Peubey, C., Radu, R., Schepers, D., Simmons, A., Soci, C., Abdalla, S., Abellan, X., Balsamo, G., Bechtold, P., Biavati, G., Bidlot, J., Bonavita, M., Chiara, G., Dahlgren, P., Dee, D., Diamantakis, M., Dragani, R., Flemming, J., Forbes, R., Fuentes, M., Geer, A., Haimberger, L., Healy, S., Hogan, R. J., Hólm, E., Janisková, M., Keeley, S., Laloyaux, P., Lopez, P., Lupu, C., Radnoti, G., Rosnay, P., Rozum, I., Vamborg, F., Villaume, S. and Thépaut, J.: The ERA5 global reanalysis, *Q. J. R. Meteorol. Soc.*, 146(730), 1999–2049, <https://doi.org/10.1002/qj.3803>, 2020.
- 850 Hogg, A. E. and Gudmundsson, G. H.: Impacts of the Larsen-C Ice Shelf calving event, *Nat. Clim. Change*, 7(8), 540–542, <https://doi.org/10.1038/nclimate3359>, 2017.
- Holt, T. O., Glasser, N. F., Quincey, D. J. and Siegfried, M. R.: Speedup and fracturing of George VI Ice Shelf, Antarctic Peninsula, *The Cryosphere*, 7(3), 373–417, 2013.
- Howat, I. M., Joughin, I., Fahnestock, M., Smith, B. E. and Scambos, T. A.: Synchronous retreat and acceleration of southeast Greenland outlet glaciers 2000–06: ice dynamics and coupling to climate, *J. Glaciol.*, 54(187), 646–660, <https://doi.org/10.3189/002214308786570908>, 2008.
- Hughes, T.: The weak underbelly of the West Antarctic ice sheet, *J. Glaciol.*, 27(97), 518–525, 1981.
- 860 Humbert, A., Gross, D., Müller, R., Braun, M., van de Wal, R. S. W., van den Broeke, M. R., Vaughan, D. G. and van de Berg, W. J.: Deformation and failure of the ice bridge on the Wilkins Ice Shelf, Antarctica, *Ann. Glaciol.*, 51(55), 49–55–49–55, <https://doi.org/10.3189/172756410791392709>, 2010.
- IMBIE: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 558(7709), 219–222, <https://doi.org/10.1038/s41586-018-0179-y>, 2018.
- 865 Jacobs, S. S., Jenkins, A., Giulivi, C. F. and Dutrieux, P.: Stronger ocean circulation and increased melting under Pine Island Glacier ice shelf, *Nat. Geosci.*, 4(8), 519–523, 2011.
- Jenkins, A., Shoosmith, D., Dutrieux, P., Jacobs, S., Kim, T. W., Lee, S. H., Ha, H. K. and Stammerjohn, S.: West Antarctic Ice Sheet retreat in the Amundsen Sea driven by decadal oceanic variability, *Nat. Geosci.*, 11(10), 733–738, <https://doi.org/10.1038/s41561-018-0207-4>, 2018.
- 870 Jezek, K. C.: RADARSAT-1 Antarctic Mapping Project: change-detection and surface velocity campaign, *Ann. Glaciol.*, 34(1), 263–268, 2002.

- Joughin, I. and Alley, R. B.: Stability of the West Antarctic ice sheet in a warming world, *Nat. Geosci.*, 4(8), 506–513, 2011.
- Joughin, I. and MacAyeal, D. R.: Calving of large tabular icebergs from ice shelf rift systems, *Geophys. Res. Lett.*, 32(2), L02501, <https://doi.org/10.1029/2004GL020978>, 2005.
- 875 Joughin, I., Smith, B. E. and Medley, B.: Marine Ice Sheet Collapse Potentially Under Way for the Thwaites Glacier Basin, West Antarctica, *Science*, 344(6185), 735–738, <https://doi.org/10.1126/science.1249055>, 2014.
- Khazendar, A., Rignot, E. and Larour, E.: Larsen B Ice Shelf rheology preceding its disintegration inferred by a control method, *Geophys. Res. Lett.*, 34(19), 2007.
- Kim, K. T., Jezek, K. C. and Sohn, H. G.: Ice shelf advance and retreat rates along the coast of Queen Maud Land, Antarctica, *J. Geophys. Res.-OCEANS*, 106(C4), 7097–7106, <https://doi.org/10.1029/2000JC000317>, 2001.
- 880 Konrad, H., Shepherd, A., Gilbert, L., Hogg, A. E., McMillan, M., Muir, A. and Slater, T.: Net retreat of Antarctic glacier grounding lines, *Nat. Geosci.*, 11(4), 258–258, 2018.
- Kopp, R. E., DeConto, R. M., Bader, D. A., Hay, C. C., Horton, R. M., Kulp, S., Oppenheimer, M., Pollard, D. and Strauss, B. H.: Evolving Understanding of Antarctic Ice-Sheet Physics and Ambiguity in Probabilistic Sea-Level Projections, *Earth's Future*, 5(12), 1217–1233, <https://doi.org/10.1002/2017EF000663>, 2017.
- 885 Kwok, R. and Comiso, J. C.: Spatial patterns of variability in Antarctic surface temperature: Connections to the Southern Hemisphere Annular Mode and the Southern Oscillation: Antarctic Surface Temperature, *Geophys. Res. Lett.*, 29(14), 50-1-50-4, <https://doi.org/10.1029/2002GL015415>, 2002.
- Larour, E.: Modelling of rift propagation on Ronne Ice Shelf, Antarctica, and sensitivity to climate change, *Geophys. Res. Lett.*, 31(16), L16404, <https://doi.org/10.1029/2004GL020077>, 2004.
- 890 Lavergne, T., Sørensen, A. M., Kern, S., Tonboe, R., Notz, D., Aaboe, S., Bell, L., Dybkjær, G., Eastwood, S., Gabarro, C., Heygster, G., Killie, M. A., Brandt Kreiner, M., Lavelle, J., Saldo, R., Sandven, S. and Pedersen, L. T.: Version 2 of the EUMETSAT OSI SAF and ESA CCI sea-ice concentration climate data records, *The Cryosphere*, 13(1), 49–78, <https://doi.org/10.5194/tc-13-49-2019>, 2019.
- 895 Leeson, A. A., Van Wessem, J. M., Ligtenberg, S. R. M., Shepherd, A., Van Den Broeke, M. R., Killick, R., Skvarca, P., Marinsek, S. and Colwell, S.: Regional climate of the Larsen B embayment 1980–2014, *J. Glaciol.*, 63(240), 683–690, <https://doi.org/10.1017/jog.2017.39>, 2017.
- Leeson, A. A., Forster, E., Rice, A., Gourmelen, N. and Wessem, J. M.: Evolution of Supraglacial Lakes on the Larsen B Ice Shelf in the Decades Before it Collapsed, *Geophys. Res. Lett.*, 47(4), <https://doi.org/10.1029/2019GL085591>, 2020.
- 900 Lipovsky, B. P.: Ice shelf rift propagation: stability, three-dimensional effects, and the role of marginal weakening, *The Cryosphere*, 14(5), 1673–1683, <https://doi.org/10.5194/tc-14-1673-2020>, 2020.
- Liu, H. and Jezek, K. C.: A complete high-resolution coastline of Antarctica extracted from orthorectified Radarsat SAR imagery, *Photogramm. Eng. Remote Sens.*, 70(5), 605–616, <https://doi.org/10.14358/pers.70.5.605>, 2004.
- 905 Liu, Y., Moore, J. C., Cheng, X., Gladstone, R. M., Bassis, J. N., Liu, H., Wen, J. and Hui, F.: Ocean-driven thinning enhances iceberg calving and retreat of Antarctic ice shelves, *Proc. Natl. Acad. Sci.*, 112(11), 3263–3268, 2015.

- Lovell, A. M., Stokes, C. R. and Jamieson, S. S. R.: Sub-decadal variations in outlet glacier terminus positions in Victoria Land, Oates Land and George V Land, East Antarctica (1972–2013), *Antarct. Sci.*, 29(5), 468–483–468–483, <https://doi.org/10.1017/S0954102017000074>, 2017.
- Lucchitta, B. K. and Rosanova, C. E.: Retreat of northern margins of George VI and Wilkins Ice Shelves, Antarctic Peninsula, vol. 27, edited by W. F. Budd, pp. 41–46, , 1998.
- Luckman, A., Benn, D. I., Cottier, F., Bevan, S., Nilsen, F. and Inall, M.: Calving rates at tidewater glaciers vary strongly with ocean temperature, *Nat. Commun.*, 6(1), 8566, <https://doi.org/10.1038/ncomms9566>, 2015.
- Ludescher, J., Yuan, N. and Bunde, A.: Detecting the statistical significance of the trends in the Antarctic sea ice extent: an indication for a turning point, *Clim. Dyn.*, 53(1–2), 237–244, <https://doi.org/10.1007/s00382-018-4579-3>, 2019.
- MacAyeal, D. R., Padman, L., Drinkwater, M., Fahnestock, M., Gotis, T. T., A., Gray, L., Kerman, B., Lazzara, M., Rignot, E., Scambos, T. and Stearns, C.: Effects of Rigid Body Collisions and Tide-Forced Drift on Large Tabular Icebergs of the Antarctic, http://ocp.ldeo.columbia.edu/res/div/ocp/projects/anslope/Icebergs_of_Y2k.pdf, last access: 3 December 2020, 2001.
- Marshall, G. J.: Half-century seasonal relationships between the Southern Annular mode and Antarctic temperatures, *Int. J. Climatol.*, 27(3), 373–383, <https://doi.org/10.1002/joc.1407>, 2007.
- Massom, R., Reid, P., Stammerjohn, S., Raymond, B., Fraser, A. and Ushio, S.: Change and Variability in East Antarctic Sea Ice Seasonality, 1979/80–2009/10, edited by P. Strutton, *PLoS ONE*, 8(5), e64756, <https://doi.org/10.1371/journal.pone.0064756>, 2013.
- Massom, R. A., Giles, A. B., Warner, R. C., Fricker, H. A., Legresy, B., Hyland, G., Lescarmontier, L. and Young, N.: External influences on the Mertz Glacier Tongue (East Antarctica) in the decade leading up to its calving in 2010, *J. Geophys. Res. Earth Surf.*, 120(3), 490–506, <https://doi.org/10.1002/2014JF003223>}, 2015.
- Massom, R. A., Scambos, T. A., Bennetts, L. G., Reid, P., Squire, V. A. and Stammerjohn, S. E.: Antarctic ice shelf disintegration triggered by sea ice loss and ocean swell, *Nature*, 558(7710), 383–389, <https://doi.org/10.1038/s41586-018-0212-1>, 2018.
- Mercer, J. H.: West Antarctic ice sheet and CO2 greenhouse effect: a threat of disaster, *Nature*, 271(5643), 321–325, 1978.
- Miles, B. W. J., Stokes, C. R. and Jamieson, S. S. R.: Pan-ice-sheet glacier terminus change in East Antarctica reveals sensitivity of Wilkes Land to sea-ice changes, *Sci. Adv.*, 2(5), e1501350, <https://doi.org/10.1126/sciadv.1501350>, 2016.
- Miles, B. W. J., Stokes, C. R., Jenkins, A., Jordan, J. R., Jamieson, S. S. R. and Gudmundsson, G. H.: Intermittent structural weakening and acceleration of the Thwaites Glacier Tongue between 2000 and 2018, *J. Glaciol.*, 1–11, <https://doi.org/10.1017/jog.2020.20>, 2020.
- Milillo, P., Rignot, E., Rizzoli, P., Scheuchl, B., Mouginot, J., Bueso-Bello, J. and Prats-Iraola, P.: Heterogeneous retreat and ice melt of Thwaites Glacier, West Antarctica, *Sci. Adv.*, 5(1), eaau3433, <https://doi.org/10.1126/sciadv.aau3433>, 2019.
- Mosbeux, C., Wagner, T. J. W., Becker, M. K. and Fricker, H. A.: Viscous and elastic buoyancy stresses as drivers of ice-shelf calving, *J. Glaciol.*, 1–15, <https://doi.org/10.1017/jog.2020.35>, 2020.
- Mouginot, J.: MEaSURES Antarctic Boundaries for IPY 2007-2009 from Satellite Radar, Version 2, , <https://doi.org/10.5067/AXE4121732AD>, 2017.

- Mouginot, J., Rignot, E. and Scheuchl, B.: Continent-Wide, Interferometric SAR Phase, Mapping of Antarctic Ice Velocity, *Geophys. Res. Lett.*, 46(16), 9710–9718, <https://doi.org/10.1029/2019GL083826>, 2019.
- 945 Nakamura, K., Doi, K. and Shibuya, K.: Why is Shirase Glacier turning its flow direction eastward?, *Polar Sci.*, 1(2), 63–71, <https://doi.org/10.1016/j.polar.2007.09.003>, 2007.
- Nicholls, K. W., Østerhus, S., Makinson, K., Gammelsrød, T. and Fahrbach, E.: Ice-ocean processes over the continental shelf of the southern Weddell Sea, Antarctica: A review, *Rev. Geophys.*, 47(3), 2009.
- NSIDC: Natl. Snow Ice Data Cent. RAMP AMM-1 SAR Image Mosaic Antarct. Version 2 <https://nsidc.org/data/NSIDC-0103/versions/2>, last access: 3 January 2020, 1997.
- 950 NSIDC: Natl. Snow Ice Data Cent. MODIS Mosaic Antarct. 2008-2009 MOA2009 Image Map Version 1 <https://nsidc.org/data/NSIDC-0593/versions/1>, last access: 3 May 2020, 2009.
- Osisaf: Ocean and Sea Ice Satellite Application Facility. High Latitude Products. Sea Ice Concentration., <http://osisaf.met.no/p/ice/>, last access: 3 February 2020, 2020.
- 965 Paeth, H. and Pollinger, F.: Enhanced evidence in climate models for changes in extratropical atmospheric circulation, *Tellus Dyn. Meteorol. Oceanogr.*, 62(5), 647–660, <https://doi.org/10.1111/j.1600-0870.2010.00455.x>, 2010.
- Paolo, F. S., Fricker, H. A. and Padman, L.: Volume loss from Antarctic ice shelves is accelerating, *Science*, 348(6232), 327–331, 2015.
- Parizek, B. R., Christianson, K., Anandakrishnan, S., Alley, R. B., Walker, R. T., Edwards, R. A., Wolfe, D. S., Bertini, G. T., Rinehart, S. K., Bindshadler, R. A. and Nowicki, S. M. J.: Dynamic (in)stability of Thwaites Glacier, West Antarctica: THWAITES DYNAMICS, *J. Geophys. Res. Earth Surf.*, 118(2), 638–655, <https://doi.org/10.1002/jgrf.20044>, 2013.
- 970 Parkinson, C. L.: A 40-y record reveals gradual Antarctic sea ice increases followed by decreases at rates far exceeding the rates seen in the Arctic, *Proc. Natl. Acad. Sci.*, 116(29), 14414–14423, <https://doi.org/10.1073/pnas.1906556116>, 2019.
- Pattyn, F., Ritz, C., Hanna, E., Asay-Davis, X., DeConto, R., Durand, G., Favier, L., Fettweis, X., Goelzer, H., Golledge, N. R., Munneke, P. K., Lenaerts, J. T. M., Nowicki, S., Payne, A. J., Robinson, A., Seroussi, H., Trusel, L. D. and Broeke, M. van den: The Greenland and Antarctic ice sheets under 1.5 °C global warming, *Nat. Clim. Change*, 1, <https://doi.org/10.1038/s41558-018-0305-8>, 2018.
- 965 Pollard, D., DeConto, R. M. and Alley, R. B.: Potential Antarctic Ice Sheet retreat driven by hydrofracturing and ice cliff failure, *Earth Planet. Sci. Lett.*, 412, 112–121, 2015.
- 970 Rack, W. and Rott, H.: Pattern of retreat and disintegration of the Larsen B ice shelf, Antarctic Peninsula, *Ann. Glaciol.*, 39(1), 505–510, 2004.
- Rankl, M., Fürst, J. J., Humbert, A. and Braun, M. H.: Dynamic changes on the Wilkins Ice Shelf during the 2006–2009 retreat derived from satellite observations, *The Cryosphere*, 11(3), 1199–1211, <https://doi.org/10.5194/tc-11-1199-2017>, 2017.
- Rignot, E.: Ice-shelf changes in Pine Island Bay, Antarctica, 1947-2000, *J. Glaciol.*, 48(161), 247–256, <https://doi.org/10.3189/172756502781831386>, 2002.
- 975 Rignot, E., Mouginot, J. and Scheuchl, B.: Ice Flow of the Antarctic Ice Sheet, *Science*, 333(6048), 1427–1430, <https://doi.org/10.1126/science.1208336>, 2011.

- Rignot, E., Jacobs, S., Mouginot, J. and Scheuchl, B.: Ice-Shelf Melting Around Antarctica, *Science*, 341(6143), 266–270, <https://doi.org/10.1126/science.1235798>, 2013.
- 980 Rignot, E., Mouginot, J., Morlighem, M., Seroussi, H. and Scheuchl, B.: Widespread, rapid grounding line retreat of Pine Island, Thwaites, Smith, and Kohler glaciers, West Antarctica, from 1992 to 2011, *Geophys. Res. Lett.*, 41(10), 3502–3509, <https://doi.org/10.1002/2014GL060140>, 2014.
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J. and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proc. Natl. Acad. Sci.*, 116(4), 1095–1103, <https://doi.org/10.1073/pnas.1812883116>, 2019.
- 985 Robel, A. A., Seroussi, H. and Roe, G. H.: Marine ice sheet instability amplifies and skews uncertainty in projections of future sea-level rise, *Proc. Natl. Acad. Sci.*, 116(30), 14887–14892, <https://doi.org/10.1073/pnas.1904822116>, 2019.
- Rosier, S. H. R., Green, J. A. M., Scourse, J. D. and Winkelmann, R.: Modeling Antarctic tides in response to ice shelf thinning and retreat, *J. Geophys. Res. Oceans*, 119(1), 87–97, <https://doi.org/10.1002/2013JC009240>, 2014.
- 990 Rott, H., Müller, F., Nagler, T. and Floricioiu, D.: The imbalance of glaciers after disintegration of Larsen-B ice shelf, *Antarctic Peninsula, The Cryosphere*, 5(1), 125–125, 2011.
- Royston, S. and Gudmundsson, G. H.: Changes in ice-shelf buttressing following the collapse of Larsen A Ice Shelf, Antarctica, and the resulting impact on tributaries, *J. Glaciol.*, 62(235), 905–911, 2016.
- Scambos, T. A., Hulbe, C., Fahnestock, M. and Bohlander, J.: The link between climate warming and break-up of ice shelves in the Antarctic Peninsula, *J. Glaciol.*, 46(154), 516–530, <https://doi.org/10.3189/172756500781833043>, 2000.
- 995 Scambos, T. A., Haran, T. M., Fahnestock, M. A., Painter, T. H. and Bohlander, J.: MODIS-based Mosaic of Antarctica (MOA) data sets: Continent-wide surface morphology and snow grain size, *Remote Sens. Environ.*, 111(2), 242–257, <https://doi.org/10.1016/j.rse.2006.12.020>, 2007.
- Scambos, T. A., Fricker, H. A., Liu, C.-C., Bohlander, J., Fastook, J., Sargent, A., Massom, R. and Wu, A.-M.: Ice shelf disintegration by plate bending and hydro-fracture: Satellite observations and model results of the 2008 Wilkins ice shelf break-ups, *Earth Planet. Sci. Lett.*, 280(1), 51–60, <https://doi.org/10.1016/j.epsl.2008.12.027>, 2009.
- 1000 Scheuchl, B., Mouginot, J., Rignot, E., Morlighem, M. and Khazendar, A.: Grounding line retreat of Pope, Smith, and Kohler Glaciers, West Antarctica, measured with Sentinel-1a radar interferometry data, *Geophys. Res. Lett.*, 43(16), 8572–8579, <https://doi.org/10.1002/2016GL069287>, 2016.
- Seehaus, T., Marinsek, S., Helm, V., Skvarca, P. and Braun, M.: Changes in ice dynamics, elevation and mass discharge of Dinsmoor–Bombardier–Edgeworth glacier system, Antarctic Peninsula, *Earth Planet. Sci. Lett.*, 427, 125–135, <https://doi.org/10.1016/j.epsl.2015.06.047>, 2015.
- Spence, P., Griffies, S. M., England, M. H., Hogg, A. McC., Saenko, O. A. and Jourdain, N. C.: Rapid subsurface warming and circulation changes of Antarctic coastal waters by poleward shifting winds: Antarctic subsurface ocean warming, *Geophys. Res. Lett.*, 41(13), 4601–4610, <https://doi.org/10.1002/2014GL060613>, 2014.
- 1010 Stewart, A. L., Klocker, A. and Menemenlis, D.: Circum-Antarctic Shoreward Heat Transport Derived From an Eddy- and Tide-Resolving Simulation, *Geophys. Res. Lett.*, 45(2), 834–845, <https://doi.org/10.1002/2017GL075677>, 2018.

Tedesco, M. and Monaghan, A. J.: An updated Antarctic melt record through 2009 and its linkages to high-latitude and tropical climate variability, *Geophys. Res. Lett.*, 36(18), L18502, <https://doi.org/10.1029/2009GL039186>, 2009.

1015 Tetzner, D., Thomas, E. and Allen, C.: A Validation of ERA5 Reanalysis Data in the Southern Antarctic Peninsula—Ellsworth Land Region, and Its Implications for Ice Core Studies, *Geosciences*, 9(7), 289, <https://doi.org/10.3390/geosciences9070289>, 2019.

Thoma, M., Jenkins, A., Holland, D. and Jacobs, S.: Modelling Circumpolar Deep Water intrusions on the Amundsen Sea continental shelf, *Antarctica, Geophys. Res. Lett.*, 35(18), L18602, <https://doi.org/10.1029/2008GL034939>, 2008.

1020 Trusel, L. D., Frey, K. E., Das, S. B., Munneke, P. K. and van den Broeke, M. R.: Satellite-based estimates of Antarctic surface meltwater fluxes: Satellite-Based Antarctic Melt Fluxes, *Geophys. Res. Lett.*, 40(23), 6148–6153, <https://doi.org/10.1002/2013GL058138>, 2013.

Vaughan, D. G. and Doake, C. S. M.: Recent atmospheric warming and retreat of ice shelves on the Antarctic Peninsula, *Nature*, 379(6563), 328–328, 1996.

1025 Verdy, A., Marshall, J. and Czaja, A.: Sea surface temperature variability along the path of the Antarctic Circumpolar Current., *J. Phys. Oceanogr.*, 36(7), 1317–1331, 2006.

Walker, C. C. and Gardner, A. S.: Rapid drawdown of Antarctica’s Wordie Ice Shelf glaciers in response to ENSO/Southern Annular Mode-driven warming in the Southern Ocean, *Earth Planet. Sci. Lett.*, 476, 100–110, <https://doi.org/10.1016/j.epsl.2017.08.005>, 2017.

1030 Walker, C. C., Bassis, J. N., Fricker, H. A. and Czerwinski, R. J.: Structural and environmental controls on Antarctic ice shelf rift propagation inferred from satellite monitoring: ANTARCTIC ICE SHELF RIFTING, *J. Geophys. Res. Earth Surf.*, 118(4), 2354–2364, <https://doi.org/10.1002/2013JF002742>, 2013.

Walker, C. C., Gardner, A. S., Neumann, T., Fricker, H. A., Bassis, J. N. and Paolo, F. S.: Iceberg, right ahead!: The surprising and ongoing collapse of an East Antarctic ice shelf in response to changes in the ocean environment, in *AGU Fall Meeting Abstracts*, vol. 2019, pp. C13A-06, , 2019.

1035 Wang, G., Cai, W. and Purich, A.: Trends in Southern Hemisphere wind-driven circulation in CMIP5 models over the 21st century: Ozone recovery versus greenhouse forcing, *J. Geophys. Res. Oceans*, 119(5), 2974–2986, <https://doi.org/10.1002/2013JC009589>, 2014.

1040 Wouters, B., Martin-Espanol, A., Helm, V., Flament, T., van Wessem, J. M., Ligtenberg, S. R. M., van den Broeke, M. R. and Bamber, J. L.: Dynamic thinning of glaciers on the Southern Antarctic Peninsula, *Science*, 348(6237), 899–903, <https://doi.org/10.1126/science.aaa5727>, 2015.

Wuite, J., Nagler, T., Gourmelen, N., Escorihuela, M. J., Hogg, A. E. and Drinkwater, M. R.: Sub-Annual Calving Front Migration, Area Change and Calving Rates from Swath Mode CryoSat-2 Altimetry, on Filchner-Ronne Ice Shelf, Antarctica, *Remote Sens.*, 11(23), 2761, <https://doi.org/10.3390/rs11232761>, 2019.