

1 A pilot study about microplastics and mesoplastics in an Antarctic glacier:  
2 the role of atmospheric dry deposition

3  
4 Miguel González-Pleiter<sup>1,2†</sup>, Gissell Lacerot<sup>3</sup>, Carlos Edo<sup>1</sup>, Juan Pablo Lozoya<sup>4</sup>, Francisco  
5 Leganés<sup>2</sup>, Francisca Fernández-Piñas<sup>2</sup>, Roberto Rosal<sup>1</sup>, Franco Teixeira-de-Mello<sup>5†</sup>

6  
7 <sup>1</sup>Department of Analytical Chemistry, Physical Chemistry and Chemical Engineering,  
8 University of Alcala, Alcalá de Henares, E-28871 Madrid, Spain

9  
10 <sup>2</sup>Departament of Biology, Faculty of Sciences, Universidad Autónoma de Madrid,  
11 Cantoblanco, E-28049 Madrid, Spain

12  
13 <sup>3</sup>Ecología Funcional de Sistemas Acuáticos, Centro Universitario Regional del Este  
14 (CURE), Universidad de la República, Ruta nacional N°9 y ruta N°15, 27000 Rocha,  
15 Uruguay

16 <sup>4</sup>Centro Interdisciplinario de Manejo Costero Integrado del Cono Sur (C-MCISur), Centro  
17 Universitario Regional del Este (CURE), Universidad de la República, Tacuarembó entre  
18 Av. Artigas y Aparicio Saravia, 20000 Maldonado, Uruguay

19  
20 <sup>5</sup>Departamento de Ecología y Gestión Ambiental, Centro Universitario Regional del Este  
21 (CURE), Universidad de la República, Tacuarembó entre Av. Artigas y Aparicio Saravia,  
22 20000 Maldonado, Uruguay

23  
24 †Corresponding authors:

25 Miguel González-Pleiter, email: [mig.gonzalez@uam.es](mailto:mig.gonzalez@uam.es)

26 Franco Teixeira-de-Mello, email: [frantei@cure.edu.uy](mailto:frantei@cure.edu.uy)

27  
28 **Abstract**

29 Plastics have been found in several compartments in Antarctica. However, there is  
30 currently no evidence of their presence in Antarctic glaciers. Our pilot study investigated  
31 plastic occurrence on two ice surfaces (one area close to Uruguay lake and another one  
32 close to Ionosferico lake) that constitute part of the ablation zone of Collins Glacier (King  
33 George Island, Antarctica). Our results showed that expanded polystyrene (EPS) was  
34 ubiquitous ranging from 0.17 to 0.33 items m<sup>-2</sup> whereas polyester was found only on the  
35 ice surface close to Uruguay lake (0.25 items m<sup>-2</sup>). Furthermore, we evaluated the daily  
36 changes in the presence of plastics in these areas in the absence of rainfall to clarify the  
37 role of the wind in their transport. We registered an atmospheric dry deposition rate  
38 between 0.08 items m<sup>-2</sup> day<sup>-1</sup> on the ice surface close to Uruguay lake and 0.17 items m<sup>-2</sup>  
39 day<sup>-1</sup> on the ice surface close to Ionosferico lake. Our pilot study is the first report of  
40 plastic pollution presence in an Antarctic glacier, possibly originated from local current  
41 and past activities, and the first to assess the effect of wind in its transport.

46 **Introduction**

47 The cryosphere is the frozen water part of the Earth system that consists of areas in  
48 which the temperatures are below 0°C for at least part of the year (NOAA, 2019). Most  
49 of the cryosphere in terms of volume of ice is in Antarctica. Despite the increasing rate  
50 of ice loss during last decades (Rignot et al., 2019), it has been estimated that the  
51 Antarctic cryosphere holds around 90% of Earth's ice mass (Dirscherl et al., 2020).  
52 Furthermore, the Antarctic cryosphere represents the majority of the world's  
53 freshwater, representing the largest freshwater ecosystem on the planet (Shepherd et  
54 al., 2018).

55

56 Plastics, especially microplastics (plastic items < 5 mm long; MPs), have been detected  
57 in several specific locations of the cryosphere including mountain glaciers (Ambrosini et  
58 al., 2019; Cabrera et al., 2020; Materić et al., 2020), snow (Bergmann et al., 2019;  
59 Österlund et al., 2019) and sea ice (Geilfus et al., 2019; Kelly et al., 2020; La Daana et al.,  
60 2020; Obbard et al., 2014; Peeken et al., 2018; von Friesen et al., 2020). The occurrence  
61 of MPs in snow ranged from 0 to  $1.5 \times 10^5$  MP L<sup>-1</sup> of melted snow (Bergmann et al., 2019),  
62 although it should be noted that a part of this study was conducted near urban areas.  
63 Regarding sea ice, concentrations of up to  $1.2 \times 10^4$  MP L<sup>-1</sup> have been reported, although  
64 there are large differences between studies even from the same region (Peeken et al.,  
65 2018; von Friesen et al., 2020). The use of different units in reporting MP concentrations  
66 in mountain glaciers such as the number of items per mass of ice weight ( $78.3 \pm 30.2$   
67 MPs kg<sup>-1</sup> of sparse and fine supraglacial debris; Ambrosini et al., 2019) and mass of MPs  
68 per volume (0 to  $23.6 \pm 3.0$  ng of MPs mL<sup>-1</sup>; Materić et al., 2020), makes comparisons  
69 between studies difficult (101.2 items L<sup>-1</sup>; Cabrera et al., 2020). Regarding the shape of  
70 the MPs found in the cryosphere, fibers seem to be dominant in mountain glaciers (65  
71 %) and sea ice (79 %), followed by fragments (Ambrosini et al., 2019; La Daana et al.,  
72 2020). Concerning the size of MPs, it has been reported a broad size distribution in sea  
73 ice, with 67 % of MPs in the 500-5000 µm range (La Daana et al., 2020). Other studies  
74 found lower sizes, however, with significant amounts (up to 90 %) of MPs smaller than  
75 100 µm in snow and sea ice (Ambrosini et al., 2019; Bergmann et al., 2019; Bergmann et  
76 al., 2017; Kelly et al., 2020; Peeken et al., 2018). The differences between these studies  
77 may be due to the different analytical methods used, particularly methodologies such  
78 as micro Fourier transform infrared spectroscopy (µFTIR, which can identify smaller  
79 sized MPs). In general, the presence of plastics > 5mm has not been reported in the  
80 cryosphere, probably because they occur at lower concentrations and evade detection.  
81 µFTIR revealed that polyethylene terephthalate (PET), polyamide (PA), polyester (PE),  
82 varnish (acrylates/polyurethane), several synthetic rubbers, polypropylene (PP), and  
83 polyurethane (PU) are the most common types of MPs in the cryosphere (Ambrosini et  
84 al., 2019; Bergmann et al., 2019; Bergmann et al., 2017; La Daana et al., 2020; Materić  
85 et al., 2020; Obbard et al., 2014; Peeken et al., 2018). The sources of MPs detected in the  
86 cryosphere, however, remain poorly understood. It has been suggested that they could  
87 be transported by the wind before being deposited by both wet and dry deposition in  
88 remote areas such as polar regions (Halsband and Herzke, 2019). In fact, it has been  
89 reported that air masses can transport MPs through the atmosphere over distances of

90 at least 100 km and that they can be released from the marine environment into the  
91 atmosphere by sea-spray (Allen et al., 2020; Allen et al., 2019; González-Pleiter et al.,  
92 2020a).

93

94 So far, plastics have been found in specific parts of the cryosphere (mountain glacier,  
95 snow, and sea ice) and Antarctica (seawater, freshwater, sediments, and organisms). We  
96 hypothesize that plastics have also reached freshwater glaciers in Antarctica and that  
97 atmospheric dry deposition plays a crucial role in this process. To test this hypothesis,  
98 we carried out a pilot study to investigate the presence of plastics on two ice surfaces  
99 (one area close to Uruguay lake and another one close to Ionosferico lake) that  
100 constitute part of the ablation zone of Collins Glacier in Maxwell Bay in King George  
101 Island (Antarctica). Furthermore, the daily changes in the presence of plastics in these ice  
102 surfaces was evaluated in the absence of rainfall, to clarify the role of wind in their  
103 transport.

104

## 105 **Materials and Methods**

### 106 2.1 Study area

107 Collins Glacier is located on the northeast of Fildes Peninsula (King George Island,  
108 Antarctica; Figure 1A) and has a total surface area of 15 km<sup>2</sup> (Simoes et al., 2015). Our  
109 study was carried out on the ice surface of the glacier ablation areas close to two lakes  
110 (Uruguay or Profound, and Ionosferico) in Maxwell Bay (Figure 1B). Uruguay lake (S 62°  
111 11' 6.54", O 58° 54' 42.23") is located in the proximity of the Artigas Antarctic Scientific  
112 Base and its access road (~300 m) is subjected to human transit (Figure 1B). The lake is  
113 used for drinking and domestic water supply. The glacier surface studied in this lake  
114 covered 1680 m<sup>2</sup>. Ionosferico lake (62° 11' 59.41", O 58° 57' 44.17") is located ~600 m  
115 from Artigas Base and has minimal human activity. The glacier surface studied in this  
116 lake covered 537 m<sup>2</sup> (Figure 1B). It should be noted that there were no visible footpaths  
117 through or nearby the glacier surfaces of both lakes during the duration of our study  
118 (except our own footprints).

119

### 120 2.2 Experimental assessment of plastic concentration

121 To evaluate the concentration of plastics, twelve squares were marked on the ice  
122 surface close to Uruguay lake (Figure 1C) and six squares on the ice surface close to  
123 Ionosferico lake (Figure 1D), which constitute part of the ablation zone of Collins Glacier,  
124 on 18/2/2020. The first square of 1m<sup>2</sup> on the ice surface close to each lake was randomly  
125 marked. After that, the rest of the squares of 1m<sup>2</sup> were distributed every ten meters  
126 covering the entire ice surface in each lake (Figure 1E). All items visually resembling  
127 plastic (suspected plastic) inside the squares were registered (Figure 1F). It should be  
128 noted that our sampling strategy excluded the plastics non-detectable by the naked eye  
129 (i.e. small plastics such as fibers). Thus, we probably underestimated the concentration  
130 of small plastics on the ice surface.

131

### 132 2.3 Experimental assessment of atmospheric dry deposition of plastics

133 After the initial sampling, we selected six squares on the ice close to each lake for  
134 subsequent daily monitoring. Additional sampling was performed every twelve hours for  
135 two days (18/02/2020 and 20/02/2020) after the initial sampling. No rainfall occurred  
136 during the duration of the experiment.

137

#### 138 2.4 Characterization and identification of plastics

139 Every item visually resembling plastic detected in the squares was collected with  
140 stainless-steel tweezers, placed into 100 mL ISO reagent bottles, and stored at 4 °C until  
141 analysis. All collected items were photographed, measured and their composition was  
142 identified by ATR-FTIR using an Agilent Cary 630 FTIR spectrometer or by  $\mu$ FTIR on a  
143 Perkin-Elmer Spotlight 200 Spectrum Two apparatus equipped with a MCT detector  
144 (depending on the size of the item). The spectra were taken using the following  
145 parameters in micro-transmission mode: spot 50  $\mu$ m, 32 scans, and spectral range 550-  
146 4000  $\text{cm}^{-1}$  with 8  $\text{cm}^{-1}$  resolution. The spectra were processed using Omnic software  
147 (Thermo Fisher). Items with matching values > 60% were considered plastic materials.  
148 The results of concentration and atmospheric dry deposition of plastics reported in this  
149 study include only items positively identified as plastics according to the FTIR analysis  
150 and were expressed as number of items per surface unit and items per surface unit and  
151 day respectively.

152

#### 153 2.5 Prevention of procedural contamination

154 To avoid sample contamination, all materials used were previously cleaned with MilliQ  
155 water, wrapped in aluminum foil, and heated to 300 °C for 4 h to remove organic matter.  
156 The use of any plastic material during sampling was avoided. Furthermore, possible  
157 contamination from our clothes was controlled throughout the sampling, by checking  
158 fibers and fragments extracted from the clothes against the MPs and MePs found in the  
159 samples, and by positioning us against the wind during sampling. Given their size,  
160 plastics found in this study were detected by the naked eye and their traceability could  
161 be easily maintained during quantification and identification of the samples.

162

### 163 **Results**

#### 164 3.1 Characterization and identification of the plastics

165 In total, 45 items preliminarily identified as plastics were collected, of which 29 items  
166 were confirmed as plastic by FTIR or  $\mu$ FTIR analyses (matching > 60%). The size of  
167 plastics ranged from 2292 to 12628  $\mu$ m length and from 501 to 11334  $\mu$ m width (Figure  
168 2A). According to their size, 13 mesoplastic items (plastic items between 5-25 mm long;  
169 MeP) and 3 MP items were found on the ice close to Uruguay lake, and 12 MeP items  
170 and 1 MP item on the ice close to Ionosferico lake (Figure 2B). Meso and MPs  
171 (hereinafter referred to as plastics) of expanded polystyrene (EPS) were found on the  
172 ice close to both lakes: 8 plastic items on the ice close to Uruguay lake and 13 plastic  
173 items on the ice close to Ionosferico lake (Figure 2 B, C, and D). Polyester (n = 7 items;  
174 Figure 2B, E, and F) and polyurethane (n = 1 item; Figure 2B, G and H) items were present  
175 only on the ice close to Uruguay lake. It should be noted that spectra of the polyester

176 (Figure 2F) showed a high similarity with alkyd resin, a thermoplastic polyester widely  
177 used in synthetic paints.

178

### 179 3.2 Plastic concentration

180 EPS items were ubiquitous on the ice with concentrations ranging from 0.17 items m<sup>-2</sup>  
181 on the ice close to Uruguay lake to 0.33 items m<sup>-2</sup> on the ice close to Ionosferico lake  
182 (Table S1). The concentration of polyester, which was found only on the ice close to  
183 Uruguay lake, was 0.25 items m<sup>-2</sup> (Table S1). Polyurethane items were not observed in  
184 Ionosferico lake (Table S1).

185

### 186 3.3 Atmospheric dry deposition of plastics

187 The dry deposition rate of EPS was 0.08 EPS items m<sup>-2</sup> day<sup>-1</sup> and 0.17 EPS items m<sup>-2</sup> day<sup>-1</sup>  
188 on the ice close to Uruguay and Ionosferico lakes, respectively (Table S2 and Figure 3).  
189 Polyester was only deposited on the ice close to Uruguay lake at a rate of 0.08 items m<sup>-2</sup>  
190 day<sup>-1</sup>. Polyurethane items were not observed in Ionosferico lake during the duration  
191 of the experiment (Table S2). The plastics deposited on the ice of Ionosferico lake during  
192 the experiment were exclusively EPS (Table S2 and Figure 3).

193

## 194 Discussion

195 The presence of plastics has been documented in different places in Antarctica: marine  
196 surface waters (Cincinelli et al., 2017; Isobe et al., 2017; Jones-Williams et al., 2020;  
197 Lacerda et al., 2019; Suaria et al., 2020), marine sediments (Cunningham et al., 2020;  
198 Munari et al., 2017; Reed et al., 2018), zooplankton samples from ocean water (Absher  
199 et al., 2019), marine benthic invertebrates (Sfriso et al., 2020), Antarctic Collembola  
200 (Bergami et al., 2020b), penguins (Bessa et al., 2019), seabirds (Ibañez et al., 2020) and  
201 freshwater (González-Pleiter et al., 2020b). However, there was only one study showing  
202 the occurrence of plastics in the Antarctic cryosphere, which was carried out on sea ice  
203 (Kelly et al., 2020). Thus, this is the first report on the presence of MPs and MePs in  
204 Antarctic freshwater glaciers. Furthermore, our findings provide an insight into the role  
205 of wind in the transport of this material.

206

207 In this sense, winds (especially high-speed ones) appear to be a key element in the  
208 transport of plastics to Antarctic glaciers. The prevailing winds in the study area (Figure  
209 1B) blow predominantly from the west (Figure 4A). However, strong winds (Figure 4B),  
210 wind gusts (Figure 4C), and strong wind gusts (Figure 4D) blow mainly from the east and  
211 southeast directions, and could be responsible for the spreading of plastics from the  
212 different origins to the surface of the glacier ablation areas. These strong winds would  
213 explain the presence of MePs despite their size (Figure 2A). In fact, the low density of  
214 the MePs found (mainly EPS; Figure 2B) would have allowed their easy dispersion by  
215 wind.

216

217 Our results on the dry deposition of plastics support the hypothesis that the role of the  
218 wind is essential for the transport of MPs and MePs in (and among) different areas of  
219 Antarctica. The dry deposition of plastics (Table S2) was closely related to the wind

220 regimes during the study period (Figure S1). Based on information available on the  
221 meteorological conditions during the study dates (18/02/2020 - 20/02/2020) in La Villa  
222 de la Estrellas (Figure S1A), which is located near the Artigas Beach (Figure S2B), the  
223 wind blew from the northeast veering to the south with a speed between 10 and 30  
224 km/h (Figure S1A). These wind conditions suggest a possible link with marine  
225 environment, which can act as a source of plastics (Allen et al., 2020), and potentially  
226 explain the presence of plastics on the glacier ablation areas. However, considering the  
227 low intensity of the winds recorded during those days (Figure S1A) and the presence of  
228 MePs, it is also possible that the predominant high-speed winds transported MePs from  
229 other adjacent areas of the Fildes Peninsula to the vicinity of the lakes, in the days prior  
230 to our study (Figure 4B, C, and D) and then, the milder winds registered during the  
231 sampling days (Figure S1A) deposited these MePs on the ice.

232

233 The chemical composition of the plastics found (Figure 2D, F, and H) supports the fact  
234 that the source of the plastics could be of marine and/or land-based origin. The types of  
235 plastics found (Figure 2B) are related to human activities in the Fildes Peninsula that  
236 could generate plastic debris such as tourism, leaks in waste management at scientific  
237 bases or the presence of abandoned infrastructures. Considering the location of Collins  
238 Glacier and the main human activities on the Fildes Peninsula (e.g. airfield, scientific  
239 bases), the prevailing winds from the west could have transported small and lightweight  
240 plastics to the study area. In fact, EPS is widely used in packaging and as insulation  
241 material in old buildings in this area and polyester is also a component of old buildings  
242 paints. In the same way, some of these plastics could be released from the marine  
243 environment to Artigas beach area and, then, be transported by the wind to the glaciers.  
244 In this sense, polyurethane MePs (which are similar to those found in this work) have  
245 already been reported in sea surface waters in the Antarctic (Jones-Williams et al., 2020)  
246 and EPS MePs have been found on Artigas beach (Laganà et al., 2019). These findings  
247 highlight a potential threat to the fragile Antarctic ecosystem, since the presence of  
248 these plastics (e.g. polystyrene particles) has been shown to affect Antarctic biota  
249 (Bergami et al., 2019; Bergami et al., 2020a).

250

251 The role of the atmospheric dry deposition on the presence of plastics on glaciers is  
252 supported by recent studies suggesting that MPs can be transported, up to hundreds of  
253 kilometres, through the atmosphere before being deposited (González-Pleiter et al.,  
254 2020a). Our results showed that the atmospheric deposition of plastics on glaciers is still  
255 low, with figures between two and four orders of magnitude lower than values reported  
256 in populated areas (Brahney et al., 2020; Cai et al., 2017; Dris et al., 2016; Klein and  
257 Fischer, 2019; Roblin et al., 2020; Wright et al., 2020). Our results also show that plastic  
258 pollution, even if only in small quantities, reaches remote areas with few human  
259 settlements. The occurrence of plastic pollution in Antarctica represents the spreading  
260 of anthropogenic pollutants in the last pristine environment on the Earth. Further  
261 research is needed then to elucidate the occurrence, sources, fate, and impact of plastics  
262 in such remote places.

263

264 Taken together, our research indicates that human activities in sensitive remote areas  
265 such as Antarctica leave a footprint that includes plastic pollution. Since the early reports  
266 of litter pollution on the seafloor (Dayton and Robilliard, 1971) and ,subsequently, on  
267 beaches and seabirds of Antarctica (Convey et al., 2002; Creet et al., 1994; Fijn et al.,  
268 2012; Lenihan et al., 1990; Sander et al., 2009) the handling of waste has been improved  
269 by the implementation of the Antarctic Treaty System, Annex III ‘Waste Disposal and  
270 Waste Management’. The Treaty forces to remove all plastic from Antarctica, with the  
271 only exception of plastics that can be incinerated without producing harmful emissions  
272 (Antarctic Treaty Secretariat, 1998). However, once plastics are broken down into  
273 smaller fractions and dispersed throughout the continent and nearby waters,  
274 management measures become very difficult to address, as evidenced by our data.  
275 Thus, a more rigorous management of plastics is essential for preserving a clean  
276 environment within the Treaty Area (Zhang et al., 2020).

277

## 278 **Conclusion**

279 This is the first report of the presence of both MePs and MPs in an Antarctic glacier,  
280 which were probably transported by wind from local sources such as beach areas. In  
281 total, three types of plastics (EPS, PU and polyester) were found on two glacier surfaces  
282 that constitute part of the ablation zone of Collins Glacier (King George Island,  
283 Antarctica). EPS was ubiquitous in the two glacier surfaces studied. Our study showed  
284 that the management of plastic contamination in Antarctica should be improved,  
285 focusing on the waste generated by current and past anthropogenic activities that occur  
286 in that area.

287

## 288 **Author contribution**

289 **Miguel González-Pleiter**: identified the research question, formulated the hypothesis,  
290 developed the experimental design, planned the experiments, performed the  
291 experiments in the field, performed the experiments in the laboratory, compiled the  
292 data sets, analyzed the data, discussed the results, prepared graphical material, wrote  
293 the paper (original draft) and provided financial support. **Gissell Lacerot**: identified the  
294 research question, formulated the hypothesis, developed the experimental design,  
295 planned the experiments, checked the field data, discussed the results, wrote the paper  
296 (final version). **Carlos Edo**: performed the experiments in the laboratory, compiled the  
297 data sets, analyzed the data, discussed the results, prepared graphical material and  
298 review final manuscript. **Juan Pablo Lozoya**: developed the experimental design,  
299 checked the field data, discussed the results, review final manuscript and provided  
300 financial support. **Francisco Leganés**: discussed the results, review final manuscript and  
301 provided financial support. **Francisca Fernández-Piñas**: checked the field data, checked  
302 the laboratory data, discussed the results, review final manuscript and provided  
303 financial support. **Roberto Rosal**: checked the field data, checked the laboratory data,  
304 discussed the results, review final manuscript and provided financial support. **Franco**  
305 **Teixeira de Mello**: identified the research question, formulated the hypothesis,  
306 developed the experimental design, planned the experiments, performed the  
307 experiments in the field, checked the field data, prepared graphical material, discussed  
308 the results, review final manuscript and provided financial support.

309

### 310 **Acknowledgements**

311 This research was funded by the Government of Spain (CTM2016-74927-C2-1/2-R) and  
312 the Uruguayan Antarctic Institute. MGP thanks the Carolina Foundation for the award  
313 of a postdoctoral grant (SEGIB). CE thanks the Spanish Government for the award of a  
314 predoctoral grant. The authors gratefully acknowledge the support of Fiorella Bresesti,  
315 Evelyn Krojmal and Barbara De Feo from the Centro Universitario Regional del Este,  
316 Universidad de la República for their assistance during sampling, of Marta Elena  
317 González Mosquera from University of Alcalá for providing access to the Agilent Cary  
318 630 FTIR spectrometer, and of Gastón Manta from Facultad de Ciencias, Universidad de  
319 la República for providing historical wind analysis at the Artigas Antarctic Research Base.  
320 FTM, GL and JPL thanks the Sistema Nacional de Investigadores (SNI) and the Programa  
321 de Desarrollo de las Ciencias Básicas (PEDECIBA).

322

### 323 **Declaration of competing interest**

324 The authors declare no conflict of interest.

325

### 326 **References**

- 327 Absher, T. M., Ferreira, S. L., Kern, Y., Ferreira, A. L., Christo, S. W., and Ando, R. A.: Incidence  
328 and identification of microfibers in ocean waters in Admiralty Bay, Antarctica, *Environmental*  
329 *Science and Pollution Research*, 26, 292-298, 2019.
- 330 Allen, S., Allen, D., Moss, K., Le Roux, G., Phoenix, V. R., and Sonke, J. E.: Examination of the  
331 ocean as a source for atmospheric microplastics, *PLoS One*, 15, e0232746, 2020.
- 332 Allen, S., Allen, D., Phoenix, V. R., Le Roux, G., Jiménez, P. D., Simonneau, A., Binet, S., and Galop,  
333 D.: Atmospheric transport and deposition of microplastics in a remote mountain catchment,  
334 *Nature Geoscience*, 12, 339-344, 2019.
- 335 Ambrosini, R., Azzoni, R. S., Pittino, F., Diolaiuti, G., Franzetti, A., and Parolini, M.: First evidence  
336 of microplastic contamination in the supraglacial debris of an alpine glacier, *Environmental*  
337 *pollution*, 253, 297-301, 2019.
- 338 Antarctic Treaty Secretariat, 1998. Annex III to the Protocol on Environmental Protection to the  
339 Antarctic Treaty. Waste disposal and waste management. Accessed at <http://www.ats.aq> on 9  
340 Feb 2021.
- 341 Bergami, E., Emerenciano, A. K., González-Aravena, M., Cárdenas, C., Hernández, P., Silva, J., and  
342 Corsi, I.: Polystyrene nanoparticles affect the innate immune system of the Antarctic sea urchin  
343 *Sterechinus neumayeri*, *Polar Biology*, 42, 743-757, 2019.
- 344 Bergami, E., Manno, C., Cappello, S., Vannuccini, M., and Corsi, I.: Nanoplastics affect moulting  
345 and faecal pellet sinking in Antarctic krill (*Euphausia superba*) juveniles, *Environment*  
346 *International*, 143, 105999, 2020a.
- 347 Bergami, E., Rota, E., Caruso, T., Birarda, G., Vaccari, L., and Corsi, I.: Plastics everywhere: first  
348 evidence of polystyrene fragments inside the common Antarctic collembolan *Cryptopygus*  
349 *antarcticus*, *Biology letters*, 16, 20200093, 2020b.
- 350 Bergmann, M., Mützel, S., Primpke, S., Tekman, M. B., Trachsel, J., and Gerdt, G.: White and  
351 wonderful? Microplastics prevail in snow from the Alps to the Arctic, *Science Advances*, 5,  
352 eaax1157, 2019.
- 353 Bergmann, M., Wirzberger, V., Krumpfen, T., Lorenz, C., Primpke, S., Tekman, M. B., and Gerdt,  
354 G.: High quantities of microplastic in Arctic deep-sea sediments from the HAUSGARTEN  
355 observatory, *Environmental science & technology*, 51, 11000-11010, 2017.
- 356 Bessa, F., Ratcliffe, N., Otero, V., Sobral, P., Marques, J. C., Waluda, C. M., Trathan, P. N., and  
357 Xavier, J. C.: Microplastics in gentoo penguins from the Antarctic region, *Scientific reports*, 9, 1-  
358 7, 2019.



359 Brahney, J., Hallerud, M., Heim, E., Hahnenberger, M., and Sukumaran, S.: Plastic rain in  
360 protected areas of the United States, *Science*, 368, 1257-1260, 2020.

361 Cabrera, M., Valencia, B. G., Lucas-Solis, O., Calero, J. L., Maisincho, L., Conicelli, B., Moulatlet,  
362 G. M., and Capparelli, M. V.: A new method for microplastic sampling and isolation in mountain  
363 glaciers: A case study of one antisana glacier, *Ecuadorian Andes, Case Studies in Chemical and*  
364 *Environmental Engineering*, 2, 100051, 2020.

365 Cai, L., Wang, J., Peng, J., Tan, Z., Zhan, Z., Tan, X., and Chen, Q.: Characteristic of microplastics  
366 in the atmospheric fallout from Dongguan city, China: preliminary research and first evidence,  
367 *Environmental Science and Pollution Research*, 24, 24928-24935, 2017.

368 Cincinelli, A., Scopetani, C., Chelazzi, D., Lombardini, E., Martellini, T., Katsoyiannis, A., Fossi, M.  
369 C., and Corsolini, S.: Microplastic in the surface waters of the Ross Sea (Antarctica): occurrence,  
370 distribution and characterization by FTIR, *Chemosphere*, 175, 391-400, 2017.

371 Convey, P., Barnes, D., and Morton, A.: Debris accumulation on oceanic island shores of the  
372 Scotia Arc, Antarctica, *Polar Biology*, 25, 612-617, 2002.

373 Creet, S., Van Franeker, J., Van Spanje, T., and Wolff, W.: DIET OF THE PINTADO PETREL DAPTION  
374 CAPENSE AT KING GEORGE ISLAND, ANTARCTICA, 1990-91, *Marine Ornithology*, 22, 221-229,  
375 1994.

376 Cunningham, E. M., Ehlers, S. M., Dick, J. T., Sigwart, J. D., Linse, K., Dick, J. J., and Kiriakoulakis,  
377 K.: High abundances of microplastic pollution in deep-sea sediments: evidence from Antarctica  
378 and the Southern Ocean, *Environmental Science & Technology*, 54, 13661-13671, 2020.

379 Dayton, P. K. and Robilliard, G. A.: Implications of pollution to the McMurdo Sound benthos,  
380 1971. 1971.

381 Dirscherl, M., Dietz, A. J., Dech, S., and Kuenzer, C.: Remote sensing of ice motion in Antarctica–  
382 A review, *Remote Sensing of Environment*, 237, 111595, 2020.

383 Dris, R., Gasperi, J., Saad, M., Mirande, C., and Tassin, B.: Synthetic fibers in atmospheric fallout:  
384 a source of microplastics in the environment?, *Marine pollution bulletin*, 104, 290-293, 2016.

385 Fijn, R. C., Van Franeker, J. A., and Trathan, P. N.: Dietary variation in chick-feeding and self-  
386 provisioning Cape Petrel Daption capense and Snow Petrel Pagodroma nivea at Signy Island,  
387 South Orkney Islands, Antarctica, *Marine Ornithology*, 40, 81-87, 2012.

388 Geilfus, N.-X., Munson, K., Sousa, J., Germanov, Y., Bhugaloo, S., Babb, D., and Wang, F.:  
389 Distribution and impacts of microplastic incorporation within sea ice, *Marine pollution bulletin*,  
390 145, 463-473, 2019.

391 González-Pleiter, M., Edo, C., Aguilera, Á., Viúdez-Moreiras, D., Pulido-Reyes, G., González-Toril,  
392 E., Osuna, S., de Diego-Castilla, G., Leganés, F., and Fernández-Piñas, F.: Occurrence and  
393 transport of microplastics sampled within and above the planetary boundary layer, *Science of*  
394 *The Total Environment*, 2020a. 143213, 2020a.

395 González-Pleiter, M., Edo, C., Velázquez, D., Casero-Chamorro, M. C., Leganés, F., Quesada, A.,  
396 Fernández-Piñas, F., and Rosal, R.: First detection of microplastics in the freshwater of an  
397 Antarctic Specially Protected Area, *Marine Pollution Bulletin*, 161, 111811, 2020b.

398 Halsband, C. and Herzke, D.: Plastic litter in the European Arctic: What do we know?, *Emerging*  
399 *Contaminants*, 5, 308-318, 2019.

400 Ibañez, A., Morales, L., Torres, D., Borghello, P., Haidr, N., and Montalti, D.: Plastic ingestion risk  
401 is related to the anthropogenic activity and breeding stage in an Antarctic top predator seabird  
402 species, *Marine Pollution Bulletin*, 157, 111351, 2020.

403 Isobe, A., Uchiyama-Matsumoto, K., Uchida, K., and Tokai, T.: Microplastics in the southern  
404 ocean, *Marine Pollution Bulletin*, 114, 623-626, 2017.

405 Jones-Williams, K., Galloway, T., Cole, M., Stowasser, G., Waluda, C., and Manno, C.: Close  
406 encounters-microplastic availability to pelagic amphipods in sub-antarctic and antarctic surface  
407 waters, *Environment International*, 140, 105792, 2020.

408 Kelly, A., Lannuzel, D., Rodemann, T., Meiners, K., and Auman, H.: Microplastic contamination in  
409 east Antarctic sea ice, *Marine Pollution Bulletin*, 154, 111130, 2020.

410 Klein, M. and Fischer, E. K.: Microplastic abundance in atmospheric deposition within the  
411 Metropolitan area of Hamburg, Germany, *Science of the Total Environment*, 685, 96-103, 2019.

412 La Daana, K. K., Gardfeldt, K., Krumpfen, T., Thompson, R. C., and O'Connor, I.: Microplastics in  
413 sea ice and seawater beneath ice floes from the Arctic Ocean, *Scientific reports*, 10, 1-11, 2020.

414 Lacerda, A. L. d. F., Rodrigues, L. d. S., Van Sebille, E., Rodrigues, F. L., Ribeiro, L., Secchi, E. R.,  
415 Kessler, F., and Proietti, M. C.: Plastics in sea surface waters around the Antarctic Peninsula,  
416 *Scientific reports*, 9, 1-12, 2019.

417 Laganà, P., Caruso, G., Corsi, I., Bergami, E., Venuti, V., Majolino, D., La Ferla, R., Azzaro, M., and  
418 Cappello, S.: Do plastics serve as a possible vector for the spread of antibiotic resistance? First  
419 insights from bacteria associated to a polystyrene piece from King George Island (Antarctica),  
420 *International journal of hygiene and environmental health*, 222, 89-100, 2019.

421 Lenihan, H. S., Oliver, J. S., Oakden, J. M., and Stephenson, M. D.: Intense and localized benthic  
422 marine pollution around McMurdo Station, Antarctica, *Marine Pollution Bulletin*, 21, 422-430,  
423 1990.

424 Materić, D. a., Kasper-Giebl, A., Kau, D., Anten, M., Greilinger, M., Ludewig, E., van Sebille, E.,  
425 Röckmann, T., and Holzinger, R.: Micro- and nanoplastics in Alpine Snow: a new method for  
426 chemical identification and (semi) quantification in the nanogram range, *Environmental science  
427 & technology*, 54, 2353-2359, 2020.

428 Munari, C., Infantini, V., Scoconi, M., Rastelli, E., Corinaldesi, C., and Mistri, M.: Microplastics in  
429 the sediments of Terra Nova Bay (Ross Sea, Antarctica), *Marine pollution bulletin*, 122, 161-165,  
430 2017.

431 NOAA, 2019. What is the cryosphere?. National Ocean Service website, 290  
432 <https://oceanservice.noaa.gov/facts/cryosphere.html>

433 Obbard, R. W., Sadri, S., Wong, Y. Q., Khitun, A. A., Baker, I., and Thompson, R. C.: Global  
434 warming releases microplastic legacy frozen in Arctic Sea ice, *Earth's Fu  
435 ture*, 2, 315-320, 2014.

436 Österlund, H., Renberg, L., Nordqvist, K., and Viklander, M.: Micro litter in the urban  
437 environment: sampling and analysis of undisturbed snow, 2019.

438 Peeken, I., Pimpke, S., Beyer, B., Gütermann, J., Katlein, C., Krumpfen, T., Bergmann, M.,  
439 Hehemann, L., and Gerdts, G.: Arctic sea ice is an important temporal sink and means of  
440 transport for microplastic, *Nature communications*, 9, 1-12, 2018.

441 Reed, S., Clark, M., Thompson, R., and Hughes, K. A.: Microplastics in marine sediments near  
442 Rothera research station, Antarctica, *Marine pollution bulletin*, 133, 460-463, 2018.

443 Rignot, E., Mouginot, J., Scheuchl, B., Van Den Broeke, M., Van Wesseem, M. J., and Morlighem,  
444 M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proceedings of the  
445 National Academy of Sciences*, 116, 1095-1103, 2019.

446 Roblin, B., Ryan, M., Vreugdenhil, A., and Aherne, J.: Ambient Atmospheric Deposition of  
447 Anthropogenic Microfibers and Microplastics on the Western Periphery of Europe (Ireland),  
448 *Environmental Science & Technology*, 54, 11100-11108, 2020.

449 Sander, M., Costa, E. S., Balbão, T. C., Carneiro, A. P. B., and dos Santos, C. R.: Debris recorded in  
450 ice free areas of an Antarctic Specially Managed Area (ASMA): Admiralty Bay, King George Island,  
451 Antarctic Peninsula, *Neotropical Biology and Conservation*, 4, 36-39, 2009.


452 Sfriso, A. A., Tomio, Y., Rosso, B., Gambaro, A., Sfriso, A., Corami, F., Rastelli, E., Corinaldesi, C.,  
453 Mistri, M., and Munari, C.: Microplastic accumulation in benthic invertebrates in Terra Nova Bay  
454 (Ross Sea, Antarctica), *Environment international*, 137, 105587, 2020.

455 Shepherd, A., Fricker, H. A., and Farrell, S. L.: Trends and connections across the Antarctic  
456 cryosphere, *Nature*, 558, 223-232, 2018.


457 Simoes, C. L., Rosa, K. K. d., Czapela, F. F., Vieira, R., and Simoes, J. C.: Collins Glacier retreat  
458 process and regional climatic variations, King George Island, Antarctica, *Geographical Review*,  
459 105, 462-471, 2015.

460 Suaria, G., Perold, V., Lee, J. R., Lebouard, F., Aliani, S., and Ryan, P. G.: Floating macro-and  
461 microplastics around the Southern Ocean: Results from the Antarctic Circumnavigation  
462 Expedition, *Environment international*, 136, 105494, 2020.  
463 von Friesen, L. W., Granberg, M. E., Pavlova, O., Magnusson, K., Hassellöv, M., and Gabrielsen,  
464 G. W.: Summer sea ice melt and wastewater are important local sources of microlitter to  
465 Svalbard waters, *Environment international*, 139, 105511, 2020.  
466 Wright, S., Ulke, J., Font, A., Chan, K., and Kelly, F.: Atmospheric microplastic deposition in an  
467 urban environment and an evaluation of transport, *Environment international*, 136, 105411,  
468 2020.  
469 Zhang, M., Haward, M., and McGee, J.: Marine plastic pollution in the polar south: Responses  
470 from Antarctic Treaty System, *Polar Record*, 56, 2020.


471  
472  
473  
474  
475  
476  
477  
478  
479  
480



481  
 482 **Figure 1.** (A) General view of Antarctica and location of King George Island. The blue  
 483 circle indicates the Fildes Peninsula. Collins Glacier is located on the northeast of Fildes  
 484 Peninsula. (B) A detailed view of Ionosferico lake, Uruguay lake, Artigas Research Station  
 485 and Collins Glaciers in the Fildes Peninsula. (C) and (D) ablation zone of Collins Glacier  
 486 close to Ionosferico lake and Uruguay lake, respectively. (E) Photograph of the glacier  
 487 surface close to Uruguay lake that constitute part of the ablation zone of Collins Glacier  
 488 taken on 18/02/2020. (F) A representative square on the glacier surface used in this  
 489 study.




490  
 491 **Figure 2.** (A) Size of the plastics collected on the glacier surface. (B) Total number of the  
 492 mesoplastics and microplastics found on the glacier surface close to Uruguay lake and  
 493 lonosferico. Representative photographs of expanded polystyrene (C), polyester (E) and  
 494 polyurethane (G) found on the glacier surface. The red arrows indicate the plastics. FTIR  
 495 representative spectra of expanded polystyrene (D), polyester (F) and polyurethane (H)  
 496 found on the glacier surface.



497  
498

499 **Figure 3.** Changes in the presence of plastics into the squares marked on ice surface close to Uruguay lake (A) and close to Ionosferico lake (B)  
500 that constitute part of the ablation zone of Collins Glacier in Maxwell Bay in King George Island (Antarctica). Plastics were monitored every 12 hours  
501 for two days (18/2/2020 and 20/2/2020) in the absence of rainfall. Asterisks indicate squares different from those used to the assessment of  
502 plastic concentration.



503  
504 **Figure 4.** Wind Roses obtained for the area of BCAA based on historical data of the  
505 Uruguayan National Institute of Meteorology (January 1998 - May 2016; 24,698  
506 records). Based on the speed of winds considered (A) and (B) refer to *Winds* and *Strong*  
507 *winds*, and (C) and (D) to *Wind Gusts* and *Strong wind gusts*, respectively.