Improved ELMv1-ECA Simulations of Zero-Curtain Periods and Cold-season CH₄ and CO₂ Emissions at Alaskan Arctic Tundra Sites

Jing Tao^{1,2}, Qing Zhu¹, William J. Riley¹, Rebecca B. Neumann²

¹Climate and Ecosystem Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA, 94720, USA

5 ²Department of Civil and Environmental Engineering, University of Washington, Seattle, WA, 98195, USA

Correspondence to: Jing Tao (JingTao@lbl.gov)

Abstract. Field measurements have shown that cold-season methane (CH₄) and carbon dioxide (CO₂) emissions contribute a substantial portion to the annual net carbon emissions in permafrost regions. However, most earth system land models do not accurately reproduce cold-season CH₄ and CO₂ emissions, especially over the shoulder (i.e., thawing and freezing) seasons.

- 10 Here we use the Energy Exascale Earth System Model (E3SM) land model version 1 (ELMv1-ECA) to tackle this challenge and fill the knowledge gap of how cold-season CH₄ and CO₂ emissions contribute to the annual totals at Alaska Arctic tundra sites. Specifically, we improved the ELMv1-ECA soil water phase-change scheme, environmental controls on microbial activity, and cold-season methane transport module. Results demonstrate that both soil temperature and the duration of zerocurtain periods (i.e., the fall period when soil temperatures linger around 0°C) simulated by the updated ELMv1-ECA were
- 15 greatly improved, e.g., the Mean Absolute Error (MAE) in zero-curtain durations at 12 cm depth was reduced by 62% on average. Furthermore, the MAE of simulated cold-season carbon emissions at three tundra sites were improved by 72% and 70% on average for CH₄ and CO₂, respectively. Overall, CH₄ emitted during the early cold season (Sep. and Oct.), which often includes most of the zero-curtain period in Arctic tundra, accounted for more than 50% of the total emissions throughout the entire cold season (Sep. to May) in the model, compared with around 49.4% (43-58%) in observations. From 1950 to 2017,
- 20 both CO_2 emissions during the zero-curtain period and during the entire cold season showed increasing trends, for example, of 0.17 gC m⁻² year⁻¹ and 0.36 gC m⁻² year⁻¹ at Atqasuk. This study highlights the importance of zero-curtain periods in facilitating CH₄ and CO₂ emissions from tundra ecosystems.

1 Introduction

- 25 Cold-season carbon emissions from the Arctic tundra could potentially offset warm-season net carbon uptake under 21st century warming climate (Commane et al., 2017; Oechel et al., 2014; Oechel et al., 2000; Koven et al., 2011; Piao et al., 2008; Natali et al., 2019; Belshe et al., 2013; Fahnestock et al., 1998; Jones et al., 1999). Field measurements have indicated large cold-season CO₂ losses over Arctic tundra ecosystems (Oechel et al., 2014; Natali et al., 2019). Also, CH₄ emitted from September to May were found to contribute more than 50% of the annual total CH₄ emissions from Alaska upland tundra sites
- 30 (Zona et al., 2016; Taylor et al., 2018). Despite the importance of cold-season carbon emissions and their sensitivity to changing climate, prevailing earth system land models do not accurately reproduce cold-season CH₄ and CO₂ emissions and their contributions to the annual budgets, largely because of the poorly understood mechanisms of cold-season soil heterotrophic respiration and therefore uncertain numerical representations (Natali et al., 2019; Zona et al., 2016; Wang et al., 2019; Commane et al., 2017). Thus, it remains challenging to assess the response of permafrost carbon dynamics to Arctic
- 35 warming and to predict future annual carbon budgets with current Earth System Models (ESMs).

In ESM land models, soil environment influences soil microbial heterotrophic respiration (HR) and decomposition of soil organic carbon (SOC) mainly through applying prescribed temperature and moisture functions to modify base decomposition rates. These functions, however, rely heavily on empirical or semi-empirical relationships which are highly uncertain (Sierra et al., 2017; Sierra et al., 2015; Yan et al., 2018; Moyano et al., 2013; Tang and Riley, 2019; Rafique et al., 2016; Bhanja and

- Wang, 2020; Kim et al., 2019). Specifically, the temperature sensitivities of soil carbon decomposition is often represented with a Q_{10} value (i.e., the increase in respiration rate from a 10°C increase in temperature) that is fixed at 1.5 or 2.0 (Meyer et al., 2018). However, the values of Q_{10} are controversial (Davidson and Janssens, 2006). Some studies found a uniform Q_{10} across biomes and climate zones, e.g., as 1.4 (Mahecha et al., 2010). Other studies demonstrated that Q_{10} varies with
- 45 environmental conditions, ecosystem types, and soil texture (Meyer et al., 2018; Graf et al., 2011; Kim et al., 2019), showing a large spatial heterogeneity with generally higher values in the high-latitudinal regions (Zhou et al., 2009). In addition, Wilkman et al. (2018) reported a temporal heterogeneity in Q₁₀ over the Alaskan Arctic Tundra and suggested a higher value (e.g., 2.45) for early summer (e.g., June) but lower value (e.g., 1.58 to 1.67) for the peak growing season (e.g., July). Dynamic decomposition temperature sensitivities are also consistent with theory of microbial dynamics (Tang and Riley, 2015). Also,
- 50 the response of HR to changes in soil moisture is commonly expressed by empirical relationships in ESMs, which vary substantially (Sierra et al., 2015; Yan et al., 2018; Moyano et al., 2013). Although in-situ measurements reveal that microbial respiration occurs under very cold conditions (e.g., even when soil temperature is lower than -15 °C) (Natali et al., 2019; Zona et al., 2016), most process-based models completely shut down microbial activity due to limited liquid water in freezing and subfreezing soils, and few modelling studies have closely investigated the HR-moisture relationships in frozen conditions.

55

The strong dependency of CO_2 and CH_4 emission on soil temperature and moisture in ESM land models (Riley et al., 2011; Koven et al., 2017; Lawrence et al., 2015) requires accurate estimates of these two closely related soil variables, especially in cold regions where both increases and decreases in soil temperature could lead to soil "drying" due to drainage or freezing processes. However, current land models tend to significantly underestimate soil temperature during the cold season over

- 60 permafrost regions (Dankers et al., 2011; Tao et al., 2017; Nicolsky et al., 2007; Yang et al., 2018b). One possible reason is that while many land models account for latent heat released during soil water freezing, they do not treat and distribute this heat appropriately or/and do not simulate soil moisture correctly (Yang et al., 2018a; Nicolsky et al., 2007). Latent heat released during freezing might be sufficient to offset heat conduction towards the surface, thus maintaining the subsurface soil temperature around the freezing point (i.e., 0°C) for weeks or even months during the fall (i.e., the so-called Zero-Curtain
- 65 Period; ZCP) (Outcalt et al., 1990). The ZCP conditions allow for continued soil heterotrophic respiration at notable rates, and thus CO₂ and CH₄ production and emissions from subsurface soils (Kittler et al., 2017; Arndt et al., 2019; Commane et al., 2017). For instance, Zona et al. (2016) reported that a substantial portion of cold season CH₄ emissions occurred during the ZCP from Alaskan upland tundra sites. Nevertheless, many land models cannot accurately capture the ZCP length due to inaccurately simulated soil moisture and/or inadequate representation of latent heat, thus underestimating soil temperature and
- 70 cold-season emissions of CO₂ (Commane et al., 2017) and CH₄ (Zona et al., 2016). We note that snow representation can also play a major role in correctly simulating winter soil temperatures (Slater et al., 2017; Lawrence and Slater, 2010), although we do not focus on this process here.

We hypothesize that the underestimation of modelled cold-season CO₂ and CH₄ emissions in ESMs land models primarily results from underestimated soil temperatures during the cold season, the poor representations of environmental controls on heterotrophic respiration in subfreezing soils, and the lack of appropriate representation of cold-season methane transport processes. Here we apply the Energy Exascale Earth System Model (E3SM) land model version 1, the Equilibrium Chemistry Approximation configuration (ELMv1-ECA) (Golaz et al., 2019; Zhu et al., 2019; Burrows et al., 2020) to explore these hypotheses. We apply ELMv1-ECA to (i) improve simulations of subsurface soil temperatures, ZCPs, and CO₂ and CH₄ emissions over the permafrost tundra ecosystem; (ii) investigate the underlying processes that influence cold-season carbon emissions from freezing and subfreezing soils, including source characterization and transport pathways; and (iii) estimate

historical trends (from 1950 to present) of cold-season CO_2 and CH_4 emissions at multiple Alaskan tundra sites.

The paper is organized as follows: (1) We describe the study sites and the data used in the study. (2) We present the theoretical background of essential modules of ELMv1-ECA relevant to this study and our modifications to the model's representations of phase-change, SOC decomposition, and methane dynamics. (3) We then describe the model configuration and experimental design. (4) We assess the modified phase-change scheme by comparing simulated soil temperatures and ZCPs against observations. (5) With the revised phase-change scheme, we analyze how the parameterization of decomposition schemes and methane module impact simulated CO₂ and CH₄ emissions at the site scale. (6) Finally, we summarize the main findings and

90 discuss needed observations and model development to further improve predictability.

2 Study Sites and Data

We assembled daily observations of CO₂ and CH₄ fluxes from 2013 to 2017 at five eddy-covariance flux tower sites in Alaska's North Slope tundra (Figure 1) from the Arctic-Boreal Vulnerability Experiment (ABoVE) project (2015 - 2017) (Oechel and Kalhori, 2018) and CH₄ fluxes from the Carbon in Arctic Reservoirs Vulnerability Experiment (CARVE) flight campaign (2013 - 2014) (Zona et al., 2016). The CARVE CO₂ measurements were not available at the data archive we used here; therefore, monthly winter-time CO₂ flux data from 2013 to 2014 at the same towers assembled by Natali et al. (2019) are included to complete CO₂ observations. The five sites include three eddy covariance (EC) towers at Barrow (i.e., the Barrow Environmental Observatory (BEO) tower, the Biocomplexity Experiment South (BES) tower, and the Climate Monitoring and Diagnostics Laboratory (CMDL) tower), one tower at Atqasuk (ATQ) and another at Ivotuk (IVO) which is located at the foothills of the Brooks Range. BES and CMDL are collocated with each other with sensors installed at different heights (i.e., 2 m for BES and 5 m for CMDL). Vegetation at Barrow is mainly moist acidic tundra. Instrument height at ATQ and IVO is 2 m and 4 m, respectively. ATQ is a well-drained upland site, and the vegetation consists of moist-wet coastal sedge tundra and moist-tussock tundra surfaces. Vegetation at IVO is polar tundra. Table S1 provides basic information including geolocations, vegetation mosaic, and climatologic air temperature at the sites. (Tables numbered with a prefix "S" are include

105 in the supplementary file, which will not be repeated in the following context throughout the manuscript.)

ABoVE and CARVE provide soil temperature and moisture measurements at various depths from 5 cm to 40 cm. The Permafrost Laboratory, Geophysical Institute of University of Alaska Fairbanks (GIPL-UAF), provides daily subsurface soil temperature observations down to various depths at permafrost sites across Alaska(http://permafrost.gi.alaska.edu/sites_map) (Romanovsky et al., 2009). We used the GIPL-UAF permafrost sites that are collocated with the ABoVE sites to complement the ABoVE observations at deeper depths, including BR2 (down to 15 m) and IV4 (down to 1 m). We first filled missing gaps vertically by fitting a polynomial to the soil temperature profile (Kurylyk and Hayashi, 2016) on a daily scale, then screened out outliers by examining the daily time series. Further, we aggregated both the ABoVE and the GIPL-UAF soil temperature measurements to ELMv1-ECA soil layer node depths using the inverse distance weighting method (Tao et al., 2017), and then averaged the two sets of aggregated observations. We used the assembled subsurface temperature observation datasets to evaluate the ELMv1-ECA simulated soil temperature profiles and the zero-curtain periods.

The observed soil moisture is only available at two or three depths that are quite different from model layer node-depths, and also show discontinuities in time. Thus, evaluation of ELMv1-ECA simulated liquid water content was limited. We matched

liquid water content at layers for time periods during which observations were available. In addition, we used ABoVE observed maximum soil moisture to infer site-scale soil porosity and then organic carbon content at IVO (see Section 3.2), which is used to prescribe thermal and hydraulic soil properties. Note that carbon substrate for respiration is simulated dynamically in the model (see Appendix B).

3 Methodology

3.1 Modifications to E3SM Land Model (ELM)

The E3SM land model version 1 (ELMv1-ECA) couples essential biogeophysical and biogeochemical processes that solve terrestrial ecosystem energy, water, carbon, and nutrient dynamics (Golaz et al., 2019; Zhu et al., 2019). In the appendix, we
describe in detail its subsurface soil thermodynamics, the carbon decomposition module, and the methane module that are of particular relevance to our study. Here we identify the potential problems of ELMv1-ECA that are responsible for the underestimation of cold-season CH₄ and CO₂ emissions and summarize the modifications made to ELMv1-ECA, emphasizing the model enhancements.

3.1.1 Phase Change Scheme

- 135 We first improved ELMv1-ECA's numerical representation of coupled water and heat transport with freeze-thaw processes via improving the phase-change scheme. The freeze-thaw processes of soil water within ELMv1-ECA is simulated in a decoupled way, i.e., it solves soil temperatures ignoring the latent heat associated with phase change, determines the mass change of soil water required to adjust the initially solved soil temperature to the freezing point (i.e., 0°C; T_f), adjusts the soil liquid and ice content by mass and energy conservation, and then readjusts temperatures after accounting for the heat deduction
- or compensation resulted from melting or freezing (see the detailed description in the Appendix A). The underlying assumption here is, taking the freezing process as an example, the available liquid water at the initially solved temperature (T_i^{n+1}) will be completely frozen, releasing latent heat (H_i) to bring up T_i^{n+1} back to T_f . Then, the estimated phase-change rate will be tuned down and the current temperature (i.e., T_f) will be readjusted if the to-be-increased ice mass is larger than the required mass change $(-H_m)$ (see (Eq. A4) in the Appendix A), which, however, only occasionally occurs. When the liquid water available
- to be frozen becomes small enough, the released latent heat is not sufficient to compensate for the required energy deficit $(T_f T_i^{n+1})$, and then the freezing process stops. Consequently, the model freezes soil water quickly, resulting in an underestimated duration of the soil water phase-change processes and the zero-curtain periods, and also cold-biased winter temperatures (Nicolsky et al., 2007; Yang et al., 2018a).

- Here, we employed a phase-change efficiency and the temperature of the freezing-point depression to effectively solve the problem of overestimating phase-change rates within the current ELMv1-ECA modelling structure. These modification factors are explained below. The phase-change efficiency, introduced by Le Moigne et al. (2012) and adopted by Masson et al. (2013) and Yang et al. (2018a), introduces the dependency of available liquid water on the phase-change rate (Le Moigne et al., 2012). The phase-change efficiency for freezing, εⁿ_{liq,i} (see (Eq. A9)), is identical to the degree of moisture saturation, or the volumetirc fraction of soil liquid water content (i.e., Sfⁿ_{liq,i} = θⁿ_{liq,i}/θ_{sat,i} where θⁿ_{liq,i} is soil liquid water content and θ_{sat,i} is
- porosity). We applied the phase-change efficiency to the initially estimated energy and mass change involved, i.e., H_i and H_m (see (Eq. A4) in the Appendix) when freezing or thawing process occur.
- As in Nicolsky et al. (2007) and Yang et al. (2018a), the occurrence of a phase-change process is then determined by the 160 temperature of the freezing point depression (i.e., an virtual temperature, see (Eq. A10)) instead of T_f . The virtual freezing point depression temperature is reversely derived from the freezing point temperature-depression equation (Fuchs et al., 1978; Cary and Mayland, 1972). With an upper limit as T_f , the virtual temperature describes the lowest temperature that can hold current liquid water content in the freezing soils. That is, the soil temperature has to be lower than the current virtual temperature to allow the freezing process to occur further.

165

170

175

We describe in detail the revised phase-change scheme in the Appendix A. In short, we improved the phase-change scheme of ELMv1-ECA by incorporating two modifications: 1) applying a phase-change efficiency to implicitly account for the heat compensation/deduction to the system from latent heat released/absorbed by soil water freezing/melting, and 2) replacing the constant freezing point with the temperature of the freezing point depression, as a virtual temperature, to determine the occurrence of phase change in subfreezing soils.

3.1.2 Environmental Modifiers to the Decomposition Rate

We revisited ELMv1-ECA's representation for soil heterotrophic respiration dynamics in subfreezing soils and then scrutinized the environmental scalars of soil temperature and moisture. Within ELMv1-ECA's decomposition cascade model, the environmental factors that impact the decomposition rates of soil organic matter include soil temperature (f_T) , soil moisture (f_W) , oxygen stress (f_0) and a depth scalar (f_D) (See Appendix B). Within freezing and subfreezing soils, the soil water

- potential is related to temperature through the freezing point depression equation (Niu and Yang, 2006). The current moisture factor f_W , therefore, predicts zero respiration rates for subfreezing soils given a specific lower limit of soil water potential ψ_{min} (-10 MPa; (Eq. B13)) (Oleson et al., 2013), as shown by Figure S1a in the supplementary file. We thus decreased the ψ_{min} further to prevent zero respiration within the active layer when soil becomes subfreezing during cold-season months (Figure S1b) as large as the soil water potential the support of the prescribed of
- 180 (Figure S1b) as long as the soil water potential ψ_i exceeds the prescribed ψ_{min} .

For wet soils, the factor that primarily limits the decomposition rates is oxygen availability (Sierra et al., 2017; Yan et al., 2018), since increases in soil moisture result in decreased dissolved oxygen. ELMv1-ECA approximates oxygen stress (f_0) as a ratio of available oxygen to the demand by decomposers, which, however, is highly uncertain and unstable (Oleson et al.,

- 185 2013). Some existing moisture scalars incorporate the oxygen stress together to account for the inhibition of decomposition in wet anoxic conditions, e.g., a moisture function proposed by Yan et al. (2018) and several functions tested in Sierra et al. (2015), including Standcarb (Harmon and Domingo, 2001), Daycent (Kelly et al., 2000), Skopp (Skopp et al., 1990), and Moyano (Moyano et al., 2013). We thus also tested these existing moisture functions by replacing the original moisture scalar with them in the ELMv1-ECA. Particularly for the moisture function of Yan et al. (2018), we implemented it for each soil
- 190 layer using the soil properties (i.e., porosity and clay content) of each layer, and also tested it with different shape parameters and optimal wetness thresholds. When using the moisture scalars with built-in oxygen stress within ELMv1-ECA, the total environmental impacts on decomposition, *i.e.*, $f_{total} = f_T f_W f_O f_D$ will be modified accordingly as $f_{total} = f_T f_W f_D$ to avoid double-counting of the oxygen stress.
- 195 ELMv1-ECA uses a Q₁₀-based standard exponential function to account for the temperature effect on SOC decomposition (Eq. B12), with Q_{10} as 1.5 and T_{ref} as 25°C. Here, rather than striving for a single value of Q_{10} or a spatial map of Q_{10} as discussed in the introduction, or a particular individual temperature function, we seek a group of environmental modifiers (f_{total}) that can correctly represent moisture and temperature sensitivity on heterotrophic respiration. Specifically, we assembled and tested 814 cases of f_{total} using the newly modified moisture scalars and a variety of other empirical moisture and temperature 200 functions, as documented by Sierra et al. (2015) and Yan et al. (2018). A full list of the specific moisture and temperature scalars tested is provided in Table S2.

3.1.3 Cold-season Methane Process

- The ELMv1-ECA methane model solves the reaction and diffusion equation for CH₄ and O₂ fluxes with the Crank-Nicholson 205 method. It includes the representations of CH₄ production, oxidation, and three pathways of transport, including aerenchyma tissues, ebullition, aqueous and gaseous diffusion (Riley et al. (2011)). A short description of the ELMV1-ECA methane module is provided in Appendix C. The ELMv1-ECA methane model has been found to underestimate cold-season methane emissions over northern wetlands (Xu et al., 2016). The modifications to the phase-change scheme impact simulations of soil water and heat transfer (3.1.1); the changes in environmental scaler affect substrate availability (3.1.2). Both (3.1.1) and (3.1.2)
- 210 influence soil heterotrophic respiration, and could potentially lead to improvements in simulated CO₂ and CH₄ production, but not necessarily CH₄ emissions which are also controlled by oxidation and transport mechanisms.

Here, we first modified the ELMV1-ECA CH₄ transport mechanism in cold seasons by mimicking possible pathways for CH₄ emissions from freezing and subfreezing soils. Specifically, we mimicked the emissions from ice cracks by plant aerenchyma

- 215 transport (Zona et al., 2016), approximating the gas diffusion through ice cracks to the similar mechanism of diffusion through the aerenchyma tissues. Although in-situ experiments demonstrated that during winter, produced CH₄ in frozen soils is predominately emitted to the atmosphere through vascular plants aerenchyma tissues (e.g., Kim et al., 2007), here we integrate the possible transport pathways including ice cracks and remnants of aerenchyma tissues together through equation (Eq. C16). Also, during the cold season over the tundra ecosystem, snow on the land surface provides strong resistance to CH₄ transport
- 220 to the atmosphere in ELMv1-ECA. But in reality, studies have shown methane can diffuse through snowpack at varying rates (Kim et al., 2007). We thus decreased snow resistance at the upper boundary by introducing a new scale factor when snow is present (Appendix C).

Table 1 summarized all the specific modifications made to ELMv1-ECA. These modifications involve new parameters that are all tuneable and can be systematically optimized via calibration. Here, we seek to reproduce the first-order cold-season process relevant to this study with these default formation and values listed in Table 1. We also conducted sensitivity tests on three variables three key parameters related to CH₄ oxidation and transport processes and tested seven parameterizations (Table S3). The three CH₄ process-related parameters include two key variables in the original CH₄ model that have been reported having large uncertainty (Riley et al., 2011), i.e., f_{CH_4} (a fraction of anaerobically mineralized carbon atoms becoming CH₄;

Eq. C14) and $R_{o,max}$ (the maximum oxidation rate constant; Eq. C17), and the newly introduced variable ε_{aere} (a factor representing remnants of aerenchyma tissues during cold seasons and possible pathways via ice cracks; Eq. C16). The sensitivity tests on CH₄ process-related parameters were applied to model with identified carbon decomposition schemes that predicted good simulations of CO₂ flux (see section 3.3).

235 3.2 Climate Forcing, Model Configuration, and Experiment Design

We conducted transient simulations at 30-minute temporal resolution driven by climate forcing from 0.5°×0.5° CRU JRA (Harris, 2019) from 1901 to 2017 at the four Alaska tundra site locations. Before the transient simulation, we conducted a 200year Accelerated Decomposition (AD) spin-up period followed by a 200-year regular spin-up period (Koven et al., 2013b; Zhu et al., 2019) to initialize land carbon pools. Spin-up simulations start from a wet and cold condition. Specifically, sub-surface temperatures were initialized as 274 K for the 1st to 5th soil layers, 273 K for the 6th to 10th layer, and 272 K for the 11th to the 15th layer, and volumetric soil water content was initialized fully saturated for all layers. In this manner, consistent vertical soil water content profiles were built in over the permafrost regions.

Baseline simulations were conducted with ELMv1-ECA default physics, parameters, and surface datasets, i.e., OriPC_OriDecom_OriCH4 using original phase-change scheme, original decomposition scheme and methane module (Table 2). To improve the model representation of the site-level soil environment, we first examined the global soil organic matter data at the ABoVE sites by evaluating ELMv1-ECA simulated subsurface soil temperature with the topsoil temperature prescribed to observations (as did in Tao et al., 2017). Using the top soil layer as the upper boundary, the modelling system excluded potential errors induced by inaccurate meteorological forcing and vegetation cover that impact the simulation of heat

- transfer from the atmosphere to the shallow soil (Tao et al., 2017). Then, the accuracy of simulated soil subsurface temperature is directly determined by the factors impacting heat transfer along the "shallow-to-deep soil" gradient (Koven et al., 2013a), e.g., soil thermal properties which are mostly determined by SOC content (Tao et al., 2017; Lawrence and Slater, 2008). Results well reproduced the subsurface soil temperatures except at IVO, where summer soil temperatures were notably overestimated (see Figure S2a). This result indicates that the SOC content at IVO was too small, leading to a large thermal
 conductivity, small soil porosity, and small heat capacity, altogether resulting in fast penetration of heat into the subsurface soil during summer (Tao et al., 2017; Lawrence and Slater, 2008). Thus, we derived the organic matter density at IVO based on ABoVE soil moisture data through a linear relationship between SOC content was porosity (i.e., Equation 3 in Lawrence and Slater (2008)), assuming the observed maximum volumetric water content was porosity (see Figure S3 for details). With the newly derived profile of soil organic matter density at IVO, the simulation showed large improvements in summer soil
- consistent with the organic layer thickness reported in Davidson and Zona (2018). Hereafter, the simulations at IVO presented in this paper use the newly derived organic carbon data without repeated clarification.

The representative spatial scale of the eddy flux tower is small compared to the grid cell of global surface datasets and the climate forcing data used by ELMv1-ECA, although the forcing dataset was interpolated to the site scale with a bilinear or nearest-neighbor method. The site-scale vegetation cover also shows a large diversity of vegetation types according to the detailed vegetation survey at ABoVE flux tower footprints obtained in 2014 (Davidson and Zona, 2018). The ELMv1-ECA's default plant type function (PFT) dataset was derived from satellite-based data by Lawrence et al. (2007). We analyzed the

vegetation composition from the closet survey plot to the flux tower and examined the rationality of ELMv1-ECA's percentage

- 270 of PFT for the site-scale simulation through testing different PFT datasets derived from this vegetation survey (Davidson and Zona, 2018). We found that these PFT datasets generally are not superior to the original PFT dataset, which generally reproduced satellite-based GPP (Figure S4). We thus confirmed that ELMv1-ECA's PFT dataset was a good compromise between representing the site-scale ecosystem and other global parameters and surface datasets within the model. The surface CH₄ emission is a weighted average of simulated saturated and unsaturated components using predicted inundation and non-
- inundation fractions. To compare simulated CH₄ emissions with ABoVE measurements at the site scale, we use the estimated inundation fractions at the footprint of ABoVE eddy-covariance flux towers (see details in (Xu et al., 2016)).

Table 2 lists the experiments conducted in this study. We modified each model component (i.e., the heat transfer model, carbon decomposition model, and methane model) serially. All the experiments ran through 1901 to 2017 with spin up as described earlier, although the evaluation and optimization were conducted only using results from 2013 to 2017. We first ran simulations

with the 814 environmental modifiers together with the modified methane model with default parametrization (Table S3).

Then, we selected the environmental modifiers that provided satisfactory performance in simulating CO_2 flux, and repeated simulations with the seven CH_4 parameterizations (Table S3). Among all the simulations results, we identified an optimal simulation for each site (see details in section 3.3).

285

3.3 Evaluation Metrics, Optimization Method, and Trend Analysis

We define the early cold season as September and October, the cold-season period as September to May which includes the two shoulder seasons (both thawing and freezing) as consistent with Zona et al. (2016), and the warm season from June to August. We define the zero-curtain period (ZCP) as the set of successive days when the soil temperature is within the range of [-0.75°C, 0.75°C] starting in fall (i.e., the freezing season) based on Zona et al. (2016). We computed the ZCP duration for each soil layer every year from 1950 to 2017 and estimated the historical trend as the regression slope between ZCP duration and time. Similarly, we estimated the trends of cold-season CH₄ and CO₂ emissions through linear regression analysis. A p-value of 0.05 is used to determine if the computed trend is statistically significant. Results for ZCP duration and trend vary with soil depths; thus, we choose a common modelling depth at which the ZCPs show significant trends for all the sites, to give an example.

To evaluate ELMv1-ECA simulated soil temperature and moisture, we calculated the RMSE for each soil layer, i.e., $\sqrt{\sum_{t=1}^{N} (\hat{E}_t - O_i)^2 / N}$ where the \hat{E}_t and O_t is simulated and observed soil temperature or moisture, respectively, and t is a daily time step. We used the Mean Absolute Error (MAE, *i.e.*, $\frac{1}{N} \sum_{t=1}^{N} |\hat{E}_t - O_i|$ to assess the simulated duration of ZCP of each soil layer. Note that, depending on the amount of soil liquid water content, the whole course of the freezing process may or may not entirely fall into the ZCP, i.e., the ending time of ZCP does not necessarily align with the end of the freezing process. The onset of freezing, though, is always later than the starting day of the ZCP, and the main course of the freezing process is still within the ZCP.

- 305 Here the modelled active layer thickness (ALT), i.e., maximum thaw depth during an annual cycle, is computed as the bottom depth of the deepest thawed soil layer (i.e., with a maximum annual temperature above 0°C) further extended down to the possible non-frozen fraction of the layer below, as in Tao et al. (2019; 2017). We only derived the length of ZCP for soil layers with a maximum annual temperature above 0°C since limited phase-change processes occur in deeper layers. Then, the soil layers containing or below the permafrost table have a zero-day ZCP. We computed the MAE of ALT simulated with both
- 310 original (OriPC) and the new phase-change (NewPC) scheme. Also, we computed the relative improvement in simulated soil temperature (Ts) and ZCP compared to the baseline results. Specifically, we calculated 100% × (RMSE_Ts_OriPC RMSE_Ts_NewPC) / RMSE_Ts_OriPC and 100% × (MAE_ZCP_OriPC MAE_ZCP_NewPC) / MAE_ZCP_OriPC to quantify the enhancement by employing the new phase-change scheme.

315 We used Nash-Sutcliffe Efficiency (NSE) (Nash and Sutcliffe, 1970) to examine the performance of the ELMv1-ECA simulated time series of CH₄ and CO₂ net fluxes in comparison with assembled observations (Section 2) at the monthly time scale. The NSE ranges from negative infinity to one, calculated as Eq. (1):

$$NSE = 1 - \left(\frac{1}{N}\sum_{t=1}^{N} (\hat{E}_t - O_t)^2\right) / \sigma_o^2,$$
(1)

where t means monthly time step, N is the total number of time steps, \hat{E}_t and O_t is simulated and observed flux at time step t, respectively; and σ_o is the standard deviation of observations. Note we only used observed monthly averages when the number

of daily observations was more than 20 days. The model performance is generally considered satisfactory with an NSE > 0.50 (Moriasi et al., 2007), and perfect with an NSE as one. To simultaneously evaluate CH₄ and CO₂ fluxes, we combined both NSE_{CH4} and NSE_{CO2} in the form of $dist = \sqrt{(1 - NSE_{CH4})^2 + (1 - NSE_{CO2})^2}$, representing the distance from (NSE_{CH4} , NSE_{CO2}) to (1, 1) in a coordinate plane with x-axis as NSE_{CH4} and y-axis as NSE_{CO2} . Then, the optimal simulation thereby is the one having the shortest distance to the ideal scenario (1, 1). We also define a satisfactory model performance in terms of simulating CH₄ and CO₂ fluxes as the case with both NSE_{CH4} and NSE_{CO2} larger than 0.5.

We optimized the model simulations through two steps. Specifically, we first evaluated the simulations using (814) environmental modifiers to the base decomposition rate that assembled commonly used empirical soil temperature- and moisture-dependency functions (Table S2). These simulations used the newly modified methane model with the default parameters (Table S3). We selected the common decomposition schemes that provided satisfactory results of CO₂ flux for all the sites (i.e., $NSE_{CO2} > 0.5$). Then, we iteratively repeated simulations with the common carbon decomposition schemes along with the seven CH₄ parameterizations (Table S3). Among all these simulations ("NewPC_NewDecomNewCH4"; Table 2), we identified an optimal simulation for each site that has the smallest distance from (NSE_{CH4} , NSE_{CO2}) to (1, 1) (i.e., *dist*);

335 the optimized parameterization for this site.

Further, among the common parameterizations of environmental modifiers and CH₄ parameterizations that show satisfactory performance both in CH₄ and CO₂ fluxes for all the sites, we identified a generic scheme as the one providing the minimum Euclidean distance in a site-performance space, calculated as $\sqrt{\sum_{i=1}^{n} dist_i^2}$ where n is number of sites. The generic scheme

the environmental modifier to the base decomposition rate and the methane parameterization used in the optimal simulation is

340 then is the common satisfactory scheme that provides the best overall performance for all the sites and can be applied for the regional simulation over Alaskan North Slope tundra. In general, we use NSE to evaluate the model's performance in capturing seasonality (i.e., time series) of CH_4 and CO_2 net fluxes and optimize CH_4 and CO_2 simulations. We use RMSE and MAE to assess the model's capability in simulating the

345 magnitudes of soil temperature, moisture saturation, ZCP durations, and cumulative CH₄ and CO₂ emissions.

4 Results and Discussion

350

4.1 Evaluation of Soil Temperature and Zero-curtain Period

We first evaluated the simulated daily soil temperature profiles against the observations from ABoVE and GIPL-UAF at the four site locations. Then, we examined improvements in simulations of soil temperature, soil moisture, and the durations of ZCPs by employing the newly revised phase-change scheme (i.e., "NewPC OriDecomOriCH4"; Table 2).

Results for the BES/CMDL and IVO site are shown in Figure 2; results for other sites are shown in supplementary Figure S4. At BES/CMDL, the baseline (i.e., "OriPC_OriDecomOriCH4"; Table 2) simulated soil temperatures (Ts) with the default phase-change scheme (Ts_OriPC; blue lines; Figure 2a) decrease rapidly in fall due to the overestimated freezing rate (i.e., the slope of decreasing liquid water fraction), notably underestimating the duration of the ZCP (bluish shaded area). Consequently, liquid water saturation (S_f_OriPC, green lines; Figure 2a) quickly drops to a lower bound (i.e., the supercooled liquid water content divided by porosity), and the freezing process generally completes within a short period (days for top layers to one month at the most for deeper layers). The baseline model soil temperature drops (Ts_OriPC) sharply after the freezing process ends (i.e., S_f_OriPC decreases to the lower bound). In contrast, the new phase-change scheme effectively slows freezing rates, showing relatively smaller slopes of decreasing liquid water saturation (S_f_OriPC; green lines) especially in the 4th and 5th layer. Hence, the gradually released latent heat maintains soil temperatures around the freezing point for a longer period (Ts_NewPC; red lines; Figure 2a), effectively

extending the ZCPs (reddish shaded area) which agree better with observations (grey shaded area) than the baseline results.
The ZCP duration increases with depth and can extend into December for deep soil layers. Similarly, improved performance
was found at the BEO and ATQ sites (supplementary Figure S4). At IVO, however, while the new phase-change scheme

- greatly improved simulated results relative to the baseline simulation (Figure 2b), the model still slightly underestimated ZCP durations and also underestimated winter (December to April) soil temperature (red vs. black). This result at IVO is consistent with the underestimation of late-season soil liquid water available to be frozen, and thereby to release sufficient latent heat (Figure S5). In general, the improvements in ZCP are larger in deeper layers than topsoils, with the top layer showing only
- 370 marginal improvement.

Simulated ZCP durations with the revised phase-change scheme (NewPC) demonstrated notable improvements over the baseline (original) phase-change scheme (OriPC) (solid circles vs. open diamonds) (Figure 3), showing greatly reduced mean absolute errors (MAEs) (Table 3). For example, at 12 cm depth (4th layer), the relative improvements in MAE of the ZCP

- 375 durations were 65%, 65%, 66%, and 50% for the four site locations (Table 3). The largest improvement in MAE was as large as 65 days for the 6th layer at BES/CMDL, with a relative improvement of 84% (Table 3). This large improvement stems from the better-estimated ALT at this site; the OriPC simulated 6th layer temperature remained below freezing, leading to a zeroday ZCP (diamonds on the x-axis in Figure 3). The new phase-change scheme not only improved simulation of the ZCP and cold-season soil temperatures, but also affected the warm season dynamics and thus ALT estimates. As Figure 3 indicates, the
- 380 NewPC improved simulated ALTs at all four site locations with reduced bias in multi-year averaged ALT, resulting in more reasonable ZCP durations for the 6th layer (and also the 7th layer for IVO), while the baseline results were zero days.

The deeper active layer simulated by NewPC implies more soil water storage capacity, resulting in lower soil moisture in shallow soil layers and higher soil water in deep layers (S_f NewPC; magenta lines; Figure 2) compared to baseline results.

- The changes in soil liquid water content, in turn, impact phase-change and soil temperature simulations. Comparison with the observed soil liquid water content reveals a better agreement with observations (Table S5). For example, at ATQ (Figure S7), the RMSEs of the liquid water content were reduced by 5.4%, 35.3%, 42.6%, and 25.4% for the 3rd through 6th layers, respectively (Table S5).
- 390 The changes to model representations of phase change led to large reductions in soil temperature bias. The relative improvements in RMSE of simulated soil temperatures during Sep. and Oct. (i.e., the two months that the ZCPs usually cover), generally increased with depth for surface layers (within about 20 cm of the surface, i.e., 1st to 4th layer), and were above 80% for the intermediate layers (5th to 8th) at all the sites (Figure 4). At the two Barrow sites where observed soil temperatures were available, the relative improvements for the deepest (13th) layer were 72.6% and 71.1%, on average, for the early winter and
- 395 annual cycle, respectively. Therefore, incorporating the new phase-change scheme also resulted in improved bottom temperature boundary conditions, which is critical for accurately simulating permafrost dynamics (Sapriza-Azuri et al., 2018). Improvements between Septemper and December and the whole annual cycle also increased with soil depth, showing site-averaged reductions in RMSEs ranging from 47% to 63% and from 36% to 46% for the two periods, respectively. The whole cold-season period (Sep. to May) showed, on average, 44% to 53% reduction in RMSEs from the 1st to 6th layer at relatively
- 400 warmer sites (i.e., ATQ and IVO), and from 19% to 69% for the top 13 layers for the two Barrow sites. Also, after the freezing process ends, simulated deeper soil layer temperatures were underestimated (e.g., December through April). This bias might be caused by underestimated snow depth (Figure S9) possibly resulting from inaccurate forcing (particularly snowfall), land cover, microtopography, and/or wind-blown snow redistribution.
- 405 Simulations with the new phase change scheme also show improved agreements between simulated and observed soil temperatures during the spring thawing season compared to the baseline results (red vs. blue in Figure 2). Compared to observations, the newly simulated soil temperatures were still slightly underestimated during the thawing season (i.e., May) at all four sites, showing later onset of thawing indicated by the timing when warming soil temperatures cross 0°C and soil

moisture starts to rise (Figure 2). One possible reason for this bias is the lack of representation of advective heat transport.

410 That is, the model does not represent the heat of spring rain that is advectively transported into soils (Neumann et al., 2019; Mekonnen et al., 2020); nor does it account for advective heat transport associated with water fluxes in subsurface soils after the spring-rainwater mix with existing cold liquid water in soils.

The improved simulations of soil temperature, liquid water content, and ZCP duration greatly impacted soil HR and methane

415 production but did not necessarily guarantee improvements in CO₂ and CH₄ emissions. In the next section, we closely evaluate simulated CH₄ and CO₂ fluxes with different parameterizations of environmental modifiers and the modified CH₄ parameterizations as described in Section 3.1.

4.2 Evaluation of CO2 and CH4 Fluxes

- 420 Here we evaluate the simulated monthly CO₂ and CH₄ fluxes at the site scale against EC tower observations. Figure 5 displays the NSEs of ELMv1-ECA simulations using different carbon decomposition schemes and CH₄ process-related parameters, i.e., "NewPC_NewDecomNewCH4" (grey dots) (see configurations in Table 2). (Time series of all the simulations are provided in Figure S7). The failure of simulated CH₄ emissions to capture the methane seasonality at IVO (as indicated by Figures S8 and S10) might occur because of the lack of 1) a reasonable wetland module that can adequately account for inundated hydro-ecological dynamics, 2) advective heat transport at the air-ground interface through rainfall infiltration and within subsurface soils through water transfer, 3) a representation of microbial dynamics, and 4) the geological micro-seepage emission of CH₄, as reported in previous studies (Anthony et al., 2012; Etiope and Klusman, 2010; Russell et al., 2020). For
- instance, Lyman et al. (2020) showed large temporal variability of CH₄ at natural gas well pad soils, similar to the observations at IVO (Anthony et al., 2012). The advective heat transport not only impacts soil temperature, but also affects soil moisture
 redistribution, substrate availability, and microbial activity. Also, methanogen seasonal dynamics would cause hysteretic effects on CH₄ emission response to soil temperatures (Chang et al., 2020; 2021; Chadburn et al., 2020). In the future, we will incorporate a representation of methanogen seasonal dynamics and simulate microbial population and activity levels to address the hysteresis of CH₄ emissions with temperature. We will also explore more on the contribution of geological micro-seepage emission. The four mechanisms discussed above (i.e., wetland dynamics, advective heat transport, microbial dynamics, and
- 435 geological micro-seepage CH₄ emission) currently missing in our model are likely necessary to simulate CH₄ emissions at this site, and we therefore do not include CH₄ analysis at IVO in the following sections.

The improved phase-change scheme, and thus improved simulations of ZCP durations and soil temperature and moisture, resulted in greatly improved performance for CO₂ emissions at BES/CMDL and BEO, and slightly better performance for CH₄

440 emissions at ATQ, compared to the baseline (blue for "NewPC_OriDecomOriCH4" vs. green for baseline; Figure 5), even though the carbon decomposition and methane model remained the same as the baseline. Incorporating the revised CH₄ model

with the default parameter (discussed in section 3.1.3) improved simulated CH₄ emissions at BES/CMDL, BEO, and ATQ (magenta for "NewPC_OriDecomNewCH4" vs. blue for "NewPC_OriDecomOriCH4"), especially during the cold season (Figure S8). The improved NSEs for CH₄ emissions mainly resulted from increased emissions over early winter (Sep. and

- Oct.) and slight but persistent enhancements throughout the rest of the cold season (magenta in Figure S8), which were related to our modifications to CH₄ transport mechanisms. Further, with the identified optimal parameterization of environmental modifiers to the base decomposition rate and methane parameters, results demonstrate substantial improvements to the simulation of CO₂ net flux and CH₄ emissions compared to baseline results (red vs. others; i.e., shortest distance from (*NSE_{CH4}*, *NSE_{CO2}*) to (1, 1)). Among the 121 common schemes providing good performance for both CO₂ and CH₄ emissions (i.e., both *NSE_{CH4}* and *NSE_{CO2}* larger than 0.5, indicated by the grey dots within the boxes in Figure 5), we identified a generic scheme by selecting the common parameterization that provided the best overall performance for all the sites (except IVO) (cyar; Figure 5). The specific environmental modifier functions and methane parameters for the optimal and generic scheme are provided in Table S6.
- 455 Figure S8 illustrates the uncertainty associated with the model representations of environmental influences on heterotrophic respiration and methane parameters. The optimal simulations at the study sites either used the modified ELMv1-ECA moisture scalar or Yanetal (see Table S6), i.e., two groups of moisture-dependency functions implemented for each soil layer. For the Sierra et al. (2015) empirical moisture functions, the influence of liquid moisture content on heterotrophic respiration is uniformly applied to all active soil layers, even though the soil properties (e.g., porosity and saturated soil water potential) are
- 460 quite different vertically. ELMv1-ECA's moisture scalars (including the original scheme) that use soil water potential, in contrast, reasonably explained the varying influence along with the vertical soil profile (i.e., relationships between soil liquid water content and soil temperature varies with soil clay fraction as demonstrated by Fig.1 in Niu and Yang, 2006). The Yanetal moisture functions also used soil layer-dependent porosity and clay content to calculate relevant parameters (Yan et al., 2018). The simulations with moisture functions documented in Sierra et al. (2015) generally overestimated CO₂ and CH₄ emissions,
- 465 especially during the warm season when the thaw depth is deep and soil wetness is high, thus permitting large moisture modifier scalar applied to the base decomposition rates for all the soil layers regardless of soil properties.

Reducing the minimum soil water potential ψ_{min} for moisture scalar effectively prevents the possibility of zero respiration in subfreezing soils during wintertime (Figure 6). This change exerts more impact on cold sites, such as the two Barrow sites, 470 due to the smaller supercooled liquid water under the colder temperature. Thus, the improved NSEs for CO₂ and CH₄ emissions at BES/CMDL and BEO were larger than those at ATQ (Figure 5). Since the temperature at ATQ was not cold enough to make the supercooled liquid water content small enough to give a zero moisture scalar, the microbial respiration was not completely shut down with the original decomposition modifier at this site. Indeed, at ATQ, where cold-season temperatures are relatively warmer than at BES/CMDL and BEO, simulations with the original ELMv1-ECA environmental modifier (i.e.,

475 "NewPC_OriDecomNewCH4" in Figure S8; discussed in Section 3.1.2), already released much more CO₂ and CH₄ throughout

the cold season than in the baseline simulations, owning to the improved simulations of soil temperature and moisture, and the modifications for CH₄ transport. At IVO, although generally showing low NSEs for CH₄, some new simulations have greatly improved NSE_{CO2} that are larger than 0.5 (Figure 5), compared with -0.3 for baseline. Indeed, the best result at IVO (with a NSE_{CO2} =0.78) significantly improved the simulation of summer CO₂ sink compared to baseline result (Figure 6).

480

The optimal simulations used Davcent2 temperature-dependency function at ATO and O₁₀-based temperature functions at BES/CMDL and BEO with high Q₁₀ values (e.g., 2.0 and 2.5, respectively) (Table S6), mutually mediating the response of microbial respiration with moisture functions discussed above. At all three sites, the optimized parameterizations used a higher ε_{aere} (i.e., 0.05; Table S6), representing possible cold-season CH₄ emissions through ice cracks and remnants of aerenchyma 485 tissues. This newly introduced variable is highly uncertain, though; it can be calibrated at any other sites against cold-season measurements. At BES/CMDL, the optimized parameterization used a decreased maximum CH₄ oxidation rate constant which has been reported highly uncertain, especially over high latitudes (Riley et al., 2011). The generic scheme (i.e., carbon decomposition + CH₄ parameters) overlaps with the optimized parameterization at BES/CMDL, which provided the best overall performance at all three sites (Table S6). Despite the small site number and the limited spatial representativeness of 490 each site, the identified generic scheme might be applied to the Alaska North Slope tundra. Nevertheless, the generic scheme might induce uncertainty in simulations and might not be the optimal regional scheme over other ecosystems or given different climate forcing and soil conditions. Still, we conclude that the generic scheme can serve as a reasonable initial scheme for estimating CO₂ and CH₄ emissions over other high-latitude areas (e.g., Figure S11). In the future, we will explore more sites from newly published CO₂ and CH₄ datasets from pan-Arctic ecosystems, e.g., BAWLD-CH4 (Kuhn et al., 2021) and 495 FLUXNET-CH4 (Delwiche et al., 2021; Knox et al., 2019).

The extended ZCPs, the revised environmental modifier to decomposition, and the modified CH_4 transport mechanism and oxidation parameter together resulted in large improvements for both CO_2 and CH_4 emissions, especially over the cold season. Nevertheless, the optimal simulations still overestimated the contribution of the early cold season (Sep. and Oct.) CO_2

- 500 emissions at BEO and ATQ (top panel; Figure 7), and underestimated CH₄ emissions during post-ZCP months (e.g., Oct. to Dec.) (bottom panel; Figure 7). Many reasons are responsible for the early cold-season CO₂ overestimations, including model deficiencies, prescribed land parameters, and possibly inaccurate forcing. As for the underestimations of post-ZCP carbon emissions, one critical reason is the lack of sudden bursts of CO₂ and CH₄ within the model, i.e., the gases are pushed out of freezing soils during the freeze-up period (Mastepanov et al., 2008; Pirk et al., 2017). Currently, the ELMv1-ECA mimics this
- 505 sudden burst mechanism by preventing CO₂ and CH₄ from dissolving in the soil ice fraction (Riley et al., 2011), which could capture some burst emissions (e.g., CH₄ emissions in Oct. and Sep. of 2013 at ATQ; Figure 6); but it still shows an overall underestimation for sudden-burst emissions especially at colder sites (e.g., BES/CMDL and BEO; Figures 6 and 7). We will improve this mechanism in the future by explicitly simulating ice encroaching soil pores and pushing out gases and liquid

water out of the soil matrix. In the next section, we quantify the cold season contribution of CO2 and CH4 emissions and then

510 estimate the historical trends of seasonal CO₂ and CH₄ emissions from 1950 to 2017.

515

4.3 Cold-season Contribution of CH4 and CO2 emissions and Historical Trends

Throughout this section, we only retain and discuss the identified optimal simulation results (i.e., ELM_NewPC_Optimized) for each site. To better verify the cold-season contribution of CH₄ and CO₂ emissions to the annual budget, a multi-year average approach was taken because of discontinuity in the observed time series.

The new simulation results with the optimal parameterization showed greatly enhanced performance in terms of capturing the averaged seasonal cycle (red; Figure 7), especially for the cold-season months (Sep. to May; Figure 7), reducing site-averaged MAEs in cold-season total CH₄ and CO₂ emissions by 72% and 70% (Figure 8, Table 4), respectively. Specifically, compared to baseline results which significantly underestimated the cold-season carbon emissions, the optimized simulation results showed 0.79 gC m⁻² and 44.0 gC m⁻² increases in site-averaged cold-season CH₄ and CO₂ emissions, respectively. The optimized simulations reduced biases in early cold-season (cold-season) CH₄ emissions by 80% (74%), 86% (76%), and 77% (61%) for BES/CMDL, BEO, and ATQ (Table 4), respectively. The observed cold-season CH₄ emissions contributed at least ~40% to the annual total at three of the study sites, of which about half occurred in early cold-season months (Sep. and Oct.) CH₄ emissions to the cold-season total were 51%, 65%, and 55% for the

three sites, in comparison with the observed 47%, 58%, and 43%, showing slight overestimations. Compared to the baseline-simulated percentage of cold-season contributions to the annual total CH₄ emissions (i.e., only 5%, 6%, and 15%), the optimized simulation showed greatly improved agreements with observed contributions, i.e., 35%, 35%, 33% vs. 45%, 42%, 45% for BES/CMDL, BEO, and ATO, respectively.

The optimized simulations showed larger improvements in cold-season CO_2 emissions (Figure 8; Table 4) for cold sites (i.e., BES/CMDL and BEO) than for the warmer site (i.e., ATQ and IVO). Specifically, compared to baseline results, the updated ELMv1-ECA reduced the biases in simulated cold-season CO_2 emissions from -56.1 gC m⁻² (64% of the observation) to -12.1

- 535 gC m⁻² (14% of the observation) for BES/CMDL and from -65.0 gC m⁻² (68% of the observation) to -12.4 gC m⁻² (13% of the observation) for BEO. In contrast, the optimized simulation showed slight overestimations for cold-season CO₂ emissions at ATQ and IVO (Table 4). Nevertheless, the optimized ELMv1-ECA provided greatly improved warm-season CO₂ net flux for all the four sites, reducing biases by 110%, 78%, 37%, and 102% compared to baseline results at BES/CMDL, BEO, ATQ, and IVO, respectively. Indeed, the updated model switched warm-season net CO₂ flux from baseline-simulated net emissions
- 540 (positive CO₂ net flux) to net uptake (negative CO₂ net flux) at BES/CMDL, BEO, and IVO, correctly matching with observed warm-season CO₂ net flux (Figure 8).

The observed multi-year averaged annual CO₂ net flux was 19.9 gC m⁻² (source), 31.8 gC m⁻² (source), -3.8 gC m⁻² (sink), and -16.7 gC m⁻² (sink) at BES/CMDL, BEO, ATQ, and IVO, respectively. However, due to the large discontinuity in CO₂

545 observations (Figure 6), the calculated annual CO₂ budget is uncertain. Still, we can characterize the CO₂ budget with simulated results using the updated ELMv1-ECA. We find that the simulated cold-season CO₂ emissions were larger than the warm-season CO₂ net uptake during the analysing period (2013-2017) at all four sites (Figure 8), showing annual CO₂ net flux as 1.1 gC m⁻² (source), 36.6 gC m⁻² (source), 36.5 gC m⁻² (source), and 18.2 gC m⁻² (source) at BES/CMDL, BEO, ATQ, and IVO, respectively. The simulated CO₂ emissions over the early cold season (Sep. and Oct.) accounted for 50%, 56%, 66%, and 35%

550 of the total emissions throughout the cold season for BES/CMDL, BEO, ATQ, and IVO, respectively.

Through trend analysis between 1950 and 2017, we found that the ZCP durations showed increasing trends at all three sites, with ZCP trends increasing with depth (Table 5). At ATQ, a warmer site than BES/CMDL and BEO, the trends of ZCP durations increase from 0.12 to 0.49 days yr⁻¹ along with the vertical soil profile. At BES/CMDL and BEO, only soil layers at 3 cm and 6 cm show statistically significant increasing trends, ranging from 0.10 to 0.13 days yr⁻¹. The CO₂ emissions during the 6 cm ZCP and during cold-season months (September to May) both showed increasing trends at all three sites (Table 6), ranging from 0.12 to 0.17 gC m⁻² yr⁻¹ for the 6 cm ZCP, and from 0.30 to 0.40 gC m⁻² yr⁻¹ for the entire cold season period. Annual CH₄ emissions showed a nonsignificant increasing trends at other sites. In the future, we will examine the generic model parameterization at more sites over the pan-Arctic; we will also optimize regional simulations against spatial datasets of CO₂ and CH₄ upscaled from in-situ measurements over pan-Arctic permafrost domain (Natali et al., 2019; Virkkala et al., 2021; Zeng et al., 2020; Peltola et al., 2019), and discuss the uncertainty of estimated trends of the spatially averaged CO₂ and CH₄ emissions associated with snow impact and model parameterizations.

5 Summary and Discussion

- 565 In this study, we improved ELMv1-ECA simulated subsurface soil temperature, zero-curtain period durations, and cold-season CH₄ and CO₂ net emissions at Alaskan North Slope tundra sites. We first improved the numerical representation of coupled water and heat transport with freeze-thaw processes via modifying ELMv1-ECA's phase-change scheme. Then, we revised the dependency of soil decomposition rates on soil temperature and moisture. We further refined the cold-season methane processes by mimicking emission pathways through ice cracks and remnants of aerenchyma tissues, reducing the maximum
- 570 oxidation rate constant, and reducing upper boundary (snow) resistance that allows CH₄ to be emitted from frozen soils through snow to the atmosphere. We also used the updated ELMV1-ECA to estimate historical trends of cold-season CH₄ and CO₂ net emissions at the Alaskan tundra sites from 1950 to 2017. This study is among the first efforts toward improving simulations of zero-curtain periods and cold-season carbon emissions over the Arctic tundra by ESMs. The strategy of improving ELMv1-

ECA phase-change scheme, environmental controls on microbial activity, and methane parameterizations can be easily applied to other global land models.

With the revised phase-change scheme, the updated ELMv1-ECA greatly improved site-scale simulations of soil temperature, soil moisture, and zero-curtain period. Specifically, the RMSE of daily subsurface soil temperature was substantially reduced compared to the baseline simulation, showing site-averaged improvements ranging from 58% to 87% over the early cold

580 season (Sep. to Oct.) and from 36% to 46% over the annual cycle for soil layers within the active layer. The evaluation of simulated liquid water content with the new phase-change scheme, although limited by the availability of observations, showed a relative reduction in RMSE as high as 43% for the 5th layer at ATQ, and site-averaged improvements of 15% and 21% for the 4th and 5th layer, respectively. Simulated ZCP durations were also greatly improved, with, e.g., relative reductions in MAEs of 65%, 66%, and 50% for the 4th layer (about 12 cm) at BES/CMDL, BEO, ATQ, and IVO, respectively.

585

575

Based upon the improved simulations of soil temperature and moisture with the new phase-change scheme, the optimized parameterization for SOC decomposition scheme and the revised methane module, the site-averaged mean annual errors of cold-season emissions were reduced by 72% and 70% for CH₄ and CO₂, respectively. We also found that CH₄ and CO₂ emissions over the early cold season, i.e., September and October, which usually account for most of the zero-curtain period, contributed more than 50% of the total emissions throughout the cold season (September to May). Zero-curtain period durations showed increasing trends from 1950 to 2017, with larger trends in deeper soil layers. Also, both CO₂ emissions during the 6 cm depth zero-curtain period and the entire cold-season period (Sep. to May) showed increasing trends. Note that the optimized parameterizations would be biased if there is a bias in simulated soil carbon, and therefore should not be taken directly to other models without further analysis. Instead, the optimization procedure described in this study provides a roadmap that can be directly adopted to calibrate other models at different sites.

Although showing improvements compared to baseline results, the new simulations generally overestimated the contribution of the early cold season (Sep. and Oct.) CO₂ emissions at BEO and ATQ. Many reasons could contribute to the overestimations, including poor representation of coupled biogeochemical and hydrological processes in the localized permafrost soil
environment, the lack of accurate representation of inundated hydro-ecological dynamics, underestimation of snow accumulation due to micro-topographic effects, and thus the snow insulation to the ground (e.g., Bisht et al., 2018), among others. Strong microtopographic impacts on CO₂ and CH₄ emissions across seven landscape types in Barrow, Alaska, were recently reported (Wang et al. (2019); Grant et al., 2017a; Grant et al., 2017b). Sensitivity analysis demonstrates large impacts of snow depth on simulated winter soil temperature, summer soil moisture, heterotrophic respiration, and CO₂ fluxes (Figure S9); therefore, the simulation of snow should be the subject of future investigations

The underestimated emissions during post-ZCP months (Oct. to Nov.) may be caused by the lack of sudden bursts of CO_2 and CH_4 during the freeze-up period. In addition, the single static multiplicative function used to parameterize the impact of environmental conditions on respiration might not be appropriate because the environmental impact also depends on maximum

- 610 respiration rate, soil texture, soil carbon content and quality, and microbial biomass (Tang and Riley, 2019). Moreover, due to lacking representations of wetland hydro-ecological dynamics, the model uses simulated upland heterotrophic respiration to estimate CH₄ production (Riley et al., 2011), which might cause underestimations of CH₄ emissions, especially under wet conditions. Also, inappropriately prescribed land cover at the site scale or inaccurate climate forcing (particularly air temperature and precipitation; Chang et al. (2019)) could all impact snow accumulation processes (Tao et al., 2017), which
- 615 can significantly impact CO₂ and CH₄ emission simulations. Customizing the complex local ecosystem vegetation community might be feasible at the site scale, however, it is less possible for regional or global land model simulations. This issue calls for the importance of upscaling methods to model (e.g., Pau et al., 2016; Liu et al., 2016) and measure (e.g., Natali et al., 2019; Virkkala et al., 2019) carbon and water cycle dynamics at the regional and global scales.
- 620 Given the persistent warming and the continued more severe warming in the cold season (Box et al., 2019), we envision continuing increases in cold-season CO₂ and CH₄ emissions from the permafrost tundra ecosystem. The increasing rate of cold-season heterotrophic respiration (releasing CO₂) may become larger than the trend of warm-season vegetation CO₂ uptake under future climate. To accurately characterize cold-season emissions and warm-season net uptake, models have to correctly simulate both components, which, however, few models can do. The updated ELMv1-ECA, with the enhanced capacity to
- 625 reproduce cold-season CO₂ and CH₄ emissions proven by this study, can serve as a starting point to better predict permafrost carbon responses to future climate. Finally, the complex water-carbon interactions require modeling systems with fully coupled hydrological-thermal-biogeochemical processes to better predict the carbon budget in permafrost regions under future climate.

Code/Data availability

The observations used in this study are available at http://dx.doi.org/10.3334/ornldaac/1300 and https://doi.org/10.3334/ornldaac/1562. The UAF observations are available at http://permafrost.gi.alaska.edu. This paper resulted in two updated versions of ELMv1-ECA, i.e., one optimized version from the discussion version (ELMv1a) and another version from the finalized paper (ELMv1b), which are all available at GitHub (https://github.com/jingtao-lbl/)

Author contributions

JT assembled observations, developed the methodology, conducted model simulations, analysed results, and wrote and revised

635 the paper. QZ and WJR contributed to experiment design, editing the original and revised manuscript. RBN edited the original and revised manuscript, and provided valuable discussion and guidance.

Competing interests

The authors declare that they have no conflict of interest.

Acknowledgments

640 We are grateful for valuable discussions with Jinyun Tang, Roisin Commane, Xiyan Xu, Kai Yang, and Chenghai Wang. We thank the anonymous reviewers for their helpful comments. This material is based upon work supported by the U.S. Department of Energy, Office of Science, Office of Biological and Environmental Research under Award Number DE-SC0019063.

Appendices: Description of Relevant Modules within ELMv1-ECA 645

Here we describe the heat transfer in subsurface soils, the carbon decomposition, and the methane module within the ELMv1-ECA that are of particular relevance to our study.

Appendix A Subsurface Heat Transfer

ELMv1-ECA approximates the subsurface heat transfer process with a one-dimensional heat diffusion equation:

$$c\frac{\partial T}{\partial t} = \frac{\partial}{\partial z} \left(\lambda \frac{\partial T}{\partial z} \right), \tag{Eq. A1}$$

- where T is the soil temperature (K), c is the volumetric soil heat capacity (J m⁻³ K⁻¹), λ is soil thermal conductivity (W m⁻¹ K⁻¹) 650 ¹), and z is the soil depth (m) of the ELMv1-ECA soil layers. The ELMv1-ECA soil column consists of 15 layers, with soil thickness increasing exponentially with depth. The bottom of soil column is down to 42 m, and the top 10 layers are hydrologically active with layer node depth as 0.0071 m, 0.0279 m, 0.0623 m, 0.1189 m, 0.2122 m, 0.3661 m, 0.6198 m, 1.0380 m, 1.7276 m, 2.8646 m, respectively. The soil heat capacity and thermal conductivity is updated at each time step based
- 655 on the fractions of soil matrix components, i.e., liquid water content, ice content, and soil solids. The impact of organic carbon on soil thermal and hydraulic properties was incorporated as a linear combination of the counterparts properties of mineral soil and organic matter (Lawrence and Slater, 2008). To solve the (Eq. A1), ELMv1-ECA employs the Crank-Nicholson method, resulting in a tridiagonal system equation (Oleson et al., 2013). We assume a zero-flux bottom boundary condition. The top boundary condition is estimated by solving the energy balance equation at the air and ground interface, with additionally an 660 overlying five-layer snow model and a one-layer surface water model in between. When snow and surface water present, ELMv1-ECA incorporates the snow layers and surface water layer into the tridiagonal system to solve the heat transfer along the entire column.

665

670

ELMv1-ECA incorporates freeze-thaw processes of soil water in a decoupled way. Specifically, the model determines the onset of melting or freezing by soil temperature initially solved at time step n + 1 without consideration of the phase change process, denoted as T_i^{n+1} , i.e.,

$$T_i^{n+1} > T_f \text{ and } w_{ice,i}^n > 0 \qquad \text{melting} \\ T_i^{n+1} < T_f \text{ and } w_{liq,i}^n > w_{liq,max,i}^{n+1} \qquad \text{thawing} \qquad , \qquad (\text{Eq. A2})$$

where T_f is the freezing temperature of water (0°C in Kelvin, i.e., 273.15 K), $w_{ice,i}^n$ and $w_{liq,i}^n$ is the mass of ice and liquid water (kg m⁻²) of layer i, and $w_{liq,max,i}^{n+1}$ (kg m⁻²) is the supercooled liquid water that is allowed to coexist with ice given the subfreezing soil temperature T_i^{n+1} . This $w_{lia,max,i}^{n+1}$ varies with soil texture and temperature and is calculated by the freezing point depression equation (Niu and Yang, 2006),

$$w_{liq,max,i}^{n+1} = \Delta z_i \theta_{sat,i} \left[\frac{10^3 L_f (T_f - T_i^{n+1})}{g T_i^{n+1} \psi_{sat,i}} \right]^{-1/B_i} , \qquad (Eq. A3)$$

where Δz_i is the soil thickness of the *i*th layer (in mm), $\theta_{sat,i}$ represents the soil porosity (i.e., the saturated volumetric water content), L_f is the latent heat of fusion (J kg⁻¹), B_i is the Clapp and Hornberger exponent (Clapp and Hornberger, 1978), g is the gravitational acceleration (m s⁻²), and $\psi_{sat,i}$ is the soil texture-dependent saturated matric potential (mm).

675 The rate of phase change is initially assessed from the heat excess (or deficit) needed to change the estimated temperature to the freezing point. Specifically, the model first computes the energy (H_i) needed for adjusting current soil temperature (T_i^{n+1}) to T_f :

$$H_{i} = -c_{i} \frac{\Delta z_{i}}{\Delta t} T_{inc} + (1 - f_{sno} - f_{h2osfc}) \frac{\partial h}{\partial T} T_{inc} \quad i = 1$$

$$H_{i} = -c_{i} \frac{\Delta z_{i}}{\Delta t} T_{inc} \qquad i > 1$$
(Eq. A4)

where $T_{inc} = T_f - T_i^{n+1}$, *h* is ground heat flux, f_{sno} and f_{h2osfc} is the snow and surface water fraction within the grid cell, respectively. The mass change involved then is computed as $H_m = \frac{H_i \Delta t}{L_f}$ (i.e., $-c_i \frac{\Delta z_i}{L_f} T_{inc}$ for soils below the top interface

680 layer). That is, the mass of ice increased/decreased by freezing/melting is $-H_m$, releasing/absorbing energy H_i to bring up/down the current soil temperature to T_f . Accordingly, the ice and liquid mass are adjusted as:

$$w_{ice,i}^{n+1} = \begin{cases} \min(w_{ice,i}^{n} + w_{liq,i}^{n} - w_{liq,max,i}^{n+1*}, w_{ice,i}^{n} - H_{m}) & w_{liq,i}^{n} + w_{ice,i}^{n} \ge w_{liq,max,i}^{n+1*} \\ 0 & w_{liq,i}^{n} + w_{ice,i}^{n} \le w_{liq,max,i}^{n+1*} \end{cases}$$
(Eq. A5)

 $w_{liq,i}^{n+1} = \max\left(w_{liq,i}^{n} + w_{ice,i}^{n} - w_{ice,i}^{n+1}, 0\right)$

The H_i then is adjusted to H_{i^*} , calculated as $H_{i^*} = H_i - \frac{L_f(w_{ice,i}^n - w_{ice,i}^{n+1})}{\Delta t}$. The H_{i^*} then is the ultimately determined latent heat and is used to further readjust soil temperature as in equation (Eq. A6),

$$T_{i}^{n+1*} = T_{f} + \frac{\Delta t}{c_{i}\Delta z_{i}} H_{i^{*}} = T_{f} - \frac{L_{f}(w_{ice,i}^{n} - w_{ice,i}^{n+1})}{c_{i}\Delta z_{i}},$$
(Eq. A6)

in which the temperature adjusted to T_f is further increased by $-\frac{L_f(w_{ice,i}^n - w_{ice,i}^{n+1})}{c_i \Delta z_i}$ due to soil freezing since $w_{ice,i}^{n+1} \ge w_{ice,i}^n$, or 685 decreased due to melting when $w_{ice,i}^{n+1} < w_{ice,i}^n$.

To improve this scheme, we can incorporate soil-water freezing phase change into equation (Eq. A1) and rewrite the heat transfer equation as (Eq. A7) or (Eq. A8),

$$c\frac{\partial T}{\partial t} = \frac{\partial}{\partial z} \left(\lambda \frac{\partial T}{\partial z} \right) - L_f \rho_{liq} \frac{\partial \theta_{liq}}{\partial t}, \tag{Eq. A7}$$

$$\left(c + L_f \rho_{liq} \frac{\partial \theta_{liq}}{\partial T}\right) \frac{\partial T}{\partial t} = \frac{\partial}{\partial z} \left(\lambda \frac{\partial T}{\partial z}\right),\tag{Eq. A8}$$

where L_f is the latent heat of fusion (J kg⁻¹), θ_{liq} is soil liquid water content (m³ m⁻³), and ρ_{liq} is the density of liquid water (kg m⁻³). To solve (Eq. A8), we need to compute the derivative of the soil freezing characteristic curve ($\theta_{liq}(T)$) with respect 690 to temperature $\left(\frac{\partial \theta_{liq}}{\partial \tau}\right)$. As discussed above, we approximate the $\theta_{liq}(T)$ curve by combining the freezing point temperaturedepression equation (Fuchs et al., 1978) and the soil water retention curve (Clapp and Hornberger, 1978). This leads to the supercooled water formulation (Eq. A3) (Niu and Yang, 2006). Computing $\frac{\partial \theta_{liq}}{\partial T}$ requests the soil freezing curves $\theta_{liq}(T)$ to be continuous and differentiable for a range of temperatures during the freezing process (Kurylyk and Watanabe, 2013; Hansson 695 et al., 2004). Here, we follow the existing ELM framework discussed above to solve (Eq. A8). The original numerical representation for readjusting soil temperature (Eq. A6) obtained by the uncoupled two-step implementation (Eq. A2 to A5) significantly overestimates soil water freezing rates. Two reasons are responsible for the overestimation. First, the freezing point ($T_f = 0^{\circ}$ C) is used to determine the occurrence of soil water phase change under all conditions. To further freeze supercooled soil liquid water, however, the soil temperature has to be colder than a virtual soil temperature (as we described below). Second, due to the steep slope of $\frac{\partial \theta_{liq}}{\partial \tau}$ (especially close to $T_f = 0^{\circ}$ C), the estimated ice mass increase (i.e., w_{ice}^{n+1} – 700 w_{ice}^n or $w_{liq}^n - w_{liq,max}^{n+1}$; see (Eq. A5)) most often exceeds the required mass change, i.e., $H_m = -c_i \frac{\Delta z_i}{L_f} (T_f - T_i^{n+1})$, and thus soil liquid water freezes quickly in a large chunk. Soon, the liquid water available to be frozen becomes too small to release sufficient latent heat to compensate for the required energy deficit $(T_f - T_i^{n+1})$.

Thus, we revised the phase-change scheme mainly through incorporating a phase-change efficiency (ε) and replacing the constant freezing point T_f with the temperature of the freezing point depression in (Eq. A2). The phase-change efficiency, introduced by Le Moigne et al. (2012) and adopted by Masson et al. (2013) and Yang et al. (2018a), is calculated as,

$$\varepsilon = \begin{cases} \varepsilon_{liq,i}^{n} = \frac{\theta_{liq,i}^{n}}{\theta_{sat,i}} & \text{for freezing} \\ \varepsilon_{ice,i}^{n} = \frac{\theta_{ice,i}^{n}}{\theta_{sat,i}} & \text{for melting} \end{cases},$$
(Eq. A9)

where $\theta_{liq,i}^n$ and $\theta_{ice,i}^n$ is the soil liquid and ice volumetric water content of layer *i* at previous time step *n*, respectively, and $\theta_{sat,i}$ represents the soil porosity (i.e., the saturated volumetric water content). The temperature of the freezing point depression, as a virtual temperature (*Tv*) reversely derived from the freezing point temperature-depression equation, i.e.,

$$\psi(T) = \frac{10^3 L_f(T_f - T_i)}{gT_i}$$
 (Fuchs et al., 1978; Cary and Mayland, 1972), is calculated as,

$$Tv_i^{n+1} = \frac{10^3 L_f T_f}{10^3 L_f + g\psi_i^n}$$
, (Eq. A10)

where L_f is the latent heat of fusion (J kg⁻¹) and g is the gravitational acceleration (m s⁻²). ψ_i^n is the soil water potential (mm), calculated as the soil water retention curve of Clapp and Hornberger (1978), i.e., $\psi_i^n = \psi_{sat,i} \left(\frac{\theta_{liq,i}^n}{\theta_{sat,i}}\right)^{-B_i}$, where $\theta_{liq,i}^n = w_{liq,i}^n / \Delta z_i$ as in (Eq. A3), B_i is the Clapp and Hornberger exponent, and $\psi_{sat,i}$ is the soil texture-dependent saturated matric potential (mm).

Then, through multiplying the initially estimated mass change (H_m) by the phase change efficiency (ε) , we replace the freezing point with an efficiency-weighted average of the initially solved soil temperature (T_i^{n+1}) and the freezing point,

$$T_{i}^{n+1*} = T_{f} + \frac{\Delta t}{c_{i}\Delta z_{i}} \left(-c_{i} \frac{\Delta z_{i}}{\Delta t} T_{inc} \varepsilon_{i} + \frac{L_{f} \left(w_{ice,i}^{n+1} - w_{ice,i}^{n} \right)}{\Delta t} \right)$$

$$= T_{f} - \left(T_{f} - T_{i}^{n+1} \right) \varepsilon_{i} + \frac{L_{f} \left(w_{ice,i}^{n+1} - w_{ice,i}^{n} \right)}{c_{i}\Delta z_{i}} \qquad .$$
(Eq. A11)
$$= (1 - \varepsilon_{i}) T_{f} + \varepsilon_{i} T_{i}^{n+1} + \frac{L_{f} \left(w_{ice,i}^{n+1} - w_{ice,i}^{n} \right)}{c_{i}\Delta z_{i}}$$

Here, $w_{ice,i}^{n+1}$ is calculated by (Eq. A5) as well, but with updated H_m (i.e., $-\varepsilon_i c_i \frac{\Delta z_i}{L_f} (Tv_i^{n+1} - T_i^{n+1})$). The two changes can effectively improve the soil water freezing process simulations and prevent soil becoming irreversibly too cold quickly as simulated by the baseline phase change scheme.

Appendix B Decomposition Cascade Model

- 725 ELMv1-ECA explicitly simulates carbon cycle dynamics (both plant and soil) and accounts for the limitation of nutrient (i.e., nitrogen and phosphorus) availability for plant growth and the nutrient competition between plants and microbes (Burrows et al., 2020; Zhu et al., 2019; Golaz et al., 2019; Zhu et al., 2020). The ELMv1-ECA uses a Century-like soil carbon decomposition cascade model with vertically resolved soil biogeochemistry (Koven et al., 2013b), and explicitly accounts for the influence of substrate and nutrient availability on soil respiration (both root and microbes) (Zhu et al., 2019).
- 730

715

Within the ELMv1-ECA Century decomposition cascade model, the respiration fractions are parameterized as the fraction of the decomposition carbon flux out of each carbon pool, including litter and soil organic matter. The base decomposition rate is modified by a function representing environmental controls on soil decomposition which accounts for the impacts of

individual factors including temperature (f_T) and moisture (f_W) , an oxygen scalar (f_O) , and a depth scalar (f_D) , in a 735 multiplicative way, i.e., $f_{total} = f_T f_W f_O f_D$.

We use a Q_{10} -based standard exponential function to account for the temperature effect on decomposition,

$$f_T = Q_{10} \Big(\frac{T - T_{ref}}{10} \Big),$$
 (Eq. B12)

where $Q_{10} = 1.5$ by default, which is consistent with ecosystem-level observations (Mahecha et al., 2010), and T_{ref} is the reference temperature (25°C). During cold seasons when soil temperature becomes subfreezing, respiration continues but with more controls from liquid water stress. The original moisture scalar (f_W) within ELMV1-ECA is given in the formulation, calculated as,

$$f_{W} = \begin{cases} 0 & For \, \psi_{i} < \psi_{min} \\ \frac{\log(\psi_{min}/\psi_{i})}{\log(\psi_{min}/\psi_{max})} & For \, \psi_{min} \le \psi_{i} \le \psi_{max} \\ 1 & For \, \psi_{i} > \psi_{max} \end{cases}$$
(Eq. B13)

where $\psi_i = \psi_{max} \left(\frac{\theta_{liq,i}}{\theta_{sat,i}}\right)^{-B_i}$ is the soil water potential, where B_i is the Clapp and Hornberger exponent (Clapp and Hornberger, 1978). In frozen soil, the soil water potential is related to soil temperature through the freezing point depression equation, i.e., $\psi_i = \frac{L_f(T_f - T_i)}{10^3 T}$ (Fuchs et al., 1978; Cary and Mayland, 1972) in the supercooled water formulation (Niu and Yang, 2006). Thus, the liquid water stress on decomposition is translated into dependency on temperature when soil temperature is below the freezing point. By default, ψ_{min} is -10MPa, which predicts zero f_W under frozen conditions since ψ_i under a subfreezing soil temperature easily gets smaller than -10MPa (Figure S1a). We thus reduced the ψ_{min} to -10³ MPa and -10⁶ MPa to alleviate the zero respiration problem in the frozen soils (see Figure S1a). ELMv1-ECA approximates oxygen stress (f_0) as the ratio of available oxygen to that demanded by decomposers, and has a minimum value of 0.2 (Oleson et al., 750 2013).

The depth scalar $(f_D = exp\left(-\frac{z_i}{z_\tau}\right))$ represents unresolved other depth-dependent processes (e.g., soil microbial dynamics, priming effects, etc.) (Koven et al., 2013b; Lawrence et al., 2015; Koven et al., 2015). Applying the depth scalar to decomposition rates would exponentially decrease the respiration fluxes along with the vertical soil layers. The Z_τ is the e-

folding depth for decomposition, and by default Z_{τ} is 0.5 m (Oleson et al., 2013).

Appendix C Methane Model

760

765

The ELMv1-ECA methane model includes the representations of CH₄ production, oxidation, and three pathways of transport (i.e., aerenchyma tissues, ebullition, aqueous and gaseous diffusion), and solves the transient reaction diffusion equation for CH₄. ELMv1-ECA estimates CH₄ production (*P*; mol m⁻³ s⁻¹) in the anaerobic portion of the soil column based on the upland heterotrophic respiration (HR; mol C m⁻² s⁻¹) from soil and litter, further adjusted by factors representing influence from soil temperature (*f_T*), pH (*f_{pH}*), redox potential (*f_{pE}*), and seasonal inundation condition (*S*) (Riley et al., 2011), expressed as,

$$P = HR \times f_{CH_4} \times f_T \times f_{pH} \times f_{pE} \times S.$$
(Eq. C14)

The f_{CH_4} is a fraction of anaerobically mineralized carbon atoms becoming CH₄ and is 0.2 by default. Detailed explanation on these factors can be found in Riley et al. (2011). The methane production *P* is directly related to the estimated HR and impacted by soil temperature, and thus the changes in the carbon decomposition model (Appendix B) and water and heat transfer model (Appendix A) directly influence methane production simulations. Besides, ELMv1-ECA considers the availability of carbon substrate as an important driver of methanogenesis activity and methane production (Riley et al., 2011; Xu et al., 2016).

The ultimately estimated CH₄ emissions are controlled by oxidation, transport mechanisms (i.e., aerenchyma transport, ebullition, and diffusion), and the upper boundary resistance. Detailed descriptions on CH₄ oxidation and transport mechanisms are provided in (Riley et al., 2011). Here we modified CH₄ transport mechanisms for facilitating reasonable cold-season CH₄ emissions.

Vascular plant aerenchyma tissues serve as diffusive pathways to transport CH_4 from soil to the atmosphere. The CH_4 transport 775 via aerenchyma from soil layer z (A(z), mol m⁻² s⁻¹) is calculated as:

$$A(z) = (C(z) - C_a) / \left(\frac{r_L z}{D p T_{aere} \rho_r(z)} + r_a\right) \quad , \tag{Eq. C15}$$

where C(z) and C_a is the gaseous CH₄ concentration (mol m⁻³) in soil depth *z* and in the atmosphere, respectively; r_a is the aerodynamic resistance (s m⁻¹); *D* is the gas diffusion coefficient (m² s⁻¹); *p* is aerenchyma porosity (-); r_L is the ratio of root length to vertical depth (i.e., root obliquity); and $\rho_r(z)$ is the root fraction in soil depth *z* (-). T_{aere} is the specific aerenchyma area (m² m⁻²), and is expressed as,

$$T_{aere} = \frac{f_N N_a LAI}{0.22} \pi R^2,$$
 (Eq. C16)

where *R* represents the aerenchyma radius (= 2.9×10^{-3} m); N_a is the annual net primary production (NPP), and f_N is the belowground fraction of current NPP; and the factor 0.22 represents average observed tiller biomass (gC per tiller) (Wania et al., 2010; Schimel, 1995). Here, tillers mean segmented stems of plants in the Order of Poales, including grasses (Poaceae) and sedges (Cyperaceae) (Wania et al., 2010).

In ELMv1-ECA, methane emissions through aerenchyma were turned off when the soil temperature is below 0°C. We first removed this temperature limitation; then, we integrated the emissions from ice cracks and remnants of aerenchyma tissues with (Eq. C16) by applying a small ε_{aere} during winter time. That is, $T_{aere} = \frac{f_N N_a \varepsilon_{aere}}{0.22} \pi R^2$ when soil temperature is below the freezing point, where ε_{aere} represents possible ice crack fractions and remnants of aerenchyma tissues, and is 0.01 by default. We also tested a larger value for this parameter together with f_{CH_4} and the maximum CH₄ oxidation rate constant 790 $R_{0,max}$ (Table S3).

The $R_{o,max}$ is a key variable controlling CH₄ oxidation rate (R_{oxic}), calculated as,

$$R_{oxic} = R_{o,max} \left[\frac{c_{CH4}}{\kappa_{CH4} + c_{CH4}} \right] \left[\frac{c_{O2}}{\kappa_{O2} + c_{O2}} \right] Q_{10} F_{\nu}, \tag{Eq. C17}$$

where C_{CH4} and C_{02} are CH₄ and O₂ concentrations, respectively; K_{CH4} and K_{02} are the half saturation coefficients (mol m⁻³) for C_{CH4} and C_{02} , respectively. Details about the CH₄ oxidation is provided in Riley et al. (2011). The maximum oxidation rate $R_{0,\text{max}}$ (mol m⁻³ s⁻¹) by default is 1.25E-05 and 1.25E-06 for saturated and unsaturated conditions, respectively. We tested the $R_{0,\text{max}}$ with the smaller value (1.25E-06) for saturated condition as well (Table S3).

Another key variable that is highly uncertain is snow resistance to gas emissions. When snow is present, the upper boundary layer resistance to gas emissions is added by a snow resistance accounting for diffusion through the snow based on the Millington-Quirk expression (Riley et al. (2011). Specifically, the gaseous and aqueous diffusivity in snow is calculated by

800

795

$$D_{snow} = \begin{cases} \varepsilon_{diff} D_{gas} \frac{(\theta_{air}/(\theta_{air} + \theta_{water}))^{10/3}}{(\theta_{air} + \theta_{water})^2} & for gaseous \\ \varepsilon_{diff} D_{liq} (\theta_{water})^2 & for aqueous \end{cases},$$
(Eq. C18)

where D_{gas} and D_{liq} are gaseous and aqueous diffusion coefficients (m² s⁻¹) for CH₄, respectively (see Table 2 in Riley et al. (2011)), $\varepsilon_{\text{diff}}$ is a scale factor (-) for the diffusion coefficients, θ_{water} is water-filled fraction of snow volume (-), and θ_{air} is air fraction in snow (-). The D_{snow} is calculated for each snow layer; then, the total snow resistance is estimated as $R_{snow} =$ $\sum_{i=1}^{n} \frac{dz(i)}{D_{snow}(i)}$ where dz(i) is the thickness of i^{th} snow layer. The top boundary layer conductance for CH₄ then is calculated as $\frac{1}{R_{snow}+R_{pond}+1/C_g}$, where R_{pond} is ponding resistance and C_g is ground conductance (details can be found in Oleson et al., 2013). We found the computed snow resistance generally was too large. We thus decreased snow resistance by introducing new scale factor, i.e., $\varepsilon_{snowdiff}$ (10² by default) in replace with ε_{diff} in (Eq. C18) to increase the conductance at the upper boundary when snow is present.

References

- Anthony, K. M. W., Anthony, P., Grosse, G., and Chanton, J.: Geologic methane seeps along boundaries of Arctic permafrost thaw and melting glaciers, Nat Geosci, 5, 419-426, 2012.
- Arndt, K. A., Oechel, W. C., Goodrich, J. P., Bailey, B. A., Kalhori, A., Hashemi, J., Sweeney, C., and Zona, D.: Sensitivity
 of Methane Emissions to Later Soil Freezing in Arctic Tundra Ecosystems, J Geophys Res-Biogeo, 124, 2595-2609, 10.1029/2019JG005242, 2019.
 - Belshe, E. F., Schuur, E. A. G., and Bolker, B. M.: Tundra ecosystems observed to be CO2 sources due to differential amplification of the carbon cycle, Ecol Lett, 16, 1307-1315, 10.1111/ele.12164, 2013.
 - Bhanja, S. N., and Wang, J. Y.: Estimating influences of environmental drivers on soil heterotrophic respiration in the Athabasca River Basin, Canada, Environ Pollut, 257, 2020.
 - Bisht, G., Riley, W. J., Wainwright, H. M., Dafflon, B., Yuan, F. M., and Romanovsky, V. E.: Impacts of microtopographic snow redistribution and lateral subsurface processes on hydrologic and thermal states in an Arctic polygonal ground ecosystem: a case study using ELM-3D v1.0, Geosci Model Dev, 11, 61-76, 2018.
- Box, J. E., Colgan, W. T., Christensen, T. R., Schmidt, N. M., Lund, M., Parmentier, F. J. W., Brown, R., Bhatt, U. S.,
 Euskirchen, E. S., Romanovsky, V. E., Walsh, J. E., Overland, J. E., Wang, M. Y., Corell, R. W., Meier, W. N., Wouters,
 B., Mernild, S., Mard, J., Pawlak, J., and Olsen, M. S.: Key indicators of Arctic climate change: 1971-2017,
 Environmental Research Letters, 14, 2019.
 - Burrows, S. M., Maltrud, M., Yang, X., Zhu, Q., Jeffery, N., Shi, X., Ricciuto, D., Wang, S., Bisht, G., Tang, J., Wolfe, J., Harrop, B. E., Singh, B., Brent, L., Baldwin, S., Zhou, T., Cameron-Smith, P., Keen, N., Collier, N., Xu, M., Hunke, E.
- C., Elliott, S. M., Turner, A. K., Li, H., Wang, H., Golaz, J. C., Bond-Lamberty, B., Hoffman, F. M., Riley, W. J.,
 Thornton, P. E., Calvin, K., and Leung, L. R.: The DOE E3SM v1.1 Biogeochemistry Configuration: Description and
 Simulated Ecosystem-Climate Responses to Historical Changes in Forcing, J Adv Model Earth Sy, 12, 2020.
 - Cary, J. W., and Mayland, H. F.: Salt and Water Movement in Unsaturated Frozen Soil, Soil Sci Soc Am Pro, 36, 549-&, 1972.
 - Chadburn, S. E., Aalto, T., Aurela, M., Baldocchi, D., Biasi, C., Boike, J., Burke, E. J., Comyn-Platt, E., Dolman, A. J., Duran-
- Rojas, C., Fan, Y. C., Friborg, T., Gao, Y., Gedney, N., Gockede, M., Hayman, G. D., Holl, D., Hugelius, G., Kutzbach,
 L., Lee, H., Lohila, A., Parmentier, F. J. W., Sachs, T., Shurpali, N. J., and Westermann, S.: Modeled Microbial
 Dynamics Explain the Apparent Temperature Sensitivity of Wetland Methane Emissions, Global Biogeochemical
 Cycles, 34, 2020.
- Chang, K. Y., Riley, W. J., Crill, P. M., Grant, R. F., Rich, V. I., and Saleska, S. R.: Large carbon cycle sensitivities to climate across a permafrost thaw gradient in subarctic Sweden, Cryosphere, 13, 647-663, 2019.
 - Chang, K. Y., Riley, W. J., Crill, P. M., Grant, R. F., and Saleska, S. R.: Hysteretic temperature sensitivity of wetland CH4 fluxes explained by substrate availability and microbial activity, Biogeosciences, 17, 5849-5860, 2020.

Chang, K. Y., Riley, W. J., Knox, S. H., Jackson, R. B., and al., e.: Substantial hysteresis in emergent temperature sensitivity of global wetland CH4 emissions, (Under Review), 2021.

- 845 Clapp, R. B., and Hornberger, G. M.: Empirical equations for some soil hydraulic properties, Water Resources Research, 14, 601-604, 1978.
 - Commane, R., Lindaas, J., Benmergui, J., Luus, K. A., Chang, R. Y. W., Daube, B. C., Euskirchen, E. S., Henderson, J. M., Karion, A., Miller, J. B., Miller, S. M., Parazoo, N. C., Randerson, J. T., Sweeney, C., Tans, P., Thoning, K., Veraverbeke, S., Miller, C. E., and Wofsy, S. C.: Carbon dioxide sources from Alaska driven by increasing early winter respiration from Arctic tundra, P Natl Acad Sci USA, 114, 5361-5366, 2017.
 - Dankers, R., Burke, E. J., and Price, J.: Simulation of permafrost and seasonal thaw depth in the JULES land surface scheme, Cryosphere, 5, 773-790, 2011.

850

- Davidson, E. A., and Janssens, I. A.: Temperature sensitivity of soil carbon decomposition and feedbacks to climate change, Nature, 440, 165-173, 2006.
- 855 Davidson, S. J., and Zona, D.: Arctic Vegetation Plots in Flux Tower Footprints, North Slope, Alaska, 2014, ORNL DAAC, Oak Ridge, Tennessee, USA. https://doi.org/10.3334/ORNLDAAC/1546, 2018.
 - Delwiche, K. B., Knox, S. H., Malhotra, A., Fluet-Chouinard, E., McNicol, G., Feron, S., Ouyang, Z., Papale, D., Trotta, C., and Canfora, E.: FLUXNET-CH4: A global, multi-ecosystem dataset and analysis of methane seasonality from freshwater wetlands, Earth System Science Data Discussions, 1-111, 2021.
- 860 Etiope, G., and Klusman, R. W.: Microseepage in drylands: Flux and implications in the global atmospheric source/sink budget of methane, Global and Planetary Change, 72, 265-274, 2010.
 - Fahnestock, J. T., Jones, M. H., Brooks, P. D., Walker, D. A., and Welker, J. M.: Winter and early spring CO2 efflux from tundra communities of northern Alaska, Journal of Geophysical Research-Atmospheres, 103, 29023-29027, Doi 10.1029/98jd00805, 1998.
- 865 Fuchs, M., Campbell, G., and Papendick, R.: An analysis of sensible and latent heat flow in a partially frozen unsaturated soil, Soil Sci Soc Am J, 42, 379-385, 1978.
 - Golaz, J. C., Caldwell, P. M., Van Roekel, L. P., Petersen, M. R., Tang, Q., Wolfe, J. D., Abeshu, G., Anantharaj, V., Asay-Davis, X. S., Bader, D. C., Baldwin, S. A., Bisht, G., Bogenschutz, P. A., Branstetter, M., Brunke, M. A., Brus, S. R., Burrows, S. M., Cameron-Smith, P. J., Donahue, A. S., Deakin, M., Easter, R. C., Evans, K. J., Feng, Y., Flanner, M.,
- Foucar, J. G., Fyke, J. G., Griffin, B. M., Hannay, C., Harrop, B. E., Hoffman, M. J., Hunke, E. C., Jacob, R. L., Jacobsen, D. W., Jeffery, N., Jones, P. W., Keen, N. D., Klein, S. A., Larson, V. E., Leung, L. R., Li, H. Y., Lin, W. Y., Lipscomb, W. H., Ma, P. L., Mahajan, S., Maltrud, M. E., Mametjanov, A., McClean, J. L., McCoy, R. B., Neale, R. B., Price, S. F., Qian, Y., Rasch, P. J., Eyre, J. E. J. R., Riley, W. J., Ringler, T. D., Roberts, A. F., Roesler, E. L., Salinger, A. G., Shaheen, Z., Shi, X. Y., Singh, B., Tang, J. Y., Taylor, M. A., Thornton, P. E., Turner, A. K., Veneziani, M., Wan, H., Wang, H. L., Wang, S. L., Williams, D. N., Wolfram, P. J., Worley, P. H., Xie, S. C., Yang, Y., Yoon, J. H., Zelinka, M. D., Zender, C. S., Zeng, X. B., Zhang, C. Z., Zhang, K., Zhang, Y., Zheng, X., Zhou, T., and Zhu, Q.:

The DOE E3SM Coupled Model Version 1: Overview and Evaluation at Standard Resolution, J Adv Model Earth Sy, 11, 2089-2129, 2019.

- Graf, A., Weihermuller, L., Huisman, J. A., Herbst, M., and Vereecken, H.: Comment on "Global Convergence in the Temperature Sensitivity of Respiration at Ecosystem Level", Science, 331, 1265-+, 2011.
- Grant, R. F., Mekonnen, Z. A., Riley, W. J., Arora, B., and Torn, M. S.: Mathematical Modelling of Arctic Polygonal Tundra with Ecosys: 2. Microtopography Determines How CO2 and CH4 Exchange Responds to Changes in Temperature and Precipitation, J Geophys Res-Biogeo, 122, 3174-3187, 2017a.
- Grant, R. F., Mekonnen, Z. A., Riley, W. J., Wainwright, H. M., Graham, D., and Torn, M. S.: Mathematical Modelling of Arctic Polygonal Tundra with Ecosys: 1. Microtopography Determines How Active Layer Depths Respond to Changes
 - in Temperature and Precipitation, J Geophys Res-Biogeo, 122, 3161-3173, 2017b.

880

885

900

- Harmon, M., and Domingo, J.: A users guide to STANDCARB version 2.0: a model to simulate the carbon stores in forest stands, Dep. of For. Sci., Oreg. State Univ., Corvallis, OR, 2001.
- Harris, I.: CRU JRA v1. 1: A forcings dataset of gridded land surface blend of Climatic Research Unit (CRU) and Japanese
- reanalysis (JRA) data; Jan. 1901-Dec. 2017. Published by: University of East Anglia Climatic Research Unit, Centre for Environmental Data Analysis, 2905 doi: 10.5285/13f3635174794bb98cf8ac4b0ee8f4ed, 2019.
 - Jones, M. H., Fahnestock, J. T., and Welker, J. M.: Early and late winter CO2 efflux from arctic tundra in the Kuparuk River watershed, Alaska, USA, Arct Antarct Alp Res, 31, 187-190, Doi 10.2307/1552607, 1999.
- Kelly, R., Parton, W., Hartman, M., Stretch, L., Ojima, D., and Schimel, D.: Intra-annual and interannual variability of
 ecosystem processes in shortgrass steppe, Journal of Geophysical Research: Atmospheres, 105, 20093-20100, 2000.
 - Kim, D., Lee, M. I., and Seo, E.: Improvement of Soil Respiration Parameterization in a Dynamic Global Vegetation Model and Its Impact on the Simulation of Terrestrial Carbon Fluxes, Journal of Climate, 32, 127-143, 2019.
 - Kim, Y., Ueyama, M., Nakagawa, F., Tsunogai, U., Harazono, Y., and Tanaka, N.: Assessment of winter fluxes of CO2 and CH4 in boreal forest soils of central Alaska estimated by the profile method and the chamber method: a diagnosis of methane emission and implications for the regional carbon budget, Tellus B: Chemical and Physical Meteorology, 59, 223-233, 2007.
 - Kittler, F., Heimann, M., Kolle, O., Zimov, N., Zimov, S., and Gockede, M.: Long-Term Drainage Reduces CO2 Uptake and CH4 Emissions in a Siberian Permafrost Ecosystem, Global Biogeochemical Cycles, 31, 1704-1717, 2017.

Knox, S. H., Jackson, R. B., Poulter, B., McNicol, G., Fluet-Chouinard, E., Zhang, Z., Hugelius, G., Bousquet, P., Canadell,

- J. G., Saunois, M., Papale, D., Chu, H., Keenan, T. F., Baldocchi, D., Torn, M. S., Mammarella, I., Trotta, C., Aurela, M., Bohrer, G., Campbell, D. I., Cescatti, A., Chamberlain, S., Chen, J., Chen, W., Dengel, S., Desai, A. R., Euskirchen, E., Friborg, T., Gasbarra, D., Goded, I., Goeckede, M., Heimann, M., Helbig, M., Hirano, T., Hollinger, D. Y., Iwata, H., Kang, M., Klatt, J., Krauss, K. W., Kutzbach, L., Lohila, A., Mitra, B., Morin, T. H., Nilsson, M. B., Niu, S., Noormets, A., Oechel, W. C., Peichl, M., Peltola, O., Reba, M. L., Richardson, A. D., Runkle, B. R. K., Ryu, Y., Sachs,
- 910 T., Schafer, K. V. R., Schmid, H. P., Shurpali, N., Sonnentag, O., Tang, A. C. I., Ueyama, M., Vargas, R., Vesala, T.,

Ward, E. J., Windham-Myers, L., Wohlfahrt, G., and Zona, D.: FLUXNET-CH4 Synthesis Activity: Objectives, Observations, and Future Directions, Bulletin of the American Meteorological Society, 100, 2607-2632, 10.1175/Bams-D-18-0268.1, 2019.

- Koven, C. D., Ringeval, B., Friedlingstein, P., Ciais, P., Cadule, P., Khvorostyanov, D., Krinner, G., and Tarnocai, C.:
 Permafrost carbon-climate feedbacks accelerate global warming, P Natl Acad Sci USA, 108, 14769-14774, 2011.
 - Koven, C. D., Riley, W. J., and Stern, A.: Analysis of Permafrost Thermal Dynamics and Response to Climate Change in the CMIP5 Earth System Models, Journal of Climate, 26, 1877-1900, 10.1175/Jcli-D-12-00228.1, 2013a.
 - Koven, C. D., Riley, W. J., Subin, Z. M., Tang, J. Y., Torn, M. S., Collins, W. D., Bonan, G. B., Lawrence, D. M., and Swenson, S. C.: The effect of vertically resolved soil biogeochemistry and alternate soil C and N models on C dynamics of CLM4, Biogeosciences, 10, 7109-7131, 2013b.

920

- Koven, C. D., Lawrence, D. M., and Riley, W. J.: Permafrost carbon-climate feedback is sensitive to deep soil carbon decomposability but not deep soil nitrogen dynamics, P Natl Acad Sci USA, 112, 3752-3757, 10.1073/pnas.1415123112, 2015.
- Koven, C. D., Hugelius, G., Lawrence, D. M., and Wieder, W. R.: Higher climatological temperature sensitivity of soil carbon
 in cold than warm climates, Nat Clim Change, 7, 817-+, 10.1038/Nclimate3421, 2017.
 - Kuhn, M. A., Varner, R. K., Bastviken, D., Crill, P., MacIntyre, S., Turetsky, M., Walter Anthony, K., McGuire, A. D., and Olefeldt, D.: BAWLD-CH 4: A Comprehensive Dataset of Methane Fluxes from Boreal and Arctic Ecosystems, Earth System Science Data Discussions, 1-56, 2021.
- Kurylyk, B. L., and Hayashi, M.: Improved Stefan Equation Correction Factors to Accommodate Sensible Heat Storage during
 Soil Freezing or Thawing, Permafrost Periglac, 27, 189-203, 2016.
 - Lawrence, D. M., Thornton, P. E., Oleson, K. W., and Bonan, G. B.: The partitioning of evapotranspiration into transpiration, soil evaporation, and canopy evaporation in a GCM: Impacts on land-atmosphere interaction, J. Hydrometeorol., 8, 862-880, 2007.
- Lawrence, D. M., and Slater, A. G.: Incorporating organic soil into a global climate model, Clim Dynam, 30, 145-160, 10.1007/s00382-007-0278-1, 2008.
 - Lawrence, D. M., and Slater, A. G.: The contribution of snow condition trends to future ground climate, Clim Dynam, 34, 969-981, 2010.
 - Lawrence, D. M., Koven, C. D., Swenson, S. C., Riley, W. J., and Slater, A. G.: Permafrost thaw and resulting soil moisture changes regulate projected high-latitude CO2 and CH4 emissions, Environmental Research Letters, 10, 10.1088/1748-9326/10/9/094011, 2015.
 - Le Moigne, P., Boone, A., Belamari, S., Brun, E., Calvet, J., Decharme, B., Faroux, S., Gibelin, A., Giordani, H., Lafont, S., Lebeaupin, C., Le Moigne, P., Mahfouf, J., Martin, E., Masson, V., Mironov, D., Morin, S., Noilhan, J., Tulet, P., Van den Hurk, B., and Vionnet, V.: SURFEX Scientific Documentation, Note de centre (CNRM/GMME), Météo-France, Toulouse, France, 2012.

- 945 Liu, Y. N., Bisht, G., Subin, Z. M., Riley, W. J., and Pau, G. S. H.: A Hybrid Reduced-Order Model of Fine-Resolution Hydrologic Simulations at a Polygonal Tundra Site, Vadose Zone J, 15, 2016.
 - Lyman, S. N., Tran, H. N., Mansfield, M. L., Bowers, R., and Smith, A.: Strong temporal variability in methane fluxes from natural gas well pad soils, Atmospheric Pollution Research, 2020.
 - Mahecha, M. D., Reichstein, M., Carvalhais, N., Lasslop, G., Lange, H., Seneviratne, S. I., Vargas, R., Ammann, C., Arain,
- 950 M. A., Cescatti, A., Janssens, I. A., Migliavacca, M., Montagnani, L., and Richardson, A. D.: Global Convergence in the Temperature Sensitivity of Respiration at Ecosystem Level, Science, 329, 838-840, 2010.
 - Masson, V., Le Moigne, P., Martin, E., Faroux, S., Alias, A., Alkama, R., Belamari, S., Barbu, A., Boone, A., Bouyssel, F., Brousseau, P., Brun, E., Calvet, J. C., Carrer, D., Decharme, B., Delire, C., Donier, S., Essaouini, K., Gibelin, A. L., Giordani, H., Habets, F., Jidane, M., Kerdraon, G., Kourzeneva, E., Lafaysse, M., Lafont, S., Brossier, C. L., Lemonsu,
- A., Mahfouf, J. F., Marguinaud, P., Mokhtari, M., Morin, S., Pigeon, G., Salgado, R., Seity, Y., Taillefer, F., Tanguy,
 G., Tulet, P., Vincendon, B., Vionnet, V., and Voldoire, A.: The SURFEXv7.2 land and ocean surface platform for coupled or offline simulation of earth surface variables and fluxes, Geosci Model Dev, 6, 929-960, 2013.
 - Mekonnen, Z. A., Riley, W. J., Grant, R. F., and Romanovsky, V.: Changes in precipitation and air temperature contribute comparably to permafrost degradation in a warmer climate, Environ Res Lett, 2020.
- 960 Meyer, N., Welp, G., and Amelung, W.: The Temperature Sensitivity (Q10) of Soil Respiration: Controlling Factors and Spatial Prediction at Regional Scale Based on Environmental Soil Classes, Global Biogeochemical Cycles, 32, 306-323, 2018.
 - Moriasi, D. N., Arnold, J. G., Van Liew, M. W., Bingner, R. L., Harmel, R. D., and Veith, T. L.: Model evaluation guidelines for systematic quantification of accuracy in watershed simulations, Transactions of the Asabe, 50, 885-900, 2007.
- 965 Moyano, F. E., Manzoni, S., and Chenu, C.: Responses of soil heterotrophic respiration to moisture availability: An exploration of processes and models, Soil Biol Biochem, 59, 72-85, 2013.
 - Nash, J. E., and Sutcliffe, J. V.: River flow forecasting through conceptual models part I A discussion of principles, J. Hydrol., 10, 282-290, 10.1016/0022-1694(70)90255-6, 1970.
- Natali, S. M., Watts, J. D., Rogers, B. M., Potter, S., Ludwig, S. M., Selbmann, A.-K., Sullivan, P. F., Abbott, B. W., Arndt,
 K. A., Birch, L., Björkman, M. P., Bloom, A. A., Celis, G., Christensen, T. R., Christiansen, C. T., Commane, R.,
 Cooper, E. J., Crill, P., Czimczik, C., Davydov, S., Du, J., Egan, J. E., Elberling, B., Euskirchen, E. S., Friborg, T.,
 Genet, H., Göckede, M., Goodrich, J. P., Grogan, P., Helbig, M., Jafarov, E. E., Jastrow, J. D., Kalhori, A. A. M., Kim,
 Y., Kimball, J. S., Kutzbach, L., Lara, M. J., Larsen, K. S., Lee, B.-Y., Liu, Z., Loranty, M. M., Lund, M., Lupascu, M.,
 Madani, N., Malhotra, A., Matamala, R., McFarland, J., McGuire, A. D., Michelsen, A., Minions, C., Oechel, W. C.,
 Olefeldt, D., Parmentier, F.-J. W., Pirk, N., Poulter, B., Quinton, W., Rezanezhad, F., Risk, D., Sachs, T., Schaefer, K.,
 Schmidt, N. M., Schuur, E. A. G., Semenchuk, P. R., Shaver, G., Sonnentag, O., Starr, G., Treat, C. C., Waldrop, M.
 P., Wang, Y., Welker, J., Wille, C., Xu, X., Zhang, Z., Zhuang, Q., and Zona, D.: Large loss of CO2 in winter observed across the northern permafrost region, Nat Clim Change, 10.1038/s41558-019-0592-8, 2019.

Neumann, R. B., Moorberg, C. J., Lundquist, J. D., Turner, J. C., Waldrop, M. P., McFarland, J. W., Euskirchen, E. S., Edgar,

- 980 C. W., and Turetsky, M. R.: Warming effects of spring rainfall increase methane emissions from thawing permafrost, Geophys Res Lett, 2019.
 - Nicolsky, D. J., Romanovsky, V. E., Alexeev, V. A., and Lawrence, D. M.: Improved modeling of permafrost dynamics in a GCM land-surface scheme, Geophys Res Lett, 34, 2007.
 - Niu, G. Y., and Yang, Z. L.: Effects of frozen soil on snowmelt runoff and soil water storage at a continental scale, J.
- 985
 - Hydrometeorol., 7, 937-952, 2006.
 - Oechel, W. C., Vourlitis, G. L., Hastings, S. J., Zulueta, R. C., Hinzman, L., and Kane, D.: Acclimation of ecosystem CO2 exchange in the Alaskan Arctic in response to decadal climate warming, Nature, 406, 978-981, 2000.
 - Oechel, W. C., Laskowski, C. A., Burba, G., Gioli, B., and Kalhori, A. A. M.: Annual patterns and budget of CO2 flux in an Arctic tussock tundra ecosystem, J Geophys Res-Biogeo, 119, 323-339, 2014.
- 990 Oechel, W. C., and Kalhori, A.: ABoVE: CO2 and CH4 Fluxes and Meteorology at Flux Tower Sites, Alaska, 2015-2017, https://doi.org/10.3334/ornldaac/1562, 2018.
 - Oleson, K. W., Lawrence, D., Bonan, G., Drewniak, B., Huang, M., Koven, C., Levis, S., Li, F., Riley, W., and Subin, Z.: Technical Description of version 4.5 of the Community Land Model (CLM)(NCAR Technical Note No. NCAR/TN-503+ STR). Citeseer, National Center for Atmospheric Research, PO Box, 3000, 2013.
- 995 Outcalt, S. I., Nelson, F. E., and Hinkel, K. M.: The Zero-Curtain Effect Heat and Mass-Transfer across an Isothermal Region in Freezing Soil, Water Resources Research, 26, 1509-1516, 1990.
 - Pau, G. S. H., Shen, C. P., Riley, W. J., and Liu, Y. N.: Accurate and efficient prediction of fine-resolution hydrologic and carbon dynamic simulations from coarse-resolution models, Water Resources Research, 52, 791-812, 2016.
 - Peltola, O., Vesala, T., Gao, Y., Raty, O., Alekseychik, P., Aurela, M., Chojnicki, B., Desai, A. R., Dolman, A. J., Euskirchen,
- E. S., Friborg, T., Gockede, M., Helbig, M., Humphreys, E., Jackson, R. B., Jocher, G., Joos, F., Klatt, J., Knox, S. H., Kowalska, N., Kutzbach, L., Lienert, S., Lohila, A., Mammarella, I., Nadeau, D. F., Nilsson, M. B., Oechel, W. C., Peichl, M., Pypker, T., Quinton, W., Rinne, J., Sachs, T., Samson, M., Schmid, H. P., Sonnentag, O., Wille, C., Zona, D., and Aalto, T.: Monthly gridded data product of northern wetland methane emissions based on upscaling eddy covariance observations, Earth Syst Sci Data, 11, 1263-1289, 10.5194/essd-11-1263-2019, 2019.
- 1005 Piao, S. L., Ciais, P., Friedlingstein, P., Peylin, P., Reichstein, M., Luyssaert, S., Margolis, H., Fang, J. Y., Barr, A., Chen, A. P., Grelle, A., Hollinger, D. Y., Laurila, T., Lindroth, A., Richardson, A. D., and Vesala, T.: Net carbon dioxide losses of northern ecosystems in response to autumn warming, Nature, 451, 49-U43, 2008.
- Rafique, R., Xia, J. Y., Hararuk, O., Asrar, G. R., Leng, G. Y., Wang, Y. P., and Luo, Y. Q.: Divergent predictions of carbon storage between two global land models: attribution of the causes through traceability analysis, Earth Syst Dynam, 7, 2016.

- Riley, W. J., Subin, Z. M., Lawrence, D. M., Swenson, S. C., Torn, M. S., Meng, L., Mahowald, N. M., and Hess, P.: Barriers to predicting changes in global terrestrial methane fluxes: analyses using CLM4Me, a methane biogeochemistry model integrated in CESM, Biogeosciences, 8, 1925-1953, 10.5194/bg-8-1925-2011, 2011.
- Romanovsky, V. E., Kholodov, A. L., Cable, W. L., Cohen, L., Panda, S., Marchenko, S., Muskett, R. R., and Nicolsky, D.:
 1015 Network of Permafrost Observatories in North America and Russia, 10.18739/A2SH27, 2009.
 - Russell, S. J., Bohrer, G., Johnson, D. R., Villa, J. A., Heltzel, R., Rey-Sanchez, C., and Matthes, J. H.: Quantifying CH4 concentration spikes above baseline and attributing CH4 sources to hydraulic fracturing activities by continuous monitoring at an off-site tower, Atmos Environ, 117452, 2020.
- Sapriza-Azuri, G., Gamazo, P., Razavi, S., and Wheater, H. S.: On the appropriate definition of soil profile configuration and initial conditions for land surface-hydrology models in cold regions, Hydrol Earth Syst Sc, 22, 3295-3309, 10.5194/hess-22-3295-2018, 2018.
 - Schimel, J. P.: Plant-Transport and Methane Production as Controls on Methane Flux from Arctic Wet Meadow Tundra, Biogeochemistry, 28, 183-200, 1995.
- Sierra, C. A., Trumbore, S. E., Davidson, E. A., Vicca, S., and Janssens, I.: Sensitivity of decomposition rates of soil organic matter with respect to simultaneous changes in temperature and moisture, J Adv Model Earth Sy, 7, 335-356, 2015.
 - Sierra, C. A., Malghani, S., and Loescher, H. W.: Interactions among temperature, moisture, and oxygen concentrations in controlling decomposition rates in a boreal forest soil, Biogeosciences, 14, 703-710, 2017.
 - Skopp, J., Jawson, M., and Doran, J.: Steady-state aerobic microbial activity as a function of soil water content, Soil Sci Soc Am J, 54, 1619-1625, 1990.
- 1030 Slater, A. G., Lawrence, D. M., and Koven, C. D.: Process-level model evaluation: a snow and heat transfer metric, Cryosphere, 11, 989-996, 2017.
 - Tang, J. Y., and Riley, W. J.: Weaker soil carbon-climate feedbacks resulting from microbial and abiotic interactions, Nat Clim Change, 5, 56-60, 10.1038/Nclimate2438, 2015.
- Tang, J. Y., and Riley, W. J.: A Theory of Effective Microbial Substrate Affinity Parameters in Variably Saturated Soils and an Example Application to Aerobic Soil Heterotrophic Respiration, J Geophys Res-Biogeo, 124, 918-940, 2019.
 - Tao, J., Reichle, R. H., Koster, R. D., Forman, B. A., and Xue, Y.: Evaluation and Enhancement of Permafrost Modeling With the NASA Catchment Land Surface Model, J Adv Model Earth Sy, 9, 2771-2795, 2017.
 - Tao, J., Koster, R. D., Reichle, R. H., Forman, B. A., Xue, Y., Chen, R. H., and Moghaddam, M.: Permafrost variability over the Northern Hemisphere based on the MERRA-2 reanalysis, The Cryosphere, 13, 2087-2110, 2019.
- 1040 Taylor, M. A., Celis, G., Ledman, J. D., Bracho, R., and Schuur, E. A. G.: Methane Efflux Measured by Eddy Covariance in Alaskan Upland Tundra Undergoing Permafrost Degradation, J Geophys Res-Biogeo, 123, 2695-2710, 2018.
 - Virkkala, A.-M., Aalto, J. A., Tagesson, T., Treat, C. C., Lehtonen, A., Rogers, B. M., Natali, S., and Luoto, M.: High-latitude terrestrial regions remain a CO 2 sink over 1990-2015, AGUFM, 2019, B43E-01, 2019.

Virkkala, A. M., Aalto, J., Rogers, B. M., Tagesson, T., Treat, C. C., Natali, S. M., Watts, J. D., Potter, S., Lehtonen, A., and

- 1045 Mauritz, M.: Statistical upscaling of ecosystem CO2 fluxes across the terrestrial tundra and boreal domain: regional patterns and uncertainties, Global Change Biol, 2021.
 - Wang, Y. H., Yuan, F. M., Yuan, F. H., Gu, B. H., Hahn, M. S., Torn, M. S., Ricciuto, D. M., Kumar, J., He, L. Y., Zona, D., Lipson, D. A., Wagner, R., Oechel, W. C., Wullschleger, S. D., Thornton, P. E., and Xu, X. F.: Mechanistic Modeling of Microtopographic Impacts on CO2 and CH4 Fluxes in an Alaskan Tundra Ecosystem Using the CLM-Microbe Model, J Adv Model Earth Sv, 11, 4288-4304, 2019.
 - Wania, R., Ross, I., and Prentice, I. C.: Implementation and evaluation of a new methane model within a dynamic global vegetation model: LPJ-WHyMe v1.3.1, Geosci Model Dev, 3, 565-584, 2010.

1050

- Wilkman, E., Zona, D., Tang, Y. F., Gioli, B., Lipson, D. A., and Oechel, W.: Temperature Response of Respiration Across the Heterogeneous Landscape of the Alaskan Arctic Tundra, J Geophys Res-Biogeo, 123, 2287-2302, 2018.
- 1055 Xu, X. Y., Riley, W. J., Koven, C. D., Billesbach, D. P., Chang, R. Y. W., Commane, R., Euskirchen, E. S., Hartery, S., Harazono, Y., Iwata, H., McDonald, K. C., Miller, C. E., Oechel, W. C., Poulter, B., Raz-Yaseef, N., Sweeney, C., Torn, M., Wofsy, S. C., Zhang, Z., and Zona, D.: A multi-scale comparison of modeled and observed seasonal methane emissions in northern wetlands, Biogeosciences, 13, 5043-5056, 10.5194/bg-13-5043-2016, 2016.
- Yan, Z. F., Bond-Lamberty, B., Todd-Brown, K. E., Bailey, V. L., Li, S. L., Liu, C. Q., and Liu, C. X.: A moisture function of soil heterotrophic respiration that incorporates microscale processes, Nat Commun, 9, 2018.
 - Yang, K., Wang, C. H., and Li, S. Y.: Improved Simulation of Frozen-Thawing Process in Land Surface Model (CLM4.5), Journal of Geophysical Research-Atmospheres, 123, 13238-13258, 2018a.
 - Yang, Q., Dan, L., Wu, J., Jiang, R., Dan, J., Li, W., Yang, F., Yang, X., and Xia, L.: The Improved Freeze-Thaw Process of a Climate-Vegetation Model: Calibration and Validation Tests in the Source Region of the Yellow River, Journal of Geophysical Research-Atmospheres, 123, 13346-13367, 2018b.
 - Zeng, J. Y., Matsunaga, T., Tan, Z. H., Saigusa, N., Shirai, T., Tang, Y. H., Peng, S. S., and Fukuda, Y.: Global terrestrial carbon fluxes of 1999-2019 estimated by upscaling eddy covariance data with a random forest, Scientific Data, 7, 2020.
 - Zhou, T., Shi, P. J., Hui, D. F., and Luo, Y. Q.: Global pattern of temperature sensitivity of soil heterotrophic respiration (Q(10)) and its implications for carbon-climate feedback, J Geophys Res-Biogeo, 114, 2009.
- 1070 Zhu, Q., Riley, W. J., Tang, J. Y., Collier, N., Hoffman, F. M., Yang, X. J., and Bisht, G.: Representing Nitrogen, Phosphorus, and Carbon Interactions in the E3SM Land Model: Development and Global Benchmarking, J Adv Model Earth Sy, 11, 2238-2258, 2019.
 - Zhu, Q., Riley, W. J., Iversen, C. M., and Kattge, J.: Assessing Impacts of Plant Stoichiometric Traits on Terrestrial Ecosystem Carbon Accumulation Using the E3SM Land Model, J Adv Model Earth Sy, 12, 2020.
- 1075 Zona, D., Gioli, B., Commane, R., Lindaas, J., Wofsy, S. C., Miller, C. E., Dinardo, S. J., Dengel, S., Sweeney, C., and Karion, A.: Cold season emissions dominate the Arctic tundra methane budget, Proceedings of the National Academy of Sciences, 113, 40-45, 2016.

| | Part 1 – P | hase-change | Part 2 – E | nvironmental | Part 3 – Methane module | | | |
|---|----------------------------|---|-------------------------|---------------------------------------|-------------------------|--|--|--|
| | scheme wi | thin the heat | modifier to the base | | | | | |
| | transfer m | odule | decomposition rate | | | | | |
| Relevant | Water and | heat transfer, plant | Plant and s | oil respiration, | CH4 emissions | | | |
| processes | and soil res | spiration, plant | plant productivity, CO2 | | | | | |
| influenced | productivit | y, CO ₂ fluxes and | fluxes and CH4 | | | | | |
| | CH ₄ emissions. | | emissions. | | | | | |
| | Original | New | Original | New | Original | New | | |
| Variables or equations influenced | Eq. A2-6 | Imposing Eq. A7 and Eq. A8 to Eq. A2-A6 | Eq. B9- B10 | Eq. B11 and changes in Table S2 | Eq. C13- C18 | Introducing a new factor ε_{aere}, representing possible pathways via ice crack fractions and remnants of aerenchyma tissues in frozen soils, and thus permitting transport even when temperature is below 0°C. Introducing new scale factors for snow resistance ε_{snowdiff}. Seven parameterizations combining tested values for three highly uncertain variables, i.e., f_{CH4}, ε_{aere}, and R_{0,max} (Table S3). | | |

| Experiment Name | Part 1 - P Scheme v Transfer | Phase Change vithin Heat Model | Part 2 – Envir Modifier withi Decomposition | onmental n Carbon 1 Model | Part 3 – Methane Model | |
|--|------------------------------------|--------------------------------------|---|---------------------------------|---------------------------|--------------|
| | Original | New | Original | New | Original | New |
| OriPC_OriDecomOriCH4 (Baseline) | \checkmark | | \checkmark | | \checkmark | |
| NewPC_OriDecomOriCH4 | | \checkmark | \checkmark | | \checkmark | |
| NewPC_OriDecomNewCH4 ^{&} | | \checkmark | \checkmark | | | \checkmark |
| NewPC_NewDecomNewCH4* (NewPC_Optimized) [#] (NewPC_Generic) ^{\$} | | \checkmark | | \checkmark | | \checkmark |

& "NewCH4" uses the newly modified methane module with default parameterization (see Table S3 in the supplementary file).

* "NewDecom" means replacing the original temperature- and moisture-dependency functions on decomposition rates with (814) new functions of environmental modifiers as listed in Table S2 in the supplementary file. "NewCH4" here means the newly modified methane module with seven parameterizations (Table S3). The seven CH₄ parametrizations were iteratively applied to the model together with 160 common carbon decomposition schemes that provide good performance for CO_2 flux (i.e., NSE_ $CO_2 > 0.5$) at all the sites. This resulted

in 1934 "NewPC_NewDecomNewCH4" simulations in total (i.e., $814+160\times7 = 1934$).

"NewPC_Optimized" is the optimal simulation among the 1934 "NewPC_NewDecomNewCH4" cases at each site.

\$ "NewPC_Generic" means the simulation with the identified generic parameterization that can be applied to regional simulation. The generic scheme is the common satisfactory scheme that provides the best overall performance for all the sites.

1095

Table 3: Mean absolute error (MAE) of simulated ZCP (days) with the original phase-change scheme (Ori_PC) and newly resized phase-change scheme (NewPC), and the relative improvement (%) of using the new phase-change scheme compared to the baseline results, calculated as 100% × (MAE_ZCP_OriPC – MAE_ZCP_NewPC) / MAE_ZCP_OriPC.

| | BES&CMDL | | | BEO | | | ATQ | | | IVO | | |
|------------|---------------------------------|---------------------------------|----------------------------|---------------------------------|---------------------------------|----------------------------|------------------------------|---------------------------------|-------------------------|------------------------------|---------------------------------|-------------------------|
| | MAE_ ZCP_ OriPC (days) | MAE_ ZCP_N ewPC (days) | Impr ove ment (%) | MAE_ ZCP_O riPC (days) | MAE_Z CP_Ne wPC (days) | Impr ovem ent (%) | MAE_Z CP_OriP C (days) | MAE_Z CP_Ne wPC (days) | Impro veme nt (%) | MAE_Z CP_OriP C (days) | MAE_ ZCP_N ewPC (days) | Impro veme nt (%) |
| Layer 1 | 38.80 | 31.40 | 19.0 7 | 37.60 | 33.20 | 11.70 | 26.33 | 13.33 | 49.37 | 54.00 | 51.50 | 4.63 |
| Layer 2 | 29.20 | 14.20 | 51.3 7 | 27.40 | 12.60 | 54.01 | 24.33 | 5.67 | 76.71 | 50.50 | 37.50 | 25.74 |
| Layer 3 | 35.20 | 18.40 | 47.7 3 | 33.60 | 16.80 | 50.00 | 28.00 | 9.33 | 66.67 | 55.75 | 30.25 | 45.74 |
| Layer 4 | 29.60 | 10.40 | 64.8 6 | 30.60 | 10.60 | 65.36 | 28.67 | 9.67 | 66.28 | 61.50 | 30.50 | 50.41 |
| Layer 5 | 18.00 | 11.40 | 36.6 7 | 17.60 | 10.80 | 38.64 | 27.67 | 17.33 | 37.35 | 54.50 | 22.00 | 59.63 |
| Layer 6 | 77.40 | 12.20 | 84.2 4 | 77.40 | 13.00 | 83.20 | 61.67 | 36.67 | 40.54 | 68.00 | 14.75 | 78.31 |
| Layer 7 | NaN | NaN | NaN | NaN | NaN | NaN | NaN | NaN | NaN | 151.33 | 46.67 | 69.16 |

Table 4: Total CH4 emissions and CO2 net flux over three seasonal periods, including the early cold season, cold season, and the
warm season. The "ELM_Optimized" here means ELM_NewPC_Optimized (Table 2). The percentage of each seasonal total CH4
emissions to the annual total is included in the brackets.

| | BES/CMDL | | | BEO | | ATQ | | | |
|---|----------------------------|----------------------------|----------------------------|----------------------------------|----------------------------|----------------------------|----------------------------------|----------------------------|----------------------------|
| | Early Cold | Cold Season | Warm Season | Early Cold Season | Cold Season | Warm Season | Early Cold Season | Cold Season | Warm Season |
| Total CH4 Emissions (gC m ⁻²) | and Oct.) | (Sep. to May) | (Jun. to Aug.) | (Sep. and Oct.) | (Sep. to May) | (Jun. to Aug.) | (Sep. and Oct.) | (Sep. to May) | (Jun. to Aug.) |
| ELM_Baseline | 0.08 (4.7%) | 0.08 (5.1%) | 1.53 (94.9%) | 0.09 (5.5%) | 0.10 (6.2%) | 1.54 (93.8%) | 0.16 (15.1%) | 0.16 (15.3%) | 0.89 (84.7%) |
| ELM_Optimized | 0.51 (17.8%) | 1.00 (34.8%) | 1.88 (65.2%) | 0.73 (22.8%) | 1.11 (35.0%) | 2.07 (65.0%) | 0.18 (18.2%) | 0.58 (33.4%) | 1.16 (66.6%) |
| Observation | 0.63 (21.0%) | 1.32 (44.5%) | 1.65 (55.5%) | 0.83 (24.4%) | 1.43 (41.9%) | 1.97 (58.1%) | 0.36 (19.2%) | 0.85 (44.7%) | 1.04 (55.3%) |
| Bias of ELM_Baseline | 0.55 | 1.24 | 0.13 | 0.74 | 1.32 | 0.43 | 0.20 | 0.69 | 0.16 |
| Bias of ELM_Optimized | 0.11 | 0.32 | -0.23 | 0.10 | 0.31 | -0.09 | 0.05 | 0.27 | -0.11 |
| Bias Reduction | 79.7% | 74.2% | - | 86.1% | 76.5% | - | 77.4% | 61.3% | - |
| | BES/CMDL | | BEO | | | ATQ | | | |
| | Early Cold Season (Sep. | Cold Season (Sep. to | Warm Season (Jun. to | Early Cold Season (Sep. | Cold Season (Sep. to | Warm Season (Jun. to | Early Cold Season (Sep. | Cold Season (Sep. to | Warm Season (Jun. to |
| Total CO ₂ Net Flux (gC m ⁻²) | and Oct.) | May) | Aug.) | and Oct.) | May) | Aug.) | and Oct.) | May) | Aug.) |
| ELM_Baseline | 31.27 | 31.38 | 2.03 | 30.99 | 31.14 | 13.91 | 40.46 | 40.86 | -26.05 |
| ELM_Optimized | 37.97 | 75.36 | -74.27 | 47.14 | 83.71 | -47.15 | 50.51 | 76.18 | -39.64 |
| Observation | 43.60 | 87.50 | -67.61 | 28.20 | 96.14 | -64.33 | 24.29 | 58.64 | -62.41 |
| Bias of ELM_Baseline | 12.33 | 56.12 | -69.64 | -2.79 | 64.99 | -78.24 | -16.16 | 17.78 | -36.36 |
| Bias of ELM_Optimized | 5.63 | 12.14 | 6.66 | -18.94 | 12.42 | -17.19 | -26.21 | -17.54 | -22.76 |
| Bias Reduction | 54.4% | 78.4% | 109.6% | - | 80.9% | 78.0% | - | - | 37.39% |

| <u></u> | | | | | | | |
|----------------------------|--------------------------------|---------|--------------------------------|---------|--------------------------------|---------|--|
| | BES/CMD | L | BEO | | ATQ | | |
| | Trend (days yr ⁻¹) | p Value | Trend (days yr ⁻¹) | p Value | Trend (days yr ⁻¹) | p Value | |
| ZCP Duration of Layer 1 | 0.02 | 0.74 | -0.02 | 0.68 | 0.07 | 0.39 | |
| ZCP Duration of Layer 2 | 0.10 | 0.02 | 0.10 | 0.03 | 0.12 | 0.09 | |
| ZCP Duration of Layer 3 | 0.13 | 0.01 | 0.10 | 0.04 | 0.15 | 0.04 | |
| ZCP Duration of Layer 4 | 0.09 | 0.14 | 0.10 | 0.07 | 0.21 | 0.01 | |
| ZCP Duration of Layer 5 | 0.07 | 0.28 | 0.10 | 0.10 | 0.22 | 0.03 | |
| ZCP Duration of Layer 6 | 0.98 | 0.39 | 0.35 | 0.56 | 0.49 | 0.00 | |

Table 5: Historical trend of ZCP durations (days year⁻¹) for each soil layer from 1950 to 2017. (Trends with p > 0.05 are not statistically significant.)

Table 6: Historical trend (1950 - 2017) in site-scale heterotrophic respiration, CH₄ emission, and CO₂ flux during the ZCP duration at 6 cm (3^{rd} layer), cold-season months (Sep. - May), and the whole annual cycle (Sep. - Aug.). (Trends with p > 0.05 are not statistically significant.)

1115

BES/CMDL BEO ATQ **Trend of Heterotrophic Respiration** Trend Trend Trend р р р $(g C m^{-2} yr^{-1})$ Value $(g C m^{-2} yr^{-1})$ Value $(g C m^{-2} yr^{-1})$ Value 0.09 **ZCP duration at 6** 0.14 0.00 0.04 0.15 0.02 cm **Cold Season** 0.42 0.00 0.31 0.00 0.41 0.00 (Sep.-May) 0.81 0.80 1.06 Annual (Sep.-0.00 0.00 0.00 Aug.) **Trend of CH4 Emission** Trend Trend Trend р р р $(mg C m^{-2} yr^{-1})$ Value $(mg C m^{-2} yr^{-1})$ Value $(mg C m^{-2} yr^{-1})$ Value -0.20 -0.71 -1.69 0.04 **ZCP duration at 6** 0.37 0.13 cm **Cold Season** -0.63 0.16 -2.01 0.00 -1.68 0.22 (Sep.-May) -1.37 0.10 -4.71 0.01 0.52 0.82 Annual (Sep.-Aug.) Trend of CO₂ Net Flux Trend Trend Trend р р р (g C m⁻² yr⁻¹) Value $(g C m^{-2} yr^{-1})$ Value $(g C m^{-2} yr^{-1})$ Value **ZCP duration at 6** 0.15 0.00 0.12 0.17 0.00 0.00 cm **Cold Season** 0.40 0.00 0.30 0.00 0.36 0.00 (Sep.-May) 0.21 0.15 0.43 0.30 0.29 0.19 Annual (Sep.-Aug.)



Figure 1: Red dots indicate the five ABoVE flux tower sites used in this study. Cyan circles are GIPL-UAF permafrost sites.



Figure 2: Comparison of multi-year (2013 - 2017) averaged daily soil temperatures observed (Ts_Obs, black) and simulated with the original (Ts_OriPC, blue) and improved (Ts_NewPC, red) phase-change schemes at BES/CMDL (a) and IVO (b). Simulated moisture saturation with the original (Sf OriPC; green) and improved (Sf NewPC; magenta) schemes are shown on the right hand

axes. Here, the moisture saturation means soil unfrozen (liquid) water content. The horizontal axes indicates days from July to June, with ticks represent the first day of each month. Hatched areas represent durations of zero-curtain periods observed (ZCP_Obs, gray) and simulated (ZCP_OriPC, blue; ZCP_NewPC, red). No baseline ZCP is shown in the 6th layer for BES/CMDL and the 7th layer for IVO because the maximum annual temperature is below 0°C.



Figure 3: Comparison between observed and ELMv1-ECA simulated durations of ZCP for the original (OriPC; open diamonds) and improved (NewPC; solid circles) phase-change schemes over four annual cycles (July to June) from 2013 to 2017. "ly" means

1135

model layer. Simulated ZCP durations with NewPC demonstrate significant improvements compared to OriPC (solid dots vs. open diamonds), especially for the 4th to the deepest layer above permafrost. Note, a zero days ZCP means that the maximum daily temperature during an annual cycle is below 0°C. The pairs of zero vs. non-zero days ZCP (e.g., OriPC ly 7 at IVO and OriPC ly 6 at other sites) indicate that baseline results underestimated ALT. The bias (simulation - observation) of multi-year averaged ALT simulated by the two experiments is provided in each panel.



Figure 4: Relative improvement in the RMSE of simulated soil temperature with the new phase-change scheme (RMSE_Ts_NewPC) compared to that with the original scheme (RMSE_Ts_OriPC), calculated as $100\% \times (RMSE_Ts_OriPC - RMSE_Ts_NewPC) / RMSE_Ts_OriPC$.



Figure 5: Scatter plot between the Nash–Sutcliffe Efficiency (NSE) of simulated monthly CH4 and CO2 emissions. An ideal simulation has both NSEs of CH4 and CO2 as one (i.e., the upper right corner). The boxes encompass simulations with satisfactory performance (NSE > 0.5). Optimized (red) – the best simulation for each site; Generic (cyan) – the simulation with a common parameterization of carbon decomposition scheme and CH4 parameter scheme that provides best overall performance for all the sites. See Table 2 for the configuration for each experiment. The grey dots represent all the tested (1934) simulations indicated in the annotation of Table 2. Symbols outside the plotting ranges indicate poor performance, e.g., (-34.9, -0.3) for baseline at IVO, thus are not shown in the figure.



1155

Figure 6: Observed and simulated monthly CO₂ (left) and CH₄ (right) net flux with the baseline model (ELM_Baseline) and the experiments with updated models (See Table 2 for the configuration for each experiment). Shaded grey areas indicate the minimum-to-maximum bound of simulations within the good performance zone (as shown in Figure 5). Black open circles are observed monthly averages with the number of daily observations less than 10 days, which are not used for the computation in Figure 5. "ELM_NewPC_Optimized" means the best simulation for each site (red diamonds in Figure 5). Intermediate results including

1160 "ELM_NewPC_Optimized" means the best simulation for each site (red diamonds in Figure 5). Intermediate results including "ELM_NewPC_OriDecomNewCH4" are included in supplementary Figure S8.



Figure 7: Comparison of multi-year (2013-2017) averaged monthly mean CO₂ net flux (top) and CH₄ emissions (bottom) from simulations and measurements at the study sites. The error bars represent standard deviation of monthly mean.



Figure 8: Multi-year (2013-2017) averaged total CH4 emissions (bottom) and CO2 net flux (top) during the early cold season (Sep. and Oct.), cold-season period (Sep. to May), warm-season period (Jun. to Aug.), and the annual cycle (Sep. to Aug.) at our study sites. Due to the large discontinuity in CO₂ observations, especially over the warm season (shown in Figure 6), the observed annual CO₂ budget is highly uncertain. Still, the cold-season contributions of both CH₄ and CO₂ emissions are greatly improved by the optimized ELMv1-ECA (i.e., ELM NewPC Optimized).