

# Significant additional Antarctic warming in atmospheric bias-corrected ARPEGE projections with respect to control run

Julien Beaumet<sup>1</sup>, Michel Déqué<sup>2</sup>, Gerhard Krinner<sup>1</sup>, Cécile Agosta<sup>3</sup>, Antoinette Alias<sup>2</sup>, and Vincent Favier<sup>1</sup>

<sup>1</sup>Univ. Grenoble Alpes, CNRS, Institut des Géosciences de l'Environnement, F-38000, Grenoble, France

<sup>2</sup>CNRM, Université de Toulouse, Météo-France, CNRS, Toulouse, France

<sup>3</sup>Laboratoire des Sciences du Climat et de l'Environnement, LSCE-IPSL, CEA-CNRS-UVSQ, Université Paris-Saclay, F-91190 Gif-sur-Yvette, France

*Correspondence to:* Julien Beaumet (Julien.Beaumet@univ-grenoble-alpes.fr)

**Abstract.** In this study, we use run-time bias-correction to correct for ARPEGE atmospheric model systematic errors on large-scale atmospheric circulation. The bias correction terms are built using the climatological mean of the adjustment terms on tendency errors in an ARPEGE simulation relaxed towards ERA-Interim reanalyses. The ~~improvements~~ bias reduction with respect to the AMIP-style uncorrected control run for the general atmospheric circulation in the Southern Hemisphere ~~are~~ is significant for mean state and daily variability. Comparisons for the Antarctic Ice Sheet with the polar-oriented regional atmospheric models MAR and RACMO2 and *in-situ* observations also suggest substantial bias reduction for near-surface temperature and precipitation in coastal areas. Applying the method to climate projections for the late 21<sup>st</sup> century (2071-2100) leads to large differences in the projected changes of the atmospheric circulation in the Southern high latitudes and of the Antarctic surface climate. The projected poleward shift and strengthening of the southern westerly winds are greatly reduced. These changes result in a significant 0.7 to 0.9 K additional warming and a 6 to 9% additional increase in precipitation over the grounded ice sheet. The sensitivity ~~to temperature~~ of precipitation increase to temperature increase (+7.7 and +9%.K<sup>-1</sup>) found is also higher than previous estimates. ~~Highest~~ The highest additional warming rates are found over East Antarctica in summer. In winter, there is a dipole of weaker warming and weaker precipitation increase over West Antarctica, contrasted by a stronger warming and a concomitant stronger precipitation increase from Victoria to Adélie Land, associated with a weaker intensification of the Amundsen Sea Low.

*Copyright statement.* TEXT

## 1 Introduction

The Antarctic Ice Sheet (AIS) contribution to sea-level rise (SLR) has increased ~~dramatically~~ substantially since the 1990s (Velicogna, 2009; Shepherd et al., 2018). The largely positive AIS surface mass balance (SMB), for which positive but ~~insignificant~~ generally not significant trends have been reported during the second part of the 20<sup>th</sup> century (Lenaerts et al., 2016;

Palermé et al., 2017; King and Watson, 2020), ~~now fails to compensate for~~ is now largely overtaken by the increasing ice losses of the West Antarctic Ice Sheet (WAIS) (Velicogna, 2009; Pritchard et al., 2012; Shepherd et al., 2018) (WAIS, Velicogna, 2009; Pritchard et al., 2012; Shepherd et al., 2018). During the course of the 21<sup>st</sup> century, the AIS contribution to SLR is expected to increase (Ritz et al., 2015), possibly dramatically (Pollard et al., 2015). Regarding AIS SMB, ~~there is currently a good agreement on the expectation that it will increase at a~~ rate existing studies agree on expected increasing rates of  $5 \pm 1 \text{ \% K}^{-1}$  (Agosta et al., 2013; Ligtenberg et al., 2013; Frieler et al., 2015; Krinner et al., 2014), ~~mainly~~ as a result of increase in snowfall ~~linked due to~~ higher water vapour saturation pressure in a warmer climate. ~~In this regard~~ Therefore, it is crucial to reduce ~~the uncertainties about the evolution of the future Antarctic climate, and particularly remaining uncertainties on Antarctic regional warming and changes in~~ SMB, in order to assess ~~its the~~ SMB negative contribution to SLR and to better constrain surface forcings for ice dynamics, and ocean and ice-shelves interactions studies. Main source of uncertainties arise from poorly represented sea surface conditions and changes in atmospheric general circulation over Southern high latitudes in most climate models (Turner et al., 2013; Bracegirdle et al., 2013). Due to a lack of *in-situ* measurements and the difficulty to measure SMB from space (Favier et al., 2013; Thomas et al., 2017), polar-oriented regional climate models (RCM) driven by climate reanalyses are deemed the most reliable, and are the most commonly used method to provide estimates of the current Antarctic SMB (e.g., Agosta et al., 2019; van Wessem et al., 2018). For future Antarctic SMB projections, RCMs are usually driven by output from coarser resolution coupled Atmosphere-Ocean General Circulation Model (AOGCMs), such as those involved in the Coupled Model Intercomparison Project, phase 5 (CMIP5, Taylor et al., 2012) or the ongoing phase 6 (CMIP6, Eyring et al., 2016). Since all these models show substantial biases (Gleckler et al., 2008; Flato et al., 2013), there are large uncertainties associated with the dynamical downscaling using RCMs of their future projections. This is particularly relevant for Antarctica, firstly because many state-of-the-art AOGCMs fail to reproduce Southern Hemisphere Sea-Ice Extent (SIE) seasonal cycle and recent trends (Turner et al., 2013; Mahlstein et al., 2013). This is concerning as Sea Surface Conditions (SSC) around Antarctica were shown to have larger instantaneous control on future Antarctic climate than increases in Greenhouse Gases (GHG) concentration (Krinner et al., 2014). Secondly, because in the Southern mid-latitudes, the CMIP5 ensemble mean shows a large ( $\sim 3^\circ$ ) equatorward bias on the position of the surface westerly winds maximum (or “jet”) (Bracegirdle et al., 2013) and there are large uncertainties associated with the atmospheric circulation change signals suggested by these models, as there is an historical state dependence: models with a larger equatorward bias show a larger poleward shift in a warmer climate (Bracegirdle et al., 2013). For Antarctic climate change assessment, it is of prime importance to evaluate the poleward shift of the westerlies and the variations of their intensity, which are among the most salient expected consequences of GHG forcing for the Southern Hemisphere atmospheric circulation (Arblaster and Meehl, 2006; Miller et al., 2006; Fyfe and Saenko, 2006). The associated storm track changes have influenced regional warming (~~Nicolas and Bromwich, 2014~~) and SMB changes (~~Medley and Thomas, 2019~~) in Antarctica and other southern ~~high latitude regions (e.g., Verfaillie et al., 2019)~~ mid and high latitudes (Jones et al., 2019; Medley and Thomas, 2019; Verfaillie et al., 2019), and will continue to do so (~~Bracegirdle et al., 2013~~) in the future (Bracegirdle et al., 2018). Besides, wind-driven oceanic currents in the Amundsen Sea sector influence the rate of ice shelf basal melt (Rignot et al., 2013), which can enhance ice discharge in this sector (Pritchard et al., 2012; Fürst et al., 2016) where large ice losses have been reported recently (e.g., Shepherd et al., 2018).

Another possibility for the downscaling of climate model outputs is the use of variable-resolution or stretched-grid atmosphere-only GCMs (VarGCM : e.g., Fox-Rabinovitz et al., 2006; McGregor, 2015). For projections, the anomaly method, which consists in driving the atmospheric model with observed SSC for historical climate and bias-corrected SSC coming from AOGCM scenarios for future projections, has been extensively used with such models (e.g., Gibelin and Déqué, 2003; Déqué, 2007; Krinner et al., 2008; Beaumet et al., 2019a). Using these methods, the uncertainties on base-line historical climate as well as on climate change signals are reduced. However, even when driven by observed SSC, biases in atmospheric models remain substantial. For instance, the CMIP5 AMIP (Atmospheric Model Intercomparison Project, Gates, 1992) ensemble mean still shows the classical double ~~ITCZ-Intertropical Convergence Zone (ITCZ)~~ problem, even if it is reduced with respect to the CMIP5 coupled model ensemble (Li and Xie, 2012).

Because all regional and global climate ~~model-models~~ bear some biases, and provide information at horizontal resolution too coarse for impact studies, outputs from climate ~~scenarios-model projections~~ are generally bias-corrected (or bias-adjusted) and downscaled *a posteriori* using statistical methods (Hall, 2014; Maraun and Widmann, 2018). However, such methods fail to correct for biases associated with atmospheric general circulation errors (Eden et al., 2012; Stocker et al., 2015; Maraun et al., 2017) or for biases due to poorly represented feedback processes in a warming climate (Maraun et al., 2017). ~~Bias-stationarity is a strong hypothesis needed to justify bias-correction of model errors on atmospheric general circulation. Due to the lack of such justifications, empirical-~~

Empirical run-time bias correction of systematic errors on atmospheric general circulation using the statistics of a nudged simulation towards climate reanalysis ~~such as done in Guldberg et al. (2005) or in Kharin and Seinocea (2012) has been restricted so far to has been applied for instance in~~ seasonal forecasting applications ~~;-Krinner and Flanner (2018) have shown (Guldberg et al., 2005; Kh~~

This type of empirical bias correction has been implemented in this study and is similar to flux corrections methods which have been used in many studies over the last decades (Manabe and Stouffer, 1988; Schneider, 1996; Collins et al., 2006), even though it is not currently deployed in state-of-the-art AOGCMs. Dommenges and Rezny (2018) argued that well-documented flux correction is more valuable than model tuning in future climate projection, as it is “cheaper, simpler, more transparent and does not introduce artificial error interactions between submodels”. Bias stationarity is a strong hypothesis needed to justify the application of such methods for future climate projections. Recently, Krinner and Flanner (2018) have demonstrated that state-of-the-art coupled AOGCMs show striking stationarity of large-scale atmospheric circulation biases under strong warming scenarios. ~~Recently, (Krinner et al., 2020) demonstrated~~ Furthermore, (Krinner et al., 2020) have evidenced that the added-value of run-time bias correction ~~for future projections using is mostly preserved in future projections (at least until the end of current century) using a set~~ perfect model tests. Taking advantage of these findings ~~;-we here apply and justifications, we apply in this~~

study run-time bias correction using the statistics on tendency errors of atmospheric variables in an ARPEGE (Déqué et al., 1994) simulation nudged towards ERA-Interim reanalysis (Dee et al., 2011), following closely the method described in Guldberg et al. (2005). The method is presented in section 2. In section 3.1, we present the ~~improvement-bias reductions~~ obtained for the representation of the Southern Hemispheric atmospheric general circulation as well as for Antarctic surface climate and SMB. In section 3.2, we present and discuss the climate change obtained for the late 21<sup>st</sup> century and compare them with climate changes obtained in the control simulations performed without atmospheric bias-correction. The ~~bias-reduction-obtained~~

~~for historical climate and the difference in Antarctic projected climate change for late 21<sup>st</sup> century perspectives associated with empirical bias correction for future climate projections and the realism of the changes projected for Antarctica~~ are discussed in section 4.

## 2 Data and Methods

### 5 2.1 CNRM-ARPEGE set-up

In this study, the configuration of the CNRM-ARPEGE model is the same as in Beaumet et al. (2019a). The 6.2.4 version of ARPEGE (Déqué et al., 1994), a spectral primitive equation AGCM, is used with a T255 truncation, a stretching pole on the centre of the East Antarctic Plateau (80°S, 90°E) and a 2.5 ~~zoom-factor~~ stretching factor, which means that the horizontal resolution is increased by a factor 2.5 with respect to a regular grid of equal size at the stretching pole. With this setting, the  
10 horizontal resolution over Antarctica varies between 30 km at the stretching pole and 45 km over the northernmost parts ~~on~~ of the Antarctic Peninsula (AP). More details on the model grid are given in the supplementary material (section A1). The horizontal resolution at the antipodes in the Northern Hemisphere is 135 km. The atmosphere is discretized into 91 sigma-pressure vertical levels. The surface processes are solved by SURFEX-ISBA-ES surface scheme (Noilhan and Mahfouf, 1996). Over snow-covered surfaces, a 3-layer intermediate complexity snow scheme (Boone and Etchevers, 2001) is used, which  
15 explicitly accounts for the evolution of the surface albedo, heat transfer through the snow layers and refreezing of liquid water. Over the ocean, a 1D version (that is, without sea-ice advection) of the sea-ice model GELATO (Mélia, 2002) is used in order to correctly model surface energy balance (SEB) over sea-ice. In each simulation, a spin-up phase of two years for the atmosphere is considered and these two years are dismissed for the analysis. ARPEGE set-up and prescribed oceanic surface conditions used in this study are identical to those used in Beaumet et al. (2019a).

### 20 2.2 Empirical bias correction of the AGCM

Following the method presented by Guldberg et al. (2005), we use the climatological mean correction terms of a nudged ARPEGE simulation to build a climatology of the tendency errors of the atmospheric model in a first step. ~~A recent study applying a similar method has been performed for Antarctic climate change (Krinner et al., 2019) using the LMDZ model, the atmospheric component of the IPSL coupled climate model.~~ The second step consists in adding a term derived from the  
25 climatology of the model drift to the prognostic equations of the model, in order to correct in-line (at each time step) selected atmospheric state variables (see below). A recent study applying a similar method has been performed for Antarctic climate change (Krinner et al., 2019) using the LMDZ model, the atmospheric component of the Institut Pierre Simon Laplace (IPSL) coupled climate model.

More precisely, in the first nudged experience, ARPEGE was relaxed (Jeuken et al., 1996) towards ERA-Interim reanalysis  
30 (Dee et al., 2011) over 18 years (1993-2010) in order to use the most reliable period of the reanalysis over the Southern Hemisphere. In this simulation, initial first guess ~~prognostic variables for a given prognostic variables  $\psi$~~  are relaxed towards

the reanalysis reference data following Guldberg et al. (2005) :

$$\psi(t) = \psi^*(t) + \Delta t \frac{\psi^{REF}(t) - \psi^*(t)}{\tau}. \quad (1)$$

The upper index  $\star$  stands for the prognostic solution of the atmospheric model dynamics and physics for the corresponding time step, while  $REF$  indicates the reference variable, from ERA-Interim reanalysis in this case, towards which the model is nudged. The relaxation time ( $\tau$ ) for the nudging is 72 hours for each variable, with an update every 6 hours using a linear interpolation. The nudged variables are the following : air temperature, air specific humidity, logarithm of surface pressure, divergence and vorticity. The term  $[\psi^{REF}(t) - \psi^*(t)]/\tau$  in (1) is the estimate of the tendency residual and is stored in memory at each time step in order to build the correction terms :

$$G = \left[ \frac{\psi^{REF}(t) - \psi^*(t)}{\tau} \right]^{AC} \quad (2)$$

- 10 In (2), the  $AC$  exponent indicates that climatological mean is applied to the tendency residuals in order to produce a seasonally and spatially varying correction term which can be seen as a climatology of the free atmospheric model bias with respect to the reference data set. In a second experiment, this correction term  $G$  is then added at each time step to the atmospheric model prognostic equations for the variables mentioned above :

$$\psi^C(t) = \psi^*(t) + G \quad (3)$$

- 15 which yields the empirically bias-corrected solution  $\psi^C$ . The 72 h value for  $\tau$  in the first nudged experiment was chosen after a few sensitivity tests, as this value yields acceptably small errors in the corrected experiment and weak bias correction terms of the order of a few percent of typical physical tendencies (see Fig.B3). Moreover, Guldberg et al. (2005) found that small values of  $\tau$  (e.g., 6 hours) are not recommended for variables that are poorly constrained in the climate reanalyses, such as the divergence. In the bias-corrected experiment, variables are corrected only above the planetary boundary layer (around 1500 m above sea level), with a progressive transition to uncorrected variables towards the lowest layers (around 100 m).

Some examples of the values obtained for the correction terms are given in the supplementary materials. Order of magnitude of the correction terms with respect to other tendencies associated with model physics are briefly discussed (Appendix B1).

### 2.3 Simulation setup

- In this work, we use a set of 6 simulations, summarized in Table 1. Three simulations use atmospheric bias correction, whereas the three other simulations are the uncorrected reference. Both of these subsets consist of one AMIP-type present-day simulation using observed sea-surface conditions and atmospheric boundary conditions (e.g. greenhouse gas concentrations etc.), and of two simulations for the 2071-2100 period under the RCP8.5 forcing scenario, using oceanic surface anomalies from coupled CMIP5 projections.

- Observed Sea Surface Temperatures (SST) are used for the AMIP-type control run (denoted ARP-AMIP in the following) and for the present-day run with atmospheric bias correction (denoted ARP-AMIP-AC). As already mentioned above, the 1D version of sea-ice model GELATO is used over the sea-ice area. However the sea-ice concentration (SIC) is nudged towards

**Table 1.** Summary of the period, sea surface conditions, greenhouse gases concentration and reference historical simulation for each ARPEGE future projections and historical simulation presented in this study. AC, OC, and AOC acronyms stand for Atmospheric and/or Oceanic Correction. \* ARP-AMIP, ARP-MIR-21-OC and ARP-NOR-21-OC are also described in Beaumet et al. (2019a)

Simulations	Period	Atm.	SSC	<del>GES</del> <u>GHG</u>	Reference for hist. climate
ARP-AMIP	1981-2010	Uncorr.	Observed	historical	-
ARP-AMIP-AC	1981-2010	Bias-corr.	Observed	historical	-
ARP-NOR-21-OC	2071-2100	Uncorr.	Bias-corr. NorESM1-M RCP8.5	RCP8.5	ARP-AMIP
ARP-MIR-21-OC	2071-2100	Uncorr.	Bias-corr. MIROC-ESM RCP8.5	RCP8.5	ARP-AMIP
* ARP-NOR-21-AOC	2071-2100	Bias-corr.	Bias-corr. NorESM1-M RCP8.5	RCP8.5	ARP-AMIP-AC
* ARP-MIR-21-AOC	2071-2100	Bias-corr.	Bias-corr. MIROC-ESM RCP8.5	RCP8.5	ARP-AMIP-AC

observed or bias corrected (for future projections) SIC. In order to obtain a consistent sea-ice thickness (SIT) with concentration, especially between recent climate simulations and projected climate, the SIT is prescribed using a simple parametrization (Krinner et al., 1997, 2010)~~presented in Beaumet et al. (2019b)~~. SST and SIC in future projections are bias-corrected following methods and recommendations from Beaumet et al. (2019b).

5 For ~~the~~ future projections, prescribed SST and SIC changes come from MIROC-ESM and NorESM1-M model under their radiative concentration pathways RCP8.5 (Moss et al., 2010). The reason for this choice and a more complete analysis are presented in Beaumet et al. (2019a). To summarize, these two models were chosen among the CMIP5 ensemble because they display very different changes in winter Sea-Ice Extent (SIE) in their RCP8.5 projection for the late 21<sup>st</sup> century (2071-2010) around Antarctica (respectively -45 % and -14 %). Since we bias-correct SSC from the AOGCM scenarios, our choice of model  
10 is guided by the desire to cover a large range of possible future evolution of SSC around Antarctica rather than by their skills for SSC in present climate. The use of bias-corrected SSC in our future projections is justified by the need to remain consistent with the bias correction terms for the atmospheric model derived in the present climate with an experiment using observed SSC. We combine corrected SSC from the two chosen AOGCMs and bias-corrected or uncorrected atmospheric model to produce four different future climate projections, which are presented and compared in this study. The suffix “OC” in the simulation  
15 names (see Table 1) indicates that only the oceanic boundary conditions are bias-corrected, whereas the suffix “AOC” indicates that in addition to using bias-corrected oceanic boundary conditions, the atmospheric correction is also applied.

## 2.4 Evaluation method for historical climate

In section 3.1, we present the ~~improvement in differences between~~ ARP-AMIP-AC ~~with respect to and~~ ARP-AMIP for the representation of the atmospheric general circulation in the Southern Hemisphere (south of 20°S) and for the Antarctic surface  
20 climate over the 1981-2010 period. For atmospheric general circulation, both simulations are compared to the ERA-Interim reanalyses (ERA-I in the following, Dee et al., 2011), with evaluation of temperature, geopotential heights and specific humidity at different pressure levels for the representation of the mean state. The relative root mean square error (RMSE, denoted  $E$  in the following) reduction obtained for the new ARP-AMIP-AC simulation with respect to the previous uncorrected reference is

calculated using :

$$\Delta_r E = 1 - \frac{E_{\text{ARP-AMIP-AC}}}{E_{\text{ARP-AMIP}}} \quad (4)$$

5 Besides, an assessment of the high frequency variability in ARP-AMIP-AC, ARP-AMIP and ERA-I has been performed using an artificial neural network also called self-organizing map (SOM, Kohonen, 1990, 2013), as already done in other climate studies (e.g., Reusch et al., 2007; Sheridan and Lee, 2011; Krinner et al., 2014). The unsupervised machine-learning algorithm was given as input the daily sea-level pressure (SLP) maps of the first 10 years (1981-1990) of each ARPEGE simulation presented in this study and from the corresponding period in ERA-I. The 20 typical circulation patterns (also called Best Matching  
10 Units, BMU) that were identified after this first step are presented on a 5x4 hexagonal grid in Fig. 2. In a second step, each daily SLP map from each simulation is attributed to the closest BMU using the same distance metric as used to determine the BMU in the first step.

For surface climate, ERA-I reanalyses were shown to have substantial biases in Antarctic near-surface atmospheric temperatures ( $T_{2m}$  hereafter), especially over the East Antarctic Plateau (Fréville et al., 2014; Dutra et al., 2015). On the other hand,  
15 Antarctic  $T_{2m}$  and SMB in polar-oriented RCMs such as MAR and RACMO2 (Van Meijgaard et al., 2008) have been successfully validated against *in situ* observations in many studies (Van Wessem et al., 2014; Agosta et al., 2019; van Wessem et al., 2018) and were found to generally outperform climate reanalyses for Antarctic precipitation and SMB. Therefore, ARPEGE  $T_{2m}$ , precipitation and SMB are evaluated against MAR (Agosta et al., 2019) and RACMO2 (van Wessem et al., 2018) ERA-I  
20 driven simulations and *in situ* data (for temperature only) from the SCAR READER data base (Turner et al., 2004), as in Beaumet et al. (2019a). Significance of the differences is assessed using double-sided t-tests as in Beaumet et al. (2019a). The limitations and potential improvement of the method are also addressed.

### 3 Results and discussion

#### 3.1 Evaluation for Present Climate

In this section, the results of the evaluation for the representation of atmospheric general circulation mean state and **high**  
25 **frequency-daily** variability are first presented and are followed by the results of the evaluation for surface climate. The results are briefly commented and discussed. Where relevant, we also discuss possible biases compensation suppression associated the systematic errors correction for atmospheric general circulation.

##### 3.1.1 Atmospheric General Circulation : mean state

The errors with respect to winter and summer ERA-I ~~mean-sea-level-pressure (SLP)~~ SLP for ARP-AMIP and ARP-AMIP-  
30 AC over the 1981-2010 period can be seen in Fig. 1. As already presented in Beaumet et al. (2019a), the mean SLP of the uncorrected ARP-AMIP control run is low biased in the Southern mid-latitudes, especially in the Pacific sector, while ~~it~~



~~underestimates~~ the depth of the circum-antarctic troughs is underestimated (that is, there is a positive pressure bias), particularly the Amundsen Sea Low (ASL). In ARP-AMIP-AC, most of these errors are removed and only a slight positive SLP bias around the Antarctic coasts remains in winter (JJA, Fig. 1a). The magnitude of the bias is substantially reduced in the Amundsen Sea sector. The slight remaining positive bias in winter around the Antarctic coasts for ARP-AMIP-AC is also found in the 850 and 500 hPa geopotentials height (not shown).

We evaluate the ~~improvement in ARPEGE skills~~ decrease in ARPEGE model errors associated with run-time bias correction through the RMSE reduction ( $\Delta_r E$ ) south of 20°S with respect to ERA-I over the whole period 1981-2010 (Table 2). In order to assess the method independently from the period from which the correction terms were derived (1993-2010), we also assessed the  $\Delta_r E$  for a few key variables in ARP-AMIP-AC for the earliest and independent 1981-1992 period (Table ~~2C1~~). The results of the evaluation using any of the two periods are similar and large RMSE reduction are found. For SLP, the RMSE reduction in ARP-AMIP-AC with respect to ARP-AMIP ranges between 50 and 90% with the lowest ~~improvement in winter~~ reductions in spring. Reduction of the seasonal RMSE between 50 to 70 % with respect to ARP-AMIP are found for many mid and upper tropospheric variables. The largest improvements are found for mid and upper tropospheric temperatures with RMSE decrease around 90% in all seasons. At 200 hPa, a large cold bias, increasing with height, was found in the southern Tropics and mid-latitudes in the ARP-AMIP simulation. In ARP-AMIP-AC, these biases are completely suppressed. ~~However, a~~ In this regard, it is interesting to note that Guldberg et al. (2005) also found large (20-30 %) improvement in the seasonal forecasting skills of their corrected model in the Southern Hemisphere while none were reported on average in the Northern Hemisphere. A larger warm bias at 200 hPa ( $\sim 2K$ ) is present over Antarctica in winter and spring (Fig. ~~??~~). ~~RMSE scores C1b~~.  $\Delta_r E$  (Table 2) indicate little bias reductions for 850 hPa ~~temperatures (T850, ??~~ geopotentials in summer (Z850), 850 and 500 hPa specific humidity (Q850 and Q500, ~~not shown~~). Below 850 hPa, ~~the atmospheric state is progressively uncorrected~~ we remind that tendencies on atmospheric variables are progressively less corrected in ARP-AMIP-AC, while the lowest 100 m near the surface are totally uncorrected. The new bias patterns appearing at this level in ARP-AMIP-AC over southern mid-latitudes land masses and the adjacent oceans (see Appendix C) are most likely linked to surface processes (e.g., development of marine stratocumulus, convective boundary layer). The typical time scale for the development of such processes is shorter than the 72 h relaxation time used in the nudged simulation realized to build the correction terms. As a consequence, this advocates for further sensitivity tests using different relaxation times depending on the variable, especially for specific humidity, in order to improve the skills of the corrected model for this variable in future similar experiment. Re-tuning the atmospheric model during the nudging step could also be an option to ensure consistency of the free physical parameters with the modified circulation characteristics.

The position and the value of the maximum of 850 hPa zonal wind component, referred to as westerlies maximum position (WMPOS) and strength (WMSTR) henceforth are presented in Table 3. In the uncorrected ARP-AMIP simulation, the westerlies maximum is characterized by significant large equatorward bias (3.4°) and underestimation of its strength ( $-1.4 \text{ m.s}^{-1}$ ). The agreement for WMPOS and WMSTR is much better in ARP-AMIP-AC, and remaining errors are insignificant ( $p < 0.05$ ). The annual variabilities of WMPOS and WMSTR decrease in ARP-AMIP-AC with respect to AMIP. This is beneficial, because ARP-AMIP shows a larger variability in WMPOS than what is found in the ERA-I reanalysis.



**Table 2.** Relative seasonal root mean square error reduction  $\Delta_r E$  (in %, [see Eq. 4](#)) south of 20°S with respect to ERA-Interim for ARP-AMIP-AC with respect to ARP-AMIP [during](#) the 1981-2010 period for different surface and tropospheric variables at constant pressure levels :

<i>SimulationsVariable</i>	JJA	SON	DJF	MAM
<b>SLP</b>	<del>87</del> <u>64</u>	<del>69</del> <u>74</u>	<del>52</del> <u>80</u>	<del>75</del> <u>57</u>
<b>T200</b>	95	93	90	86
<b>T500</b>	95	95	96	95
<b>T700</b>	88	90	88	87
<b>T850</b>	95	93	90	86
<b>Z500</b>	93	87	77	90
<b>Z850</b>	90	71	58	87
<b>Q500</b>	13	14	81	23
<b>Q850</b>	42	7	-57	-7

### 3.1.2 Atmospheric General Circulation : daily variability

The relative frequencies for each BMU for AMIP, ARP-AMIP-AC and ERA-I are presented in Fig. 3. ARP-AMIP-AC simulation shows a clearly better distribution of daily SLP with reduced RMSE compared to ARP-AMIP and a much better correlation with the ERA-I distribution. The BMU frequencies in ARP-AMIP are generally overestimated for circulation patterns with a


5 low meridional pressure gradient and low pressures center located relatively far off the Antarctic coasts (1, 2, 3, 6, 11) and/or pressure ridges over the Pacific sector (1, 2, 3, 5). Conversely, frequencies of patterns with large meridional pressure gradient and low pressure systems closer to the Antarctic coasts are mostly underestimated in ~~AMIP~~[ARP-AMIP](#). These errors are consistent with the biases evidenced in the analysis of the errors of the mean state in previous paragraph. For ARP-AMIP-AC, although it is also clearly the most frequent pattern in ERA-I, there is a large overestimation of the 20<sup>th</sup> BMU. The large over-

10 estimation of pattern 20 probably reflects the fact that a certain number of circulation patterns present in ARP-AMIP-AC are not correctly represented in the 20 BMU derived using the daily SLP from all simulations [and reanalysis](#). BMU 20 represents synoptic situations with very high meridional pressure gradient and strongly zonal circulation. As indicated by its position at the fringe of the figure, it represents patterns that are in this sense extreme among the situations appearing in the ARP-AMIP simulation, and it apparently best represents the presumably even more zonal circulation patterns with stronger meridional

15 gradients only present in ARP-AMIP-AC.

[Overall, the analysis of the best matching unit frequencies in the self-organizing maps analysis evidence a much better representation of daily variability in ARP-AMIP-AC which is not an expected results as by construction the empirical bias correction is only expected to improve mean state.](#)

### 3.1.3 Near-surface temperatures



**Figure 1.** ~~AMIP~~Mean SLP difference between ARP-AMIP-AC (a) and ARP-AMIP (b) simulations minus ERA-I for the reference period 1981-2010 in winter (JJA, left) and summer (DJF, right). Value of the RMSE are given below the plots.


#### ARP-AMIP-AC

~~Difference between ARP-AMIP and ARP-AMIP-AC simulations with ERA-I mean SLP for the reference period 1981-2010 in winter (JJA, left) and summer (DJF, right). Value of the RMSE are given below the plots.~~


**Table 3.** Mean annual 850 hPa westerly maximum strength (WMSTR) and position (WMPOS) in ERA-Interim, ARP-AMIP and ARP-AMIP-AC  $\pm$  one standard deviation of the annual mean. Differences significant at  $p=0.05$  with respect to ERA-I are presented in **bold**.

Simulation/Data set	WMSTR ( $\text{m.s}^{-1}$ )	WMPOS ( $^{\circ}$ )
ERA-I	$12.5 \pm 0.6$	$-51.4 \pm 0.8$
ARP-AMIP	<b><math>11.1 \pm 0.5</math></b>	<b><math>-48.0 \pm 1.4</math></b>
ARP-AMIP-AC	$12.2 \pm 0.3$	$-51.0 \pm 0.6$

~~In this section, The effect of the atmospheric correction on ARPEGE mean near-surface air temperatures ( $T_{2m}$ ) from ARPEGE simulations are can be seen in the ARP-AMIP-AC minus ARP-AMIP difference (Fig. 4a). Here,  $T_{2m}$  from the corrected ARPEGE simulation are also compared with those from a MAR RCM simulation forced at its MAR (Agosta et al., 2019) and RACMO2 (van Wessem et al., 2018) RCMs simulations driven at their lateral boundary by ERA-I (Agosta et al., 2019) and those from weather station of the READER data base (Fig. 4ab and c). In this analysis, weather stations for which less than 80 % of valid data were available for the season and the period considered have been discarded from the analysis. The same figure for the ARP-AMIP simulation already presented in Beaumet et al. (2019a) can be seen in the supplementary material (Fig. ??D1). The errors for each station and each season, as well as the mean error and RMSE per regions bias for each Antarctic regions for ARP-AMIP-AC and ARP-AMIP are also presented in the supplementary material for ARP-AMIP-AC~~



**Figure 2.** Mean Sea Level Pressure (SLP) map for the twenty best matching unit (BMU) obtained after a self-organizing map analysis on daily SLP fields. SLP ranges from 1030 hPa (red) to 960 hPa (purple) with 10 hPa intervals.



**Figure 3.** Best matching unit (BMU) relative frequency (%) of ARP-AMIP (red), ARP-AMIP-AC (blue) and ERA-I (black) on daily SLP map over the 1981-2010 period. Root mean square errors and pearson correlation coefficient with respect to ERA-I BMU frequencies are shown to the right of the legend.

(Table ??) and ARP-AMIP (Table ??). The effect of the atmospheric correction on ARPEGE mean  $T_{2m}$  can be seen in the ARP-AMIP-AC minus ARP-AMIP difference (Fig. 4b (Table D1)).

On the East Antarctic Plateau (EAP), the impact of the atmospheric bias-correction is a winter warming (1-3 °C) over large parts of the centre of the Plateau. The warm bias with respect to MAR during this season increases, which is confirmed for instance by a decrease of ARPEGE skills at Vostok in all season but summer (see Table ??). This bias increase seems to result from the removal of a bias compensation that was present in ARP-AMIP. We investigated the value of the near-surface temperature inversion in both historical ARPEGE simulation (not shown) as the difference between surface temperature ( $T_s$ ) and the temperature of the first atmospheric layer, located around 6 to 10 meters height in this ARPEGE setting (level at  $0.9988 \times$  surface pressure). The value of this near-surface temperature inversion around (-15°C) does not decrease in ARP-AMIP-AC. It even increases locally, suggesting small changes in surface boundary layer processes. However, there was a substantial negative bias with respect to MAR in incoming long wave radiation (LWD) in ARP-AMIP, which disappears in ARP-AMIP-AC due to considerable warming at 500 hPa in winter (Fig. C3). The warm bias in winter over the East Antarctic Plateau with respect to MAR or READER weather stations (between 3 and 5 °C), whereas smaller differences are found in the comparison with RACMO2  $T_{2m}$ , is to be put into perspective with even larger biases sometimes found over the East Antarctic Plateau in climate models or in even climate reanalyses (Dutra et al., 2015; Fréville et al., 2014; Bracegirdle and Marshall, 2012).

For coastal East Antarctic READER stations, the cold bias present in every season, particularly in winter, is greatly reduced in ARP-AMIP-AC. The improvement is even dramatic bias reduction is very large for some stations in eastern East Antarctic (e.g.: McMurdo, Dumont D'Urville, Casey, Davis). The effect of the bias correction is also a cooling of some margins of the eastern East Antarctic Plateau in summer, where the warm bias with respect to MAR decreases. However, the errors remain substantial and significant ( $p < 0.05$ ) in many stations and seasons, especially in the winter (mean error  $\approx -2$  °C). No improvement is found for the warm bias on the ice shelves and coastal regions of Dronning Maud Land. In this perspective, the comparison with McMurdo and Scott Base is interesting. These stations, distant by only 3 kilometers and located at the same altitude, belong to the same ARPEGE grid point. While ARP-AMIP-AC bias with respect to Scott Base is small and statistically insignificant in all season but summer, the cold bias is  $\sim 3$  °C (significant) throughout the year with respect to Mc Murdo. This example shows the limited spatial representativity of weather stations in coastal Antarctica, and therefore the comparisons with a 35 km horizontal resolution atmospheric model should be interpreted carefully.

Over West Antarctica and the Peninsula, the effect of the atmospheric bias-correction is a warming over much of coastal and central West Antarctica and on the southern and western parts of the Peninsula in winter. In summer, this warming is restricted to the south-western part of the Peninsula, while there is a cooling of the eastern coasts, particularly marked on the Larsen Ice Shelf. The systematic cold bias with respect to MAR and READER stations of the southern and western part of the Antarctic Peninsula is greatly reduced, with the largest improvements found in the southernmost stations (Rothera and Faraday). Warm and moist advection from the North-West over this part of the Peninsula is indeed underestimated in ARP-AMIP. However, the errors remain significant in summer (mean error  $\approx -1.5$  °C). Moreover and in winter for the comparison with RACMO2.

Besides, ARP-AMIP-AC is cold biased with respect to MAR over the Larsen Ice Shelf in summer, which was not the case of ARP-AMIP.

~~Finally, no substantial improvement is to be reported~~ No reduction is found for the warm bias on the ice shelves and coastal regions of Dronning Maud Land or

5 ~~AMIP-style~~ AMIP-style control run was already fairly high.

The warm temperature bias with respect to MAR-ERA-I over Antarctic ice shelves does not decrease in ARP-AMIP-AC, it even increases over Neumayer and Halley stations, especially in winter. ARPEGE model deficiencies for surface temperatures over the large Ronne-Filchner and Ross ice-shelves have been widely addressed in Beaumet et al. (2019a). They are mostly due to misrepresented stable boundary layers processes and excess (with respect to MAR) in long-wave downward radiation in winter associated with higher cloud cover. Large disagreement have also been evidenced between MAR and RACMO2 for these areas. Indeed, we note that this large winter warm bias is not found over the Ronne-Filchner and is much reduced over the Ross Ice Shelf in the comparison with RACMO2-ERA-I.

### 3.1.4 ~~Surface mass balance and precipitation~~ Precipitation

~~The SMB and its components~~ Total precipitation integrated over the ~~whole grounded~~ grounded Antarctic ice sheet (GIS) for

15 the 1981-2010 reference period are presented in Table 4 for the two historical ARPEGE simulations presented in this study.


~~The table also displays the corresponding values for ERA-I driven simulations with two ARPEGE simulation over the historical period as well as for~~ MAR and RACMO2 (Agosta et al., 2019; van Wessem et al., 2018). ~~The total precipitation driven by ERA-I reanalysis are shown in Table 4. The spatial distributions of precipitation in ARP-AMIP-AC, ARP-AMIP, MAR-ERA-I and RACMO2-ERA-I are compared in Fig. 5. Precipitation over the GIS significantly decreases in ARP-AMIP-AC with respect~~

20 to ARP-AMIP ( $-284 \text{ Gt yr}^{-1}$ ,  $\sim 3$  times the inter-annual variability) and is about 10% lower than the estimates from the two polar RCMs, whereas ~~total~~ precipitation in ARP-AMIP is in good agreement with the latter. ~~The integrated GIS-SMB is about four times MAR interannual variability ( $4\sigma$ ) lower in ARP-AMIP-AC than in MAR-ERA-I and RACMO2-ERA-I as a consequence of lower precipitation and largely overestimated surface sublimation rates (and, to a lesser extent, run-off). In the atmosphere-corrected simulation, there is a significant decrease of run-off and surface snow sublimation with respect~~

25 to previously uncorrected simulation, yet insufficient to match with the estimates from the two polar-oriented RCMs. The overestimation is still large for the comparison with RACMO2 simulation where sublimation include surface and blowing snow sublimation (see Table 4). Estimations of surface snow melt ~~Unsurprisingly, the precipitation minus evaporation difference (P-E), which is a good approximation of moisture transport, has also decreased in ARP-AMIP-AC are however within the interannual variability uncertainty range ( $1\sigma$ ) when compared to RACMO2 and MAR.~~

30 The spatial distributions of total precipitation, surface snow sublimation and SMB of ( $-221 \text{ Gt yr}^{-1}$ ). The main change in atmospheric general circulation in ARP-AMIP-AC ~~are compared with MAR-ERA-I~~ with respect to ARP-AMIP is an increase of SLP at mid-latitude, and a decrease around Antarctica, together with a poleward position of the westerly winds maximum, which can be seen as a larger dominance of SAM<sup>+</sup>-type patterns (see BMU 15,17,18,19,20 in Fig. ??). ~~The same comparison for the 2 and 3). We note that changes in precipitation in ARP-AMIP-AC with respect to ARP-AMIP simulation was already~~





**Figure 4.** (a) ARP-AMIP-AC minus ~~MAR-ERA-I~~ ARP-AMIP  $T_{2m}$  in winter (left) and summer (right). (b) Same as (a) but for ARP-AMIP-AC minus MAR-ERA-I. Circles represent mean bias for weather stations from the monthly READER data base. Black contour lines represents where the difference is one standard deviation of MAR  $T_{2m}$ . (bc) Same as (ab) but for ARP-AMIP-AC minus ~~ARP-AMIP~~ RACMO2-ERA-I.

presented in Beaumet et al. (2019a) and is reproduced in the supplementary material here for convenience (Fig. ??). The effect of the in-line correction of errors on atmospheric general circulation on ARPEGE Antarctic precipitation can be seen in Fig. ??5c) bear many similarities with the signature of a positive SAM pattern on Antarctic precipitation assessed in a RACMO2 ERA-Interim driven simulation in Marshall et al. (2017). It is noteworthy that the link between circulation changes induced by the bias correction and ensuing precipitation changes seen in our simulations is very similar to the effect of bias corrections in the LMDZ model reported by Krinner et al. (2019). It shows mainly a drying in many parts of Antarctica such as Marie-Byrd Land, Dronning Maud Land, Victoria Land and the Transantarctic Mountains, a large part of the East Antarctic Plateau up to Adélie Land, and the eastern side of the AP. Conversely, precipitation increases in ARP-AMIP-AC over central and western West Antarctica and over the western side of the Antarctic Peninsula. These changes in ~~precipitations~~ precipitation result in a better agreement for the spatial distribution of precipitation with MAR-ERA-I (Fig. 5e and f) over large parts of West Antarctica, Dronning Maud Land and the Peninsula. However, the disagreement between the two model is still considerable (> 20%) in many places, and the dry bias with respect to MAR-ERA-I present in ARP-AMIP over the East Antarctic Plateau and the Transarctic Mountains increases. ~~Moreover, the agreement in terms of statistics (mean error and RMSE) does not show the increase in skills expected for ARP-AMIP-AC. Therefore, we also compared precipitation in the ARPEGE simulation~~ In the comparison of ARPEGE precipitation with those from ~~RACMO2 and assessed the impact of the atmospheric correction on the the comparison between the two models~~ RACMO2-ERA-I (Fig. ??a and Fig. ??b)5h and i), the increase in the agreement for ARP-AMIP-AC is much more significant. Many differences between ARP-AMIP with RACMO2 are the same as those seen in the comparison with MAR, and these are also ~~improved~~ reduced in ARP-AMIP-AC. Moreover, ARP-AMIP-AC and RACMO2 agree remarkably well (errors below 20%) in many areas with rather complex topography such as Dronning Maud Land, coastal West Antarctica or the Transantarctic Mountains. The systematic dry bias with respect to MAR over the Transantarctic Mountains and Victoria Land is not found in the comparison with RACMO2. This seems to confirm a wet bias in MAR over this area, such as suggested by the comparison with sparse *in-situ* observations evidenced in Agosta et al. (2019).

The widespread dry bias in ARPEGE over the eastern part of the East Antarctic Plateau and the ridges of the western parts of the Plateau is confirmed in the comparison with RACMO2 and explains most of the ~10% precipitation deficit at the continental scale in the ARP-AMIP-AC simulation with respect to both RCMs. Further investigations are needed in order to identify possible causes of this error, such as deficiencies in moisture transport over the high continental interior or misrepresented clear-sky precipitation processes, which bring a large share of total precipitation in this area (Walden et al., 2003). Improvement for the representation of precipitation, cloudiness and therefore radiative budget for the high continent interior of Antarctica could be obtained for ARPEGE by accounting for saturation with respect to the ice for the formation of cloud condensates and by tuning cloud microphysics to the cold context of Antarctica such as done in Van Wessem et al. (2014).

The better agreement (decrease in RMSE) with precipitation modelled by the polar-oriented RCMs MAR and RACMO2 precipitation increase the confidence in the reliability of spatial distribution of Antarctic precipitation modeled in ARP-AMIP-AC. Both RCMs have been widely validated against *in-situ* measurements of Antarctic SMB (e.g., Agosta et al., 2019; van Wessem et al., 2018). The closer agreement between the ARP-AMIP-AC simulation and RACMO2 in coastal areas offers an interesting opportunity to further investigate causes of remaining disagreement between MAR and RACMO2 in Antarctic precipitation and SMB



Simulation	SMB	Precip	Subli.	Run-Off	Rain	Melt
ARP-AMIP	1970±96	2268±94	277±17	22±14	10±2	52±32
ARP-AMIP-AC	<b>1758±119</b>	<b>1994±117</b>	<b>222±11</b>	<b>14±4</b>	<b>5±2</b>	50±18
MAR-ERA-I <sup>1</sup>	2158±106	2260±104	84±10	3±2	16±3	45±15
RACMO2-ERA-I <sup>1,2</sup>	2117±92	2268±99	136±4	2±2	3±1	–
Vaughan et al. (1999)	1811	–	–	–	–	–


**Table 4.** Mean GIS Surface Mass Balance and its components ( $\text{Gt yr}^{-1}$ )  $\pm$  one standard deviation of the annual value for the reference period 1981-2010. ARPEGE values are integrated over the original model grid and take into account the model land-mask. ARPEGE, <sup>1</sup>MAR and RACMO2 ERA-Interim driven, statistics for 1981-2010 for the Antarctic GIS using MAR grounded ice-masks (GIS area =  $12.37 \cdot 10^6 \text{ km}^2$ ). <sup>2</sup>RACMO2 data original source is van Wessem et al. (2018). Statistics in **bold** for ARP-AMIP-AC are statistically different from ARP-AMIP at  $p=0.05$

identified in Agosta et al. (2019). In this paper, it is argued that RACMO2 does not account for horizontal transport of falling precipitation and therefore misses some of the sublimation of snowfall in the dry katabatic layer such as shown in Grazioli et al. (2017). The same issue could be present in ARPEGE simulations due to the relatively large model physics time step used (15 min).

### 3.1.5 Surface mass balance

- 5 The SMB and its components integrated over the whole grounded ice sheet (GIS) for the 1981-2010 reference period are also presented in Table 4 for the two historical ARPEGE simulations and for ERA-I driven MAR and RACMO2 simulations. The spatial distribution of SMB for the different model and associated differences can be seen in Fig. 6. The same figure for surface sublimation is shown in the supplementary material (Fig. E1). The integrated GIS SMB is about four times the inter-annual variability uncertainty range ( $\sigma$ ) lower in ARP-AMIP-AC than in MAR-ERA-I and RACMO2-ERA-I as a
- 10 consequence of lower precipitation and largely overestimated surface sublimation rates (and, to a lesser extent, run-off). In the atmosphere-corrected simulation, there is a significant decrease of run-off and surface snow sublimation with respect to previously uncorrected simulation, yet insufficient to match with the estimates from the two polar-oriented RCMs. The overestimation is still large for the comparison with RACMO2 simulation where sublimation includes surface and blowing snow sublimation (see Table 4). The overestimation of surface sublimation in ARPEGE is consistent with the warm bias
- 15 evidenced previously in winter over the Antarctic Plateau and in summer at the fringe of the Plateau (see Fig. 4). The model deficiencies in capturing the frequent formation of very stable boundary layer in Antarctica is indeed expected to cause an overestimation of turbulent mixing and heat fluxes near the surface. These biases are also consistent with an overestimation of surface run-off and snow melt in summer. Estimations of surface snow melt in ARP-AMIP-AC are nevertheless within the  $1\sigma$  uncertainty range when compared to RACMO2 and MAR.


## Yearly mean Precipitation ( $\text{mmWe yr}^{-1}$ ), 1981-2010



**Figure 5.** Total Yearly mean total precipitation in ( $\text{mmWe yr}^{-1}$ ), surface snow sublimation for ARP-AMIP-AC (b) and SMB, ARP-AMIP (b) for ARP-AMIP-AC minus, MAR-ERA-I ( $\text{mmWe yr}^{-1}$ ) (d) and RACMO-ERA-I (g) for the reference period 1981-2010. Pink Difference ( $\text{mmWe yr}^{-1}$ ) for ARP-AMIP-AC minus ARP-AMIP (c), ARP-AMIP-AC minus MAR-ERA-I (e), ARP-AMIP minus MAR-ERA-I (f), ARP-AMIP-AC minus RACMO2-ERA-I (h) and blue ARP-AMIP minus RACMO2-ERA-I (i). Blue (magenta) hatched contour lines represents areas where ARPEGE-MAR differences are the positive (negative) difference is larger than 20%. Mean error (ME) and RMSE statistics ( $\text{mmWe mmWe yr}^{-1}$ ) for the comparison with MAR and RACMO2 are presented in upper-left corner shown below the subplot



# Yearly mean SMB ( $\text{mmWe yr}^{-1}$ ), 1981-2010



**Figure 6.** (a) ARP-AMIP minus RACMO2-ERA-I Yearly mean total precipitation-SMB ( $\text{mmWe yr}^{-1}$ ) for ARP-AMIP-AC (a), ARP-AMIP (b), MAR-ERA-I (d) and RACMO-ERA-I (g) for the reference period 1981-2010. Difference ( $\text{bmmWe yr}^{-1}$ ) same as for ARP-AMIP-AC minus ARP-AMIP (ac) but for, ARP-AMIP-AC minus AMIPMAR-ERA-I (e), ARP-AMIP minus MAR-ERA-I (f), ARP-AMIP-AC minus RACMO2-ERA-I (h) and ARP-AMIP minus RACMO2-ERA-I (i). Blue (magenta) hatched contour lines represents areas where the positive (negative) difference is larger than 20%. Mean error (ME) and RMSE statistics ( $\text{mmWe yr}^{-1}$ ) for the comparison with MAR and RACMO2 are presented in upper-left corner, shown below the subplot

## 3.2 Climate change signal


In this section, we present and compare the climate change signals for late 21<sup>st</sup> century obtained in the different ARPEGE RCP8.5 projections presented in this study. Climate change signals are obtained by computing the difference with their reference simulation in present-day climate (see Table 1). Differences in changes in atmospheric general circulation, near-surface temperature, precipitation and SMB obtained when using atmospheric bias-correction and their consistency with previous studies, are more specifically emphasized and discussed.

### 3.2.1 Atmospheric General Circulation

Changes in atmospheric general circulation are summarized by representing the Southern Hemisphere latitudinal profiles of sea-level pressure for each of the present-day simulation and future projections (Fig. 7). The simulated climate change signal is represented by the difference between each projection (coloured lines) and their reference simulation for present-day climate (dashed or plain lines). It can be seen that each future projection is characterized by a strengthening of the mid-latitude highs and a deepening of the circum-antarctic troughs and by a poleward movement of these features with respect to their reference historical simulation. This corresponds to an increasingly positive phase of the ~~SAM-index~~ Southern Annular Mode (SAM) index, the main mode of variability of atmospheric general circulation in the southern high-latitudes, which is in good agreement with the generally expected consequences of the increase in GHG concentration on the Southern Hemisphere's high latitude atmospheric circulation for the late 21<sup>st</sup> century (Arblaster and Meehl, 2006; Miller et al., 2006; Fyfe and Saenko, 2006).

However, it is noteworthy that both projections realized with atmospheric bias corrections show ~~much~~ substantially weaker changes in this ~~high-pressure-increase-(resp.-low-pressure-decrease)-pattern~~ increase of the meridional surface pressure gradient as well as a weaker poleward shift. This is confirmed by statistics from the changes in 850 hPa westerly maximum strength and position ( $\Delta$ WMPOS and  $\Delta$ WMSTR, Table 5). Each future projection displays ~~an-increase-in-WMSTR~~ a strengthening and poleward movement of the westerly maximum, but the magnitude of these changes are about 50 % weaker in projections realized with atmospheric bias correction. In similar experiments conducted with LMDZ model (Krinner et al., 2019) with different oceanic forcings, similar results were found. The magnitude of this difference is however much more reduced when compared to the results with ARPEGE. The realism of this reduced poleward shift and strengthening is discussed more extensively in section 4.1 and in the supplementary materials. Future projections realized with MIROC-ESM SSC (higher decrease in sea-ice) show a larger southward displacement of the westerlies than the one realised with NorESM1-M SSC, which is consistent with the impact of sea-ice extent on the position of the westerly winds maximum evidenced in previous studies (Krinner et al., 2014; Bracegirdle et al., 2018).

Another main difference in projected changes in atmospheric general circulation is the largely reduced deepening (not shown) of the Amundsen Sea Low (ASL), the main pressure climatological minimum, which is located at the fringe of the Amundsen and the Ross Sea in winter and off the Bellingshausen Sea, west of the Antarctic Peninsula in summer (Raphael et al., 2016). Blocking activity in the Amundsen Sea region and a negative phase of the SAM, which both correlate with El-Niño conditions



**Figure 7.** Yearly mean Sea-Level Pressure (hPa) for ARPEGE historical simulation (1981-2010) and RCP8.5 projections (2071-2100). Uncorrected control run are displayed in dashed lines and run with atmospheric bias-correction are presented in plain lines. Historical simulations realized with observed SSC are displayed in black (ARP-AMIP-AC) or gray (AMIP). ~~Projections~~ Future projections (RCP8.5) driven by bias-corrected SSC from NorESM1-M (MIROC-ESM) are shown in red (green).

**Table 5.** Anomalies in Southern westerlies maximum strength ( $\Delta\text{WMSTR}$ , m/s) and position ( $\Delta\text{WMPOS}$ , °) for the different ARPEGE projections

Simulations	$\Delta\text{WMSTR}$ (m/s)	$\Delta\text{WMPOS}$ (°)
ARP-NOR-21-OC	1.5	-2.2
ARP-NOR-21-AOC	0.8	-0.8
ARP-MIR-21-OC	2.0	-3.8
ARP-MIR-21-AOC	0.9	-1.5

(Scott Yiu and Maycock, 2019), have been found to influence warming rates in West Antarctica (Scott et al., 2019). Since the SAM pattern and the deepening of the ASL were found to be different in ARPEGE bias-corrected projections, this will likely impact blocking activity in the Amundsen Sea Region and therefore warming in West Antarctica. A detailed analysis in our ARPEGE simulation of this relationship between regional warming and blocking activity associated with the propagation of

5 Rossby-waves in the Pacific sector is however beyond the scope of this study.

### 3.2.2 Near-surface temperature

The increase in mean yearly  $T_{2m}$  averaged over the GIS are respectively  $3.5 \pm 1.0$  and  $5.0 \pm 1.3$  °C for ARP-NOR-21-AOC and ARP-MIR-21-AOC . This represents respectively a +0.7 and +0.9 °C additional warming (significant at  $p < 0.05$ ) compared to the corresponding projections without atmospheric bias correction. Differences in warming rate per season are presented in Table 6. Differences in warming for the atmospheric-corrected simulations are the largest in summer and are significant for both projections, while they are smaller and not significant in winter. For ARP-NOR-21-AOC , the larger warming in autumn (MAM) is also significant ~~, while and so~~ it is in spring for ARP-MIR-21-AOC .


The spatial distribution of the increase in  $T_{2m}$  and corresponding differences in winter (JJA) and summer (DJF) warming are presented in Fig. 8. The two sets of projections show very similar patterns in terms of differences in regional warming. The larger surface warming in summer in the atmospheric-corrected experiment is essentially the consequence of a stronger temperature increase over East Antarctica. This larger additional warming found for the surface of the East Antarctic Plateau in summer (+1 to +2 °C) is also found at higher altitudes (500 hPa), which results in increased LWD. This additional warming is consistent with the weaker increase of the pressure gradient and a reduced poleward shift of the westerly winds in these projections, as this corresponds to a lower increase towards a more positive phase of the SAM in future climate. The link between negative (positive) anomalies of the SAM and positive (negative) temperatures anomalies over the East Antarctic Plateau has been established in many previous studies (e.g., Marshall and Thompson, 2016; Kwok and Comiso, 2002) and was found to be stronger in summer and autumn (Marshall, 2007; Clem et al., 2016). However, following this hypothesis and the findings from Marshall and Thompson (2016), less pronounced positive phase of the SAM should also result in a larger warming over West Antarctica and a weaker warming over the north-eastern Peninsula in atmosphere-corrected projections, which is not the case here.

In winter, there is a well-marked dipole with lower warming over West Antarctica and the Ross Ice Shelf and higher warming over southern Victoria and Adélie Land. The much weaker deepening of the Amundsen Sea Low found in our atmosphere-corrected experiments is consistent with these differences in regional warming. The mean position of the Amundsen Sea Low was indeed shown to be located over the east-side of the Ross Sea in winter (Raphael et al., 2016), and so it is in ARPEGE simulations.

Systematically, near-surface temperature increases are the strongest where sea ice is lost. This is particularly noticeable over the northern part of the Weddell Sea. This area (together with Larsen and Ronne-Filchner ice shelves) shows a large additional warming in atmosphere-corrected experiments. ~~In winter, there is a well-marked dipole with lower warming over West Antarctica and the Ross Ice Shelf and higher warming over southern Victoria and Adélie Land.~~

Overall, most of the differences in near-surface temperatures warming found are consistent with corresponding differences in large-scale atmospheric circulation changes and the relation found between pressure and temperature anomalies in previous studies.





**Figure 8.**  $T_{2m}$  anomaly for ARPEGE RCP8.5 projection for the late 21<sup>st</sup> century (reference period : 1981-2010) with atmospheric bias-correction (AOC, left), uncorrected atmosphere (OC, center) and difference (right). Anomalies for winter (JJA) are displayed at the top of the subfigures and for summer (DJF) at the bottom. Results for projections with bias-corrected SSC from NorESM1-M (resp. MIROC-ESM) are shown in a) (resp. b). Grey contour lines indicate where differences in anomaly are  $> 25\%$  with respect to the uncorrected control run.



**Table 6.** Mean seasonal  $T_{2m}$  increase (K) for the Antarctic GIS for the different ARPEGE RCP8.5 projection for the late 21<sup>st</sup> century with respect to their historical reference simulation. Anomalies in projections with bias-corrected atmosphere significantly different ( $p < 0.05$ ) from the anomaly in the uncorrected control run level are shown in *bold*.

Simulations	DJF	MAM	JJA	SON
ARP-NOR-21-OC	3.0±1.4	2.6±1.4	3.1±1.4	2.6±1.0
ARP-NOR-21-AOC	<b>3.8±1.2</b>	<b>3.7±1.2</b>	3.3±1.9	3.4±1.9
ARP-MIR-21-OC	3.6±1.5	4.6±1.7	4.6±1.4	3.8±1.5
ARP-MIR-21-AOC	<b>5.1±1.4</b>	5.2±1.7	5.1±1.7	<b>4.8±2.0</b>

### 3.2.3 Surface mass balance Precipitation and precipitationSurface Mass Balance

Absolute and relative increase in SMB and for its different components for the Antarctic GIS are presented in Table 7. All projections show an increase in surface mass balance, resulting from the absolute increase in precipitation still being much larger than the corresponding increases in surface sublimation and run-off. This is in agreement with previous Antarctic climate change studies for the late 21<sup>st</sup> century (e.g., Lenaerts et al., 2016; Krinner et al., 2014; Frieler et al., 2015). Additional increases in total precipitation of +78 and +90 Gt yr<sup>-1</sup>, which corresponds respectively to a +6 and +9 % additional relative increase (not significant at  $p < 0.05$ ), are found respectively for ARP-NOR-21-AOC and ARP-MIR-21-AOC atmospheric corrected projections. In ARP-MIR-21-AOC, significant higher increase in surface sublimation mitigates slightly the increase in SMB with respect to ARP-MIR-21-OC. Despite a larger warming at the continental scale, increases and cumulated amounts of precipitation are significantly lower in projections with atmospheric bias correction.

There is an additional increase in moisture transport towards the AIS (approximated through P-E) of +3.5 and +5.8 % respectively with respect to uncorrected control run. The sensitivity to temperature of the increase in precipitation ( $\alpha$ ) in ARP-NOR-21-AOC and ARP-MIR-21-AOC are respectively +7.7%.K<sup>-1</sup> and +9.1%.K<sup>-1</sup>, whereas these were respectively +5.2% and +8.8% in the uncorrected control run. This suggests that a significant part of the additional increase in precipitation is due to different changes in the atmospheric general circulation, particularly in the projection driven by NorESM1-M oceanic boundary conditions, while a remaining part is of course due to increased moisture holding capacity of the atmosphere due to higher warming. The latest values found for  $\alpha$  are somewhat higher than previous estimates (Krinner et al., 2008; Frieler et al., 2015; Ligtner et al., 2015) which usually range between +5 and +7 %.K<sup>-1</sup>. However, Bracegirdle et al. (2015) also evidenced that  $\alpha$  tends to be higher in models projecting a larger decrease in sea-ice extent, which is the case for experiments forced by MIROC-ESM sea-ice anomaly.

The spatial distribution of precipitation changes for each projection are shown in Fig. 9 along with the differences between the corrected simulations and the uncorrected control runs. Projections with atmospheric bias correction show a smaller precipitation increase over most of West Antarctica, the Transantarctic Mountains and western Dronning Maud Land. Conversely, the increase in precipitation is larger on the coastal AP, on the Ross side of Marie-Byrd Land and on Adélie and southern Victoria Land, where uncorrected projections suggest a precipitation decrease. The higher precipitation increase over

**Table 7.** Absolute values, absolute (Abs.) and relative (Rel.) anomalies for mean SMB and its components ( $\text{Gt yr}^{-1}$ ) for the Antarctic GIS in the different ARPEGE RCP8.5 projections (reference period: 1981-2019). Anomalies and absolute values significantly different ( $p < 0.05$ ) in projections realized with bias-corrected atmosphere with respect to control run are displayed in bold.

Simulations	SMB	Tot. PCP	Surf. Sublim.	Rainfall	Melt
<b>ARP-NOR-21-OC</b>	<b>2334±181</b>	<b>2742±176</b>	331±21	27±7	184±82
<i>Abs. anomaly (<math>\text{Gt yr}^{-1}</math>)</i>	364±195	474±179	55±26	17±8	132±137
<i>Rel. anomaly</i>	19%	21%	20%	171%	252%
<b>ARP-NOR-21-AOC</b>	<b>2172±143</b>	<b>2534±158</b>	<b>284±19</b>	<b>21±2</b>	210±79
<i>Abs. change (<math>\text{Gt yr}^{-1}</math>)</i>	422±169	540±176	62±18	<b>16±8</b>	<b>160±64</b>
<i>Rel. change</i>	24%	27%	28%	320%	320%
<b>ARP-ARP-MIR-21-OC</b>	<b>2637±156</b>	<b>3108±202</b>	345±29	52±15	306±144
<i>Abs. change (<math>\text{Gt yr}^{-1}</math>)</i>	<b>667±202</b>	<b>840±227</b>	68±23	42±15	254±118
<i>Rel. change</i>	34%	37%	25%	416%	484%
<b>ARP-MIR-21-AOC</b>	<b>2460±197</b>	<b>2903±222</b>	<b>308±23</b>	<b>45±14</b>	359±118
<i>Abs. change (<math>\text{Gt yr}^{-1}</math>)</i>	709±218	909±229	<b>86±23</b>	<b>40±15</b>	309±104
<i>Rel. change</i>	40%	46%	39%	800%	618%

Adélie and southern Victoria Land, and the corresponding weaker increase over most of West Antarctica can be related, as for the corresponding dipole in differences of warming in winter, to the weaker deepening to the Amundsen Sea Low in bias-corrected projections. Winter is indeed, with autumn, the season of highest precipitation rates over peripheral Antarctica (Palerm et al., 2017).


- 5 Additionally, Palerm et al. (2017) found that CMIP5 models which agree better with Antarctic snowfall derived from CloudSat show a larger warming ( $+0.4\text{ }^{\circ}\text{C}$ ) and a higher precipitation increase ( $+4.8\%$ ) with respect to the ensemble mean in their RCP8.5 projections. In light of our results, it would be interesting to investigate whether models agreeing with CloudSat snowfall are doing so because of a better representation of the atmospheric general circulation and whether the higher increase in temperature and precipitation can be generally linked to the better representation of atmospheric general circulation in the
- 10 context of CMIP ensembles such as found in our bias corrected experiment.

## 4 Discussion

### 4.1 Representation-Realism of projected changes

In future works, the remaining uncertainties on the less pronounced strengthening of the pressure gradient and the historical

- 15 climateIn this section, we discuss following our comparisons with ERA-I reanalyses, *in-situ* observations and polar-oriented RCMs, the improvement in skills associated with atmospheric empirical in-line bias correction of ARPEGE model for the



**Figure 9.** Late 21<sup>st</sup> century anomalies in yearly total precipitation (%) for ARPEGE RCP8.5 projections with atmospheric bias-correction (*left*), uncorrected reference simulation (*center*) and difference (*right*). Results for projections with bias-corrected SSC from NorESM1-M (resp. MIROC-ESM) are presented in a) (resp. b).

atmospheric general circulation in the high southern latitudes as well as for the surface climate of the Antarctic ice sheet. Where relevant, we also discuss possible biases compensation suppression associated the systematic errors correction for atmospheric general circulation. The limitations and potential improvement of the method are also addressed.

#### 4.1.1 Large-scale atmospheric circulation

- 5 Compared to ERA-I, the simulated large-scale atmospheric circulation South of 20°S using atmospheric bias correction in ARP-AMIP-AC is dramatically improved with respect to the uncorrected ARP-AMIP control run. Reduction of the seasonal RMSE between 50 to 70 % with respect to ARP-AMIP are found for many mid and upper tropospheric variables. Improvement for the position and strength of the mid-latitude surface westerly jet are also clear and the difference with ERA-I are insignificant in ARP-AMIP-AC. The analysis of the best matching unit frequencies in the self-organizing maps analysis also evidenced much
- 10 better representation of daily variability in ARP-AMIP-AC. In this regard, it is noteworthy that Guldberg et al. (2005) also found large (20-30 %) improvement in the seasonal forecasting skills of their corrected model in the Southern Hemisphere while no improvement was reported on average in the Northern Hemisphere.

In our experiment, we found no bias reduction in tropospheric humidity (Q850 and Q500) and lower tropospheric temperature (T850). As a reminder, 850 hPa is the level at which the correction on the tendency for the atmospheric variables starts to decrease towards the surface. The new bias patterns appearing at this level in ARP-AMIP-AC over southern mid-latitudes land masses and the adjacent oceans (see Section. ??) are most likely linked to surface processes (e.g., development of marine stratocumulus, convective boundary layer). The typical time scale for the development of such processes is shorter than the 72 h relaxation time used in the nudged simulation realized to build the correction terms. As a consequence, this advocates for further sensitivity tests using different relaxation times depending on the variable, especially for specific humidity, in order to improve the skills of the corrected model for this variable in future similar experiment. Re-tuning the atmospheric model during the nudging step could also be an option to ensure consistency of the free physical parameters with the modified circulation characteristics.

#### 4.1.1 Near-surface temperature

The correction of large-scale circulation errors in ARP-AMIP-AC has clearly lead to a reduced cold bias found in ARP-AMIP control run over the western and southern part of the Antarctic Peninsula, especially in winter. The skill of the model improves dramatically for stations such as Faraday and Rothera and for most stations of the AP in general. Warm and moist advection from the North-West over this part of the Peninsula is indeed underestimated in ARP-AMIP. However, the east coast of the Peninsula has cooled dramatically in ARP-AMIP-AC and there is now a substantial cold bias with respect to MAR. The skill of ARP-AMIP-AC also exceeds that of ARP-AMIP for coastal stations of East Antarctica (e.g., Mc Murdo, Scott Base, Casey and Davis). However, for many stations, the cold bias remains substantial and statistically significant. In this perspective, the comparison with Mc Murdo and Scott Base is interesting. These stations, distant by only 3 kilometers and located at the same altitude, belong to the same ARPEGE grid point. While ARP-AMIP-AC bias with respect to Scott Base is small and statistically insignificant in all season but summer, the cold bias is  $\sim 3^\circ\text{C}$  (significant) throughout the year with respect to Mc Murdo. This example shows the limited spatial representativity of weather stations in coastal Antarctica, and therefore the comparisons with a 35 km horizontal resolution atmospheric model should therefore be interpreted carefully. On the East Antarctic Plateau, the warm bias in winter increases in ARP-AMIP-AC in both comparisons with weather station and MAR-ERA-I. This bias increase seems to result from the removal of a bias compensation that was present in RP-AMIP. We investigated the value of the near-surface temperature inversion in both present-day ARPEGE simulation (not shown) as the difference between surface temperature ( $T_S$ ) and the temperature of the first atmospheric layer, located around 6 to 10 meters height in this ARPEGE setting (level at  $0.9988 \times \text{surface pressure}$ ). The value of large magnitude of the reduced poleward shift found in atmospheric bias-corrected experiments should be reduced with additional experiments using either other set oceanic surface forcing or other atmospheric models, considering the large impacts of these on the projected climate change for the Southern Hemisphere high-latitudes. These results are however consistent with results from Barnes and Hartmann (2012) who found, using GCMs future projections and an idealised-case barotropic model, that there is a theoretical limit to how far South the maximum of cyclonic wave breaking can move as these dynamical changes are constrained by the absolute vorticity gradient. Their results suggested that the currently observed Southern Hemisphere general circulation might already be close

to this limit. Investigating these same physical mechanism such as done in Barnes and Hartmann (2012) in our ARPEGE bias-corrected experiments is out of the scope of this near-surface temperature inversion around ( $-15^{\circ}\text{C}$ ) does not decrease in ARP-AMIP-AC. It even increases locally, suggesting small changes in surface boundary layer processes. However, there was a substantial negative bias with respect to MAR in incoming long-wave radiation (LWD) in ARP-AMIP, which disappears in ARP-AMIP-AC due to considerable warming at 500 hPa in winter study and generalising results from a set of only two pair of thirty years experiments would be impossible due to climate's internal variability. Nevertheless, when we put them in the context of previous CMIP5 large ensemble analyses (Bracegirdle et al., 2013), we confirm that these processes are likely at play in our simulations (Fig. ??). The warm bias in winter with respect to MAR, which generally varies between 3 and  $5^{\circ}\text{C}$  (similar biases are found when considering READER weather stations or RACMO2  $T_{2m}$ ) is to be put into perspective with even larger biases sometimes found over the East Antarctic Plateau in climate models or in even climate reanalyses (Dutra et al., 2015; Fréville et al., 2014; Bracegirdle and Marshall, 2012) [B4](#).

The warm temperature bias over Antarctic ice shelves does not decrease in ARP-AMIP-AC, it even increases over Neumayer and Halley stations, especially in winter. ARPEGE model deficiencies for surface temperatures over the large Ronne-Filchner and Ross ice shelves have been widely addressed in Beaumet et al. (2019a). They are mostly due to misrepresented stable boundary layers processes and excess (with respect to MAR) in long-wave downward radiation in winter associated with higher cloud cover. Large discrepancies have also been evidenced between polar-oriented RCMs MAR and RACMO2 for these areas.

#### 4.1.1 Surface mass balance and precipitation

Over the GIS, the atmospheric bias correction caused a decrease of total precipitation of  $285 \text{ Gt yr}^{-1}$  ( $\sim 3$  times the interannual variability). Unsurprisingly, the precipitation minus evaporation difference (P-E), which is a good approximation of moisture transport, has also decreased, by  $221 \text{ Gt yr}^{-1}$ . The main feature of the atmospheric bias correction in ARP-AMIP-AC with respect to ARP-AMIP is an increase of SLP at mid-latitude, and a decrease around Antarctica, which can be seen as a larger dominance of  $\text{SAM}^+$ -type patterns (see Fig. 3). In this regard, we note that changes in precipitation in ARP-AMIP-AC with respect to ARP-AMIP (Fig. ??b) bear many similarities with the signature of a positive SAM pattern on Antarctic precipitation assessed in a RACMO2 ERA-Interim driven simulation in Marshall et al. (2017). It is noteworthy that the link between circulation changes induced by the bias correction and ensuing precipitation changes seen in our simulations is very similar to the effect of bias corrections in the LMDZ model reported by Krinner et al. (2019). The correction of atmospheric circulation errors has caused a drying over Dronning Maud and Marie Byrd Land and an increase in precipitation over central and eastern West Antarctica as well as on the western side of the Peninsula. In these regions, the agreement with MAR and RACMO2 has increased substantially. The improvement of the agreement (RMSE) is much larger for the comparison with RACMO2, and except for some parts the AP and of [Even though the realism of projected changes in bias-corrected experiments still and will always bear uncertainties \(as by construction future climate projections are impossible to verify\), the uncorrected projections are expected to be of limited use for impact assessment studies since biases in the historical reference experiment are of the same order of magnitude as projected changes. In such impact assessment studies, absolute future state of the climate is](#)

more relevant than relative projected changes. An interesting example is given with the impact of the position and depth of the Amundsen Sea Low on the regional changes in temperature and precipitation over western West Antarctica, Victoria and Adélie Land; the agreement is quite high (errors below 20 %) for many coastal areas. The comparison with both RCMs also reveals that ARP-AMIP-AC is likely too dry over much of the East Antarctic Plateau. Further investigations are needed in order to identify possible causes of this error, such as deficiencies in moisture transport over the high continental interior or misrepresented clear-sky precipitation processes, which bring a large share of total precipitation in this area (Walden et al., 2003). Improvement for the representation of precipitation, cloudiness and therefore radiative budget for the high continent interior of Antarctica could be obtained for ARPEGE by accounting for saturation. While the more reliable deepening and displacement of the ASL in the bias corrected projections with respect to non-corrected reference can still to some extent be questioned, the impact of this deepening and displacement on regional temperature and precipitation changes are by construction more realistic in the bias-corrected projections as the ice for the formation of cloud condensates and by tuning cloud microphysics to the cold context of Antarctica such as done in Van Wessem et al. (2014); van Wessem et al. (2018). The better agreement with precipitation modelled by the polar-oriented RCMs MAR and RACMO2 precipitation increase the confidence in the reliability of spatial distribution of Antarctic SMB modeled in ARP-AMIP-AC. Both RCMs have been widely validated against *in-situ* measurements of Antarctic SMB (e.g., Agosta et al., 2019; van Wessem et al., 2018). The closer agreement between the ARPEGE ARP-AMIP-AC simulation and RACMO2 in coastal areas offers an interesting opportunity to further investigate causes of remaining disagreement between MAR and RACMO2 in Antarctic precipitation and SMB identified in Agosta et al. (2019). In this paper, it is argued that RACMO2 struggles to represent sublimation of snowfall in the dry katabatic layer present at the ice sheet margins and in valleys such as shown in Grazioli et al. (2017). The same issue could be present in ARPEGE simulations due to the relatively large model physics timestep used (15 min). Over the lee-side of the Transantarctic mountains (Victoria Land), the good agreement between ARPEGE and RACMO2 seems to confirm a wet bias in MAR over this area, which has also been identified after comparison with and sparse *in-situ* observations in Agosta et al. (2019).

## 4.2 Climate change signals

In this section, we discuss the differences in projected climate change for atmospheric general circulation of the southern high-latitudes and for Antarctic surface climate between corrected and uncorrected projections and their consistency with previous studies.

### 4.1.1 Large-scale atmospheric circulation

The increase of the surface pressure gradient between mid and high Southern latitudes for the late 21<sup>st</sup> century is substantially weaker in the atmosphere-corrected projections. The strengthening and the poleward shift of the 850 hPa westerly wind maximum are reduced by about 50 %. This weaker climate change signal for polarward shift in the corrected simulations can be related to the dependence on historical state of projected changes in atmospheric circulation evidenced in Bracegirdle et al. (2013). However, concerning the strength of the westerlies, Bracegirdle et al. (2013) did not find a significant link between the simulated historical state and projected changes. Larger differences in sea-level pressure projected changes found over the Pacific sector



in our atmosphere-corrected experiment are also consistent with this results from Bracegirdle et al. (2013) who found that the historical state dependence was stronger in this area. In similar experiments conducted with LMDZ model (Krinner et al., 2019) with different oceanic forcings, a smaller decrease (increase) of the high latitudes low (mid-latitudes highs) pressure is also found. The magnitude of these difference is however much more reduced when compared to the results with ARPEGE. In future works, the less pronounced strengthening of the pressure gradient and the large magnitude of this reduced poleward shift found for atmospheric-corrected ARPEGE experiment however need to be confirmed with supplementary experiments using either other set oceanic surface forcing or other atmospheric models, considering the large impacts of these on the projected climate change for the Southern Hemisphere high latitudes biased position and depth of the ASL in the historical uncorrected simulation already induce incorrect regional temperature and precipitation patterns for present time.

#### 4.1.1 Near-surface temperatures

The additional 0.7 to 0.9 °C warming of the Antarctic GIS found in both atmosphere-corrected projections is largely due to the much larger warming of East Antarctica, especially in summer

Assessing the impact of extreme events in a changing climate using projections that have highly biased representation of Antarctic's climate mean state and variability in their historical reference is unrealistic. In our study, we found that climate's

variability at the daily time scale is better represented in ARPEGE bias corrected experiment. This is consistent with a weaker increase of the pressure gradient in these projections, as this corresponds to a lower increase towards a more positive phase of the SAM in future climate. The link between negative (positive) anomalies of the SAM and positive (negative) temperatures anomalies over the East Antarctic Plateau has been established in many previous studies (e.g., Marshall and Thompson, 2016; Kwok and Co. However, following this hypothesis and findings from Marshall and Thompson (2016), less pronounced positive phase of the SAM should also result in a larger warming over West Antarctica and a weaker warming over the northern Peninsula in atmosphere-corrected projections, which is not the case here. Nevertheless, we note that in Marshall and Thompson (2016), temperature variability over West Antarctica and over the Peninsula is also strongly linked to the polarities of the first and second Pacific South American modes (PSA1 and PSA2), the second and third leading empirical functions of geopotential heights south of 20°S. Moreover, the much weaker deepening of the Amundsen Sea Low found in our atmosphere-corrected experiments is consistent with weaker increase in winter temperatures over West Antarctica and higher increase over southern Adélie and Victoria Land. The mean position of the Amundsen Sea Low was indeed shown to be located over the east-side of the Ross Sea in winter (Raphael et al., 2016), and so it is in ARPEGE simulations. The large additional warming found for the surface of the East Antarctic Plateau in summer (+1 to +2 °C) is also found higher up in altitude (500 hPa), which results in increased LWD. Overall, most of the differences in near-surface temperatures warming found are consistent with corresponding differences in large-scale atmospheric circulation changes and the relation found between pressure and temperature anomalies in previous studies.

#### 4.1.1 Precipitation and surface mass balance



Both atmospheric bias-corrected projections suggest a higher, yet not significant, precipitation increase of respectively +6 and +9 % in ARP-NOR-21-AOC and ARP-MIR-21-AOC. There is an additional increase in moisture transport towards the AIS (approximated through P-E) of +3.5 and +5.8 % respectively with respect to uncorrected control run. The sensitivity to temperature of the increase in precipitation ( $\alpha$ ) in ARP-NOR-21-AOC and ARP-MIR-21-AOC are respectively +7.7%.K<sup>-1</sup> and +9.1%.K<sup>-1</sup>, whereas these were respectively +5.2% and +8.8% in the uncorrected control run. This suggests that a large part of the additional increase in precipitation is due to different changes in the atmospheric general circulation, particularly in the projection driven by NorESM1-M oceanic boundary conditions, while a remaining part is of course due to increased moisture holding capacity of the atmosphere. The latest values found for  $\alpha$  are somewhat higher than previous estimates (Krinner et al., 2008; Frieler et al., 2015; Ligtenberg et al., 2013; Palerme et al., 2017), which usually range between +5 and +7 %.K<sup>-1</sup>. However, Bracegirdle et al. (2015) also evidenced that  $\alpha$  tends to be higher in models projecting a larger decrease in sea-ice extent, which is the case for experiments forced by MIROC-ESM sea-ice anomaly.

In addition to this, Palerme et al. (2017) found that CMIP5 models which agree better with Antarctic snowfall derived from CloudSat show a larger warming (+0.4 °C) and a higher precipitation increase (+4.8 %) with respect to the ensemble mean in their RCP8.5 projections. Therefore, it would be interesting to investigate whether models agreeing with CloudSat snowfall are doing so because of an important result as by construction the empirical bias correction is expected to reduce biases on mean state only. Krinner et al. (2019, 2020) found similar results in their application with LMDZ and other atmospheric models for the representation of climate variability ranging from daily to inter-decadal time scale. This is another argument in favour of using bias-corrected projections for impact assessment studies as a better representation of the atmospheric general circulation. At the regional scale, the higher precipitation increase over Adélie and southern Victoria Land, and the corresponding weaker increase over most of West Antarctica can be related, as for the corresponding dipole in differences of warming in winter, to the weaker deepening to the Amundsen Sea Low. Winter is indeed, with autumn, the season of highest precipitation rates over peripheral Antarctica (Palerme et al., 2017). climate variability is expected to bring more useful information on extreme events.

## 4.2 Implication of bias-correction and perspectives

The large bias reductions for large-scale atmospheric circulation and surface climate obtained in this study should not be perceived as an argument against pursuing the efforts to improve the dynamics and physics of coupled and atmospheric models in a physically consistent and comprehensive way, nor as a loss of confidence in these tools. These are crucial in order to explore some feedbacks and interactions between the different component of the Earth system in a warming climate. Yet, as long as biases of state-of-the-art climate models are of about the same order of magnitude as projected changes at the end of current century (Flato et al., 2013), *a posteriori* statistical bias-correction (Hall, 2014; Maraun and Widmann, 2018) will be applied to future projections before they are used as input for impact studies-assessment-assessment studies. However, these methods fail to correct for biases due to atmospheric circulation errors (Eden et al., 2012; Stocker et al., 2015; Maraun et al., 2017). Therefore, the method presented in this study offers an excellent opportunity to circumvent this drawback. Bias stationarity is a strong hypothesis needed to support the application of run-time atmospheric bias-correction in climate projections.

However, the evidence of large stationarity in biases of coupled models on large-scale atmospheric circulation evidenced in Krinner and Flanner (2018) and the preservation of the added-value of the value until late 21st century climate evidenced in Krinner et al. (2020) using the perfect model test framework supports this application under strong climate change.

~~Applying this-~~

- 5 Bias correction can be easily turned off and its impact on projected climate change can easily be identified. Therefore, applying these bias correction method also allows to assess remaining uncertainties on projected climate change in coupled and atmospheric climate models ~~-, and it could help to identify~~ such as suggested by Dommenges and Reznay (2018) and could help identifying which efforts one should focus on in order to reduce these uncertainties. ~~Because~~ Considering it removes most of the biases on large-scale atmospheric circulation, the bias-correction method implemented in this study also offers the
- 10 opportunity to test the tuning of the parametrizations of smaller scale processes (e.g. boundary layer) in a different context (freely evolving simulation) than nudged simulations often used in this purpose. This would allow to test whether introduced modifications in new parametrizations are detrimental to the model skills on large-scale atmospheric circulation when the model is not constrained by atmospheric reanalysis.

- Even though they still bear some uncertainties, the significant differences in large-scale atmospheric circulation and surface
- 15 climate changes reported in this study could have large impacts on projected changes of the Antarctic ice-sheet mass balance. Therefore, we suggest to use surface forcing provided by our ~~new~~ bias-corrected projections to drive ice-dynamics or ocean and ice shelves interactions studies. Downscaling the projections presented in this study with polar oriented RCMs such as MAR and RACMO2 could also help to identify remaining uncertainties associated with ARPEGE biases on Antarctic surface climate and SMB mostly associated with its less sophisticated surface snow scheme ~~-, and these new downs-called projections~~
- 20 would help better constraining future ice dynamics or ice-ocean-atmosphere interactions.

## 5 Summary and Conclusion

- In this study, we used empirical run-time bias correction following the method described in Guldberg et al. (2005) or in Kharin and Scinocca (2012). In order to build correction terms, we used the climatology of the adjustment term on tendency errors
- 25 coming from an ERA-Interim driven ARPEGE simulation over the 1993-2010 period. In this experiment, nudged variables were air temperatures, air specific humidity, logarithm of surface pressure, divergence and vorticity with a relaxation time of 72 hours.

- The application of this method over present climate (1981-2010) yielded a ~~substantially improved~~ substantial bias reduction in the representation of large-scale atmospheric circulation in the Southern Hemisphere. The biases of westerly wind maximum
- 30 position and strength were almost completely suppressed. This improvement of Southern Hemispheric general circulation produced a decrease of the biases on near-surface temperature over the Antarctic Peninsula, while we found a slightly increased warm winter bias on the East Antarctic Plateau. Regarding precipitation, the agreement with polar-oriented RCMs MAR and RACMO2 has increased, especially in the comparison with the latest, where differences below 20 % are reported over most

coastal areas. A dry bias in the atmosphere-corrected experiment over the summit of the East Antarctic Plateau was also evidenced by this comparison with polar-oriented RCMs.

The application of the method for future climate projections (RCP8.5) using bias-corrected oceanic forcing from MIROC-ESM and NorESM1-M has revealed considerable differences in projected changes in large-scale atmospheric circulation. The strengthening and the poleward shift of the westerly wind maximum are reduced by about 50% with respect to uncorrected reference projections. These differences in change of atmospheric general circulation have caused significant additional warming of +0.7 to 0.9 K, resulting essentially from the much larger warming of East Antarctica in summer. A dipole with higher warming and increase in precipitation over southern Victoria and Adélie Land and corresponding lower warming and increase in precipitation over most of West Antarctica is also found in winter. This is attributed to a reduced deepening of the Amundsen Sea Low in the atmosphere-corrected projections.

The magnitude of the difference in changes of large-scale atmospheric circulation needs to be confirmed with experiments using other oceanic forcing or other atmospheric models, as these would have large impacts on the evolution of the Antarctic ice-sheet mass balance. However, a reduced poleward shift of the westerlies maximum in the bias-corrected experiments is consistent with the state dependence on historical biases in CMIP5 coupled and atmospheric models projections evidenced by ~~Bracegirdle et al. (2013)~~ in previous studies. Many of the differences found in projected changes in temperature and precipitation are also consistent with previously evidenced signatures of pressure anomalies, especially considering a weaker SAM+ anomaly (Kwok and Comiso, 2002; Marshall and Thompson, 2016; Marshall et al., 2017).

Because statistical bias corrections, usually applied to climate model output before their use for impact studies, generally fails to correct biases associated with errors on atmospheric general circulation (e.g., Hall, 2014; Maraun et al., 2017; Maraun and Widmann, 2017), the method proposed in this study offers interesting perspectives. The downscaling of atmosphere-corrected projections with polar-oriented RCMs could help to better constrain the evolution of the future Antarctic ice-sheet surface mass balance and evaluate remaining uncertainties in this study associated with ARPEGE biases on surface climate. The potentially large effect on the Antarctic ice-sheet of the differences in snow accumulation and surface climate changes suggested in this study should be explored using surface forcings coming from atmosphere-corrected projections to drive ice-dynamics or ocean and ice shelves interactions impact studies.

*Data availability.* Historical run and future projections presented in this study are available on Antarctic Cordex grid at daily time step for atmospheric surface temperature (mean, min and max), precipitation, snowfall, surface snow melt, surface run-off and surface snow sublimation at the following address: [https://zenodo.org/record/4059193#.X4mUgCU6\\_RZ](https://zenodo.org/record/4059193#.X4mUgCU6_RZ) (DOI : 10.5281/zenodo.4059193). Data and metadata mostly follow Cordex recommendations.

*Competing interests.* The authors have no competing interests.

*Acknowledgements.* This publication was supported by PROTECT. This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 869304. This is PROTECT contribution number XX.

We acknowledge the World Climate Research Programme's Working Group on Coupled Modelling, which is responsible for CMIP, and we thank the climate modeling groups participating to CMIP5 for producing and making available their model output. For CMIP the U.S.

5 Department of Energy's Program for Climate Model Diagnosis and Intercomparison provides coordinating support and led development of software infrastructure in partnership with the Global Organization for Earth System Science Portals.

The Centre National de Recherches Météorologique (Météo-France, CNRS) and associated colleagues are warmly thanks for providing resources and help to run ARPEGE model.

We also thank the Scientific Committee on Antarctic Research, SCAR and the British Antarctic Survey for the availability of the SCAR-  
10 READER data base.

We kindly acknowledge Michiel van den Broeke for providing RACMO2 data and helpful discussions.

Finally, we want to thanks the two anonymous reviewers for their constructive remarks and the editor for helping in improving the quality of this manuscript.

*Author contributions.* AA prepared ARPEGE version and scripts that were used to produce ARPEGE runs used in this study. MD performed  
15 the sensitivity tests and ran the bias-corrected simulations. JB ran the non-corrected reference runs, analysed the output and prepared the manuscript and the figures. CA provided MAR outputs and usefull scripts for the interpolation and comparisons of MAR, RACMO2 and ARPEGE output. GK read and corrected the manuscript many times. All the authors have provided complete reviews and relevant remarks for the wement of the manuscript.

## References

- Agosta, C., Favier, V., Krinner, G., Gallée, H., Fettweis, X., and Genthon, C.: High-resolution modelling of the Antarctic surface mass balance, application for the twentieth, twenty first and twenty second centuries, *Climate Dynamics*, 41, 3247–3260, <https://doi.org/10.1007/s00382-013-1903-9>, 2013.
- 5 Agosta, C., Amory, C., Kittel, C., Orsi, A., Favier, V., Gallée, H., van den Broeke, M. R., Lenaerts, J., van Wessem, J. M., van de Berg, W. J., et al.: Estimation of the Antarctic surface mass balance using the regional climate model MAR (1979–2015) and identification of dominant processes, *The Cryosphere*, 13, 281–296, <https://doi.org/10.5194/tc-2018-76>, 2019.
- Arblaster, J. M. and Meehl, G. A.: Contributions of External Forcings to Southern Annular Mode Trends, *Journal of Climate*, 19, 2896–2905, <https://doi.org/10.1175/JCLI3774.1>, 2006.
- 10 Barnes, E. A. and Hartmann, D. L.: Detection of Rossby wave breaking and its response to shifts of the midlatitude jet with climate change, *Journal of Geophysical Research: Atmospheres*, 117, <https://doi.org/https://doi.org/10.1029/2012JD017469>, 2012.
- Beaumet, J., Déqué, M., Krinner, G., Agosta, C., and Alias, A.: Effect of prescribed sea surface conditions on the modern and future Antarctic surface climate simulated by the ARPEGE atmosphere general circulation model, *The Cryosphere*, 13, 3023–3043, 2019a.
- Beaumet, J., Krinner, G., Déqué, M., Haarsma, R., and Li, L.: Assessing bias corrections of oceanic surface conditions for atmospheric  
15 models, *Geoscientific Model Development*, 12, 321–342, 2019b.
- Boone, A. and Etchevers, P.: An Intercomparison of Three Snow Schemes of Varying Complexity Coupled to the Same Land Surface Model: Local-Scale Evaluation at an Alpine Site, *Journal of Hydrometeorology*, 2, 374–394, [https://doi.org/10.1175/1525-7541\(2001\)002<0374:AIOTSS>2.0.CO;2](https://doi.org/10.1175/1525-7541(2001)002<0374:AIOTSS>2.0.CO;2), 2001.
- Bracegirdle, T. J. and Marshall, G. J.: The Reliability of Antarctic Tropospheric Pressure and Temperature in the Latest Global Reanalyses,  
20 *Journal of Climate*, 25, 7138–7146, <https://doi.org/10.1175/JCLI-D-11-00685.1>, 2012.
- Bracegirdle, T. J., Shuckburgh, E., Sallee, J.-B., Wang, Z., Meijers, A. J. S., Bruneau, N., Phillips, T., and Wilcox, L. J.: Assessment of surface winds over the Atlantic, Indian, and Pacific Ocean sectors of the Southern Ocean in CMIP5 models: historical bias, forcing response, and state dependence, *Journal of Geophysical Research: Atmospheres*, 118, 547–562, <https://doi.org/10.1002/jgrd.50153>, 2013.
- Bracegirdle, T. J., Stephenson, D. B., Turner, J., and Phillips, T.: The importance of sea ice area biases in 21st century multimodel projections  
25 of Antarctic temperature and precipitation, *Geophysical Research Letters*, 42, 10,832–10,839, <https://doi.org/10.1002/2015GL067055>, 2015.
- Bracegirdle, T. J., Hyder, P., and Holmes, C. R.: CMIP5 Diversity in Southern Westerly Jet Projections Related to Historical Sea Ice Area: Strong Link to Strengthening and Weak Link to Shift, *Journal of Climate*, 31, 195–211, <https://doi.org/10.1175/JCLI-D-17-0320.1>, 2018.
- Clem, K. R., Renwick, J. A., McGregor, J., and Fogt, R. L.: The relative influence of ENSO and SAM on Antarctic Peninsula climate, *Journal  
30 of Geophysical Research: Atmospheres*, 121, 9324–9341, 2016.
- Collins, M., Booth, B. B., Harris, G. R., Murphy, J. M., Sexton, D. M., and Webb, M. J.: Towards quantifying uncertainty in transient climate change, *Climate dynamics*, 27, 127–147, 2006.
- Dee, D. P., Uppala, S. M., Simmons, A., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M., Balsamo, G., Bauer, d. P., et al.: The ERA-Interim reanalysis: Configuration and performance of the data assimilation system, *Quarterly Journal of the royal meteorological  
35 society*, 137, 553–597, 2011.
- Déqué, M.: Frequency of precipitation and temperature extremes over France in an anthropogenic scenario: Model results and statistical correction according to observed values, *Global and Planetary Change*, 57, 16 – 26,

- <https://doi.org/https://doi.org/10.1016/j.gloplacha.2006.11.030>, <http://www.sciencedirect.com/science/article/pii/S0921818106002748>, extreme Climatic Events, 2007.
- Déqué, M., Dreveton, C., Braun, A., and Cariolle, D.: The ARPEGE/IFS atmosphere model: a contribution to the French community climate modelling, *Climate Dynamics*, 10, 249–266, <https://doi.org/10.1007/BF00208992>, 1994.
- 5 Dommenget, D. and Rezny, M.: A Caveat Note on Tuning in the Development of Coupled Climate Models, *Journal of Advances in Modeling Earth Systems*, 10, 78–97, <https://doi.org/https://doi.org/10.1002/2017MS000947>, 2018.
- Dutra, E., Sandu, I., Balsamo, G., Beljaars, A., Freville, H., Vignon, E., and Brun, E.: Understanding the ECMWF winter surface temperature biases over Antarctica, *European Centre for Medium-Range Weather Forecasts*, 2015.
- Eden, J. M., Widmann, M., Grawe, D., and Rast, S.: Skill, correction, and downscaling of GCM-simulated precipitation, *Journal of Climate*, 25, 3970–3984, 2012.
- 10 Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., and Taylor, K. E.: Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization, *Geoscientific Model Development (Online)*, 9, 2016.
- Favier, V., Agosta, C., Parouty, S., Durand, G., Delaygue, G., Gallee, H., Drouet, A., Trouvilliez, A., and Krinner, G.: An updated and quality controlled surface mass balance dataset for Antarctica, *The Cryosphere*, 7, p. 583 – p. 597, <https://doi.org/10.5194/tc-7-583-2013>, 2013.
- 15 Flato, G., Marotzke, J., Abiodun, B., Braconnot, P., Chou, S. C., Collins, W. J., Cox, P., Driouech, F., Emori, S., Eyring, V., et al.: Evaluation of Climate Models. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, *Climate Change 2013*, 5, 741–866, 2013.
- Fox-Rabinovitz, M., Côté, J., Dugas, B., Déqué, M., and McGregor, J. L.: Variable resolution general circulation models: Stretched-grid model intercomparison project (SGMIP), *Journal of Geophysical Research: Atmospheres*, 111, <https://doi.org/10.1029/2005JD006520>, 2006.
- 20 Fréville, H., Brun, E., Picard, G., Tatarinova, N., Arnaud, L., Lanconelli, C., Reijmer, C., and van den Broeke, M.: Using MODIS land surface temperatures and the Crocus snow model to understand the warm bias of ERA-Interim reanalyses at the surface in Antarctica, *The Cryosphere*, 8, 1361–1373, <https://doi.org/10.5194/tc-8-1361-2014>, <https://www.the-cryosphere.net/8/1361/2014/>, 2014.
- Frieler, K., Clark, P. U., He, F., Buizert, C., Reese, R., Ligtenberg, S. R. M., van den Broeke, M., Winkelmann, R., and Levermann, A.: Consistent evidence of increasing Antarctic accumulation with warming, *Nature Climate Change*, 5, <https://doi.org/10.1038/nclimate2574>, <http://dx.doi.org/10.1038/nclimate2574>, 2015.
- 25 Fürst, J. J., Durand, G., Gillet-Chaulet, F., Tavard, L., Rankl, M., Braun, M., and Gagliardini, O.: The safety band of Antarctic ice shelves, *Nature Climate Change*, 6, 479, 2016.
- Fyfe, J. C. and Saenko, O. A.: Simulated changes in the extratropical Southern Hemisphere winds and currents, *Geophysical Research Letters*, 33, 2006.
- 30 Gates, W. L.: AN AMS CONTINUING SERIES: GLOBAL CHANGE–AMIP: The Atmospheric Model Intercomparison Project, *Bulletin of the American Meteorological Society*, 73, 1962–1970, 1992.
- Gibelin, A.-L. and Déqué, M.: Anthropogenic climate change over the Mediterranean region simulated by a global variable resolution model, *Climate Dynamics*, 20, 327–339, 2003.
- 35 Gleckler, P. J., Taylor, K. E., and Doutriaux, C.: Performance metrics for climate models, *Journal of Geophysical Research: Atmospheres*, 113, <https://doi.org/10.1029/2007JD008972>, 2008.

- Grazioli, J., Madeleine, J.-B., Gallée, H., Forbes, R. M., Genthon, C., Krinner, G., and Berne, A.: Katabatic winds diminish precipitation contribution to the Antarctic ice mass balance, *Proceedings of the National Academy of Sciences*, 114, 10858–10863, <https://doi.org/10.1073/pnas.1707633114>, 2017.
- Guldborg, A., Kaas, E., Déqué, M., Yang, S., and Vester, T.: Reduction of systematic errors by empirical model correction: impact on seasonal prediction skill, *Tellus A*, 57, 575–588, <https://doi.org/10.1111/j.1600-0870.2005.00120.x>, 2005.
- Hall, A.: Projecting regional change, *Science*, 346, 1461–1462, 2014.
- Jeuken, A., Siegmund, P., Heijboer, L., Feichter, J., and Bengtsson, L.: On the potential of assimilating meteorological analyses in a global climate model for the purpose of model validation, *Journal of Geophysical Research: Atmospheres*, 101, 16 939–16 950, 1996.
- Jones, M. E., Bromwich, D. H., Nicolas, J. P., Carrasco, J., Plavcová, E., Zou, X., and Wang, S.-H.: Sixty Years of Widespread Warming in the Southern Middle and High Latitudes (1957–2016), *Journal of Climate*, 32, 6875 – 6898, <https://doi.org/10.1175/JCLI-D-18-0565.1>, 2019.
- Kharin, V. and Scinocca, J.: The impact of model fidelity on seasonal predictive skill, *Geophysical Research Letters*, 39, 2012.
- King, M. A. and Watson, C. S.: Antarctic Surface Mass Balance: Natural Variability, Noise, and Detecting New Trends, *Geophysical Research Letters*, 47, e2020GL087493, <https://doi.org/10.1029/2020GL087493>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2020GL087493>, 2020.
- Kohonen, T.: The self-organizing map, *Proceedings of the IEEE*, 78, 1464–1480, 1990.
- Kohonen, T.: Essentials of the self-organizing map, *Neural networks*, 37, 52–65, 2013.
- Krinner, G. and Flanner, M. G.: Striking stationarity of large-scale climate model bias patterns under strong climate change, *Proc. Natl. Acad. Sci.*, 15, 9462–9466, <https://doi.org/10.1073/pnas.1807912115>, 2018.
- Krinner, G., Genthon, C., Li, Z.-X., and Le Van, P.: Studies of the Antarctic climate with a stretched-grid general circulation model, *Journal of Geophysical Research: Atmospheres*, 102, 13 731–13 745, <https://doi.org/10.1029/96JD03356>, <http://dx.doi.org/10.1029/96JD03356>, 1997.
- Krinner, G., Guicherd, B., Ox, K., Genthon, C., and Magand, O.: Influence of Oceanic Boundary Conditions in Simulations of Antarctic Climate and Surface Mass Balance Change during the Coming Century, *Journal of Climate*, 21, 938–962, <https://doi.org/10.1175/2007JCLI1690.1>, 2008.
- Krinner, G., Rinke, A., Dethloff, K., and Gorodetskaya, I. V.: Impact of prescribed Arctic sea ice thickness in simulations of the present and future climate, *Climate Dynamics*, 35, 619–633, <https://doi.org/10.1007/s00382-009-0587-7>, 2010.
- Krinner, G., Largeron, C., Ménégot, M., Agosta, C., and Brutel-Vuilmet, C.: Oceanic Forcing of Antarctic Climate Change: A Study Using a Stretched-Grid Atmospheric General Circulation Model, *Journal of Climate*, 27, 5786–5800, <https://doi.org/10.1175/JCLI-D-13-00367.1>, 2014.
- Krinner, G., Beaumet, J., Favier, V., Déqué, M., and Brutel-Vuilmet, C.: Empirical Run-Time Bias Correction for Antarctic Regional Climate Projections With a Stretched-Grid AGCM, *Journal of Advances in Modeling Earth Systems*, 11, 64–82, <https://doi.org/10.1029/2018MS001438>, 2019.
- Krinner, G., Kharin, V., Roebrig, R., Scinocca, J., and Codron, F.: Historically-based run-time bias corrections substantially improve model projections of 100 years of future climate change, *Communications Earth Environment*, 1, <https://doi.org/10.1038/s43247-020-00035-0>, 2020.
- Kwok, R. and Comiso, J. C.: Spatial patterns of variability in Antarctic surface temperature: Connections to the Southern Hemisphere Annular Mode and the Southern Oscillation, *Geophysical Research Letters*, 29, 50–1, 2002.



- Lenaerts, J. T. M., Vizcaino, M., Fyke, J., van Kampenhout, L., and van den Broeke, M. R.: Present-day and future Antarctic ice sheet climate and surface mass balance in the Community Earth System Model, *Climate Dynamics*, 47, 1367–1381, <https://doi.org/10.1007/s00382-015-2907-4>, 2016.
- Li, G. and Xie, S.-P.: Origins of tropical-wide SST biases in CMIP multi-model ensembles, *Geophysical Research Letters*, 39, <https://doi.org/10.1029/2012GL053777>, 2012.
- Ligtenberg, S. R. M., van de Berg, W. J., van den Broeke, M. R., Rae, J. G. L., and van Meijgaard, E.: Future surface mass balance of the Antarctic ice sheet and its influence on sea level change, simulated by a regional atmospheric climate model, *Climate Dynamics*, 41, 867–884, <https://doi.org/10.1007/s00382-013-1749-1>, 2013.
- Mahlstein, I., Gent, P. R., and Solomon, S.: Historical Antarctic mean sea ice area, sea ice trends, and winds in CMIP5 simulations, *Journal of Geophysical Research: Atmospheres*, 118, 5105–5110, <https://doi.org/10.1002/jgrd.50443>, 2013.
- Manabe, S. and Stouffer, R.: Two stable equilibria of a coupled ocean-atmosphere model, *Journal of Climate*, 1, 841–866, 1988.
- Maraun, D. and Widmann, M.: Statistical downscaling and bias correction for climate research, Cambridge University Press, 2018.
- Maraun, D., Shepherd, T. G., Widmann, M., Zappa, G., Walton, D., Gutiérrez, J. M., Hagemann, S., Richter, I., Soares, P. M., Hall, A., et al.: Towards process-informed bias correction of climate change simulations, *Nature Climate Change*, 7, 764, 2017.
- Marshall, G. J.: Half-century seasonal relationships between the Southern Annular mode and Antarctic temperatures, *International Journal of Climatology*, 27, 373–383, <https://doi.org/10.1002/joc.1407>, 2007.
- Marshall, G. J. and Thompson, D. W. J.: The signatures of large-scale patterns of atmospheric variability in Antarctic surface temperatures, *Journal of Geophysical Research: Atmospheres*, 121, 3276–3289, <https://doi.org/10.1002/2015JD024665>, 2016.
- Marshall, G. J., Thompson, D. W., and van den Broeke, M. R.: The signature of Southern Hemisphere atmospheric circulation patterns in Antarctic precipitation, *Geophysical research letters*, 44, 2017.
- McGregor, J. L.: Recent developments in variable-resolution global climate modelling, *Climatic Change*, 129, 369–380, 2015.
- Medley, B. and Thomas, E.: Increased snowfall over the Antarctic Ice Sheet mitigated twentieth-century sea-level rise, *Nature Climate Change*, 9, 34–39, 2019.
- Mélia, D. S.: A global coupled sea ice-ocean model, *Ocean Modelling*, 4, 137 – 172, [https://doi.org/10.1016/S1463-5003\(01\)00015-4](https://doi.org/10.1016/S1463-5003(01)00015-4), <http://www.sciencedirect.com/science/article/pii/S1463500301000154>, 2002.
- Miller, R., Schmidt, G., and Shindell, D.: Forced annular variations in the 20th century intergovernmental panel on climate change fourth assessment report models, *Journal of Geophysical Research: Atmospheres*, 111, 2006.
- Moss, R. H., Edmonds, J. A., Hibbard, K. A., Manning, M. R., Rose, S. K., Van Vuuren, D. P., Carter, T. R., Emori, S., Kainuma, M., Kram, T., et al.: The next generation of scenarios for climate change research and assessment, *Nature*, 463, 747, 2010.
- Nicolas, J. P. and Bromwich, D. H.: New Reconstruction of Antarctic Near-Surface Temperatures: Multidecadal Trends and Reliability of Global Reanalyses, *Journal of Climate*, 27, 8070–8093, <https://doi.org/10.1175/JCLI-D-13-00733.1>, 2014.
- Noilhan, J. and Mahfouf, J.-F.: The ISBA land surface parameterisation scheme, *Global and Planetary Change*, 13, 145 – 159, [https://doi.org/10.1016/0921-8181\(95\)00043-7](https://doi.org/10.1016/0921-8181(95)00043-7), <http://www.sciencedirect.com/science/article/pii/0921818195000437>, soil Moisture Simulation, 1996.
- Palmer, C., Genthon, C., Claud, C., Kay, J. E., Wood, N. B., and L'Ecuyer, T.: Evaluation of current and projected Antarctic precipitation in CMIP5 models, *Climate Dynamics*, 48, 225–239, <https://doi.org/10.1007/s00382-016-3071-1>, <https://doi.org/10.1007/s00382-016-3071-1>, 2017.

- Pollard, D., DeConto, R. M., and Alley, R. B.: Potential Antarctic Ice Sheet retreat driven by hydrofracturing and ice cliff failure, *Earth and Planetary Science Letters*, 412, 112 – 121, <https://doi.org/10.1016/j.epsl.2014.12.035>, 2015.
- Pritchard, H., Ligtenberg, S., Fricker, H., Vaughan, D., Van den Broeke, M., and Padman, L.: Antarctic ice-sheet loss driven by basal melting of ice shelves, *Nature*, 484, 502, 2012.
- 5 Raphael, M., Marshall, G., Turner, J., Fogt, R., Schneider, D., Dixon, D., Hosking, J., Jones, J., and Hobbs, W.: The Amundsen sea low: variability, change, and impact on Antarctic climate, *Bulletin of the American Meteorological Society*, 97, 111–121, 2016.
- Reusch, D. B., Alley, R. B., and Hewitson, B. C.: North Atlantic climate variability from a self-organizing map perspective, *Journal of Geophysical Research: Atmospheres*, 112, <https://doi.org/10.1029/2006JD007460>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006JD007460>, 2007.
- 10 Rignot, E., Jacobs, S., Mouginot, J., and Scheuchl, B.: Ice-shelf melting around Antarctica, *Science*, 341, 266–270, 2013.
- Ritz, C., Tamsin, E. L., Durand, G., Payne, A. J., Peyaud, V., and Hindmarsh, R. C. A.: Potential sea-level rise from Antarctic ice-sheet instability constrained by observations, *Nature*, 528, 115, <https://doi.org/http://dx.doi.org/10.1038/nature16147>, 2015.
- Schneider, E. K.: Flux correction and the simulation of changing climate, in: *Annales Geophysicae*, vol. 14, pp. 336–341, Copernicus GmbH, 1996.
- 15 Scott, R. C., Nicolas, J. P., Bromwich, D. H., Norris, J. R., and Lubin, D.: Meteorological drivers and large-scale climate forcing of West Antarctic surface melt, *Journal of Climate*, 32, 665–684, 2019.
- Scott Yiu, Y. Y. and Maycock, A. C.: On the seasonality of the El Niño teleconnection to the Amundsen Sea region, *Journal of Climate*, 32, 4829–4845, 2019.
- Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., Krinner, G., et al.: Mass balance of the Antarctic Ice Sheet from 1992 to 2017, *Nature*, 556, pages219–222, 2018.
- Sheridan, S. and Lee, C.: The self-organizing map in synoptic climatological research, *Progress in Physical Geography*, 35, 109 – 119, 2011.
- Stocker, T. F., Dahe, Q., Plattner, G.-K., and Tignor, M.: IPCC Workshop on Regional Climate Projections and their Use in Impacts and Risk Analysis Studies, in: *Workshop Report*, vol. 15, p. 18, Citeseer, 2015.
- Taylor, K. E., Stouffer, R. J., and Meehl, G. A.: An overview of CMIP5 and the experiment design, *Bulletin of the American Meteorological Society*, 93, 485–498, 2012.
- 25 Thomas, E. R., van Wessel, J. M., Roberts, J., Isaksson, E., Schlosser, E., Fudge, T., Vallelonga, P., Medley, B., Bertler, N., van de Broeke, M., et al.: Review of regional Antarctic snow accumulation over the past 1000 years, *Climate of the Past Discussion*, <http://dx.doi.org/10.5194/cp-2017-18>. [www.clim-past-discuss.net/cp-2017-18](http://www.clim-past-discuss.net/cp-2017-18), 2017.
- Turner, J., Colwell, S. R., Marshall, G. J., Lachlan-Cope, T. A., Carleton, A. M., Jones, P. D., Lagun, V., Reid, P. A., and Iagovkina, S.: The SCAR READER Project: Toward a High-Quality Database of Mean Antarctic Meteorological Observations, *Journal of Climate*, 17, 2890–2898, [https://doi.org/10.1175/1520-0442\(2004\)017<2890:TSRPTA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<2890:TSRPTA>2.0.CO;2), [https://doi.org/10.1175/1520-0442\(2004\)017<2890:TSRPTA>2.0.CO;2](https://doi.org/10.1175/1520-0442(2004)017<2890:TSRPTA>2.0.CO;2), 2004.
- 30 Turner, J., Bracegirdle, T. J., Phillips, T., Marshall, G. J., and Hosking, J. S.: An Initial Assessment of Antarctic Sea Ice Extent in the CMIP5 Models, *Journal of Climate*, 26, 1473–1484, <https://doi.org/10.1175/JCLI-D-12-00068.1>, 2013.
- 35 Van Meijgaard, E., Van Uft, L., Van de Berg, W., Bosveld, F., Van den Hurk, B., Lenderink, G., and Siebesma, A.: The KNMI regional atmospheric climate model RACMO version 2.1, *Koninklijk Nederlands Meteorologisch Instituut*, 43, 2008.
- Van Wessel, J., Reijmer, C., Lenaerts, J., Van de Berg, W., Van den Broeke, M., and Van Meijgaard, E.: Updated cloud physics in a regional atmospheric climate model improves the modelled surface energy balance of Antarctica, *The Cryosphere*, 8, 125–135, 2014.

- van Wessem, J. M., van de Berg, W. J., Noël, B. P. Y., van Meijgaard, E., Amory, C., Birnbaum, G., Jakobs, C. L., Krüger, K., Lenaerts, J. T. M., Lhermitte, S., Ligtenberg, S. R. M., Medley, B., Reijmer, C. H., van Tricht, K., Trusel, L. D., van Uft, L. H., Wouters, B., Wuite, J., and van den Broeke, M. R.: Modelling the climate and surface mass balance of polar ice sheets using RACMO2 – Part 2: Antarctica (1979–2016), *The Cryosphere*, 12, 1479–1498, <https://doi.org/10.5194/tc-12-1479-2018>, 2018.
- 5 Vaughan, D. G., Bamber, J. L., Giovinetto, M., Russell, J., and Cooper, A. P. R.: Reassessment of Net Surface Mass Balance in Antarctica, *Journal of Climate*, 12, 933–946, [https://doi.org/10.1175/1520-0442\(1999\)012<0933:RONSMB>2.0.CO;2](https://doi.org/10.1175/1520-0442(1999)012<0933:RONSMB>2.0.CO;2), 1999.
- Velicogna, I.: Increasing rates of ice mass loss from the Greenland and Antarctic ice sheets revealed by GRACE, *Geophysical Research Letters*, 36, 2009.
- Verfaillie, D., Favier, V., Gallée, H., Fettweis, X., Agosta, C., and Jomelli, V.: Regional modeling of surface mass balance on the Cook Ice Cap, Kerguelen Islands (49°N, 69°E, *Climate Dynamics*, 53, 5909–5925, 2019.
- 10 Walden, V. P., Warren, S. G., and Tuttle, E.: Atmospheric ice crystals over the Antarctic Plateau in winter, *Journal of Applied Meteorology*, 42, 1391–1405, 2003.

### Large-scale atmospheric circulation

In this section, we present some of the results mentioned in the results or discussion section for large-scale atmospheric circulation in order to facilitate the comprehension of the discussion. In Fig. ??, we can see the large bias reduction with respect to ERA-I for 200 hPa temperatures in ARP-AMIP-AC. However, we can see the slight increase in spring and winter of the warm bias over the South Pole already present in spring for AMIP. In Fig. ??, we can see the remaining bias on 850 hPa temperatures in ARP-AMIP-AC with respect to ERA-I. The bias is close to zero in most places except for relatively small warm bias ( $\sim 1$ –2K) over mid-latitudes land masses (South America, South Africa and Australia). Wet or dry biases are found over the same places in 850 and 500 hPa specific humidity, but their sign vary depending on the season or the level considered (*figures not shown*). These biases were absent in ARP-AMIP (*figures not shown*) and probably results from errors on planetary boundary layer or clouds processes (i.e. convection). In Fig. ??, we can see that there is substantial warming in winter 500 hPa temperatures in ARP-AMIP-AC with respect to AMIP. This warming resulted in an increase downward longwave radiation over the East Antarctic Plateau and explains the increase of the winter warm bias in this area in near-surface temperatures in ARP-AMIP-AC.

25

ARP-AMIP – ERA-I ARP-AMIP-AC – ERA-I Seasonal ARP-AMIP (top) and ARP-AMIP-AC (bottom) bias on 200 hPa temperature (K) with respect to ERA-Interim over 1981–2010.

Seasonal ARP-AMIP-AC bias on 850 hPa temperatures with respect to ERA-I over 1981–2010.

ARP-AMIP-AC – ARP-AMIP winter (JJA) 500 hPa temperatures

30 Relative seasonal root mean square error reduction  $\Delta_r E$  (in %) south of 20°S with respect to ERA-Interim for ARP-AMIP-AC with respect to ARP-AMIP the 1981–1992 period for different surface and tropospheric variables at constant pressure levels ÷ Simulations JJA SON DJF MAM SLP 78 55 48 68

T500 91 93 94 89 Z500 87 83 72 81 Q500 3 1 77 1

## Near-Surface Temperatures

In this section, we present the difference with ERA-I driven MAR simulation in  $T_{2m}$  for the ARP-AMIP simulation (Fig. ??, already shown in Beaumet et al. (2019a)). The differences with *in-situ* stations from READER data base are presented in Tab ?? for ARP-AMIP-AC and ARP-AMIP Tab ??. Statistics are gathered per Antarctic regions.

- 5 ARP-AMIP minus MAR-ERA-I  $T_{2m}$  in winter (*left*) and summer (*right*). Circles represent differences with stations from the monthly READER data base. *Black contour line* represents where the difference is one standard deviation of MAR  $T_{2m}$ .
- Error on READER weather station  $T_{2m}$  in the ARP-AMIP-AC simulation for the reference period 1981-2010. Errors significant at  $p < 0.05$  are presented in **bold**. Stations DJF MAM JJA SON EAP Amundsen Scott 0.76 **3.15** 1.22 0.7 Vostok **-1.12 4.45 5.44 2.24** Mean error -0.18 3.80 3.33 1.47 RMSE 0.96 3.86 3.94 1.66 Coastal EA Casey **-0.94 -3.5 -3.92 -3.38** Davis **-1.28 -2.03 -1.56 -1.29** Dumont Durville **-0.53 -3.21 -3.56 -2.76** Mawson -0.28 **-2.62 -2.84 -2.53** McMurdo **-3.45 -2.33 -3.12 -3.38** Mirny **-1.57 -0.32 -0.01 -0.08** Novolazarevskaya 0.38 **-2.33 -1.78 -1.33** Scott Base **-1.36** 1 0.43 0.01 Syowa **-2.31** -0.53 -1.43 -0.75 Mean error -0.91 -1.76 -1.98 -1.70 RMSE 1.65 2.26 2.44 2.13 Ice shelves Halley **2.68 6.84 7.54 5.38** Neumayer **3.21 5.45 6.58 5.25** Mean error .95 6.15 7.06 5.32 RMSE 2.96 6.18 7.08 5.32 Peninsula Bellingshausen **-0.86** 0.3 0.11 0.08 Esperanza **-1.66** 1.32 -0.76 -0.9 Faraday **-1.79 -1.23 -2.24 -2.12** Marambio **-2.34** 1.6 -1 -1.62 Marsh **-0.64** 0.36 0.06 0.13
- 10 Oreadas **-0.92** 0.2 0.19 -0.64 Rothera **-2 -0.99 -3.14 -2.63** Mean error -1.46 0.22 -0.97 -1.1 RMSE 1.58 1.54 1.48 Southern Ocean Gough **-1.05** -0.3 0.12 **-0.71** Macquarie **-0.47** 0.09 **0.39** -0.25 Marion **-0.92 -0.43** 0.01 **-0.46** Mean error -0.81 -0.21 0.17 -0.47 RMSE 0.85 0.31 0.24 0.51

- Error on READER weather station  $T_{2m}$  in the ARP-AMIP simulation for the reference period 1981-2010. Errors significant at  $p = 0.05$  are presented in **bold**. Stations DJF MAM JJA SON EAP Amundsen Scott 0.47 **2.4** 1.06 0.94 Vostok **-1.46 3.21 3.22 1.89** Mean error -0.50 2.81 2.14 1.42 RMSE 1.08 2.83 2.40 1.49 Coastal EA Casey **-3.97 -5.72 -6.88 -5.41** Davis **-1.61 -4.19 -5.98 -3.31** Dumont Durville -0.45 **-2.82 -4.07 -2.24** Mawson **-2.24 -4.32 -5.67 4.26** McMurdo **-7.13 -6.48 -8.11 -8.38** Mirny **-1.24 -2.21 -2.97 -1.98** Novolazarevskaya **2.49** 0.58 -1.02 0.58 Scott Base **-5.03 -3.15 -4.56 -4.98** Syowa -0.17 -0.58 **-1.49** 0.04 Mean error -2.15 -3.34 -4.53 -3.33 RMSE 3.46 3.86 5.06 4.25 Ice shelves Halley **1.27 2.45** 1.21 0.88 Neumayer **2.18 1.21** 0.9 **1.41** Mean error 1.73 1.83 1.06 1.15 RMSE 1.78 1.93 1.07 1.18 Peninsula Bellingshausen **-1.02** -0.42 -0.24 -0.08 Esperanza **-1.1** 0.5 -1.33 -0.88 Faraday **-2.66 -4.66 -5.74 -3.66** Marambio **-1.87** 1.04 -1.27 -1.6 Marsh **-0.81** -0.36 -0.29 -0.03 Oreadas **-1.13** -0.04 0.61 -0.76 Rothera **-5.55 -7.88 -8.72 -6.13** Mean error -2.02 -1.69 -2.43 -1.88 RMSE 2.55 3.49 4.02 2.80 Southern Ocean Gough **-0.98** -0.34 0.02 **-0.79** Macquarie **-0.71** -0.35 0.2 **-0.45** Marion **-1.15 -0.43** -0.05 **-0.68** Mean error -0.95 -0.37 0.06 -0.64 RMSE 0.96 0.38 0.12 0.66

## Surface Mass Balance

- 30 In this section, we present the ARP-AMIP comparison with MAR-ERA-I for total precipitation, surface sublimation and surface mass balance (Fig. ??) already presented in Beaumet et al. (2019a). The similar comparison for ARP-AMIP-AC is presented in Section 3.1 (Fig. ??). Total Precipitations (*top*), surface snow sublimation (*centre*) and SMB (*bottom*) for ARP-AMIP minus

MAR-ERA-1 yearly cumul difference ( $\text{mmWe yr}^{-1}$ ) for the reference period 1981-2010. Pink (brown) and blue (green) contour lines represents areas where ARPEGE-MAR differences are larger than 20%.