# Response to reviewers' comments on the manuscript "Geothermal flux beneath the Antarctic Ice Sheet derived from measured temperature profiles in deep boreholes" submitted to The Cryosphere

5 First of all, we would like to thank the Editor Alex Robinson and both anonymous reviewers for fruitful comments and advices. We tried to consider all mentioned issues and, in order to address comments (only critical ones), you will find here our answers point-by-point. The comments are in **brown**, and our answers are in black. The list of all relevant changes made in the manuscript and the marked-up manuscript version are placed at the end of the response letter.

10

# Anonymous Referee #1

# 1. Heat flow model assumptions

The Antarctic ice sheet has an exceedingly long thermal memory and the slowest response time of the ice sheet is on a timescale exceeding 10 kyr (Ackert, 2003). The ice sheet is continuously in a

15 transient state responding to past changes as well as contemporary forcings. The ice sheet is in disequilibrium, therefore, the assumption that the system is in a thermodynamical steady state must be properly justified and quantified. Otherwise, how are the results of this study meant to be interpreted against the literature (e.g. Martos et al., 2017; Passalacqua et al., 2017).

Over the last several glacial cycles, ice thickness, surface temperatures, and accumulation rates

- 20 have varied across the ice core sites. Within the scope of a 1D time dependent heat flow model, these boundary conditions (BCs) directly impact the thermal profile of the ice. Many ice core records offer reconstructions of both temperature and accumulation rates through time. These could directly be applied as BCs into a time-dependent heat flow model rather then constant model parameters.
- 25 The structural uncertainty affiliated with the assumption of a steady state heat flow model should be quantified. Time-dependent transient experiments should be conducted with proper timedependent BCs wherever appropriate to assess the impact of a steady state assumption on the GHF results. Supplemented with a proper uncertainty analysis, this would contextualize the results with the literature.
- 30 Three drill sites (Dome C, Dome F, Vostok) are in close vicinity to ice divides where horizontal advection and horizontal heat conduction are assumed to be minimal and the environment approximates a steady state (Cuffey and Paterson, 2010). To first order, we also assume WAIS Divide is in a steady state. Byrd and Kohnen are in the interior slow-moving areas of the Antarctic Ice Sheet with a relatively smooth bed, where horizontal conduction is much lower than vertical
- 35 conduction (Hindmarsh, 1999, 2018), as well as horizontal advection and horizontal heat conduction can be neglected (Robin, 1955; Van Liefferinge et al., 2018). Thus, we assume that the temperature measured at the six drill sites is at thermal steady state in the near base portion.

We modified the design method and chose the form factor for concrete site according with the best fitting between modeled and measured depth-age scales. The best value for the form factor was

40 selected on the basis of the nonlinear correlation analysis. We think that the corrected design method gave more exact results. In addition, we added into manuscript section 2.3 "Uncertainties" and section 4.1 "Transient model vs. steady-state model".

# 2. Surface forcing of heat flow model

- 45 The heat flow model uses four model parameters: surface temperature, surface accumulation rate, basal melt, and basal temperature gradient. This seems to suggest that the surface temperature and accumulation rate are constant values and not time dependent. What are the resultant optimal GA temperature and accumulation forcings for each ice core site and how do they compare to present day observed values? There is a passing mention of the accumulation rate being time dependent in
- 50 Section 2.2 to calculate vertical velocities at each ice core site. What is this study using, constant surface accumulation rates (model parameter), time-dependent accumulation rates (vertical velocity inference), or both? How does the accumulation rate used in the vertical velocity calculations compare against the optimal rate inferred from the GA? The study should be consistently using time-dependent surface temperature and accumulation rates. No reference is
- 55 provided for the accumulation time-series mentioned at line 99, rendering this work not reproducible by other researchers.

In our calculations, we do not use time-dependent values of the surface temperature and accumulation rate. These parameters are changed within GA in a wide range to fit the measured temperature. Then "equivalent" vertical velocity, modern accumulation rate and temperature can be calculated from the GA results (please, see section 3.3 "Indirect results").

- 60 be calculated from the GA results (please, see section 3.3 "Indirect results"). In general, the computational details that need to be captured and shared for reproducible research include: (1) the data that were used in the analysis; (2) written statements in a programming language (i.e., the source code of the software used in the analysis or to generate data products); (3) numeric values of all configurable settings for software; (4) detailed specification of computational
- 65 environment including system software and hardware requirements, including the version number of each software used; and (5) computational workflow<sup>1</sup>. All these are extremely extensive. We tried to provide baseline that can guarantee reproducibility of our scientific findings and will be happy to provide other data (if considered necessary).
- 70 3. Understated uncertainties

The GHF results come with uncertainty estimates that only represent one source of uncertainty affiliated with the initial parameter choices going into the GA. This significant underrepresents the overall uncertainties in their GHF estimates, which compromises the interpretation of their results with respect to the literature. The study does not account for structural uncertainties associated

75 with their assumptions (steady state and no horizontal advection). Moreover, it is unclear if the ice thickness in the analysis is kept constant at present day values, this is not explicitly state. It appears

<sup>&</sup>lt;sup>1</sup> National Academies of Sciences, Engineering, and Medicine (2019). Reproducibility and replicability in science. Washington, DC: The National Academies Press.

the study uses constant ice thickness at each ice core site and does not attempt to estimate GHF uncertainties affiliated with this assumption. The heat flow model does not apply time-dependent surface temperature and accumulation rates, these time-series come with uncertainties which should also be propagated into the uncertainty model of the GHF estimations.

Furthermore, the uncertainty of the power law exponent (form factor) for the vertical velocity profile from Fischer et al. (2013) is not considered. The form factor could be anywhere from m = 0.5 to 1, with the former being favoured by Fischer et al. (2013). The study chooses m=1 without justifying that choice. The analysis should be conducted again using m=0.5 and 0.75 to quantify the

85 impact of the form factor on the GHF estimates. This would propagate parametric uncertainties of the vertical velocity parametrization to the GHF estimates.

The GA manages to identify parameter choices that produce a strong fit to the observed borehole temperatures. However, given the unquantified impact of model assumptions and model weaknesses, it is possible the model is overfitting the data. Therefore, the study would greatly

90 benefit from more robust confidence intervals that incorporate parametric uncertainties and structural errors in the assumptions made in the heat flow model. Upon achieving this, the study would be able to assess the robustness of the anomalous GHF values at Kohnen and WAIS Divide.

For uncertainty analysis, please, see section 2.3 in the revised manuscript below.

95 Minor comments:

In Figure 1. a GHF comparison is shown at each ice core site. A legend showing which reference is affiliated with which color would clean up the figure and caption. This would remove all the subscript a-e appended onto each GHF bar graph.

Figure 1 is corrected.

100

80

# Anonymous Referee #2

... there are crucial aspects that are unclear from the text such as, why the results are important, what is the new gained knowledge, how these results compared with other local GHF values obtained through modeling in the same drill sites by other authors?

- 105 The importance of these studies is emphasized at the beginning of "Introduction". Obtained GHF values are compared with five modellings (Shapiro and Ritzwoller, 2004; Fox Maule et al., 2005 Van Liefferinge and Pattyn, 2013; An et al., 2015; Martos et al., 2017) using bar graphs on the Fig. 1. Further comparison with this data and data from other references for specific sites is given in the section 3.2 "Data comparison and divergences".
- 110

The manuscript lacks of a proper discussion section. The manuscript should separate results from discussion and conclusions. Additionally, a more detailed discussion is necessary.

"Discussion" and "Conclusions" sections are added into revised version of the manuscript.

115 In addition, key components of the methods are not adequately described or are missing. In particular, uncertainties are not adequately addressed which makes it difficult to evaluate the results and conclusions of this study.

We added section 2.3 "Uncertainties".

120 Below are my comments, suggestions and concerns that I hope will be useful for the authors to improve the manuscript:

- I suggest to change the title as it is not accurately representing the content of the manuscript.

The title of the manuscript is changed to: "Geothermal heat flux from measured temperature profiles in deep ice boreholes in Antarctica".

125

- Regarding the discrepancy between the high values obtained in Kohnen and WAIS Divide in comparison with Antarctic-wide maps:

One thing to consider is that the Antarctic-wide geothermal heat flow maps are representing the heat flow of a region, while a heat flow value derived using borehole measurements is representing

- 130 a specific local value. Therefore, probably these higher than predicted heat flow values obtained for Kohnen and WAIS Divide are only representing local values, not necessarily hot spots. The higher values could be consequence of, for example, a higher concentration of a particular radiogenic material in that spot, or a consequence of some particularity of the subglacial topography or the parameters and assumptions that are involved in the solutions of the model to obtain the local value.
- 135 For these reasons, understanding the uncertainty sources and quantifying them is extremely important and it is necessary.

We recalculated GHF in Kohnen and WAIS Divide according to the best value for the form factor that was selected on the basis of the nonlinear correlation analysis between modeled and measured age scales.

140

- L69: The manuscript should demonstrate the temperature measurement precision in a robust and scientific way

We added section 2.3.1 "Temperature measurements" with explanations of temperature measurement precision.

145

- L78-80: Where is this shown? Quantify the good agreement. This is important for the uncertainties of the estimated local geothermal heat flow

This is shown in Table 1 (the line "Ice thickness according with radar/seismic survey (m)") and Table 2 (the line "Ice thickness according with depth of pressure melting point (m)").

- Figure 1: The drill sites as well as other local values are plotted in this figure together with a geological map for the Antarctic continent. However, the geology is not mentioned in the text, there is no discussion about results and the subglacial geology. What is the purpose of the geological map if it is not used in the manuscript? I recommend to either include some discussion about it or select another background data to plot the drill sites and discuss the results in that context.

- At the first stage of the paper writing, we planned to connect revealed GHF values with Antarctic subglacial geology but then, because of the insufficiency of data, we dropped this idea. We redrew this figure and added location of the Antarctic ice divides.
- 160 Regarding uncertainties I have two main comments/concerns:

1. How uncertainties are calculated is not adequately explained and more information and details are needed to evaluate the GHF estimates.

2. A substantial discussion about which parameters are contributing to the uncertainty is necessary. In addition, there are assumptions made in the thermodynamic model and also parameters that are

165 assumed to be constant. These assumptions also carry uncertainties and they need to be properly quantified and included in the final uncertainty budget. For example, one important aspect to quantify would be the contribution to the uncertainty budget of considering steady-state condition. We added uncertainty considerations into revised version of the paper.

# 170

155

# List of all relevant changes made in the manuscript

1. Title of the manuscript has been specified in a narrower sense.

2. The method of GHF estimation is modified: now we use the published depth-age scales at the studied sites and estimated the best value for the form factor.

175 3. We added tables with polynomial approximations of borehole temperature as a function of true vertical depth and "equivalent" thermophysical parameters.

4. Figure 1 is modified and location of the Antarctic ice divides is added.

5. Figure with comparison of the measured and modeled age scales with different form factors has been added.

180 6. The section with uncertancities evaluation has been added.

7. "Discussion" and "Conclusions" sections have been added.

On behalf of co-authors:

Pavel Talalay

185 Yazhou Li

# Geothermal <u>heat</u> flux from measured temperature profiles in deep <u>ice</u> boreholes <u>in Antarctica</u>

Pavel Talalay<sup>1</sup>, Yazhou Li<sup>1</sup>, Laurent Augustin<sup>2</sup>, Gary D. Clow<sup>3</sup>, Jialin Hong<sup>1</sup>, Eric Lefebvre<sup>4</sup>, Alexey Markov<sup>1</sup>, Hideaki Matoyama<sup>5</sup>, Catherine Ritz<sup>4</sup>

<sup>1</sup>Polar Research Center, Institute for Polar Science and Engineering, Jilin University, 130021 Changchun, China
 <sup>2</sup>Division Technique de l'INSU, CNRS, 83507 La Seyne sur Mer, France
 <sup>3</sup>Institute of Arctic and Alpine Research, University of Colorado Boulder, Boulder, Colorado, USA
 <sup>4</sup>Université Grenoble Alpes, CNRS, IRD, IGE, 38000 Grenoble, France
 <sup>5</sup>National Institute of Polar Research, Tokyo, Japan

195 Correspondence to: Pavel Talalay (ptalalay@yahoo.com) and Yazhou Li (jluyazhouli@163.com)

Abstract. The temperature at the Antarctic ice sheet bed and the temperature gradient in subglacial rocks have been directly measured only a few times, although extensive thermodynamic modeling has been used to estimate the geothermal heat flux (GHF) under the ice sheet. During the last five decades, deep ice-core drilling projects at six sites – Byrd, WAIS Divide, Dome C, Kohnen, Dome F, and Vostok – have succeeded in reaching to, or nearly to, the bed at inland locations in Antarctica. When temperature profiles in these boreholes and steady-state heat flow modeling are combined with estimates of vertical velocity, the heat flow at the ice-sheet base is translated to a geothermal heat flux of 57.9±6.4 mW m<sup>-2</sup> at Dome C, 78.9±5.0 mW m<sup>-2</sup> at Dome F, and 86.9±16.6 mW m<sup>-2</sup> at Kohnen, all higher than the predicted values at these sites. This warm base under the East Antarctic Ice Sheet (EAIS) could be caused by radiogenic heat effects or hydrothermal circulation not accounted for by the models. The GHF at the base of the ice sheet base. Correlation analyses between modeled and measured depth-age scales at the EAIS sites indicate that all of them can be adequately approximated by a steady-state model. Horizontal velocities and their variation over ice-age cycles are much greater for the West Antarctic Ice Sheet than for the interior EAIS sites; a steady-state model cannot precisely describe the temperature distribution here. Even the correlation factors for the best fitting age-depth

210 mW m<sup>-2</sup> at WAIS Divide as references before more precise estimates are made on the subject.

### **1** Introduction

The Antarctic geothermal heat flux (GHF), an important boundary condition for ice sheet behavior, is associated with sea level changes (Golledge et al., 2015) considering its significant influence on the viscosity of basal ice and meltwater content at the ice-base interface. What are the basal ice temperature and mechanical properties? How does GHF control basal melt and affect

curve are only moderate for the West Antarctic sites, we can use the GHF values of 88.4±7.6 mW m<sup>-2</sup> at Byrd and 113.3±16.9

215 the internal deformation of the ice sheet? How old is ice at different locations? These questions can be answered only by applying reliable GHF measurements or estimates.

The average global surface GHF is  $\sim$ 86 mW m<sup>-2</sup>, which varies from 64.7 mW m<sup>-2</sup>, the mean continental heat flow (including arcs and continental margins), to 95.9 mW m<sup>-2</sup>, the mean oceanic heat flow (Davies, 2013). However, several geologic factors including heat from the mantle, heat production in the crust by radioactive decay, and tectonic history, cause spatially variable

220 GHF in Antarctica (Burton-Johnson et al., 2020).

- Most studies of GHF in Antarctica rely on thermal models (Pattyn, 2010; Van Liefferinge and Pattyn, 2013). Modeling studies based on a global seismic survey and the structural similarity of crust and upper mantle showed that the West Antarctic Ice Sheet (WAIS) has a GHF three times higher than that of the East Antarctic ice sheet (EAIS) (Shapiro and Ritzwoller, 2004). For a central point in the WAIS (78S, 110W), the average GHF is expected to be 110 mW m<sup>-2</sup>. The GHF can also be estimated
- on the basis of geologic information, where uniform values are attributed to large geologically homogeneous areas (An et al., 2015; Goodge, 2018; Llubes et al., 2006; Martos et al., 2015; Pollard et al., 2005).
  - Some studies use remote methods to estimate <u>the</u> GHF underneath the Antarctic ice sheet. For example, satellite magnetic data showed that the GHF underneath the ice sheet varies from 40 to 185 mW m<sup>-2</sup> and that areas of high GHF coincide with known current volcanism and some areas known to have ice streams (Fox Maule et al., 2005). In the central part of the EAIS, the
- 230 average GHF was estimated to be in the range of 50 to 60 mW m<sup>-2</sup>; however, elevated GHFs were found along the WAIS– EAIS boundary and around the Siple Coast. Similarly, high GHFs were found around Victoria Land, Oates Land, and George V Land. Observations of crustal heat production within the continental crust underneath the Lambert-Amery glacial system in East Antarctica also show high heat flux of at least 120 mW m<sup>-2</sup> (Pittard et al., 2016).

Direct temperature measurement obviously produces the most reliable GHF estimates and can be used to verify results of

- preliminary thermal modeling and geological-geophysical studies. While over 10,000 heat flow measurements have been made globally, 90% are from Europe, North America, and southern Africa. South America, Asia, and Australia have far fewer measurements, <u>while</u> Antarctica has virtually none (<u>Davies, 2013</u>). Drilling through thick ice is extremely complicated, time-consuming, and expensive; therefore, direct temperature measurements in Antarctic subglacial till/bedrock environments have only been conducted twice so far, both under the WAIS: at the subglacial Lake Whillans (285±80 mW m<sup>-2</sup>) (Fisher et al., 2015)
- 240 and near the grounding zone of the Whillans Ice Stream (88±7 mW m<sup>-2</sup>) (Begeman et al., 2017), ~100 km apart (Fig. 1). The tremendous difference in the values of GHF between these two adjacent sites suggests\_high spatial variability\_in West Antarctica.

More reliable GHF estimates under the Antarctic ice sheet can be made from available temperature profiles in ice boreholes. During the last five decades, deep ice-core drilling projects at six sites – Byrd (Ueda and Garfield, 1970), WAIS Divide

(Slawny et al., 2014), Dome C (Augustin et al., 2007), Kohnen (Wilhelms et al., 2014), Dome F (Motoyama, 2007), and Vostok (Lukin and Vasiliev, 2014; Vasiliev et al., 2011) – have succeeded in reaching, or nearly reaching, the ice sheet bed at inland locations in Antarctica. Reported drill site conditions – snow accumulation rate, mean annual surface air temperature, <u>ice sheet surface velocity</u>, ice thickness, and drilling depth – are summarized in Table 1.

The Byrd and Kohnen holes encountered water at the base, which welled up into the holes. The borehole at Vostok penetrated the subglacial Lake Vostok at 3769.3 m, and here, water rose from the lake to a height of more than 340 m. Drilling of the other holes was stopped within 10–50 m of the bed. All these holes were temperature logged and provide a good opportunity to fill the gap in our knowledge of the GHF under the Antarctic ice.

#### 2 Methods

#### 2.1 Temperature and temperature gradient at the base of Antarctic Ice Sheet

- 255 Temperatures in the Byrd, WAIS Divide, Vostok, Dome C, Kohnen, and Dome F boreholes were measured using different devices and different methods. All boreholes were mechanically drilled and filled with kerosene-based drilling fluid. Temperature profiles were then obtained by logging with custom-made borehole loggers (Dome C, Kohnen, Dome F, WAIS Divide, and Vostok) or a thermistor embedded in the drill (Byrd).
- Measured temperature profiles in four of the boreholes (Vostok, Dome C, Kohnen, and Dome F) increase <u>almost</u> linearly with depth, as expected <u>at</u> locations with minimal <u>annual snow</u> accumulation and hence small vertical velocities (Fig. 2). <u>In contrast</u>, vertical advection is much greater at the Byrd and WAIS Divide <u>sites</u> in West Antarctica; <u>at</u> these locations the upper part of the ice sheet is nearly isothermal, but at depth the temperature gradient is nearly the same as <u>that</u> at the other sites. Temperature gradients at the bed are 2.02–3.12 °C/100 m at Dome C, Kohnen, Dome F and Vostok and slightly higher in West Antarctica, 3.70–3.88 °C/100 m at Byrd and WAIS Divide (Table 2).
- 265 <u>Temperature profiles in deep ice-core drilling boreholes are approximated closely by polynomials with correlation factors of >0.99 (Table 3), indicating a positive relationship between temperature and vertical depth. Ice thicknesses generated by extrapolating the temperature profile to the depth of the pressure melting point assuming a Clausius-Clapeyron slope of 0.0742 K/MPa (Cuffey and Paterson, 2010) are in good agreement with radar data, except for WAIS Divide where the difference is ~30 m. This could be attributed to scintillations on the melted ice-bedrock interface or other effects. However, in-depth temperature extrapolation has limited accuracy and thus often does not provide a correct estimate of GHF. Thus, a steady-state model and genetic algorithm (GA) are applied herein to fit the measured temperatures.</u>

#### 2.2 GHF estimation model

275

A one-dimensional time-dependent energy-balance equation (Dahl-Jensen et al., 2003; Johnsen et al., 1995) is usually used to model the temperature distribution through the ice as a function of the climate conditions on the surface and the GHF from the bedrock:

$$\rho c \frac{\partial T}{\partial t} = \frac{\partial}{\partial z} \left( k \frac{\partial T}{\partial z} \right) - \rho c w \frac{\partial T}{\partial z} - \rho c u \frac{\partial T}{\partial x},\tag{1}$$

where *T* is the temperature as a function of *z*, *x* and *t*; *z* represents the vertical coordinate (at the ice sheet base,  $z_{=0}$ , while at ice sheet surface,  $z_{=}H$ ); *H* is the ice thickness and is assumed to be constant in time; *x* is the horizontal coordinate; *t* is the time; *k* is the <u>thermal</u> conductivity of ice dependent on *T*;  $\rho$  is the density of ice; *c* is the specific heat capacity of ice dependent on *T*; *w* and *u* are respectively, the vertical velocity and horizontal velocity of the ice sheet dependent on *z* and *t*.

The temperature measured at the six drill sites can be considered at thermal steady state in their near base portion. Three drill sites (Dome C, Dome F, Vostok) are in close vicinity to ice divides where horizontal advection and horizontal heat conduction are assumed to be minimal and the environment approximates a steady state (Cuffey and Paterson, 2010). To first order, we also assume WAIS Divide is in a steady state. Byrd and Kohnen are in the interior slow-moving areas of the Antarctic Ice

285 Sheet with a relatively smooth bed, where horizontal conduction is much lower than vertical conduction (Hindmarsh, 1999, 2018), as well as horizontal advection and horizontal heat conduction can be neglected (Robin, 1955; Van Liefferinge et al., 2018). This assumption reduces the non-steady-state heat-transfer equation to

$$\frac{\partial}{\partial z} \left( k \frac{\partial T}{\partial z} \right) - \rho c w \frac{\partial T}{\partial z} = 0, \tag{2}$$

which can be rewritten as

$$290 \quad \frac{1}{k}\frac{\partial k}{\partial z}\frac{\partial T}{\partial z} + \frac{\partial^2 T}{\partial z^2} - \frac{\rho c}{k}w\frac{\partial T}{\partial z} = 0.$$
(3)

Using  $\frac{\partial k}{\partial z} = \frac{\partial k}{\partial T} \frac{\partial T}{\partial z}$ , Eq. (3) becomes

$$\frac{\partial^2 T}{\partial z^2} + \left(\frac{1}{k}\frac{\partial k}{\partial T}\frac{\partial T}{\partial z} - \frac{\rho c}{k}w\right)\frac{\partial T}{\partial z} = 0.$$
(4)

We approximate the vertical velocity in the ice by

$$w(z) = -w_{melt} - (Acc - w_{melt}) \left(\frac{z}{H}\right)^{m+1},$$
(5)

where  $w_{melt}$  is the basal melt rate; *Acc* is the surface accumulation rate dependent on *t*; and *m* is the form factor that accounts for the variation of horizontal velocity.

Substitution of Eq. (5) into Eq. (4) and integrating on the assumption that k is constant gives the following temperature distribution in ice sheet at steady state:

$$T = T_s - \left[\frac{\partial T}{\partial z}\right]_B \int_0^z exp\left(\frac{(w_{melt} - Acc)z^{m+2}}{\alpha_T(m+2)H^{m+1}} - \frac{w_{melt}}{\alpha_T}z\right) dz + \left[\frac{\partial T}{\partial z}\right]_B \int_0^H exp\left(\frac{(w_{melt} - Acc)z^{m+2}}{\alpha_T(m+2)H^{m+1}} - \frac{w_{melt}}{\alpha_T}z\right) dz,\tag{6}$$

300 where  $T_s$  is the surface temperature;  $\left[\frac{\partial T}{\partial z}\right]_B$  is the temperature gradient at the ice sheet base;  $\alpha_T = k/\rho c$  is the thermal diffusivity of ice.

<u>The least squares</u> method was used to fit measured borehole temperatures with this equation. In fitting, the initial values of the unknown parameters  $T_s$ ,  $\begin{bmatrix} \frac{\partial T}{\partial z} \end{bmatrix}_B$ , Acc and  $w_{melt}$  can only be guessed and this results in unavoidable uncertainty of fitting. To overcome this large uncertainty, a common genetic algorithm was designed to find the optimal global solution of temperature

305 fitting by <u>constraining</u> these unknown parameters to a predetermined range (Reeves and Rowe, 2002). <u>GA can solve</u> optimization problems by limiting the unknown parameters to a predetermined range with any type of constraints, including integer constraints. In general, GA generates high-quality solutions for optimization problems and search problems.

In the GA, the crossover fraction is set to be 0.9 while the migration fraction is 0.2 (Reeves and Rowe, 2002). To obtain an accurate solution and save calculation time, we set the population size to be 8000 and the <u>number of generations</u> to be 20. 310 Usually, after 15 generations of iteration, the optimal solution can be found. All the calculations were performed <u>using</u>

MATLAB software. <u>GA provides results for the first generation of optimal solution in a wide range based on a random</u> combination of the fitting parameters. Thus, for each deep borehole, the fitting experiments were trialed five times to avoid random error of the GA caused by the initial random parameter combination. Then, the average value from the five fitting experiments was used as the GHF from bedrock into <u>the</u> ice <u>sheet</u> at <u>the</u> selected site.

#### B15 Equation (4) can also be <u>re-</u>expressed as <u>follows:</u>

$$w(z) = \left[\alpha_T \left(\frac{\partial^2 T}{\partial z^2}\right) - \alpha_T \left(\frac{1}{k}\frac{\partial k}{\partial T}\right) \left(\frac{\partial T}{\partial z}\right)^2\right] / \frac{\partial T}{\partial z}.$$
(7)

Note that the vertical velocity is markedly affected by  $\frac{\partial^2 T}{\partial z^2}(z)$  and  $\frac{\partial T}{\partial z}(z)$ . At the base of the ice sheet, the melt/freezing rate is  $w_{melt} = w(0)$  while the gradient is  $\left[\frac{\partial T}{\partial z}\right]_B = \frac{\partial T}{\partial z}(0)$ . The geothermal heat flux  $Q_{geo}$  from below the ice is balanced by the conductive flux in the ice  $q = k \left[\frac{\partial T}{\partial z}\right]_B$  and the rate at which energy is used to melt/freeze ice,  $J = \rho L w_{melt}$ . Thus, the GHF will be:

$$Q_{geo} = \rho L w_{melt} - k \left[ \frac{\partial T}{\partial z} \right]_{B}, \tag{8}$$

where L is the specific latent heat for melting of ice.

#### **2.3 Uncertainties**

320

In our method, the temperature in the lower portion of the ice sheets is assumed to be in steady state and the GA algorithm is used to fit the measured temperatures in deep ice-core drilling boreholes by varying the four key parameters influencing the temperature distribution: the surface temperature, surface accumulation rate, basal melt, and basal temperature gradient. All these parameters are suggested by algorithm in order to obtain the best-fitting curve. We assume that the main uncertainties in our fitting model are coming from temperature measurements, variability of the form factor *m*, ice thickness estimation and GA algorithm itself. It must be noted that the uncertainties we state are lower limits. There are some additional unexamined

330 <u>uncertainties that were missing from our model including transient effects associated with climate change and ice-sheet dynamics, the horizontal velocity field, the form of the vertical velocity field, the temperature dependence of the thermal conductivity, unaccounted for thermal disturbances due to drilling processes, 2D effects, and some other phenomena.</u>

#### 2.3.1 Temperature measurements

Interpretation of temperature measurements in mechanically drilled deep boreholes filled with drilling fluids is complicated by several factors (Clow, 2008). First, the temperature is measured in the borehole fluid, not in the surrounding ice; therefore, an important consideration is the need for thermal equilibration of the ice wall and the borehole fluids following drilling and prior to measurement. Second, the heat produced during drilling needs to be dissipated from the borehole or the thermal drilling disturbance needs to be accounted for (Clow, 2015). Third, increasing temperature with depth can cause convective mixing in the borehole. Fourth, the depth of temperature measurements has an inherent uncertainty due to cable slippage in the counting

- 340 assembly and cable elongation. Thus, all successful temperature measurements in deep boreholes obey a logging protocol in terms of logger tripping speed, measurement direction, borehole settling time, and so on to minimize the effects of these complicating factors. Temperature measurement errors from sensor accuracy and calibration are found to be within the tolerance for large-scale GHF estimates for our six boreholes to interpret ice-sheet basal dynamics. The temperature in the Byrd borehole was measured with an accuracy of 0.1 °C (H. Ueda, personal communication).
- Motoyama et al. (2013) reported that temperature measurements at Dome F were carried out with a precision of 0.05 °C. The absolute temperature measurement error at Vostok was estimated to be 0.07 °C (Salamatin et al., 1998a). The resolution of the temperature measurements in the Dome C borehole was 0.015 °C, while the precision was found to be 0.05 °C (Leferbre et al., 2002). The logger used in the borehole at Kohnen was calibrated to 0.03 °C (Gundestrup et al., 1994). Prior to drilling, a detailed study of the expected temperature-measurement uncertainties was made for the WAIS Divide site to optimize the
- B50 logging system setup (Clow, 2008). The standard uncertainty (accuracy) of the subsequent WAIS Divide temperature logs was ~0.0053 °C (Cuffey et al., 2016). In general, temperature measurement accuracy in the studied boreholes is more than adequate, and the measured drilling depth was recalculated to true vertical depth using available borehole inclinations.

#### 2.3.2 Form factor m

Selection of the appropriate form factor *m* is a challenging task. Classically, vertical velocity depends linearly on *z*/*H* (Cuffey

- and Paterson, 2010) and m = 0. However, at an ice divide, the downward flow of ice is slower, for the same depth, than at locations away from the divide (Raymond, 1983). This reduces the cooling influence of vertical advection and increases the basal temperature. Therefore, Raymond (1983) suggested the use of m = 1.0 for deformation in the vicinity of ice divides. To set up the vertical velocity profile at Dome C, Fischer et al. (2013) performed three runs with m = 0.3, m = 0.5 and m = 0.7and found that the temperature profile is only slightly affected by this choice. However, the form factor m exhibited a strong
- influence on the age profile of the ice. That was the reason why the authors used *m* = 0.5, which is in good agreement with the EDC3 age scale (Parrenin et al., 2007b). Following the method of Fischer et al., (2013), we use the published depth-age scales at the studied sites and estimated the best value for the form factor *m*. In order reduce the run time of multilevel calculations, we examine the form factor *m* at only five levels 0, 0.25, 0.50, 0.75, and 1.00. The best value for the form factor *m* is selected on the basis of the nonlinear correlation analysis between modeled and measured age scales. To calculate the correlation factor *R*<sup>2</sup> we first found the average value of the measured age;

B65 
$$\frac{R^2}{n}$$
, we first found the average value of the measured age:  
 $\bar{A} = \frac{1}{n} \sum_{i=1}^{n} A_{i,i}$ 
(9)  
where *n* is the number of measured ice ages  $A_i$ . Then the total sum of squares  $SS_{tot}$  and the sum of squares of residuals  $SS_{res}$ 

 $\frac{\text{were calculated:}}{SS_{tot} = \sum_{i} (A_i - \bar{A})^2},$   $370 \quad SS_{res} = \sum_{i} (A_i - \hat{A}_i)^2,$ 

where  $\hat{A}_i$  is the modeled ice age. The correlation factor  $R^2$  was estimated by

(10)

(11)

$$R^2 = 1 - \frac{SS_{res}}{SS_{tot}}.$$
(12)

Finally, the results of the nonlinear correlation analysis were checked by evaluating the root mean squared error (RMSE):

$$RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^{n} (A_i - \hat{A}_i)^2}.$$
(13)

#### 375 <u>2.3.3 Ice thickness</u>

We assume that the ice sheet thickness at the studied sites has kept constant at the present day height; however, it has varied in the past. The 3D thermo-mechanical model and the simple 1D model showed that the maximum variation of ice sheet thickness at Dome C and Dome F was less than 250 m in the past (Parrenin et al., 2007a). In general, the typical difference in the ice thickness in the glacial and interglacial periods at Dome C was 150 m (Passalacqua et al., 2017). At the Kohnen site,

- 380 the local elevation variation is on the order of 100 m (Huybrechts et al., 2007). The ice thickness variation at Vostok, located in the central part of East Antarctica plateau, exhibits the similar range as at Dome F and Dome C (Ritz et al., 2001). The best evidence for ice-sheet elevation change in the interior of the West Antarctic ice sheet comes from the Ohio Range, to the south of the WAIS Divide site at a height of 1600 m a.s.l., and from Mt. Waesche to the north of the WAIS Divide site at a height of 2000 m a.s.l. (Ackert et al., 1999, 2007). Moraines at Mt. Waesche were ~50 m higher and trimlines in the Ohio
- 385 Range were ~125 m higher, between 12 and 10 ka. The thinning of ~100 m throughout the Holocene occurred as the grounding line retreated by hundreds of km and the accumulation rates were relatively stable (Anderson et al., 2002; Conway et al., 1999). Cuffey et al. (2016) presented a model which indicates a more likely scenario of 200 m thickening at WAIS Divide when the accumulation rate rose after the last glacial maximum, followed by 300 m of thinning to the mid-Holocene. The elevation change is comparable to the amount of elevation change inferred for interior East Antarctic sites.
- 390 Comparison with the modern ice thickness value indicates that the variation of ice thickness is small and its influence on ice temperature distribution can be neglected, in particular, on lower portion of the ice borehole. For example, assuming a 150 m thickness increase from the LGM to 15 ka leads to the change in the reconstructed LGM temperature by less than 0.2 °C compared to a constant thickness in WAIS ice core (Buizert et al., 2015). This is the reason why constant ice thickness is also used by other researchers for GHF estimates (Dahl-Jensen et al., 2003; Engelhardt, 2004; Mony et al., 2020).

## 395 <u>2.3.4 Genetic algorithm</u>

For each deep borehole, the fitting experiments were repeated five times for the best value of the form factor m and the average value obtained from the five fitting experiments was used as the representative value of GHF from bedrock into ice. Thus, the uncertainty ranges came from the difference between the maximum/minimum and the average GHF values.

#### **3** Results

## 400 <u>3.1 Initial conditions and GHF estimates</u>

<u>GHF estimates</u> were made <u>using</u> the following ice parameters:

- density 918 kg m<sup>-3</sup>;
- specific heat capacity  $c = 152.5 + 7.122 (T + 273.15) \text{ J kg}^{-1} \text{ K}^{-1}$  (Yen, 1981);
- thermal conductivity  $k = 9.828e^{-0.0057(T+273.15)}$  W m<sup>-1</sup> K<sup>-1</sup> (Yen, 1981);
- specific latent heat  $L = 333.5 \text{ kJ kg}^{-1}$  (Cuffey and Paterson, 2010).

In our model, we assume that k and c are constant and equal to <u>their</u> values at the temperature of <u>the</u> pressure melting point; th<u>is can provide a</u> better estimate of <u>the</u> basal melting rate at the base of ice sheet (Fischer et al., 2013). In this case,  $\frac{1}{k} \frac{\partial k}{\partial T} = -5.7e-3 \text{ K}^{-1}$ . Fig<u>ure 3</u> shows the fitted temperature profiles compared with measured temperatures.

- We performed five runs for estimating GHF with m = 0, m = 0.25, m = 0.5, m = 0.75, and m = 1.0 for each site and compared modeled and measured age scales. As illustrated in Fig. 4, the form factor has a strong influence on the age profile of the ice.
- The results of our estimates for GHF for different *m* values are summarized in Table 4. Surprisingly, on three separate occasions the correlation factor is negative. This may occur when  $SS_{res}$  is far beyond that of  $SS_{tot}$  (see Eqs. (10)-(12)). That is to say that there is no correlation between modeled and measured age scales in these cases. The results of the correlation analyses were confirmed by evaluating how close the modeled lines are to the data points with the aid of the RMSE. The smaller the RMSE,
- 415 the closer the model is to the data. The GHF and *m* values with the highest correlation factor and smallest RSME were selected for further processing and trialed five times. The average GHF values for selected *m* are added into Table 2. The precision of the GHF estimates and basal melt/freezing rate are also specified here.

The temperature profiles show that the heat flow through the ice at six deep drilling sites in Antarctica must be >42.6– $\frac{77.1}{1000}$  mW m<sup>-2</sup> in order to match the observed temperatures in the boreholes. The basal ice at all sites is at the pressure-melting point,

420 and the amount of melt cannot be constrained by the energy-balance equation alone. When the heat flow model is combined with vertical velocity estimates, the estimated heat flow can be translated to a GHF of 57.9–113.3 mW m<sup>-2</sup>, except for Vostok with GHF of -3.3 mW m<sup>-2</sup>.

## 3.2 Data comparison and divergences

*Vostok*. The surface temperature-time curve for the upper bound of the present-day accumulation rate at Vostok corresponds to a GHF of 53 mW m<sup>-2</sup> (Salamatin et al., 1998b). We calculated that at the base of the ice sheet, the conductive flux is  $42.6\pm0.4$  mW m<sup>-2</sup> while the latent heat flux from refreezing of the lake water is  $46.3\pm5.6$  mW m<sup>-2</sup>. Thus, the GHF heat flux at the base of the ice sheet has a negative value of  $-3.6\pm5.3$  mW m<sup>-2</sup>. This is in good agreement with the isotope studies that showed that the Vostok ice core consists of ice refrozen from Lake Vostok water, from 3539 m below the surface of the Antarctic ice sheet to its bottom (Jouzel et al., 1999). Sufficiently high correlation factor (0.75) between modeled and measured age scales at *m* = 430 <u>1 indicates that ice above Lake Vostok reasonably fits Raymond's (1983) arguments for deformation in the vicinity of ice divides.</u>

At this stage we are not yet able to predict GHF at the bed of 600-m thick subglacial Lake Vostok because <u>the</u> temperature profile in the lake is still indefinite. However, the DNA detection of thermophile bacterium in the near-base accretion ice suggests the existence of near-bottom warm waters with temperatures as high as 50 °C (Bulat et al., 2012). If so, the GHF in

- 435 the lake sediments can reach 200-240 mW m<sup>-2</sup>. These values can be considered as paleo-GHF because microorganisms were picked up thousands of years ago but still actual accounting for long duration of geological processes.
  - **Dome C**. The inverse approach to retrieving GHF from radar inferred distribution of wet and dry beds at the EPICA drilling site (Passalacqua et al., 2017) gave 54.5 $\pm$ 3.5 mW m<sup>-2</sup>, <u>slightly</u> lower than estimates derived from borehole temperature profiling (<u>57.9 $\pm$ 6.4 mW m<sup>-2</sup></u>). The modeled GHF range (43–55 mW m<sup>-2</sup> obtained by An et al., 2015; Fox Maule et al., 2005;
- 440 Shapiro and Ritzwoller, 2004; Van Liefferinge and Pattyn, 2013) is also <u>a little less</u> than our estimates. <u>The high value for the correlation factor (0.997) indicates a perfectly strong relationship between modeled and measured depth-ages scales meaning that there no horizontal advection of heat and drill site is located at a perfect dome position. Perhaps, that is the reason that the core from Dome C contains the oldest continuous climate record obtained from ice cores so far (Parrenin et al., 2007b). However, a high spatial variation of GHF at Dome C area was found from radar-sounding data (Carter et al., 2009). The values</u>
- of nearly 100 mW m<sup>-2</sup> inferred for the southern shore of Concordia Subglacial Lake, approximately 50 km to the south of the drilling site, are also well outside modeled estimates.
  - *Kohnen*. The model with a standard GHF of 54.6 mW m<sup>-2</sup> predicted a basal temperature 0.3 °C below the pressure melting point at Kohnen (Huybrechtset al., 2007). <u>GHF obtained by non-thermal geophysical models are in the range of 46-62 W m<sup>-2</sup></u>. <u>Our estimate (86.9±16.6 mW m<sup>-2</sup>) is higher than the modeled GHF values suggested by Fox Maule et al. (2005), Martos et al.</u>
- 450 (2017) and other referenced models. However, subglacial water entering the borehole indicated that the actual GHF should be much higher than that indicated by the regional models. Under these circumstances, our estimate is likely to be closer to the real heat flux. Surprisingly, the depth-age scale is only slightly affected by the choice of the form factor indicating that the variation of horizontal velocity is low at this site.

**Dome F**. A previously estimated GHF of 59 mW m<sup>-2</sup> neglected the bottom ice melt rate (Hondoh et al., 2002) and thus is lower than our estimate (78.9±5.0 mW m<sup>-2</sup>). Mony et al. (2020) estimated the GHF in Dome F borehole to be even lower at 50.4 mW m<sup>-2</sup>. As the drill approached the base (approx. 10 m above), subglacial meltwater leaked into the borehole and froze onto the drill, directly indicating that ice reaches the pressure melting point, placing a lower bound on the GHF. <u>GHF values obtained</u> by non-thermal geophysical methods are in the range of 48-65 W m<sup>-2</sup>, also lower than our estimates. The correlation factor between modeled and measured depth-age scales is quite high (0.83) at m = 1 indicating ice at the site can be adequately

460 <u>approximated by the steady-state model. Thus, the slightly elevated heat flow at this location appears to represent a regional value.</u>

*Byrd*. Unfortunately, age scales for the Byrd borehole for all modeled *m* values are quite far from the measured depth-age data. Tilting measurements (Garfield and Ueda, 1976; Hansen et al., 1989) and modeling (Whillans, 1979) showed that the relative horizontal velocity of ice at this borehole reaches  $\sim$ 3 m a<sup>-1</sup> at the 1500 m depth. Thus, horizontal conduction in the

- 465 <u>bottom of the ice sheet is quite high at this site, producing a high divergence from the steady-state model. Even the correlation factor for the best fitting age-depth curve with m = 0.75 is only 0.58, we can use the GHF value of  $88.4\pm7.6 \text{ mW m}^{-2}$  at this location as a reference until more precise estimates are obtained. This value is higher than the first estimate made immediately after temperature logging (75.4 mW m<sup>-2</sup> referenced by Ueda, 2007), primarily because the latter one did not account for the basal ice melt. The latest modeling by Martos et al. (2017) revealed a high GHF at the location of Byrd Station (132 mW m<sup>-2</sup>)</u>
- 470 with an error of ±5 mW m<sup>-2</sup>) when compared with <u>values obtained from</u> previous models (An et al., 2015; Fox Maule et al., 2005; Van Liefferinge and Pattyn, 2013). <u>Generally, our approximate estimate is in the range of GHF values suggested by previous models</u>.

*WAIS Divide*. <u>A preliminary estimate</u> of the GHF at this site suggested a high value in the range of 200-230 mW m<sup>-2</sup>, depending on the actual ice thickness (Clow et al., 2012). This is more than twice that derived using non-thermal geophysical methods.

- 475 There is no depression in the local surface topography or drawdown in the subsurface layers detected by ice-penetrating radar, as would be expected over a local hot spot. Using airborne magnetic data, Martos et al. (2017) estimated the highest value for this area to be ~120 mW m<sup>-2</sup>. Our new estimate is slightly lower at 113.3±16.9 mW m<sup>-2</sup>. Mony et al. (2020) also estimated GHF from the borehole temperature profile at WAIS Divide by combining a heat-transfer equation and the physical properties of the ice sheet in a numerical model. Based on a truncated temperature profile, they estimate a GHF of 90.5 mW m<sup>-2</sup> which
- 480 is less than ours and fairly corresponds to the latest GHF map for Antarctica constructed by empirically relating the upper mantle seismic structure (Shen et al., 2020). The low correlation factor (0.59) between modeled and measured depth-age scales in our present estimate indicates that there are some important uncertainties that the steady-state model does not account for. Most likely, these are the same unaccounted effects that affect the Byrd borehole temperature profile, i.e., horizontal flow and climate-related transient effects.
- The preliminary GHF estimate (Clow et al., 2012) was based on the first temperature log in 2011 in the borehole before it reached its final depth. The reasons why the preliminary GHF estimate may be so high are that: (i) temperatures in the borehole were still thermally disturbed in 2011, and (ii) the bottom of the 2011 temperature log was still far from the base of the ice sheet. The borehole was relogged in 2014, and temperature data were obtained much closer to the bed. In addition, Clow et al. (2012) also did not account for horizontal flow effects and the GHF estimation could have been lower than the one they produced. Further investigations on ice dynamics through WAIS Divide borehole tilt measurements can allow us to determine
- in-depth stress and velocity distributions and estimate horizontal flow effects on temperature.

## **3.3 Indirect results**

Although a steady-state model is used in the lower portion of the boreholes to describe the temperature distribution, it is worth noting that the measured modern temperature is the cumulative effect of historical climate forcing. Therefore, the best fitting

- 495 parameters obtained by GA are not the real parameters occurring during the ice sheet's history. They can be considered as "equivalent" parameters which are used for calculating the modern temperature profile by eliminating the historical climate changes. Processing back, the "equivalent" vertical velocity, modern accumulation rate and temperature can be calculated from the GA results. Estimated vertical velocity profiles are shown on Fig. 5. Table 5 lists values of "equivalent" snow accumulation rate and temperature at ice sheet surface which were derived from GA calculations. In all cases, "equivalent" accumulation
- 500 rates are higher than the modern rates while the "equivalent" surface temperatures are very close to the modern ones. This can be explained by the fact that the high "equivalent" accumulation rates are used by GA to eliminate the colder climate effects on the ice temperature profile during the glacial period.

#### **4 Discussion**

#### 4.1 Transient model vs. steady-state model

- 505 Both, transient thermal models (e.g., Dahl-Jensen et al., 2003; Engelhardt, 2004; Martos et al., 2017; Passalacqua et al., 2017; Van Liefferinge et al., 2018) and steady-state models (e.g., Martin and Gudmundsson, 2012; Mony et al., 2020; Parrenin et al., 2017; Price et al., 2002; Zagorodnov et al., 2012) were used intensively in the past and are still used for GHF estimates in Antarctica. Obviously, an exact steady state never occurs in reality and thus transient models would be expected to give more precise results than steady-state models. However, the answer is not as simple as it is supposed to be.
- 510 It is important to recognize that, first, in both cases the models will produce GHF "estimates", not "measurements", and second, the thermal gradient can be affected by processes other than GHF, creating local anomalies that may coincide with the point estimate. In order to use a transient model, the accumulation rate and surface temperature in the past should be known. For some of the discussed drill sites these data are available from ice-core studies, while for other sites they are not. To evaluate the possibility of using a transient model, the GHF at WAIS Divide was estimated by using the accumulation rate
- 515 and surface temperature in the past provided by Buizert et al. (2015). In these calculations, we assume that the history of the ice sheet at WAIS Divide is about 68 ka long. The governing equation for the transient model was solved using the finite difference method. The equation was discretized by both the central-difference and upwind-difference methods and then solved using Matlab. To find the best solution, the GA algorithm was still used. The central-difference method and upwind-difference method demonstrated the same temperature profile. Therefore, here we present the calculation results obtained via the upwind upwind-difference method upwind-difference upwind-difference
- 520 <u>difference method.</u> <u>Unfortunately, the calculation results with the transient model showed the best-fit GHF value of ~500 mW m<sup>-2</sup> when m = 1, which seems to be unrealistic. Moreover, after running the model, we found that after about 4-8 ka, the influence of initial</u>

temperature on temperature profile can be ignored. Later, we assumed the form factor m = 0 and a GHF value of 235 mW m<sup>-2</sup> which showed a good fit with the measured temperatures, although the GHF is much higher than our earlier estimate and

- 525 estimates from regional models. The temperature distribution in history was modeled 61.2 ka, 54.4 ka, ... 6.8 ka ago until modern day (Fig. 6). As expected, the modeled temperature in the upper part of the ice sheet grossly changes with time but in the lower portion (~1000 m above ice sheet base) these variations are much smaller. This indicates that the heat disturbance related to atmosphere forcing (temperature and snow precipitation) from the ice-sheet surface gradually decays with the depth. From all appearances, the near-basal portion is close to a steady state.
- 530 Both models lack additional heat sources (i.e., shear heating, heat advection and basal frictional heating) that might be generated at the bottom of the ice sheet. Thus, the results of both modeling approaches strongly depend on the selected initial parameters, in particular, from the selected value for the form factor *m*. Experimental validation of both models for adequacy is extremely difficult and recently both of them have rights to exist if assumptions are examined analytically.

#### **4.2 Implications of elevated heat flux**

- Most numerical models of the EAIS basal conditions assume the GHF to be 42-65 mW m<sup>-2</sup>. However, the presence of basal meltwater beneath most of the Antarctic ice sheet requires  $GHF \ge 80 \text{ mW m}^{-2}$  (Budd et al., 1984). In support of this conjecture, processing of available temperature profiles in ice boreholes shows that at sites where subglacial water exists bedrocks are quite warm. Currently, a warm base beneath the EAIS was also confirmed by tectonic reconstructions (Carson et al., 2014). Additional evidence of high GHF at EAIS locations comes from ice-penetrating radar data that revealed a ~100 km long and
- 540 <u>50 km wide area near South Pole with GHF of 120±20 mW m<sup>-2</sup>, more than double the values expected for this cratonic sector of East Antarctica (Jordan et al., 2018). This warm base could be caused by radiogenic heat effects or hydrothermal circulation, but a coherent explanation for this phenomenon is still required.</u>

Variability of crustal thickness, hydrothermal circulation (Seroussi et al., 2017), magmatic intrusion (Van Wyk De Vries et al., 2017), and thermal conductivity variability are the main contributors to the elevated and highly variable <u>values of GHF</u> in West Antarctica (Begeman et al., 2017). One of the first pieces of evidence for an "unreasonably high" GHF (>100 mW m<sup>-2</sup>) under WAIS came from temperature-depth profiles in a 480-m-deep borehole drilled at Crary Ice Rise, Antarctica (Bindschadler et al., 1990). <u>Measured GHF in Subglacial Lake Whillans was found to be 285 ± 80 mW m<sup>-2</sup>, significantly higher than the continental and regional averages estimated for this site by using geophysical and glaciological models (Fisher
</u>

et al., 2015). The GHF at the vents of subglacial volcanoes in West Antarctica can be as high as 25 W m<sup>-2</sup> and one such 23-

550 km-wide caldera was revealed ~100 km to the south of the WAIS Divide drill site (Blankenship et al., 1993). <u>Undeniably</u>, more systematic explorations are still required to study how far this heat flow high extends into the interior of the West Antarctic Rift System.

## **5** Conclusions

Prediction of the future behavior of the Antarctic Ice Sheet undeniably requires accurate ice-sheet models. However, GHF

- 555 models based on seismic tomography, radar data, magnetic field observations, the tectonic age and geological structure of the bedrock yiel<u>ds</u> mixed results at sites of deep ice-core drilling in Antarctica. We suggested to estimate GHF from ice-borehole temperature profiles using a one-dimensional steady-state energy-balance equation and the genetic algorithm (GA) for determining the optimal solution of temperature fitting. To our knowledge, we used GA approach for the first time in ice thermodynamics. Comparison of modeled and measured depth-age scales show that our model is able to assess the variation
- 560 in GHF estimates from ice-borehole temperature profiles if in-depth horizontal ice velocities are low and can be ignored. The correlation analyses at the EAIS sites indicates that all of them can be adequately approximated by the steady-state model. However, horizontal velocities and their variation over ice-age cycles are much greater at WAIS than at the EAIS sites. Thus, the steady-state model cannot precisely describe temperature distribution here.
- At three studied EAIS sites (Dome C, Dome F, and Kohnen), the GHF is higher than that predicted by other models. We assume that this elevated GHF can represent regional value and can be used as a reference point for regional modelings. More precise GHF estimates and explanations for an elevated GHF would be possible after temperature logging and subglacial rock studies from deep boreholes that are required to drill in Antarctica in the distant future. Finally, the proposed method of GHF estimates can be used at other sites in Antarctica and Greenland where the steady-state model is acceptable.

#### Acknowledgments

570 This work was supported by grant Nos. 41327804 and 41806220 from the National Natural Science Foundation of China and the Program for Jilin University Science and Technology Innovative Research Team (Fundamental Research Funds for the Central Universities, Project No. 2017TD-24). We are grateful to H. Ueda (retired from USA CRREL) for providing original Byrd borehole temperature-log data. We thank D. Dahl-Jensen (Centre for Ice and Climate, University of Copenhagen, Denmark) for fruitful discussion and useful comments. We also would like to thank the Editor Alex Robinson and both anonymous reviewers for fruitful comments and advices.

#### References

- Ackert, R.P., David, J.B., Harold, W.B., Parker, E.C., Mark, D.K., James, L.F., and Eric, J.: Measurements of past ice sheet elevations in interior West Antarctica. Science, 286(5438), 276-280, doi:10.1126/science.286.5438.276, 1999.
   Ackert, R.P., Mukhopadhyay, S., Parizek, B.R., and Borns, H.W.: Ice elevation near the West Antarctic Ice Sheet divide during the Last Glaciation, Geophys. Res. Lett., 34 (21), L21506, doi:10.1029/2007GL031412, 2007.
- Ahn, J. and Brook E.J.: Atmospheric CO<sub>2</sub> and Climate on Millennial Time Scales During the Last Glacial Period. Science, 322, 83-85, doi: 10.1126/science.1160832, 2008.

Anderson, J.B., Shipp, S.S., Lowe, A.L., Wellner, J.S., and Mosola, A.B.: The Antarctic Ice Sheet during the Last Glacial Maximum and its subsequent retreat history: a review. Quat. Sci. Rev., 21, 49-70, doi:10.1016/S0277-3791(01)00083-X, 2002.

- 585 An, M., Wiens, D.A., Zhao, Y., Feng, M., Nyblade, A., Kanao, M., Li, Y., Maggi, A., and Lévêque, J.-J.: Temperature, lithosphere-asthenosphere boundary, and heat flux beneath the Antarctic Plate inferred from seismic velocities. Journal of Geophysical Research: Solid Earth, 120, 8720–8742, doi:10.1002/2015JB011917, 2015. Augustin, L., Panichi, S., and Frascati, F.: EPICA Dome C 2 drilling operations: performances, difficulties, results. Ann.
  - Glaciol., 47, 68-72, doi:10.3189/172756407786857767, 2007.
- 590 Bazin, L., Landais, A., Lemieux-Dudon, B., Toyé Mahamadou Kele, H., Veres, D., Parrenin F., Martinerie, P., Ritz, C., Capron, E., Lipenkov, V., Loutre, M.-F., Raynaud, D., Vinther, B., Svensson, A., Rasmussen, S.O., Severi, M., Blunier, T., Leuenberger, M., Fischer, H., Masson-Delmotte, V., Chappellaz, J., and Wolff, E.: An optimized multi-proxy, multi-site Antarctic ice and gas orbital chronology (AICC2012): 120-800 ka. Clim. Past, 4, 1715-1731, doi: 10.5194/cp-9-1715-2013, 2013.
- Begeman, C.B., Tulaczyk, S.M., and Fisher, A.T.: Spatially variable geothermal heat flux in West Antarctica: Evidence and implications. Geophys Res Lett., 44, 9823–9832, doi:10.1002/2017gl075579, 2017.
   Bereiter, B., Lüthi, D., Siegrist, M., Schüpbach, S., Stocker, T.F., Fischer, H.: Mode change of millennial CO<sub>2</sub> variability during the last glacial cycle associated with a bipolar marine carbon seesaw. Proc. Natl. Acad. Sci. USA. 109 (25), 9755-9760, doi:10.1073/pnas.1204069109, 2012.
- Bindschadler, R.A., Roberts, E.P., and Iken, A.: Age of Crary Ice Rise, Antarctica, determined from temperature-depth profiles. Ann Glaciol., 14, 13-16, doi:10.1017/S0260305500008168, 1990.
  Blankenship, D.D., Bell, R.E., Hodge, S.M., Brozena, J.M., Behrengt, J.C., and Finn, C.A.: Active volcanism beneath the West Antarctic ice sheet and implications for ice-sheet stability. Nature, 361, 526-529, doi:10.1038/361526a0, 1993.
  <u>Blunier, T. and Brook, E.J.: Timing of millennial-scale climate change in Antarctica and Greenland during the last glacial</u> period. Science, 291, 109-112, doi: 10.1126/science.291.5501.109, 2001.
  - Budd, W.F., Jenssen, D., and Smith I.N.: A three-dimensional time-dependent model of three West Antarctic ice-sheet. Ann Glaciol., 5, 29–36, doi:10.1017/S026030550000344X, 1984.

Buizert, C., Cuffey, K.M., Severinghaus, J.P., Baggenstos, D., Fudge, T.J., Steig, E.J., Markle, B.R., Winstrup, M., Rhodes, R.H., Brook, E.J., Sowers, T.A., Clow, G.D., Cheng, H., Edwards, R.L., Sigl, M., McConnell, J.R., and Taylor, K.C.: The

610 WAIS Divide deep ice core WD2014 chronology – Part 1: Methane synchronization (68–31 ka BP) and the gas age-ice age difference. Clim. Past, 11, 153–173, doi:10.5194/cp-11-153-2015, 2015.

Bulat, S.A., Marie, D., and Petit, J.-R.: Prospects for life in the subglacial Lake Vostok. Ice and Snow, 4(120), 92-96, doi:10.15356/2076-6734-2012-4-92-96, 2012.

Burton-Johnson, A., Dziadek, R., and Martin, C.: Geothermal heat flow in Antarctica: current and future directions. Cryosphere

615 <u>Discuss., in review, doi:10.5194/tc-2020-59, 2020.</u>

Carter, S.P., Blankenship, D.D., Young, D.A., and Holt, J.W.: Using radar-sounding data to identify the distribution and sources of subglacial water: application to Dome C, East Antarctica. J Glaciol., 55(194), 1025-1040, doi:10.3189/002214309790794931, 2009.

Carson, C.J., McLaren, S., Roberts, J.L., Boger, S.D., and Blankenship, D.D.: Blankenship, hot rocks in a cold place: High sub-glacial heat flow in East Antarctica. J Geol Soc London, 171(1), 9–12, doi:10.1144/jgs2013-030, 2014.

Clow, G.D.: USGS Polar temperature logging system, description and measurement uncertainties: U.S. Geological Survey Techniques and Methods 2–E3, 2008.

Clow, G.D.: A Green's function approach for assessing the thermal disturbance caused by drilling deep boreholes in rock or ice. Geophys. J. Int., 203, 1877–1895, doi: 10.1093/gji/ggv415, 2015.

625 Clow, G., Waddington, E., Hawley, R., and Dahl-Jensen, D.: Subglacial heat flow measurements in Greenland and Antarctica. European Geosciences Union General Assembly, 3-8 April, 2011, Vienna, Austria. Geophysical Research Abstracts 13, abstract id. EGU2011-6629, 2011.

Clow, G.D., Cuffey, K.M., and Waddington, E.D.: High heat-flow beneath the central portion of the West Antarctic Ice Sheet. American Geophysical Union, Fall Meeting 2012, 3-7 December, 2012, San-Francisco, USA, abstract id. C31A-0577, 2012.

- <u>Conway, H., Hall, B.L., Denton, G.H., Gades, A.M., and Waddington, E.D.: Past and future grounding-line retreat of the West Antarctic Ice Sheet. Science, 286, 280-283, doi:10.1126/science.286.5438.280, 1999.</u>
   <u>Conway, H. and Rasmussen L.A. (2009). Recent thinning and migration of the Western Divide, central West Antarctica.</u>
   <u>Geophys.Res. Lett., 36, L12502, doi: 10.1029/2009GL038072, 2009.</u>
   Cuffey, K.M. and Paterson, W.S.B. The physics of glaciers, 4th edn. Butterworth-Heinemann, Oxford, 2010.
- 635 Davies, J.H.: A global map of solid Earth surface heat flow. Geochemistry, Geophysics and Geosystems, 14, 4608-4622, doi:10.1002/ggge.20271, 2013.

<u>Cuffey, K.M., Clow, G.D., Steig, E.J., Buizert, C., Fudge, T.J., Koutnik, M., Waddington, E.D., Alley, R.B., and Severinghaus, J.P.: Deglacial temperature history of West Antarctica. PNAS, 113 (50), 14249–14254, doi:10.1073/pnas.1609132113, 2016.
 <u>Gundestrup, N.S., Clausen, H.B., and Hansen, B.L.: The UCPH borehole logger. Mem. Nat. Inst. Polar Res., 49, 224-233,</u>
</u>

640 <u>1994.</u>

Dahl-Jensen, D., Morgan, V.I., and Elcheikh, A.: Monte Carlo inverse modelling of the Law Dome (Antarctica) temperature profile. Ann. Glaciol., 29, 145-150, doi:10.3189/172756499781821102, 1999.

Dahl-Jensen, D., Gundestrup, N., Gogineni, S.P., and Miller H.: Basal melt at NorthGRIP modeled from borehole, ice-core and radio-echo sounder observations. Ann. Glaciol., 37, 217-212, doi:10.3189/172756403781815492, 2003.

545 Davis, J.H.: Global map of solid Earth surface heat flow. Geochemistry, Geophysics, Geosystems, 14(10), 4608-4622, doi:10.1002/ggge.20271, 2013.

Decker, E.R. and Bucher, G.J.: Geothermal studies in the Ross Island-Dry Valley region. Antarct Geosci, 4, 887-894, 1982.

Ekaykin, A.A., Lipenkov, V.Ya., and Shibaev, Yu.A.: Spatial distribution of the snow accumulation rate along the ice flow lines between Ridge B and Lake Vostok. Lyed i Sneg [Ice and Snow], 4(120), 122-128, doi:10.15356/2076-6734-2012-4-122-

650 128, 2012.

Engelhardt, H.: Ice temperature and high geothermal flux at Siple Dome, West Antarctica, from borehole measurements. J Glaciol., 50(169), 251-256, doi:10.3189/172756504781830105, 2004.

Fischer, H., Severinghaus, J., Brook, E., Wolff, E., Albert, M., Alemany, O., Arthern, R., Bentley, C., Blankenship, D., Chappellaz, J., Creyts, T., Dahl-Jensen, D., Dinn, M., Frezzotti, M., Fujita, S., Gallee, H., Hindmarsh, R., Hudspeth, D., Jugie,

655 G., Kawamura, K., Lipenkov, V., Miller, H., Mulvaney, R., Parrenin, F., Pattyn, F., Ritz, C., Schwander, J., Steinhage, D., van Ommen, T., and Wilhelms, F.: Where to find 1.5 million yr old ice for the IPICS "Oldest-Ice" ice core. Clim. Past, 9, 2489– 2505, doi:10.5194/cp-9-2489-2013, 2013.

Fisher, A.T., Mankoff, K.D., Tulaczyk, S.M., Tyler, S.W., Foley, N., and the WISSARD Science Team: High geothermal heat flux measured below the West Antarctic Ice Sheet. Sci. Adv., 1(6), e1500093, doi:10.1126/sciadv.1500093, 2015.

660 Fox Maule, C.F., Purucker, M.E., Olsen, N., and Mosegaard, K.: Heat flux anomalies in Antarctica revealed by satellite magnetic data. Science, 309, 464–467, doi:10.1126/science.1106888, 2005.

Garfield, D.E. and Ueda, H.T.: Resurvey of the "Byrd" Station, Antarctica, drill hole. J. Glaciol., 17(75). 29-34, doi: 10.3189/S0022143000030689, 1976.

<u>Golledge, N.R., Kowalewski, D.E., Naish, T.R., Levy R.H., Fogwill, C.J., and Gasson, E.G.W.: The multi-millennial Antarctic</u> commitment to future sea-level rise. Nature, 526(7573), 421–425, doi:10.1038/nature15706, 2015.

Goodge, J.W.: Crustal heat production and estimate of terrestrial heat flow in central East Antarctica, with implications for thermal input to the East Antarctic ice sheet. Cryosphere, 12, 491–504, doi:10.5194/tc-12-491-2018, 2018.

Gow, A.J.: Deep core studies of the accumulation and densification of snow at Byrd Station and Little America V, Antarctica. CRREL Res. Rep. 197, 1968.

 Hansen, B.L., Kelty, J.R., and Gundestrup, N.S.: Resurvey of Byrd Station drill hole, Antarctica, Cold Reg. Sci. Tech., 17 (1), 1-6, doi: 10.1016/S0165-232X(89)80011-4, 1989. Hindmarsh, R.C.A.: On the numerical computation of temperature in an ice sheet. J. Glaciol., 45, 568–574, doi: 10.1017/S0022143000001441, 1999.

Hindmarsh, R.: Ice-sheet and glacier modelling, in: Past Glacial Environments, Elsevier, 605-661, 2018.

- Hondoh, T., Shoji, H., Watanabe, O., Salamatin, A.N., and Lipenkov, V.Y.: Depth-age and temperature prediction at Dome Fuji station, East Antarctica. Ann. Glaciol. 35, 384–390, doi:10.3189/172756402781817013, 2002.
  Huybrechts, P., Rybak, O., Pattyn, F., Ruth, U. and Steinhage D.: Ice thinning, upstream advection, and non-climatic biases for the upper 89% of the EDML ice core from a nested model of the Antarctic ice sheet. Clim. Past, 3, 577–589, 10.5194/cp-3-577-2007, 2007.
- 580 Johnsen, S., Dahl-Jensen, D., Dansgaard, W., and Gundestrup N.: Greenland palaeotemperatures derived from GRIP bore hole temperature and ice core isotope profiles. Tellus, 47B, 624-629, doi:10.3402/tellusb.v47i5.16077, 1995.

Jordan, T.A., Martin, C., Ferraccioli, F., Matsuoka, K., Corr, H., Forsberg, R., Olesen, A., and Siegert M.: Anomalously high geothermal flux near the South Pole. Sci. Rep., 8, 16785, doi:10.1038/s41598-018-35182-0, 2018.

Jouzel, J., Petit, J.R., Souchez, R., Barkov, N.I., Lipenkov, V.Y., Raynaud, D., Stievenard, M., Vassiliev, N.I., Verbeke, V.,

and Vimeux F.: More than 200 meters of lake ice above subglacial Lake Vostok, Antarctica. Science, 286(5447), 2138-2141, doi:10.1126/science.286.5447.2138, 1999.

Kawamura, K., Parrenin, F., Lisiecki, L., Uemura, R., Vimeux, F., Severinghaus, J.P., Hutterli, M.A., Nakazawa, T., Aoki, S., Jouzel, J., Raymo, M.E., Matsumoto, K., Nakata, H., Motoyama, H., Fujita, S., Goto-Azuma, K., Fujii, Y., and Watanabe O.: Northern Hemisphere forcing of climatic cycles in Antarctica over the past 360,000 years. Nature, 448, 912-916, doi:10.1038/nature06015, 2007.

- Koutnik, M.R., Fudge, T.J., Conway, H., Waddington E.D., Neumann T.A., Cuffey K.M., Buizert C., and Taylor K.C.: Holocene accumulation and ice flow near the West Antarctic Ice Sheet Divide ice core site, J. Geophys. Res. Earth Surf., 121, 907–924, doi:10.1002/2015JF003668, 2016.
   Lefebvre, E., Augustin, L., and Maitre, M.: The EPICA borehole logger. Mem. Nat. Inst. Polar Res., 56, 264-274, 2002.
- Llubes, M., Lanseau, C., and Rémy, F.: Relations between basal condition, subglacial hydrological networks and geothermal flux in Antarctica. Earth Planet. Sc. Lett. 241, 655–662, doi:10.1016/j.epsl.2005.10.040, 2006.
   Lukin, V.V. and Vasiliev, N.I.: Technological aspects of the final phase of drilling borehole 5G and unsealing Vostok Subglacial Lake, East Antarctica. Ann. Glaciol., 55(65), 83-89, 10.3189/2014AoG65A002, 2014.
   Martin, C. and Gudmundsson, G.H.: Effects of nonlinear rheology, temperature and anisotropy on the relationship between
- <u>age and depth at ice divides. Cryosphere, 6, 1221–1229, doi:10.5194/tc-6-1221-2012, 2012.</u>
   Martos, Y.M., Catalán, M., Jordan, T.A., Golynsky, A., Golynsky, D., Eagles, G., and Vaughan, D.G.: Heat flux distribution of Antarctica unveiled. Geophys Res Lett., 44, 11417–11426, doi:10.1002/2017GL075609, 2017.
   <u>Mony, L., Roberts, J.L., and Halpin, J.A.: Inferring geothermal heat flux from an ice-borehole temperature profile at Law Dome, East Antarctica. J. Glaciol., 66(257), 509-519, doi:10.1017/jog.2020.27, 2020.</u>
- 705 Motoyama, H.: The second deep ice coring project at Dome Fuji, Antarctica. Sci. Drilling, 5, 41-43, doi:10.2204/iodp.sd.5.05.2007, 2007.

Motoyama, H., Furukawa, T., and Nishio, F. (2008). Study of ice flow observations in Shirase drainage basin and around
 Dome Fuji area, East Antarctica by differential GPS method. Nankyoku Shiryo (Antarctic Record, 52, 216-231, 2008.
 Motoyama, H., Furusaki, A., Takahashi, A., Tanaka, Y., Miyahara, M., Takata, M., Sawagaki, T., Matoba, S., Sugiyama, S.,

710 Shinbori, K., and Mori, S.: Deep borehole logging at Dome Fuji Station, Antarctica. Abstracts of the Fourth Symposium on Polar Science, 12-15 November, 2013, National Institute of Polar Research, Tachikawa, Tokyo, Japan, 2013. Neftel, A., Oeschger, H., Staffelbach, T., and Stauffer, B.: CO<sub>2</sub> record in the Byrd ice core 50000-5000 years BP. Nature, 331, 609-611, doi: 10.1038/331609a0, 1988.

Nicholls, K.W. and Paren, J.G.: Extending the Antarctic meteorological record using ice-sheet temperature profiles. J Clim.,

715 6, 141-150, doi:10.1175/1520-0442(1993)006<0141:ETAMRU>2.0.CO;2, 1993.

Parrenin, F., Dreyfus, G., Durand, G., Fujita, S., Gagliardini, O., Gillet, F., Jouzel, J., Kawamura, K., Lhomme, N., Masson-Delmotte, V., Ritz, C., Schwander, J., Shoji, H., Uemura, R., Watanabe, O., and Yoshida, N.: 1-D-ice flow modelling at EPICA Dome C and Dome Fuji, East Antarctica. Clim. Past, 3, 243–259, doi:10.5194/cp-3-243-2007, 2007<u>a</u>.

Parrenin, F., Barnola, J.-M., Beer, J., Blunier, T., Castellano, E., Chappellaz, J., Dreyfus, G., Fischer, H., Fujita, S., Jouzel, J.,

720 Kawamura, K., Lemieux-Dudon, B., Loulergue, L., Masson-Delmotte, V., Narcisi, B., Petit, J.-R., Raisbeck, G., Raynaud, D., Ruth, U., Schwander, J., Severi, M., Spahni, R., Steffensen, J.P., Svensson, A., Udisti, R., Waelbroeck, C., and Wolff E.: The EDC3 chronology for the EPICA Dome C ice core. Clim. Past, 3, 485-497, doi: 10.5194/cp-3-485-2007, 2007b. Parrenin, F., Cavitte, M.G.P., Blankenship, D.D., Chappellaz, J., Fischer, H., Gagliardini, O., Masson-Delmotte, V.,

Passalacqua, O., Ritz, C., Roberts, J., Siegert, M.J., Young, D.A.: Is there 1.5-million-year-old ice near Dome C, Antarctica? 725 Cryosphere, 11, 2427–2437, doi:10.5194/tc-2017-69, 2017.

- Passalacqua, O., Ritz, C., Parrenin, F., Urbini, S., and Frezzotti, M.: Geothermal flux and basal melt rate in the Dome C region inferred from radar reflectivity and heat modelling. Cryosphere, 11, 2231–2246, doi:10.5194/tc-11-2231-2017, 2017.
  Pattyn, F.: Antarctic subglacial conditions inferred from a hybrid ice sheet/ice stream model. Earth Planet. Sci. Lett., 295, 451–461, doi:10.1016/j.epsl.2010.04.025, 2010.
- Pittard, M.L., Galton-Fenzi, B.K., Roberts, J.L., and Watson, C.S.: Organization of ice flow by localized regions of elevated geothermal heat flux. Geophys Res Lett., 43, 3342–3350, doi:10.1002/2016GL068436, 2016.
  Pollard, D., DeConto, R.M., and Nyblade, A.A.: Sensitivity of Cenozoic Antarctic ice sheet variations to geothermal heat flux. Global Planet. Change, 49, 63–74, doi:10.1016/j.gloplacha.2005.05.003, 2005.
- Price, P.B., Nagornov, O.V., Bay, R., Chirkin, D., He, Y., Miocinovic, P., Richards, A., Woschnagg, K., Koci, B., and
- 735 Zagorodnov, V.: Temperature profile for glacial ice at the South Pole: Implications for life in a nearby subglacial lake. PNAS 99(12), 7844–7847, doi:10.1073/pnas.082238999, 2002

Raymond, C.F.: Deformation in the vicinity of ice divides. J. Glaciol., 29, 357–373, doi:10.1017/S0022143000030288, 1983. Reeves, C.R. and Rowe, J. E.: Genetic Algorithms: Principles and Perspectives. A Guide to GA Theory. Kluwer Academic Publishers, Boston (USA), 2002.

740 <u>Rignot, E., Mouginot, J., and Scheuchl, B.: Ice flow of the Antarctic Ice Sheet. Science 333, 1427-1430, doi:</u> 10.1126/science.1208336, 2011.

Risk, G.F. and Hochstein, R.: Heat flow at Arrival Heights, Ross Island, Antarctica. New Zeal J Geol Geop., 17, 629–644, doi:10.1080/00288306.1973.10421586, 1974.

 <u>Ritz, C., Rommelaere, V., and Dumas, C.: Modeling the evolution of Antarctic ice sheet over the last 420,000 years:</u>
 <u>Implications for altitude changes in the Vostok region. J. Geophys. Res. Atmos. 106 (D23), 31943 – 31964,</u> doi:10.1029/2001JD900232, 2001.

Robin, G. de Q.: Ice movement and temperature distribution in glaciers and ice sheets. J. Glaciol., 2, 523–532, doi:10.3189/002214355793702028, 1955.

Salamatin, A.N., Vostretsov R.N., Petit J.R., Lipenkov, V.Y. and Barkov, N.I.: Geofisicheskiye i paleoklimaticheskie

- 750 prilozheniya sostavnogo temperaturnogo profilya iz glubokoi skvazhyni na stantsii Vostok (Antaktida) [Geophysical and palaeoclimatic implications of the stacked temperature profile from the deep borehole at Vostok station, Antarctica]. Mater. Glyatsiol. Issled., 85, 233–240 (in Russian with English summary), 1998a.
- Salamatin, A.N., Lipenkov, V.Y., Barkov, N.I., Jouzel, J., Petit, J R., and Raynaud, D.: Ice core age dating and paleothermometer calibration based on isotope and temperature profiles from deep boreholes at Vostok Station (East Antarctica). J Geophys Res., 103(D8), 8963–8977, doi:10.1029/97JD02253, 1998b.
- Seroussi, H., Ivins, E.R., Wiens D.A., and Bondzio J.: Influence of a West Antarctic mantle plume on ice sheet basal conditions.
   J Geophys. Res. Solid Earth, 122, 7127–7155, doi:10.1002/2017jb014423, 2017.
   Shapiro, N.M. and Ritzwoller, M.H.: Inferring surface heat flux distributions guided by a global seismic model: particular application to Antarctica. Earth Planet. Sci. Lett., 223, 213-224, doi:10.1016/j.epsl.2004.04.011, 2004.
- 760 Shen, W., Wiens, D., Lloyd, A., and Nyblade, A.: A geothermal heat flux map of Antarctica empirically constrained by seismic structure. Geophys. Res. Lett., 47, e2020GL086955, doi: 10.1029/2020GL086955, 2020. Sigl, M., Fudge, T.J., Winstrup, M., Cole-Dai, J., Ferris, D., McConnell, J.R., Taylor, K.C., Welten, K.C., Woodruff, T.E., Adolphi, F., Bisiaux, M., Brook, E.J., Buizert, C., Caffee, M.W., Dunbar, N.W., Edwards, R., Geng, L., Iverson, N., Koffman, B., Layman, L., Maselli, O.J., McGwire, K., Muscheler, R., Nishiizumi, K., Pasteris, D.R., Rhodes, R.H. and Sowers, T.A.:
- 765 The WAIS Divide deep ice core WD2014 chronology Part 2: Annual-layer counting (0-31 ka BP), Clim. Past, 3, 769-786, doi: 10.5194/cp-12-769-2016, 2016.
  - Slawny, K.R., Johnson, J.A., Mortensen, N.B., Gibson, C.J., Goetz, J.J., Shturmakov, A.J., Lebar, D.A., and Wendricks, A.W.: Production drilling at WAIS Divide. Ann. Glaciol., 55(68), doi:10.3189/2014AoG68A018, 147-155, 2014.
- Staffelbach, T., Stauffer, B., Sigg, A., and Oeschger, H.: CO<sub>2</sub> measurements from polar ice cores: more data from different sites, Tellus, 43B, 91-96, doi: 10.3402/tellusb.v43i2.15251, 1991.
  - Ueda, H.T. and Garfield, D.E.: Deep core drilling at Byrd Station, Antarctica. In: A.J. Gow (Ed.) Proceedings of International Symp. on Antarctic Glaciological Exploration (ISAGE), Hanover, New Hampshire, USA, September 3-7, 1968. Association of Scientific Hydrology, Publ. No 86, Cambridge, UK, 53-62, 1970.

Ueltzhöffer, K.J., Bendel, V., Freitag, J., Kipfstuhl, S., Wagenbach, D., Faria, S.H., and Garbe, C.S.: Distribution of air bubbles
in the EDML and EDC (Antarctica) ice cores, using a new method of automatic image analysis. J Glaciol., 56(196), 339–348, doi:10.3189/002214310791968511, 2010.

Van Wyk De Vries, M., Robert G. Bingham R.G., and Hein A.S.: A new volcanic province: An inventory of subglacial volcanoes in West Antarctica. In: Siegert, M.J., Jamieson, S.S.R., and White, D.A. (eds) Exploration of Subsurface Antarctica: Uncovering Past Changes and Modern Processes. Geological Society, London, Special Publications 461, doi:10.1144/SP461.7, 2017.

Van Liefferinge, B. and Pattyn, F.: Using ice-flow models to evaluate potential sites of million year-old ice in Antarctica. Clim. Past, 9, 2335–2345, doi:10.5194/cpd-9-2859-2013, 2013.

Van Liefferinge, B., Pattyn, F., Cavitte, M.G.P., Karlsson, N.B., Young, D.A., Sutter, J., and Eisen, O. Promising Oldest Ice sites in East Antarctica based on thermodynamical modelling. Cryosphere, 12, 2773–2787, doi:10.5194/tc-12-2773-2018, 2018.

Vasiliev, N.I., Talalay, P.G., and Vostok Deep Ice Core Drilling Parties: Twenty years of drilling the deepest hole in ice. Sci. Drilling, 11, 41-45, doi:10.5194/sd-11-41-2011, 2011.

Veres, D., Bazin, L., Landais, A., Toyé Mahamadou Kele, H., Lemieux-Dudon, B., Parrenin, F., Martinerie, P., Blayo, E., Blunier, T., Capron, E., Chappellaz, J., Rasmussen, S.O., Severi, M., Svensson, A., Vinther, B., and Wolff, E.W.: The Antarctic

- ice core chronology (AICC2012): an optimized multi-parameter and multi-site dating approach for the last 120 thousand years. Clim. Past, 1733-1748, doi: 10.5194/cp-9-1733-2013, 2013.
   Vittuari, L., Vincent, C., Frezzotti, M., Mancini, F., Gandolfi, S., Bitelli, G., and Capra, A: Space geodesy as a tool for measuring ice surface velocity in the Dome C region and along the ITASE traverse, Ann. Glaciol., 39, 402–408, doi: 10.3189/172756404781814627, 2004.
- 795 Wexler, H.: Growth and thermal structure of the deep ice in Byrd Land, Antarctica. J Glaciol., 3(30), 1075-1087, doi:10.1017/S0022143000017482, 1961.

WAIS Divide Project Members: Onset of deglacial warming in West Antarctica driven by local orbital forcing. Nature, 500(7463), 440–444, doi: 10.1038/nature12376, 2013.

- Wendt, J., Dietrich, R., Fritsche, M., Wendt, A., Yuskevich, A., Kokhanov, A., Senatorov, A. Lukin, V., Shibuya K., and Doi
- K.: Geodetic observations of ice flow velocities over the southern part of subglacial Lake Vostok, Antarctica, and their glaciological implications. Geophys. J. Int., 166(3), 991–998, doi: 10.1111/j.1365-246X.2006.03061.x, 2006.
   Wesche, C., Eisen, O., Oerter, H., Schulte, D., Steinhage, D.: Surface topography and ice flow in the vicinity of the EDML deep-drilling site, Antarctica. J. Glaciol. 53, 442–448, doi:10.3189/002214307783258512, 2007.
   Whillans, I.M.: Ice flow along the Byrd Station strain network, Antarctica. J. Glaciol., 24(90), 15-28, doi:
- 805 <u>10.3189/S0022143000014611, 1979.</u> Whillans, I.M.: The equation of continuity and its application to the ice sheet near "Byrd" Station, Antarctica. J. Glaciol.,

785

<u>18(80)</u>, <u>359-371</u>, <u>doi:</u> <u>10.3189/S0022143000021055</u>, <u>1977</u>. Wilhelms, F., Miller, H., Gerasimoff, M.D., Drücker, C., Frenzel, A., Fritzsche, D., Grobe, H., Hansen, S.B., Hilmarsson,

S.Æ., Hoffmann, G., Hörnby, K., Jaeschke, A., Jakobsdóttir, S.S., Juckschat, P., Karsten, A., Karsten, L., Kaufmann, P.R., 810 Karlin, T., Kohlberg, E., Kleffel, G., Lambrecht, A., Lambrecht, A., Lawer, G., Schärmeli, I., Schmitt, J., Sheldon, S.G.,

- Takata, M., Trenke, M., Twarloh, B., Valero-Delgado, F., <u>and Wilhelms-Dick</u>, D.: The EPICA Dronning Maud Land deep drilling operation. Ann. Glaciol, 55(68), 355-366, doi:10.3189/2014AoG68A189, 2014. Yen, Y.: Review of thermal properties of snow, ice and see ice. CRREL Rep. 81-10, 1981.
- Zagorodnov, V., Nagornov, O., Scambos, T.A., Muto, A., Mosley-Thompson, E., Pettit, E.C., and Tyuflin, S.: Borehole 815 temperatures reveal details of 20th century warming at Bruce Plateau, Antarctic Peninsula. Cryosphere, 6, 675–686, 10.5194/tcd-5-3053-2011, 2012.

Zotikov, I.A.: Izmerenie geotermicheskogo potoka tepla v Antarktide [Measurement of the geothermal heat flow in Antarctica]. Sovetskaia antarkticheskaia ekspeditsiia. Informatsionnyi biulleten [Soviet Antarctic Expedition, Information Bulletin] 29, 30–32 [Text in Russian], 1961.

## 820 Table 1: Information for Antarctic deep ice-drilling sites

Parameters	W	AIS	EAIS				
	Byrd	WAIS	Vostok	Dome C	Kohnen	Dome F	
		Divide					
Coordinates	80°01′ S,	79°28′ S,	78°28′ S,	75°06′ S,	75° S,	77°19′ S,	
	119°31′ W	112°05′ W	106°48′ E	123°24′ E	0° E	39°40' E	
Years drilled	1966-1968 <sub>a</sub>	2006-2011 <sub>d</sub>	1990–1998,	1999-2004 <sub>i</sub>	2002-2006 <sub>k</sub>	2003-2007 <sub>n</sub>	
			$2005-2014_{f,g}$				
Surface elevation (m a.s.l.)	1530 <sub>a</sub>	1766 <sub>e</sub>	$3488_{\rm f}$	3233 <sub>i</sub>	2892 <sub>1</sub>	3810 <sub>i</sub>	
Drilled depth (m)	2193	3405 <sub>d</sub>	3769.3 <sub>g</sub>	3270.2 <sub>i</sub>	2774.2 <sub>k</sub>	3035.2 <sub>n</sub>	
Ice thickness according with	2300 <sub>b</sub>	3455 <sub>e</sub>	$3750\pm20_{g}$	3273±5 <sub>j</sub>	2750±501	3028±15 <sub>j</sub>	
radar/seismic survey (m)							
Snow accumulation at surface	169.5c	220 <sub>e</sub>	24.8 <sub>h</sub>	28.4j	70 <sub>m</sub>	29.9 <sub>j</sub>	
(mm ice $a^{-1}$ )							
Ice sheet surface horizontal	<u>12.7</u> <sub>o</sub>	<u>~3.0</u> <sub>t,u</sub>	<u>2.00±0.01s</u>	0.015±0.01 <sub>p</sub>	<u>0.74</u> <sub>r</sub>	<u>Negligible<sub>v</sub></u>	
velocity, m a <sup>-1</sup>				-			
Mean surface snow temperature	-28 <sub>a</sub>	-30 <sub>e</sub>	-57 <sub>h</sub>	-54.6 <sub>j</sub>	-44 <sub>1</sub>	-57.3 <sub>j</sub>	
(°C)							

<sup>a</sup>Ueda, 2007; <sup>b</sup>Wexler, 1961; <sup>c</sup>Gow, 1968; <sup>d</sup>Slawny et al., 2014; <sup>e</sup>WAIS Divide Project Members, 2013; <sup>f</sup>Vasiliev et al., 2011; <sup>g</sup>Lukin and Vasiliev, 2014; <sup>h</sup>Ekaykin et al., 2012; <sup>j</sup>Augustin et al., 2007; <sup>j</sup>Parrenin et al., 2007<u>a</u>; <sup>k</sup>Wilhelms et al., 2014; <sup>l</sup>Ueltzhöffer et al., 2010; <sup>m</sup>Huybrechts et al., 2007; <sup>n</sup>Motoyama, 2007<u>c</u>Whillans, 1977; <sup>p</sup>Vittuari et al., 2004; <sup>r</sup>Wesche et al., 2007; <sup>s</sup>Wendt et al., 2006; <sup>r</sup>Conway and Rasmussen, 2009; <sup>u</sup>Koutnik et al., 2016; <sup>v</sup>Motoyama et al., 2008

I

Table 2: Thermophysical properties at the base of Antarctic Ice Sheet at sites of deep ice-drilling estimated in this study
-----------------------------------------------------------------------------------------------------------------------------

Parameters	WAIS		EAIS			
	Byrd	WAIS Divide	Vostok	Dome C	Kohnen	Dome F
Temperature, °C	-1.43	-2.30	-2.49	-2.15	-1.85	-1.99
Temperature gradient (°C 100 m <sup>-1</sup> )	3.70	3.88	2.02	2.42	3.12	2.66
Ice thickness according with depth	2164	3485	3759	3257	2770	3016
of pressure melting point (m)						
Basal melt rate (mm a <sup>-1</sup> )	<u>1.2±0.8</u>	<u>3.7±1.7</u>	$-4.8\pm0.6$	1.08±0.27	<u>2.8±1.6</u>	2.5±0.5
GHF (mW m <sup>-2</sup> )	<u>88.4±7.6</u>	<u>113.3±16.9</u>	-3. <u>6</u> ±5. <u>3</u>	<u>57.9±6.4</u>	<u>86.9±16.6</u>	7 <u>8.9</u> ±5.0

Table 3: Polynomial approximations of borehole temperature T (°C) as a function of true vertical depth z and correlation factors

Drill sites	<u>Polynomial</u>	$\underline{R}^2$
Byrd	$\underline{T} = -28.343 + 0.8367 \times 10^{-3} z - 6.7651 \times 10^{-6} z^2 + 6.1339 \times 10^{-9} z^3$	<u>0.997</u>
WAIS Divide	$\underline{T} = -31.799 + 8.8595 \times 10^{-3} z - 9.4649 \times 10^{-6} z^2 + 2.657 \times 10^{-9} z^3$	<u>0.997</u>
<u>Vostok</u>	$\underline{T} = -56.034 + 2.9889 \times 10^{-3}z + 3.888 \times 10^{-6}z^2 + 0.2419 \times 10^{-9}z^3$	<u>0.999</u>
Dome C	$\underline{T} = -54.316 + 5.2978 \times 10^{-3}z + 4.4141 \times 10^{-6}z^2 - 0.368 \times 10^{-9}z^3$	<u>0.999</u>
Kohnen	$\underline{T} = -44.428 + 1.7384 \times 10^{-3} z + 4.4124 \times 10^{-6} z^2 + 0.184 \times 10^{-9} z^3$	<u>0.999</u>
Dome F	$\underline{T} = -55.016 + 5.839 \times 10^{-3} z + 5.188 \times 10^{-6} z^2 - 0.446 \times 10^{-9} z^3$	0.998

Table 4. GHF (mW m<sup>-2</sup>) calculated for different form factors m in the steady-state model, the correlation factor  $R^2$  between modeled and measured age scales and RSME

Parameters	WAIS		EAIS					
	<b>Byrd</b>	WAIS Divide	<b>Vostok</b>	Dome C	Kohnen	Dome F		
$\underline{\text{GHF for } m = 0}$	8.72	72.9	-16.16	134.6	66.04	<u>92.78</u>		
$\underline{R^2}$	<u>-0.37</u>	-2.46	0.65	0.69	<u>0.978</u>	0.78		
RMSE	28.19	36.40	58.95	107.51	4.132	40.62		
$\mathbf{GHF \ for} \ m = 0.25$	28.44	<u>100.0</u>	-35.28	105.3	<u>60.95</u>	130.9		
$\underline{R^2}$	<u>0.12</u>	<u>0.59</u>	0.50	0.72	<u>0.986</u>	-1.62		
RMSE	22.60	<u>12.93</u>	70.60	102.14	<u>2.431</u>	140.91		
$\mathbf{GHF \ for} \ m = 0.50$	<u>55.05</u>	207.4	-20.33	70.06	156.4	<u>92.96</u>		
$\underline{R^2}$	0.46	0.41	0.51	0.87	0.976	0.47		
RMSE	<u>17.59</u>	<u>15.05</u>	<u>69.74</u>	<u>69.48</u>	4.338	<u>62.91</u>		
$\underline{\text{GHF for } m = 0.75}$	<u>95.84</u>	240.3	-25.39	<u>57.40</u>	106.0	104.8		
$\underline{R^2}$	<u>0.57</u>	0.32	0.39	<u>0.997</u>	0.984	0.53		
RMSE	<u>14.86</u>	<u>16.15</u>	78.01	22.34	<u>3.477</u>	<u>59.25</u>		
$\underline{\text{GHF for } m = 1.00}$	<u>117.8</u>	251.3	<u>-8.90</u>	<u>67.3</u>	<u>161.5</u>	<u>79.20</u>		
$\underline{R^2}$	<u>0.45</u>	0.29	<u>0.75</u>	<u>0.95</u>	0.982	<u>0.83</u>		
RMSE	<u>17.83</u>	<u>16.45</u>	<u>49.71</u>	<u>43.31</u>	<u>3.692</u>	<u>35.87</u>		

GHF values with the highest correlation factor and smallest RSME are highlighted by bold.

835 Table 5: Equivalent thermophysical parameters used by GA in comparison with published data

Parameters	Byrd	WAIS Divide	<u>Vostok</u>	Dome C	Kohnen	Dome F
"Equivalent" snow accumulation at	<u>52.8</u>	48.8	<u>8.95</u>	3.87	<u>6.92</u>	3.00
surface (cm ice $a^{-1}$ )						
Modern snow accumulation at surface	16.9	22.0	2.48	2.84	7.00	2.99
$(\text{cm ice } a^{-1})$						
"Equivalent" surface temperature (°C)	<u>-29</u>	<u>-30.9</u>	<u>-56.5</u>	<u>-54.7</u>	<u>-44.9</u>	<u>-56.5</u>
Modern surface temperature (°C)	-28	-30	-57	-54.6	-44	-57.3



Figure 1: <u>GHF derived in the present study (P.S.) from basal temperature gradients in deep ice boreholes (green bars) compared with modeling.</u> Red circles show locations of deep ice drilling sites (Byrd, WAIS Divide, Vostok, Dome C, Kohnen, and Dome F)
discussed in the present study. Black squares show locations of boreholes drilled in Antarctic margins, in which borehole temperature measurements were carried out and GHF values were estimated (<u>a</u>Zagorodnov et al., 2012; <u>b</u>Nicholls and Paren, 1993; <u>c</u>Fisher et al., 2015; <u>d</u>Begeman et al., 2017; <u>c</u>Engelhardt, 2004; <u>f</u>Risk and Hochstein, 1974; <u>g</u>Decker and Bucher, 1982; <u>b</u>Clow et al., 2011; <u>i</u>Dahl-Jensen et al., 1999; <u>i</u>Zotikov, 1961). Drill sites in ocean/sub-shelf sedimentaries are not shown. <u>Location of the Antarctic ice divides is shown according with Rignot et al., 2011.
</u>



Figure 2: Smoothed measured temperature profiles in Antarctic deep ice boreholes. Pressure-melting point temperature  $T_{melt}(z)$  is shown in the assumption of Clausius-Clapeyron slope of 0.0742 K/MPa.



Figure <u>3</u>. Temperatures measured in Antarctic deep ice boreholes compared with best-fit temperature profiles for the deepest 1500 m.



Figure 4. Comparison of the measured age scales (Ahn and Brook, 2008; Bazin et al., 2013; Bereiter et al., 2012; Blunier and Brook, 2001; Kawamura et al., 2007; Neftel et al., 1988; Parrenin et al., 2007b; Sigl et al., 2016; Staffelbach et al., 1991; Veres et al., 2013) and modeled age scales with m = 0, m = 0.25, m = 0.5, m = 0.75, and m = 1.0 (correlation factors between modeled and measured age scales for each run are stated in Table 4).



Figure <u>5</u>. Estimated vertical velocities at drilling sites in West Antarctica (a) and East Antarctica (b). In East Antarctica snow accumulation and thus vertical ice velocities are far less than in West Antarctica.

![](_page_36_Figure_0.jpeg)

Figure 6. Paleo temperature profiles at WAIS Divide based on transient model (*m* = 0).