

Dear Olaf Eisen,

thank you for taking another detailed look at the manuscript.

Below please find detailed responses to some of the questions you raised, as well as a new marked-up version of the manuscript.

Dear Martin & coauthors,

thank you for your revision, in which you addressed most of the criticism raised by the reviewers. In the attached pdf I made a number of small corrections which I ask you to incorporate.

Most all comments were incorporated, although we did not quite know how you want to see the sentence on page 6 line 55ff ("To be able to disentangle these two effects,...") to be changed.

The word "timing" was indeed not very descriptive and has been substituted with "distribution of the propagation delay of individual photons". Individual photons can be time-stamped with nanosecond precision using for example photomultipliers. For details of that method please see for example doi:10.1029/2005JD006687.

On page 6, line 45 we are indeed talking about an absorption based anisotropy in contrast to the more general attenuation based anisotropy. Attenuation includes effects from both absorption and scattering. As mentioned in the manuscript, modelling the anisotropy through either absorption or scattering is badly motivated (for details see 10.18154/RWTH-2019-09941), but both approaches can result in an effective description of the effect seen in IceCube. In both cases less scattering or absorption is required for photons travelling along the flow axis to match the IceCube measurement. In the case of the dustlogger this would result in a larger return intensity in the case of the absorption (as also observed) and a smaller return signal in the case of the scattering model.

One point I would ask you to clarify is the comment and your respond on page C5 by Jan Eichler: "p.10 l.10-16: I also wonder what ..."

In principle this is correct, with a few constrains. I would appreciate if you would elaborate on this a little bit further on this in the manuscript (although you don't have to go into the details of fabric/grain development, which are not the main focus in your manuscript):

- For a girdle, the connection of mean grain elongation to be perpendicular to fabric has been described early, I think Gow. Could you add a reference? This is only true, to my understanding, for the right ratio of deformation and recrystallization, e.g. fast and/or cold.

- "assumed to be an ellipsoid" To you have any reference/data to this?

We know from private communication of yet unpublished material from SP14 that the average elongation is indeed aligned with the girdle normal vector, as one would expect from a simple deformation scenario. (This we can obviously not add to the manuscript.)

Averaging over an ensemble of randomly shaped polyhedra with an average elongation will always result in an ellipsoid. That the ellipsoid well describes the properties of the ensemble of polyhedra has for example for the surface orientation density been studied in 10.18154/RWTH-2019-09941 by comparing the ellipsoid expectation to surface orientations from crystal tessellation simulations.

In general we believe that adding the explanations given above would clutter the manuscript. From the current explanations it should be clear that both grain shape, orientation and fabric can have an effect on the propagation of light. As neither the current data, nor the current simulation offer the possibility to distinguish the different crystal properties, we prefer to leave these details for for a future paper.

Please provide a reference for your azimuth (i.e. where is  $0^\circ$  pointing to?) - I did not see that defined.

Done. The azimuth is with respect to Local Grid bearings. Grid North ( $0^\circ$ ) aligns with the Greenwich meridian.

At some point you mention "water filled borehole". I guess that is a typo for SP?

We are not quite sure what SP stands for in this context. But that is surprisingly not a typo. As part of the construction of the IceCube Neutrino Observatory (2004-2010), 86 2500 m deep, 60 cm diameter holes were drilled using hot water drilling (see doi:10.3189/2014AoG68A03). These holes took about 2 days to drill each and required two to three weeks to fully refreeze. This time not only allowed us to instrument the holes with our photosensors, but also left some time for logger deployments.

Are the data available for others, according to TC guidelines, e.g. in a data repository?

There is currently one data release available at <https://doi.org/10.15784/601222>, for which we have added a data availability section in the manuscript. It includes a single pre-processed stratigraphy. In the context of this paper it gives a good idea about the data quality, but can not be used to reproduce the anisotropy signature, as only a single stratigraphy and no heading information is provided. A data release of the full set of logs is currently being discussed, but has not yet been prepared.

# Observation of an optical anisotropy in the deep glacial ice at the geographic South Pole using a laser dust logger

Martin Rongen<sup>1</sup>, Ryan Carlton Bay<sup>2</sup>, and Summer Blot<sup>3</sup>

<sup>1</sup>RWTH Aachen University, Institute for Particle Physics III B, 52074 Aachen, Germany

<sup>2</sup>Department of Physics, University of California, Berkeley, CA, USA

<sup>3</sup>DESY, D-15738 Zeuthen, Germany

**Correspondence:** Martin Rongen (rongen@physik.rwth-aachen.de)

**Abstract.** We report on a depth-dependent observation of a directional anisotropy in the recorded intensity of back-scattered light as measured by an oriented laser dust logger. The measurement was performed in a drill hole at the geographic South Pole, about a kilometer away from the IceCube Neutrino Observatory. The drill hole remains open for access, after the SPICEcore collaboration had retrieved a 1751 m ice core. We find the ~~measured~~ anisotropy axis of  $126 \pm 3^\circ$  as measured below 1100 m to be compatible with the local flow direction. The observation is discussed in comparison to a similar anisotropy observed in data from the IceCube Neutrino Observatory and favours a birefringence based scenario over previously suggested Mie scattering based explanations. In the future, the measurement principle, when combined with a full-chain simulation, may have the potential to provide a continuous record of fabric properties along the entire depth of a drill hole.

## 1 Introduction

The viscosity of an individual ice crystal strongly depends on the direction of the applied strain. As a hexagonal crystal, ice will most readily deform as shear is applied orthogonal to the c-axis, which leads to slip of the basal planes (Petrenko and Whitworth (2002)). Thus, individual grains elongate, with the major axis being aligned perpendicular to the c-axis.

In large-scale systems, such as glaciers or ice sheets, ice is compressed under its own weight and as a result flows away from the accumulation region. This leads to preferential c-axis orientations (see Alley, 1988), most commonly girdle fabrics, where c-axis are predominantly found on a

plane, with the plane's normal vector being aligned with the flow direction. Ice fabric can not only be observed through macroscopic imaging of ice cores (Weikusat et al., 2016), but also leads to a directionality in the propagation of mechanical and electromagnetic radiation, in principle allowing for remote-sensing of the ice fabric.

The mechanical anisotropy of ice means that the speed of sound depends on the fabric realization. This has for example been derived and measured by Kluskiewicz et al. (2017). Ice crystals are also a birefringent material, with any incoming electromagnetic radiation being separated into an ordinary and extra-ordinary ray of perpendicular polarization with respect to the c-axis, and which propagate with different refractive indices. This is classically observed as a direction-dependent delay in the propagation of radio waves, as for example described by Fujita et al. (2006).

Recently, as part of ice calibration measurements for the IceCube Neutrino Observatory (Aartsen et al., 2017), Chirkin (2013) described the observation of an optical anisotropy, where about twice as much light is observed along the glacial flow axis versus orthogonal to the flow axis, at a receiver 125 m away from an isotropic emitter. The effect was originally modelled as a direction dependent modification to Mie scattering quantities, either through a modification of the scattering function as proposed by Chirkin (2013) or through the introduction of a direction dependent absorption as introduced by Rongen (2019). As also shown by Rongen (2019), both parameterizations lack a thorough theoretical justification and resulted in an incomplete description of the IceCube data.

As the wavelength of  $\sim 400$  nm employed in the IceCube studies is significantly smaller than the average grain size, the effect is challenging to derive from first principles. First attempts have been made by Chirkin and Rongen (2019) by attributing the effect to the cumulative diffusion that a light beam experiences as it is refracted or reflected on many grain boundary crossings in a birefringent polycrystal with a preferential c-axis distribution.

In this scenario the diffusion is found to be strongest when photons initially propagate along the flow and smallest when initially propagating orthogonal to the flow. In addition photons are, on average, deflected towards the flow axis.

The deflection per unit distance increases for stronger girdle fabrics, a larger average crystal elongation or a smaller average crystal size. For crystal realizations where the deflection outweighs the additional diffusion along the flow axis compared to the diffusion along the orthogonal direction, the photon flux along the flow axis will increase with distance compared to the photon flux along the orthogonal axis.


We add to the body of anisotropy observations by providing the first direction-dependent measurement of the intensity of back-scattered, optical light returning to the oriented dust logger deployed down a glacial bore hole. If the anisotropy is caused by Mie scattering a reduced return signal is expected when the light source points along the flow, while more light is expected to return in case of the birefringence and absorption explanations.

## 2 The oriented dust logger

The dust logger, as sketched in Figure 1 and introduced by Bramall et al. (2005), consists of a 404 nm laser line source, emitting a 2 mm thin, horizontal fan of light about  $60^\circ$  across. A small fraction ( $10^{-10}$  to  $10^{-6}$ ) of all emitted photons is back-scattered or reflected and returns to the bottom section of the dust logger where a 1" Hamamatsu photon-counter module is located.

Scattering and absorption on air bubbles, soot and other impurities as described by Mie scattering theory is traditionally thought to be the dominant contribution to the return signal. However, taking into consideration the findings of Chirkin and Rongen (2019), diffusion on grain boundaries may also contribute non-negligibly to the signal.

The intensity of the light source can be adjusted throughout the logging process. To avoid stray light contamination from reflections on the hole-ice interface, multiple sets of black nylon baffles are attached to the side of the pressure housing. These also sweep ice crystals and debris out of



**Figure 1.** Sketch of the laser dust logger. Light emitted by a 404 nm diode laser in a  $60^\circ$  horizontal fan can only reach the photon counter below through scattering on nearby impurities.

the source beam. Spring-loaded calipers keep the logger centered in the hole.

The depth of the logger is monitored through the cable payout and on-board pressure sensors. During offline analysis multiple logs from the same site are further aligned to achieve centimeter depth precision, using characteristic features such as volcanic [lineshorizons in the ice](#), as described by Aartsen et al. (2013).

This device has previously been deployed in West Antarctica, East Antarctica and Greenland. Due to excellent imaging properties, deployments down the water-filled drill holes of the IceCube Neutrino Observatory resulted in one of the highest resolution particulate stratigraphies of any glacier available to-date, as described by Aartsen et al. (2013).


To measure a potential directionality of the return signal relative to the direction of the emitted light fan, an optional extension consisting of an Applied Physics Systems Model 547 Directional Sensor<sup>1</sup> has been fitted to the top of the logger. By measuring the local magnetic field it deduces the absolute orientation with an azimuthal accuracy of  $\pm 1.2^\circ$  for magnetic latitudes  $< \pm 40^\circ$ . For our application at the geographic South Pole we estimate the azimuthal accuracy to be  $\pm 3^\circ$ .

<sup>1</sup>[https://www.appliedphysics.com/\\_main\\_site/wp-content/uploads/model-547-micro-orientation-sensor.pdf](https://www.appliedphysics.com/_main_site/wp-content/uploads/model-547-micro-orientation-sensor.pdf)

### 3 SPICEcore deployments

The South Pole Ice Core, SPC14 (see Casey et al., 2014), was drilled by the SPICEcore project in 2014–2016 at a location 2.7 km from the Amundsen–Scott station, using the Intermediate Depth Drill designed and deployed by the U.S. Ice Drilling Program (IDP) (Johnson et al., 2014). It reached a final depth of 1751 m (Winski et al., 2019), surpassing the original 1500 m goal.

10 The core has been retrieved in 2 m segments with a diameter of 98 mm. The resulting 126 mm diameter drill hole was filled with the non-freezing drilling fluid Estisol-140 and has been preserved for future logging access.



**Figure 2.** Depth development of c-axis distributions (Lambert azimuthal equal-area projections) measured in SPICEcore. Depths (in meters) are indicated atop each diagram. They show a clean girdle fabric below ~1000m. Adapted from: Voigt (2017)

15 Unlike most ice coring sites, the Geographic South Pole is not near an ice divide, but rather on a flank site with a local flow velocity of 10 m per year. The associated accumulation site for the deepest ice is believed by Lilien et al. (2018) to be Titan Dome (Lilien et al. (2018)), meaning that the ice has been transported as far as 200 km. The experienced strain stress experienced below ~800 m depth has resulted in a very prominent and continuously strengthening girdle fabric as measured by Voigt (2017). Figure 2 shows example c-axis distributions from the SPC14 ice core at various depths.

The oriented dust logger has been deployed down the SPICEcore hole twice during the 162016/17–2017 season, both times using the Intermediate Depth Logging Winch provided by the IDP. Due to a limited available cable length, it was only able to reach a depth of 1577 m of the 1751 m cored. During the first log the laser intensity was not yet optimized, leading to saturated and thus unusable data above 1000 m.


Two further deployments down to ~1700 m were performed during the 182018/19–2019 season. Due to mechanical problems with the winch cable payout, depth readings from the winch itself are inaccurate. Only one deployment could be depth aligned to the required precision using characteristic features as previously discussed. This deployment includes an additional round-trip between 1354-1703 m. Table 1 summarizes the properties of the logs

Log	Depth [m]	Note
Down1	130 - 1577	Saturated above 1000 m
Up1	1577 - 130	Saturated above 1000 m
Down2	130 - 1580	-
Up2	1580 - 130	-
Down3-Leg1	130 - 1703	-
Up3-Leg1	1703 - 1354	-
Down3-Leg2	1354 - 1704	Near identical orientations to Down3-Leg1
Up3-Leg2	1704 - 130	-

**Table 1.** Summary of usable logs obtained within the 162016/17–2017 and 182018/19–2019 logging seasons.

used for the measurements presented here.

As shown in Fig.3, the logger rotates as it descends and ascends the hole, mainly due to the residual twist in the logging cable. On ascent, as the cable is pulling the tool up, it undergoes a smooth rotation of slightly varying angular velocity. On descent the logger sinks under its own weight and the rotation is not continuous. The most likely explanation is that the logger is repeatedly stuck on the wall before slipping.



**Figure 3.** Rotation of the tool descending and ascending the bore hole. A smooth and continuous rotation of slightly varying angular velocity is observed when ascending. During descent the rotational movement is erratic. (The azimuthal orientation is with respect to Local Grid bearings. Grid North (0°) aligns with the Greenwich meridian.)


### 4 Anisotropy signature

The data obtained from the oriented dust logger consists of orientation, depth and optical return signal measurements at 10 ms intervals. At the usual deployment speed this is

equivalent to a sampling distance of  $\sim 2.5$  mm.

While the photomultiplier is located  $\sim 850$  mm below the laser light source, the depth resolution, as measured by Bramall et al. (2005) as the smearing of an ash layer, is dominated by the vertical extent of the laser beam and is ~~no worse~~ less than a few mm. This allows for a continuous record of optical properties down the entire depth of a drill hole at the vertical resolution of less than a year of deposition, assuming an annual layer thickness of 1-2 cm in the deep ice as reported by Aartsen et al. (2013).

Above the transition region of air bubbles to clathrate hydrate (Miller, 1969) at 700-1300 m the return signal is dominated by scattering on air bubbles. Below, the return signal is primarily proportional to the concentration of impurities contributing to scattering. The resulting high resolution stratigraphy is exemplified in Fig. 4.




**Figure 4.** Example detail of optical stratigraphies obtained by logging the same depth range three times.

This figure also shows that the optical return signals at the same depths are not consistent between logs. Instead the signal depends on the absolute orientation of the logger. We extract this anisotropy signature by taking the ratio of two logs. Usually, the ratios of raw data are on average non-unity and show slow, continuous variations. These systematic offsets, caused for example by the changing clarity of the drilling fluid or grime accumulation on the logger, are corrected using a second-degree polynomial fitted to the ratio.

Example ratios for  $\sim 100$  m depth slices and after fitting and correcting the offset for each depth slice are given in Fig. 5. When the device's orientation between logs is out of phase, pointing in different directions at the same depth, the ratio in these examples becomes as large as 1.5. When the logs are in phase, the observed intensities are equal and thus

the ratio is unity.


Analysis of the 18/19 logging season reveals that two logs, *Down3-Leg1* and *Down3-Leg2*, exhibit strongly correlated orientations. As exemplified in Fig. 6, the orientations of the two descending segments in the depth range between 1354 m and 1703 m show near identical orientations, suggesting that the rotation of the tool may have been governed by the hole geometry itself. As a result, the ratio of the two logs is consistent with unity where the logs align and the spread of 2% standard deviation indicates the typical short term intensity fluctuations seen in the measurement.



**Figure 5.** Example fits to the intensity ratio fits-in  $\sim 100$  m depth slices. The red and black dotted lines denote the orientations of the two used logs. (The orientation is defined as the cosine of the azimuth angle.) Blue is the intensity ratio. Green is the fitted intensity ratio using eq. (1).

In the following analysis, log *Down3-Leg2* is excluded from the analysis as it is fully correlated with *Down3-Leg1*. The number of remaining usable ratios at each depth range available from  $N$  logs is given by the binomial coefficient





**Figure 6.** The intensity ratio of two log segments with near identical orientations is unity and shows the typical spread of the data.

$\binom{N}{2}$  and varies between 3 and 21.

Lacking a mature first-principle explanation and simulation of the anisotropy effect and experimental setup, the ratios have instead been found to be well-described by the following empirical relationship:

$$\text{intensity ratio} = \frac{1 + a \cdot \cos(2 \cdot (\alpha_1 - \phi))}{1 + a \cdot \cos(2 \cdot (\alpha_2 - \phi))}. \quad (1)$$

Here  $\alpha_1$  and  $\alpha_2$  denote the azimuthal orientations of the logger during the two logs.  $\phi$  denotes the azimuthal phase angle of the anisotropy effect, also called anisotropy axis and is limited to  $0^\circ - 180^\circ - 0^\circ - 180^\circ$ .  $a$  is a measure of the strength of the observed effect.

The orientations and the intensity ratio are given by the dust logger data. The free parameters  $a$  and  $\phi$  can be determined by fitting eq. +(1) to the data from a given depth range.

Note that the chosen relationship, while being very robust and easy to extract, implicitly assumes that the anisotropy causes a relative modulation to the total signal. In case the signal is additive on top of a contribution caused by Mie scattering on impurities, as would be expected from a fabric driven birefringence scenario, the derived strength parameter  $a$  can not directly be interpreted as the strength of the underlying effect. For example, assuming an overall constant return signal from the anisotropy, the strength parameter  $a$  would be seen to increase as the overall return signal decreases as a function of depth.

## 5 Depth evolution


To study the depth evolution of the anisotropy signature the data is binned into 100m slices. While this only allows for a rather coarse depth resolution, it ensures that at least a few rotations are seen in each ratio. Otherwise the correction polynomial could bias the signal introduced by the anisotropy and the fit would not be able to reliably determine the phase and amplitude of each log. The systematic shift introduced by the correction polynomial was further accessed by varying its degree between 1 and 3 and was found to be below the error on the mean, which is introduced below, for all depth-slices.

In the future, a finer depth resolution may experimentally be achieved by not relying on the natural rotation of the logger as it is deployed but by artificially inducing a fast rotation, or by including several azimuthally offset light sources.


The depth evolution of the fitted anisotropy axis  $\phi$  and strength  $a$  as a function of depth are seen in Fig. 7. The spread of these fitted quantities between different ratios far outweighs the statistical error of the fit of each ratio. The errors on the means of each depth bin are thus constructed from the standard deviation of all ratios. While  $\binom{N}{2}$  ratios can be constructed, only  $N - 1$  are statistically independent. The error on each mean is thus given as  $\sigma_{\text{mean}} = \sigma / \sqrt{N - 1}$ .

Individual ratios are seen to yield consistent anisotropy axes in the deep ice. In the shallow ice above  $\sim 1100$  m, where the mean strength of the observed anisotropy signal vanishes, the phase angle is unconstrained. The average axis in Local Grid bearings (Grid North ( $0^\circ$ ) aligns with the Greenwich meridian) in the deep ice is  $126 \pm 1$  (stat)  $^\circ - 126 \pm 1$  (statistical)  $^\circ$ . Considering the  $3^\circ$  systematic uncertainty of the orientation sensor, this direction is in good agreement with the local ice flow direction as measured by Lilien et al. (2018) as well as the optical axis of  $126^\circ$  or  $130^\circ$  as fitted by Chirkin (2013) and Rongen (2019) respectively, both using IceCube data.

No While no anisotropy signatures are seen above 1100 m, with the observed strength parameter  $a$  continuously increasing in the deeper ice. It is currently unclear which fraction of the increase in observed anisotropy strength versus depth is caused by the continuously stronger girdle fabric or by the decrease in overall scattering. However, no anisotropy signal is observed at 1000m where the girdle fabric is already clearly developed (see Fig. 2) but bubbles still dominate the scattering at this depth. Therefore, we suspect that the anisotropy signature is smeared out to some extent due strong local diffusion.



(a) Anisotropy strength. An example log is superimposed as reference.



(b) Anisotropy axis with respect to Local Grid bearings. Grid North ( $0^\circ$ ) aligns with the Greenwich meridian.

**Figure 7.** Depth-dependence of fitted quantities. Gray dots denote values obtained from individual ratios. The blue markers indicate the average at each depth.

## 6 Conclusions

We have presented the first direction-dependent measurement of the intensity of back-scattered, optical light in deep glacial ice. The measurement has been performed using an oriented dust logger deployed down the SPICEcore drill hole.

Below  $\sim 1100$  m a consistent increase in received intensity is observed when the laser is aligned with the local flow axis.

This is consistent with the birefringence explanation offered by Chirkin and Rongen (2019) where more diffusion is observed along the flow, thus leading to a higher return intensity and inconsistent with the previous explanation given by Chirkin (2013) where the effect was attributed to reduced Mie scattering along this flow. The observed sign is qualitatively also consistent with an absorption-based

anisotropy anisotropy based on absorption.

The amplitude of the intensity modulation increases with depth. This is in part likely caused by the strengthening of the girdle fabric as well as the strong reduction in overall scattering, as bubbles are transformed to clathrate hydrate hydrates.

To be able to disentangle these two effects, and to potentially transition from the presented experimental ratios to a quantitative measurement of fabric properties, will require a full photon propagation simulation incorporating both Mie scattering on impurities as well as the diffusion introduced through the polycrystalline, birefringent fabric. While the basics for such a simulation have been outlined by Chirkin and Rongen (2019), more work will be required for the simulation to be efficient enough to be used in-for this application. It is currently also unclear if the intensity ratio alone will be sufficient to constrain the different fabric properties, namely the Woodcock parameters and the average grain size and shape, or if more information such as timing the distribution of propagation delays of individual photons may be required.

*Data availability.* A single stratigraphy is currently available from <https://doi.org/10.15784/601222> (Bay, R. (2019)). The full set of logs may be released in the future.

*Competing interests.* The authors declare that they have no competing interests (both financial or non-financial).

*Author contributions.* RB designed the dust logger. The logging was carried out by RB and SB. MR and RB developed the data processing and analysis. The manuscript was prepared by MR with contributions from all co-authors.

*Acknowledgements.* The authors would like to thank the SPICEcore collaboration for providing the borehole, the U.S. Ice Drilling Program, the Antarctic Support Contractor and the NSF National Science Foundation for providing the equipment to perform the described measurement and for their support at South Pole. This work has been achieved under the NSF grant #1443566 and was in part supported by BMBF, Verbundforschung. We would also like to thank the IceCube collaboration for supporting the 18/19 logging activities.

## References

Aartsen, M., Abbasi, R., et al.: South Pole glacial climate reconstruction from multi-borehole laser particulate stratigraphy, *Journal of Glaciology*, 59, 1117–1128, 2013.



- Aartsen, M., Ackermann, M., Adams, J., et al.: The IceCube Neutrino Observatory: instrumentation and online systems, *Journal of Instrumentation*, 12, P03 012, <http://stacks.iop.org/1748-0221/12/i=03/a=P03012>, 2017.
- 5 Alley, R. B.: Fabrics in Polar Ice Sheets: Development and Prediction, *Science*, 240, 493–495, <https://doi.org/10.1126/science.240.4851.493>, <https://science.sciencemag.org/content/240/4851/493>, 1988.
- Bramall, N. E. et al.: A deep high-resolution optical log of dust, ash, and stratigraphy in South Pole glacial ice, *Geophys. Res. Lett.*, 32, <https://doi.org/10.1029/2005GL024236>, 2005.
- Casey, K., Fudge, T., Neumann, T., Steig, E., Cavitte, M., and Blankenship, D.: The 1500 m South Pole ice core: recovering a 40 ka environmental record, *Annals of Glaciology*, 55, 137–146, <https://doi.org/10.3189/2014aog68a016>, 2014.
- 15 Chirkin, D.: Evidence of optical anisotropy of the South Pole ice, in: Proceedings, 33rd International Cosmic Ray Conference (ICRC2013): Rio de Janeiro, Brazil, July 2-9, 2013, p. 0580, <http://inspirehep.net/record/1412998/files/icrc2013-0580.pdf>, 2013.
- 20 Chirkin, D. and Rongen, M.: Light diffusion in birefringent polycrystals and the IceCube ice anisotropy, in: 36th International Cosmic Ray Conference (ICRC 2019) Madison, Wisconsin, USA, July 24-August 1, 2019, 2019.
- Fujita, S., Maeno, H., and Matsuoka, K.: Radio-wave depolarization and scattering within ice sheets: a matrix-based model to link radar and ice-core measurements and its application, *Journal of Glaciology*, 52, 407–424, <https://doi.org/10.3189/172756506781828548>, 2006.
- Johnson, J. A., Shturmakov, A. J., Kuhl, T. W., Mortensen, N. B., and Gibson, C. J.: Next generation of an intermediate depth drill, *Annals of Glaciology*, 55, 27–33, <https://doi.org/10.3189/2014aog68a011>, 2014.
- 30 Kluskiewicz, D., Waddington, E. D., et al.: Sonic methods for measuring crystal orientation fabric in ice, and results from the West Antarctic ice sheet (WAIS) Divide, *Journal of Glaciology*, 63, 603–617, <https://doi.org/10.1017/jog.2017.20>, 2017.
- Lilien, D. A., Fudge, T. J., Koutnik, M. R., et al.: Holocene Ice-Flow Speedup in the Vicinity of the South Pole, *Geophysical Research Letters*, 45, 6557–6565, <https://doi.org/10.1029/2018gl078253>, <https://doi.org/10.1029/2018gl078253>, 2018.
- 40 Miller, S. L.: Clathrate Hydrates of Air in Antarctic Ice, *Science*, 165, 489–490, <https://doi.org/10.1126/science.165.3892.489>, 1969.
- Petrenko, V. F. and Whitworth, R. W.: *Physics of Ice*, Oxford University Press, <https://doi.org/10.1093/acprof:oso/9780198518945.001.0001>, 2002.
- Rongen, M.: Calibration of the IceCube Neutrino Observatory, Ph.D. thesis, RWTH Aachen University, <https://doi.org/10.18154/RWTH-2019-09941>, 2019.
- 50 Voigt, D. E.: c-Axis Fabric of the South Pole Ice Core, SPC14, <https://doi.org/10.15784/601057>, <http://www.usap-dc.org/view/dataset/601057>, 2017.
- Weikusat, I., Jansen, D., et al.: Physical analysis of an Antarctic ice core—towards an integration of micro- and macrodynamics of polar ice, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 375, 20150 347, <https://doi.org/10.1098/rsta.2015.0347>, 2016.
- Winski, D. A., Fudge, T. J., Ferris, D. G., et al.: The SP19 chronology for the South Pole Ice Core – Part 1: volcanic matching and annual layer counting, *Climate of the Past*, 15, 1793–1808, <https://doi.org/10.5194/cp-15-1793-2019>, 2019.