

Response to Anonymous Referee #1

We would like to thank the reviewer for the detailed and useful comments, which have helped to improve the quality and readability of our manuscript. In the following, we provide a reply to the points discussed by the reviewer as well as the changes in the manuscript.

The comments of the reviewer are written in **bold**, the extracts of the manuscript in *italics* with changes highlighted in *blue*.

General comments:

This paper presents a four-year measurement series of near rock surface temperatures at coastal cliffs or canyon walls close to the Kongsfjord in the Ny Ålesund region, Svalbard and uses the well-known model CryoGrid 3 to calculate the energy balance at the observation sites and discusses the results. The authors show that the coastal sites are in general warmer than the sites a little bit further away from the fjord. This paper is carefully elaborated and shows interesting data and model results, which is an important contribution to a better understanding of the surface energy balance and related heat fluxes at a place where until now not many studies have been carried out in steep cliffs closely situated to a fjord. In general, before publishing this valuable article some major changes should be considered, which are outlined below:

We appreciate that the reviewer identifies the relevance of our work and states it as an important contribution to a better understanding of thermal regimes in high Arctic rock walls. We understand the concerns of the reviewer with respect to neglecting the influence of snow, the assumption of a neutral atmospheric stratification and other points. We made changes to our manuscript including new model runs and clarifications in the text. Therefore, we are now confident that the revised manuscript is an improved version of our study and suitable for publication.

Line 167, Neglecting influence of snow:

It is somehow understandable that the influence of snow is neglected for the investigated sites as the cliffs are steep and the influence of snow seems according to the authors until now not to be very important. However, there are two important points, which the authors do not consider enough in their paper, although they discuss the influence of snow in the paragraph of the discussing section on line 421 but this discussion cannot be used to justify their main approach of neglecting snow completely in their model approach.

- Firstly, recent studies show clearly that even in steep rock walls, small or even larger amounts of snow can accumulate on rock ledges and influence rock wall temperatures and change heat fluxes considerable (e.g. Haberkorn et al. 2015, 2016 and 2017) and their logger RW01 and their comment in the caption of table 5 shows that snow seems already today have some influence at certain cliff sites!

- Secondly, performing climate scenarios coupled to model runs of CryoGrid 3 not taking into account snow for future conditions is not at all reasonable as in the future probably snow cover is may going to

increase considerable during winter time particularly at sites which are close to a large moisture source (North Atlantic Ocean).

Therefore, I suggest to a) include in CryoGrid 3 a scenario including snow, which should not be a big problem as CryoGrid 3 is already well prepared and b) omit the future scenario runs in this paper (mainly chapter 4.5 and 5.4), as this part does not add any important additional information or only results, which are extremely speculative as the authors admit themselves on line 474. Particularly as the authors cannot really convincingly show and explain that snow is not going to play an important role in the future. However, a future snow cover at the small cliff sites can strongly change the whole thermal regime (e.g. the canyon site could be filled by more snow and corresponding snow drift by wind in the future).

We agree with the reviewer that snow cover can play a significant role in the thermal regime of rock walls. To account for this, we included a discussion about snow cover with additional model runs and explanations. The model runs were performed by following the approach of Magnin et al. (2017) and taking the snowfall data of the AROME-Arctic dataset as an input. We present one simulation with maximal snow height (the snowfall accumulates completely in the rock wall) and another run with a maximum height of 20 cm snow cover.

The simulations result in an overestimation of snow cover in early winter and an underestimation in late spring when comparing to the snow-covered RW01 logger. This can be explained by the fact that snow cover in the rock walls is mainly due to snowdrift and a buildup from the foot of the rock wall, until it covers the temperature logger. The buildup of snow needs time so that the snow cover is first overestimated. In late spring, the snowdrift needs a long time to melt, so that the snow cover is then underestimated. Our model is able to represent snow cover, which results from snowfall, however, the complex buildup by snowdrift cannot be simulated adequately. Therefore, we decided to exclude snow cover in our main study and focus on generally snow-free sites, which is the case for the majority of the measurement sites. Yet, we agree with the reviewer that this a very important point to discuss and included the snow cover in a comprehensive supplement as well as a short discussion in the main manuscript.

Please find the changes in our manuscript below:

Line 188-189: Further evaluations on the possible influence of a snow cover are given in the supplement, where we present model runs considering snow cover in the rock walls following the model approach of Magnin et al. (2017).

Line 332-334: Simulations with snowfall provided by the forcing data cannot represent the temporarily occurring snow cover in the rock walls adequately. A distinct overestimation of snow cover in early winter and a clear underestimation in late spring result in significant deviations from the measured data. Results of simulations including a snow cover are provided in the supplement.

Line 448-449: Model runs in the supplement show that the temporarily occurring snow cover in the rock walls poorly correlates with snowfall, but can mainly be traced back to snowdrift and in some cases the buildup of snow from the foot of the rock wall.

A detailed discussion about snow cover is provided in a supplement to the manuscript:

Supplement, Line 35-85: **Supplement 2: Simulations considering snow cover**

The measured RST data indicate temporarily occurring snow cover in some rock walls. As RW01 shows the most pronounced and regularly occurring dampening of the temperature signal, additional model runs were performed for this rock wall. Figure S2.1 displays a comparison of measured RST in RW01, a model run without snow cover (no snow scenario), a model run taking the maximum snowfall into account (snow scenario, max) and a model run restricting snow depth to a maximum of 20 cm (snow scenario, 20 cm) to account for the limited accumulation capacity of the steep rock wall. The simulations considering the snow cover were performed with CryoGrid 3, following the model approach of Magnin et al. (2017).

In January to mid of February, both snow scenarios overestimate the snow cover, leading to up to 10 °C higher RST, while the no snow scenario matches the measured data significantly better. This leads to the assumption that RW01 was not covered by snow in early winter despite the occurring snowfall. In March and April, both snow scenarios represent the measured data better than the no snow scenario. This is especially true for the snow scenario, max, which implies a thicker snow cover than expected for the vertical rock wall. In May and June, the measured data in RW01 still indicates the presence of a snow cover, while both snow scenarios underestimate the influence of snow in the rock wall. Consequently, the measurements suggest that the rock wall RW01 had a significantly thicker snow cover than calculated from the snowfall produced by the forcing data.

This leads to the conclusion that our model is able to represent snow cover and calculate a dampening of the temperature signal in general. However, the snowfall provided by the forcing data leads to high errors and cannot represent the conditions under snow cover in the rock wall. This can be explained by the fact that the snow cover in the rock wall does not directly correlate with the snowfall of the forcing data. Instead, it is likely that a snowdrift builds up from the foot of the rock wall in winter until it eventually covers the temperature logger. Therefore, both snow scenarios overestimate the snow cover in early winter, when the snow at the foot of the rock wall does not reach the temperature logger yet, and underestimates the snow cover in late spring and early summer, when the massive snow drift needs more time for melting than. As our model can only represent snow cover produced by the snowfall of the forcing data, but not the effect of snowdrift in complex terrain, we excluded periods with snow cover from the analysis in this study and focussed on measurement sites, which were found to be largely snow-free.




Figure S2.1: Comparison of measured daily RST in RW01 to a simulation without snow cover (no snow scenario), a model run taking the maximum snowfall into account (snow scenario, max) and a model run limiting snow depths to a maximum of 20 cm (snow scenario, 20 cm). The no snow scenario shows no dampening of the temperature signal, while both snow scenarios represent the snow cover well in late winter, but show an overestimation in early winter and an underestimation in late spring.

Future snow conditions will be characterized by a reduction of days with snow cover all over Svalbard. The snow water equivalent will be highly dependent on the emission scenario: Simulations of RCP4.5 show that snow water equivalent will be the same or slightly more, while simulations of RCP8.5 indicate a reduced maximum snow water equivalent of 50 % or more (Hanssen-Bauer et al., 2019). While the development of a snow cover in the analysed rock walls and its influence on RST is highly uncertain, we compare two future runs for RCP4.5 with no snow accumulation in the rock wall and snow accumulation to a maximum of 20 cm in the rock wall, respectively (Fig. S2.2). In the simulations, the snow cover reduces MARST by 0.25 °C to 1.21 °C. The lowering of MARST by the snow cover is due to the albedo increase in spring, which cools the surface by reflecting short-wave radiation. However, these results must be regarded as a rough approximation, as snowdrift and not snowfall seems to be the main driver of snow cover in these rock walls. As in most cases no significant snow cover could be detected in our measurement data, MARST might be closer to the scenario neglecting the snow cover, but might be reduced at sites where snowdrifts influence RST.




Figure S2.2: MARST of past and future simulations of the near-coastal scenario with RCP4.5 comparing two model runs neglecting the snow cover and taking potential snow cover up to 20 cm into account.

Line 121, assumption neutral atmospheric stratification:

It is understandable that the authors try to simplify the very complex processes of the turbulent fluxes. However, assuming that the conditions at the cliff walls could be approached by using for all cases neutral atmospheric stratification is not justified. This assumption would mean that an air parcel

moving close to the cliff would always have the same temperature (and density) as the surroundings at this position. This seems to be justified only for certain conditions during the year. The authors also justify in the discussion section their approach by tuning the roughness length until the model fits the observed values. This is reasonable to do. However, their estimated roughness length is, at least after my knowledge of such values and having seen the pictures of the measurement sites in the paper, at least one order of magnitude smaller what should be probably taken as roughness length for these specific observation sites.

We agree that the assumption of a neutral atmospheric stratification cannot represent the conditions at the rock walls throughout the year. However, fitting the roughness length is a pragmatic approach as used by Magnin et al. (2017) and as the reviewer states. We estimated the roughness length to 0.018 m. As a general rule of thumb, the roughness length should be selected as 1/10 of the “roughness elements” of the surface. Given the strong variability of possible roughness elements (small-scale heterogeneity on the order of centimeters to decimeters, larger-scale roughness of protruding wall parts on the scale of meters), it is not possible to exactly determine a roughness length with this concept, and we resorted to fitting an “effective roughness length”. However, we note that our chosen value is within the (admittedly large) range of values that are in agreement with the rule of thumb.

To clarify this point in our manuscript, we made the following changes:

Line 200-204: An important fitting parameter was the roughness length z_0 as performed in Magnin et al. (2017). We set it to a value of 0.018 m, which represents roughly 1/10 of the height of the surface roughness elements. This fitted well to the [small-scale variations on the rugged rock surface characterized by joint systems \(Fig. 2\)](#), [but uncertainties regarding the different spatial scale of roughness elements in the rock walls remain.](#)

Line 429-432: A critical point is the assumption of a neutral atmospheric stratification perpendicular to the vertical rock wall, which must be regarded as a first order approximation [as it does not account for the complex wind field and boundary layer conditions near the rock wall.](#) This leads to uncertainties in the near surface turbulent exchange of the vertical wall as micro-topography and changing weather conditions can influence the movement of air parcels.

Line 107, table 1 setting of surface temperature loggers:

The temperature loggers are located all at expositions of NE (except two of them ENE and N). It would be interesting to see the influence of the different expositions, which could be easily modelled by CryoGrid 3. I would assume that at this latitude the expositions do not play a very important role but it would be an interesting question which could be answered by CryoGrid 3.

We thank the reviewer for this suggestion and included it in our revised manuscript. To do so, we ran simulations for north- and south-facing rock walls for the open fjord scenario and the non-coastal scenario. Both show variations up to 0.7 °C, which is substantially less than in lower latitudes like in the European Alps. In our manuscript, we changed the following:

Line 327-331: The model results show differences in MARST according to the exposition of the rock wall: In the open fjord scenario, the lowest MARST in 2017/18 is found on the north-facing rock wall RW05 (-

0.9°C), while the highest MARST is calculated for RW06 facing east-north-east (-0.3 °C). Model runs for north- and south-facing rock walls suggest that differences in MARST due to exposition are only 0.7 °C or less. While no effect is detectable in winter, higher RST variations up to 1.6 °C are calculated for the spring season.

Line 442-447: Svalbard lies in the continuous permafrost zone (Brown et al., 1997; Obu et al., 2019) and deep permafrost is observed in the area, e.g. within the abandoned mine shafts (Liestøl, 1977) and in boreholes (Christiansen et al., 2010). We found that the exposition of the rock wall only leads to small differences in MARST. In the winter season, polar night conditions suppress any dependence on exposition, while the effect is most pronounced in spring season for low sun angles. In general, the influence of exposition in the high Arctic is small compared to sites at lower latitudes, like the European Alps (Gruber et al., 2004b; Magnin et al., 2015; Noetzli and Gruber, 2009).

Line 477, coastal cliffs in the high Arctic – a future geohazard:

This chapter does not add any important new information to the main topic of the paper. Please delete this section.

We follow the suggestion of the reviewer and deleted the chapter 5.5. Instead, we included a significantly shorter paragraph in 5.4 to account for the main message of the deleted chapter.

Line 516-522: *Our model results indicate a significant warming of permafrost temperatures and a deepening of ALT in the 21st century, a trend that can lead to destabilization of rock slopes (Krautblatter et al., 2013). Besides, loss of sea ice and correlated longer duration of open-water season can enhance coastal erosion (Barnhart et al., 2014). In this study, the thermal regime of relatively low coastal cliffs is investigated. Indeed, similar processes can also affect much higher cliffs. Failures of coastal rock slopes can impact the water body and trigger displacement waves along shorelines, as happened in Paatuut / Greenland in 2000 (Dahl-Jensen et al., 2004; Hermanns et al., 2006). Due to permafrost degradation, rock slope failures in the high Arctic might become more likely in future, which should be taken into account for risk assessment of settlements and infrastructure.*

Specific comments:

1. Line 31: better use: warming of atmosphere than warming of climate. The atmosphere can warm but the climate can only change but not warm.

Fair enough! We have changed that sentence:

Line 34: As a response to *climate change*, degradation of permafrost...

2. Line 46: may add the new literature from Etzelmüller et al. 2020; Etzelmüller, B., Guglielmin, M., Hauck, C., Hilbich, C., Hoelzle, M., Isaksen, K., Noetzli, J., Oliva, M., Ramos, M. (2020) Twenty years of European Mountain Permafrost Dynamics – the PACE Legacy. Environmental Research Letters 15, 14.

We added the suggested literature in the manuscript.

Line 48-50: The climatic changes are also apparent in permafrost temperatures on Svalbard as observed in boreholes over the last decades (Boike et al., 2018; Christiansen et al., 2010; Etzelmüller et al., 2020; Isaksen et al., 2007).

3. Line 50: May add some more literature here such as: Gislén, K., Westermann, S., Schuler, T.V., Melvold, K., Etzelmüller, B. (2016) Small-scale variation of snow in a regional permafrost model. The Cryosphere 10, 1201-1215. Gislén, K., Westermann, S., Schuler, T.V., Litherland, T., Isaksen, K., Boike, J., Etzelmüller, B. (2014) A statistical approach to represent small-scale variability of permafrost temperatures due to snow cover. The Cryosphere 8, 2063-2074. Haberkorn, A., Wever, N., Hoelzle, M., Phillips, M., Kenner, R., Bavay, M., Lehning, M. (2017) Distributed snow and rock temperature modelling in steep rock walls using Alpine3D. The Cryosphere 11, 585-607.

We added the suggested literature in the manuscript.

Line 55-56: ...effectively modulated in presence of insulating snow cover (Gislén et al., 2014, 2016; Haberkorn et al., 2015a, 2017).

4. Line 51: Exposition is not only important for steep rock walls. It is in general important also for less inclined slopes particularly at lower latitudes; already much older literature has shown this.

We agree with the reviewer, that exposition plays as well an important role for less inclined slopes. To not exclude them, we changed the following sentence in our manuscript:

*Line 56-57: The terrain exposure induces significant spatial variability of short-wave radiation that should be considered when modelling thermal conditions in **inclined slopes**.*

5. Line 82: please give a value for the altered net short-wave radiation through the decrease in reflection so that a comparison to the value given for the change in downward long-wave radiation can be done.

The value for changes in downward long-wave radiation is given for the winter season to explain the rise in winter temperatures. For a better comparison, we cite now the value for changes in net long-wave radiation in winter season. Furthermore, we added the largest changes in net short-wave radiation, which occur in summer season due to the decrease in reflection. The following changes were made in the manuscript:

*Line 86-89: The winter warming is linked to a change in **net** long-wave radiation of $+3.9 \pm 3.9 \text{ W/m}^2$ per decade (Maturilli et al., 2015). The net short-wave radiation **is mainly altered in summer season by $+12.0 \pm$***

12.0 W/m² per decade due to the decrease in reflection caused by a reduced snow cover duration (Hop and Wiencke, 2019; Maturilli et al., 2015).

6. Line 88: Is there no mean annual precipitation available in Ny-Ålesund after 2000?

We replaced the number with the mean precipitation of the years 2000-2019:

Line 93-94: Measured mean annual precipitation in Ny-Ålesund in the period 2000-2019 was 484 mm (Norwegian Meteorological Institute, 2021).

7. Line 101: Please give some information about temperature logger calibration.

In the revised manuscript, we provide uncertainty estimates for the temperature measurements based on duplicate loggers, which were installed in close vicinity of the main logger to mitigate the risk of loss and failure. These backup loggers were generally placed within 10 cm of the main sensor in exactly the same aspect, but in different cracks or different (e.g. wider/narrower/differently shaped) parts of the same crack. These data were not used in the original manuscript, but for the revised manuscript we have used them to evaluate the combined uncertainty of the sensor+logger system and the effect of the placement in the walls. For all rock wall sites, the differences in MARST between adjacent temperature loggers (i.e. between the backup and the main temperature time series) were found to be less than 0.1 °C for annual averages, while seasonal RST showed differences of less than 0.2 °C between adjacent temperature loggers.

The following changes were done in the manuscript:

Line 115-120: The temperature sensors were placed in deep cracks in the rock wall so that both sides of the iButton are in direct thermal contact with the rock surface and the sensor is protected from sunlight (Fig. 2c). At each measurement site, we installed at least one more iButton, generally placed within 10 cm of the main sensor in exactly the same aspect, but often in different cracks or different parts of the same crack. We used these duplicate measurements to evaluate the combined uncertainty of the sensor/logger system and the placement in the rock walls. For all sites, the differences between the two sensors were found to be less than 0.1 °C for annual averages, while seasonal averages showed differences of less than 0.2 °C.

8. Line 104: the expression 'non-coastal rock walls' seems not very adequate chosen as this canyon cliffs are only about 600 m from the coast. In my view a 'non-coastal rock wall' would be several kilometers away from the fjord. Please change the wording.

We understand the concern of the reviewer that the expression can lead to misunderstandings. Therefore, we changed the wording in the entire manuscript from "non-coastal" to "near-coastal". An example would be:

Line 42-43: In this study, we will focus on rock surface temperatures in steep coastal and near-coastal cliffs at a high Arctic site close to Ny-Ålesund, Svalbard.

9. Line 114/115: how is the latent heat effect considered. Please explain or give at least a reference where the reader could get more information.

The calculation of the latent heat effect is explained in detail in Westermann et al. (2016) and we have added the reference to the statement:

Line 127-129: CryoGrid 3 calculates rock temperatures by solving the heat equation, uses the surface energy balance as an upper boundary condition, and considers latent heat effects depending on water content of the substrate as performed in Westermann et al. (2015).

10. Line 176/177: please give a source for this volumetric ice and mineral content percentages.

Park et al. (2020) analysed rock samples of carbonate rocks close to Ny-Ålesund and found a porosity of circa 0.5 % for fresh rock samples without cracks or large pores. In our study, we are not looking at undisturbed material, but have to take fractures in the rock into account. Therefore, we increased the literature value up to 3 % for the volumetric water content to account for this. As the value of 3% is a coarse estimate, it was included in the sensitivity study in supplement 1. We added an explanation to the text and added the source in table 2:

Line 194-199: We considered the bedrock to have a volumetric mineral content of 97 % and a volumetric water content of 3 %, which implied saturated conditions during the entire simulation. The assumed porosity was selected higher than measurements of 0.5 % of fresh carbonate samples without cracks in the Ny-Ålesund region (Park et al., 2020), with the goal to account for the fractured nature of the rock walls. Due to the high uncertainty of this value, a sensitivity study was performed for the volumetric mineral and water content.

11. Line 185, table 2: please give information about the source of your values you show in table 2. You could add a column in the table and show the references.

We followed the suggestion of the reviewer and added a column in table 2 to show the references. Furthermore, we added the missing references in the text.

Line 194-199: We considered the bedrock to have a volumetric mineral content of 97 % and a volumetric water content of 3 %, which implied saturated conditions during the entire simulation. The assumed porosity was selected higher than measurements of 0.5 % of fresh carbonate samples without cracks in the Ny-Ålesund region (Park et al., 2020), with the goal to account for the fractured nature of the rock walls. Due to the high uncertainty of this value, a sensitivity study was performed for the volumetric mineral and water content.

Line 204-209: We set the albedo for the horizontal ground surface to 0.15 (Westermann et al., 2009) and for water surfaces to 0.1, which is in the range of the surface ocean albedo for the typical high solar zenith angles in Svalbard (Li et al., 2006; Robertson et al., 2006). The albedo for ice and snow was set to a relatively low value of 0.55, as the highest influence of reflected short-wave radiation was expected for spring, when snowmelt decreases the albedo. This is in line with the reported decrease in albedo from 0.8 to 0.5 in Westermann et al. (2009). All values can be found in Table 2 and a sensitivity study for selected parameters is provided in the supplement.

Table 1: Model parameters assumed in the simulations.

Parameter		Value	Unit	Reference
Albedo rock wall	α	0.30	[-]	Blumthaler and Ambach (1988)
Albedo ground	α_g	0.15	[-]	Westermann et al. (2009)
Albedo open water	α_w	0.1	[-]	Li et al. (2006)
Albedo melting snow / ice	α_s	0.55	[-]	Westermann et al. (2009)
Emissivity	ϵ	0.97	[-]	Bussieres (2002)
Roughness length	z_0	0.018	[m]	-
Mineral fraction	mineral	0.97	[-]	modified after Park et al. (2020)
Water and ice fraction	waterIce	0.03	[-]	modified after Park et al. (2020)
Water bucket depth	d	0.001	[m]	-

12. Line 194/195: Is the effect (transition between fjord and land) you describe here really resolved? I can hardly believe this!

The effect is not resolved, but as the grid cell covers both the fjord and land surface, it is the best possible solution to use it as forcing data for the shoreline. To avoid misunderstandings, we phrased the sentence differently:

Line 220-222: The selected grid cell covers both parts of the fjord and the adjacent land surface and therefore provides suitable forcing data for the loggers located directly or within a short distance to the shoreline.

13. Line 203: why do you use the radiation data from AROME-Arctic dataset when you have much better data from the BSRN stations. Please clarify?

We stayed with the data of AROME-Arctic throughout our study (using radiation, wind speed, rain/snowfall etc.) to use coherent forcing data and to avoid mixing data of different sources. The AROME-Arctic dataset was used as forcing data before also in other studies in the surroundings of Ny-Ålesund like in Zweigel et al. (2021). In our study, the only exception was air temperature to resolve local air temperature gradients.

Zweigel, R., Westermann, S., Nitzbon, J., Langer, M., Boike, J., Etzelmüller, B. and Schuler, T. V.: Simulating snow redistribution and its effect on the ground thermal regime at a high-Arctic site on Svalbard, J. Geophys. Res.-Earth, in press, 2021.

14. Line 240, table 4: RW01 is modeled according to table 4, but it is shown that this site is covered by snow in figure 3. Therefore, it is mandatory to include snow in the model scenarios for this logger otherwise you contradict yourself in the paper (see also the general comments about snow).

As described in the reply to the comment “neglecting the influence of snow”, the snow cover at the rock walls are mainly due to snowdrift and therefore, it is difficult to represent in our model. Consequently, we decided to focus our analysis on the periods without snow cover in the rock walls. To analyse the effect of snow, we added a supplement in our manuscript. The changes in the manuscript can be found under the comment “neglecting the influence of snow”.

15. Line 255, table 5: your model does not include snow but you wrote that only one logger is snow covered RW01 (figure 3). Please clarify this as it is very important for your assumption that there is no snow cover at the sites.

We clarified in our manuscript which rock walls were affected by snow cover. To do so, we added a sentence in the paragraph of figure 3 and distinguished in table 5 between missing data and snow cover. While snow cover seems to occur at RW01 on a regular basis in spring and over a period of two to three months (in all years except for 2018 in our measurement period), the other rock walls are only affected occasionally and over a shorter time period: in May 2019 RW05 was affected, in May 2020 RW05, RW06 and RW08. We completely agree with the reviewer, that this information has to be given more clearly in our manuscript. The explanation why we excluded snow cover from our analysis is given in supplement 2.

We made the following changes in our manuscript:


Line 184-189: We did not consider snow cover in the model, which was adequate for most of the measurement data in the analyzed time period from 2016 to 2020. An exception is displayed in Fig. 3, showing the damped signal of RW01 due to snow cover. Besides spring 2017, RW01 was influenced by snow cover over a period of two to three months in spring 2019 and 2020. In the other measurement sites, snow cover was only shortly observed in May 2019 (RW05) and in May 2020 (RW05, RW06, RW08). Further evaluations on the possible influence of a snow cover are given in the supplement, where we present model runs considering snow cover in the rock walls following the model approach of Magnin et al. (2017).

Table 5: Measured MARST and mean RST in the winter season (Dec-Feb) for all locations RW01 to RW08. Lack of data results from either (1) snow cover or (2) missing records. MARST are coldest in near-coastal settings (RW01 - RW03). Mean RST in winter season are found to be coldest in near-coastal settings, closely followed by settings in the bay (RW08), while stings at the open fjord show highest RST (RW04 – RW07).

Location	Site	Entire year				Winter: Dec-Feb			
		2016/17	2017/18	2018/19	2019/20	2016/17	2017/18	2018/19	2019/20
RW01	Near-coastal	- (1)	-1.5	- (1)	- (1)	-8.1	-6.7	-9.0	-13.1
RW02	Near-coastal	-2.2	-1.8	-2.4	-4.3	-8.5	-6.6	-9.6	-13.5
RW03	Near-coastal	-1.8	-2.0	-2.4	-4.1	-9.2	-7.2	-10.0	-13.8
RW04	Open fjord	- (2)	-1.0	-2.1	-3.6	- (2)	-5.1	-8.7	-12.0
RW05	Open fjord	-0.9	-0.8	- (1)	- (1)	-6.6	-5.2	-7.2	-10.4
RW06	Open fjord	- (2)	-0.6	- (2)	- (1)	- (2)	-5.0	-8.5	-11.5
RW07	Open fjord	- (2)	-0.8	- (2)	-3.6	- (2)	-4.6	-7.6	-11.1
RW08	Bay	-1.4	-0.9	-2.1	- (1)	-8.6	-5.8	-9.7	-13.1

16. Line 275, figure 4: why is the variability (daily values?) not higher in comparison to figure 3 where there is much more variability in the same data. please clarify.

Figure 3 does not show significantly higher variability in daily mean values. It might appear because the plot contains temperature data of more loggers and the time period is longer. Please find here a plot with similar time scales (both over 3 months) and the same loggers (RW02, RW04, RW08). RW04 is not included in the first plot, as it was not installed yet.



17. Line 335, figure 7: What is Fub in the figures? Is this not G as noted in the figure caption?

We thank the reviewer for this comment. The correct label in figure 7 is of course “G”. We changed it in the revised manuscript:




Figure 1: Surface energy balance for the seasons of the year for a vertical rock wall with an aspect of 40° (RW02, RW04 and RW08). Most pronounced differences in the scenarios are found in winter and spring, while summer and fall show similar fluxes. SW = net short-wave radiation; LW = net long-wave radiation; Qe = latent heat flux; Qh = sensible heat flux; G = ground heat flux.

Literature: Haberkorn, A., Wever, N., Hoelzle, M., Phillips, M., Kenner, R., Bavay, M., Lehning, M. (2017) Distributed snow and rock temperature modelling in steep rock walls using Alpine3D. *The Cryosphere* 11, 585-607. Haberkorn, A., Phillips, M., Kenner, R., Rhyner, H., Bavay, M., Galos, S.P., Hoelzle, M. (2016) Thermal regime of rock and its relation to snow cover in steep alpine rock walls: gemsstock, central swiss alps. *Geografiska Annaler: Series A, Physical Geography* 97, 579-597. Haberkorn, A., Hoelzle, M., Phillips, M., Kenner, R. (2015) Snow as a driving factor of rock surface temperatures in steep rough rock walls. *Cold Regions Science and Technology* 118, 64-75.

The following literature was already included in our manuscript:

Haberkorn, A., Phillips, M., Kenner, R., Rhyner, H., Bavay, M., Galos, S. P. and Hoelzle, M.: Thermal regime of rock and its relation to snow cover in steep alpine rock walls: gemsstock, central swiss alps, *Geografiska Annaler: Series A, Physical Geography*, 97(3), 579–597, <https://doi.org/10.1111/geoa.12101>, 2015.

Haberkorn, A., Wever, N., Hoelzle, M., Phillips, M., Kenner, R., Bavay, M. and Lehning, M.: Distributed snow and rock temperature modelling in steep rock walls using Alpine3D, *The Cryosphere*, 11(1), 585–607, <https://doi.org/10.5194/tc-11-585-2017>, 2017.

In addition, we added the suggested literature

Haberkorn, A., Hoelzle, M., Phillips, M. and Kenner, R.: Snow as a driving factor of rock surface temperatures in steep rough rock walls, *Cold Regions Science and Technology*, 118, 64–75, <https://doi.org/10.1016/j.coldregions.2015.06.013>, 2015a.

in the introduction of our revised manuscript:

Line 55-56: ...effectively modulated in presence of insulating snow cover (Gisnås et al., 2014, 2016; Haberkorn et al., 2015a, 2017).