

PISM-LakeCC: Implementing an adaptive proglacial lake boundary into an ice sheet model

Sebastian Hinck¹, Evan J. Gowan^{1,2}, Xu Zhang^{1,3}, and Gerrit Lohmann^{1,2}

¹Alfred Wegener Institute Helmholtz Centre for Polar and Marine Research, Am Handelshafen 12, 27570 Bremerhaven, Germany.

²MARUM - Center for Marine Environmental Sciences, University Bremen, Leobener Strasse 8, 28359 Bremen, Germany.

³College of Earth and Environmental Science, Center for Pan-Third Pole Environment, Lanzhou University, 222 Tianshui South Road, Lanzhou City, China

Correspondence: Sebastian Hinck (sebastian.hinck@awi.de)

Abstract. Geological records show that vast proglacial lakes existed along the land terminating margins of palaeo ice sheets in Europe and North America. Proglacial lakes impact ice sheet dynamics by imposing ~~marine-like boundary conditions at the ice boundary conditions analogous to a marine terminating~~ margin. These lacustrine boundary conditions include changes in the ice sheet's geometry, stress balance and frontal ablation and therefore affect the entire ice sheet's mass balance. ~~This interaction, however, has not been rigorously implemented in ice sheet models~~ Nevertheless, proglacial lakes have rarely been considered in detail in ice sheet modelling endeavors. In this study, the implementation of an adaptive lake boundary into the Parallel Ice Sheet Model (PISM) is described and applied to the glacial retreat of the Laurentide Ice Sheet (LIS). The results show that the presence of proglacial lakes locally enhances the ice flow. ~~Along the continental ice margin, ice streams and,~~ which leads to a lowering of the ice lobes can be observed. Lacustrine terminating ice streams cause immense thinning of the ice sheet's interior and thus sheet and emergence of ice lobes. ~~Due to increased melting at the lowered ice surface and enhanced mass loss due to calving at the lake boundary, the lacustrine interactions~~ play a significant role ~~in the demise of the LIS. Due to during deglaciation. For the LIS,~~ the presence of lakes ~~triggers~~ a process similar to the marine ice sheet instability, which causes the collapse of the ice saddle over Hudson Bay, ~~which blocked drainage via the Hudson Strait~~. In control experiments without ~~a the~~ lake model, Hudson Bay is still glaciated ~~at the end of the simulation. Future studies should target the development of parametrizations that better describe the glacial-lacustrine interactions. when the climate reaches present day conditions.~~

Copyright statement.

1 Introduction

During the Last Glacial Maximum (LGM), ice sheets covered parts of the Eurasian and North American continent. As they retreated, the topography was left deeply depressed due to delayed glacial isostatic adjustment (GIA). ~~Impounded between depressions and retreating ice, vast~~ Vast lakes formed along the ice margins due to a combination of this depression, and the

blocking of drainage routes by the ice sheets. Examples of these palaeo lakes are Lake Agassiz in North America (Teller and Leverington, 2004) and the Baltic Ice Lake in Eurasia (Björck, 1995). ~~Geological evidence, such as lacustrine~~ Lacustrine sediments and shorelines ~~provide~~ provide geological evidence of their presence and extent. Reconstructions of Lake Agassiz, ~~for which bordered the Laurentide Ice Sheet (LIS) for several hundred km,~~ for example, suggest areal extents of several hundred thousand km² ~~and~~ and water depths of several hundred meters ~~and bordering the Laurentide Ice Sheet (LIS) for several hundred km~~ (Teller et al., 2002; Leverington et al., 2002; Teller and Leverington, 2004).

The ~~lake basins~~ basins of proglacial lakes constantly evolved, because the ice sheet margin and topography was ~~dynamienot~~ static. Reorganization of the lakes' drainage networks and sudden drainage events due to the opening of lower spillways may have impacted the global climate by perturbing the ~~thermo-haline~~ thermohaline circulation system of the oceans (Broecker et al., 1989; Teller et al., 2002).

(Broecker et al., 1989; Teller et al., 2002; Peltier et al., 2006; Condon and Winsor, 2012). Furthermore, the presence of a lake at the ice margin impacts the ice dynamics by adding a marine-like boundary ~~conditions~~ condition to the ice sheet. This changes the boundary conditions of terrestrial ice margins: modification of the thermal regime at the submerged ice base, formation of ice shelves, increased ice loss due to melting and calving, and enhanced basal sliding near the ~~grounding-line~~ grounding line due to decreased effective pressure at the ice base (e.g. Carrivick and Tweed, 2013). These processes can lead to the formation of ice streams (Stokes and Clark, 2003; Margold et al., 2015), which ~~impacts~~ impact the mass balance of large parts of the ice sheet. ~~Due to various differences between the freshwater and ocean water, the interactions might be different than at a marine boundary (?)~~ Interactions between ice and lake resemble processes at the marine boundary, but their magnitude might differ due to various differences at lacustrine and marine boundaries (Benn et al., 2007).

In ~~times when there was the production of~~ periods with large amounts of meltwater that caused the formation of proglacial lakes ~~such as during the deglaciation of North America, the~~ the lake-ice interactions ~~are~~ might have been a key factor to explain rapid glacial retreat. As an example, ~~the one hypothesis for the cause of the~~ the 8.2ka event, a period characterized by a sudden drop in Northern Hemispheric mean temperatures, ~~was caused by~~ is the rapid demise of the central LIS (Carlson et al., 2008; Gregoire et al., 2012; Matero et al., 2017; Lochte et al., 2019) leading to a large freshwater input into the Labrador Sea. Although the retreat is assumed to be governed by the negative surface mass balance due to enhanced melting in a warming climate (Carlson et al., 2009; Gregoire et al., 2012), other dynamical effects, such as marine and lacustrine interactions, might have further amplified this effect (Matero et al., 2017, 2020). Matero et al. (2020) state the lack of proper representation of Lake Agassiz/Ojibway along the southern ice margin in their modeling study as a source of uncertainty. Lake reconstructions of this time suggest water depths up to several hundred meters (Teller et al., 2002; Leverington et al., 2002). By adding a simple parameterization for lacustrine calving to their energy balance type box model, Fowler et al. (2013) could show that this feedback can potentially explain the 100 kyr climate oscillations since the mid-Pleistocene.

~~Apart from their relevance for understanding processes that lead to the demise of the late Pleistocene and early Holocene ice sheets, interest in contemporary proglacial lakes and their role in glacial retreat is growing (Carrivick and Tweed, 2013). Motivations for these studies range from predicting and managing water resources under a warming climate and recognizing possible risks due to glacial outburst floods (Carrivick et al., 2020).~~

In most previous numerical studies, glacio-lacustrine interactions have not been considered ~~or are applied in an unrealistic way~~. Often ice dynamical models apply marine boundary conditions at places with surface elevation below global mean sea level (e.g. the PISM authors, 2015). This can lead to inner-continental ocean basins which can be considered as 'fake' lakes, with a water level that can greatly differ from the true level of a ponded proglacial lake (Matero et al., 2020). Cutler et al. (2001) and Tsutaki et al. (2019) investigated the impact of a prescribed lake level on ice dynamics using two-dimensional flow-line models. Tarasov et al. (2012) included the effects of proglacial lakes in an ice sheet model through a calving parameterization and thermodynamic refreezing scheme. Recent work of Sutherland et al. (2020) present a novel approach analyzing the impact of lake boundary conditions on ice dynamics using the three-dimensional thermo-mechanically coupled ice sheet model BISICLES (Cornford et al., 2013). Their regional survey treats the post-LGM retreat of a mountain glacier in the Southern Alps in New Zealand, terminating in glacial Lake Pukaki. They subtracted the reconstructed water level from model topography and adapted parameters controlling marine boundary interactions for lakes. This approach, however, only allows one fixed water level and is thus not applicable for more complex ice sheet scenarios featuring multiple, time variable lakes. ~~The~~ In another recent study Quiquet et al. (2021) applied the GRISLI model (Quiquet et al., 2018) to simulate the glacial retreat of the North American ice sheets during the last glacial cycle. At the southern ice margin they simulated the effect of a proglacial lake by locally elevating the sea level. As in the aforementioned study, this water level is fixed. The model results show an accelerated demise of the LIS caused by the occurrence of a proglacial lake ice sheet instability (PLISI). The need for including glacio-lacustrine interactions into numerical ice sheet modeling attempts was recently highlighted in review articles by Margold et al. (2018) and Carrivick et al. (2020). The latter ~~reference discusses concerns when implementing~~ referenced article discusses challenges for the implementation of such a lake-ice boundary condition for ice sheet models.

In this study, we describe the implementation of an adaptive lake boundary into a 3D thermo-mechanically coupled ice sheet model. Implementation into the Parallel Ice Sheet Model (PISM; the PISM authors, 2015; Bueler and Brown, 2009; Winkelmann et al., 2011) is done using a generalization of the model's marine boundary condition. The fundamental algorithm that determines the lake basins is based on the standalone model LakeCC (Hinck et al., 2020), so the model is called PISM-LakeCC. 'CC' stands for connected components, the ~~approach~~ algorithm the model is based on.

We demonstrate the model's impact on the ice dynamics ~~with an application to~~ through the application of a simple post-LGM deglaciation scenario of the North American ice complex. A comparison with control runs shows that lakes induce strong dynamical effects on the ice sheet. Increased mass loss and reduced basal strength lead to ~~the formation of ice streams along an acceleration of ice flow upstream of~~ the lake boundary, which effectively drains mass from the ice sheet interior. During the deglaciation, lakes form with water depths up to several hundreds meters. ~~Where the water is deep enough, ice shelves form.~~ The differences from the control experiments become apparent with the demise of the remainder of LIS in the Hudson Bay ~~area. In places~~ Area. In regions where there is ice-inward sloping topography, the ~~lake can rapidly expand, as the water is deep enough that the ice begins to float. This further~~ grounding line can retreat in a self-amplified manner, which corresponds to a rapid expansion of the lake. This triggers the occurrence of the PLISI, which is mainly driven by the surface elevation feedback. The instability drastically accelerates the demise of the LIS and finally leads to the drainage of the lake through Hudson Strait.

90 In the control ~~experiment~~experiments, Hudson Bay is still glaciated ~~at the end of the run~~when the present day conditions are achieved.

2 Methodology

2.1 Ice sheet model

All implementations described in this work are based on the stable release v.1.2.1 of the Parallel Ice Sheet Model (PISM; 95 the PISM authors, 2015; Bueller and Brown, 2009; Winkelmann et al., 2011). PISM is a 3-dimensional thermo-mechanically coupled ice sheet model, whose modular implementation grants the user freedom to choose between different realizations of various sub-models. In the following we give a short overview about the configuration used for our experiments. All details that diverge from the PISM defaults are listed in Tables A1 and A2 in the Appendix.

PISM's computational grid is horizontally equally spaced. We use a resolution of 20 km. In the vertical direction, the 100 computational box has a height of 5750 m and is divided into 101 layers, with height that increases quadratically from the ice base. The thermal layer within the ~~bedrock~~ground has a thickness of 2000 m and is divided into 11 levels. An adaptive time-stepping mechanism determines the shortest time step needed by any of PISM's sub-models. ~~In this study we set an upper bound of 0.25 yr for the time step.~~

The stress balance is modeled using a hybrid scheme based on the Shallow Ice (SIA) and Shallow Shelf Approximations 105 (SSA) of the full Stokes equations (Bueller and Brown, 2009). We use an energy conserving flow law, the Glen-Paterson-Budd-Lliboutry-Duval law ~~(??)~~(Lliboutry and Duval, 1985; Aschwanden et al., 2012). This models the ice softness as a function of temperature and liquid water fraction.

The basal resistance is determined using a model that assumes that the base of the ice sheet is underlain by deformable till. It only allows sliding when the driving stresses exceed the yield stress of the till. This threshold value is determined by a 110 Mohr-Coulomb formulation (Cuffey and Paterson, 2010) using a constant till-friction angle and the effective pressure in the till. The latter is estimated from the amount of water in the till, which comes from the non-water conserving hydrology model of PISM (Tulaczyk et al., 2000), and the ice thickness. The till below the water level next to the ~~ice margin and grounding line~~grounding line are assumed to be saturated, which reduces the basal strength. Bed deformation due to ice load is modeled in PISM using the ~~Lingle-Clark~~Lingle-Clark model (Lingle and Clark, 1985; Bueller et al., 2007). It uses an idealized two-layered 115 Earth model, approximating the Earth as a viscous upper mantle overlain by an elastic plate lithosphere.

The surface mass balance (SMB) is estimated from monthly means of precipitation and surface air temperature fields using the positive degree-day (PDD) approach (Calov and Greve, 2005). To prescribe the atmospheric forcing for the transient experiment conducted here, two additional models were implemented. Precipitation and temperature fields are interpolated between two distinct climatic states, which are weighted according to a glacial index (see Appendix D1). To prevent the ice 120 sheet from expanding into regions and high elevations where a more advanced approach would limit precipitation, the second model sets precipitation to zero above a threshold height or accordingly to a given mask (see Appendix D2).

~~The Marine regions of the ice sheet are defined in PISM via a flotation criterion, which describes whether the ice in areas below sea level is grounded or floating. The sea level is defined via a 2D map, which allows it to be spatially variable. In general, however, it is set to a global mean value prescribed by a scalar time series. The marine boundary treatment is described in Winkelmann et al. (2011) and Martin et al. (2011). It includes a sub-shelf-melting ~~parametrization~~ parameterization (Beckmann and Goosse, 2003), and sub-grid parameterizations of the ice shelf advance (Albrecht et al., 2011) and the ~~grounding-line~~ grounding line (Feldmann et al., 2014). To ~~parametrize~~ parameterize the effect of calving at the ice shelf front, a combination of the Eigen-calving¹ (Levermann et al., 2012) and thickness calving mechanisms, which removes ice thinner than a given threshold thickness, is applied. ~~The temporal evolution of the sea level elevation is prescribed by a scalar time series (see Fig. A1b).~~~~

2.2 LakeCC

The environment along the ice sheet edge is ~~quite particularly~~ dynamic. Ice margin migration and ongoing GIA significantly impact shape and size of proglacial lakes that form within this environment. ~~Therefore, when dynamically coupling~~ For the dynamical coupling of ice sheets and proglacial lakes, ~~steady updating of the~~ continuous updating of lake basin geometry is necessary. Depending on the complexity of the lake model, computational overhead can drastically increase. For application on ~~continental-sized~~ continental-sized ice sheets, the use of lightweight algorithms is necessary. To reduce the complexity, trade-offs have to be ~~donemade~~. This section describes how the LakeCC model computes lake reconstructions and prepares them for use in PISM. Details on how this lake boundary condition affects the ice dynamics is provided in Sect. 2.3. Limitations of the current implementation are discussed in Sect. 2.4.

For the model described here, we assume that lake basins tend to be entirely filled. This assumption might be valid for proglacial lakes during times of glacial retreat, when ~~melt-water~~ meltwater is entering the lake. However, rapid changes in the boundary conditions resulting from this approach often cause numerical instabilities ~~which that~~ cause the model to crash ~~(see Sec. 2.4.4)~~. To overcome this, changes in the lake geometry need to be monitored, and the water level needs to be adjusted gradually. Our implementation uses two different fields to realize this: the target level and the lake level (see Figure 1).

The target level field holds information about the maximum water level of all lake basins of the domain, while in the other field the actual water level is stored. These fields resemble a spatial map whose cells hold the respective information of the associated lake basin; cells outside such basin are set to be invalid. To determine the target level, a simple lake filling algorithm is used that is adapted from Hinck et al. (2020). Changes in this field, i.e. the appearance or disappearance of lake basins, are monitored by comparing the target level against the lake levels from the previous time step. This is necessary because the gradual filling algorithm relies on new lake cells being properly initialized in the lake level field. In general, new lakes are initialized with zero water depth. However, there are special cases, such as adding a lake basin to an existing lake or adding a basin that has previously been connected to the ocean, that need more advanced treatment. For more details on this, see Appendix B.

¹~~Although the Eigen-calving mechanism was activated for all experiments, the model was found to not have impacted the results.~~




Figure 1. Sketch of the PISM LakeCC and SL2DCC models. The target water level indicates the maximum water height of a lake basin before it overflows ~~into the ocean~~. The actual lake level, as it is seen from the rest of the ice model, is slowly raised (or lowered) towards the target level. The dotted lines and hatched areas indicate where the respective water levels extend into the ~~bedrock~~ground. To prevent the formation of inland ocean basins, the SL2DCC model sets the sea level in those basins to invalid. Note that lake level and the gap in the sea level field are extended by one cell, respectively. For details on this the reader is referred to the main text.

After initialization, the lake level is gradually adapted towards the target level. Determining a rate γ , at which the water levels ~~are adapted~~change is not trivial. For simplicity the fill-rates in our model are assumed to be constant. See Sect. 2.4 for more details on this.

Merging of lakes or adding a new basin to an existing lake might result in an unbalanced water level. To quickly overcome this unrealistic scenario and balance the lake level, water levels for each lake are adapted from a common level. When water level is rising, this common level is chosen to be the lowest water level of that lake, h_{\min} , while the highest level, h_{\max} , is selected for the falling water level. The potential new water level, h , is determined by adding or subtracting the respective change for a given time-step, dt :

$$h = h_{\min/\max} \pm \gamma \cdot dt \quad (1)$$

Only when h has exceeded the ~~current (local)~~ lake level, does its value get updated. This makes sure that the lake level is slowly raised or lowered towards the target level, while aligning the water levels. When a lake disappears, ~~i.e.~~i.e., the target level is set to be invalid, the lake is not immediately removed, as this could also lead to numerical instabilities. The water level is gradually lowered until it is below the bed elevation, or the ice sheet is grounded. If a basin disappears because it merged with the ocean, the lake level is gradually changed until sea level is reached, and then removed.

In the final step, the lake basins are all extended by one grid cell in each direction. This treatment serves two purposes: (i) providing information about the presence of a neighboring lake to the ice sheet model to possibly adapt its boundary conditions, and (ii) helping to close gaps in between ice margin and adjacent lake when the ice is retreating in the following time-step. It should be noted that this does not artificially enlarge the lake, as the ice sheet model independently computes the lake geometry

based on the information of the lake model and the water level in these cells is below the flotation threshold. The dotted lines in Fig. 1 illustrates this.

Another issue that we resolve here is the way how PISM and other ice sheet models (e.g., e.g. BISICLES, as noted in Matero et al. (2020) and Sutherland et al. (2020)) handle sea level and how it affects the LakeCC model. ~~Sea-~~The sea level elevation is usually assumed to be globally constant and hence is set by a scalar (time series). Differentiation between ocean and land is only done by checking if the bed elevation is above or below the sea level. Consequently, isolated basins can occur within the continent, and if they are below sea level, are falsely regarded as ocean. As described in Hinck et al. (2020), the LakeCC algorithm relies on a proper land-sea mask to properly determine lake basins. ~~Therefore, the use of a more advanced sea level model is necessary. This sea level model implemented here~~This mask can be computed by the SL2DCC model (for more details see Appendix B4), which is ~~named PISM-SL2DCC, is~~ also based on Hinck et al. (2020) and uses the connected components (CC) ~~approach to determine the two-dimensional sea level field~~algorithm. It checks ~~ocean cells on a 2D field~~potential ocean cells for connectivity with the domain boundary and marks isolated basins as invalid. The corresponding cells are recognized by the ice sheet and the lake model as land cells, which are potentially available for lake formation. ~~Further details on the SL2DCC model can be found in Appendix B4.~~

2.3 Lacustrine impact on ice dynamics

In this implementation, lakes are treated as a generalization of the existing marine boundary condition in PISM that dynamically adapts to changing environments. Most parts of the ice sheet model that access the sea level elevation do this to obtain information about the current geometry, e.g. to determine water depth. The code is adapted in a way that, if a lake is present, the lake level is used in the calculation instead of sea level and the water density is adjusted accordingly. Generally, the parameterizations provided by PISM for a marine boundary are applied analogously at the lake interface. This, however, may not be the optimal treatment in every case. The model might benefit from future implementations of advanced or more specialized lake boundary treatment. Such limitations are discussed in Sect. 2.4.

Figure 2 shows the various locations where an aquatic environment impacts the ice dynamics in PISM. In the following we will describe the parameterizations used in this setup to describe the lacustrine impact on ice dynamics. The numbers refer to their respective label in Fig. 2.

- 1 ~~---~~--- Lubrication of ice base at grounding line: Cells with grounded ice below water level next to a lake are assumed to have saturated till. This reduces the effective pressure at the base of the ice sheet and reduces basal resistance ~~in~~at this location.
- 2 ~~---~~--- Sub-shelf melting/ refreezing: In the current configuration, the ice sheet model does not differentiate between ice shelves in marine and lacustrine environments, i.e. the same parameterization for ice base temperature and sub-shelf mass-flux are used. Ice temperature at the base is set to the pressure-melting point, which is a function of pressure, hence ice thickness. Sub-shelf mass-flux is calculated based on the ~~parameterization~~parameterization from Beckmann and Goosse (2003). This formulation explicitly assumes a marine environment. It models the mass-flux proportional to the difference between the pressure-dependent freezing point of saline water and the temperature of the ambient ocean. Salinity and ambient ocean temperature are prescribed




Figure 2. Different ways how the ice dynamics are impacted by a marine or lacustrine environment in PISM. 1 — Till in cells at the grounding line is assumed to be water-saturated, which reduces basal friction. 2 — Sub-shelf melting/refreezing. 3 — Pressure of the water column against the submerged part of the grounded or floating part of the ice margin. It can be further increased by modeling the ice mélange. ~~Parametrization~~ Parameterization of mass-loss due to 4 — calving at the shelf edge and due to 5 — frontal melt at the submerged ice margin. 6 — Where ice fulfills the flotation criterion it becomes buoyant, hence lowering ice surface slope, eliminating basal friction and exposing the ice to shelf processes. Sub-grid treatment of shelf edge and ~~grounding-line~~ grounding line position are not explicitly marked here.

205 in this model implementation and thus can not be adjusted for freshwater environments. The model does not allow for refreezing at the shelf base.

3 — Ice-marginal pressure difference and mélange back-pressure: The pressure difference against the submerged ice margin is done analogously to the marine boundary, but evaluated for freshwater density. Ice mélange, which is not modeled here, can have a buttressing effect on the ice sheet.

210 4 & 5 — Calving and frontal melt: Frontal retreat is a complex but also very sensitive factor in modeling both, marine and lacustrine terminating ice sheets. In this study we only parameterize calving, with no frontal melt schemes applied. At the lake boundary, the same calving mechanisms are applied as for the ocean (compare with Sect. 2.1). The implementation of the thickness threshold calving was adapted to accept a distinct threshold value that is used at lacustrine boundaries (Δh_L).

215 6 — Formation of ice shelves: Where ice thickness is below a certain threshold, determined by depth and density of the adjacent water body, the ice floats. Formation of an ice shelf not only gives rise to the appearance of the previously mentioned effects (1-5), but also directly impacts ice dynamics. Due to geometric changes of the ice sheet, the flow regime is altered. Furthermore, friction at the shelf base is negligible.

By impacting the ice sheet geometry and mass balance further, secondary mechanisms are triggered that feed back onto the ice dynamics. These include changes in GIA and affecting the local climate due to temperature elevation feedback.

220 2.4 Limitations

For the PISM-LakeCC model, several assumptions were made to reduce the complexity of the problem and adaptively updating the lake boundary condition within the ice sheet model, without adding too much computational overhead and destabilizing the numerical system. These include both aspects of the model, the lake reconstruction and the coupling to the ice sheet. In the following, known limitations are listed and discussed.

225 2.4.1 Lake reconstruction

The PISM-LakeCC model uses the same numerical grid as the ice sheet model. For hydrological applications, such as lake basin reconstructions, the resolution an ice sheet model usually operates on is too coarse to resolve spillways through the terrain. ~~Even more important than data resolution is~~ At low resolutions, lake volumes are potentially overestimated (Berends and van de Wal, 2016). ~~Since we are not so much interested in lake volume,~~ the ice margin position and bed deformation due to GIA are, in our case, expected to be more important than resolution (Hinck et al., 2020). Considering the uncertainties of these fields retrieved from an ice sheet model, ~~the resolution issue~~ resolution is regarded as a secondary issue. A method, proposed by Hinck et al. (2020), is implemented into the LakeCC model to adapt to the low resolution. In that study, satisfactory results were obtained by applying the LakeCC algorithm to a low resolution PD topography map of North America after applying a ~~secondary field. This correction. This correction~~ field was obtained from a high resolution dataset in a preprocessing step by applying a minimum
235 filter that retains the lowest surface elevation within a certain distance of each grid cell, and calculating the difference with the lower resolution topography. The PISM-LakeCC model reads this field from an input file and applies it to the low resolution topography prior to the model update. If higher resolved input fields are available, e.g. from an external GIA model, the LakeCC model could be modified to do the calculations on that field instead and interpolate the output back onto the ice sheet model grid. ~~It is questionable if the results are worth the additional computational expense.~~

240 We further assume that all lakes tend to fill to the brim, i.e. there is no accounting for conservation of water mass. Doing this properly would require ~~permanent updating~~ updating and accounting of the hydrological network and lake basins, accumulating all water fluxes and finally iteratively redistributing overflow down-gradient. This requires highly resolved hydrology, including water fluxes of climatological and glacial origin. The required data is not available in our framework and furthermore, such algorithm would dramatically increase computational cost. During glacial retreat, when ~~melt water~~ meltwater is pervasive, the
245 assumption of filled lakes is likely valid.

~~Sudden~~ To avoid numerical instabilities due to sudden jumps in water level ~~can trigger numerical instabilities in the ice sheet model. To avoid such jumps,~~ the water level is gradually ~~adjusted with a constant rate. Setting this value should not imply that there is a common valid filling constant for proglacial lakes, as this obviously relies on individual fluxes and the areal extent of the lake. Future implementations could possibly adapt a volumetric rate instead of fixing the rate of change of water level. By limiting the time step, sudden changes in water level could be performed quicker.~~

250 ~~As a consequence of the slowly adapted water level, unphysical situations can arise~~ adapted, which can lead to unphysical situations. When lakes merge, for example, this can lead to an unbalanced water level. Furthermore, small, shallow lakes

can be overrun by the forward advancing ice margin, which creates an artificial sub-glacial lake, until the water level dropped below the floating threshold. These issues can not be avoided using such a simple model. However, the algorithm automatically
255 resolves these problems within a few time steps.

2.4.2 Glacio-lacustrine interactions

The lake-ice boundary is treated ~~in general~~, in general, the same as the marine ice boundary (see Sect. 2.3). Some ~~of the~~ parameterizations are explicitly formulated for the marine ice margin and other processes are reported to substantially differ ~~in strength~~ between lacustrine and oceanic environments (Carrivick et al., 2020). Implementation of specific lacustrine processes
260 could yield a more appropriate treatment of lake boundaries. In the following we discuss the lake-ice interactions with respect to the validity at the lacustrine ice margin. The numbers refer to the processes in Fig. 2.

For our experiments, the so-called slippery grounding line parameterization (1) has been used, which reduces the basal friction in the cells upstream the grounding line. This simple model therefore adds a direct dependency to the grid resolution. Furthermore, an increased sensitivity of the grounding line is reported when using this parameterization (Golledge et al., 2015)
265 . For this reason, sensitivity tests were run that confirm these findings; these results are presented in the supplementary material and discussed in Sect. 4.3. Due to lack of a more advanced implementation to model lubrication at the grounding line, we apply the slippery grounding line treatment.

At the shelf base (2), mass flux is parameterized using the model proposed by Beckmann and Goosse (2003). The model relates the mass flux to the difference between the pressure dependent freezing point of saline water and the temperature of the
270 ambient ocean. Freshwater, however, behaves differently from saline water, as its temperature is always above the melting point of ice, and due to the ~~anomaly in the density properties~~ of freshwater, the densest waters at the lake bottom are around 4°C. Due to the difference in density between fresh and ocean water, the layer of melt water underneath the ice shelf experiences a ~ 200 times higher buoyancy in ~~sea water~~ (Funk and Röthlisberger, 1989). ~~This seawater~~ (Funk and Röthlisberger, 1989). In saline water this increases the flux and mixing of ambient warmer water along the ice base and increases melting. Based of
275 these considerations, the melt flux was scaled accordingly in the MR sensitivity run, which is documented in the supplementary material. Because of these differences it will be important to implement a sub-shelf melting model for lacustrine environments in future studies.

A lacustrine model is also needed for frontal ablation (calving (4) and frontal melt (5)). Observations at contemporary proglacial lakes show calving rates an order of magnitude below the rates of tidewater glaciers (~~Funk and Röthlisberger, 1989; Warren et al.~~
280 ~~Funk and Röthlisberger, 1989; Warren et al., 1995; Skvarca et al., 2002; Warren and Kirkbride, 2003; Haresign, 2004; Benn et al., 2007~~). However, it should be noted that alpine proglacial lakes differ fundamentally in size and depth from the major palaeo proglacial lakes. In order to have some control on that, the current implementation accepts ~~an a~~ unique parameter for the threshold thickness calving mechanism at the lacustrine boundary. Implementation of a more physically based calving model capable of accurately parameterization of mass losses in both lacustrine and marine environments, will be needed (?)(Benn et al., 2007).

285 Another important issue that is ignored in our model is the effect of ice mélange (3), especially in smaller lakes. This is expected to have an a buttressing effect on the ice sheet and reduce mass loss. Seasonal ice cover might even increase this effect (Mallalieu et al., 2020).

2.4.3 Further lacustrine interactions

290 Our lake model explicitly only interacts with the ice dynamics, as described above. However, there are ways a lake impacts the various subsystems of the Earth system, which can not be solved for within an ice sheet model. Some of these mechanisms, which are ignored here are briefly mentioned in this section.

The presence of vast proglacial lakes impacts the local climate by reducing temperatures and increasing precipitation patterns (Krinner et al., 2004; Peyaud et al., 2007). This can locally increase the ice sheet's SMB and counteract the accelerated mass loss observed in this study. Also, the potential impact on ocean circulation and global climate due to redistribution of freshwater (Broecker et al., 1989; Teller et al., 2002; Condrón and Winsor, 2012) is ignored here.

The location where proglacial lakes appear strongly depends on the GIA signal. The GIA signal is also partially dependent on the mass held by the lakes. This feedback, however, is not taken into account here.

2.4.4 Gradual filling & numerical instability

300 When running the model, we observed that PISM occasional crashes when there is rapidly rising or dropping the water levels between time steps. Unfortunately, we can not tell the exact reason why the numerical solver failed, but we found that these crashes only appear shortly after the new lake or ocean basins, which are in contact with the ice, appear or vanish. From a numerical point of view, however, this is not surprising, as the numerical representation of a physical system requires the underlying equations to be smooth functions. These crashes, due to local unsteadiness in the fields, are known by the developers of PISM to appear².

305 With all the simplifications described above, the *LakeCC* model is not designed to exactly reconstruct the water level of proglacial lakes, nor to estimate freshwater fluxes. It rather gives a crude approximation of where potential lake basins are located and what their maximum water level is. Therefore, we assume the gradual filling mechanism using a (rather arbitrarily chosen) constant fill rate does not result in lake levels that are less valid.

310 If an almost instantaneous filling or draining of lake basins is desired, this could be achieved by setting a high fill rate γ . This would most likely require a reduction of the duration of the time steps, though, which increases the computational cost. In sensitivity tests, which are found in the supplementary material, several higher values were chosen. For these tests, no model crashes were observed. This might either be because we restricted the time step to 0.25yr for all experiments, or simply because no critical situations were triggered. The results of these experiments show no fundamental difference to the *LAKE* experiment. Future implementations could target a more realistic treatment of basin filling.

²<https://pism-docs.org/wiki/doku.php?id=kspdiverged>, accessed: 2021/08/09

To test the impact of the LakeCC model on the ice dynamics, we choose to simulate the glacial retreat of the North American ice sheets after the LGM at 21 kaBP. The computational domain covers the North American continent north of $\sim 35^\circ\text{N}$ and Greenland and spans a rectangle of 7800 km \times 6600 km (see Fig. 3). The resolution of the grid is 20 km, and has dimensions of 390 \times 330.




Figure 3. Map giving an overview of the experiment domain of the current study. Present day coastline and the major North American lakes are shown in black, whereas Canadian and U.S. American states political boundaries are gray. The outlines of selected palaeo proglacial lakes are shown in red: Ag — Lake Agassiz/Ojibway (Teller and Leverington, 2004), MC — Lake McConnell (Smith, 1994) and Pe — Lake Peace (Mathews, 1980; Hickin et al., 2015). Initial ice extent at LGM (Gowan et al., 2016) is shown in blue. Black rectangles visualize the map sections discussed in the text (the numbers refer to the respective figures). The shaded area shows the regions attributed to the (continental) Laurentide Ice Sheet (LIS) in this study.

320 Initial conditions are taken from the NAICE palaeo ice and GIA reconstructions from Gowan et al. (2016). To obtain the palaeo topography, pre-computed bed deformation is added to the present day topography dataset RTopo-2 (Schaffer et al., 2016). ~~The From the~~ temporal evolution of the topography provided by NAICE ~~can be used to calculate the uplift rates, uplift rates can be calculated~~ that are used to ~~used to initialize the Lingle-Clark initialize the Lingle-Clark~~ bed deformation model of PISM. Geothermal heat flux is considered constant in the domain over time and interpolated from the dataset of Davies (2013).

325 As described in Sect. 2.1, the atmospheric forcing model needs two climatic states between which the transient climate is interpolated. Both climate states are monthly means of the surface air temperature and precipitation fields from steady state experiments with the climate model COSMOS. Details on these runs can be found in Zhang et al. (2013). The PD climate state is the control run from that reference, while the LGM experiment is described in Hossain et al. (2018). It is determined using the same protocol, but uses the LGM reconstruction from Gowan et al. (2016). When using a glacial index derived from
330 the NGRIP $\delta^{18}\text{O}$ measurements (North Greenland Ice Core Project Members, 2007), the modeled deglaciation is too rapid to identify contributions from the lake model. Instead, the deglacial climate signal was crudely approximated by a linear model (see Fig. A1a). The transition from LGM to PD climate takes place over 12 kyr (e.g. from 21 to 9 kaBP). To prevent ice sheet growths in eastern Siberia and above 3500 m elevation, ~~ice accumulation is prevented by setting (relative to PD sea level), the~~ precipitation to zero in these regions. ~~The maximum elevation threshold was chosen in an ad-hoc fashion. However, current~~
335 ~~LGM reconstructions (e.g. NAICE (Gowan et al., 2016), and ICE-6G (Peltier et al., 2015)) exhibit higher surface elevations only in few mountainous regions.~~

Transient sea level forcing is applied accordingly to the glacial index. It is linearly ~~raised increased~~ from -120 m at the LGM to the contemporary level 0 m (see Fig. A1b).

Before running the experiments, the model needs to be spun-up. This is done by keeping all boundary conditions fixed at
340 LGM conditions and running PISM in a fixed-geometry mode for several thousand years until the temperature field within the ice has equilibrated. The final output is then used to start the experiments.

In total there are ~~four different three main~~ experiment setups: with lakes (~~LakeCC-LCC~~), ~~with lakes and increased calving to reduce the formation large ice shelves (Lakes, No Shelf-LNS), the LAKE~~, the control run without lakes but using the ~~advanced sea level model (Control-Ctrl)~~ ~~SL2DCC model for calculating a proper land sea mask (CTRL)~~, and the PISM default
345 run (~~Default-Def~~ ~~DEF~~). An overview of all experiments is given in Table 1. ~~To control lacustrine calving, the For the LAKE experiment, the lacustrine calving~~ threshold thickness Δh_L , ~~used by the thickness calving mechanism, is varied. It should be noted that these values are chosen rather arbitrarily, and are not based on any observations. For the LCC experiment, Δh_L is set to a quarter of its marine counterpart ($\Delta h_O = 200\text{m}$) in order to reflect the fact that calving rates for freshwater terminating glaciers are reported to be an order of magnitude lower than rates observed for tidewater glaciers (Funk and~~
350 ~~Röthlisberger, 1989; Warren et al., 1995; Skvarca et al., 2002; Warren and Kirkbride, 2003). In the LNS experiment Δh_L was greatly increased to remove almost any floating ice, as large ice shelves like those seen in the LCC experiment are unlikely to have existed.~~ Hereafter, the experiments referenced, if not explicitly stated otherwise, are ~~LC~~ ~~LAKE~~, including lakes, and ~~Ctrl~~ ~~CTRL~~, without lakes. The PISM default run ~~Def~~ ~~DEF~~ produces ocean basins within the continent, which will behave like lakes, but with water level restricted to sea level.

Table 1. Overview of the experiments done for this study. The ~~lake experiments have a different calving thickness threshold Δh_L . The LakeCC model requires use of the SL2DCC model, which prevents ocean basins from forming within the continent.~~

Exp. name	LakeCC	SL2DCC	Notes
LCC* LAKE*)	X	X	$\Delta h_L = 50\text{m}$
LNS CTRL*)	X	X	$\Delta h_L = 500\text{m}$
Ctr* DEF		X	Def -PISM default

*) default experiments being discussed

355 ~~Additionally, a number of sensitivity experiments were run, exploring the impact of different parameters on the results. These experiments test different values for the lacustrine calving thickness threshold, reduced sub-shelf melting, different Earth model parameters for PISM’s bed deformation model and various lake fill rates. More information on these can be found in the supplementary material.~~

360 ~~For all experiments, the maximum time step of the ice sheet model was limited to 0.25 yr. We want to stress that this setting is not requisite by the LakeCC model, but we had several reasons to do so. Estimating the computational overhead of an algorithm (as done in Sec. C) requires the time steps of the compared runs to be similar. Another benefit from limiting the time step is that the occurrence of numerical instabilities due excessively large jumps in water level (see Sec. 2.4.4) is potentially reduced.~~

3 Results

365 Here we present the results of our PISM-LakeCC ~~experiments~~ ~~experiment~~ by focusing on selected aspects of the modeled glacial retreat. Comparison ~~to~~ ~~with~~ the no-lake control experiments emphasizes the impact of proglacial lakes on the ice dynamics and ice sheet reconstruction. The ~~continental portion of the LIS (which is highlighted~~ ~~highlighted area~~ in Fig. 3) ~~is marks~~ the region of interest ~~of~~ ~~for~~ this study. In the following, ~~if not stated otherwise~~ ~~when referring to the LIS~~, we will ~~refer to it as LIS~~ ~~consider only the parts of the ice sheet that are within this region~~. Overview maps of the entire domain are provided
370 in the ~~Supplementary materials to present~~ ~~supplementary materials to show~~ the temporal evolution of all experiments. These plots show the topographic ~~setting~~ ~~configuration~~ for time slices every 500 yr.

~~In the following, all expressions of time are given in model years, relative to the start of the experiments, and ice sheet volume is quantified as sea level equivalent (SLE). SLE denotes the rise of the water level, if the ice volume V_{IS} would melt and the equivalent freshwater volume V_{fw} is added into a basin of the mean ocean area $A_O = 3.625 \cdot 10^8 \text{km}^2$:~~

$$375 \quad \Delta_{\text{SLE}} = V_{\text{fw}}/A_O = V_{\text{IS}} \frac{\rho_i}{\rho_{\text{fw}}}/A_O, \quad (2)$$

~~where $\rho_{i, \text{fw}}$ are the respective densities of ice and freshwater. Note, this formulation neglects any changes in the ocean surface area and the geoid due to mass redistribution. Furthermore, the ice sheet volume accounts for both grounded and floating ice.~~

Figure 4 shows the temporal evolution of the ~~sea-level-equivalent (SLE)~~ SLE ice volume of the LIS for all experiments. Starting from the LGM initial state after the spin-up, the ice sheets laterally expand and gain mass for about 6 kyr. Within this
 380 time, the LIS almost doubles its volume, before it retreats during the rest of the simulation. As the ice margin laterally expands, some shallow lakes that formed immediately after the simulation started are quickly overrun by the ice and are removed.




Figure 4. Temporal evolution of the ice volume of the continental LIS in the different experiments. The spatial extent that is attributed to the LIS is shaded in Fig. 3. Ice volume is given in sea level equivalent (SLE) ~~, and also includes ice shelves~~ units; see main text for definition. ~~SLE gives the rise of the water column if the melt water of the ice volume is distributed over the mean ocean surface area.~~ Note that both control experiments were run 5 kyr longer than the ~~lake experiments~~ LAKE experiment.

~~Fig.~~ Figure 5 shows the temporal evolution of the ice margins and grounding lines at 90° W to compare the northward retreat of the LIS between the different experiments. The ~~southward-shifted~~ southward-shifted grounded ice margin between about 1 and 3.5 kyr is due to formation of ~~small~~ ice lobes that advance into small lakes (compare also with Fig. 6). At ~ 3.5 kyr the ice margin is ~~furthest south before the furthest south, after which~~ it rapidly retreats ~~northwards~~ northward. The presence of a lake initiates the retreat about 1 kyr before the control runs. Topography revealed by the retreating ice is left deeply depressed and is immediately occupied by proglacial lakes following the ice margin. From about 6 kyr, an inner-continental ocean basin shows up in the PISM default experiment (~~Def~~ DEF), where the ~~bed-topography~~ bed-topography is below sea level, which increases the ice retreat compared to the ~~Ctrl~~ CTRL experiment, closely following the LCC ice margin. ~~The LNS run exhibits the highest retreat rates, which becomes even more pronounced when the ice margins pass the Great Lakes region, where water depth at the grounding line reaches up to 750 m~~ ice margin of the LAKE experiment.
 385
 390

Figure 7 shows a comparison of the ~~LCC and Ctrl~~ LAKE and CTRL experiments at 7 kyr. The lakes' impact on the ice sheet dynamics and geometry is obvious. Where the ice sheet margin terminates ~~, even in small proglacial lakes, the ice surface velocity anomaly (Fig. 7b) shows a strong increase.~~ The increased ice flow and mass loss results in a drastically lowered ice




Figure 5. Temporal evolution of the ice margins and grounding lines at 90°W for the different experiments. Ice retreat is relative to the initial LGM ice front position. The vertical dashed lines at 2.2 and 13 kyr highlight the timing of the plots in Figs. 6 and 9, respectively. To provide some geographical context, the locations of Lake Superior’s and Hudson Bay’s contemporary basins are marked by the blueish stripes.




Figure 6. Formation of ice lobes at the southern ice margin (at 2.2 kyr) where lake boundaries promote accelerated ice flow. Ice extent of the control experiment is shown in red. The pixelated appearance of this plot comes from the 20 km model resolution.

395 profile (Fig. 7a) and accelerated retreat of the ice margin. These effects can be traced back several hundreds of kilometers upstream. ~~Where ice streams diverge, the surface velocity anomaly partially shows a slowdown of ice flow.~~




Figure 7. Comparison of the standard-lake (LCC)-LAKE and no-lake (Ctrl)-CTRL experiments at 7 kyr. Both plots show anomalies (LCC LAKE - CtrlCTRL) of (a) ice thickness and (b) surface velocity. Formation-Acceleration of ice streams-flow at the proglacial lake boundaries and the resulting impact on ice geometry can clearly be seen. Ice margins of both simulations are shown in black.

From 7 to 9 kyr, the Great Lakes region is covered by a single proglacial lake, which shares an up to 2000 km long boundary with the ice sheet (compare with Fig. 8a). At around 9 kyr the water level of this lake rapidly dropped, as a lower outlet to the Atlantic became ice-free. The successor lake occupies the basin of Lake Agassiz/Ojibway (see Fig. 8b and c). Where the water depth is sufficient, the ice margin floats. Figure 9 shows profiles of the ice margin of the LCC-LAKE and the two control experiments at 13 kyr along the 90°W transect. Note that only a few kilometers behind the grounding line, the bed elevation declines into the Hudson Bay basin. As soon as the grounding line retreats into this deep basin, the lake quickly expands underneath the ice sheet (compare with Fig. 5 and Fig. 8c and d), forming an enormous ice shelf. This dramatically accelerates the glacial retreat over Hudson Bay, which can also clearly be seen in the ice volume evolution of the LIS (Fig. 4), as the lake and no-lake experiments rapidly diverge. At around 17.9 kyr, the ice saddle over Hudson Strait breaks apart and allows the lake to drain into the Labrador Sea (Fig. 8e). In the LNS experiment this happens roughly 2 kyr earlier. We want to note that the sudden jump of the ice margin (~ 500 km), seen at this time in Fig. 5, stems from the ice margin vanishing laterally out of the 90°W profile. Due to GIA processes, the-an ice-free Hudson Bay basin eventually rises-emerges above sea level (see Fig. 8e and f). For both control experiments, the Hudson Bay area-Area is still covered by a thick ice dome.

410 The drainage route towards the Arctic is blocked until the ice saddle ~~;~~ connecting the northern LIS and the [CIS, Cordilleran Ice Sheet \(CIS\)](#) collapses. Along the southern margin of this ice saddle, a vast proglacial lake forms (see Fig. 8d and e). At around 21 kyr, the saddle collapses, which drains most of the lake (Fig. 8f). ~~Again, for the LNS experiment this happens about 2 kyr earlier.~~ For the control experiments, ~~Def and Ctrl~~[DEF and CTRL](#), this separation occurs at about 24 kyr and 26 kyr, respectively. This event marks the final retreat of the LIS ~~for both lake experiments~~ [in the LAKE experiment](#) (Fig. 4).

415 4 Discussion

4.1 Transient experiments

Starting from LGM initial conditions, the ice sheet takes several thousand years to come to a dynamical equilibrium, as the initial ice sheet was reconstructed using geological evidence of ice margin history and GIA observations, and no ice dynamics were included (Gowan et al., 2016). This growth can be seen in all experiments: the ice sheet accumulates mass and the ice margins advance. After around 6 kyr the maximum ice volume is achieved for the LIS (see Fig. 4). The reason for the ice accumulation is assumed to be due to the climate forcing. By linearly interpolating between the warm, humid PD and the cold, dry LGM climate states, unrealistically high accumulation is produced, especially in cold regions and on top of the ice sheet. Only by limiting precipitation above 3500 m elevation and over Siberia does further expansion of the ice sheets stop. For a more realistic behavior, coupling with a climate model might be needed. As the climate is shifted from LGM towards PD conditions, the southern ice margin retreats ~~northwards~~ [northward](#). Reconstructions of the retreat of the western LIS show that it proceeds in an east to north-eastern direction, and the separation from the CIS happened relatively early ~~(see e.g. Dyke, 2004; Gregoire et al., 2012; Peltier et al., 2015; Gowan et al., 2016)~~ [\(see e.g. Dyke, 2004; Gregoire et al., 2012; Peltier et al., 2015\)](#). Our deglacial scenario fails at simulating realistic ice margin positions for the western LIS.

Without adding more advanced climatic feedbacks a realistic deglacial reconstruction is not expected. This, however, has not been the focus of this study, which is to test the PISM-LakeCC model and studying its impact on ~~the~~ ice dynamics and ~~the~~ glacial retreat. For these purposes the experimental setup is sufficient. Analyzing the interplay between ice sheets and proglacial lakes in more realistic setups, ~~e.g. e.g.~~ fully coupled to a climate model, and comparing against various geological proxy data is an interesting topic for future research.

4.2 Lake reconstructions

435 ~~Reconstruction~~ [The reconstruction](#) of palaeo proglacial lakes depends mainly on GIA and ice margin locations, as these define the extent [and depth](#) of lake basins. Resolution issues are assumed to be secondary in this context and are tackled by applying the preprocessing method described in Hinck et al. (2020). As already discussed in the previous section, the ice margin retreat modeled in our experiments does not match well with reconstructions based on geological evidence. For this reason, the lake reconstructions are not expected to match well with observations. Drainage towards the Arctic, for example, is blocked until 440 the ice saddle connecting the LIS and the CIS collapses, which results in a large proglacial lake in western Canada between 15

and 21 kyr. This lake is located where glacial lakes McConnell (Smith, 1994) and Peace (Mathews, 1980; Hickin et al., 2015) existed, but its basin does not match the outlines of the palaeo lakes (see Fig. 8d and e).

Another issue, ~~that~~ which is also partly related to drainage, is the immense size and depth of some lakes that appear along the southern ice margin. From around 6 to 9 kyr one large lake occupies the entire Great Lakes region. Later, it transitions into the basin of Lake Agassiz/Ojibway (Teller and Leverington, 2004) before expanding into Hudson Bay (~ 13 – 18 kyr) (Fig. 8a-d). Maximum water depths close to the grounding-line grounding line are up to 1000 m. ~~The main reason for this, we assume, is the GIA response modeled by~~ As a result of the increased accumulation of ice due to the simplified climate forcing, the topography is further depressed. The deeply depressed topography in combination with lowered ice thickness, due to dynamic thinning, allows the lake to further expand underneath the ice sheet.

450 Furthermore, for such deglacial scenario, use of a more advanced GIA model might be needed. PISM's ~~bed-deformation model. This simple, two-layered Earth model is not capable of handling the extreme deglaciation scenario of an entire continent. Since the climate forcing tends to accumulate too much ice, this situation is further exacerbated in our simulation. During glacial retreat,~~ default model is based on a simple two-layered Earth model. Viscosity variations in the upper and lower Earth mantle are not included in this model, even though these variations significantly contribute to the GIA signal in central Canada
455 (Wu, 2006).

Also, the initialization of the ~~bed was deeply depressed and the rebound was not quick enough. After deglaciation, the Hudson Bay region is over-relaxed and above PD sea level~~ Lingle-Clark model from a glaciated state is problematic here. For the model to calculate a relief topography, to which bed deformation is applied, it should ideally be initialized from an interglacial state when the residual GIA from previous glaciations is limited. Test runs, comparable to Niu et al. (2019)
460 , however, suffered from the fact that the bed deformation along the southern ice margin was so deep, that the basin was connected to the Atlantic Ocean, which consequently inhibited the formation of lakes. We therefore chose to initiate the experiments from NAICE LGM reconstructions. The mismatch in calculated relief topography, results in ice free regions to over-relax. Hudson Bay, for example, is elevated above sea level at PD (see Fig. 8f).

4.3 Lacustrine impact on ice dynamics

465 ~~Despite the discrepancy between model results and geologically inferred deglacial history~~ In comparison with the CTRL experiment, the modeled ice dynamics reveals a ~~large response~~ high sensitivity to the presence of proglacial lakes. The combination of modified stress regime, changed ice geometry and increased frontal ablation leads to accelerated ice flow upstream of the lake boundary. ~~The formation of ice streams~~ This acceleration and the associated lowering of the ice surface is shown in the ice thickness and surface velocity anomaly plots of Fig. 7. It should be noted that ~~, due to the changes in ice dynamics, the different experiments constantly diverge and features visible in these plots might therefore only indirectly be triggered by the lake boundary, but rather be a result of differences in ice sheet geometry.~~ some features in these plots also arise from the fact that the ice geometries of the experiments diverge. Sensitivity runs (see supplementary material) have shown that the observed response is strongly impacted by the grounding line treatment. If the basal resistance at the grounding line is not reduced, the

grounding line flux is strongly decreased. This becomes aparent in reduced mass loss and slower retreat of the ice margins.
475 Nevertheless, when compared to the CTRL experiment, the impact on ice flux and margin retreat is clear.

Another feature that can be observed in the model results is the formation of ice lobes (Fig. 6). ~~We assume that this is due~~
Due to the enhanced sliding at the lake boundary, ~~and~~ an advance is locally promoted. However, these lobes only appear at
shallow lakes and are not comparable in size with major ice lobes of the LIS (e.g. Hooyer and Iverson, 2002; Dyke, 2004).
When water depth of the proglacial lake becomes too deep, the thin advancing ice front is ~~presumably~~ lost due to calving. A
480 calving law adapted for lakes, as mentioned in Sect. 2.4, could help ~~overcoming this problem.~~ improve modeling this aspect.
The application of a more realistic GIA model may also aid in the development of these lobes, as the lake depth at the grounding
line margin will be reduced.

The increased thinning of the ice sheet interior gains special importance at a later stage of the glacial retreat. ~~At places where~~
~~ice is thin enough that it starts floating, the proglacial lake can,~~ as it allows the lake to expand underneath the ice sheet, ~~where~~
485 the ice is thin enough to float. Where bed topography is deeply depressed, this process is further accelerated (Fig. 8c - e). On the
reverse sloping bed of Hudson Bay, no stable grounding line position can be achieved (see Fig. 5), leading to the rapid disinte-
gration of the ice saddle blocking drainage through Hudson Strait. This process is very similar to the marine ice sheet instabil-
ity ~~(MISI, Weertman, 1974; Thomas and Bentley, 1978).~~ (MISI, Weertman, 1974; Thomas and Bentley, 1978; Schoof, 2007).
Sensitivity experiments (see supplementary material) show that this rapid disintegration of the LIS is mainly driven by the
490 strongly negative mass balance along the southern ice margin. Calving and sub-shelf melt rates play only a secondary role
here. These findings are in agreement with Quiquet et al. (2021), who refer to this instability as PLISI.

We want to note that the vast ice shelves that can be observed at this late phase of the collapse of the LIS (Fig. 5 & 8c-d)
might be unrealistic. At least we are not aware of any geological evidence or reconstruction, indicating such ice shelves existed.
It is possible that the simulated grounding line flux is too high or the calving rate too low. Two sensitivity runs checking
495 these parameters (IncCalv an nSG in the supplementary material) do not exhibit such large lacustrine shelves. However, more
research is necessary to decide how to best parameterize these issues.

4.4 Implications on the demise of the LIS

Although the modeled glacial retreat of the North American ice sheets does not match up with reconstructions based on
geological observations, the importance of proglacial lakes on the glacial retreat is evident. Comparing to the two no-lake
500 experiments with the ~~LCC and LNS experiments,~~ a LAKE experiment, totally different ice ~~geometry exists~~ geometries exist
at the end of the ~~experiment runs.~~ The presence of lakes triggers the rapid collapse of the LIS over Hudson Bay. Furthermore, it
accelerates the disintegration of the ice saddle connecting LIS and CIS. ~~When~~ Even after extending the simulations for both
control runs ~~were extended~~ by another 5kyr, Hudson Bay still remained ~~ice covered~~ ice-covered.

~~During the simulated glacial retreat, the climatic forcing is the dominant factor that affects the ice margin position. Lakes,~~
505 ~~however, do impact the early retreat by inducing the formation of ice streams, which drain the ice sheet interior. As soon~~
~~as the grounding line retreated into deeper inward-sloping topography, the proglacial lakes become the dominant factor that~~
~~destabilizes the ice sheet.~~

4.5 Outlook

510 ~~The simple implementation described has potential for further improvement. Reconstruction of lakes could benefit from more realistic accounting of water fluxes. Also, instead of assuming the marine parameterizations to be valid at the lacustrine boundary, formulation of mechanisms adapted for a lake-ice boundary, such as a more physically motivated calving model valid for grounded and floating ice termini or a lacustrine sub-shelf melting model, would improve the ice-dynamical response to lakes. Nevertheless, applying this simple model instead of a dry or a marine boundary at sea level (possibly hundreds of meters below the actual water level) will produce a more realistic depiction of the ice sheet retreat. Coupling the lake~~
515 ~~reconstructions into climate and GIA models would also yield a more realistic deglacial scenario.~~

5 Conclusions

In this study we have described the implementation of an adaptive lake boundary condition into a 3D thermo-mechanically coupled ice sheet model. To the best of the authors' knowledge this has not been done in any other study. Application of this model to a simple North American deglacial scenario, starting at ~~the LGM~~ LGM conditions, shows the importance of proglacial
520 lakes for the proper modeling of the ice dynamics. The lake model promotes the ~~formation of continental ice streams, which play an important role in the acceleration of ice sheet's mass balance. By triggering a feedback cycle similar to the marine ice sheet instability, the flow where the ice margin is bounded by a lake, which leads to a lowering of the upstream ice surface. In a warming climate, the lowered ice surface is exposed to higher temperatures and thus higher melt. This feedback mechanism becomes most obvious once the grounding line enters a deep basin on a retrograde bed. The ice sheet exhibits an instability similar to the MISI, which is called accordingly PLISI. Occurrence of the PLISI results in the~~ final break-up of the LIS ~~occurs~~.
525 Our experiments show a rapid expansion of the lake into the still glaciated basin of Hudson Bay, ~~quickly breaking up followed by a quick disintegration of~~ the ice barrier that blocks the lake's drainage through Hudson Strait into the Labrador Sea. ~~We would therefore~~

Due to the significant impact of the LakeCC model on the ice sheet dynamics, we would recommend the use of an adaptive
530 lake boundary condition when modeling the deglaciation of land terminating ice sheets.

Code availability. All implementations described in this study are based on PISM v.1.2.1. The modified code can be found in SH's GitHub repository (<https://github.com/sebhinck/pism-pub>). Implementations of the various independent models are found in separate branches. The code of the modified PISM version, as it was used in this study, is archived online (Hinck and PISM Authors, 2020).

Appendix A: Configuration parameters used for experiments

535 Experiments for this study were done using a modified version of the Parallel Ice Sheet Model (PISM; the PISM authors, 2015; Bueler and Brown, 2009; Winkelmann et al., 2011). Sections 2.1 and 2.5 describes the used configuration of the ice

sheet model. The hierarchy of the parameterizations for PISM’s different sub-models is listed in Table A1. Configuration

Table A1. PISM sub-models used for the experiments.

Coupler	Model(s) used
atmosphere	index, precip_cutoff ^{*)} , <u>precip_cutoff</u> ^{*)}
bed_def	lc
calving	eigen_calving,thickness_calving
hydrology	null
lake_level	(lakecc) ^{*)}
ocean	pik
sea_level	constant,delta_sl(<u>sl2dcc</u>) ^{*)}
stress_balance	sia+ssa
surface	pdd

*) Models added for this study.

parameters that diverge from PISM’s default configuration are shown in Table A2. Details about PISM’s sub-models and the default configuration can be found in the user manual.

540 The temporal evolution of glacial index, used to interpolate climate forcing between LGM and PD states, and the global sea level are, as described in Sec. 2.5, approximated using a linear transition from LGM to PD climate within 12 kyr. Both curves are shown in Fig. A1 and are shown next to commonly used proxy data.

Table A2. Configuration parameters used for the PISM experiments that diverge from the defaults.

Option	Default value	Used value	Units
-lakecc_dz ^{*)}	1	5	m
-lakecc_zmin ^{*)}	0	-300	m
-Lbz	1000	2000	m
-Lz	4000	5750	m
-max_dt	60	0.25	yr
-Mbz	1	11	-
-meltfactor_pik	0.005	0.01	-
-Mz	31	101	-
-pik	false	true	-
-precip_cutoff_height ^{*)}	3000	3500	m
-pseudo_plastic	false	true	-
-sia_e	1	5	-
-stress_balance.sia.max_diffusivity	100	200	m ² s ⁻¹
-tauc_slippery_grounding_lines	false	true	-
-temp_lapse_rate	0	7.9	K km ⁻¹
-thickness_calving_threshold	50	200	m
-till_effective_fraction_overburden	0.02	0.01	-
-use_precip_cutoff_height ^{*)}	false	true	-

^{*)} Commandline options added by one of the models described in this study.




Figure 8. Snapshots for different stages of the lake-LAKE experiment (LCC). Ice-Lacustrine ice shelves are shown in light blue. The red outlines show the maximum extent of selected palaeo lakes (Fig. 3). Modeled spill point and drainage routes for Lake Agassiz are shown in green in panels a-d.




Figure 9. Profiles of the ice sheets of the (a) ~~standard lake (LCC)~~ *LAKE*, (b) ~~standard no lake (Ctrl)~~ *CTRL* and (c) ~~PISM default (Def)~~ *DEF* experiments at 13 kyr. The transects shown in (d) are along 90°W , between 53°N (A) and 58°N (B) and are marked in the upper panels by a red line. In panels (a) - (c) ice shelves are shown in light blue, lakes in blue, and ocean in gray. The blue line in panel (d) marks the lake level elevation in the *LAKE* experiment, while the gray line marks the sea level elevation of the *DEF* experiment.




Figure A1. The black lines show the temporal evolution of the (a) glacial index and (b) the mean global sea level used in the experiments. Both time series are simplified by a linear evolution from their LGM to PD values between 21 and 9 kaBP and represent a crude approximation of proxy records. For visualization, such proxy records are shown in gray. These are (a) the index derived from the NGRIP $\delta^{18}\text{O}$ measurements (North Greenland Ice Core Project Members, 2007), and (b) the global mean sea level reconstruction from Lambeck et al. (2014).

Appendix B: PISM-LakeCC

This Appendix adds a more technical overview of the model description of the main text. At first, a short description of the general `lake_level` interface within PISM is given. As the LakeCC model, and also the SL2DCC model, make extensive use of functions based on the ~~Connected Components Algorithm~~ connected components (CC) algorithm (see also Hinck et al., 2020), its implementation is delineated in the following section. The last two sections address details of the implementation of the LakeCC and SL2DCC models. The names of model components in PISM and of configuration parameters are shown in ~~monospace~~ monospaced font.

550 B1 PISM's lake interface

PISM is designed in a modular way, that prescribes the interface of all sub-models. These modules inherit certain properties and functionality from C++ base classes and extent upon these. This way, different combinations of sub-models can be easily realized via configuration parameters. Furthermore, this facilitates the implementation of new sub-models. For the sub-systems acting as boundary conditions on the ice-sheet, such as atmosphere, surface and ocean, PISM distinguishes between models and modifiers. A model parameterizes all aspects needed for the treatment of the respective boundary, and can be combined with combination of different modifiers, which can apply changes to the model's parameterization.

A lake, as seen by an ice sheet model, is very similar to a locally elevated sea level. It would be possible to implement the lake interface as a sea level modifier in PISM. However, physical differences (e.g. density, temperature distributions) require different treatment of marine and lacustrine environments. To distinguish between both cases, at least two spatial maps are required. This can either be done by providing the lake and sea level combined in one field and additionally providing a mask, or providing lake and sea level elevations as separate maps. For the implementation of the PISM-LakeCC model, we chose the latter case.

The `lake_level` interface is based on PISM's `sea_level` interface. It provides a 2D field `lake_level_elevation` to the ice sheet model, which contains each cell's lake level elevation in meters above the reference geoid. Where no lake level is set, the cell is set to invalid, and a fill value is inserted, which is a large negative number (it is set by the configuration parameter `output.fill_value`, which has a default value of $-2 \cdot 10^9$). If no lake model is chosen, the entire returned field is set to invalid.

As the ice sheet and lake geometry is dynamic, it turned out that it is beneficial to spread information about the presence of lakes to neighboring cells. Lake models can therefore ask the lake interface to expand lakes one cell beyond their reconstructed basin, which can be regarded as kind of ghost cells. This does not change the actual lake geometry within the ice sheet model, because cells are considered dry, where the water level is below bed elevation or the ice sheet is grounded. The lake level elevation field can therefore rather be regarded as a virtual lake level.

Where the lake level is valid and above bed elevation, a grid cell is treated as a lake. The lacustrine boundary is essentially treated the same as the ocean, except of the different water level and adapted water density. A more advanced boundary treatment would need to be implemented for ocean (or rather wet boundary) and calving parameterizations. So far, only the thickness

calving mechanism accepts a distinct lacustrine parameter for the thickness threshold (`calving.thickness_calving.threshold_lakes`). Note, contrarily to marine ice shelves, lacustrine ice shelves are not excluded from the calculation of the potential sea level rise potential.

For restarting, the variable `effective_lake_level_elevation` is added to each restart file. ~~Furthermore, two~~
 580 ~~additional~~ Optionally, the two 2D diagnostic variables ~~, which lake depth and lake level real~~ can be added to PISM's output ~~, were added~~ (see Table B1). Outside ~~of~~ where the lake level is actually above bed elevation, both fields are set invalid.

Table B1. Newly added diagnostic variables. Some diagnostic fields are generally available, independent of the sea-level or lake model chosen.

Name	Model	Description
<code>lake_depth</code>	general	Water depth of lakes.
<code>lake_level_real</code>	general	Water level of lakes, where above bed elevation.
<code>lakecc_gradual_target</code>	LakeCC	Field <code>target_level</code> , as used internally by the LakeCC model.
<code>ocean_depth</code>	general	Water depth of the ocean, where sea-level above bedrock.
<code>sl2dcc_gradual_target</code>	SL2DCC	Field <code>target_level</code> , as used internally by the SL2DCC model.

B2 Connected Components

PISM-LakeCC and PISM-SL2DCC make extensive use of the connected components algorithm. The algorithm and its use
 585 in the LakeCC standalone model is described in Hinck et al. (2020). Parts of the models are implemented analogously, i.e. the algorithm is used to determine the target lake level (using the LakeCC method) and the ocean mask (using the SL2DCC method). However, implementation into a dynamic ice sheet model needs a more advanced treatment than the standalone tool.

In several parts of the algorithm, non-local information of the entire lake is needed. The connected component algorithm, when adapted accordingly, is capable of gathering different information and attributing it to the respective lake basins. In
 590 the following we shortly introduce the C++ classes based on this algorithm, which are used in the implementation of the PISM-LakeCC and PISM-SL2DCC models.

The `LakeLevelCC` class determines the maximum fill height of lake basins by iteratively checking the entire domain for a set of increasing water levels, as described in Hinck et al. (2020). The `SeaLevelCC` class is also implemented as described in that reference. It identifies basins in the topography, which are below the global sea level but are not connected to either
 595 of the domain margins ~~, (compare Fig. B1a).~~ The `IsolationCC` class (Fig. B1b) also marks cells as invalid that are either ~~ice-covered~~ ice-covered or not connected by an ice-free corridor with the domain margin. It is used to restrict the formation of lakes in the ice sheet interior and of subglacial lakes where thin ice covers a deep basin. To remove narrow lakes, which are often related to under-resolved topography, the `FilterLakesCC` class checks the lakes' geometry. Only lakes, which contain




Figure B1. Sketches depicting the different algorithms implemented. a) `SeaLevelCC` – only basins below sea level and connected to either of the domain margins is considered ocean. b) `IsolationCC` – Lake basins fully enclosed by the ice are considered invalid. c) `FilterLakesCC` – Lakes are only retained if they contain at least one cell with a certain amount of lake neighbors. In the example $N=4$ was chosen. d) `LakePropertiesCC` – This algorithm collects the minimum and maximum water levels for each lake basin. e) `FilterExpansionCC` – This algorithm compares the current lake mask with the previous mask and labels newly appeared and vanished lake cells. It distinguishes between narrow strips (purple) and wider basins (dark blue).

one (or more) cells that have at least a certain amount of neighbors that also are part of that lake, are retained –(Fig. B1c).
 600 `LakePropertiesCC` collects the minimum and maximum current water level of each lake basin –(compare Fig. B1d). The
 class `FilterExpansionCC` (Fig. B1e) is used to compare the new lake basins with the state of the previous time step.
 This method returns a mask that marks cells that were newly added or have vanished, but it also distinguishes the basin shape,
 similarly to `FilterLakesCC`. Furthermore, for each new lake basin the minimum bed elevation and, if it was an ocean basin
 in the previous time step, sea level are returned. This information is needed to catch different scenarios when initializing the
 605 lake level and treat them accordingly.

The difference of the implementation of the underlying connected components algorithm to Hinck et al. (2020) is the use
 of PISM’s parallelized data types. 2D fields are partitioned and distributed over several processors using the library *PETSc*
 (Balay et al., 1997, 2019). This implementation requires some extra steps, as processors need to exchange information with the
 adjacent processor domain.

The PISM-LakeCC model utilizes the LakeCC algorithm (Hinck et al., 2020) to determine the maximum water level of lake basins. The topographic and glacial state, as needed for the lake reconstruction, is provided by PISM. Options used by the model are set by configuration parameters (see Table B2) and are explained in the following. Cells with ice thinner than δ_{if} are considered ice-free.

Table B2. Configuration parameters read by the LakeCC model. Prepend `lake_level.lakecc.` to the parameter to get the full name.

Parameter	Type	Default	Symbol	Description
<code>dz</code>	Number	1m	Δz	Spacing between successive water levels.
<code>filter_size</code>	Integer	4	N_{filter}	Number of neighboring cells used by filter algorithm. See text for details.
<code>ice_free_thickness</code>	Number	10m	δ_{if}	Threshold ice thickness below which a cell is considered ice-free.
<code>init_filled</code>	Boolean	<code>false</code>	-	Bootstrap lakes as filled.
<code>keep_existing_lakes</code>	Boolean	<code>true</code>	-	Keep existing lakes, even though they are surrounded by ice.
<code>max_fill_rate</code>	Number	1m yr^{-1}	γ	Fill rate used by LakeCC gradual fill algorithm.
<code>topg_overlay_file</code>	File	-	-	File containing field <code>topg_overlay</code> .
<code>zmax</code>	Number	1000m	z_{max}	Maximum water level to check.
<code>zmin</code>	Number	0m	z_{min}	Minimum water level to check.

615 At initialization of the model, the lake level is read from the PISM input file (`effective_lake_level_elevation`). This is necessary to guarantee a smooth continuation of the simulation after model restart. When this field is not available, lakes are either initialized empty, or filled to the brim (configuration flag `init_filled`).

In every time step of the ice sheet model, the ice and topography configuration changes, thus the lake model needs to be updated. Fig. B2 shows a sketch of the update sequence of the PISM-LakeCC model. The basic principle of how the model works was given in Sect. 2. Both fields, target level and lake level are successively refreshed.

620 The process to update the target level (Fig. B2b) resembles that described in Hinck et al. (2020). If a filename is provided using the `topg_overlay_file` option, the field `topg_overlay` is read and applied to PISM's topography before proceeding. Using this processed topography field and PISM's sea level elevation data (which should have been computed using the SL2DCC model) a mask is created that marks all the ocean cells. Another mask is prepared using the `IsolationCC` method. It is called and marks cells as valid if they are ice-free and are connected by an ice free corridor to the domain margin. 625 The LakeCC method only keeps lakes that contain at least one valid cell. This prevents isolated lakes from forming in the ice sheet interior or entirely beneath the ice. It might, however, happen that lakes suddenly marked as invalid when the ice sheet advances and cuts them off from the exterior. If this is not desired, the option `keep_existing_lakes` labels all existing lakes as valid. With everything prepared, the lake basins are computed using the LakeCC algorithm and their maximum water level is kept as the target level. 630 The water levels that the algorithm iteratively checks are equally distributed between z_{min}




Figure B2. (a) Flowchart diagram showing the steps of the Pism-LakeCC update process. The red and blue boxes expand to the diagrams on the right, (b) and (c), with correspondingly colored outline.

and z_{\max} , with a spacing of Δz . As the target levels are computed using the modified topography, the lake basins have to be transferred back onto PISM's topography. Finally, to get rid of narrow lakes, which are often caused by the under-resolved topography, the target level is filtered by applying the `FilterLakesCC` method using a filter size of N_{filter} . The target level can be added to the output for diagnostic purposes (see Tab. B1).

635 In the next step, the lake level, which is the field that is accessed by other parts of the ice sheet model, is updated (Fig. B2c). Before gradually filling each basin to its maximum fill level, which is set by the target level, newly added lake cells need to be initialized. Therefore, information about existing lakes within the lake basins and about patches of newly added lake cells is collected using the `LakePropertiesCC` and `FilterExpansionCC` methods. In the initialization process (sketched in Fig. B3), these data are used to ensure that all cells of a lake start at the same water level. However, this is not always-always

640 possible. New lake basins should ideally be initialized empty, with a water level at its deepest point. Another case is when an existing lake is enlarged by just a few cells (e.g. because the ice margin retreats). This gap should be initialized at the actual lake level. To identify this case, the `FilterExpansionCC` method considers the shape of the newly added patch similarly to the `FilterLakesCC` method: if at least one cell of the patch is entirely surrounded by other cells, also belonging to that patch, it is marked as 'wide', otherwise as 'narrow'. To omit instabilities due to rapid change of the water level, 'wide' basins are

645 treated as new lakes and initialized empty. In case the lake basin was previously part of the ocean, the lake level is initialized at the sea level, to omit a rapid jump in the boundary conditions. Once the lake level is initialized, the water level can be gradually adapted towards the target level using the constant fill rate γ , as described in the main text.




Figure B3. Flowchart diagram of the initialization process, which is executed for each cell of the grid. Cells that have just been assigned a lake level (i.e. it was set invalid in the previous time step) are initialized. Interconnected clusters of these cells are grouped as patches by the `FilterExpansionCC` method and are labeled accordingly to their shape ('narrow' or 'wide', see main text). Lake level is initialized either to the minimal bed elevation of the new basin (`min_bed`), to the lake level of a connected existing lake (`ex_lake_level`) or to sea level (`sea_level`).

B4 SL2DCC

Table B3. Configuration parameters read by the SL2DCC model. Prepend `sea_level.sl2dcc.` to the parameter to get the full name.

Parameter	Type	Default	Symbol	Description
<code>max_fill_rate</code>	Number	1m yr^{-1}	γ^{SL}	Fill rate used by SL2DCC gradual fill algorithm.
<code>sl_offset</code>	Number	10m	δ_{SL}	Sea level offset.
<code>topg_overlay_file</code>	File	-	-	File containing field <code>topg_overlay</code> .

In PISM the sea level field is provided as a 2D field, which usually is set to the global mean sea level. Here, we present the
650 implementation of a sea level modifier, which takes advantage of ~~possibility~~the possibilities of a spatially variable sea level
field and removes isolated ocean basins. The model is based on the connected components (CC) algorithm as it was described
in Hinck et al. (2020) and Sect. B2. Using the `SeaLevelCC` class a mask is determined with all isolated ocean basins marked.
These basins are removed from the sea level field by setting the respective cells to a fill value, analogously to the lake interface
described in Sect. B1. The configuration parameters read by the model are listed in Table B3.

655 Similarly to the `LakeCC` model (Sect. B3), the low-resolution topography field from the ice sheet model can be overlain
by an input field read from `topg_overlay_file`. These fields are then used by the `SL2DCC` model to identify isolated
ocean basins. Potential ocean cells (cells that fulfill the flotation criterion) are grouped into inter-connected patches using the
`SeaLevelCC` method. Only patches that are connected to the margin of the computational domain are considered to be part of
the ocean. All other patches are treated as isolated inland ocean basins. When categorizing all ocean basins, the `SeaLevelCC`
660 method internally raises the water level by an offset δ_{SL} . It is only used internally to raise the water level when checking for
connected ocean basins. This treatment effectively extends the ocean mask slightly beyond the coastline. Basins near the coast,
which would otherwise be labeled as isolated, are identified as regular ocean basins. This is done to account for the relatively
low resolution topography.

Defining sea level spatially constant over the entire domain has the advantage that no sudden jumps in water level occur, that
665 cause numerical problems. Using the `SL2DCC` model, however, ocean basins are dynamically added and removed, depending
on geometric considerations using the evolving glacial topography. For this reason, the implementation of the `SL2DCC` model
needs to take care of smoothly applying changes to the water level. When using the `LakeCC` model (Sect. B3), the lake model
takes care of this. If the `LakeCC` is not used, the `SL2DCC` model gradually adjusts the water level using a constant fill rate γ^{SL} .
This algorithm is very similar to the gradual filling algorithm used in the `LakeCC` model.

670 Appendix C: Efficiency of the PISM-LakeCC model

Figure C1 shows the efficiency for all four experiments. Usage of the LakeCC model reduces the efficiency by $\sim 40\%$ compared to the PISM default run. However, as the numbers shown in the figure depend strongly on the adaptive time steps chosen by PISM, the plot does not show the pure computational burden of the LakeCC model. At times ~~where~~when ice shelves are present, smaller time steps are chosen, which can directly be seen in the model's efficiency. This is the reason why the *Ctrl* run is generally more efficient than the *Def* run, although an additional model needs to be evaluated: as no inland ocean basins are present, no ice-shelves appear and ice velocities in general are slower.




Figure C1. The plot shows the efficiency of PISM in model years per processor hour averaged over 200 yr chunks for the different experiments: ~~Ctrl~~CTRL – no lakes but with the SL2DCC model to prevent inland ocean basins; ~~Def~~DEF – PISM default, no lakes, no advanced ocean treatment; ~~LCC~~LAKE – standard lake experiment; ~~LNS~~IncCalv – lakes, but ~~strong-increased~~strongly-increased calving ~~to-reduce-threshold-such-that~~to-reduce-threshold-such-that ice ~~shelf-formations~~shelves are effectively removed (for more details see supplementary material). ~~variability~~Variability is mainly due to the adaptive time stepping mechanism of PISM which hits especially hard when big ice shelves are present. A maximum time step of 0.25 yr is used for all experiments.

Appendix D: Climate forcing

D1 Glacial index

Table D1. Command-line options for use with the PISM atmosphere model `index`.

Command-line option	Comment
<code>-atmosphere_index_file</code>	File containing climate index time series (<code>glac_index</code>).
<code>-atmosphere_index_climate_file</code>	File containing climate forcing fields for both climate states (<code>i</code> needs to be replaced with the corresponding index (0,1)): air temperature (<code>airtemp_i</code>), precipitation (<code>precip_i</code>), surface elevation (<code>usurf_i</code>).
<code>-temp_lapse_rate</code>	Temperature lapse rate γ_T used in Eq. (D1). Default value: 0.0 K km^{-1}
<code>-atmosphere.precip_exponential_factor_for_temperature</code>	Exponential factor C used in Eq. (D2). Default value: $7.04167 \cdot 10^{-2} \text{ K}^{-1}$

The climate forcing applied for the ~~testrun~~ test run carried out for this study ~~uses~~ uses a custom implementation that is based on the concept of a glacial index. It approximates the temperature T_i and precipitation fields P_i between two climate states, for example between LGM and PI. For simplicity, it is assumed that the spatially varying fields can be linearly interpolated between these states, which are each assigned an index of 0 and 1, respectively. Interpolation between these states is done by weighting them according to a time-dependent climatic index $i(t)$. It is usually derived from measurements of, for example, $\delta^{18}\text{O}$ (e.g. NGRIP (North Greenland Ice Core Project members, 2004), EDML (Augustin et al., 2004)), which is assumed to give a rough approximation of the global mean surface temperature. However, this index can also be artificially designed, for example as a simple linear transition from one state to the other (see Fig. A1a).

The basic idea has already been used in a variety of studies (e.g. Niu et al., 2019). The difference is only how the final interpolation method for T and P is implemented.

For both climate states a reference surface elevation $h_{0,1}^{\text{ref}}$ must be provided. This is then used to account for elevation changes, similarly to the `elevation_change` atmosphere model of PISM, before combining both climate states. Temperature is assumed to change linearly with elevation change, $\Delta T = -\gamma_T \cdot \Delta h$, where γ_T is the temperature lapse rate.

$$T^* = T + \Delta T = T - \gamma_T \cdot \Delta h \quad (\text{D1})$$

Precipitation is scaled using an exponential factor C for temperature:

$$P^* = P \cdot \exp(C \cdot \Delta T(\Delta h)) = P \cdot \exp(-C \cdot \gamma_T \cdot \Delta h). \quad (\text{D2})$$

Using these equations, the temperature and precipitation fields are scaled to a common reference surface (i.e. sea level), before the actual interpolation is applied. As it applies to both fields, T and P , we write V instead:

$$V^{\text{SL}}(t) = (V_1^{\text{SL}} - V_0^{\text{SL}}) \cdot i(t) + V_0^{\text{SL}}. \quad (\text{D3})$$

After this step, it needs to be checked that the result for precipitation is positive. In a final step, Eqs. (D1) and (D2) are applied again to transfer the results back onto the current surface elevation $h(t)$.

700 It should be noted that the `index` atmosphere model, as it was described above, also accepts seasonal (i.e. time varying, one year periodic) input fields. The climate states can thereby be described by monthly mean values as it is used by the Positive Degree Day (PDD) `surface` model to calculate the surface mass balance of the ice sheet. All command-line options used by this model are listed in Table D1.

D2 Precipitation Cut-off

705 The PISM `precip_cutoff` atmosphere modifier sets precipitation to zero where one of the following conditions apply:

- the surface elevation exceeds a threshold height h_{\max} (and this method is not disabled),
- a user-defined mask has a value of 1.

This method is a simple approach to limit ice sheet growth in regions where other climate forcing methods fail to restrict precipitation. Table D2 shows all command-line options that are used by this PISM modifier.

Table D2. Command-line options for use with the PISM atmosphere modifier `precip_cutoff`.

Command-line option	Comment
<code>-use_precip_cutoff_height</code>	Boolean variable to determine if the threshold height should be used. Default value: <i>false</i> .
<code>-precip_cutoff_height</code>	Defines the threshold height h_{\max} above which precipitation is set to zero. Default value: 3000 m
<code>-precip_cutoff_file</code>	File containing the mask <code>precip_cutoff_mask</code> , which is 1 where precipitation should be eliminated, and 0 elsewhere. If not set, the mask is initialized with zeroes.

710 *Author contributions.* The concept of this study was developed by all authors. SH developed the tool, conducted the experiments and did the analysis. EJG provided LGM palaeo-geography reconstructions used for model initialization. XZ provided LGM and PD climate reconstructions used for climate forcing of the model. The manuscript was written by SH with contributions from all co-authors.

Competing interests. The authors have no competing interests to declare.

715 *Acknowledgements.* SH received funding from the Federal Ministry of Education and Science (BMBF) through the project PalMod. EJG was funded by Helmholtz Exzellenznetzwerk “The Polar System and its Effects on the Ocean Floor (POSY)” and Helmholtz Climate

Initiative REKLIM (Regional Climate Change), a joint research project at the Helmholtz Association of German research centres (HGF). XZ was funded by the National Science Foundation of China (Nr. 42075047) and the Helmholtz Postdoc Program (PD-301). GL received funding from Helmholtz through the program PACES and Changing Earth — Sustaining our Future. Development of PISM is supported by NSF grants PLR-1603799 and PLR-1644277 and NASA grant NNX17AG65G. All computations were done at the AWI computing center. Furthermore, we want to thank Constantine Khroulev from University of Alaska Fairbanks for the helpful advises concerning the implementation into PISM. [Torsten Albrecht, Tijn Berends and an anonymous reviewer helped improve this manuscript with their thoughtful comments.](#)

References

- Albrecht, T., Martin, M., Haseloff, M., Winkelmann, R., and Levermann, A.: Parameterization for Subgrid-Scale Motion of Ice-Shelf Calving
725 Fronts, *The Cryosphere*, 5, 35–44, <https://doi.org/10.5194/tc-5-35-2011>, 2011.
- Aschwanden, A., Bueler, E., Khroulev, C., and Blatter, H.: An Enthalpy Formulation for Glaciers and Ice Sheets, *Journal of Glaciology*, 58,
441–457, <https://doi.org/10.3189/2012JoG11J088>, 2012.
- Augustin, L., Barbante, C., Barnes, P. R. F., Marc Barnola, J., Bigler, M., Castellano, E., Cattani, O., Chappellaz, J., Dahl-Jensen, D.,
730 Delmonte, B., Dreyfus, G., Durand, G., Falourd, S., Fischer, H., Flückiger, J., Hansson, M. E., Huybrechts, P., Jugie, G., Johnsen, S. J.,
Jouzel, J., Kaufmann, P., Kipfstuhl, J., Lambert, F., Lipenkov, V. Y., Littot, G. C., Longinelli, A., Lorrain, R., Maggi, V., Masson-Delmotte,
V., Miller, H., Mulvaney, R., Oerlemans, J., Oerter, H., Orombelli, G., Parrenin, F., Peel, D. A., Petit, J.-R., Raynaud, D., Ritz, C., Ruth,
U., Schwander, J., Siegenthaler, U., Souchez, R., Stauffer, B., Peder Steffensen, J., Stenni, B., Stocker, T. F., Tabacco, I. E., Udisti, R., van
de Wal, R. S. W., van den Broeke, M., Weiss, J., Wilhelms, F., Winther, J.-G., Wolff, E. W., Zucchelli, M., EPICA community members,
and EPICA community members (participants are listed alphabetically): Eight Glacial Cycles from an Antarctic Ice Core, *Nature*, 429,
735 623–628, <https://doi.org/10.1038/nature02599>, 2004.
- Balay, S., Gropp, W. D., McInnes, L. C., and Smith, B. F.: Efficient Management of Parallelism in Object Oriented Numerical Software
Libraries, in: *Modern Software Tools in Scientific Computing*, edited by Arge, E., Bruaset, A. M., and Langtangen, H. P., pp. 163–202,
Birkhäuser Press, 1997.
- Balay, S., Abhyankar, S., Adams, M. F., Brown, J., Brune, P., Buschelman, K., Dalcin, L., Dener, A., Eijkhout, V., Gropp, W. D., Karpeyev,
740 D., Kaushik, D., Knepley, M. G., May, D. A., McInnes, L. C., Mills, R. T., Munson, T., Rupp, K., Sanan, P., Smith, B. F., Zampini, S.,
Zhang, H., and Zhang, H.: PETSc Web Page, <https://www.mcs.anl.gov/petsc>, 2019.
- Beckmann, A. and Goosse, H.: A Parameterization of Ice Shelf–Ocean Interaction for Climate Models, *Ocean Modelling*, 5, 157–170,
[https://doi.org/10.1016/S1463-5003\(02\)00019-7](https://doi.org/10.1016/S1463-5003(02)00019-7), 2003.
- Benn, D. I., Warren, C. R., and Mottram, R. H.: Calving Processes and the Dynamics of Calving Glaciers, *Earth-Science Reviews*, 82,
745 143–179, <https://doi.org/10.1016/j.earscirev.2007.02.002>, 2007.
- Berends, C. J. and van de Wal, R. S. W.: A Computationally Efficient Depression-Filling Algorithm for Digital Elevation Models, Applied
to Proglacial Lake Drainage, *Geosci. Model Dev.*, 9, 4451–4460, <https://doi.org/10.5194/gmd-9-4451-2016>, 2016.
- Björck, S.: A Review of the History of the Baltic Sea, 13.0–8.0 Ka BP, *Quaternary International*, 27, 19–40, [https://doi.org/10.1016/1040-6182\(94\)00057-C](https://doi.org/10.1016/1040-6182(94)00057-C), 1995.
- 750 Broecker, W. S., Kennett, J. P., Flower, B. P., Teller, J. T., Trumbore, S., Bonani, G., and Wolfli, W.: Routing of Meltwater from the Laurentide
Ice Sheet during the Younger Dryas Cold Episode, *Nature*, 341, 318, <https://doi.org/10.1038/341318a0>, 1989.
- Bueler, E. and Brown, J.: Shallow Shelf Approximation as a “Sliding Law” in a Thermomechanically Coupled Ice Sheet Model, *Journal of
Geophysical Research: Earth Surface*, 114, <https://doi.org/10.1029/2008JF001179>, 2009.
- Bueler, E., Lingle, C. S., and Brown, J.: Fast Computation of a Viscoelastic Deformable Earth Model for Ice-Sheet Simulations, *Annals of
755 Glaciology*, 46, 97–105, <https://doi.org/10.3189/172756407782871567>, 2007.
- Calov, R. and Greve, R.: A Semi-Analytical Solution for the Positive Degree-Day Model with Stochastic Temperature Variations, *Journal of
Glaciology*, 51, 173–175, <https://doi.org/10.3189/172756505781829601>, 2005.
- Carlson, A. E., LeGrande, A. N., Oppo, D. W., Came, R. E., Schmidt, G. A., Anslow, F. S., Licciardi, J. M., and Obbink, E. A.: Rapid Early
Holocene Deglaciation of the Laurentide Ice Sheet, *Nature Geoscience*, 1, 620–624, <https://doi.org/10.1038/ngeo285>, 2008.

- 760 Carlson, A. E., Anslow, F. S., Obbink, E. A., LeGrande, A. N., Ullman, D. J., and Licciardi, J. M.: Surface-Melt Driven Laurentide Ice Sheet Retreat during the Early Holocene, *Geophysical Research Letters*, 36, <https://doi.org/10.1029/2009GL040948>, 2009.
- Carrivick, J. L. and Tweed, F. S.: Proglacial Lakes: Character, Behaviour and Geological Importance, *Quaternary Science Reviews*, 78, 34–52, <https://doi.org/10.1016/j.quascirev.2013.07.028>, 2013.
- Carrivick, J. L., Tweed, F. S., Sutherland, J. L., and Mallalieu, J.: Toward Numerical Modeling of Interactions Between Ice-Marginal Proglacial Lakes and Glaciers, *Frontiers in Earth Science*, 8, <https://doi.org/10.3389/feart.2020.577068>, 2020.
- 765 Condon, A. and Winsor, P.: Meltwater Routing and the Younger Dryas, *Proceedings of the National Academy of Sciences*, 109, 19928–19933, <https://doi.org/10.1073/pnas.1207381109>, 2012.
- Cornford, S. L., Martin, D. F., Graves, D. T., Ranken, D. F., Le Brocq, A. M., Gladstone, R. M., Payne, A. J., Ng, E. G., and Lipscomb, W. H.: Adaptive Mesh, Finite Volume Modeling of Marine Ice Sheets, *Journal of Computational Physics*, 232, 529–549, <https://doi.org/10.1016/j.jcp.2012.08.037>, 2013.
- 770 Cuffey, K. and Paterson, W. S. B.: *The Physics of Glaciers*, Butterworth-Heinemann/Elsevier, Burlington, MA, 4th ed edn., 2010.
- Cutler, P. M., Mickelson, D. M., Colgan, P. M., MacAyeal, D. R., and Parizek, B. R.: Influence of the Great Lakes on the Dynamics of the Southern Laurentide Ice Sheet: Numerical Experiments, *Geology*, 29, 1039–1042, [https://doi.org/10.1130/0091-7613\(2001\)029<1039:IOTGLO>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<1039:IOTGLO>2.0.CO;2), 2001.
- 775 Dalton, A. S., Margold, M., Stokes, C. R., Tarasov, L., Dyke, A. S., Adams, R. S., Allard, S., Arends, H. E., Atkinson, N., Attig, J. W., Barnett, P. J., Barnett, R. L., Batterson, M., Bernatchez, P., Borns, H. W., Breckenridge, A., Briner, J. P., Brouard, E., Campbell, J. E., Carlson, A. E., Clague, J. J., Curry, B. B., Daigneault, R.-A., Dubé-Loubert, H., Easterbrook, D. J., Franzi, D. A., Friedrich, H. G., Funder, S., Gauthier, M. S., Gowan, A. S., Harris, K. L., Hétu, B., Hooyer, T. S., Jennings, C. E., Johnson, M. D., Kehew, A. E., Kelley, S. E., Kerr, D., King, E. L., Kjeldsen, K. K., Knaeble, A. R., Lajeunesse, P., Lakeman, T. R., Lamothe, M., Larson, P., Lavoie, M., Loope, H. M., Lowell, T. V., Lusardi, B. A., Manz, L., McMartin, I., Nixon, F. C., Occhietti, S., Parkhill, M. A., Piper, D. J., Pronk, A. G., Richard, P. J., Ridge, J. C., Ross, M., Roy, M., Seaman, A., Shaw, J., Stea, R. R., Teller, J. T., Thompson, W. B., Thorleifson, L. H., Utting, D. J., Veillette, J. J., Ward, B. C., Weddle, T. K., and Wright, H. E.: An Updated Radiocarbon-Based Ice Margin Chronology for the Last Deglaciation of the North American Ice Sheet Complex, *Quaternary Science Reviews*, 234, 106 223, <https://doi.org/10.1016/j.quascirev.2020.106223>, 2020.
- 780 785 Davies, J. H.: Global Map of Solid Earth Surface Heat Flow, *Geochemistry, Geophysics, Geosystems*, 14, 4608–4622, <https://doi.org/10.1002/ggge.20271>, 2013.
- Dyke, A. S.: An Outline of North American Deglaciation with Emphasis on Central and Northern Canada, in: *Developments in Quaternary Sciences*, vol. 2, pp. 373–424, Elsevier, [https://doi.org/10.1016/S1571-0866\(04\)80209-4](https://doi.org/10.1016/S1571-0866(04)80209-4), 2004.
- Feldmann, J., Albrecht, T., Khroulev, C., Pattyn, F., and Levermann, A.: Resolution-Dependent Performance of Grounding Line Motion in a Shallow Model Compared with a Full-Stokes Model According to the MIS2 Intercomparison, *Journal of Glaciology*, 60, 353–360, <https://doi.org/10.3189/2014JoG13J093>, 2014.
- 790 Fowler, A. C., Rickaby, R. E. M., and Wolff, E. W.: Exploration of a Simple Model for Ice Ages, *GEM - International Journal on Geomathematics*, 4, 227–297, <https://doi.org/10.1007/s13137-012-0040-7>, 2013.
- Funk, M. and Röthlisberger, H.: Forecasting the Effects of a Planned Reservoir Which Will Partially Flood the Tongue of Unteraargletscher in Switzerland, *Annals of Glaciology*, 13, 76–81, <https://doi.org/10.3189/S0260305500007679>, 1989.
- 795 Golledge, N. R., Kowalewski, D. E., Naish, T. R., Levy, R. H., Fogwill, C. J., and Gasson, E. G. W.: The Multi-Millennial Antarctic Commitment to Future Sea-Level Rise, *Nature*, 526, 421–425, <https://doi.org/10.1038/nature15706>, 2015.

- Gowan, E. J., Tregoning, P., Purcell, A., Montillet, J.-P., and McClusky, S.: A Model of the Western Laurentide Ice Sheet, Using Observations of Glacial Isostatic Adjustment, *Quaternary Science Reviews*, 139, 1–16, <https://doi.org/10.1016/j.quascirev.2016.03.003>, 2016.
- 800 Gregoire, L. J., Payne, A. J., and Valdes, P. J.: Deglacial Rapid Sea Level Rises Caused by Ice-Sheet Saddle Collapses, *Nature*, 487, 219–222, <https://doi.org/10.1038/nature11257>, 2012.
- Haresign, E. C.: Glacio-Limnological Interactions at Lake-Calving Glaciers, Thesis, University of St Andrews, 2004.
- Hickin, A. S., Lian, O. B., Levson, V. M., and Cui, Y.: Pattern and Chronology of Glacial Lake Peace Shorelines and Implications for Isostasy and Ice-Sheet Configuration in Northeastern British Columbia, Canada, *Boreas*, 44, 288–304, <https://doi.org/10.1111/bor.12110>, 2015.
- 805 Hinck, S. and PISM Authors: PISM-LakeCC, Zenodo, <https://doi.org/10.5281/ZENODO.4304671>, 2020.
- Hinck, S., Gowan, E. J., and Lohmann, G.: LakeCC: A Tool for Efficiently Identifying Lake Basins with Application to Palaeogeographic Reconstructions of North America, *Journal of Quaternary Science*, 35, 422–432, <https://doi.org/10.1002/jqs.3182>, 2020.
- Hooyer, T. S. and Iverson, N. R.: Flow Mechanism of the Des Moines Lobe of the Laurentide Ice Sheet, *Journal of Glaciology*, 48, 575–586, <https://doi.org/10.3189/172756502781831160>, 2002.
- 810 Hossain, A., Zhang, X., and Lohmann, G.: A Model-Data Comparison of the Last Glacial Maximum Surface Temperature Changes, *Climate of the Past Discussions*, pp. 1–18, <https://doi.org/10.5194/cp-2018-9>, 2018.
- Krinner, G., Mangerud, J., Jakobsson, M., Crucifix, M., Ritz, C., and Svendsen, J. I.: Enhanced Ice Sheet Growth in Eurasia Owing to Adjacent Ice-Dammed Lakes, *Nature*, 427, 429–432, <https://doi.org/10.1038/nature02233>, 2004.
- Lambeck, K., Rouby, H., Purcell, A., Sun, Y., and Sambridge, M.: Sea Level and Global Ice Volumes from the Last Glacial Maximum to the Holocene, *Proceedings of the National Academy of Sciences*, 111, 15 296–15 303, <https://doi.org/10.1073/pnas.1411762111>, 2014.
- 815 Leverington, D. W., Mann, J. D., and Teller, J. T.: Changes in the Bathymetry and Volume of Glacial Lake Agassiz between 9200 and 7700 14C Yr B.P., *Quaternary Research*, 57, 244–252, <https://doi.org/10.1006/qres.2001.2311>, 2002.
- Levermann, A., Albrecht, T., Winkelmann, R., Martin, M. A., Haseloff, M., and Joughin, I.: Kinematic First-Order Calving Law Implies Potential for Abrupt Ice-Shelf Retreat, *The Cryosphere*, 6, 273–286, <https://doi.org/10.5194/tc-6-273-2012>, 2012.
- 820 Lingle, C. S. and Clark, J. A.: A Numerical Model of Interactions between a Marine Ice Sheet and the Solid Earth: Application to a West Antarctic Ice Stream, *Journal of Geophysical Research: Oceans*, 90, 1100–1114, <https://doi.org/10.1029/JC090iC01p01100>, 1985.
- Lliboutry, L. and Duval, P.: Various Isotropic and Anisotropic Ices Found in Glaciers and Polar Ice Caps and Their Corresponding Rheologies: *Ann Geophys V3, N2, March–April 1985, P207–224*, *International Journal of Rock Mechanics and Mining Sciences & Geomechanics Abstracts*, 22, 198, [https://doi.org/10.1016/0148-9062\(85\)90267-0](https://doi.org/10.1016/0148-9062(85)90267-0), 1985.
- 825 Lochte, A. A., Repschläger, J., Kienast, M., Garbe-Schönberg, D., Andersen, N., Hamann, C., and Schneider, R.: Labrador Sea Freshening at 8.5 Ka BP Caused by Hudson Bay Ice Saddle Collapse, *Nature Communications*, 10, 586, <https://doi.org/10.1038/s41467-019-08408-6>, 2019.
- Mallalieu, J., Carrivick, J. L., Quincey, D. J., and Smith, M. W.: Calving Seasonality Associated With Melt-Undercutting and Lake Ice Cover, *Geophysical Research Letters*, 47, <https://doi.org/10.1029/2019GL086561>, 2020.
- 830 Margold, M., Stokes, C. R., and Clark, C. D.: Ice Streams in the Laurentide Ice Sheet: Identification, Characteristics and Comparison to Modern Ice Sheets, *Earth-Science Reviews*, 143, 117–146, <https://doi.org/10.1016/j.earscirev.2015.01.011>, 2015.
- Margold, M., Stokes, C. R., and Clark, C. D.: Reconciling Records of Ice Streaming and Ice Margin Retreat to Produce a Palaeogeographic Reconstruction of the Deglaciation of the Laurentide Ice Sheet, *Quaternary Science Reviews*, 189, 1–30, <https://doi.org/10.1016/j.quascirev.2018.03.013>, 2018.

- 835 Martin, M. A., Winkelmann, R., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., and Levermann, A.: The Potsdam Parallel Ice Sheet Model (PISM-PIK) – Part 2: Dynamic Equilibrium Simulation of the Antarctic Ice Sheet, *The Cryosphere*, 5, 727–740, <https://doi.org/10.5194/tc-5-727-2011>, 2011.
- Matero, I. S. O., Gregoire, L. J., Ivanovic, R. F., Tindall, J. C., and Haywood, A. M.: The 8.2 Ka Cooling Event Caused by Laurentide Ice Saddle Collapse, *Earth and Planetary Science Letters*, 473, 205–214, <https://doi.org/10.1016/j.epsl.2017.06.011>, 2017.
- 840 Matero, I. S. O., Gregoire, L. J., and Ivanovic, R. F.: Simulating the Early Holocene Demise of the Laurentide Ice Sheet with BISICLES (Public Trunk Revision 3298), *Geoscientific Model Development*, 13, 4555–4577, <https://doi.org/10.5194/gmd-13-4555-2020>, 2020.
- Mathews, W. H.: Retreat of the Last Ice Sheets in Northeastern British Columbia and Adjacent Alberta, Tech. Rep. 331, <https://doi.org/10.4095/102160>, 1980.
- Niu, L., Lohmann, G., Hinck, S., Gowan, E. J., and Krebs-Kanzow, U.: The Sensitivity of Northern Hemisphere Ice Sheets to Atmospheric Forcing during the Last Glacial Cycle Using PMIP3 Models, *Journal of Glaciology*, pp. 1–17, <https://doi.org/10.1017/jog.2019.42>, 2019.
- 845 North Greenland Ice Core Project members: High-Resolution Record of Northern Hemisphere Climate Extending into the Last Interglacial Period, *Nature*, 431, 147–151, <https://doi.org/10.1038/nature02805>, 2004.
- North Greenland Ice Core Project Members: 50 Year Means of Oxygen Isotope Data from Ice Core NGRIP, Supplement to: North Greenland Ice Core Project Members (2004): High-Resolution Record of Northern Hemisphere Climate Extending into the Last Interglacial Period.
- 850 *Nature*, 431, 147-151, <https://doi.org/10.1594/PANGAEA.586886>, 2007.
- Peltier, W. R., Vettoretti, G., and Stastna, M.: Atlantic Meridional Overturning and Climate Response to Arctic Ocean Freshening, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2005GL025251>, 2006.
- Peltier, W. R., Argus, D. F., and Drummond, R.: Space Geodesy Constrains Ice Age Terminal Deglaciation: The Global ICE-6G_C (VM5a) Model, *Journal of Geophysical Research: Solid Earth*, 120, 450–487, <https://doi.org/10.1002/2014JB011176>, 2015.
- 855 Peyaud, V., Ritz, C., and Krinner, G.: Modelling the Early Weichselian Eurasian Ice Sheets: Role of Ice Shelves and Influence of Ice-Dammed Lakes, *Climate of the Past*, 3, 375–386, <https://doi.org/10.5194/cp-3-375-2007>, 2007.
- Quiquet, A., Dumas, C., Ritz, C., Peyaud, V., and Roche, D. M.: The GRISLI Ice Sheet Model (Version 2.0): Calibration and Validation for Multi-Millennial Changes of the Antarctic Ice Sheet, *Geoscientific Model Development*, 11, 5003–5025, <https://doi.org/10.5194/gmd-11-5003-2018>, 2018.
- 860 Quiquet, A., Dumas, C., Paillard, D., Ramstein, G., Ritz, C., and Roche, D. M.: Deglacial Ice Sheet Instabilities Induced by Proglacial Lakes, *Geophysical Research Letters*, 48, e2020GL092141, <https://doi.org/10.1029/2020GL092141>, 2021.
- Schaffer, J., Timmermann, R., Arndt, J. E., Kristensen, S. S., Mayer, C., Morlighem, M., and Steinhage, D.: A Global, High-Resolution Data Set of Ice Sheet Topography, Cavity Geometry, and Ocean Bathymetry, *Earth System Science Data*, 8, 543–557, <https://doi.org/10.5194/essd-8-543-2016>, 2016.
- 865 Schoof, C.: Ice Sheet Grounding Line Dynamics: Steady States, Stability, and Hysteresis, *Journal of Geophysical Research*, 112, <https://doi.org/10.1029/2006JF000664>, 2007.
- Skvarca, P., Angelis, H. D., Naruse, R., Warren, C. R., and Aniya, M.: Calving Rates in Fresh Water: New Data from Southern Patagonia, *Annals of Glaciology*, 34, 379–384, <https://doi.org/10.3189/172756402781817806>, 2002.
- Smith, D. G.: Glacial Lake McConnell: Paleogeography, Age, Duration, and Associated River Deltas, Mackenzie River Basin, Western
- 870 Canada, *Quaternary Science Reviews*, 13, 829–843, [https://doi.org/10.1016/0277-3791\(94\)90004-3](https://doi.org/10.1016/0277-3791(94)90004-3), 1994.

- Stokes, C. R. and Clark, C. D.: The Dubawnt Lake Palaeo-Ice Stream: Evidence for Dynamic Ice Sheet Behaviour on the Canadian Shield and Insights Regarding the Controls on Ice-Stream Location and Vigour, *Boreas*, 32, 263–279, <https://doi.org/10.1111/j.1502-3885.2003.tb01442.x>, 2003.
- 875 Sutherland, J. L., Carrivick, J. L., Gandy, N., Shulmeister, J., Quincey, D. J., and Cornford, S. L.: Proglacial Lakes Control Glacier Geometry and Behavior During Recession, *Geophysical Research Letters*, 47, e2020GL088 865, <https://doi.org/10.1029/2020GL088865>, 2020.
- Tarasov, L., Dyke, A. S., Neal, R. M., and Peltier, W. R.: A Data-Calibrated Distribution of Deglacial Chronologies for the North American Ice Complex from Glaciological Modeling, *Earth and Planetary Science Letters*, 315-316, 30–40, <https://doi.org/10.1016/j.epsl.2011.09.010>, 2012.
- 880 Teller, J. T. and Leverington, D. W.: Glacial Lake Agassiz: A 5000 Yr History of Change and Its Relationship to the $\Delta 18\text{ O}$ Record of Greenland, *Geological Society of America Bulletin*, 116, 729–742, <https://doi.org/10.1130/B25316.1>, 2004.
- Teller, J. T., Leverington, D. W., and Mann, J. D.: Freshwater Outbursts to the Oceans from Glacial Lake Agassiz and Their Role in Climate Change during the Last Deglaciation, *Quaternary Science Reviews*, 21, 879–887, [https://doi.org/10.1016/S0277-3791\(01\)00145-7](https://doi.org/10.1016/S0277-3791(01)00145-7), 2002.
- the PISM authors: PISM, a Parallel Ice Sheet Model, <http://www.pism-docs.org/>, 2015.
- 885 Thomas, R. H. and Bentley, C. R.: A Model for Holocene Retreat of the West Antarctic Ice Sheet, *Quaternary Research*, 10, 150–170, [https://doi.org/10.1016/0033-5894\(78\)90098-4](https://doi.org/10.1016/0033-5894(78)90098-4), 1978.
- Tsutaki, S., Fujita, K., Nuimura, T., Sakai, A., Sugiyama, S., Komori, J., and Tshering, P.: Contrasting Thinning Patterns between Lake- and Land-Terminating Glaciers in the Bhutanese Himalaya, *The Cryosphere*, 13, 2733–2750, <https://doi.org/10.5194/tc-13-2733-2019>, 2019.
- Tulaczyk, S., Kamb, W. B., and Engelhardt, H. F.: Basal Mechanics of Ice Stream B, West Antarctica: 2. Undrained Plastic Bed Model, *Journal of Geophysical Research: Solid Earth*, 105, 483–494, <https://doi.org/10.1029/1999JB900328>, 2000.
- 890 Warren, C. R. and Kirkbride, M. P.: Calving Speed and Climatic Sensitivity of New Zealand Lake-Calving Glaciers, *Annals of Glaciology*, 36, 173–178, <https://doi.org/10.3189/172756403781816446>, 2003.
- Warren, C. R., Greene, D. R., and Glasser, N. F.: Glaciar Upsala, Patagonia: Rapid Calving Retreat in Fresh Water, *Annals of Glaciology*, 21, 311–316, <https://doi.org/10.3198/1995AoG21-1-311-316>, 1995.
- Weertman, J.: Stability of the Junction of an Ice Sheet and an Ice Shelf, *Journal of Glaciology*, 13, 3–11, <https://doi.org/10.3189/S0022143000023327>, 1974.
- 895 Winkelmann, R., Martin, M. A., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., and Levermann, A.: The Potsdam Parallel Ice Sheet Model (PISM-PIK) – Part 1: Model Description, *The Cryosphere*, 5, 715–726, <https://doi.org/10.5194/tc-5-715-2011>, 2011.
- Wu, P.: Sensitivity of Relative Sea Levels and Crustal Velocities in Laurentide to Radial and Lateral Viscosity Variations in the Mantle, *Geophysical Journal International*, 165, 401–413, <https://doi.org/10.1111/j.1365-246X.2006.02960.x>, 2006.
- 900 Zhang, X., Lohmann, G., Knorr, G., and Xu, X.: Different Ocean States and Transient Characteristics in Last Glacial Maximum Simulations and Implications for Deglaciation, *Climate of the Past*, 9, 2319–2333, <https://doi.org/10.5194/cp-9-2319-2013>, 2013.