

Brief communication: Increased glacier mass loss in the Russian High Arctic (2010-2017)

Christian Sommer¹, Thorsten Seehaus¹, Andrey Glazovsky², Matthias H. Braun¹

¹Institut für Geographie, Friedrich-Alexander-Universität Erlangen-Nürnberg, Erlangen, 91058, Germany

5 ²Institute of Geography RAS, Moscow, 119017, Russia

Correspondence to: Christian Sommer (chris.sommer@fau.de)

Abstract. Glaciers in the Russian High Arctic have been subject to extensive atmospheric warming due to global climate change, yet their contribution to sea level rise has been relatively small over the past decades. Here we show surface elevation change measurements and geodetic mass balances of 93% of all glacierized areas of Novaya Zemlya, Severnaya Zemlya and Franz Josef Land using interferometric synthetic aperture radar measurements taken between 2010 and 2017. We calculate an overall mass loss rate of $-23 \pm 6 \text{ Gt a}^{-1}$, corresponding to a sea level rise contribution of $0.06 \pm 0.02 \text{ mm a}^{-1}$. Compared to measurements prior to 2010, mass loss of glaciers on the Russian archipelagos has doubled in recent years.

1 Introduction

The Arctic has undergone large environmental changes due to polar climate change (Box et al., 2019). An increase in glacier mass loss has been observed in many polar regions (Morris et al., 2020; Zheng et al., 2018; Ciraci et al., 2020). The Russian High Arctic, including the archipelagos Novaya Zemlya, Severnaya Zemlya and Franz Josef Land, is one of these regions. Despite a glacierized area of $\sim 52,000 \text{ km}^2$, in-situ observations of glacier mass change are sparse. Previous regionwide assessments were mostly limited to the early 21st century and based on gravimetry (Gardner et al., 2013; Jacob et al., 2012; Matsuo and Heki, 2013; Wouters et al., 2019) and altimetry (Ciraci et al., 2018; Moholdt et al., 2012). Most of these studies show mass change rates ranging from -5 to -10 Gt a^{-1} . However, both methods have limitations: altimetry requires interpolation while uncertainties of the gravimetric approach might arise from the scattered ice caps and various corrections related to surrounding oceans, surface hydrology and glacial isostatic adjustment (GIA). In this study we have measured surface elevation changes of most Russian Arctic glaciers from digital elevation models (DEM) to derive geodetic mass changes between 2010 and 2017. We use synthetic aperture radar (SAR) DEMs of the TanDEM-X satellites which are independent from cloud cover. Additionally, we apply a correction for differences in SAR penetration depth due to temporal offsets between acquisitions based on backscatter intensity.

2 Data & Methods

2.1 Glacier elevation change

Glacier elevation change rates are calculated by comparing DEMs of the TerraSAR-X add-on for Digital Elevation Measurement mission (TanDEM-X), operated by the German Aerospace Center and Astrium Defence and Space. For the Russian Arctic, TanDEM-X acquisitions of winter 2010/11 (Dec-Feb, Apr) and autumn/winter 2016/17 (Sep-Feb) are available. Elevation models are derived from TanDEM-X Co-registered Single look Slant range Complex (CoSSC) data, closely following the workflow of Braun et al. (2019) and Seehaus et al. (2019). A detailed description of the interferometric DEM generation, co-registration and uncertainty assessment is provided in the supplement. Eventually, the co-registered TanDEM-X DEMs are merged to create two elevation mosaics of winter 2010/11 and 2016/17 and differenced to derive glacier elevation and mass change rates based on glacier areas of the Randolph Glacier Inventory (Pfeffer et al., 2014).

2.2 X-band surface penetration

The interferometric TanDEM-X DEMs provide a high spatial resolution and almost complete coverage of the Russian Arctic archipelagos but can suffer from SAR signal penetration into the glacier surface. The depth of signal penetration is related to the prevailing glacier surface conditions at the time of the SAR acquisition. In general, SAR penetration is close to zero for melting snow surfaces and bare glacier ice and increases during dry and frozen conditions. X-band penetration depths of several meters have been observed in different regions (e.g. Millan et al., 2015; Zhao and Floricioiu, 2017; Abdullahi et al., 2018; Li et al., 2021). When comparing elevations of different TanDEM-X scenes the relative difference in penetration depths is determined by the acquisition dates and seasons. The TanDEM-X data used at most glacierized areas of the Russian Arctic archipelagos were acquired during winter 2010/11 (94% of total glacier area) and winter 2016/17 (83%) at temperatures well below zero degrees Celsius and frozen ice surfaces. It is likely that for those acquisitions the difference in X-band penetration depth is small as the SAR data was acquired in the same season and the presence of surface melt or liquid water is very unlikely in the Arctic winter months. However, for some glacier areas of Novaya Zemlya (35%) and Franz Josef Land (6%), SAR data from September 2016 had to be included to calculate elevation changes because there were no respective winter scenes available. Using those DEMs without further correction can bias the measured surface elevation change as seasonal changes in snow and ice of the glacier surface have significant impacts on the SAR penetration depth (Abdullahi et al., 2019).

2.3 SAR backscatter intensity analysis

The SAR backscatter intensity depends on physical properties of the glacier ice, such as grain size and density, roughness and water content (Wessel et al., 2016), and has been used to estimate penetration depths by a number of studies (e.g. Abdullahi et al., 2018, 2019; Li et al., 2021). Fig. 1a shows the hypsometric backscatter distribution of TanDEM-X acquisitions between September 2016 and January 2017 on Novaya Zemlya. While average backscatter intensity of the October-January acquisitions shows similar patterns, a clear difference is observed for the September DEMs. At altitudes above ~400 m a.s.l., the September

60 data show much higher backscatter values than the respective winter scenes, indicating different surface conditions at the acquisition times. To estimate the difference in penetration depth between the September 2016 and winter 2016/17 DEMs, we derive the measured elevation difference and respective backscatter intensity from overlapping glacier areas which were covered by the September as well as winter acquisitions. Those reference areas cover a total glacier area of $\sim 2,500 \text{ km}^2$ and are equally distributed across the Novaya Zemlya ice cap (Fig. S2a). Using the local incidence angle and surface slope, the measured absolute vertical offsets between September and winter 2016/17 are converted to surface penetration depths according to EQ. 1.

65

$$d_p = \Delta h_{W-A} \times \frac{\cos \alpha}{\cos \varphi} \quad \text{EQ. 1}$$

Where d_p is the depth of surface penetration, Δh_{W-A} the height difference between Winter and Autumn (September) acquisition, α the glacier surface slope and φ the local incidence angle. Thereafter, d_p is aggregated and compared to altitude and backscatter intensity of the September acquisitions, respectively. As shown in Fig. 1c and Fig. 1d, the offset between the September and winter 2016/17 backscatter intensity increases above elevations of $\sim 400 \text{ m a.s.l.}$ while the difference in estimated signal penetration depth increases with higher backscatter intensities of the September acquisitions. Based on this relationship between penetration difference, backscatter intensity and altitude, a linear regression model can be created to estimate the penetration bias d_p (EQ. 2):

75

$$d_p = \beta_0 + \beta_1 \times Int \quad \text{EQ. 2}$$

Int is the backscatter intensity in decibel and β_0 and β_1 the regression coefficients. To fit d_p and Int , the difference between the autumn and winter TanDEM-X acquisitions and backscatter intensity of all overlapping DEM pixels is used. To predict the bias in surface penetration depth between the autumn and winter acquisitions 2016/17, the model is then applied to all glacier areas above 400 m a.s.l. on Novaya Zemlya which were only covered by September 2016 SAR ($\sim 7,800 \text{ km}^2$, Fig. S2a). Eventually, the estimated surface penetration depths are converted back to vertical differences in elevation by rearranging EQ. 1. The predicted vertical correction values (Fig. S2b) are then added to the September 2016 elevations and the adjusted elevation change rate is calculated.

85 For Franz Josef Land and Severnaya Zemlya, the elevation change rate is not adjusted, as on both archipelagos the temporal offset between DEM acquisitions is much smaller as on Novaya Zemlya. Average backscatter intensity is relatively homogeneous for all 2016/17 acquisitions on Severnaya Zemlya (Fig. S1c) while on Franz Josef Land only a very small fraction of September TanDEM-X acquisitions (6%) shows significant differences in backscatter intensity (Fig. S1e). Therefore, transferring the empirical relationship between differences in surface penetration depth and autumn backscatter intensities on Novaya Zemlya to those archipelagos would rather increase the uncertainty of the elevation change measurement.

90

3. Results

Glacier surface elevation changes of the Russian Arctic Archipelagos are shown in Fig. 2. High thinning rates are measured at elevations below 600 m a.s.l., while surface change rates in the upper accumulation areas are close to zero or slightly positive. Average elevation change rates are highest on Novaya Zemlya (signal-penetration corrected: $-0.68 \pm 0.46 \text{ m a}^{-1}$), mostly due to strong surface thinning close to the termini of the large outlet and tidewater glaciers (Fig. S3c). Regional elevation changes of glaciers in Franz Josef Land ($-0.48 \pm 0.04 \text{ m a}^{-1}$) and Severnaya Zemlya ($-0.34 \pm 0.12 \text{ m a}^{-1}$) are in general less negative and confined to a smaller number of glaciers. Average elevation changes on Severnaya Zemlya are strongly positive below 50 m a.s.l. (Fig. S3b) due to a surge event within the observation period at the Vavilov Ice Cap (Zheng et al., 2019). Slight thickening is also observed at the highest glacierized altitudes and the Academy of Sciences Ice Cap (Severnaya Zemlya), similar to the observations of (Sánchez-Gómez et al., 2019). The overall adjusted mass change of the Russian Arctic is $-23.06 \pm 6.41 \text{ Gt a}^{-1}$ (density conversion factor: 850 kg m^{-3}). Approximately 50% of the total mass loss are caused by glaciers on Novaya Zemlya, while mass changes of Severnaya Zemlya and Franz Josef Land account for about a quarter each. Table 1 summarises the measured and adjusted change rates for the Russian Arctic.

For the DEM acquisitions of 2016/17 on Novaya Zemlya, a distinct difference in backscatter intensity is visible between SAR data acquired in September 2016 and October-January 2016/17 (Fig. 1a). The average vertical difference in surface elevation on the respective overlapping glacier areas (Fig. S2a) is 2.13 m while the elevation change rates derived from all glacier areas which were acquired in September 2016 (Fig. 1b), show an average difference of 0.4 m a^{-1} in surface lowering compared to the winter 2016/17 DEMs. Furthermore, the elevation change rate of the period winter 2010/11 to winter 2016/17 is consistently more negative at all altitudes while the change rate between winter 2010/11 and September 2016 indicates elevation gains at the highest glacierized altitudes. The analysed vertical elevation differences of the overlapping glacier areas (Fig. S2a) and the respective backscatter intensity of the September datasets (Fig. 1a) indicate altitudinal differences in signal penetration depth between the autumn and winter SAR data.

When transferred to all areas on Novaya Zemlya, the glacier surface acquired by TanDEM-X in September 2016 was approximately 2.3 m higher than the surface elevations measured during the winter months 2016/17. The measured glacier mass change rate of Novaya Zemlya is therefore ~20% lower than the adjusted mass change because the elevation changes derived from DEM acquisitions of September 2016 is consistently less negative than those from the winter months.

4. Discussion

Differences in the SAR derived elevation change rates (Fig. 1b) can be related either to surface penetration of the X-Band radar or physical changes of the surface height due to accumulation or ablation of snow and ice. The TanDEM-X DEM difference on Novaya Zemlya does not fully cover the accumulation period of the last year of the observation period as the acquisitions of September 2016 do not or only partially capture the amount of winter accumulation from October to December. This potential bias in measured winter accumulation would lead to an overestimation of surface elevation loss between winter

2010/11 and September 2016. However, the analysis of surface elevation changes derived from autumn and winter DEMs indicate, that the surface measured by TanDEM-X in winter 2016/17 was below the surface heights acquired in September 2016. As the occurrence of major surface melt within the Arctic winter months is unlikely, the observed elevation offset is most likely related to differences in the relative depth of signal penetration of the X-band SAR. The analysis of backscatter intensities of different acquisition months (Fig. 1a) indicates a change in glacier surface properties between the acquisitions from September 2016 and winter 2016/17. The observed differences in backscatter could be related either to the presence of melt or fresh snow at the glacier surface which would decrease the depth of signal penetration as the amount of penetration depends on the condition of the glacier surface and is close to zero for melting snow surfaces and bare glacier ice. The majority of SAR data of the 2016/17 timestep was acquired at months with temperatures well below 0°C while average temperatures on Novaya Zemlya in September 2016 were close to the melting point (Fig. S1f). Thus, the differences in backscatter intensity could be caused by either accumulation of fresh snow at the glacier surface or days with snow melt during September 2016. It is likely that the depth of signal penetration in the winter seasons 2010/11 and 2016/17 was relatively large but similar due to comparable dry and frozen surface conditions. A similar observation was reported for Antarctic Peninsula glaciers where the measured TanDEM-X cold-season heights rather referred to the refrozen firn of the previous summer than to the actual glacier surface (Rott et al., 2014). During the September 2016 acquisitions on Novaya Zemlya, the absolute depth of signal penetration was probably smaller and the measured surface closer to the glacier surface. However, due to the change in surface conditions, the relative difference between penetration depths of the winter season 2010/11 and autumn 2016 increased. The adjusted glacier change rate of Novaya Zemlya is therefore more negative than the measured rate because the effects of different signal penetration depths probably outweigh the winter accumulation.

Over the last decades, the High Arctic has been subject to ongoing warming (Jansen et al., 2020) and glacier mass budgets have become more negative. Compared to previous studies (Fig. S5), glacier mass loss has increased in the Russian High Arctic since 2010. The glacier mass changes measured by TanDEM-X are similar or more negative than recent gravimetric records (Ciraci et al., 2020; Wouters et al., 2019), supporting their observation of increasing mass loss. A recent global study based on optical elevation models (Hugonnet et al., 2021) reported a less negative mass change ($-10.4 \pm 1.9 \text{ Gt a}^{-1}$) since 2000, yet their measurements also indicate a distinct acceleration in mass loss over the course of the 21st century.

While the regional geodetic mass change derived from TanDEM-X data of Franz Josef Land is very similar to recent gravimetric (Ciraci et al., 2020) and altimetric (Zheng et al., 2018) measurements, the estimate for Severnaya Zemlya is even more negative which might indicate recent acceleration of glacier mass loss also on this archipelago. However, the highest mass changes are mostly confined to a small number of outlet glaciers of the Vavilov and Academy of Sciences Ice Caps (RGI60-09.00915,919,920,971). For the remaining glacierized areas of the Severnaya Zemlya Archipelago, the average elevation change rate is much smaller (-0.20 m a^{-1}). The strongest local surface lowering is observed at the large outlet glaciers, most notably on Novaya Zemlya. For those glaciers, an increasing retreat in the early 21st century was attributed to fjord geometries and changes in sea-ice concentrations (Carr et al., 2014). Long-term observations also indicate a more rapid thinning during recent years, particularly at the termini of marine-terminating glaciers (Melkonian et al., 2016). Using a combination of

gravimetric and altimetric measurements, (Ciraci et al., 2018) reported a similar mass change of $-14 \pm 4 \text{ Gt a}^{-1}$ for Novaya Zemlya (2010-2016) to the signal penetration adjusted mass change rate derived by TanDEM-X. An acceleration in flow velocities for the major tidewater glaciers in the Russian Arctic was also measured by (Strozzi et al., 2017) over the course of the last decades. In contrast to the lower ablation areas, elevation gains of up to 0.4 m a^{-1} are measured for the highest altitudes of the Russian Arctic archipelagos, which does not seem to be related to potential SAR penetration because the respective measurements were acquired under similar surface conditions. This is particularly noticeable at some parts of the large accumulation areas of Novaya Zemlya, Severnaya Zemlya and Graham Bell Island (Franz Josef Land). Similar patterns can be also observed in the elevation change maps of altimetry measurements (Ciraci et al., 2018; Moholdt et al., 2012; Sánchez-Gómez et al., 2019; Zheng et al., 2018) and might be related to increased moisture transport and accumulation (Box et al., 2019). The ERA5 datasets also indicate a positive trend in temperature and total column water vapor (Fig. S4a & b) for the Russian Arctic archipelagos. However, the latter trend is not statistically significant in most regions and less pronounced than the increase in temperature, supporting our observations of an overall amplification of glacier mass loss.

5. Conclusion

Glaciers in the Russian High Arctic have shown increasingly negative glacier mass change during the 21st century and contributed 0.06 mm a^{-1} to global sea-level rise between 2010 and 2017. This observation is in line with glacier changes of other Arctic regions, showing an increasing contribution to sea-level rise in recent years compared to glacierized areas outside of the polar regions.

The acquisition date related differences in elevation change on Novaya Zemlya highlight the relevance of similar surface conditions between SAR acquisitions when using DEM-differencing. Particularly for shorter observation periods, corrections for temporal offsets between acquisitions are crucial as the measured elevation change rate can be biased by changes in surface conditions. However, acquisitions from the same season should be used whenever possible, as the measurement uncertainty increases depending on the corrected glacier area. Regarding upcoming TanDEM-X acquisitions, combined measurements with the new ICESat-2 laser altimeter have the potential to much better constrain offsets between different acquisition dates.

190 *Data availability.* Elevation change maps and raster masks with the specific observation period of each cell are provided via the World Data Center PANGAEA (<https://www.pangaea.de/>) at #link#.

Author contributions. C.S. processed the glacier elevation and mass change data, created the graphs and wrote the manuscript. The analysis code of the DEM creation and coregistration was jointly developed by C.S. and T.C.S. A.G. contributed to the
195 comparison and interpretation of measured glacier changes. M.H.B. initiated and led the study. All authors revised the paper.

Competing interests. The authors declare that they have no conflict of interest.

Acknowledgements. This study was financially supported by the grant BR2105/14-2 within the DFG Priority Program “Re-
200 gional Sea Level Change and Society”. We thank the Copernicus Climate Change Service (C3S) which is implemented by the European Centre for Medium-range Weather Forecasts (ECMWF) on behalf of the European Commission for free and open data access. TanDEM-X data was kindly provided free of charge by the German Aerospace Center (DLR) under AO mabra_XTI_GLAC0264.

205

210

215

220


References

- Abdullahi, S., Wessel, B., Leichtle, T., Huber, M., Wohlfart, C., and Roth, A.: Investigation of Tandem-x Penetration Depth Over the Greenland Ice Sheet, in: IGARSS 2018 - 2018 IEEE International Geoscience and Remote Sensing Symposium, IGARSS 2018 - 2018 IEEE International Geoscience and Remote Sensing Symposium, Valencia, 1336–1339, 225 <https://doi.org/10.1109/IGARSS.2018.8518930>, 2018.
- Abdullahi, S., Wessel, B., Huber, M., Wendleder, A., Roth, A., and Kuenzer, C.: Estimating Penetration-Related X-Band InSAR Elevation Bias: A Study over the Greenland Ice Sheet, 19, 2019.
- Box, J. E., Colgan, W. T., Christensen, T. R., Schmidt, N. M., Lund, M., Parmentier, F.-J. W., Brown, R., Bhatt, U. S., Euskirchen, E. S., Romanovsky, V. E., Walsh, J. E., Overland, J. E., Wang, M., Corell, R. W., Meier, W. N., Wouters, B., Mernild, S., Mård, J., Pawlak, J., and Olsen, M. S.: Key indicators of Arctic climate change: 1971–2017, *Environ. Res. Lett.*, 14, 045010, 230 <https://doi.org/10.1088/1748-9326/aafc1b>, 2019.
- Braun, M. H., Malz, P., Sommer, C., Farías-Barahona, D., Sauter, T., Casassa, G., Soruco, A., Skvarca, P., and Seehaus, T. C.: Constraining glacier elevation and mass changes in South America, *Nat. Clim. Change*, 9, 130–136, <https://doi.org/10.1038/s41558-018-0375-7>, 2019.
- 235 Carr, J. R., Stokes, C., and Vieli, A.: Recent retreat of major outlet glaciers on Novaya Zemlya, Russian Arctic, influenced by fjord geometry and sea-ice conditions, *J. Glaciol.*, 60, 155–170, <https://doi.org/10.3189/2014JoG13J122>, 2014.
- Ciraci, E., Velicogna, I., and Sutterley, T.: Mass Balance of Novaya Zemlya Archipelago, Russian High Arctic, Using Time-Variable Gravity from GRACE and Altimetry Data from ICESat and CryoSat-2, *Remote Sens.*, 10, 1817, <https://doi.org/10.3390/rs10111817>, 2018.
- 240 Ciraci, E., Velicogna, I., and Swenson, S.: Continuity of the Mass Loss of the World’s Glaciers and Ice Caps From the GRACE and GRACE Follow-On Missions, *Geophys. Res. Lett.*, 47, <https://doi.org/10.1029/2019GL086926>, 2020.
- Gardner, A. S., Moholdt, G., Cogley, J. G., Wouters, B., Arendt, A. A., Wahr, J., Berthier, E., Hock, R., Pfeffer, W. T., Kaser, G., Ligtenberg, S. R. M., Bolch, T., Sharp, M. J., Hagen, J. O., van den Broeke, M. R., and Paul, F.: A Reconciled Estimate of Glacier Contributions to Sea Level Rise: 2003 to 2009, *Science*, 340, 852–857, <https://doi.org/10.1126/science.1234532>, 2013.
- 245 Hugonnet, R., McNabb, R., Berthier, E., Menounos, B., Nuth, C., Girod, L., Farinotti, D., Huss, M., Dussailant, I., Brun, F., and Kääh, A.: Accelerated global glacier mass loss in the early twenty-first century, *Nature*, 592, 726–731, <https://doi.org/10.1038/s41586-021-03436-z>, 2021.
- Jacob, T., Wahr, J., Pfeffer, W. T., and Swenson, S.: Recent contributions of glaciers and ice caps to sea level rise, *Nature*, 482, 514–518, <https://doi.org/10.1038/nature10847>, 2012.
- 250 Jansen, E., Christensen, J. H., Dokken, T., Nisancioglu, K. H., Vinther, B. M., Capron, E., Guo, C., Jensen, M. F., Langen, P. L., Pedersen, R. A., Yang, S., Bentsen, M., Kjær, H. A., Sadatzki, H., Sessford, E., and Stendel, M.: Past perspectives on the present era of abrupt Arctic climate change, *Nat. Clim. Change*, 10, 714–721, <https://doi.org/10.1038/s41558-020-0860-7>, 2020.
- Li, J., Li, Z.-W., Hu, J., Wu, L.-X., Li, X., Guo, L., Liu, Z., Miao, Z.-L., Wang, W., and Chen, J.-L.: Investigating the bias of TanDEM-X digital elevation models of glaciers on the Tibetan Plateau: impacting factors and potential effects on geodetic mass-balance measurements, *J. Glaciol.*, 1–14, <https://doi.org/10.1017/jog.2021.15>, 2021.

- Matsuo, K. and Heki, K.: Current Ice Loss in Small Glacier Systems of the Arctic Islands (Iceland, Svalbard, and the Russian High Arctic) from Satellite Gravimetry, *Terr. Atmospheric Ocean. Sci.*, 24, 657, [https://doi.org/10.3319/TAO.2013.02.22.01\(TibXS\)](https://doi.org/10.3319/TAO.2013.02.22.01(TibXS)), 2013.
- 260 Melkonian, A. K., Willis, M. J., Pritchard, M. E., and Stewart, A. J.: Recent changes in glacier velocities and thinning at Novaya Zemlya, *Remote Sens. Environ.*, 174, 244–257, <https://doi.org/10.1016/j.rse.2015.11.001>, 2016.
- Millan, R., Dehecq, A., Trouve, E., Gourmelen, N., and Berthier, E.: Elevation changes and X-band ice and snow penetration inferred from TanDEM-X data of the Mont-Blanc area, in: 2015 8th International Workshop on the Analysis of Multitemporal Remote Sensing Images (Multi-Temp), 2015 8th International Workshop on the Analysis of Multitemporal Remote Sensing Images (Multi-Temp), Annecy, France, 1–4, <https://doi.org/10.1109/Multi-Temp.2015.7245753>, 2015.
- 265 Moholdt, G., Wouters, B., and Gardner, A. S.: Recent mass changes of glaciers in the Russian High Arctic: GLACIER MASS CHANGES, RUSSIAN ARCTIC, *Geophys. Res. Lett.*, 39, n/a-n/a, <https://doi.org/10.1029/2012GL051466>, 2012.
- Morris, A., Moholdt, G., and Gray, L.: Spread of Svalbard Glacier Mass Loss to Barents Sea Margins Revealed by CryoSat-2, *J. Geophys. Res. Earth Surf.*, 125, <https://doi.org/10.1029/2019JF005357>, 2020.
- 270 Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., Kienholz, C., Miles, E. S., Moholdt, G., Mölg, N., Paul, F., Radić, V., Rastner, P., Raup, B. H., Rich, J., Sharp, M. J., and The Randolph Consortium: The Randolph Glacier Inventory: a globally complete inventory of glaciers, *J. Glaciol.*, 60, 537–552, <https://doi.org/10.3189/2014JoG13J176>, 2014.
- Rott, H., Floricioiu, D., Wuite, J., Scheiblauer, S., Nagler, T., and Kern, M.: Mass changes of outlet glaciers along the Nordenskjöld Coast, northern Antarctic Peninsula, based on TanDEM-X satellite measurements: TanDEM-X Antarctic Peninsula glaciers, *Geophys. Res. Lett.*, 41, 8123–8129, <https://doi.org/10.1002/2014GL061613>, 2014.
- Sánchez-Gómez, P., Navarro, F. J., Benham, T. J., Glazovsky, A. F., Bassford, R. P., and Dowdeswell, J. A.: Intra- and inter-annual variability in dynamic discharge from the Academy of Sciences Ice Cap, Severnaya Zemlya, Russian Arctic, and its role in modulating mass balance, *J. Glaciol.*, 65, 780–797, <https://doi.org/10.1017/jog.2019.58>, 2019.
- 280 Seehaus, T., Malz, P., Sommer, C., Lippl, S., Cochachin, A., and Braun, M.: Changes of the tropical glaciers throughout Peru between 2000 and 2016 – mass balance and area fluctuations, *The Cryosphere*, 13, 2537–2556, <https://doi.org/10.5194/tc-13-2537-2019>, 2019.
- Strozzi, T., Paul, F., Wiesmann, A., Schellenberger, T., and Käab, A.: Circum-Arctic Changes in the Flow of Glaciers and Ice Caps from Satellite SAR Data between the 1990s and 2017, *Remote Sens.*, 9, 947, <https://doi.org/10.3390/rs9090947>, 2017.
- 285 Wessel, B., Bertram, A., Gruber, A., Bemm, S., and Dech, S.: A NEW HIGH-RESOLUTION ELEVATION MODEL OF GREENLAND DERIVED FROM TANDEM-X, *ISPRS Ann. Photogramm. Remote Sens. Spat. Inf. Sci.*, III–7, 9–16, <https://doi.org/10.5194/isprsannals-III-7-9-2016>, 2016.
- Wouters, B., Gardner, A. S., and Moholdt, G.: Global Glacier Mass Loss During the GRACE Satellite Mission (2002-2016), *Front. Earth Sci.*, 7, 96, <https://doi.org/10.3389/feart.2019.00096>, 2019.
- 290 Zhao, J. and Floricioiu, D.: THE PENETRATION EFFECTS ON TANDEM-X ELEVATION USING THE GNSS AND LASER ALTIMETRY MEASUREMENTS IN ANTARCTICA, *ISPRS - Int. Arch. Photogramm. Remote Sens. Spat. Inf. Sci.*, XLII-2/W7, 1593–1600, <https://doi.org/10.5194/isprs-archives-XLII-2-W7-1593-2017>, 2017.

Zheng, W., Pritchard, M. E., Willis, M. J., Tepes, P., Gourmelen, N., Benham, T. J., and Dowdeswell, J. A.: Accelerating glacier mass loss on Franz Josef Land, Russian Arctic, *Remote Sens. Environ.*, 211, 357–375, 295 <https://doi.org/10.1016/j.rse.2018.04.004>, 2018.

Zheng, W., Pritchard, M. E., Willis, M. J., and Stearns, L. A.: The Possible Transition From Glacial Surge to Ice Stream on Vavilov Ice Cap, *Geophys. Res. Lett.*, 46, 13892–13902, <https://doi.org/10.1029/2019GL084948>, 2019.



300 **Figure 1** a) Backscatter intensity of different TanDEM-X DEM acquisition months versus elevation on Novaya Zemlya. Black lines
 305 indicate average backscatter aggregated within 50m elevation bins. b) Mean elevation change rates of DEM differences between
 winter 2010/11 and winter 2016/17 (WW, triangles) and winter 2010/11 and September 2016 (WA, dots). c) Altitudinal distribution
 of mean backscatter intensity (aggregated in 10m elevation bins) of September and winter 2016/17 SAR data on overlapping glacier
 areas (i.e. areas which were acquired in September and winter 2016/17). d) Differences in estimated signal penetration between
 September and winter 2016/17 versus mean backscatter intensity of September 2016 acquisitions on overlapping glacier areas (ag-
 gregated within 0.1 db backscatter intervals between -20 to -28 db).




Figure 2: Surface elevation changes of glaciers on Franz Josef Land (a), Severnaya Zemlya (b), and Novaya Zemlya (c) between 2010 and 2017. Hatched areas indicate glaciers without coverage by TanDEM-X. Respective average elevation change rates and total/measured glacier areas within 50m elevation bins are shown in Figures 1 (d)-(f). The hypsometric distribution of Severnaya Zemlya does not include the surge of the Vavilov ice cap (RGI60-09.00971). Elevation changes of Novaya Zemlya were corrected for SAR-signal penetration.

310

315

Table 1: Overview of glacier elevation and mass change in the Russian Arctic between 2010/11 and 2016/17. Glacier areas (S) are derived from the Randolph Glacier Inventory 6.0. Its spatial coverage by elevation change measurements ($S_{mea.}$) is stated in percent. Dh/dt shows elevation change rates as measured by TanDEM-X while $dh/dt_{adj.}$ includes the SAR signal-penetration corrected elevation change rate of Novaya Zemlya. dM/dt and $dM/dt_{adj.}$ are the respective glacier mass change rates using a volume-to-mass conversion factor of 850 kg m^{-3} (a) and 900 kg m^{-3} (b).

*Acquisition date offsets adjusted for Novaya Zemlya.

Region	S [km ²]	S mea. [%]	dh/dt [m a ⁻¹]	dh/dt adj. [m a ⁻¹]	dM/dt [Gt a ⁻¹]a	dM/dt adj. [Gt a ⁻¹]a	dM/dt [Gt a ⁻¹]b	dM/dt adj. [Gt a ⁻¹]b
Franz Josef Land	12750	96	-0.48±0.04		-5.14±0.43		-5.45±0.45	
Severnaya Zemlya	16529	97	-0.34±0.12		-4.70±1.31		-4.98±1.38	
Novaya Zemlya	22117	91	-0.53±0.23	-0.68±0.46*	-9.95±3.14	-12.92±6.18	-10.54±3.31	-13.68±6.54*
Russian Arctic	51707	93	-0.46±0.15	-0.52±0.24*	-20.05±3.47	-23.06±6.41	-21.23±3.66	-24.42±6.79*

320