

Firn changes at Colle Gnifetti revealed with a high-resolution process-based physical model approach

Enrico Mattea¹, Horst Machguth¹, Marlene Kronenberg¹, Ward van Pelt², Manuela Bassi³, and Martin Hoelzle¹

¹Department of Geosciences, University of Fribourg, Fribourg, Switzerland

²Department of Earth Sciences, Uppsala University, Uppsala, Sweden

³Department of Forecasting Systems, Regional Agency for Environmental Protection of Piedmont, Turin, Italy

Correspondence: Enrico Mattea (enrico.mattea@unifr.ch)

Abstract. Our changing climate is expected to affect ice core records as cold firn progressively transitions to a temperate state. Thus, there is a need to improve understanding and further develop quantitative process modeling, to better predict cold firn evolution under a range of climate scenarios. Here we present the application of a distributed, fully coupled energy balance model, to simulate ~~high-alpine~~ cold firn at ~~Colle Gnifetti~~ the high-alpine glaciated saddle of Colle Gnifetti (Swiss/Italian Alps) over the period 2003–2018. ~~For the first time, we force such a~~ We force the model with high-resolution, long-term and extensively quality-checked meteorological data measured in closest vicinity of the firn site, at the ~~Capanna Margherita~~ (highest automatic weather station in Europe (Capanna Margherita, 4560 m a.s.l.)). The model incorporates the spatial variability of snow accumulation rates, and is calibrated using several, partly unpublished high-altitude measurements from the Monte Rosa area. The simulation reveals a very good overall agreement in the comparison with a large archive of firn temperature profiles. ~~The rate of firn warming at 20 m depth is estimated at 0.44 °C per decade.~~ Our results show that surface melt over the glaciated saddle is increasing by 3–4 mm w.e. yr⁻² depending on the location (29–36 % in 16 years), although with large inter-annual variability. Analysis of modeled melt indicates ~~a marked tendency towards~~ the frequent occurrence of small melt events (< 4 mm w.e.), which collectively represent a significant fraction of the melt totals. ~~Atmospheric humidity is found to be a prominent control over melt occurrence, with considerable amounts of sublimation happening in dry conditions. For future developments, we recommend implementing a physical simulation of meltwater infiltration, to be calibrated with more measurements of percolation in cold firn.~~ Modeled firn warming rates at 20 m depth are relatively uniform above 4450 m a.s.l. (0.4–0.5 °C per decade). They become highly variable at lower elevations, with a marked dependence on surface aspect and absolute values up to 2.5 times the local rate of atmospheric warming. Our distributed simulation contributes to the understanding of the thermal regime and evolution of a prominent site for alpine ice cores, and may support the planning of future core drilling efforts. Moreover, thanks to an extensive archive of measurements available for comparison, we also highlight the possibilities of model improvement most relevant to the investigation of future scenarios, such as the fixed-depth parametrized routine of deep preferential percolation.

1 Introduction

Cold firn and ice – defined by negative temperatures year-round – are recognized as a valuable archive of past atmospheric conditions, accessed through ice cores (Wagenbach et al., 2012) (e.g., Masson-Delmotte et al., 2006; Lüthi et al., 2008; Wolff et al., 2010; Wagenbach et al., 2012). In the recrystallization and recrystallization-infiltration firn facies, meltwater infiltration is respectively absent or limited to the near-surface layer of the firn (Shumskii, 1964; Hoelzle et al., 2011). This enables preservation of the original layering of accumulated snow, which can be dated to provide an atmospheric record including greenhouse gases, aerosols, precipitation, and isotopic temperature proxies (Wagenbach et al., 2012) (e.g., Preunkert et al., 2001; Barbante et al., 2004; Thevenon et al., 2009; Konrad et al., 2012).

The longest records are found in ice cores from the polar regions. Nonetheless, cold firn is also present in the Alps above 3400–4150 m a.s.l., depending on location and aspect (Suter et al., 2001). Such alpine cold firn is located close to major, historical sources of European anthropogenic emissions, thus providing a particularly valuable record of man-made changes to atmospheric composition (Wagenbach et al., 2012) (Jenk et al., 2006; Thevenon et al., 2009; Legrand et al., 2013). Moreover, the Alpine region features a historically high density of meteorological observations as well as other paleoclimatic records (such as tree rings and speleothems), enabling calibration and comparison of the atmospheric archives (Wagenbach et al., 2012) and climatic archives (Wagenbach et al., 2012, and references therein).

Besides its importance for ice core studies, cold firn also acts as a buffer against mass losses brought upon glaciers by a warming climate. Specifically, meltwater refreezing close to the surface does not contribute to water runoff: thus an increased input in the firn surface energy balance (SEB), with enhanced meltwater production, does not directly affect mass balance (e.g., Harper et al., 2012). As a result, rising temperatures – instead of mass losses – are the main expression of 20th-century atmospheric warming in cold firn (Haeberli and Beniston, 1998; Buri, 2013) (Gilbert and Vincent, 2013; Hoelzle et al., 2011).

Climate change is expected to trigger a progressive transition from cold to temperate firn, naturally advancing from the lower elevations towards the higher (Darms, 2009) (Lüthi and Funk, 2001; Vincent et al., 2007; Gilbert et al., 2010). Expected consequences for sites of presently cold firn are the onset of mass loss (e.g., van Pelt and Kohler, 2015) and an irremediable degradation of the climatic archive, induced by meltwater infiltration to increasing depths (Hoelzle et al., 2011; de Blasi et al., 2019). Then (Gabrielli et al., 2010; Hoelzle et al., 2011). Moreover, a change of thermal regime could affect the stability of cold-based hanging glaciers, with potentially hazardous consequences (Gilbert et al., 2015). Thus, a better understanding of this transition will become crucial to the continued viability of ice core campaigns, as well as the mitigation of glacier hazards and the prediction of future runoff regimes in high-alpine and polar catchments. Particularly valuable will be the acquisition of quantitative modeling capabilities and localized information on to estimate the timing and uncertainties of firn changes, also incorporating the regularly updated climatic scenarios.

Among alpine cold firn sites, the Colle Gnifetti saddle (CG; Fig. 1) in the Monte Rosa range (4450 m a.s.l., Swiss/Italian Alps) stands out for the dense coverage of glaciological measurements, acquired continuously over almost 50 years. The very low annual accumulation rates of the area (between 0.3 and 1.2 m w.e.) enable ice core records covering the last millennium and

potentially extending into the late Pleistocene ~~-, more (up to 19 kyr B.P.), further back~~ than any other glaciated site in the Alps (Jenk et al., 2009; Wagenbach et al., 2012; Bohleber et al., 2018). Important climatological results from CG ice cores include a 1200-year time series of air temperature, reconstructed from mineral dust proxies (Bohleber et al., 2018), as well as a glacio-chemical record of anthropogenic alterations to atmospheric composition, starting before industrialization and extending to the recent emission reductions of some pollutant species (Schwikowski, 2004; Barbante et al., 2004; More et al., 2017; Gabrieli et al., 2011).

Parallel to ice core investigations, borehole measurements starting in 1976 (Haeberli and Funk, 1991) provide a detailed picture of the thermal conditions at the CG saddle. Firn temperatures below the depth of annual fluctuations (about 18 m) ~~range from were measured in the range of~~ -13.5 °C to -10 °C ~~-, depending on aspect (Suter and Hoelzle, 2002) by Suter and Hoelzle (2002), with a clear dependence on aspect.~~ Haeberli and Funk (1991) found almost steady-state temperature conditions in ~~englacial profiles-an englacial profile~~ from 1983, noting only a weak influence of meltwater refreezing. Lüthi and Funk (2001) reported temperature profiles from 1995 having striking bends at a depth of about 30 m, and interpreted them as the first published clues to a non-steady firn warming situation. Subsequently, Hoelzle et al. (2011) found evidence of accelerated englacial warming exceeding the air temperature increase, ~~exposing-highlighting~~ an enhanced role of meltwater percolation.

Several studies have previously applied simple firn models at CG, to reproduce englacial temperatures from idealized boundary conditions (Haeberli and Funk, 1991; Lüthi, 2000; Lüthi and Funk, 2001). Suter (2002) presented a distributed model study of the area, simulating one year of SEB at daily resolution, and formulated one-dimensional firn temperature predictions according to a simple parametric model. Buri (2013) used the coupled model GeoTOP to simulate energy balance and sub-surface temperatures at several locations in the CG area.

A major challenge for firn models at CG is the complex boundary condition of surface accumulation: the low accumulation rates are due to an extreme wind scouring of the snow surface, favored by the west-east saddle orientation. This process is countered by solar radiation through melt consolidation. Thus, snow accumulation has a strong spatial gradient according to terrain aspect, and is biased towards summer ~~-, a compared to climatological precipitation.~~ A significant interannual variability is also observed ~~(Alean et al., 1983; Wagenbach et al., 2012), including years with no residual accumulation; wind erosion sometimes even affects the previous year's layer (Alean et al., 1983; Lüthi, 2000; Wagenbach et al., 2012).~~ Surface accumulation directly controls the cold content and initial stratigraphy of the firn, ~~as well as the rate of heat advection (e.g., Kuipers Munneke et al., 2014).~~ However, the complex accumulation patterns at CG have not been addressed in past modeling studies, even within models coupling the energy balance to the sub-surface. ~~Suter et al. (2001) considered accumulation only as a proxy for vertical velocities at the surface, to account for ice advection in a minimal firn temperature model. Buri (2013)-The profile calculations by Suter et al. (2001) only considered heat transfer by conduction, hence neglecting vertical advection by surface accumulation. Buri (2013) simulated individual borehole locations in the CG area, relying for all of them on weather data (including precipitation) collected at the Corvatsch automatic weather station (AWS), almost 160 km away and over 1 km lower in altitude. More recently, Licciulli (2018) established a flow model independent on-of surface accumulation,~~ and obtained distributed accumulation from the model on the assumption of steady state conditions.

Moreover, surface melt – increasingly occurring at high altitudes in a warming climate – is expected to play a central role in the transition from cold to temperate firn. Gilbert et al. (2014b) found on Mont Blanc that melt ~~patterns can events~~ strongly affect the overall energy balance of a cold firn pack in summer, because the emission of long-wave radiation from the surface becomes limited as surface temperatures reach 0 °C. The resulting energy excess is then released within the firn pack by meltwater refreezing at depth. To this day, little knowledge exists on the atmospheric conditions leading to melt events in cold firn areas, or the dynamics of such melt events. Suter et al. (2001) modeled CG firn temperatures with a 1 day time-step, too coarse compared to the time scales of melt and infiltration processes. Buri (2013), while using an hourly time-step, did not examine melt amounts and dynamics. Licciulli (2018), focusing on ice flow, modeled the firn at CG with a 1 year time-step, assuming no meltwater infiltration in the firn.

Physical models of high-alpine cold firn were also notably applied to the Col du Dôme site (CdD) in the Mont Blanc range (French Alps). At 4250 m a.s.l., CdD has a thermal regime similar to that of CG (Suter and Hoelzle, 2002); accumulation is generally higher and less affected by wind erosion (e.g., Preunkert et al., 2000). At CdD, Gilbert et al. (2014b) present a coupled surface energy balance and sub-surface heat transport model, including physical representation of meltwater percolation and refreezing, but only at a single point location. Gilbert et al. (2014a) use a simplified degree-day based approach to distribute near-surface temperatures, to force a 3D glacier thermo-mechanical model.

Here, we present the application of a distributed, physically-based coupled model to simulate firn evolution at CG, over a 16 year period on a high-resolution gridded domain. After validation against a large archive of firn temperature measurements, the model is used to investigate the conditions and dynamics of high-alpine melt events. For the first time, model forcing is based on the extensively processed hourly time series of the Capanna Margherita (CM) AWS (Martorina et al., 2003), which benefits from an exceptional location at 4560 m a.s.l. and within 400 m of the CG saddle point.

The data used within the study are described in Sect. 2, the coupled model in Sect. 3, the results are presented in Sect. 4 and discussed in Sect. 5, while the last section provides the conclusions and some perspectives for future cold firn research.

2 Data

2.1 Meteorological time series

Meteorological forcing of the coupled model consists of an hourly time series, describing air temperature, atmospheric pressure, wind speed, relative humidity, fractional cloud cover and precipitation (Fig. 2). The series was primarily assembled using data from the CM AWS, located on the Signalkuppe summit at 4560 m a.s.l. (Fig. 1a). The station records air temperature, barometric pressure, wind speed and direction, and global radiation (Table 1). The measured parameters are available as hourly instantaneous values.

The consistent availability (> 95 %) of long-term data (since mid-2002) makes the series a valuable data-set for high-alpine research. Still, an extensive evaluation and processing of the data is essential due to the extreme measuring conditions (Martorina et al., 2003). In particular, freezing of the anemometer – as well as snow and ice accumulation interfering with the pyranometer – are relatively common occurrences. Therefore, the hourly data from eight other high-altitude AWS located in

Table 1. Sensors installed at the CM AWS.

Parameter	Sensor	Notes
Air temperature	CAE TU20 thermometer	Rated accuracy 0.2 °C Rated ambient radiation influence < 0.8 °C Not artificially ventilated
Barometric pressure	CAE BA20 barometer	Rated accuracy 0.5 hPa
Global radiation	CAE HE20/K pyranometer	Rated daily accuracy 5 % Wavelength band 305–2800 nm
Wind speed	CAE VV20 cup taco-anemometer	Rated accuracy 0.07 m s ⁻¹ or 1 %
Wind direction	CAE DV20 gonio-anemometer	Rated accuracy 2.8°

the region (Fig. 1b, id 2–9 in Table 2) were used to perform quality checks, to fill gaps in the CM time series, and to provide data for parameters not measured at CM. The selection of stations to include was determined by the availability of different parameters measured at each site, by the effort to provide an unbiased geographic coverage in all directions from the CM AWS, and by the station tendency towards concurrent failures in challenging conditions, such as winter storms.

All AWS series were pre-processed to remove clock errors, detected with cross-correlation analysis. Then each series was entirely reconstructed from the data of the best correlated others using quantile mapping (Cannon et al., 2015; ?)(Cannon et al., 2015; Feigenv
, to provide a reference for robust outlier detection. High-resolution reanalysis series of the parameters measured at CM were also collected as an additional basis for comparison (COSMO-REA2 and COSMO-REA6: Wahl et al., 2017; Bollmeyer et al., 2015; Frank et al., 2018). Due to the large volume of data involved, an automated pre-filtering routine was implemented, based on objective criteria (absolute values, rates of change, comparison with reconstructed series and reanalysis) to mark single values as potential outliers, which were then manually checked. The CM AWS was always processed last in order to be compared to the highest-quality data. After quality check, all gaps in the hourly CM series were filled with the corresponding reconstructed values. The close match of the reconstructed series (Table 3) despite the large elevation differences involved (Table 2) confirms the benefit of the extensive processing described.

The CM AWS does not record series of relative humidity (RH) or cloudiness, which are required within the model input. Thus RH was entirely derived from the other quality-checked time series; cloudiness was reconstructed from the incoming SW radiation at CM when available, and otherwise computed from the incoming LW radiation and observed cloud cover, respectively at Stockhorn and Plateau Rosa. Lapse rates of temperature and pressure – used to extend the AWS record to the gridded model domain – were computed from the difference between CM and the other stations. Despite the large elevation differences involved (Table 2), a robust computation of lapse rates was possible thanks to the availability of multiple stations within a close distance; the spread of the individual rates was usually one order of magnitude smaller than their absolute values. Moreover, the close proximity of the CM AWS to the domain minimizes the amount of extrapolation needed, as does the narrow elevation range of the domain itself. The other variables (wind speed, cloud cover and RH) were held constant over the model

Table 2. AWS series included in this study. Numbering corresponds to Fig. 1b. X, Y and Z are metric CH1903/LV03 coordinates. $d_{CG-d_{SP}}$ is the horizontal distance to the CG saddle point (labeled SP in Fig. 1a). Stations 10 and 11 were deployed temporarily for energy-balance studies; permanent stations 3, 4 and 6 were installed between 2006 and 2010, while the others were already operational before the CM AWS. Further AWS information is reported in Mattea (2020).

Id	Name	X [m]	Y [m]	Z [m a.s.l.]	Variables ^a	$d_{CG-d_{SP}}$ [m]	Reference
1	Capanna Margherita	634007	86266	4560	T, P, W, S	380	ARPA Piemonte (2020)
2	Gornergrat	626900	92512	3129	T, P, H, W, S, D	9100	MeteoSwiss (2020)
3	Stockhorn	629900	92850	3415	T, H, N, W, S, L	7400	Gruber et al. (2004)
4	Monte Rosa Plattje	629149	89520	2885	T, P, H, W, S	5500	MeteoSwiss (2020)
5	Passo dei Salati	633339	80689	2970	T, H, W	5900	Visit Monte Rosa (2020)
6	Macugnaga Rifugio Zamboni	637094	88977	2075	T, P, H, N, S, R	4200	ARPA Piemonte (2020)
7	Passo del Moro	641664	94075	2820	T, H, R, W, S	11000	ARPA Piemonte (2020)
8	Plateau Rosa	620840	87123	3488	T, P, H, W, C	12900	MeteoAM (2020)
9	Bocchetta delle Pisse	635910	80543	2410	T, N, R, W, S	6400	ARPA Piemonte (2020)
10	Seserjoch	633727	85785	4292	T, H, N, W, S, L, G, F	820	Suter et al. (2004)
11	Colle del Lys	632665	85360	4236	T, H, N, S, W	1700	Rossi et al. (2000a,b)

^a T: air temperature. P: atmospheric pressure. W: wind speed and direction. S: SW radiation. L: LW radiation. H: relative humidity. D: sunshine duration. N: snow height. C: cloud cover. R: precipitation. G: surface temperature. F: firn temperature. Not all measured variables were used in the study.

domain, as no measurements were available to estimate their spatial variability. Further details on the series processing can be found in Mattea (2020). Figure 2 shows the final time series of all meteorological parameters assigned to CM and used as model input.

150 To determine some of the model calibration parameters (Table 4), we also re-analysed data from two additional weather and energy-balance stations (10 and 11 in Table 2), temporarily installed above 4000 m a.s.l. within the ALPCLIM project (Auer et al., 2001). The Seserjoch station operated from September 1998 to October 2000 for the detailed SEB investigations of Suter et al. (2004); the Colle del Lys station (Rossi et al., 2000a,b) was active between 1996 and 2000.

2.2 Sub-surface data

155 For model validation (Sect. 4.1) we used archived firn temperature profiles (Fig. 1a), measured between 2003 and 2018. **Beside** Besides the CG saddle, we included two profiles from the Grenzletscher slopes and four from Seserjoch. In total, we considered 25 temperature profiles from 18 boreholes, some locations having been measured more than once. Detailed description of the profiles is reported in Hoelzle et al. (2011) and GLAMOS (2017). Earlier information on the deep temperatures (Haeberli and Funk, 1991) was also used as reference during model set-up (Sect. 3.4).

160 Moreover, we compiled point measurements of annual accumulation (Fig 3a), derived from layer thicknesses in GPR profiles (Konrad et al., 2013) and from the stake network of Suter and Hoelzle (2002), as well as from archived ice core measurements

Table 3. ~~Performance metrics~~ Summary statistics of the ~~simulated~~ CM weather series ~~with respect~~ processing. The mean and standard deviation (SD) are computed on the final hourly series used as model input. The root mean square error (RMSE) and mean signed deviation (BIAS) refer to the comparison between the reconstructed series and the quality-checked hourly measurements (Sect. 2.1).

Variable	<u>RMS-Mean</u>	<u>SD</u>	<u>RMSE</u>	BIAS
Air temperature [°C]	<u>-12.03</u>	<u>7.27</u>	1.82	0.02
Barometric pressure [hPa]	<u>576.70</u>	<u>9.59</u>	1.26	0.00
Wind speed [m s ⁻¹]	<u>4.47</u>	<u>5.14</u>	5.77	-0.08
Global radiation [W m ⁻²]	<u>194</u>	<u>290</u>	114	-2

(densities and annual dating) acquired at CG between 1982 and 2019. In total, 14 core profiles were available; only one (core *Zumsteinkern*, from 1991) is located on the high-accumulation, south-facing Zumsteinspitze slope (Fig. 3a). Description of the individual cores can be found in Licciulli (2018) and Lier (2018).

165 Finally, we hand-drilled a 5.5 m core (*unifr-2019*, Fig. 1a) near the CG saddle point on June 25, 2019, analyzing density and stratigraphy in the field. Core description is presented in ~~the appendix~~ Appendix A.

2.3 Topography

The model grid was based on a 20 m digital elevation model (DEM). Due to the Swiss-Italian border crossing the ~~modeling~~ simulation domain, the DEM was produced by merging the stereo-photogrammetric SwissAlti3D dataset (acquired in 2015
170 with a vertical accuracy of 1–3 m) on the Swiss side, and the ICE LiDAR digital terrain model (acquired in 2011 with a vertical accuracy of 0.3–0.6 m) on the Italian side (Regione Piemonte, 2011). Elevation mismatch along the border (RMS error of 3.2 m) was corrected via smoothing, to avoid unwanted biases in slope and aspect.

3 The coupled energy-balance and firn model (EBFM)

The coupled model used in this work was introduced by van Pelt et al. (2012) to simulate mass balance of Nordenskiöldbreen
175 (Svalbard). Driven by a meteorological time series, the model computes energy fluxes on the snow surface: short-wave (*SW*) and long-wave (*LW*) radiation, sensible (Q_{SH}) and latent (Q_{LH}) turbulent fluxes, heat advection from rainfall (Q_{rain}) and heat conduction into the snow or ice (Q_g). Then the SEB (Eq. 1) is solved for surface temperature and melt amounts: these, together with the lower boundary condition of geothermal heat flux Q_{ground} , drive the sub-surface evolution.

$$Q_{melt} = SW_{net} + LW_{net} + Q_{SH} + Q_{LH} + Q_{rain} + Q_g \quad (1)$$

180 Simulation of surface processes is developed along the lines of Klok and Oerlemans (2002), while the multi-layer sub-surface snow model is based on the SOMARS approach (Simulation Of glacier surface Mass balance And Related sub-surface processes, Greuell and Konzelmann, 1994). In this work, we used the model version described by van Pelt et al. (2019) ~~was used~~,

with a parametrized water percolation routine simulating preferential flow (Marchenko et al., 2017), and an updated scheme for albedo decay based on Bougamont et al. (2005). This model participated in the firn meltwater Retention Model Intercomparison Project (RetMIP) under the designation "UppsalaUniDeepPerc" (Vandecrux et al., 2020).

In the following we highlight the main EBFM routines and their respective adaptations to the CG setting. The model was originally developed over large, polythermal Arctic glaciers: set-up for a high-alpine cold firn saddle was possible thanks to a rich archive of energy-balance measurements from the Monte Rosa area.

3.1 Radiative fluxes

The model computes incoming SW radiation as

$$SW_{in}(x, y) = TOA_{shaded}(x, y) t_{rg} t_w t_a t_{cl} \quad (2)$$

$$t_{cl}(n) = 1 - an - bn^2, \quad (3)$$

where $TOA_{shaded}(x, y)$ is the unattenuated top-of-atmosphere radiation ($W m^{-2}$) corrected for topographic shading and angle of incidence on the surface, t_{rg} , t_w , t_a and t_{cl} gaseous, water vapor, aerosol and cloud transmissivities, n fractional cloud cover, and a and b calibration parameters. Values for a and b (Table 4) were derived from Greuell et al. (1997) as calibrated in high-alpine terrain, compared to unlike the EBFM defaults which were tuned from measurements in the Arctic. ~~The choice was supported by a distribution~~ (van Pelt et al., 2012). The parameter values of Greuell et al. (1997) increase the dependence of t_{cl} on the cloud fraction, allowing a stronger decrease of the incoming SW flux under overcast conditions. Such a higher dependence was confirmed by an analysis of the ~~incoming SW flux~~, distribution of the incoming radiation flux measured at the CM AWS ~~under overcast sky conditions~~, supporting our parameter choices (Table 4).

Reflected SW radiation is controlled by a broadband, isotropic surface albedo α_{snow} (Eq. 4). Albedo evolution is modeled after Oerlemans and Knap (1998) as an exponentially decaying function of time ~~since t~~ since the last significant snowfall (Eq. 5), bounded by constant values for fresh snow ~~and firn~~, ~~Decay~~ (α_{fresh}) and firn (α_{firn}). The precipitation threshold to reset albedo is $0.1 \text{ mm w.e. h}^{-1}$ (van Pelt et al., 2019). The time-scale of albedo decay is a function of snow surface temperature (Eq. 6) to account for slower metamorphism in cold conditions (Bougamont et al., 2005; van Pelt et al., 2019):

$$SW_{out} = \alpha_{snow} SW_{in} \quad (4)$$

$$\alpha_{snow}(t) = \alpha_{firn} + (\alpha_{fresh} - \alpha_{firn}) e^{-\frac{t}{t^*}} \quad (5)$$

$$t^*(T_s) = \begin{cases} t_{wet}^* & T_s = 0 \text{ } ^\circ\text{C} \\ t_{dry}^* + K |max(T_s, T_{max, t^*})| & T_s < 0 \text{ } ^\circ\text{C}, \end{cases} \quad (6)$$

where t^* is the computed albedo decay time-scale (d), T_s snow surface temperature (K), t_{wet}^* (t_{dry}^*) time-scale (d) for a melting (dry) surface at $0 \text{ } ^\circ\text{C}$, K a calibration parameter ~~, T snow surface temperature ($d \text{ } ^\circ\text{C}^{-1}$)~~ and T_{max, t^*} a temperature cut-off value (K) for decay slow-down. ~~Values for these parameters~~ (Table 4) ~~were estimated from a new analysis of albedo evolution, measured at the Seserjoch and Colle del Lys stations respectively in 1999 and between 1996 and 2000.~~

Incoming LW radiation is computed with the Stefan-Boltzmann law for grey-body radiation (Eq. 7); sky emissivity is modeled after Konzelmann et al. (1994) as a function of cloud cover, air temperature and humidity (Eqs. 8 and 9):

$$215 \quad LW_{in} = e\sigma T^4 \quad (7)$$

$$e = e_{cs}(1 - n^2) + e_{cl}n^2 \quad (8)$$

$$e_{cs} = 0.23 + bc \left(\frac{VP}{T} \right)^{0.125}, \quad (9)$$

where e_{cs} is σ is the Stefan-Boltzmann constant ($\text{W m}^{-2} \text{K}^{-4}$), e_{cs} clear-sky emissivity, bc a calibration parameter ($\text{K}^{0.125} \text{Pa}^{-0.125}$), VP vapor pressure (Pa), T air temperature (K), e sky emissivity, n fractional cloud cover and e_{cl} cloud emissivity.

220 ~~Parameters b~~

We calibrated several parameters of the radiation routines (Table 4) to reflect the local conditions of the high-alpine CG site. We selected the parameters to be tuned based on their relevance for our site, the availability of local measurements, and the simplicity of comparison within the model result. Within the albedo routine we optimized parameters α_{fresh} , t_{wet}^* and K (Eq. 5 and 6), which refer to situations often observed at CG (respectively, a fresh snow surface, a melting surface, and a
 225 sub-freezing surface). By contrast, we kept the default values for α_{firn} and t_{dry}^* , as they correspond to conditions which are almost never encountered at our site (respectively, a bare firn surface and a non-melting surface at 0 °C). We also used the original formulations of t_{rg} , t_w and t_a (Eq. 2): this because the modeled incoming SW flux (Eqs. 2 and 3) is derived from measured radiation at CM, through the series of reconstructed cloud cover (Sect. 2.1). Thus, the effect of these transmissivity coefficients is already taken into account in the cloud cover series. Furthermore, we optimized parameters c and e_{cl} ~~were~~
 230 estimated for CG from radiation measurements at Seserjoch in the LW radiation module (Eq. 8 and 9), since locally measured values were available from the Seserjoch station.

To perform the tuning, we applied the EBFM radiation routines individually (outside the full model runs), driving them with the meteorological and energy-balance measurements of the Seserjoch and Colle del Lys stations at 10 minute resolution. Then, we adjusted the parameter values to find the best match (in terms of bias and RMSE) between the simulated and measured series
 235 of albedo and incoming LW radiation.

3.2 Turbulent heat fluxes

In the EBFM, turbulent heat exchange is modeled with the glacier katabatic wind parametrization of Oerlemans and Grisogono (2002). This was developed with a focus on large valley glaciers; notably, it computes heat fluxes which are independent ~~on~~ of
 the ambient wind field, since wind speeds are estimated from the katabatic flow model (Oerlemans and Grisogono, 2002). This
 240 was deemed inadequate for a high-alpine, wind-exposed saddle: thus the EBFM computation of turbulent heat exchange was re-implemented, following the bulk aerodynamic equations of Essery and Etchevers (2004). These were chosen due to their operational simplicity, allowing calculation of turbulent fluxes from a single measurement level of wind speeds. The fluxes are

Table 4. EBFM parameters considered for calibration. Additional model parameters not listed here were kept at the default value (van Pelt et al., 2012; van Pelt and Kohler, 2015; van Pelt et al., 2019).

Parameter	Definition	Unit	Value	Source
a	Cloud SW transmissivity coefficient	-	0.233	Greuell et al. (1997), verified with supported by CM A
b	Cloud SW transmissivity coefficient	-	0.415	Greuell et al. (1997), verified with supported by CM A
α_{fresh}	Fresh snow albedo	-	0.83	Tuned from Seserjoch and Colle del Lys radiation
α_{firm}	Firn albedo	-	0.52	EBFM default (van Pelt and Kohler, 2015)
t_{wet}^*	Decay time-scale (melting snow surface)	d	10	Tuned from Seserjoch and Colle del Lys radiation
t_{dry}^*	Decay time-scale (dry snow surface at 0 °C)	d	30	EBFM default (Bougamont et al., 2005)
K	Increase of t_{dry}^* at negative temperatures	d °C ⁻¹	14	Tuned from Seserjoch and Colle del Lys radiation
T_{max,t^*}	Temperatures Snow temperature cut-off for the t_{dry}^* increase	°C	-10	EBFM default (Reijmer et al., 2005)
b_c	Constant in LW emission formula	-	0.420	Tuned from Seserjoch LW measurements
e_{cl}	Clouds emissivity	-	0.960	Tuned from Seserjoch LW measurements
z_0	Surface roughness length	m	0.001	Suter et al. (2004) from Seserjoch wind profiles
Q_{ground}	Geothermal flux	W m ⁻²	0.040	Lüthi and Funk (2001) from CG boreholes
ρ_{fresh}	Fresh snow density	kg m ⁻³	350	Klok and Oerlemans (2002)
z_{lim}	Characteristic depth of meltwater infiltration	m	4	Tuned to CG 20 m firn temperatures

computed as

$$Q_{SH} = \rho_a c_{p,p,a} C_h V_1 (T_1 - T_s) \quad (10)$$

$$245 \quad Q_{LH} = \rho_a L_{s,v} C_h V_1 (q_1 - q_s), \quad (11)$$

where ρ_a is air density (~~ρ~~ [\(kg m⁻³\)](#)), $c_{p,a}$ specific heat of dry air ([\(J kg⁻¹ K⁻¹\)](#)), V wind speed (~~T~~ [air temperature \(m s⁻¹\)](#)), $L_{s,v}$ latent heat of sublimation or vaporization ([\(J kg⁻¹\)](#), chosen depending on the modeled surface temperature), q specific humidity ([\(kg kg⁻¹\)](#)), and the s and 1 subscripts refer respectively to the snow surface and the measurement level (2 m). Exchange coefficient C_h is defined as

$$250 \quad C_h = C_{hm} f_h, \quad (12)$$

where C_{hn} is the value under neutral conditions and f_h a correction for atmospheric stability, expressed in terms of the bulk Richardson number Ri_B :

$$C_{hn} = k^2 \left(\log \left(\frac{z_1}{z_0} \right) \right)^{-2} \quad (13)$$

$$f_h = \begin{cases} (1 + 10Ri_B)^{-1} & Ri_B \geq 0 \text{ (stable)} \\ 1 - 10Ri_B \left(1 + 10C_{hn} \frac{\sqrt{-Ri_B}}{f_z} \right)^{-1} & Ri_B < 0 \text{ (unstable)} \end{cases} \quad (14)$$

$$255 \quad Ri_B = \frac{gz_1}{V_1^2} \left(\frac{T_1 - T_s}{T_1} + \frac{q_1 - q_s}{q_1 + \epsilon(1 - \epsilon)^{-1}} \right) \quad (15)$$

$$f_z = \frac{1}{4} \left(\frac{z_0}{z_1} \right)^{0.5} \quad (16)$$

Here, k is the von Kármán constant, z_1 the measurement level (m), z_0 the surface roughness length (m), g the gravity acceleration (m s^{-2}), and ϵ the ratio of molecular weights for water and dry air.

260 At CG, surface roughness length is a poorly constrained parameter, due to frequent scouring by extreme winds which alter the snow surface. In our simulation, we used the value computed by Suter et al. (2004) from measurements of wind profiles at Seserjoch. In Appendix B we examine the sensitivity of our simulation to this parameter (Fig. B1a/b).

3.3 Precipitation model

The model precipitation routine was adapted to reproduce the extreme spatial gradient of snow accumulation distinctive of the site (Sect. 1). Since the EBFM does not include a blowing snow routine, the simple model of linear precipitation rates with altitude (van Pelt et al., 2019) was replaced by a gridded precipitation time series, already corrected for snow lost to wind scouring. This was computed with a three-phase anomaly method inspired from New et al. (2000), by combining a fixed climatological grid, an annual anomaly series, and a temporal down-scaling coefficient. Specifically, for grid cell (x, y) , at simulation time-step t , in year i , precipitation was expressed as

$$P(x, y, t) = C(x, y)A(i)D(t), \quad (17)$$

270 where C is the long-term annual accumulation climatology (m w.e. yr^{-1}), A the domain-wide annual anomaly, and D the down-scaling coefficient. The main assumption of the method is that spatial patterns of relative accumulation do not change over time. The climatological grid C was assembled by interpolating point values of long-term net accumulation, estimated from the snow mass of dated firn cores and from the mean layer thickness in GPR profiles (Fig. 3a). To extend data coverage and reduce the occurrence of extrapolation, stake measurements from Suter and Hoelzle (2002) were also used in the western and southern domain regions. ~~Accumulation measured at each stake over a single year was~~ Because single-year stake measurements are in principle not representative of the long-term means, we re-scaled to a mean annual estimate by using the overlaps with firn cores and GPR points ~~their values with a conversion factor. We computed this as the mean ratio between the stake and the firn core/GPR point values, taken at the locations of overlap.~~

280 For annual anomalies A , the multiplicative snow mass anomaly of the KCC deep core (Bohleber et al., 2018) was found to be moderately anti-correlated with wind speed measured at the CM AWS (Fig. 3b). Linear fit over 9 annual data points yielded

the formulation

$$A = 1.46 - 0.21\bar{V} \quad (18)$$

where A is core snow mass anomaly and \bar{V} the median wind speed (m s^{-1}) observed at CM over the corresponding year.

~~Anomaly from the KCC core~~ The computed time series of annual anomaly was then applied to the whole domain. Equation 18 represents a minimal model of the inter-annual variability of wind scouring at CG, accounting for increased erosion rates at higher wind speeds.

Finally, the down-scaling coefficients D were computed by normalizing to a unit sum each year of hourly precipitation, averaged over the three closest rain gauges (Passo Monte Moro, Bocchetta delle Pisse and Rifugio Zamboni; Fig. 1b, Table 2). Equation 17 produces an hourly series of gridded precipitation (already corrected for wind erosion) which was used to force the EBFM. The model uses local air temperature to compute the fraction of precipitation falling as snow: this increases linearly from 0 to 100 % within a 2 °C interval, symmetric around a threshold $T_{s/r} = 0.6$ °C.

3.4 ~~Water infiltration~~Sub-surface model

The EBFM ~~features a parametrized water infiltration routine, which routes any liquid water from the snow surface towards depth, instantly and according to a constant vertical distribution (Marehenko et al., 2017),~~ represents the sub-surface with a Lagrangian discretization: model layers move freely along the depth axis, following the addition or removal of mass at the surface. A new layer is created whenever snowfall and riming push the topmost layer thickness beyond a fixed threshold. This approach prevents numerical diffusion and at the same time accounts for heat advection towards depth (van Pelt et al., 2014).

Layer temperature evolves according to processes of heat conduction and water refreezing (van Pelt et al., 2012):

$$\rho_f c_p(T_f) \frac{\partial T_f}{\partial t} = \frac{\partial}{\partial z} \left(\kappa(\rho_f) \frac{\partial T_f}{\partial z} \right) + F L_m \quad (19)$$

where ρ_f and T_f are layer density (kg m^{-3}) and temperature (K), c_p firn heat capacity ($\text{J kg}^{-1} \text{K}^{-1}$), z depth (m), κ effective conductivity ($\text{W m}^{-1} \text{K}^{-1}$), F refreezing rate ($\text{kg m}^{-3} \text{s}^{-1}$) and L_m latent heat of melting (J kg^{-1}). Parametrizations for c_p and κ are taken respectively from Yen (1981) and Sturm et al. (1997):

$$c_p(T_f) = 152.2 + 7.122T_f \quad (20)$$

$$\kappa(\rho_f) = 0.138 - 1.01 \cdot 10^{-3} \rho_f + 3.23 \cdot 10^{-6} \rho_f^2. \quad (21)$$

Layer density is governed by gravitational settling and water refreezing (van Pelt et al., 2012):

$$\frac{\partial \rho_f}{\partial t} = K_g(\rho_f, T_f) + F \quad (22)$$

$$K_g(\rho_f, T) = b_{acc} g(\rho_{ice} - \rho_f) \exp\left(-\frac{E_c}{RT_f} + \frac{E_g}{RT_{avg}}\right) C_{Lig}(b_{acc}). \quad (23)$$

310 Here, K_g is gravitational densification ($\text{kg m}^{-3} \text{s}^{-1}$) as in Arthern et al. (2010), b_{acc} accumulation rate (mm yr^{-1}), ρ_{ice} ice density (kg m^{-3}), R universal gas constant ($\text{J mol}^{-1} \text{K}^{-1}$), E_c (60 kJ mol^{-1}) and E_g (42.4 kJ mol^{-1}) activation energies of creep by respectively lattice diffusion and grain growth, and T_{avg} year-averaged firn temperature (K). C_{Lig} is a correction based on the accumulation rate, accounting for different densification regimes above and below the critical density value of 550 kg m^{-3} (Ligtenberg et al., 2011).

315 In the EBFM the only source of water at depth is infiltration from the surface (after melt, rainfall and moisture condensation), since sub-surface melting is not simulated. The model features a parametrized routine to account for preferential percolation (Marchenko et al., 2017): as long as the modeled near-surface layers are not impermeable. ~~Water at depth can then refreeze, be held by capillary forces as irreducible water, or form slush layers which drain gradually (van Pelt et al., 2012). In this study, the water vertical distribution was chosen to follow the normal law; the maximum depth,~~ liquid water is instantly routed from the surface to a prescribed sub-surface distribution, defined by its shape along the vertical axis (constant, linear or Gaussian) and maximum depth reached (z_{lim}). Here we selected the Gaussian profile and we tuned z_{lim} ~~was tuned~~ to a value of 4 m, to 320 match the 20 m firn temperature of the ~~CG saddle point at a depth of 20 m (Haeberli and Funk, 1991)~~ SP location (Fig. 1a) as measured by Haeberli and Funk (1991). The chosen reference depth allows to minimize the impact of the annual temperature cycle and of any recent temperature trends.

325 Water in the sub-surface can subsequently refreeze until reaching an upper bound on layer density (the value of glacier ice) and temperature (the melting point). Excess water is partly retained by capillary and adhesive forces (irreducible water content), and partly routed to deeper layers until it is depleted; if an impermeable layer is reached, the resulting slush water drains gradually according to the linear reservoir model (van Pelt et al., 2012; Marchenko et al., 2017). The maximum irreducible water content θ_{mi} of a layer (kg kg^{-1}) is computed from its porosity n_p following Schneider and Jansson (2004):

$$\theta_{mi} = 0.0143 \exp(3.3n_p). \quad (24)$$

330 Appendix B presents a sensitivity analysis of the EBFM to parameters z_{lim} and θ_{mi} , among others; further details on the sub-surface model can be found in van Pelt et al. (2012) and Marchenko et al. (2017).

3.5 Model initialisation

The model grid was initialized to steady state conditions by looping eight times over the 2004–2011 weather input: the period was selected to exclude the extreme melt year of 2003 and the increasing temperature warming of the 2010s. Then two main model runs were performed, with 20 m / 1 h and 100 m / 3 h spatio-temporal resolution. The coarser version was introduced to decrease the large computational volume of sub-surface investigations (Fig. 6) and to ~~perform simple sensitivity experiments~~ examine the impact of different spatio-temporal resolutions. On the depth axis, the model grid included 250 layers up to 10 cm thick, for an effective modeling depth of about 20 m ~~(EBFM layers follow a Lagrangian discretization to prevent numerical diffusion due to layer compaction)~~.

4 Results

340 The EBFM computes and logs a wide variety of surface and sub-surface variables. In the following, we focus on firn temperatures and melt amounts as relevant descriptors of current cold firn evolution. For these variables, ~~a conspicuous~~ an extensive archive of field measurements and model estimates exists at CG, allowing ~~verification~~ validation and comparison of our results.

4.1 Firn temperatures

Comparison of modeled firn temperatures to measured borehole profiles (Table 5, Fig. 4 and 5) shows that most model deviations are similar in magnitude to the spatial variability of firn temperatures. Indeed, profiles CG08-1/08 and CG08-2/08 (acquired in the flat region, on the same day and within a radius of 20 m: Fig. 4e/f) report measured temperature differences in excess of 2 °C at all depths. Model residuals show a clear spatial pattern: simulated profiles tend to be too cold on shaded, north-facing slopes, and too warm in the flat or south-facing regions of CG and Seserjoch. As such, model bias is moderately correlated with mean annual accumulation and potential incoming solar radiation (correlation coefficients of respectively 0.42 and 0.64). Moreover, model ~~biases~~ residuals appear to be maintained over time for boreholes with repeated measurements (e.g., CG05-1 and CG13-1).

Table 5. Performance metrics of the modeled firn temperatures. For each case the number of considered profiles is reported in parentheses (also see Sect. 2.2). All metrics are computed as the arithmetic mean of the respective depth-averaged values of each profile. For a uniform comparison across different profiles, both measured and modeled values were linearly interpolated to a 1 cm vertical resolution before computing deviations.

Model run	Profiles	RMS <u>RMSE</u> [°C]	BIAS [°C]
20 m / 1 h	CG only (19)	1.3	-0.4
	All (25)	1.4	-0.3
100 m / 3 h	CG only (19)	1.4	0.0
	All (25)	1.6	0.1

The time series of modeled firn temperatures (Fig. 6) ~~shows~~ show large differences over rather small distances at the CG saddle, depending on surface aspect. Still, relative annual deviations are consistent across locations. The annual cycle on average reaches an amplitude of 40 °C at the snow surface and is fully damped at a depth of 20 m.

355 The model simulates the frequent summer occurrence of infiltration and refreezing, reflected in sudden near-surface warming events (inset in Fig. 6). Magnitude of these events shows a high variability, with especially large heat amounts simulated in the summers of 2003, 2008, 2015 and 2017. With more meltwater refreezing, the sun-exposed ZS slope commonly shows sustained near-melting temperatures in the topmost 4 m; conversely, these conditions almost never appear at the shaded SK location.

360 A slight positive temperature anomaly can be seen at depth after the extreme melt year of 2015, persisting over the following years despite non-record melt amounts. ~~Across the CG saddle, overall firn warming modeled-~~

The spatial distribution of 20 m firn temperatures (Fig. 7a) shows strong spatial gradients, reflecting surface elevation and aspect. Temperatures range from -17 °C at the shaded NE cliff to near-temperate conditions below 4300 m on the Grenzgletscher slope (Fig. 1a).

365 Modeled temperature trends are relatively uniform over the saddle area above 4400 m a.s.l., where the overall 2003-2018 warming at 20 m depth over 2003–2018 amounts to 0.64–0.75–0.75 °C .At the saddle point, this corresponds to a rate of 0.44 (Fig. 7b). The steep NE cliff constitutes an exception: there, the simulation indicates a very slight and non-uniform tendency towards decreasing temperatures. At lower elevations, temperature trends have a much higher spatial variability, ranging from 0 °C per decade yr^{-1} in the near-temperate area to 0.13 °C yr^{-1} (or 2.5 times the atmospheric warming rate) on the west-facing slopes, about 200 m away.

370

4.2 Melt amounts and dynamics

Modeled mean annual melt amounts have an extreme spatial variability (Fig. 8), broadly reflecting surface elevation, slope and aspect. Values increase from less than 1 cm w.e. yr^{-1} on the steepest slopes of the Signalkuppe, to 17 cm w.e. yr^{-1} at the saddle point, and about 23 cm w.e. yr^{-1} on the Zumsteinspitze slope. Even higher melt amounts, exceeding 30 cm w.e. yr^{-1} , are simulated for the lower elevation Grenzgletscher slopes (towards the western border of the domain) and for Seserjoch. Grid average is 21 cm w.e. yr^{-1} . Within the overall ~~energy balance~~SEB, melt represents a relatively minor component (Fig. 9): the largest mean monthly contribution, in August, is well below 10 % of the total energy turnover, and in every month sublimation is a more effective energy sink than melt. Still, ~~in-refreezing at depth transfers heat deep into the snow pack (Fig. 6), compared to the slower processes of diffusion and advection which proceed from the surface. In~~ the NE domain region – where wind scouring is strongest – annual melt amounts correspond to a significant fraction of net accumulation (Fig. 3a). With regard to temporal patterns, the entire surface was found to always refreeze at night over the modeled period. ~~Moreover~~Also, no melt is simulated between November and March, with only minor amounts in April and October (Fig. 9).

375
380

Despite the large spatial heterogeneity, and an inter-annual variability exceeding 50 %, a common trend of melt increase could be detected in the annual time series (Fig. 10): the fitted slope ranges ~~between-from~~ (3 ± 2) and-to (4 ± 3) mm w.e. yr^{-2} across the saddle. While the trend is somewhat masked by the 2003 extreme melt year at the very beginning, it becomes statistically significant over the rest of the period.

385

The EBFM shows a marked tendency towards small melt amounts: frequency of modeled melt events decays exponentially with their magnitude (Fig. 11a), and a significant fraction of total melt amounts is contributed by micro-melt events under 4 mm w.e. in a single day (Fig. 11b).

390 Investigation of the weather conditions leading to melt occurrence (Fig. 12) reveals the ~~relationship~~relationships between weather variables and surface melt. Air temperature provides a critical control over melt rates ~~and total amounts~~, unlike cloud cover which appears to have almost no effect (Fig. 12a/b-). The majority of melt happens under clear-sky conditions ~~and~~ and slightly positive temperatures (Fig. 12b); still, significant amounts are simulated between -5 and 0 °C ~~(Fig. 12b): at SP, SK~~

and ZS melt at sub-freezing air temperatures accounts for respectively 22, 34 and 17 % of the total simulated amounts. Non-zero (though minimal) melt amounts are modeled down to air temperatures of -7°C , under clear skies and moderate to high humidity. Conversely, in very dry conditions (Fig. 12c/d) sublimation losses hinder melt even at slightly positive temperatures. Finally, wind speed appears to have a minor effect on long-term mean melt rates, slightly decreasing the likelihood of melt under high winds (Fig. 12e/f).

5 Discussion

At CG, the repeated, long-term investigations enable interpretation of the model output against a rich literature background. The following sections evaluate and put into context the model results for firn temperatures, meltwater infiltration and refreezing, as well as melt amounts.

5.1 Firn temperatures

The EBFM shows considerable potential at simulating cold firn. In addition to reproducing individual borehole profiles, the model confirms broader patterns such as the strong firn temperature gradient towards the Grenzgletscher slopes (Suter and Hoelzle, 2002) and the depth of zero annual temperature oscillation, at about 20 m (Hoelzle et al., 2011). This last observation indicates a realistic simulation of heat conduction within the firn pack.

The spatial distribution of model aspect-dependent spatial pattern of temperature residuals (Fig. 5) could be affected by the lack in the modeled SEB of radiation reflected from the surrounding terrain. This process is expected to induce a net energy transfer from the more sun-exposed cells towards the more shaded ones, which could partially re-balance the aspect dependence of model residuals. Several factors. The locations of largest temperature under-estimation coincide with a positive bias of modeled density, by 5-30 % compared to measured core profiles below 2 m depth. This density bias leads to a higher thermal conductivity (Eq. 21), resulting in colder firn temperatures as shown in Appendix B. The density bias could be due to a lack of local calibration for the accumulation-dependent densification model, developed over Antarctic firn (Ligtenberg et al., 2011). Another possibility is a deep density increase caused by the percolation routine. Specifically, after each melt event water is distributed between the surface and the depth of z_{lim} and there it refreezes, increasing density over that entire vertical extent – even in case of repeated melt-freeze cycles (which would melt a same ice surface, not increasing density). We expect this effect to be especially significant at locations where melt amounts represent a large fraction of accumulation: as such, it would be amplified by the accumulation model, which computes precipitation amounts already corrected for losses from wind scouring. Indeed, accumulation at CG results from summer precipitation events (Sect. 1), but modeled precipitation is distributed more evenly throughout the year, as it is based on weather station measurements from lower elevations (Sect. 3.3). Thus, we expect an under-estimation of the strong seasonal gradient that favours summer accumulation, so that modeled melt and refreeze can temporarily approach (or even locally exceed) the low accumulated snow amounts in summer, hence exacerbating the density bias. Simple sensitivity tests support this interpretation, by showing that the simulation of temperature profiles at low-accumulation borehole CG05-1 (Fig. 4 and 5) improves substantially with an artificial increase of precipitation amounts.

~~In addition, A deficit in summer precipitation totals in the model (with a corresponding winter excess) could also introduce a systematic deviation in the advected heat, due to a different deposition temperature of snow between summer and winter. The resulting firn temperature bias would be roughly proportional to mean accumulation (Appendix B): thus, a more extreme accumulation seasonality (with no winter precipitation) could potentially amplify the spatial pattern of modeled temperature residuals (Fig. 5). Still, complete removal of winter precipitation would be problematic for the albedo decay routine (Eq. 5). A more realistic model of wind scouring should account for the timing of snow erosion, which at CG can happen several months after deposition (e.g., Alean et al., 1983).~~

Some boreholes with positive model bias ~~are often~~ (Fig. 5) are simulated with too strong near-surface temperature gradients, leading to sharp positive deviations in the profile near the surface (e.g., Fig. 4f/k/p/q). Such a behavior suggests too deep refreezing is occurring in ~~the~~ ~~their~~ simulation, which could ~~again~~ be linked to the parametrized preferential infiltration routine (Sect. 3.4). Indeed, below a depth of 0–4 m (where refreezing is occurring) the simulation appears to be unbiased at most locations. ~~The vertical distribution of refreezing is also affected by residual saturation θ_{mi} , controlling water storage after melt events and potentially affecting firn temperatures. In practice, we show in Appendix B that sensitivity to this parameter is very low in our setup, and limited to high-melt locations.~~

~~Model under-estimation of firn temperatures appears to correlate with the lowest mean annual accumulation values. The spatial pattern of model residuals (Fig. 3a and 5). At these locations wind scouring is strongest, and melt amounts represent a significant fraction of net annual accumulation. Simple sensitivity tests showed that the simulation of profiles at borehole location CG05-1 improves substantially by forcing increased precipitation amounts. This suggests that the accumulation model – which computes precipitation amounts already corrected for losses from wind scouring – 5) could in principle be affected by the lack in the modeled SEB of SW radiation reflected from the surrounding terrain. This process could induce a net energy transfer from the more sun-exposed cells towards the more shaded ones. We quantified its magnitude applying a simple Lambert reflection model (e.g., Koppal, 2014) to the modeled series of SW radiation reflected by each grid cell. We found that radiation interception can indeed introduce a mean energy flux of up to about 3 W m^{-2} in the SEB, but the net effect (taking into account mutual radiation exchange and the high surface albedo) is always smaller than $\pm 0.2 \text{ W m}^{-2}$. As such, redistribution of reflected radiation cannot be considered a major contributor to the aspect-dependent model temperature residuals. Another effect which is not captured by our SEB formulation is the penetration of SW radiation within the snow pack. This process – may be problematic whenever melt totals approach the value of net accumulation. Indeed, accumulation at CG results from summer precipitation events (Sect. 1), but modeled precipitation is distributed more evenly throughout the year, as it is based on weather station measurements (Sect. 3.3). Thus we expect modeled precipitation to be under-estimated in summer compared to the actual accumulation. As such, in summer modeled melt can temporarily approach (or even locally exceed) the low accumulated snow amounts. Then the parametrized percolation would generate dense, thick firn layers: their limited refreezing capacity could reduce the heat release at depth during subsequent melt events, inducing under-estimation of the englacial temperatures enabling sub-surface melt – challenges the EBFM assumption of water originating entirely at the snow surface, and was found to attenuate a cold bias in firn temperatures within the model of Gilbert et al. (2014b) at CdD.~~

460 The relative spatial distribution of modeled firn temperatures (Fig. 7a) is in good agreement with the interpolated result shown by Suter and Hoelzle (2002). Features such as the regular temperature gradient across the CG saddle and the near-temperate south-facing slope below 4300 m a.s.l. are reproduced well. The high spatial variability of firn temperature trends (Fig. 7b) is consistent with the borehole observations of Hoelzle et al. (2011) and the model results of Gilbert et al. (2014a) at CdD. Below 4400 m a.s.l., firn warming rates are generally high on west-facing slopes and low on south-facing ones: this could be related
465 to a prevalent role of solar radiation over air temperature in the firn thermal regime at south-facing locations. As warming progresses, it is possible that such pronounced spatial patterns migrate also towards the higher saddle region, where warming rates are at present more uniform. In the context of increasing firn temperatures and melt amounts, the large spatial variability of warming trends will likely have a growing importance for the localization of future ice core drilling campaigns.

Among past modeling efforts at CG, both Lüthi and Funk (2001) and Suter (2002) formulated independent predictions for
470 firn temperatures evolution by 2020. As the target time frame for verification is reached, a major limitation of their modeled scenarios is found in the expected magnitude of atmospheric warming: while they assumed linear air temperature increases by respectively 0.4 and 0.45 °C between 2000 and 2020, the fitted trend at over the CM AWS annual means (series in Fig. 2) amounts to 0.05 (0.05 ± 0.03) °C yr⁻²⁻¹, corresponding to a much stronger warming of 1 °C over the period. For the CG saddle region, the two studies predict firn temperature increases at 18 m by 0.42 and 1.06 °C respectively, between 2000 and 2020.
475 By comparison, the EBFM simulates 0.70 °C of warming between 2003 and 2018 at SP, which can be uniformly rescaled over 2000–2020 to 0.88 °C. Such a firn warming is consistent with the results of the two studies, but lower when compared to the assumed-measured air temperature increase. A contributing factor could be the EBFM spin-up: temperatures were initialized with repeated model runs over 2004–2011, thus the initial grid at all depths is in equilibrium with the mean forcing over that period. By contrast, an adjustment time of 2–4 years is to be expected at the considered depth (Hoelzle et al., 2011). Therefore
480 at the beginning of the simulation the EBFM may be slightly over-estimating deep firn temperatures (Fig. 4a/b), resulting in a lower trend for 2003–2018.

At the CdD site, Gilbert et al. (2014a) estimated firn warming by 2030 and 2050 using a thermo-mechanical coupled model, forced by three climate projections within the A1B emission scenario (Nakićenović, 2000). They found a dependence of the warming rate on advection velocities and percolation amounts. Their reported firn warming at 20 m depth is in the range 0.0-1.8
485 °C over 2010-2030 (depending on location and climate scenario), similar to our results modeled for 2000-2020 over the CG domain (Fig. 7b).

5.2 Meltwater infiltration

A key parameter used for model set-up is the percolation depth z_{lim} , which defines the maximum depth reached by infiltrated meltwater through preferential flow. In cold firn such a parameter is crucial: simulated firn temperatures (Fig. 6) indicate that
490 all meltwater can be expected to refreeze not far from the initial location (controlled by the parametrized vertical distribution: Sect. 3.4). Thus z_{lim} effectively determines not only the initial meltwater distribution, but also the depth of refreezing-latent heat release. The effect of this parameter on firn temperatures is indeed large at all depths (Appendix B). The calibrated value of 4 m, together with the Gaussian vertical distribution, correspond-corresponds to a mean preferential percolation depth of 1.06

m. Unfortunately, in situ quantitative measurements of infiltration depths are scarce and mostly indirect. Evidence from winter snow packs and glacier accumulation areas shows that percolation and refreezing are strongly dependent on the meltwater supply amounts and rates (Marchenko et al., 2017), the temperature of the firn matrix (Koerner, 1970) and its stratigraphy (Illangasekare et al., 1990), notably affected by the previous history of infiltration and refreezing. An ~~obvious~~ often observed consequence is the increase of percolation depths over the melting season (Marchenko et al., 2017). At CG, Alean et al. (1983) observed slight melting but no meltwater percolation during a warm spell in summer 1981. Suter (2002) attempted to directly track meltwater refreezing by continuously logging a temperature profile at Seserjoch in 1999, but the setup failed before the onset of summer melt. Hoelzle et al. (2011) reported evidence for increasing infiltration depths, indicating (at Seserjoch and on the Grenzletscher slopes) a transition to a percolation regime spanning several annual firn layers. ~~By contrast~~ On the cold firm of CdD, Gilbert et al. (2014b) tracked sub-surface temperatures over the summer of 2012, measuring percolation depths of up to 4-5 m: these would match the value of z_{lim} in our simulation. However, the stratigraphy of core *unifr-2019* (Fig. A1) revealed dry firn layers interspersed with thin ice crusts, suggesting small amounts of infiltration and refreezing, except for a thick, ice-rich layer at 4.5 m depth. Therefore in our simulation we consider z_{lim} more as a tuning parameter than a realistic percolation depth.

Stratigraphy of the firn core points to a shortcoming of the subsurface model: meltwater distributed in cold firn along the parametrized vertical profile usually refreezes in place, producing a diluted density increase in the simulation. Instead, distinct ice layers are known to form at depth after preferential percolation, affecting the mechanical, hydrological and thermodynamical properties of the snow pack (Quéno et al., 2020). In the present EBFM formulation, such ice layers would not appear even with a very fine model grid. Refreezing after parametrized infiltration also distributes heat instantly over a fixed, large vertical extent, resulting in unrealistic, frequent warm pulses at depth after each melt event – no matter how small (inset in Fig. 6). These observations ~~indicate the need for~~ provide motivation for future work on a physically based percolation routine in the EBFM, accounting for the time evolution of z_{lim} and its dependence on snow density and stratigraphy.

5.3 Melt amounts

Modeled melt amounts can be compared to the amount of refrozen ice observed in core *unifr-2019* (Fig. A1). According to the computed climatology (Fig. 3a), the core location has a mean long-term accumulation of about 50 cm w.e. yr^{-1} : thus the 5.5 m core should span an estimated period of about 5–6 years (with a fairly large uncertainty due to the inter-annual accumulation variability). The model predicts about 85–100 cm w.e. of melt over such a time span (Fig. 8); ~~while~~ Instead, the core was found to contain only 31 cm of ice layers. These correspond to just 14 cm w.e. of refrozen ice after subtraction of the mean density of the ice-free sections (such a correction is fairly uncertain due to the high density variability of the profile). Some observations can be brought forward to put the apparent discrepancy into context.

First, refrozen ice amounts recorded in the core are affected by repeated cycles of melt-refreeze. Indeed, the very small amounts of meltwater produced during less intense (but rather frequent: Fig. 11) melt events can be expected to refreeze in the very first snow centimeters. Then any subsequent melt occurring before the next snowfall would affect the same ice surface, contributing to total melt amounts but without significant increases in ice layer thickness. Such surface crusts of relatively

impermeable ice have already been observed at CG (e.g., Lier, 2018). Since melt mostly happens in clustered patterns (almost only in summer and within a specific set of weather conditions: Fig. 9 and 12), contribution of repeated melt-refreeze cycles could potentially be very large.

530 ~~With~~ Because ice-equivalent thicknesses of daily melt amounts ~~often close to are often of the same order as~~ the size of single crystals (Fig. 11a), it is suggested that detection of some refrozen forms would require a resolution not achieved during our field analysis. This is supported by the observation in the core of sections of icy firn, as opposed to well-defined ice layers (see Appendix A). Investigations of such minimal melt processes are very limited in the literature. Still, Das and Alley (2005) 535 performed hot-box experiments on the formation of thin refreeze layers in Antarctic snow, finding that a profile resolution of 1 mm was necessary to capture small-scale melt processes. Moreover, in their experiments the wetted and refrozen snow next to melt layers did not show any type of melt feature detectable in firn core stratigraphy (Das and Alley, 2005), possibly enabling some refrozen layers to remain undetected. The possibility of an overlooked vertical ice gland embedded in our core also cannot be ruled out (measured core sections were not broken up after analysis). In fact, the presence of undetected ice 540 would be consistent with the high density variability encountered in the core profile (Fig. A1). Still, this observation could also be linked to wind compaction, and its interplay with wind erosion exposing older, denser snow.

~~An Lier (2018) proposed an estimation of refreezing amounts from profile density anomalies was proposed by Lier (2018) using several new and archived cores in the CG area. The method involved sizable uncertainties due to a poorly constrained surface density value at CG from the measured density anomalies over ideal dry densification profiles.~~ At the *Sattelkern* and 545 *Zumsteinkern* cores (respectively ~~at the saddle point and on the south-facing slope~~), ~~estimated refreezing rates are respectively in the close to SP and to ZS) the reported refreezing rates have confidence intervals of~~ 1–13 and 3–33 cm ice yr⁻¹ ~~ranges~~ (Lier, 2018). The EBFM predicts approximately 19 and 25 cm ice yr⁻¹ of melt. On the shaded Signalkuppe flank, values span the range 0–15 cm ice yr⁻¹ from multiple cores; the EBFM result over the same region is between 5 and 12 cm ice yr⁻¹ of melt. Considering the very strong inter-annual variability of melt amounts (Fig. 10), the results are largely compatible.

550 The same density anomaly approach had been applied at CdD by Gilbert et al. (2014b) to estimate melt amounts of the 2011 summer season. Based on 14 firn density profiles (measured between 4230 and 4310 m a.s.l.) the authors computed melt amounts in the range 1-18 cm w.e., significantly correlated with potential incoming solar radiation. At both CG and CdD the density anomaly method shows considerable potential and consistent results, but does not account for the occurrence of repeated melt-refreeze cycles. Moreover, at CG the method involves sizable uncertainties reflecting a high variability of surface density (Lier, 2018).

The significant melt amounts modeled at sub-freezing 2 m air temperatures suggest reconsideration of degree-day models for simulating melt at high-alpine (and possibly high-latitude) locations. This is consistent with the findings of Suter et al. (2004) at Seserjoch, who observed several surface melt events but no days with positive mean temperatures over the whole 1999 summer. Indeed, laboratory experiments by Beck et al. (1988) revealed the occurrence of melt already at -4 °C with 560 475 W m⁻² of incoming SW radiation. At the CM AWS, values in excess of 1000 W m⁻² are a common summer occurrence ; ~~supporting~~ (195 hours per year on average in the 2003–2018 series). This supports the plausibility of melt under even colder conditions (Fig. 12). From a theoretical perspective, Kuhn (1987) analytically explored a standard SEB equation (neglecting

sub-surface heat conduction), in relation to common weather situations on an alpine glacier. The conclusion was that melt onset ~~likely happens can likely happen~~ at air temperatures between -10 and +10 °C; ~~very much comparable to our results: our~~ results appear to corroborate such a range (Fig. 12).

6 Conclusions and outlook

This work ~~marked-marks~~ a first effort to apply a coupled, high-resolution distributed EBFM to alpine cold firn, within a multi-year simulation forced with extensively processed meteorological data, acquired at high altitude and in closest vicinity of the study site. In both cold and near-temperate conditions, the model can achieve promising results for firn temperatures, ~~comparable in accuracy to the spatial variability of the measurements with an average RMS error below 1.5 °C within the~~ depth of annual temperature oscillations. Therefore the EBFM can be deemed suitable for further investigations of ~~future cold firn evolution~~ the present thermal regime as well as future temperatures evolution at cold firn sites, based on localized climate scenarios.

At CG, our results corroborate earlier observations on the spatial patterns of surface melt and ~~the deep firn temperatures.~~ For the first time we provide a spatial estimation of distributed firn warming at the site, showing a large variability over small distances: modeled trends of deep ~~temperatures. firn temperature range from no change to 2.5 times the atmospheric warming rate. We also report a novel trend of increasing surface melt amounts, currently close to statistical significance despite high inter-annual variability and the brevity of the time series. These observations confirm the potential for accelerated changes in the thermal regime and firn facies of the site (Hoelzle et al., 2011). Further developments should be closely monitored, since~~ in the near future they could affect the suitability of CG for retrieving climate records from ice cores.

A previously unreported feature in melt dynamics is the occurrence of micro-melt events, with daily amounts below 4 mm w.e.: these remain difficult to detect, but our analysis hints at a possibly significant impact on melt totals and firn temperatures, hence on calibration and ground-truthing of model results. More field observations are needed to verify the occurrence and improve the understanding of such events, ~~also assessing and assess~~ their potential effect on subsequent water infiltration.

Our model results point to significant melt happening at negative 2 m air temperatures, confirming earlier field observations: this would re-affirm the importance of using a full energy-balance model over a parametrized melt approach in cold conditions. Additional field investigations of the meteorological conditions at the onset of melt would be a valuable development.

~~Moreover, we found a novel trend of increasing surface melt amounts, currently on the edge of statistical significance. Further evolution of this tendency should be closely monitored, since in the near future it could affect the site suitability for~~ retrieving climate records from ice cores. Estimations of local melt amounts ~~may also and warming rates may~~ contribute to the localization of future core drilling efforts: this provides motivation to attempt model deployment at other cold firn/ice sites and potentially on a larger scale.

~~For such an~~ For this application, it ~~would be beneficial is important~~ to better constrain patterns and depths of meltwater percolation, further refining the model percolation scheme by implementing a physical infiltration routine ~~and acquiring vital~~ calibration data: this would allow to overcome the limitations of a fixed-depth parametrized percolation, which presently

carries a strong impact on simulated firn temperatures. The acquisition of site-specific calibration data will be vital to support this advancement. In addition to thermal tracking, recent developments in non-destructive analysis methods (e.g., Heilig et al., 2018; Katsushima et al., 2020) could advantageously serve this purpose.

600 *Code and data availability.* The EBFM code used in this study is available at <https://doi.org/10.5281/zenodo.4327090>. Due to their large volume, modeled grids are available on request. The meteorological time series and digital elevation models should be requested from the respective providers. Full information on our processing of meteorological and topographic data (described in Mattea, 2020) is also available on simple request, including the respective software code.

Appendix A: Core *unifr-2019*

605 The 5.5 m firn core (Fig. A1) was recovered with a manually operated Kovacs Mark II corer. Firn density was measured in 20 cm sections using a digital scale, while stratigraphy was visually inspected at a resolution of 0.5 cm. Mean core density is 474 kg m⁻³, with a high variability and no clear densification trend towards depth. Relatively dense snow (up to 600 kg m⁻³) was encountered near the surface at around 0.5 m, with no ice layers concurrently observed. The first shallowest traces of refreezing were found at a depth of about 2 m. In total, 31 cm of refrozen layers could be identified, typically less than 2 cm thick and with variable ice content. The missing 20 cm section at 3.8 m depth may have included additional ice layers. At 4.5 m several 610 ice-rich layers mixed with icy firn were found, over a contiguous thickness of 23 cm.

Appendix B: Sensitivity experiments

We investigated model sensitivity to several surface and sub-surface parameters by testing single parameter perturbations, each including the respective model spin-up as in Sect. 3.5. For performance reasons we focused on firn temperature deviations at the three points marked in Fig. 1a, comparing the perturbed model output to the baseline shown throughout the paper (Fig. B1).

615

For surface roughness length z_0 , we tested values of 0.1 mm and 10 mm, corresponding to the extreme ends of the range reported by Brock et al. (2006) over snow surfaces on mid-latitude glaciers. Our simulation is moderately sensitive to the value of z_0 , with deep temperature deviations between -1.6 and +1.2 °C (Fig. B1a/b); these changes have no clear dependence on melt amounts or accumulation rates. Firn temperatures tend to decrease with $z_0 = 10$ mm, due to a strong increase in sublimation rates and a decrease in melt amounts. Conversely, the low value of $z_0 = 0.1$ mm extends the melting season from March to October, a result not supported by field evidence at CG. Overall, the observed sensitivity provides motivation to test further refinements of the EBFM turbulent fluxes routines. These could include a time dependence of roughness length, which in snow can span more than one order of magnitude over a single season (Brock et al., 2006).

620 The rain/snow temperature threshold $T_{s/r}$ was calibrated by van Pelt et al. (2019) against mass balance measurements in Svalbard between 1967 and 2015. The influence at CG is very small (Fig. B1c/d), due to the rarity of precipitation events at

positive air temperatures: precipitation above 0 (1) °C accounts for just 1.2 (0.7) % of the total 2003-2018 amount. Such a minor role could potentially become more significant in the future, as rainfall amounts increase amidst rising air temperatures. Thus, local calibration of parameter $T_{s/r}$ could be important for an investigation of future scenarios.

630 Within the sub-surface model, we tested sensitivity to residual saturation θ_{mi} and percolation depth z_{lim} . Residual saturation appears to have almost no effect on firn temperatures, except at the high-melt ZS location (Fig. B1e/f). This is likely due to the small meltwater amounts being distributed by the percolation routine over a considerable vertical extent, within sub-freezing snow and firn (Fig. 6); thus, all water can refreeze in place and residual saturation does not play an important role. Even at ZS, temperature deviations are within ± 0.25 °C after halving or doubling the parameter value. As such, θ_{mi} is not the most essential parameter for calibration in the present CG setup. Still, scenarios including higher amounts of meltwater production
635 (Fig. 10) could be more affected by its value.

By contrast, the percolation depth parameter z_{lim} provides an important control on firn temperatures, proportional to melt amounts and relatively uniform across depths (Fig. B1g/h). Compared to the tuned value of 4 m, restricting preferential percolation to the first 2 m can reduce firn temperatures by as much as 4 °C at the ZS location, as a larger fraction of released latent heat can escape towards the surface (Fig. B1g).

640 To examine the thermal effect of summer precipitation under-estimation (Sect. 5.1), we experimented with an alternative seasonal precipitation cycle as model forcing. The complete removal of winter precipitation – to reproduce the effect of winter wind scouring – is problematic for albedo decay in the model (Eq. 5): thus, we tested an opposite change, consisting of a 50 % reduction of precipitation in May-October, redistributed over the other months to preserve the annual totals. The resulting changes in firn temperatures are strongly anti-correlated (coefficient of -0.99) with mean annual accumulation (Fig. B1i). Thus (except for albedo decay) a more pronounced accumulation seasonality in the model would likely increase firn
645 temperatures proportionally to the mean accumulation rates. In the present setup this change could amplify the spatial pattern of firn temperature biases (Fig. 5).

Finally, we tested the recent parametrization of snow/firn thermal conductivity proposed by Calonne et al. (2019), to cover within one formula the full range of densities and temperatures found on glaciers. In the density range of interest at CG,
650 conductivity is increased by about 20-50 % compared to the formula of Sturm et al. (1997). As a result, deep firn temperatures decrease by 1-2.5 °C, with some more differences in the seasonal cycle at shallower depths (Fig. B1j). Two factors contribute to the cooling. Melt amounts decrease by about 10 % because the higher conductivity delays the onset of melt, through a larger heat loss towards the glacier when the SEB approaches melting conditions. Also, modeled temperatures (up to about 2 m depth) are on average colder (by up to 3-4 °C) than deep temperatures: thus, a higher conductivity shifts the deep equilibrium
655 temperature towards colder values. The Calonne et al. (2019) parametrization of thermal conductivity will be included in an upcoming release of the EBFM.

Author contributions. EM performed the analysis and wrote the paper. HM and EM drilled the firn core. MH provided unpublished data and previous works. WvP supplied the EBFM model code and support to use it. MK contributed to the deployment of the model. MB provided the main meteorological data-set and extensive clarification on the CM AWS. All authors participated to the discussion of the results.

660 *Competing interests.* The authors declare that they have no conflict of interest.

Acknowledgements. We would like to thank Carlo Licciulli and Josef Lier (Heidelberg University) for supplying core, borehole and radar data. We would also like to thank ARPA Piemonte, the Italian Air Force Weather Service, MeteoSwiss, PERMOS and Visit Monte Rosa for providing access to the respective meteorological data. This work contains data/products of the Italian Air Force Weather Service. Swisstopo and Regione Piemonte provided the digital elevation models. Past firn temperature measurements were performed within the GLAMOS project, with funding by the Federal Office for the Environment FOEN, MeteoSwiss and the Swiss Academy of Sciences SCNAT. The legacy data collected in the Monte Rosa region was mainly organized and measured within the PhD thesis of Dr. Stephan Suter and was funded by the European Union Environment and Climate Programme under ENV4-CT97-0639 and the Swiss Government under BBW Nr. 97.0349-1, within the framework of the ALPCLIM EU project (Environmental and Climate records from high-elevation Alpine glaciers). MK, MH and HM acknowledge support by the Swiss National Science Foundation SNSF (grant 200021_169453). [WvP acknowledges funds from the Swedish National Space Agency \(project 189/18\)](#). This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (project acronym CASSANDRA, grant agreement No. 818994).

665
670

References

- Alean, J., Haeberli, W., and Schädler, B.: Snow accumulation, firn temperature and solar radiation in the area of the Colle Gnifetti core drilling site (Monte Rosa, Swiss Alps): distribution patterns and interrelationships, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 19, 131–147, 1983.
- ARPA Piemonte: Dati meteo orari, [https://www.arpa.piemonte.it/rischinaturali/accesso-ai-dati/Richieste-dati-formato-standard/richiesta-dati/Richiesta-automatica/Dati-meteo-orari.html?delta=1&\\$SCADENZA\\$=2](https://www.arpa.piemonte.it/rischinaturali/accesso-ai-dati/Richieste-dati-formato-standard/richiesta-dati/Richiesta-automatica/Dati-meteo-orari.html?delta=1&$SCADENZA$=2), last access December 2020, 2020.
- Arthern, R. J., Vaughan, D. G., Rankin, A. M., Mulvaney, R., and Thomas, E. R.: In situ measurements of Antarctic snow compaction compared with predictions of models, *J. Geophys. Res.*, 115, F03 011, <https://doi.org/10.1029/2009JF001306>, <http://doi.wiley.com/10.1029/2009JF001306>, 2010.
- Auer, I., Böhm, R., and Schöner, W.: Chapter 3: Instrumental Climate, in: Final report of EU-rtd-project ALPCLIM, Zentralanstalt für Meteorologie und Geodynamik, Vienna, Austria, <http://www.zamg.ac.at/histalp/download/abstract/Auer-etal-2001c-F.pdf>, 2001.
- Barbante, C., Schwikowski, M., Döring, T., Gäggeler, H. W., Schotterer, U., Tobler, L., Van de Velde, K., Ferrari, C., Cozzi, G., Turetta, A., Rosman, K., Bolshov, M., Capodaglio, G., Cescon, P., and Boutron, C.: Historical Record of European Emissions of Heavy Metals to the Atmosphere Since the 1650s from Alpine Snow/Ice Cores Drilled near Monte Rosa, *Environmental Science & Technology*, 38, 4085–4090, <https://doi.org/10.1021/es049759r>, <https://pubs.acs.org/doi/10.1021/es049759r>, 2004.
- Beck, N., Wagenbach, D., and Münnich, K. O.: Laboratory experiments on the formation of solar radiation induced melt layers in dry snow, *Zeitschrift für Gletscherkunde und Glazialgeologie*, 24, 31–40, 1988.
- Bohleber, P., Erhardt, T., Spaulding, N., Hoffmann, H., Fischer, H., and Mayewski, P.: Temperature and mineral dust variability recorded in two low-accumulation Alpine ice cores over the last millennium, *Climate of the Past*, 14, 21–37, <https://doi.org/10.5194/cp-14-21-2018>, <https://www.clim-past.net/14/21/2018/>, 2018.
- Bollmeyer, C., Keller, J. D., Ohlwein, C., Wahl, S., Crewell, S., Friederichs, P., Hense, A., Keune, J., Kneifel, S., Pscheidt, I., Redl, S., and Steinke, S.: Towards a high-resolution regional reanalysis for the European CORDEX domain: High-Resolution Regional Reanalysis for the European CORDEX Domain, *Quarterly Journal of the Royal Meteorological Society*, 141, 1–15, <https://doi.org/10.1002/qj.2486>, <http://doi.wiley.com/10.1002/qj.2486>, 2015.
- Bougamont, M., Bamber, J. L., and Greuell, W.: A surface mass balance model for the Greenland Ice Sheet, *Journal of Geophysical Research: Earth Surface*, 110, n/a–n/a, <https://doi.org/10.1029/2005JF000348>, <http://doi.wiley.com/10.1029/2005JF000348>, 2005.
- Brock, B. W., Willis, I. C., and Sharp, M. J.: Measurement and parameterization of aerodynamic roughness length variations at Haut Glacier d’Arolla, Switzerland, *J. Glaciol.*, 52, 281–297, <https://doi.org/10.3189/172756506781828746>, https://www.cambridge.org/core/product/identifier/S0022143000208605/type/journal_article, 2006.
- Buri, P.: Simulation of cold-firn-temperatures at an Alpine site using the model GEOtop, Master’s thesis, University of Zürich, https://uzb.swisscovery.slsp.ch/view/delivery/41SLSP_UZB/12464773910005508, 2013.
- Calonne, N., Milliancourt, L., Burr, A., Philip, A., Martin, C. L., Flin, F., and Geindreau, C.: Thermal Conductivity of Snow, Firn, and Porous Ice From 3-D Image-Based Computations, *Geophysical Research Letters*, 46, 11, <https://doi.org/10.1029/2019GL085228>, 2019.
- Cannon, A. J., Sobie, S. R., and Murdock, T. Q.: Bias Correction of GCM Precipitation by Quantile Mapping: How Well Do Methods Preserve Changes in Quantiles and Extremes?, *Journal of Climate*, 28, 6938–6959, <https://doi.org/10.1175/JCLI-D-14-00754.1>, <http://journals.ametsoc.org/doi/10.1175/JCLI-D-14-00754.1>, 2015.

- Darms, G. A.: Firntemperaturen auf dem Colle Gnifetti: Zusammenstellung und Analyse bestehender und neuer Temperaturprofile, Master's thesis, University of Zürich, 2009.
- 710 Das, S. B. and Alley, R. B.: Characterization and formation of melt layers in polar snow: observations and experiments from West Antarctica, *Journal of Glaciology*, 51, 307–312, <https://doi.org/10.3189/172756505781829395>, https://www.cambridge.org/core/product/identifier/S0022143000214937/type/journal_article, 2005.
- de Blasi, F., Gabrieli, J., Zhuravlev, A., Urbini, S., Burgay, F., Lodi, R., Dallo, F., Spolaor, A., Colucci, R. R., and Barbante, C.: Mt. Grand Combin ice core: correspondence among meteorological, morphological and geophysical evidence with ice core analysis, in: *Geophysical Research Abstracts*, vol. 21, Vienna, Austria, <https://meetingorganizer.copernicus.org/EGU2019/EGU2019-14393-2.pdf>, 2019.
- 715 Essery, R. and Etchevers, P.: Parameter sensitivity in simulations of snowmelt, *Journal of Geophysical Research*, 109, <https://doi.org/10.1029/2004JD005036>, <http://doi.wiley.com/10.1029/2004JD005036>, 2004.
- Feigenwinter, I., Kotlarski, S., Casanueva, A., Fischer, A., Schwierz, C., and Liniger, M.: Exploring quantile mapping as a tool to produce user-tailored climate scenarios for Switzerland, *Tech. Rep. 270*, MeteoSwiss, 2018.
- 720 Frank, C. W., Wahl, S., Keller, J. D., Pospichal, B., Hense, A., and Crewell, S.: Bias correction of a novel European reanalysis data set for solar energy applications, *Solar Energy*, 164, 12–24, <https://doi.org/10.1016/j.solener.2018.02.012>, <https://linkinghub.elsevier.com/retrieve/pii/S0038092X18301294>, 2018.
- Gabrieli, J., Cozzi, G., Vallelonga, P., Schwikowski, M., Sigl, M., Eickenberg, J., Wacker, L., Boutron, C., Gäggeler, H., Cescon, P., and Barbante, C.: Contamination of Alpine snow and ice at Colle Gnifetti, Swiss/Italian Alps, from nuclear weapons tests, *Atmospheric Environment*, 45, 587–593, <https://doi.org/10.1016/j.atmosenv.2010.10.039>, <https://linkinghub.elsevier.com/retrieve/pii/S1352231010009258>, 2011.
- 725 Gabrielli, P., Carturan, L., Gabrieli, J., Dinale, R., Krainer, K., Hausmann, H., Davis, M., Zagorodnov, V., Seppi, R., Barbante, C., Dalla Fontana, G., and Thompson, L.: Atmospheric warming threatens the untapped glacial archive of Ortles mountain, South Tyrol, *J. Glaciol.*, 56, 843–853, <https://doi.org/10.3189/002214310794457263>, https://www.cambridge.org/core/product/identifier/S0022143000214111/type/journal_article, 2010.
- 730 Gilbert, A. and Vincent, C.: Atmospheric temperature changes over the 20th century at very high elevations in the European Alps from englacial temperatures: EUROPEAN ALPS AIR TEMPERATURE CHANGES, *Geophys. Res. Lett.*, 40, 2102–2108, <https://doi.org/10.1002/grl.50401>, <http://doi.wiley.com/10.1002/grl.50401>, 2013.
- Gilbert, A., Wagnon, P., Vincent, C., Ginot, P., and Funk, M.: Atmospheric warming at a high-elevation tropical site revealed by englacial temperatures at Illimani, Bolivia (6340 m above sea level, 16°S, 67°W), *J. Geophys. Res.*, 115, D10 109, <https://doi.org/10.1029/2009JD012961>, <http://doi.wiley.com/10.1029/2009JD012961>, 2010.
- 735 Gilbert, A., Gagliardini, O., Vincent, C., and Wagnon, P.: A 3-D thermal regime model suitable for cold accumulation zones of polythermal mountain glaciers: A 3-D thermal regime model for glaciers, *Journal of Geophysical Research: Earth Surface*, 119, 1876–1893, <https://doi.org/10.1002/2014JF003199>, <http://doi.wiley.com/10.1002/2014JF003199>, 2014a.
- 740 Gilbert, A., Vincent, C., Six, D., Wagnon, P., Piard, L., and Ginot, P.: Modeling near-surface firn temperature in a cold accumulation zone (Col du Dôme, French Alps): from a physical to a semi-parameterized approach, *The Cryosphere*, 8, 689–703, <https://doi.org/10.5194/tc-8-689-2014>, <https://www.the-cryosphere.net/8/689/2014/>, 2014b.
- Gilbert, A., Vincent, C., Gagliardini, O., Krug, J., and Berthier, E.: Assessment of thermal change in cold avalanching glaciers in relation to climate warming, *Geophys. Res. Lett.*, 42, 6382–6390, <https://doi.org/10.1002/2015GL064838>, <http://doi.wiley.com/10.1002/2015GL064838>, 2015.
- 745

- GLAMOS: The Swiss Glaciers 2013/14 and 2014/15 Glaciological Report No. 135/136, Tech. rep., Cryospheric Commission (EKK) of the Swiss Academy of Sciences (SCNAT); Laboratory of Hydraulics, Hydrology and Glaciology (VAW), Swiss Federal Institute of Technology Zurich (ETH Zurich), https://doi.org/10.18752/GLREP_135-136, 2017.
- 750 Greuell, W. and Konzelmann, T.: Numerical modelling of the energy balance and the englacial temperature of the Greenland Ice Sheet. Calculations for the ETH-Camp location (West Greenland, 1155 m a.s.l.), *Global and Planetary Change*, 9, 91–114, [https://doi.org/10.1016/0921-8181\(94\)90010-8](https://doi.org/10.1016/0921-8181(94)90010-8), <https://linkinghub.elsevier.com/retrieve/pii/0921818194900108>, 1994.
- Greuell, W., Knap, W. H., and Smeets, P. C.: Elevational changes in meteorological variables along a midlatitude glacier during summer, *Journal of Geophysical Research: Atmospheres*, 102, 25 941–25 954, <https://doi.org/10.1029/97JD02083>, <http://doi.wiley.com/10.1029/97JD02083>, 1997.
- 755 Gruber, S., King, L., Kohl, T., Herz, T., Haerberli, W., and Hoelzle, M.: Interpretation of geothermal profiles perturbed by topography: the alpine permafrost boreholes at Stockhorn Plateau, Switzerland, *Permafrost and Periglacial Processes*, 15, 349–357, <https://doi.org/10.1002/ppp.503>, <http://doi.wiley.com/10.1002/ppp.503>, 2004.
- Haerberli, W. and Beniston, M.: Climate Change and Its Impacts on Glaciers and Permafrost in the Alps, *Ambio*, 27, 258–265, <http://www.jstor.org/stable/4314732>, 1998.
- 760 Haerberli, W. and Funk, M.: Borehole temperatures at the Colle Gnifetti core-drilling site (Monte Rosa, Swiss Alps), *Journal of Glaciology*, 37, 37–46, <https://doi.org/10.3189/S0022143000042775>, https://www.cambridge.org/core/product/identifier/S0022143000042775/type/journal_article, 1991.
- Harper, J., Humphrey, N., Pfeffer, W. T., Brown, J., and Fettweis, X.: Greenland ice-sheet contribution to sea-level rise buffered by meltwater storage in firn, *Nature*, 491, 240–243, <https://doi.org/10.1038/nature11566>, <http://www.nature.com/articles/nature11566>, 2012.
- 765 Heilig, A., Eisen, O., MacFerrin, M., Tedesco, M., and Fettweis, X.: Seasonal monitoring of melt and accumulation within the deep percolation zone of the Greenland Ice Sheet and comparison with simulations of regional climate modeling, *The Cryosphere*, 12, 1851–1866, <https://doi.org/10.5194/tc-12-1851-2018>, <https://www.the-cryosphere.net/12/1851/2018/>, 2018.
- Hoelzle, M., Darms, G., Lüthi, M. P., and Suter, S.: Evidence of accelerated englacial warming in the Monte Rosa area, Switzerland/Italy, *The Cryosphere*, 5, 231–243, <https://doi.org/10.5194/tc-5-231-2011>, <http://www.the-cryosphere.net/5/231/2011/>, 2011.
- 770 Illangasekare, T. H., Walter, R. J., Meier, M. F., and Pfeffer, W. T.: Modeling of meltwater infiltration in subfreezing snow, *Water Resources Research*, 26, 1001–1012, <https://doi.org/10.1029/WR026i005p01001>, <http://doi.wiley.com/10.1029/WR026i005p01001>, 1990.
- Jenk, T. M., Szidat, S., Schwikowski, M., Gäggeler, H. W., Brüttsch, S., Wacker, L., Synal, H.-A., and Saurer, M.: Radiocarbon analysis in an Alpine ice core: record of anthropogenic and biogenic contributions to carbonaceous aerosols in the past (1650–1940), *Atmos. Chem. Phys.*, 6, 5381–5390, <https://doi.org/10.5194/acp-6-5381-2006>, <https://acp.copernicus.org/articles/6/5381/2006/>, 2006.
- 775 Jenk, T. M., Szidat, S., Boliuss, D., Sigl, M., Gäggeler, H. W., Wacker, L., Ruff, M., Barbante, C., Boutron, C. F., and Schwikowski, M.: A novel radiocarbon dating technique applied to an ice core from the Alps indicating late Pleistocene ages, *Journal of Geophysical Research*, 114, <https://doi.org/10.1029/2009JD011860>, <http://doi.wiley.com/10.1029/2009JD011860>, 2009.
- Katsushima, T., Adachi, S., Yamaguchi, S., Ozeki, T., and Kumakura, T.: Nondestructive three-dimensional observations of flow finger and lateral flow development in dry snow using magnetic resonance imaging, *Cold Regions Science and Technology*, 170, 102 956, <https://doi.org/10.1016/j.coldregions.2019.102956>, <https://linkinghub.elsevier.com/retrieve/pii/S0165232X19301119>, 2020.
- 780 Klok, E. and Oerlemans, J.: Model study of the spatial distribution of the energy and mass balance of Morteratschgletscher, Switzerland, *Journal of Glaciology*, 48, 505–518, <https://doi.org/10.3189/172756502781831133>, https://www.cambridge.org/core/product/identifier/S0022143000209623/type/journal_article, 2002.

- Koerner, R.: The Mass Balance of the Devon Island Ice Cap, Northwest Territories, Canada, 1961-66, *Journal of Glaciology*, 9, 325–336, <https://doi.org/10.3189/S0022143000022863>, https://www.cambridge.org/core/product/identifier/S0022143000022863/type/journal_article, 1970.
- Konrad, H., Bohleber, P., Wagenbach, D., Vincent, C., and Eisen, O.: Determining the age distribution of Colle Gnifetti, Monte Rosa, Swiss Alps, by combining ice cores, ground-penetrating radar and a simple flow model, *Journal of Glaciology*, 59, 179–189, <https://doi.org/10.3189/2013JoG12J072>, https://www.cambridge.org/core/product/identifier/S00221430000203511/type/journal_article, 2013.
- Konzelmann, T., Vandewal, R., Greuell, W., Bintanja, R., Henneken, E., and Abeouchi, A.: Parameterization of global and longwave incoming radiation for the Greenland Ice Sheet, *Global and Planetary Change*, 9, 143–164, [https://doi.org/10.1016/0921-8181\(94\)90013-2](https://doi.org/10.1016/0921-8181(94)90013-2), <https://linkinghub.elsevier.com/retrieve/pii/0921818194900132>, 1994.
- Koppal, S. J.: Lambertian Reflectance, pp. 441–443, Springer US, Boston, MA, https://doi.org/10.1007/978-0-387-31439-6_534, 2014.
- Kuhn, M.: Micro-Meteorological Conditions for Snow Melt, *Journal of Glaciology*, 33, 24–26, <https://doi.org/10.3189/S002214300000530X>, https://www.cambridge.org/core/product/identifier/S002214300000530X/type/journal_article, 1987.
- Kuipers Munneke, P., M. Ligtenberg, S. R., van den Broeke, M. R., van Angelen, J. H., and Forster, R. R.: Explaining the presence of perennial liquid water bodies in the firn of the Greenland Ice Sheet, *Geophys. Res. Lett.*, 41, 476–483, <https://doi.org/10.1002/2013GL058389>, <http://doi.wiley.com/10.1002/2013GL058389>, 2014.
- Legrand, M., Preunkert, S., May, B., Guilhermet, J., Hoffman, H., and Wagenbach, D.: Major 20th century changes of the content and chemical speciation of organic carbon archived in Alpine ice cores: Implications for the long-term change of organic aerosol over Europe, *J. Geophys. Res. Atmos.*, 118, 3879–3890, <https://doi.org/10.1002/jgrd.50202>, <http://doi.wiley.com/10.1002/jgrd.50202>, 2013.
- Licciulli, C.: Full Stokes ice-flow modeling of the high-Alpine glacier saddle Colle Gnifetti, Monte Rosa: Flow field characterization for an improved interpretation of the ice-core records, Ph.D. thesis, University of Heidelberg, 2018.
- Lier, J.: Estimating the amount of latent heat released by refreezing surface melt water for the high-Alpine glacier saddle Colle Gnifetti, Swiss/Italian Alps, Master's thesis, University of Heidelberg, 2018.
- Ligtenberg, S. R. M., Helsen, M. M., and van den Broeke, M. R.: An improved semi-empirical model for the densification of Antarctic firn, *The Cryosphere*, 5, 809–819, <https://doi.org/10.5194/tc-5-809-2011>, <https://tc.copernicus.org/articles/5/809/2011/>, 2011.
- Lüthi, D., Le Floch, M., Bereiter, B., Blunier, T., Barnola, J.-M., Siegenthaler, U., Raynaud, D., Jouzel, J., Fischer, H., Kawamura, K., and Stocker, T. F.: High-resolution carbon dioxide concentration record 650,000–800,000 years before present, *Nature*, 453, 379–382, <https://doi.org/10.1038/nature06949>, <http://www.nature.com/articles/nature06949>, 2008.
- Lüthi, M. P.: Rheology of cold firn and dynamics of a polythermal ice stream: Studies on Colle Gnifetti and Jakobshavns Isbræ, Ph.D. thesis, ETH Zürich, <https://doi.org/10.3929/ethz-a-003884174>, <http://hdl.handle.net/20.500.11850/144542>, 2000.
- Lüthi, M. P. and Funk, M.: Modelling heat flow in a cold, high-altitude glacier: interpretation of measurements from Colle Gnifetti, Swiss Alps, *Journal of Glaciology*, 47, 314–324, <https://doi.org/10.3189/172756501781832223>, https://www.cambridge.org/core/product/identifier/S0022143000211027/type/journal_article, 2001.
- Marchenko, S., van Pelt, W. J. J., Claremar, B., Pohjola, V., Pettersson, R., Machguth, H., and Reijmer, C.: Parameterizing Deep Water Percolation Improves Subsurface Temperature Simulations by a Multilayer Firn Model, *Frontiers in Earth Science*, 5, <https://doi.org/10.3389/feart.2017.00016>, <http://journal.frontiersin.org/article/10.3389/feart.2017.00016/full>, 2017.
- Martorina, S., Olivero, A., Loglisci, N., and Pelosini, R.: La stazione meteo più alta d'Europa, Neve e Valanghe, 49, <https://issuu.com/aineva7/docs/nv49>, 2003.

- Masson-Delmotte, V., Dreyfus, G., Braconnot, P., Johnsen, S., Jouzel, J., Kageyama, M., Landais, A., Loutre, M.-F., Nouet, J., Parrenin, F., Raynaud, D., Stenni, B., and Tuerter, E.: Past temperature reconstructions from deep ice cores: relevance for future climate change, *Clim. Past*, 2, 145–165, <https://doi.org/10.5194/cp-2-145-2006>, <https://cp.copernicus.org/articles/2/145/2006/>, 2006.
- 825 Mattea, E.: Measuring and modelling changes in the firm at Colle Gnifetti, 4400 m a.s.l., Swiss Alps, Master's thesis, University of Fribourg, [https://bigweb.unifr.ch/Science/Geosciences/GeographyTechnical/Secretary/Pub/Publications/Geography/SelectedBachelorMasterThesis/2020/Mattea_E._\(2020\)_M_Measuring_modelling_changes_Colle_Gnifetti.pdf](https://bigweb.unifr.ch/Science/Geosciences/GeographyTechnical/Secretary/Pub/Publications/Geography/SelectedBachelorMasterThesis/2020/Mattea_E._(2020)_M_Measuring_modelling_changes_Colle_Gnifetti.pdf), 2020.
- MeteoAM: Servizio Meteorologico dell'Aeronautica Militare - Disponibilità dei dati, http://www.meteoam.it/dati_in_tempo_reale, last access December 2020, 2020.
- 830 MeteoSwiss: Data portal for experts, <https://www.meteoswiss.admin.ch/home/services-and-publications/beratung-und-service/datenportal-fuer-experten.html>, last access December 2020, 2020.
- More, A. F., Spaulding, N. E., Bohleber, P., Handley, M. J., Hoffmann, H., Korotkikh, E. V., Kurbatov, A. V., Loveluck, C. P., Sneed, S. B., McCormick, M., and Mayewski, P. A.: Next-generation ice core technology reveals true minimum natural levels of lead (Pb) in the atmosphere: Insights from the Black Death, *GeoHealth*, 1, 211–219, <https://doi.org/10.1002/2017GH000064>, <https://onlinelibrary.wiley.com/doi/abs/10.1002/2017GH000064>, 2017.
- 835 Nakićenović, N., ed.: Special report on emissions scenarios: a special report of Working Group III of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge ; New York, oCLC: ocm44652561, 2000.
- New, M., Hulme, M., and Jones, P.: Representing Twentieth-Century Space–Time Climate Variability. Part II: Development of 1901–96 Monthly Grids of Terrestrial Surface Climate, *Journal of Climate*, 13, 2217–2238, [https://doi.org/10.1175/1520-0442\(2000\)013<2217:RTCSTC>2.0.CO;2](https://doi.org/10.1175/1520-0442(2000)013<2217:RTCSTC>2.0.CO;2), <http://journals.ametsoc.org/doi/abs/10.1175/1520-0442%282000%29013%3C2217%3ARTCSTC%3E2.0.CO%3B2>, 2000.
- 840 Oerlemans, J. and Grisogono, B.: Glacier winds and parameterisation of the related surface heat fluxes, *Tellus A*, 54, 440–452, <https://doi.org/10.1034/j.1600-0870.2002.201398.x>, <http://tellusa.net/index.php/tellusa/article/view/12164>, 2002.
- Oerlemans, J. and Knap, W. H.: A 1 year record of global radiation and albedo in the ablation zone of Morteratschgletscher, Switzerland, *Journal of Glaciology*, 44, 231–238, <https://doi.org/10.1017/S0022143000002574>, https://www.cambridge.org/core/product/identifier/S0022143000002574/type/journal_article, 1998.
- 845 Preunkert, S., Wagenbach, D., Legrand, M., and Vincent, C.: Col du Dôme (Mt Blanc Massif, French Alps) suitability for ice-core studies in relation with past atmospheric chemistry over Europe, *Tellus B: Chemical and Physical Meteorology*, 52, 993–1012, <https://doi.org/10.3402/tellusb.v52i3.17081>, <https://www.tandfonline.com/doi/full/10.3402/tellusb.v52i3.17081>, 2000.
- 850 Preunkert, S., Legrand, M., and Wagenbach, D.: Sulfate trends in a Col du Dôme (French Alps) ice core: A record of anthropogenic sulfate levels in the European midtroposphere over the twentieth century, *J. Geophys. Res.*, 106, 31 991–32 004, <https://doi.org/10.1029/2001JD000792>, <http://doi.wiley.com/10.1029/2001JD000792>, 2001.
- Quéno, L., Fierz, C., van Herwijnen, A., Longridge, D., and Wever, N.: Deep ice layer formation in an alpine snowpack: monitoring and modeling, *The Cryosphere*, 14, 3449–3464, <https://doi.org/10.5194/tc-14-3449-2020>, <https://tc.copernicus.org/articles/14/3449/2020/>, 2020.
- 855 Regione Piemonte: RIPRESA AEREA ICE 2009-2011 – DTM 5, <http://www.geoportale.piemonte.it/geonetworkrp/srv/ita/metadata.show?id=2552&currTab=rndt>, last access November 2020, 2011.
- Reijmer, C. H., van Meijgaard, E., and van den Broeke, M. R.: Evaluation of temperature and wind over Antarctica in a Regional Atmospheric Climate Model using 1 year of automatic weather station data and upper air observations, *Journal of Geophysical Research*, 110, <https://doi.org/10.1029/2004JD005234>, <http://doi.wiley.com/10.1029/2004JD005234>, 2005.

- 860 Rossi, G., Johnston, P., and Maggi, V.: ALPCLIM project: Reconstruction of the monthly values of solar radiation incident over the Lys Glacier surface (Monte Rosa-Western Italian Alps), in: 26th Intl. Conference on Alpine Meteorology, ICAM, Innsbruck, 2000a.
- Rossi, G., Johnston, P., and Maggi, V.: Project ALPCLIM: Résultats de l'observation météorologique dans le site de Colle du Lys (4250 mètres), in: Réunion Annuelle Société Hydrotechnique de France, Section Glaciologie, Société Hydrotechnique de France, Grenoble, 2000b.
- 865 Schneider, T. and Jansson, P.: Internal accumulation in firn and its significance for the mass balance of Storglaciären, Sweden, *J. Glaciol.*, 50, 25–34, <https://doi.org/10.3189/172756504781830277>, https://www.cambridge.org/core/product/identifier/S0022143000213634/type/journal_article, 2004.
- Schwikowski, M.: Reconstruction of European Air Pollution from Alpine Ice Cores, in: *Earth Paleoenvironments: Records Preserved in Mid- and Low-Latitude Glaciers*, edited by DeWayne Cecil, L., Green, J. R., and Thompson, L. G., vol. 9, pp. 95–119, Kluwer Academic Publishers, Dordrecht, https://doi.org/10.1007/1-4020-2146-1_6, http://link.springer.com/10.1007/1-4020-2146-1_6, series Title: *Developments in Paleoenvironmental Research*, 2004.
- 870 Shumskii, P. A.: *Principles of Structural Glaciology: The Petrography of Fresh-water Ice as a Method of Glaciological Investigation*, Dover Publications Inc., New York, 1964.
- Sturm, M., Holmgren, J., König, M., and Morris, K.: The thermal conductivity of seasonal snow, *Journal of Glaciology*, 43, 26–
- 875 41, <https://doi.org/10.3189/S0022143000002781>, https://www.cambridge.org/core/product/identifier/S0022143000002781/type/journal_article, 1997.
- Suter, S.: *Cold firn and ice in the Monte Rosa and Mont Blanc areas: spatial occurrence, surface energy balance and climatic evidence*, Ph.D. thesis, ETH Zürich, <https://doi.org/10.3929/ethz-a-004288434>, <http://hdl.handle.net/20.500.11850/145885>, 2002.
- Suter, S. and Hoelzle, M.: Cold firn in the Mont Blanc and Monte Rosa areas, *European Alps: spatial distribution and statistical models*, *Annals of Glaciology*, 35, 9–18, <https://doi.org/10.3189/172756402781817059>, https://www.cambridge.org/core/product/identifier/S0260305500257200/type/journal_article, 2002.
- 880 Suter, S., Laternser, M., Haeberli, W., Frauenfelder, R., and Hoelzle, M.: Cold firn and ice of high-altitude glaciers in the Alps: measurements and distribution modelling, *Journal of Glaciology*, 47, 85–96, <https://doi.org/10.3189/172756501781832566>, https://www.cambridge.org/core/product/identifier/S0022143000212264/type/journal_article, 2001.
- 885 Suter, S., Hoelzle, M., and Ohmura, A.: Energy balance at a cold Alpine firn saddle, Seserjoch, Monte Rosa, *International Journal of Climatology*, 24, 1423–1442, <https://doi.org/10.1002/joc.1079>, <http://doi.wiley.com/10.1002/joc.1079>, 2004.
- Thevenon, F., Anselmetti, F. S., Bernasconi, S. M., and Schwikowski, M.: Mineral dust and elemental black carbon records from an Alpine ice core (Colle Gnifetti glacier) over the last millennium, *J. Geophys. Res.*, 114, D17 102, <https://doi.org/10.1029/2008JD011490>, <http://doi.wiley.com/10.1029/2008JD011490>, 2009.
- 890 van Pelt, W., Pohjola, V., Pettersson, R., Marchenko, S., Kohler, J., Luks, B., Hagen, J. O., Schuler, T. V., Dunse, T., Noël, B., and Reijmer, C.: A long-term dataset of climatic mass balance, snow conditions, and runoff in Svalbard (1957–2018), *The Cryosphere*, 13, 2259–2280, <https://doi.org/10.5194/tc-13-2259-2019>, <https://www.the-cryosphere.net/13/2259/2019/>, 2019.
- van Pelt, W. J. and Kohler, J.: Modelling the long-term mass balance and firn evolution of glaciers around Kongsfjorden, Svalbard, *Journal of Glaciology*, 61, 731–744, <https://doi.org/10.3189/2015JG14J223>, https://www.cambridge.org/core/product/identifier/S0022143000202542/type/journal_article, 2015.
- 895

- van Pelt, W. J., Pettersson, R., Pohjola, V. A., Marchenko, S., Claremar, B., and Oerlemans, J.: Inverse estimation of snow accumulation along a radar transect on Nordenskiöldbreen, Svalbard, *Journal of Geophysical Research: Earth Surface*, 119, 816–835, <https://doi.org/10.1002/2013JF003040>, <http://doi.wiley.com/10.1002/2013JF003040>, 2014.
- van Pelt, W. J. J., Oerlemans, J., Reijmer, C. H., Pohjola, V. A., Pettersson, R., and van Angelen, J. H.: Simulating melt, runoff and refreezing on Nordenskiöldbreen, Svalbard, using a coupled snow and energy balance model, *The Cryosphere*, 6, 641–659, <https://doi.org/10.5194/tc-6-641-2012>, <https://www.the-cryosphere.net/6/641/2012/>, 2012.
- Vandecrux, B., Mottram, R., Langen, P. L., Fausto, R. S., Olesen, M., Stevens, C. M., Verjans, V., Leeson, A., Ligtenberg, S., Kuipers Munneke, P., Marchenko, S., van Pelt, W., Meyer, C. R., Simonsen, S. B., Heilig, A., Samimi, S., Marshall, S., Machguth, H., MacFerrin, M., Niwano, M., Miller, O., Voss, C. I., and Box, J. E.: The firn meltwater Retention Model Intercomparison Project (RetMIP): evaluation of nine firn models at four weather station sites on the Greenland ice sheet, *The Cryosphere*, 14, 3785–3810, <https://doi.org/10.5194/tc-14-3785-2020>, <https://tc.copernicus.org/articles/14/3785/2020/>, 2020.
- Vincent, C., Le Meur, E., Six, D., Possenti, P., Lefebvre, E., and Funk, M.: Climate warming revealed by englacial temperatures at Col du Dôme (4250 m, Mont Blanc area), *Geophys. Res. Lett.*, 34, <https://doi.org/10.1029/2007GL029933>, <http://doi.wiley.com/10.1029/2007GL029933>, 2007.
- 910 Vincent, C., Gilbert, A., Jourdain, B., Piard, L., Ginot, P., Mikhalenko, V., Possenti, P., Le Meur, E., Laarman, O., and Six, D.: Strong changes in englacial temperatures despite insignificant changes in ice thickness at Dôme du Gôuter glacier (Mont Blanc area), *The Cryosphere*, 14, 925–934, <https://doi.org/10.5194/tc-14-925-2020>, <https://www.the-cryosphere.net/14/925/2020/>, 2020.
- Visit Monte Rosa: Stazioni meteo Monte Rosa Val d’Aosta e Piemonte, <https://www.visitmonterosa.com/stazioni-meteo/>, last access December 2020, 2020.
- 915 Wagenbach, D., Bohleber, P., and Preunkert, S.: Cold, alpine ice bodies revisited: what may we learn from their impurity and isotope content?, *Geografiska Annaler: Series A, Physical Geography*, 94, 245–263, <https://doi.org/10.1111/j.1468-0459.2012.00461.x>, <https://www.tandfonline.com/doi/full/10.1111/j.1468-0459.2012.00461.x>, 2012.
- Wahl, S., Bollmeyer, C., Crewell, S., Figura, C., Friederichs, P., Hense, A., Keller, J. D., and Ohlwein, C.: A novel convective-scale regional reanalysis COSMO-REA2: Improving the representation of precipitation, *Meteorologische Zeitschrift*, 26, 345–361, <https://doi.org/10.1127/metz/2017/0824>, http://www.schweizerbart.de/papers/metz/detail/26/87333/A_novel_convective_scale_regional_reanalysis_COSMO?af=crossref, 2017.
- 920 Wolff, E., Barbante, C., Becagli, S., Bigler, M., Boutron, C., Castellano, E., de Angelis, M., Federer, U., Fischer, H., Fundel, F., Hansson, M., Hutterli, M., Jonsell, U., Karlin, T., Kaufmann, P., Lambert, F., Littot, G., Mulvaney, R., Röthlisberger, R., Ruth, U., Severi, M., Siggaard-Andersen, M., Sime, L., Steffensen, J., Stocker, T., Traversi, R., Twarloh, B., Udisti, R., Wagenbach, D., and Wegner, A.: Changes in environment over the last 800,000 years from chemical analysis of the EPICA Dome C ice core, *Quaternary Science Reviews*, 29, 285–295, <https://doi.org/10.1016/j.quascirev.2009.06.013>, <https://linkinghub.elsevier.com/retrieve/pii/S027737910900208X>, 2010.
- Yen, Y.-C.: Review of thermal properties of snow, ice and sea ice, CRREL report 81-10, DTIC, 1981.




Figure 1. (a) [Location map of the Colle Gnifetti location map area](#). Labeled model cells [across the saddle](#) correspond to the point series in Fig. 6–10. **ZS:** Zumsteinspitze slope; **CG:** Colle Gnifetti saddle point; **SK:** Signalkuppe slope. Several borehole locations were measured more than once (also see Fig. 5). (b) Overview map of the Monte Rosa range with AWS locations (see Table 2 for more information). [The red square shows the extent of Fig. 1a](#). In both panels X and Y are metric CH1903/LV03 coordinates. Orthophoto and topographic map source: Federal Office of Topography swisstopo.




Figure 2. Quality-checked and gap-filled hourly weather series used as model input (reference elevation 4560 m a.s.l.). Black lines represent monthly means. For readability, only monthly means are shown for the precipitation down-scaling coefficient (Sect. 3.3). The reduced variability in temperature lapse rates in late 2016 is due to a 4 months data gap, where the series was entirely reconstructed from lower altitude stations. Pie charts show the contribution of the original CM AWS series to the model input. For cloudiness, the contribution is based on values reconstructed from SW radiation measurements. "Discarded" includes values rejected during quality check as well as night-time SW radiation measurements; "missing" includes data gaps and parameters not recorded at CM. All such values were replaced with the reconstructed series.




Figure 3. (a) Long-term net annual accumulation climatology at CG, serving as spatial component for the distributed accumulation model. Dense point sequences in the north-east were derived from GPR profiles. X and Y are metric CH1903/LV03 coordinates. Topographic map source: Federal Office of Topography swisstopo. (b) Linear fit of annual snow mass anomaly at core KCC versus median annual wind speed at the CM AWS.




Figure 4. Measured and modeled borehole temperature profiles, sorted chronologically. Axes range is the same in all plots. The bottom depth of 20 m corresponds to the deepest simulated values and to the depth of zero amplitude of the annual cycle. Profile codes follow the scheme XXYY-Z/WW, with XX location code, YY borehole year, Z borehole number, and WW year of the profile measurement (Hoelzle et al., 2011; GLAMOS, 2017). Letter codes correspond to Fig. 5.




Figure 5. Map of mean-depth-averaged residuals of modeled borehole temperature profiles (20 m grid, 1 h time-step). Letter codes correspond to Fig. 4. X and Y are metric CH1903/LV03 coordinates. Topographic map source: Federal Office of Topography swisstopo.




Figure 6. Time-depth plots of modeled firn temperatures at (a) EGSP, (b) SK and (c) ZS (map in Fig. 1a). 100 m grid, 3 h time-step. Inset shows 2003 summer temperatures down to 3.5 m depth. The annual cycle becomes smaller than 0.1 °C at approximately 15 m depth, and further decreases beyond the model quantization noise (due to the vertical layer discretization) by 20 m.




Figure 7. (a) Modeled 20 m firn temperatures on 31 December 2018. (b) Modeled 20 m firn temperature trends over 2003-2018. In both panels, marked cells correspond to the representative points of Fig. 1a. X and Y are metric CH1903/LV03 coordinates. Topographic map source: Federal Office of Topography swisstopo.




Figure 8. Mean (2003-2018) modeled annual melt amounts (20 m grid, 1 h time-step). Marked cells correspond to the representative points of Fig. 1a. X and Y are metric CH1903/LV03 coordinates. Topographic map source: Federal Office of Topography swisstopo.




Figure 9. Mean (2003-2018) monthly distribution of modeled energy balance components at (a) SK and (b) ZS. 20 m grid, 1 h time-step. Distribution of fluxes at [EG-SP](#) (not shown) is intermediate between the two.




Figure 10. Time series of modeled annual melt amounts at three representative grid cells (solid lines; **EGSP**: saddle point, SK: Signalkuppe slope, ZS: Zumsteinspitze slope). Linear least-squares fits (each including $n = 16$ annual values) are shown as dashed lines. Reported annual melt trends and corresponding p-values are computed both on the whole 2003–2018 modeling period (first number), and excluding the extreme melt year of 2003 (second number, in parentheses). 20 m grid resolution, 1 h time-step.




Figure 11. (a) Distribution of magnitudes of modeled melt events (defined as total daily melt amounts). Y-axis is logarithmic. (b) Cumulative melt amounts sorted by the magnitude of contributing melt events. Melt amount for each event is averaged over the whole domain of Fig. 8. 20 m grid resolution, 1 h time-step.




Figure 12. Mean melt rates (left) and cumulative melt amounts (right) modeled over 2003-2018 at the [CG-saddle-point-SP location](#) (20 m grid / 1 h time-step), sorted according to the respective weather conditions: air temperature and **(a-b)** cloud cover, **(c-d)** relative humidity, **(e-f)** wind speed. Tiles of weather conditions not encountered in the input series are not drawn.




Figure A1. Density and stratigraphy of firn core *unifr-2019*. Values between 3.7 and 3.9 m are missing due to broken core during recovery. Drilling location is in metric CH1903/LV03 coordinates.




Figure B1. Modeled firn temperature deviations ΔT_f (2003-2018 means) by month, depth and location, after single parameter perturbations. (a)-(b): roughness length z_0 from 1 mm to 10 mm and to 0.1 mm. (c)-(d): rain/snow temperature threshold $T_{s/r}$ from 0.6 °C to 1.2 °C and to 0.0 °C. (e)-(f): residual saturation θ_{mi} scaled by a factor 0.5 and by a factor 2 compared to Eq. 24. (g)-(h): percolation depth parameter z_{lim} from 4 m to 2 m and to 6 m. (i): precipitation amounts reduced by 50 % in May-October and redistributed over the rest of the year. (j): thermal conductivity parametrization changed from Sturm et al. (1997) to Calonne et al. (2019). The vertical scale is the same within each row. Locations are shown in Fig. 1a.