

# High-resolution inventory to capture glacier disintegration in the Austrian Silvretta

Andrea Fischer<sup>1</sup>, Gabriele Schwaizer<sup>2</sup>, Bernd Seiser<sup>1</sup>, Kay Helfricht<sup>1</sup>, Martin Stocker-Waldhuber<sup>1</sup>

5 <sup>1</sup> Institute for Interdisciplinary Mountain Research, Austrian Academy of Sciences, Innsbruck, Austria

<sup>2</sup> ENVEO GmbH, Fürstenweg 176, 6020 Innsbruck, Austria

*Correspondence to:* Andrea Fischer (andrea.fischer@oeaw.ac.at)

10 **Abstract.** The rapidly increasing decay of the glaciers in the Austrian Silvretta was captured for the years 2017 and 2018 by a new high-resolution glacier inventory. Identifying the glacier outlines offers a wide range of possible interpretations of glaciers that have evolved into small and now totally debris-covered cryogenic structures. In previous inventories, a high proportion of active bare ice allowed a clear delineation of the glacier margins even by optical imagery. In contrast, in the current state of the glacier only the patterns and amounts of volume change allow us to estimate the area of the buried glacier

15 remnants. Glacier outlines were mapped manually, based on LiDAR elevation models and patterns of volume change at 1 to 0.5 m spatial resolution. The vertical accuracy of the DEMs generated from 6 to 8 LiDAR points per m<sup>2</sup> is in the order of centimetres. Between 2004/06 and 2017/2018, the 46 glaciers of the Austrian Silvretta lost  $-29\pm 4\%$  of their area and now cover  $13.1\pm 0.4$  km<sup>2</sup>. This is only  $32\pm 2\%$  of their LIA extent of  $40.9\pm 4.1$  km<sup>2</sup>. The area change rate increased from  $-0.6\%/year$  (1969-2002) to  $-2.4\%/year$  (2004/06-2017/18). The Sentinel-2-based glacier inventory of 2018 deviates by just

20  $1\%$  of the area. The annual geodetic mass balance referring to the area at the beginning of the period showed a loss increasing from  $-0.2\pm 0.1$  m w.e./year (1969-2002) to  $-0.8$  m  $\pm 0.1$  w.e./year (2004/06-2017/18) with an interim peak in 2002-2004/06 of  $-1.5\pm 0.7$  m w.e./year. To keep track of the buried ice and its fate, and to distinguish increasing debris cover from ice loss, we recommend inventory repeat frequencies of three to five years and surface elevation data with a spatial resolution of one metre.

25


# 1 Introduction

As a reaction to climate change, mountain glaciers all over the world are changing at an increasing pace (IPCC, 2019) and at unprecedented rates (Zemp et al., 2015), but with significant regional variability (IPCC, 2019). It is widely accepted that in some mountain ranges a significant part of today's glaciers may disappear within this century (e.g. Zemp et al., 2019a, Huss and Fischer, 2016a). A loss of almost all mountain glaciers by 2300 seems possible (Marzeion et al., 2012). For tackling climate change using glaciers as essential climate variables (Bojinski et al., 2014), the precise and detailed monitoring of glacier recession is essential (Zemp et al., 2019b). It can also reveal regionally different response times (Zekollari et al., 2020), various uncertainties (Huss et al., 2014) and serve as a basis for specific scenarios (e.g. Zekollari et al., 2019).

Glacier inventories are amongst the most valuable tools for glacier monitoring on a regional scale (Haeberli et al., 2007, Gärtner Roer et al., 2019), and for remote and large glaciers. After the pioneering World Glacier Inventory (WGI), compiled by WGMS and NSIDC (1999), several initiatives, such as GLIMS (Kargel et al., 2014, Racoviteanu et al., 2009), have published global glacier inventories like the Randolph glacier inventory (Pfeffer et al., 2014). International consortia like Globglacier work out guidelines for mapping glaciers, e.g. Paul et al. (2009). At the same time, smaller regional studies work on new methods (e.g. Paul et al., 2020) and have responded to regional phenomena and demands, for example, debris-covered glaciers (Nagai et al., 2016) or the large and often cloudy and snow-covered Patagonian glaciers (Meier et al., 2018). Regional inventories at high resolution can also serve as validation for large-scale semi-automatic remote sensing products.

Airborne LiDAR has been a valuable tool for glacier studies for nearly 20 years (e.g. Geist and Stötter, 2002, Pellikka and Rees, 2009). Acquisition technologies, processing and analysis have been significantly enhanced since the early years and now reach a few cm nominal vertical accuracy in flat areas. High point densities allow processing gridded elevation data with a spatial resolution of 0.5 m or even higher. Early scientific work included investigating the potential of (repeat) LiDAR data to map geomorphological processes (Höfle and Rutzinger, 2011) and glacier/rock glacier extent (Abermann et al. 2010). In 2004, federal authorities in Austria initiated the compilation of the first federal DEMs based on LiDAR, which were used to update the photogrammetric Austrian glacier inventories (Fischer et al., 2015 a,b). Now a repeat federal LiDAR DEM is available for several regions in Austria.

50



**Figure 1: Overview of the glaciers, ongoing and past mass balance and fluctuation time series in the Austrian part of the Silvretta range.**


In this article, we present a new LiDAR-based glacier inventory and the respective geodetic mass balance of the Austrian Silvretta (Figure 1), located in the federal states of Tyrol and Vorarlberg. This inventory presents the glacier area and surface elevation for the years 2017 (Vorarlberg) and 2018 (Tyrol). Here, for the first time, a regional glacier inventory was derived from two high-resolution LiDAR surveys for a period of beginning glacier downwaste indicated by the extreme melt during the summer of 2003. Repeat LiDAR has the potential to tackle volume loss caused by the melt of debris-covered ice (Abermann et al., 2009). The mass balance monitoring at Jamtalferner glacier indicates extremely negative mass balances since then (Fischer et al., 2016c) in all elevation zones. As a result, the glacier darkens and disintegrates, so that the delineation of glacier outlines becomes increasingly uncertain if based on visible imagery only. The evolution of the other glaciers in the region is similar, so that it was necessary to include the interpretation of volume changes to delineate the debris-covered glacier margins. The volume change was also used to estimate the geodetic mass balance of the 46 glaciers for the period 2004/2006 2017/2018 with two different reference areas, the area at the beginning of the period and the average area.

Heavy and continuous mass losses resulted in significant changes of glacier geomorphology and topology in the Austrian Silvretta. For example, in recent years the Schnapfenkuchl V and H glaciers (Figures 2, 3 and close-ups in Figures S1 and 4) cannot at first glance be identified even during a field survey, as bare ice is rarely visible. The geomorphological structure of the totally debris-covered surface is not dominated by ice dynamics, with crevasses, steep lateral moraines causing different

70 inclinations of glacier areas and adjacent surfaces or depressions at the ice margins, as could be expected for larger debris-covered valley glaciers.




Figure 2: Aerial overview of the Schnapfenkuchl glaciers V, M, and H in an aerial photograph of 21.08.2020 (photograph by Andrea Fischer). While Schnapfenkuchl M still is a classic glacier with exposed bare ice, on Schnapfenkuchl H and V bare ice is rarely exposed (see the close-ups in Figure S2).



75 Figure 3: Overview of the Schnapfenkuchl glaciers V, M, and H in an orthophoto of 2020 with the information available prior to the new glacier inventory: the glacier margins of the former glacier inventories LIA, 1969 and 2006 and rock glacier margins (Krainer

and Ribis, 2012). While the ice margin of Schnapfenkuchl M and Larainferner is clearly visible, the delineation of Schnapfenkuchl V and H glaciers from the orthophoto can only be guessed at.



80

**Figure 4:** Bare ice exposed at Schnapfenkuchl glacier V in the orthophoto of 2015 (red arrows), with stratigraphic layers (yellow arrows) indicating sedimentary ice or firn. Orthophotos: CC 4.0 <https://www.data.gv.at/katalog/en/dataset/orthofoto>

Schnapfenkuchl V and H glaciers were clearly identified as glaciers in past inventories: they showed exposed bare ice, accumulation areas, an englacial drainage system and crevasses as indicators for ice dynamics were present. In recent years,

85

traditional and evident properties of a classical glacier became hidden by debris (see Figure 5 for the evolution of Schnapfenkuchl H glacier). In the orthophotos of 1954, 1970, and 2002, Schnapfenkuchl H glacier can be clearly identified as glaciers, while in 2009, 2015 and 2020 we would hardly map Schnapfenkuchl V and H as glaciers compiling the first glacier inventory of the region (Figure S2 A-C).

90

The increase in debris cover is not restricted to the few small glaciers like Schnapfenkuchl H and V. It is a widespread phenomenon in the Austrian Silvretta, also evident from regular field surveys. For example, length change measurements at one of the largest glaciers of Austrian Silvretta, Litzner glacier (Figure 1) were abandoned because the identification of the glacier margin was hampered by the increase in debris cover at the glacier tongue. Figure S3 illustrates that this process took only about a decade, with bare ice clearly evident at the glacier tongue in the orthophoto of 2006, but not in 2012 and 2015, where only the former firn area is still free of debris. Moreover, the higher glaciers, like Verhupf glacier, are affected in a way which potentially impacts on the accuracy of glacier delineation: With melt rates of more than 1 m w.e. recorded at the Jamtalferner even in elevations above 3000 m (Fischer et al., 2016c) even the small glaciers which remained fairly unchanged in the last inventories (Abermann et al., 2009) were affected in a way which is relevant for compiling glacier inventories. One example of this process is the evolution of Schnapfenkuchl H glacier illustrated by a time series of orthophotos (Figure 5).

95

100

Many glaciers in the Austrian Silvretta were reported to switch from the moderate glacier retreat of the last 150 years to a rapid downwaste close to a fade out of the ice (Fischer et al., 2016a, Fischer et al., 2016c, example in Figure 5). As these processes related to the rapid recession of mountain glaciers, which have potentially not only occurred in the Austrian Silvretta during the last decade, but sooner or later will in other mountain regions also, we have to validate and elaborate our monitoring strategies to capture this process. This is important on a local and global level, as even small glaciers contribute to sea level rise (Bahr and Radic, 2012) and can be significant for local hydrological and hazard management. The precise monitoring of the transition to a glacier-free landscape is also important for the interpretation and dating of paleoglacial landforms, as studied in the Austrian Silvretta by Braumann et al., 2020.

105

The now very small and rapidly downwasting glaciers of the Austrian Silvretta are a perfect test site for analysing the potentials and limitations of repeat LiDAR as a high-resolution airborne remote-sensing method for monitoring glacier fade out in qualitative and quantitative terms.

110

By definition, the new glacier inventory aims at tackling the changes in area and volume changes of all glaciers in a specific region. This raises the research questions which of the potentially transient cryogenic structures (e.g. Figure 5, year 2020) should remain part of a glacier inventory, and what the effect of neglecting ice remnants on inventory data would be.


115

To answer the research questions related to the compilation of glaciers inventories in a stage of early deglaciation, this study presents

120

- I) a new glacier inventory based on two high-resolution LiDAR DEMs to identify and quantify glacier areas in 2017/18,
- II) the rate of area and volume changes since the last glacier inventories of 1850, 1969, 2002 and 2004/2006
- III) a discussion of the potential and limitations of repeat LiDAR to map glacier changes under current glacier states
- IV) a discussion of glacier inventory strategies to monitor transient glacier states under conditions of rapid glacier decay close to complete loss of ice.

125



**Figure 5: The evolution of Schnapfenkuchl glacier V in a time series of orthophotos from 1954 to 2020 illustrates that the evolution from a ‘classical glacier’ with crevasses as visible indicators of ice dynamics took just 20 years. In the orthophotos of 2009 and 2015, bare ice is still visible at the surface. Orthophotos: CC 4.0 <https://www.data.gv.at/katalog/en/dataset/orthofoto>**

130

## 2 Data

### 2.1 Orthophotos 2015/2018 and surface elevations for 2017/2018

Surface elevation data for 2017/18 were made available by the federal administrations of the provinces of Tyrol and Vorarlberg (see the locations of the glaciers in Figure 1). The orthophotos of Tyrol date from summer 2015 for the southern part and from 135 2018 for the northern part (data.tirol.gv.at, 2020). The orthophotos of Vorarlberg date from 2018 and are available via the WMS Service geoland.at (2020). Spatial resolution of the images is 0.2-0.5 m.

The airborne LiDAR Digital Elevation Models (DEMs) used for mapping glaciers in this study date from 2017 (Vorarlberg) and 2018 (Tyrol), with horizontal resolutions of 0.5x0.5 m (Vorarlberg) and 1x1 m (Tyrol).

The LiDAR DEMs of 2017 and 2018 were both coregistered to the national GIS Grid (BEV, 2011a) with elevation (BEV, 140 2011b) as a national standard procedure. The coregistration of the high-resolution full waveform LiDAR DEMs to the earlier LiDAR DEMs is not considered state of the art of LiDAR technology (Attwenger, personal communication).

For the LiDAR data of Vorarlberg, the Silvretta was covered by 76 flight stripes during 14 and 29 August 2017. Quality control was carried out with ORIENT-LIDAR, resulting in a  $\sigma_0$  of 0.028 m, a RMS for pass area residuals of 0.004 / 0.004 / 0.016 m in x/y/z and 0.056 / 0.056 / 0.031 m for all points (Würländer, 2019). In the final report for the Tyrolean part (Rieger,2019) 145 the uncertainty estimate of the LiDAR data processed with the OPALS software is estimated by the comparison to control areas. The resulting standard deviation of elevation at the control areas is 0.032 m. The control area located in the subsample we used for the study showed a standard deviation of 0.030 m. All control areas are located on stable ground outside glaciers.

### 2.2 Areas and surface elevations in previous glacier inventories

150 For the Austrian part of the Silvretta range, we compiled glacier area (A) changes to the LIA maximum from mapping moraines, for 1969, 1996 and 2002 from orthophotos, and for 2004 (Vorarlberg) and 2006 (Tyrol) from existing glacier inventories (Fischer et al., 2015 a,b). For all inventories apart from the LIA inventory, not only glacier areas but also glacier surface elevations are available. The glacier margins were delineated manually with an uncertainty of the resulting area ( $\sigma_A$ ) of  $\pm 1.5\%$  for glaciers larger than 1km<sup>2</sup> and  $\pm 5\%$  for smaller ones (Abermann et al. 2009).

155 The uncertainty of surface elevations on glaciers as quantified by Abermann et al. (2010) is less  $\pm 0.3$  m for the DEMs based on the LiDAR flights in 2004 and 2006. The federal administration of Tyrol quantifies the single point standard deviation as 0.03 m based on the analysis of pass areas (Federal Government of Tyrol, 2020). The federal administration of Vorarlberg



160 evaluated their DEM of 2004 with 58.597 object elevations and 49.483 terrestrially measured points of different types and found a mean elevation difference of -0.04 m and -0.05 m respectively. 95% of the points meet an accuracy of  $\pm 0.4$  m (Landesvermessungsamt Feldkirch, 2004).

### 2.3 Comparison of nominal relative uncertainties of all LiDAR data used in the study

165 LiDAR surveys using different instruments and intended point densities (Table 1) produce different representations of the infinitesimally accurate ‘real surface’, even without real surface changes over time. Based on achieved point densities, the gridding method and resolution may add misalignment of different DEMs of the same area. For this, LiDAR data is validated at defined reference areas. Nevertheless, there is a tradition in glaciological remote sensing and photogrammetry to cross-check the DEM accuracy for glacier-covered areas in potentially stable areas without surface changes.

170 **Table 1: Instrument, minimum point density and pixel size of the LiDAR campaigns used in this study for the calculation of volume changes.**

Federal region	Year of survey	LiDAR instrument	Minimum point density per m <sup>2</sup>	DEM cell size m
Tyrol	2006	ALTM 3100 and Gemini	0.25	1.0
Tyrol	2018	Riegl VQ 780	8.00	1.0
Vorarlberg	2004	ALTM 2050	2.50	1.0
Vorarlberg	2017	Riegl LMS-Q780	6.20	0.5

175 To estimate the final uncertainty with respect to changing point densities and methods applied for generating DEMs (Table 2), we analysed surface elevation changes ( $\Delta z$ ) not only at glaciers, but also at rock glaciers and for a buffer of 1000 m and between 1000 m and 2000 m around all glaciers and rock glaciers. Although these areas are only partly representative for glaciers in terms of slope (Figure 6) and roughness, we consider these numbers a very conservative estimate for the uncertainty of the  $\Delta z$  at glaciers. For the rough and changing rock glaciers, we found a mean  $\Delta z$  of -0.4 m with a standard deviation of 1.1 m. At the buffer, excluding the unstable paraglacial areas (between 1000 to 2000m), we found a mean elevation difference of  $0.0 \pm 0.6$  m. The standard deviation is twice the uncertainty found by Abermann for LiDAR DEMs. We took the standard deviation as error in  $\Delta z$  for further error propagation.


180

**Table 2: Comparison of the slope and elevation change (mean, standard deviation) between 2004/06 and 2017/18 for 3 subsamples.**

Subsamples	Slope			Elevation change ( $\Delta z$ )	
	area	mean	$\sigma$	mean	$\sigma$
	km <sup>2</sup>	°	°	m	m
rock glaciers	7.35	26	10	-0.4	1.1
buffer 1000 m	112.85	33	15	-0.1	0.9
buffer 1000 2000 m	28.92	27	15	0.0	0.6

185 Studies on the derivation of DEMs from LiDAR point clouds reveal that a slope steeper than about 40° potentially exhibits larger deviation from the ‘true’ surface (Sailer et al., 2014). Although the algorithm applied to convert point clouds to gridded data plays a major role for the representation of a specific surface (elevation and shape), the representation of the smooth glacier surfaces is a bit more resilient to low resolution than very rough geomorphological features. Sailer et al. (2014) claim that the cell size for analysing glacier changes could be even between 5 and 10 m.

190 Sailer et al. (2014) recommend cell sizes below 1 m for terrain steeper than 40°, which we rarely find on the glaciers of the Austrian Silvretta. There 90% of the glacier area presents slopes below 40° (Figure 6). In any case, the spatial resolution of the LiDAR DEMs analysed in this study fulfils the criteria above.




195 **Figure 6: Distribution of slopes for glaciers, rock glaciers and the two buffer areas used for validation. The buffer regions are steeper than the glaciers, so that the elevation difference in the stable buffer zones can be considered an upper limit for the uncertainty.**

### 3 Methods

#### 200 3.1 Compilation of the glacier inventory 2017/2018

The glacier outlines were mapped manually based on the high-resolution LiDAR shaded reliefs and volume changes generated from LiDAR DEMs (Abermann et al., 2009) with 1x1 m pixel size as shown in Figure 7 for Schnapfenkuchl M terminus. The shaded reliefs allow a estimate of the glacier outline by distinguishing the smooth glacier areas from the rougher peri- and paraglacial area. The characteristic patterns of elevation changes reaching their maximum at the glacier terminus help to include debris-covered glacier areas as well.  
205



210 **Figure 7: The glacier margins were mapped as shown for Schnapfenkuchl M terminus illustrated in the orthophoto 2020 (A). The margin was mapped from thickness changes between 2004-2017 (B), as the total thickness loss of areas where the ice is gone is smaller than that for the ice margin (1,2), so that a characteristic pattern of thickness loss indicates the position of the glacier terminus and rock outcrops. The hillshade of the DEM 2018 (C) provides an additional source of information, as the glacier surface tends to be smoother than the surrounding area. Source orthophoto: CC 4.0 <https://www.data.gv.at/katalog/en/dataset/orthofoto>.**

215 The identification of areas with subsidence by melt is a clear advantage of this technology and allows mapping glacier areas that are hard to identify in orthophotos or VIS remote-sensing data. In a validation step, orthophotos (acquired at different dates) were used for a plausibility check of the LiDAR-derived outlines.

Several past glacier inventories use a minimum size threshold, as compiled by Leigh et al. (2019) based on the spatial resolution of the remote sensing data used. In this study, we applied no minimum size to mapping the glaciers. Only four of the 46 glaciers  
220 mapped were larger than 1 km<sup>2</sup> in 2002, and two of the glaciers were smaller than the 0.01 km<sup>2</sup> recommended by Paul et al. (2009) as a practical lower limit for mapping mountain glaciers by remote sensing.

### 3.2 Calculation of the volume change

The volume change for each pixel was calculated by multiplying the  $\Delta z$  with the 1x1 m<sup>2</sup> pixel area  $A$ . For the total volume change  $\Delta V$  of a glacier, the volume changes of all pixels within the glacier margins of the first date ( $t_1$ ) of the period  $t_1$  to  $t_2$   
225 (Equation 1) were summed up.

$$\Delta V = \sum A * \Delta z \quad \text{for all pixels part of the glacier area at } t_1 \quad (1)$$

The maximum uncertainty of  $\Delta z$  at one pixel is the sum of the uncertainties in the elevations of both DEMs involved. The uncertainty in area adds to the uncertainties in  $\Delta z$  for the uncertainty of the total volume change.

230 Presuming that  $\Delta z$  and  $\Delta A$  are independent variables, the uncertainties in area and thickness change propagate to the volume change uncertainty (Equation 2).

$$\sigma_{\Delta V} = |\Delta V| \sqrt{\left(\frac{\sigma_A}{A}\right)^2 + \left(\frac{\sigma_{\Delta z}}{\Delta z}\right)^2} \quad (2)$$

### 3.3 Calculation of the geodetic mass balance

The geodetic mass balance  $B_{geo}$  was calculated from volume change, assuming a constant glacier density  $\rho$  of 850 kg/m<sup>3</sup>  
235 (Equation 3).

$$B_{geo} = \Delta V * \rho \quad (3)$$

Calculating the geodetic mass balance from volume change requires assumptions on the stability of the glacier bed and on the density of the volume lost or gained. Erosion and deposition of sediments at the glacier bed was neglected for this study, as

research on the quantification of volumes is still ongoing. Previous studies on Austria's mass balance glaciers used a constant  
 240 density of 850 kg/m<sup>3</sup>, so did Fischer et al. (2015c) for the Swiss glaciers.

In recent years, the firn cover has melted. Thus the density at the surface may be higher. At the same time, ice flow velocities  
 have slowed down (Stocker-Waldhuber et al., 2019). This has resulted in the formation of englacial cavities (Stocker-  
 Waldhuber et al., 2017), which reduces the average density at the glacier tongues. The spatial and temporal variability of  
 glacier density  $\sigma\rho$  is expressed in an uncertainty of  $\pm 60$  kg/m<sup>3</sup>. Rocks, debris and sediments on and within the glacier are  
 245 treated as being part of the glacier.

$$\sigma_{B_{geo}} = |B_{geo}| \sqrt{\left(\frac{\sigma\rho}{\rho}\right)^2 + \left(\frac{\sigma\Delta V}{\Delta V}\right)^2} \quad (4)$$

The annual area-averaged specific geodetic mass balance  $b_{geo}$  is then calculated in two ways, i) dividing the geodetic mass  
 balance  $B_{geo}$  by the area of the glacier at the beginning of period  $t_1$  and by the number of years  $(t_2-t_1)$  and ii) dividing the  
 geodetic mass balance  $B_{geo}$  by the average of the glacier in the period  $t_1$  to  $t_2$  and by the number of years  $(t_2-t_1)$ .

$$250 \quad b_{geo\_total} = \frac{B_{geo}}{A(t_1) \cdot (t_2 - t_1)} \quad (5)$$

$$b_{geo\_average} = \frac{B_{geo}}{(A(t_1) + A(t_2)) / 2 \cdot (t_2 - t_1)} \quad (6)$$

The uncertainty of the annual area-averaged specific mass balance is then defined as (Equation 7)

$$\sigma_{b_{geo}} = |b_{geo}| \sqrt{\left(\frac{\sigma B_{geo}}{B_{geo}}\right)^2 + \left(\frac{\sigma A}{A}\right)^2} \quad (7)$$

255 where both potential definitions of the area, as area at the beginning of the period (Eq. 5) and as average area within the period,  
 (Eq. 6) were applied.

## 4 Results

### 4.1 Total area and volume changes

260 From the LIA maximum to 2017/18, the Austrian Silvretta lost 68% of its glacier area (Table 3, Figure S4). The mean annual  
 area loss in the latest period was 2.4%, which is more than twice the loss of the period before. The 10 totally debris-covered  
 glaciers cover an area 0.303 km<sup>2</sup> (Table S3). For three glaciers (ID 13006, Fluchthornferner S and Litzner Gletscher E), neither  
 bare ice nor signs of motion or a drainage system were visible, mean thickness changes between 2004/06 and 2017/18 were  
 smaller than 2.6 m without a clear thickness change pattern to indicate an ice margin. From that we can conclude that the  
 265 subsurface ice merely melted.

Area changes were calculated for all glaciers for the periods LIA maximum -1969 and 1969-2002 (Table 3). The dates of LiDAR campaigns differ between the federal states of Vorarlberg (2004 and 2017) and Tyrol (2006 and 2018). Later periods are thus 2002-2004/06, and 2004/06-2017/18.


270 **Table 3: Glacier area in the Austrian part of the Silvretta between LIA maximum, 1969, 2002, 2004/06 and 2017/18. For Vorarlberg, LiDAR flights were carried out in 2004 and 2017, for Tyrol in 2006 and 2018.**

Time	Area		% of LIA area		Period	Area change		
	km <sup>2</sup>					km <sup>2</sup>	%	% per yr
LIA maximum	40.9 ± 4.1		100 ± 10					
1969	24.0 ± 0.8		59 ± 2		LIA-1969	-17.0 ± 4.9	-41 ± 12	
2002	19.1 ± 0.6		47 ± 3		1969-2002	-4.8 ± 1.4	-20 ± 6	-0.6 ± 0.2
2004/06	18.5 ± 0.3		45 ± 1		2002-2004/06	-0.6 ± 0.9	-3 ± 5	-1.1 ± 1.6
2017/18	13.1 ± 0.4		32 ± 2		2004/06-2017/18	-5.4 ± 0.7	-29 ± 4	-2.4 ± 0.3

275 The annual specific geodetic mass balance (Table 4) was most negative in the short period with the extreme summer of 2003 (-1.5±0.7 m w.e./y), and least negative in the period 1969-2002 (-0.2±0.1 m w.e). That geodetic mass balance in the latest period is less negative in magnitude than in the short period before, can be attributed to the loss of ablation area within the period. The overall volume loss is reduced despite increasing ablation rates at the stakes because the areas with highest ablation rates in the past are ice-free now. The most negative glacier – wide specific direct mass balance was recorded in 2015 (Fischer et al., 2016).

**Table 4: Glacier volume loss, geodetic balance (total, specific and specific annual) in the Austrian part of the Silvretta between 1969, 2002, 2004/06 and 2017/18. For Vorarlberg, LiDAR flights were carried out in 2004 and 2017, for Tyrol in 2006 and 2018.**

	Volume loss	Geodetic balance	Specific geodetic balance	Annual specific geodetic balance
Period	km <sup>3</sup>	km <sup>3</sup> w.e.	m w.e.	m w.e./yr
1969-2002	-0.213 ± 0.092	-0.181 ± 0.079	-7.6 ± 3.3	-0.2 ± 0.1
2002-2004/06	-0.118 ± 0.048	-0.100 ± 0.042	-5.2 ± 2.2	-1.5 ± 0.7
2004/06-2017/2018	-0.237 ± 0.024	-0.201 ± 0.025	-10.9 ± 1.4	-0.8 ± 0.1



**Figure 8: Annual geodetic mass balances for the Austrian Silvretta in 1969/2002, 2002/2004/2006, 2004/2006/2017/2018.**

## 4.2 Volume changes and geodetic balance for individual glaciers

For individual glaciers, the three inventories also show strongest annual losses for the short period with the extreme year 2003 (Figure 8, Table S1), lowest annual losses for all glacier sizes in the period 1969-2002 and medium losses for the latest period. 290  
Glaciers with smaller areas present the highest variability. In contrast to the short and first warm inventory period 2002-2004/06, when some of the smaller glaciers were still quite stable, the range of geodetic mass balance for small glacier sizes is extremely high, from losses of more than 1 m w.e./ year to a few quite stable conditions, which are related to debris cover (or loss of ice).

## 295 4.3 Influence of area reference on geodetic balance

The choice of the area, either the area at the beginning of the period or the average area within the period, has significant impact on the geodetic mass balance, changing annual balance by up to 0.33 m w.e. /year for individual glaciers (Table S2). The balance becomes more negative if related to the average area, by -0.16 m w.e. /year. This also reveals that it is not only a neglect of debris-covered glacier areas, but also the wrong inclusion of debris-covered areas without ice, that can affect the 300  
uncertainty of geodetic mass balance.

## 5 Discussion

### 5.1 Challenges for mapping area changes of disintegrating glaciers

305 With ongoing glacier disintegration, mapping glacier outlines becomes ever more ambiguous even if using high-resolution volume change data. Major points to discuss are:

- what exactly are the properties that make a cryogenic feature a glacier which should be included in a glacier inventory,
- should we introduce inventories of all cryogenic features including glaciers, permafrost and rock glaciers or
- 310 - do we need to define a point of glacier disappearance?

All glaciers in this study were mapped first in previous inventories at times when bare ice was largely visible. For the now totally debris-covered glaciers, it is hard to decide using only optical data whether there is any ice left. In the absence of bare ice or surface structures, such as ponds or crevasses, and with soft slopes and low potential velocities, surface velocities do not 315  
help to distinguish buried glaciers from rock glaciers and permafrost dynamics. As measurements in bore holes in rock glaciers show, sliding debris and rocks on the ice can account for a major part of the total surface velocity (Krainer et al., 2015). If a subsurface runoff system exists at the terminus, it is necessary to analyse if it is fed by groundwater, melt of



seasonal snow, permafrost ice or glacier ice. This can be done by analysing seasonality of the amount of runoff (e.g. Brighenti et al., 2019), chemical composition and ecological properties (e.g. Tolotti et al., 2020) as well as by isotope analysis of the meltwater (e.g. Wagenbach et al., 2012).


Careful analysis is needed to decide whether a formerly well-defined glacier still fulfils the criteria of a glacier. Taking glaciers off inventories prematurely is to be avoided, as they may still contribute to glacial runoff in the basin, can force debris slide or serve as essential climate variables. We lose track of these transient states by dropping these glaciers from the inventories.

## 325 **5.2 Uncertainties in mapping totally debris-covered glaciers**

Mapping glaciers that have become fully covered with debris is uncertain for two reasons. First, mapping areas of possible ice covered with debris by volume changes only allows tackling the presence of melting ice during the inventory period, but it does not prove that any ice is left at the end of the period. Second, the presence of melting ice is not restricted to debris-covered glaciers but is also true for permafrost. The Schnapfenkuchl glaciers V and H present glacier ice at locations where debris flows exposed the ice and past inventories showed bare ice (Figure 9).

The Schnapfenkuchl glaciers are embedded in an environment adjacent to a number of rock glaciers. The delineation of buried glaciers in the presence of permafrost and mass movements upon and from the glacier needs a high temporal frequency of inventory data to arrive at the detection of glacier ice. High spatial resolution is needed to distinguish between ice loss and ice dynamics and to track the geomorphological processes and features related to volume change.

Surface elevation changes result from ice volume changes, but also from relocation, accumulation or erosion of supra-, en- or subglacial debris. The related geomorphological processes can be tackled by characteristic patterns of volume changes. The higher the temporal resolution of the repeat imagery, the more clearly can individual events be distinguished.




340


**Figure 9: Final glacier outline 2018 and volume changes 2006-2018 for the Schnapfenkuchl glaciers V, M and H and Larainferner displayed on (A) the orthophoto from 2020 and (B) the elevation changes from 2006-2018 superimposed on the hillshade of 2018. Source orthophoto: CC 4.0 <https://www.data.gv.at/katalog/en/dataset/orthofoto>.**

### 345 5.3 Comparison to remote sensing results

The Sentinel-2-based inventory of the Alps is based on data of exactly the same year, 2018, as the Tyrolean LiDAR inventory, for Vorarlberg acquisition dates differ by one year. This is an opportunity to compare the representation of different types of glaciers in the various data sets, as shown for Landsat 8 (2018-09-09, 30 m resolution), Sentinel-2 (2018-09-09, 10 m resolution), and LiDAR data (2018) as well as orthophotos (2020) for the bare ice snout of Schnapfenkuchl M (Figure 10) and the totally debris-covered Schnapfenkuchl H (Figure 11). While clean ice can be easily mapped, the identification of debris-covered glaciers like Schnapfenkuchl H is more challenging in all types of optical imagery, independent of the pixel spacing. Although higher resolution of optical imagery allows the detection of features potentially related to subsurface deformation, the potential for the quantification of the ice extent is limited. Volume change allows only the detection of the past presence of ice, and not the proof of actual presence of ice. Thus the area affected by volume change can be constrained. The regional comparison of change rates can also give a hint of the total melt within the period, as in this case the volume change rate is lower than in similar settings with permanent presence of ice.



360 **Figure 10: The comparison of the Landsat 8, Sentinel-2, LiDAR (hillshade and volume change 200620018, colour map as for Figure 9), all dating from 2018, and the orthophoto (2020) illustrate the different representation of the glacier in the various spatial resolutions for the snout of Schnapfenkuchl M. For glaciers with bare ice, glacier margins are clear at all scales of resolutions.**



**Figure 11: Representation of the debris-covered glacier Schnapfenkuchl H in the Landsat 8, Sentinel-2, LiDAR images**  
 365 **(hillshade and volume change 200620018, colour map as for Figure 9) and the orthophoto. The manual delineation of**  
**glacier outlines (red: based on LiDAR, black: based on Sentinel-2, Paul et al., 2019) is consistent for both methods.**

For the total region, the Sentinel-based glacier area is 13.035 km<sup>2</sup>, therefore only 0.098 smaller than the LiDAR-based  
 inventory, both listing 46 glaciers. Part of the consistency of the inventories results from the fact that both are based on the  
 370 same older glacier inventory, and that both observers also focused on consistency with this inventory. Therefore the  
 compilation of time series of inventories as homogenous as possible could be a major advantage for precision even in the  
 absence of high-resolution data.

375 **5.4 Should we still map small and totally debris-covered glaciers?**

Past definitions of glaciers focussed on the mass of ice. From the earliest definitions, e.g. by Walcher (1773) and Tyndall (1860), to more modern ones within glaciology (Klebensberg, 1948) and in neighbouring fields (Dexter, 2013), scientists have obviously been investigating glaciers closer to equilibrium than those in our study.

380 *“Diese Eisberge werden Ferner genennet, welches Wort... das Eis bedeutet welches mit Schnee vermengt gesammelt hat“*  
(Walcher, 1773) [These mountains of ice are called „Ferner“, a word ...that denotes ice mixed with snow which has accumulated]

*“At its origin then a glacier is snow — at its lower extremity it is ice.”* (Tyndall, 1860)

385 ***Gletscher sind Massen körnigen Firns und Eises, die aus Schneeanstimmungen hervorgehen und sich dahin bewegen, wo sie abschmelzen oder verdunsten können....scheinbarer oder auch wirklicher Mangel an Bewegung schliesst aber doch die Bezeichnung Gletscher nicht aus, übergeordnetes Merkmal ist die Körnerstruktur.*** (Klebensberg 1948)

*[Glaciers are masses of granulated firn and ice that have evolved out of accumulations of snow and which move towards where they can melt or evaporate... however, an apparent or actual lack of movement does not preclude it being labelled a glacier, the predominant characteristic is the granular structure.]*

390

*“A glacier is a mass of relatively slow moving ice created by the accumulation of snow”* (Dexter, 2013)

*“A perennial mass of ice, and possibly firn and snow, originating on the land surface by the recrystallization of snow or other forms of solid precipitation and showing evidence of past or present flow... In contrast to what is natural in dynamic glaciology and glacial geomorphology, for mass-balance purposes the glacier consists only of frozen water. Sediment carried by the glacier is deemed to be outside the glacier.”*

395

(Cogley et al., 2011)

*“A glacier or perennial snow mass, identified by a single GLIMSglacier ID, consists of a body of ice and snow that is observed at the end of the melt season, or, in the case of tropical glaciers, after transient snow melts. This includes, at a minimum, all tributaries and connected feeders that contribute ice to the main glacier, plus all debris-covered parts of it. Excluded is all exposed ground, including nunataks. .... A stagnant ice mass still in contact with a glacier is part of the glacier, even if it supports an old-growth forest...All debris-covered parts of the glacier must be included.... Rock glaciers and heavily debris-covered glaciers tend to look similar, but their geneses are different. GLIMS does not currently deal with the former, but does include the latter.”*

400

(Raup and Khalsa, 2010)

405

The more modern definitions of Cogley et al. (2011) and Raup and Khalsa (2010) address the need of the glaciological community for mapping glaciers, mainly based on remote-sensing data.

410 Although englacial and supraglacial debris was present on and in glaciers even during the LIA, ice and snow dominated at the time. In our glacier inventory of 2017/18, 10 of 43 glaciers are predominantly covered by debris, not snow, with only few remnants of bare ice visible. Mean thickness changes range from  $-1.5 \pm 0.1$  m to  $-12.3 \pm 0.6$  m for the total period. At Garnera Gletscher and glaciers #12018, #13005, no bare ice is visible. On the surface of all other glaciers, bare ice is visible in parts. Several of the debris-covered glaciers, for example, the Schnapfenkuchl glaciers (Figure 9), are located between active rock glaciers captured in the Tyrolean rock glacier inventory (Krainer and Ribis, 2012). This means that two independent groups of  
415 researchers, glaciologists and geologists, mapped the same site, for example, the easternmost Schnapfenkuchl glacier/rock glacier, in inventories of different landforms. A continuum from glacier to debris-covered glacier to rock glacier was recently discussed by Anderson et al. (2018). Kellerer-Pirklbauer and Kaufmann (2018) analysed the glacial history of an Austrian site of long-term rock glacier monitoring and found evidence of post-LIA deglaciation to permafrost formation. For the broader scientific investigation of this phenomenon, international cooperation for comparing processes and regimes in other mountain  
420 regions are vital, since a case study can only be a first step towards deriving more general empirical and theoretical knowledge. More concise definitions of the term *glacier* are needed to reflect current conditions on the ground. We suggest:

- 1) Glaciers are bodies of sedimentary ice, firn and snow, formed by densification of snow AND en- and supraglacial sediments of all grain sizes.

or

425 2) Glaciers are exposed bodies of sedimentary ice, firn and snow formed by densification of snow with signs of deformation and an en- or supraglacial drainage system.

Definition (1) includes all sizes of ice bodies which have been formed as part of glaciers, even dead debris-covered glacier ice and ice cored moraines. It understands debris as a natural part of the glacier system, also in the calculation of volume changes,  
430 and sets down a clear base for mapping glaciers in inventories and calculating geodetic balances. Mapping internal properties as englacial deformation or the existence of drainage systems often is not possible by remote sensing, and the advantage of definition (1) is that there is no need to do so. The drawback is, that including debris accepts that glacier volume changes are not necessarily ice volume changes. This accounts for the 'real world' problem that it is not possible to assess the amount of englacial debris anyway.

435

Definition (2) excludes dead and buried glacier ice, which might have advantages for mapping, especially in low resolutions, but leaves an undefined gap between glaciers and permafrost ice, thereby introducing a third class, i.e. buried glacier ice. Janke et al. (2015) propose a gradual transition between debris-covered and buried ice, with still ongoing melt and respective volume loss for the ice buried by 3-5 m of debris. As criteria for the classification as rock glacier, they apply a complete debris

440 cover with no ice visible at the surface and a less heterogeneous internal structure, with glacier ice, segregated ice and  
interstitial ice. The ice content varies between 25 to 45 %. Predominant surface features are thermokarst depressions and  
transverse ridges and furrows. This classification is definitely valuable for mapping glaciers and rock glaciers, but the  
traditional Austrian rock glacier and glacier inventories have been using other types of classifications, e.g. including parts of  
bare ice in the rock glacier inventory. The coming years or decades will show if the transient features in the Austrian Silvretta  
445 can develop ridges or furrows, or if the ice is gone before these structures evolve.

#### **5.4 Comparison of changes in the Austrian Silvretta with other glacier regions**

For the Austrian Silvretta, the mean annual geodetic balance is  $-0.8 \pm 0.1$  m w.e./yr for the period from 2004/06 to 2017/18.  
This is less negative than the  $-1.03$  m w.e. reported for the Glarus and Leopoldine Alps by Sommer et al. (2020) for the period  
450 from 2000 to 2014, but but a greater loss than than the  $-0.62$  m w.e./yr calculated by Fischer, Huss and Hoelzle (2015) for the  
Swiss Alps between 1980 and 2010. They also reported a variability of geodetic balances on the catchment scale, ranging from  
 $-0.52$  to  $-1.07$  m w.e. Our data confirm the highly variable sensitivity of small glaciers to warming as found by Huss and  
Fischer (2016).

The annual rate of area losses in the Austrian Silvretta (1969-2017/18) of  $-1.13\%$  is larger than the  $-0.52\%$  reported for a  
455 slightly shorter period (1986-2014) in the Caucasus (Tielidze et al., 2020), but similar to the  $-1.1\%$  reported by Paul et al.  
(2020) for the Alps between 2003 and 2015/16 excluding glaciers smaller than  $0.01$  km<sup>2</sup>.

Although the respective regions are very different in terms of elevation range, climate and glacier cover, the glacier mass loss  
is striking similar.

For mapping small glaciers in Norway, Leigh et al. (2019) developed a rating system, classifying glaciers from ‘certain’ to  
460 ‘probable’ based  $<1$  m resolution visible imagery. The scoring is based on visible features as ogives, crevasses, and moraines.  
In absence of volume change data, this might be a helpful approach to keep track of buried glacier ice.

#### **5.5 Additional uncertainties of geodetic mass balance**

Geodetic mass balance has been analysed for potential uncertainties, for example, unaccounted seasonal snow interpreted as  
465 ice volume change or changing glacier beds, and real mass changes differing from surface mass balance, for example, as a  
result of refreezing, firn density changes, or crevasse volume (e.g. Zemp et al., 2013). For the very small and debris-covered  
glaciers we analysed, an additional source of uncertainty is the amount of the accumulated debris on the surface. This volume  
is not part of the hydrological cycle so that, from a hydrological perspective, we would wish to exclude this volume from the  
analysis. This would be possible only if we could keep track of the erosion rates in the source areas. This would necessitate a  
470 totally different monitoring system to track the steep headwalls. In terms of geomorphology, debris and rocks are part of the  
glacier and therefore there is no need to distinguish old from new supra- and glacial debris and rock.


The rougher surface at steep debris-covered glaciers with avalanche activity seems to encourage the formation of perennial snow patches on the debris-covered glacier. This is not relevant for delineating glaciers but raises the question on the effects of these perennial snow volumes on geodetic mass balance. This question could be also answered with a specific monitoring effort tackling the now stored volumes and water equivalents. In this effort first hints on potential ice formation in such structures could be gathered.

## 6 Summary and Conclusions

The annual rates of change of area and volume change indicate an increasing pace of glacier retreat in the Austrian Silvretta. A growing number of nunataks, the disintegration of larger glaciers into smaller ones and the accumulation and relocation of debris on the glacier surface all indicate an additional need for compiling glacier inventories - if we want to keep track of buried glacier ice. Figure 12 outlines the procedure applied in this study. For most of the 46 glaciers, the procedure of delineating the glacier surface by surface roughness and volume change validated by orthophotos worked out well, even for the ten glaciers totally covered by debris. However, for some of the analysed glaciers, for example Schnapfenkuchl V, future analysis of areas will be difficult, as the surface is now totally covered with debris, and accumulation and relocation of the debris is still ongoing. Therefore, we cannot expect a distinct pattern of thickness change in the future, with a maximum close to the terminus. In the next decade we thus need a scientific discussion on how to proceed with as yet undefined subsurface glacier remnants.

The technical requirements for mapping small vanishing glaciers depend on glacier size, annual rate of thickness changes and accumulation rates of debris. The small glacier structures that remain in the Austrian Silvretta are typically just a few tens of metres wide. With increasing pixel size, the accuracy of mapping vertical changes decreases, which makes it difficult to distinguish geomorphological processes like accumulation and relocation of rocks and debris from ice volume change. Therefore, we recommend 1 m spatial resolution. The vertical accuracy needed to represent the geomorphological processes of debris relocation depends on the steepness of the area and on the temporal interval chosen. For a period of 10 years and a slope of up to 40° for Alpine glaciers, a vertical resolution of 1/10 of the spatial resolution is sufficient to distinguish volume changes by ice melt from erosion and deposition.





**Figure 12: Summary of the workflow for high-resolution glacier mapping and mass balance calculation as applied in this study. The workflow does lead to quantitative results for most of the glaciers, but also shows up the need of scientific discussion where geomorphological processes other than ice melt contribute significantly to volume changes. See Figure 9 for the colour table of the volume change maps displayed.**

500

Of the now 43 glaciers of the Austrian Silvretta, only three are larger than 1 km<sup>2</sup>, 19 are smaller than 0.1 km<sup>2</sup> (of these, 13 are smaller than 0.05 km<sup>2</sup>). Applying minimum sizes for glacier area with thresholds at 0.1 or 0.05 km<sup>2</sup> would exclude 0.82 km<sup>2</sup>/6.2% and 0.35 km<sup>2</sup>/2.7% of the total area. This is higher than /in the same magnitude as the nominal uncertainty of

505

0.4 km<sup>2</sup> in the total area. Such thresholds for very small glaciers would make many of them disappear from inventories and hamper any efforts to tackle the hazard potential of deglaciation, as even small glacier remnants could be relevant here.

510 There is probably no hard limit for surveying deglaciation in terms of glacier size, as the monitoring strategies for rock glaciers and permafrost can take over. Handing over what is left from glaciers to the scientific networks of the permafrost community could be an emerging and exciting new playground for both fields.

This regional study can only point out the specific challenges and limitations for tackling glacier change in the Austrian Silvretta. These will differ from region to region depending on the climatic regime, lithology, topography, glacier types and many other factors. We would therefore appreciate an international effort to compare the need for tackling deglaciation in other regions, as a common monitoring framework will be essential in an ever warmer future.

515

### **Author Contributions**

Andrea Fischer designed this study, worked on glaciers in the Silvretta range for over a decade and wrote the text. Kay Helfricht, Bernd Seiser and Martin Stocker-Waldhuber analysed the geodetic data, Gabriele Schwaizer contributed remote sensing data and expertise. All authors contributed to discussion and text.

### 520 **Funding**

Both LiDAR DEMs were provided by the TIRIS section of the federal administration of Tyrol and the federal administration of Vorarlberg.

### **Acknowledgments**

525 The federal administrations of Tyrol and Vorarlberg are acknowledged for providing geodata. We thank the community of Galtür, the Lorenz family at Jamtalhütte and Oswald Heis for supporting logistics. Kati Heinrich helped with the figures, Brigitte Scott checked the English - thank you!

### **Data availability statement**

The glacier inventory data are available at <https://doi.org/10.1594/PANGAEA.844988> (Fischer et al., 2015b) and <https://doi.org/10.1594/PANGAEA.XXXXXX> (submitted).

530

## References

- Abermann, J., Fischer, A., Lambrecht, A., and Geist, T.: On the potential of very high-resolution repeat DEMs in glacial and periglacial environments. *The Cryosphere* 4, 53–65. <https://doi.org/10.5194/tc-4-53-2010>, 2010.
- Abermann, J., Lambrecht, A., Fischer, A., and Kuhn, M.: Quantifying changes and trends in glacier area and volume in the  
535 Austrian Ötztal Alps (1969-1997-2006), *The Cryosphere* 3, 205–215, <https://doi.org/10.5194/tc-3-205-2009>, 2009.
- Anderson, R. S., Anderson, L. S., Armstrong, W. H., Rossi, M. W., and Crump, S. E.: Glaciation of alpine valleys: The glacier – debris-covered glacier – rock glacier continuum. *Geomorphology* 311, 127–142, <https://doi.org/10.1016/j.geomorph.2018.03.015>, 2018.
- Bahr, D. B. and Radić, V.: Significant contribution to total mass from very small glaciers. *The Cryosphere* 6, 763–770,  
540 <https://doi.org/10.5194/tc-6-763-2012>, 2012.
- BEV [https://www.bev.gv.at/portal/page?\\_pageid=713,2363167&\\_dad=portal&\\_schema=PORTAL](https://www.bev.gv.at/portal/page?_pageid=713,2363167&_dad=portal&_schema=PORTAL) 2011a  
BEV [https://www.bev.gv.at/portal/page?\\_pageid=713,2363167&\\_dad=portal&\\_schema=PORTAL](https://www.bev.gv.at/portal/page?_pageid=713,2363167&_dad=portal&_schema=PORTAL) 2011b
- Bojinski, S., Verstraete, M., Peterson, T., Richter, C., Simmons, A., and Zemp, M.: The Concept of Essential Climate Variables in Support of Climate Research, Applications, and Policy. *Bulletin of the American Meteorological Society*. 95. 1431.  
545 <https://doi.org/10.1175/BAMS-D-13-00047.1>, 2014.
- Braumann, S. M., Schaefer, J. M., Neuhuber, S. M., Reitner, J. M., Lüthgens, C., and Fiebig, M.: Holocene glacier change in the Silvretta Massif (Austrian Alps) constrained by a new <sup>10</sup>Be chronology, historical records and modern observations. *Quaternary Science Reviews*, 245, 106493, <https://doi.org/10.1016/j.quascirev.2020.106493>, 2020.
- Brighenti S., Tolotti M., Bruno M.C., Wharton G., Pusch M.T., and Bertoldi W.: Ecosystem shifts in Alpine streams under  
550 glacier retreat and rock glacier thaw: a review. *Science of the Total Environment*, 675,542-559. <https://doi.org/10.1016/j.scitotenv.2019.04.221>, 2019.
- data.tirol.gv.at: Current and historical orthoimagery of Tyrol. Open government data, [https://www.data.gv.at/katalog/dataset/land-tirol\\_orthofototirol](https://www.data.gv.at/katalog/dataset/land-tirol_orthofototirol), last access: 2020-10-11, 2020.
- Dexter, L. R., and Birkeland K. W.: Snow, Ice, Avalanches, and Glaciers. In: *Mountain Geography: Physical and Human*  
555 *Dimensions*, edited by L. W. Price et al., 1st ed., University of California Press, 2013, pp. 85–126. JSTOR, [www.jstor.org/stable/10.1525/j.ctt46n4cj.10](http://www.jstor.org/stable/10.1525/j.ctt46n4cj.10). Accessed 15 Dec. 2020.
- Federal Government of Tyrol: Landesweite Laserscanbefliegung 2006-2010, [https://www.tirol.gv.at/fileadmin/themen/sicherheit/geoinformation/Laserscandaten/ALS\\_Tirol\\_Onlinebericht.pdf](https://www.tirol.gv.at/fileadmin/themen/sicherheit/geoinformation/Laserscandaten/ALS_Tirol_Onlinebericht.pdf), last access 11.12.2020.
- 560 Fischer, A., Patzelt, G., and Kinzl, H.: Length changes of Austrian glaciers 1969-2016. Institut für Interdisziplinäre Gebirgsforschung der Österreichischen Akademie der Wissenschaften, Innsbruck, PANGAEA, <https://doi.org/10.1594/PANGAEA.821823>, 2016a.

- Fischer, A., Helfricht, K., Wiesenegger, H., Hartl, L., Seiser, B., and Stocker-Waldhuber, M.: Chapter 9 - What Future for Mountain Glaciers? Insights and Implications From Long-Term Monitoring in the Austrian Alps. In: Gregory B. Greenwood and J.F. Shroder, Editor(s), *Developments in Earth Surface Processes*, Elsevier, 21, 325-382. <http://dx.doi.org/10.1016/B978-0-444-63787-1.00009-3>, 2016b.
- Fischer, A. Markl, G., and Kuhn, M.: Glacier mass balances and elevation zones of Jamtalferner, Silvretta, Austria, 1988/1989 to 2016/2017. Institut für Interdisziplinäre Gebirgsforschung der Österreichischen Akademie der Wissenschaften, Innsbruck, PANGAEA, <https://doi.org/10.1594/PANGAEA.818772>, 2016c.
- 570 Fischer, A., Patzelt, G., AchRAINER, M., Groß, G., Lieb, G. K., Kellerer-Pirklbauer, A., and Bendler, G.: *Gletscher im Wandel: 125 Jahre Gletschermessdienst des Alpenvereins*. Springer Spektrum, 140 S. <https://doi.org/10.1007/978-3-662-55540-8>, 2018.
- Fischer, A., Seiser, B., Stocker-Waldhuber, M., Mitterer, C., and Abermann, J.: Tracing glacier changes in Austria from the Little Ice Age to the present using a lidar-based high-resolution glacier inventory in Austria. *The Cryosphere* 9, 753-766, <https://doi.org/10.5194/tc-9-753-2015>, 2015a.
- Fischer, A., Seiser, B., Stocker-Waldhuber, M., Mitterer, C., and Abermann, J.: The Austrian Glacier Inventories GI 1 (1969), GI 2 (1998), GI 3 (2006), and GI LIA in ArcGIS (shapefile) format. PANGAEA, <https://doi.org/10.1594/PANGAEA.844988> 2015b.
- Fischer, M., Huss, M., and Hoelzle, M.: Surface elevation and mass changes of all Swiss glaciers 1980–2010. *The Cryosphere*, 580 9, 525–540, <https://doi.org/10.5194/tc-9-525-2015>, 2015c.
- Gärtner-Roer, I., Nussbaumer, S.U., Hüsler, F., and Zemp, M.: Worldwide assessment of national glacier monitoring and future perspectives. *Mountain Research and Development*, 39(2). <https://doi.org/10.1659/MRD-JOURNAL-D-19-00021.1>, 2019.
- Geist, T. and Stötter, H.: First results of airborne laser scanning technology as a tool for the quantification of glacier mass balance. *Proceedings, EARSeL workshop on observing our cryosphere from space: techniques and methods for monitoring snow and ice with regard to climate change*, 11 - 13 März2002, Bern, 2002.
- geoland.at.: The free geodata portal of Austria, last access: 2020-10-12, 2020.
- Haeberli, W., Hoelzle, M., Paul, F., and Zemp, M.: Integrated monitoring of mountain glaciers as key indicators of global climate change: The European Alps. *Annals of Glaciology*, 46, 150-160. <https://doi.org/10.3189/172756407782871512>, 2007.
- 590 Höfle, B. and Rutzinger, M.: Topographic airborne LiDAR in geomorphology: A technological perspective. *Zeitschrift für Geomorphologie, Supplementary Issues*. 55. 1-29, 2011. 10.1127/0372-8854/2011/0055S2-0043.
- Huss, M., and Fischer, M.: Sensitivity of very small glaciers in the Swiss Alps to future climate change. *Frontiers in Earth Science*, 4. <https://doi.org/10.3389/feart.2016.00034>, 2016.
- Huss, M., Dhulst, L., and Bauder, A.: New long-term mass balance series for the Swiss Alps. *Journal of Glaciology* 61 (227), 595 551-562. <https://doi.org/10.3189/2015JoG15j015>, 2015.

- Huss, M., Zemp, M., Joerg, P.C., and Salzmann, N.: High uncertainty in 21st century runoff projections from glacierized basins. *Journal of Hydrology*, 510, 35-48. 2014.
- IPCC.: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M. Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)]. 2019.
- 600 Janke, J. R., Bellisario, A. C., Ferrando, F. A.: Classification of debris-covered glaciers and rock glaciers in the Andes of central Chile, *Geomorphology*, 241, 98-121, 2015. <https://doi.org/10.1016/j.geomorph.2015.03.034>.
- Kargel, J.S., Leonard, G.J., Bishop, M.P., Kaab, A., Raup, B., (Eds.): *Global Land Ice Measurements from Space* Springer, 876 pp. 2014.
- 605 Kellerer-Pirklbauer, A., and Kaufmann, V.: Deglaciation and its impact on permafrost and rock glacier evolution: New insight from two adjacent cirques in Austria. *Science of The Total Environment* 621, 1397-1414, <https://doi.org/10.1016/j.scitotenv.2017.10.087>, 2018.
- Klebsberg, R. v.: *Handbuch der Gletscherkunde und Glazialgeologie*, 1. Band, Allgemeiner Teil, Springer, Wien, 403 p.. 1948.
- 610 Krainer, K., and Ribis, M.: A rock glacier inventory of the Tyrolean Alps (Austria). *Austrian Journal of Earth Sciences* 105(2). 32-47. 2012.
- Krainer, K., Bressan, D., Dietre, B., Haas, J. N., Hajdas, I., Lang, K., Mair, V., Nickus, U., Reidl, D., Thies, H., and Tonidandel, D.: A 10,300-year-old permafrost core from the active rock glacier Lazaun, southern Ötztal Alps (South Tyrol, northern Italy). *Quaternary Research*, 83, 2, 324-335, <https://doi.org/10.1016/j.yqres.2014.12.005>, 2015.
- 615 Landesvermessungsamt Feldkirch, *Technischer Bericht zur Evaluierung der Laserscanning-Höhenmodelle Unterland & Vorderwald der Firma Topscan GesmbH (insgesamt 545 km<sup>2</sup>), Aktenzahl: LVA – 603.02.4.04.09.07, provided by the Federal Government of Vorarlberg*, 2004.
- Leigh, J., Stokes, C., Carr, R., Evans, I., Andreassen, L., and Evans, D.: Identifying and mapping very small (<0.5 km<sup>2</sup>) mountain glaciers on coarse to high-resolution imagery. *Journal of Glaciology*, 65(254), 873-888, 2019.
- 620 [doi:10.1017/jog.2019.50](https://doi.org/10.1017/jog.2019.50)
- Marzeion, B., Jarosch, A. H., and Hofer, M.: Past and future sea-level change from the surface mass balance of glaciers. *The Cryosphere*, 6, 1295–1322. <https://doi.org/doi.org/10.5194/tc-6-1295-2012>, 2012.
- Meier, W. J.-H., Griebinger, J., Hochreuther, P., and Braun, M. H.: An Updated Multi-Temporal Glacier Inventory for the Patagonian Andes With Changes Between the Little Ice Age and 2016. *Frontiers in Earth Science*, 6.
- 625 <https://doi.org/10.3389/feart.2018.00062>, 2018.
- Nagai, H., Fujita, K., Sakai, A., Nuimura, T., and Tadono, T.: Comparison of multiple glacier inventories with a new inventory derived from high-resolution ALOS imagery in the Bhutan Himalaya. *The Cryosphere*, 10, 65–85. <https://doi.org/10.5194/tc-10-65-2016>, 2016.

- Paul, F., Barry, R.G., Cogley, J.G., Frey, H., Haeberli, W., Ohmura, A., Ommanney, C.S.L., Raup, B., Rivera, A., and Zemp, M.: Recommendations for the compilation of glacier inventory data from digital sources. *Annals of Glaciology*, 50(53), 119-126. 2009.
- Paul, F., Rastner, P., Azzoni, R. S., Diolaiuti, G., Fugazza, D., Le Bris, R., Nemeč, J., Rabatel, A., Ramusovic, M., Schwaizer, G., and Smiraglia, C.: Glacier shrinkage in the Alps continues unabated as revealed by a new glacier inventory from Sentinel-2. *Earth Syst. Sci. Data*, 12, 1805–1821. <https://doi.org/10.5194/essd-12-1805-2020>, 2020.
- Pellikka, P. (ed.) and Rees, W. G. (ed.): *Remote Sensing of Glaciers: Techniques for topographic, spatial and thematic mapping of glaciers*, Leiden: CRC Press, Taylor & Francis Group, A Balkema Book. 340 p., 2009.
- Pfeffer, W. T., Arendt, A. A., Bliss, A., Bolch, T., Cogley, J. G., Gardner, A. S., Hagen, J.-O., Hock, R., Kaser, G., Kienholz, C., Miles, E. S., Moholdt, G., Mölg, N., Paul, F., Radic, V., Rastner, P., Raup, B.H., Rich, J., Sharp, M. J. and the Randolph Consortium: The Randolph Glacier Inventory: a globally complete inventory of glaciers. *Journal of Glaciology* 60, 537—552. 2014.
- Racoviteanu, A., Paul, F., Raup, B., Khalsa, S., and Armstrong, R.: Challenges and recommendations in mapping of glacier parameters from space: Results of the 2008 Global Land Ice Measurements from Space (GLIMS) workshop, Boulder, Colorado, USA. *Annals of Glaciology*, 50(53), 53-69, 2009. doi:10.3189/172756410790595804
- Raup, B. and Khalsa, S.J.S.: GLIMS Analysis Tutorial. GLIMS, Global Land Ice Measurements from Space, NSIDC, [www.GLIMS.org https://www.glims.org/MapsAndDocs/assets/GLIMS\\_Analysis\\_Tutorial\\_a4.pdf](https://www.glims.org/MapsAndDocs/assets/GLIMS_Analysis_Tutorial_a4.pdf), 2010.
- Rieger, W.: ALS Tirol 2017-18, Abschlussbericht AZ VIg-450/10 GZ/AVT: 31757, ARGE AVT-Milan, 48 p, 2019.
- Sailer, R., Rutzinger, M., Rieg, L. and Wichmann, V. : Digital elevation models derived from airborne laser scanning point clouds: appropriate spatial resolutions for multi-temporal characterization and quantification of geomorphological processes. *Earth Surf. Process. Landforms*, 39: 272-284. <https://doi.org/10.1002/esp.3490>, 2014.
- Sommer, C., Malz, P., Seehaus, T.C. et al. Rapid glacier retreat and downwasting throughout the European Alps in the early 21st century. *Nat Commun* 11, 3209, <https://doi.org/10.1038/s41467-020-16818-0>, 2020.
- Tielidze, L. G., Bolch, T., Wheate, R. D., Kutuzov, S. S., Lavrentiev, I. I., and Zemp, M.: Supra-glacial debris cover changes in the Greater Caucasus from 1986 to 2014. *The Cryosphere*, 14, 585-598. <https://doi.org/10.5194/tc-14-585-2020>, 2020.
- Tolotti M., Cerasino L., Donati C., Pindo M., Rogora M., Seppi R., and Albanese D.: Alpine headwaters emerging from glaciers and rock glaciers host different bacterial communities: Ecological implications for the future. *Science of the Total Environment*, 717:137101. <https://doi.org/10.1016/j.scitotenv.2020.137101>, 2020.
- Tyndall, J. Origin of glaciers. In *The Glaciers of the Alps: Being a Narrative of Excursions and Ascents, an Account of the Origin and Phenomena of Glaciers and an Exposition of the Physical Principles to Which They Are Related* (Cambridge Library Collection - Earth Science, pp. 248-249). Cambridge: Cambridge University Press. doi:10.1017/CBO9781139097048.034, 1860.

- Wagenbach, D., Bohleber, P., and Preunkert, S.: Cold, alpine ice bodies revisited: what may we learn from their impurity and isotope content?, *Geografiska Annaler: Series A, Physical Geography*, 94:2, 245-263. <https://doi.org/10.1111/j.1468-0459.2012.00461.x>, 2012.
- Walcher, J.: Nachrichten von den Eisbergen im Tyrol, Frankfurt und Leipzig, Permalink: <http://mdz-nbn-resolving.de/urn:nbn:de:bvb:12-bsb10011721-1>, 1773.
- WGMS, and National Snow and Ice Data Center (comps.): World Glacier Inventory, Version 1. Boulder, Colorado USA. NSIDC: National Snow and Ice Data Center. <https://doi.org/10.7265/N5/NSIDC-WGI-2012-02>, 1999, updated 2012.
- Würländer, R: Airborne Laserscanner Daten 2017, Abschlussbericht. Provided by the Federal Government of Vorarlberg, 2019.
- 670 Zekollari, H., Huss, M., and Farinotti, D.: Modelling the future evolution of glaciers in the European Alps under the EURO-CORDEX RCM ensemble. *The Cryosphere*, 13, 1125–1146. <https://doi.org/10.5194/tc-13-1125-2019>, 2019.
- Zekollari, H., Huss, M., and Farinotti, D.: On the imbalance and response time of glaciers in the European Alps. *Geophysical Research Letters*, 47(2), e2019GL085578. <https://doi.org/10.1029/2019GL085578>, 2020.
- Zemp, M., Frey, H., Gärtner-Roer, I., Nussbaumer, S.U., Hoelzle, M., Paul, F., Haeberli, W., Denzinger, F., Ahlstrom, A.P., 675 Anderson, B., Bajracharya, S., Baroni, C., Braun, L.N., Cáceres, B.E., Casassa, G., Cobos, G., Davila, L.R., Delgado Granados, H., Demuth, M.N., Espizua, L., Fischer, A., Fujita, K., Gadek, B., Ghazanfar, A., Hagen, J.O., Holmlund, P., Karimi, N., Li, Z., Pelto, M., Pitte, P., Popovnin, V.V., Portocarrero, C.A., Prinz, R., Sangewar, C.V., Severskiy, I., Sigurdsson, O., Soruco, A., Usabaliyev, R., and Vincent, C.: Historically unprecedented global glacier decline in the early 21st century. *Journal of Glaciology*, 61 (228), 745-762. <https://doi.org/10.3189/2015JoG15J017>, 2015.
- 680 Zemp, M., Huss, M., Thibert, E. et al.: Global glacier mass changes and their contributions to sea-level rise from 1961 to 2016. *Nature* 568, 382–386. <https://doi.org/10.1038/s41586-019-1071-0>, 2019a
- Zemp, M., Sajood, A.A., Pitte, P., van Ommen, T., Fischer, A., Soruco, A., Thomson, L., Schaefer, M., Li, Z., Ceballos Lievano, J.L., Cáceres Correa, B.E., Vincent, C., Tielidze, L., Braun, L.N., Ahlstrøm, A.P., Hannesdóttir, H., Dobhal, D.P., Karimi, N., Baroni, C., Fujita, K., Severskiy, I., Prinz, R., Usabaliyev, R., Delgado-Granados, H., Demberel, O., Joshi, S.P., 685 Anderson, B., Hagen, J.O., Dávila Roller, L.R., Gadek, B., Popovnin, V.V., Cobos, G., Holmlund, P., Huss, M., Kayumov, A., Lea, J.M., Pelto, M., and Yakovlev, A.: Glacier monitoring to track warming. *Nature*, 576, p. 39. [go.nature.com/34ak25y](https://www.nature.com/34ak25y), 2019b.
- Zemp, M., Thibert, E., Huss, M., Stumm, D., Rolstad Denby, C., Nuth, C., Nussbaumer, S. U., Moholdt, G., Mercer, A., Mayer, C., Joerg, P. C., Jansson, P., Hynes, B., Fischer, A., Escher-Vetter, H., Elvehøy, H., and Andreassen, L. M.: Reanalysing 690 glacier mass balance measurement series, *The Cryosphere*, 7, 1227–1245, <https://doi.org/10.5194/tc-7-1227-2013>, 2013.