

Spectral attenuation coefficients from measurements of light transmission in bare ice on the Greenland Ice Sheet

Matthew G. Cooper^{1,2}, Laurence C. Smith^{3,4}, Asa K. Rennermalm⁵, Marco Tedesco^{6,7}, Rohi Muthyala⁵, Sasha Z. Leidman⁵, Samiah E. Moustafa³, Jessica V. Fayne¹

- 5 ¹Department of Geography, University of California, Los Angeles, Los Angeles, 90027, USA
²Pacific Northwest National Laboratory, Richland, Washington, 99354, USA
³Institute at Brown for Environment and Society, Brown University, Providence, 02912, USA
⁴Department of Earth, Environmental and Planetary Sciences, Brown University, Providence, 02912, USA
⁵Department of Geography, Rutgers University, Piscataway, 08854, USA
10 ⁶NASA Goddard Institute for Space Studies, New York, 10025, USA
⁷Lamont Doherty Earth Observatory, Columbia University, New York, 10964, USA

Correspondence to: Matthew G. Cooper (guycooper@ucla.edu)

Abstract. Light transmission into bare glacial ice affects surface energy balance, bio-photochemical cycling, and light detection and ranging (LiDAR) laser elevation measurements but has not previously been reported for the Greenland Ice Sheet.

- 15 We present measurements of spectral transmittance at 350–900 nm in bare glacial ice collected at a site in the western Greenland ablation zone. Empirical irradiance attenuation coefficients at 350–750 nm are $\sim 0.9\text{--}8.0\text{ m}^{-1}$ for ice at 12–124 cm depth. Relative to theoretical values for optically pure ice, our empirical attenuation coefficients are nearly one order of magnitude larger at 350–530 nm. The estimated absorption coefficient at 400 nm suggests the ice volume contained an equivalent black carbon concentration of $\sim 1\text{--}2\text{ ng g}^{-1}$, similar to pre-industrial values found in remote polar snow. The equivalent mineral dust concentration is $\sim 300\text{--}600\text{ ng g}^{-1}$, similar to values for Northern Hemisphere warm periods with low aeolian activity inferred from ice core records. The absorption minimum is at $\sim 390\text{ nm}$, in agreement with snow transmission measurements in Antarctica and optical mapping of deep ice at the South Pole. For a layer of quasi-granular white ice extending from the surface to $\sim 10\text{ cm}$ depth, attenuation coefficients are 1.5 to 4 times larger than for interior bubbly ice. Owing to higher attenuation in this layer of near-surface granular ice, optical penetration depth at 532 nm is 14 cm (20%) lower than asymptotic attenuation lengths inferred from two-stream theory for optically pure bubbly ice. In addition to the traditional concept of light scattering on air bubbles, our results imply that the granular near-surface ice microstructure is an important control on radiative transfer in bare ice on the Greenland Ice Sheet ablation zone, and we provide new values of flux attenuation, absorption, and scattering coefficients to support model development and validation.
- 20
- 25
- 30

Deleted: attenuation

Deleted: bare ice suggest shallower subsurface radiative heating and optical penetration depth in the ablation zone

Deleted: We present in-ice solar irradiance

Deleted: measured over the spectral range 350–900 nm and

Deleted: 12–77 cm depth collected at

Deleted: The acquired spectral irradiance measurements are used to calculate flux attenuation coefficients

Deleted: using

Deleted: an exponential decay Bouguer law model and are compared to values calculated from

Deleted: two-stream radiative transfer theory

Deleted: -. Relative to asymptotic two-stream theory, our empirical attenuation coefficients are up to one order of magnitude larger in the range 350–530 nm

Deleted: Monte Carlo simulations indicate detector interference imparts a bias of +14–19% near the wavelength of minimum absorption. ...

Deleted: suggesting light absorbing particles embedded in ice enhance visible light absorption at our field site. The empirical coefficients accurately describe light attenuation in the ice interior but underestimate light attenuation near the ice surface. Consequently, Bouguer's law overestimates transmitted flux by up to 50% depending on wavelength. Refraction is unlikely to explain the discrepancy. Instead,

Deleted: vertical variation in the ice microstructure and the concentration of light absorbing particles appears to enhance near-surface attenuation at our field site. The magnitude of this near-surface attenuation implies that o

Deleted: at wavelengths relevant to visible-wavelength lidar altimetry of ice surface elevation (e.g. 532 nm for the Ice, Cloud, and Land Elevation Satellite-2) than is suggested by e-folding depths inferred from two stream theory for optically pure glacier ice. This enhanced near-surface attenuation implies shallower light transmission and therefore lower subsurface light availability for subsurface radiative heating and bio-photochemical cycling.

Deleted: We

Deleted: recommend radiative transfer models applied

Deleted: to

Deleted: in

Deleted: , and we

Deleted: values of

Deleted: .

1 Introduction

75 Understanding the transmission, absorption, and scattering of light in ice is important for snow and ice energy balance
modelling (Brandt and Warren, 1993), lidar remote sensing of snow surface elevation and grain size (Deems et al., 2013; Yang
et al., 2017), primary productivity beneath sea ice (Frey et al., 2011; Grenfell, 1979), bio-photochemical cycling in ice and
snow (France et al., 2011), and theoretical predictions of “Snowball Earth” paleoclimates (Dadic et al., 2013; Warren et al.,
2002). Each of these applications requires knowledge of the vertical distribution of light attenuation in ice, which for a medium
80 (such as glacier ice) that both absorbs and scatters light is specified by the spectral attenuation coefficient:

$$k_{\text{att}}(\lambda) = k_{\text{abs}}(\lambda) + k_{\text{sca}}(\lambda), \quad (1)$$

where k_{abs} (m^{-1}) is the absorption coefficient, k_{sca} (m^{-1}) is the scattering coefficient, and all are functions of wavelength, λ .
This study reports on the irradiance attenuation coefficient k_{att} of bare glacier ice in the Greenland Ice Sheet ablation zone, a
critical parameter needed to calculate subsurface absorption and scattering of transmitted radiation that to our knowledge has
85 received no direct field study.

Measurements of k_{att} in snowpack and sea ice indicate three main sources of variation with relevance to geophysical
applications. First, the magnitude of k_{att} is primarily controlled by ice microstructure (e.g., the size, shape, orientation, and
number of air bubbles, ice grains, and cracks), via its control on k_{sca} (Dadic et al., 2013; Libois et al., 2013; Light et al., 2004,
90 2008). For the range of air bubble sizes ($\sim 10^{-6}$) and ice grain sizes ($\sim 10^{-1}$ – 10^{-3}) observed in glacier ice, k_{sca} is effectively
independent of wavelength in the visible and near-infrared spectrum (Bohren, 1983; Dadic et al., 2013; Perovich, 1996).
Spectrally, k_{att} is low in the near-ultraviolet and blue-green, (~ 250 – 600 nm) where k_{abs} is extremely low ($< 10^{-8}$), and
progressively higher for wavelengths > 600 nm, where k_{abs} rapidly increases up to its maximum value ($\sim 10^{-2}$) at the far end of
the solar spectrum (Warren and Brandt, 2008). Vertically, k_{att} is at a maximum at the incident boundary (the snow or ice
95 surface) where a portion of upwelling radiation (i.e., transmitted flux reflected upwards) escapes the ice volume before
re-reflection downward. Within this near-surface optical boundary layer (Bohren and Barkstrom, 1974), attenuation rates
rapidly decrease with depth to an asymptotic value as multiple scattering establishes an isotropic (diffuse) radiation field
(Briegleb and Light, 2007; Warren, 1982). For fine-grained dry snow, a few cm depth is typically sufficient to reach the
asymptotic regime where k_{att} is constant (Brandt and Warren, 1993). For sea ice the depth required is typically larger and can
100 exceed > 20 cm depending on near-surface ice microstructure and the vertical location of the refractive boundary if present
(Grenfell, 1991; Grenfell and Maykut, 1977). Attenuation coefficients are also influenced by the horizontal distribution of ice
type and surface cover (Frey et al., 2011) but this source of variation is not examined here.

In addition to experimental values obtained from measurements of light transmission in ice or snow, k_{att} is obtained
105 analytically from optical theory (Bohren, 1987; Warren et al., 2006). Light attenuation in pure ice is specified analytically by
the complex refractive index;

Deleted: nearly

Deleted: spectral region

Deleted: significant

Deleted: of refraction

$$m(\lambda) = m_{\text{re}}(\lambda) - i m_{\text{im}}(\lambda), \quad (2)$$

where m_{re} is the real part of the complex refractive index (~1.31 in the visible), m_{im} is the imaginary part, and:

$$k_{\text{abs}}^{\text{ice}}(\lambda) = \frac{4\pi}{\lambda} m_{\text{im}}(\lambda) \quad (3)$$

is the absorption coefficient of pure ice (Warren et al., 2006; Warren and Brandt, 2008).

115

Light attenuation in glacier ice differs from pure ice owing to compositional and structural factors that control scattering and absorption, such as the size, geometry, and vertical distribution of embedded light absorbing particles (LAPs) and light scattering air bubbles and ice grains of size larger than wavelength (Askebjerg et al., 1997; Picard et al., 2016; Price and Bergström, 1997b; Warren et al., 2006). Analytical methods typically approximate ice and snowpack as homogeneous plane-parallel slabs of spheres having the same volume-to-surface-area ratio (i.e., optically-equivalent grain size) as the collection of non-spherical ice grains and air bubbles in realistic ice (Brandt and Warren, 1993; Grenfell and Warren, 1999; Wiscombe and Warren, 1980). Mie theory is used to calculate the single-scattering properties and two-stream radiative transfer approximations are used to calculate multiple scattering and bulk absorption in the ice volume (Bohren and Barkstrom, 1974; Mullen and Warren, 1988; Wiscombe and Warren, 1980). The single-scattering properties can also be derived from the ratio of surface area to mass (i.e., specific surface area) with or without the assumption of spherical scattering geometry (Kokhanovsky and Zege, 2004; Malinka, 2014), as applied to the highly scattering granular surface layer on sea ice (Malinka et al., 2016). Models of the prior form have been used to calculate subsurface meltwater production caused by penetration of solar radiation in ice both in Greenland (van den Broeke et al., 2008; Kuipers Munneke et al., 2009) and Antarctica (Brandt and Warren, 1993; Hoffman et al., 2014; Liston et al., 1999a, 1999b; Liston and Winther, 2005). However, theoretical values for k_{att} are rarely validated experimentally, and to our knowledge no such experimental values exist for near-surface glacier ice.

120

125

130

Deleted: >

Deleted: assume

Deleted: can be approximated

Deleted: for which

Deleted: Such m

135

140

In addition to ice surface energy balance, understanding light attenuation in ice is important for interpreting interactions between visible-wavelength light sources and ice surfaces, for example laser altimetry measurements of ice surface elevation (Deems et al., 2013; Gardner et al., 2015; Greeley et al., 2017). The reciprocal of k_{att} is the attenuation length, or the average distance travelled by a photon before attenuation by absorption or scattering (Ackermann et al., 2006). In the context of altimetry, the attenuation length is sometimes referred to as the penetration depth, or the average depth to which the electromagnetic signal penetrates before it is backscattered to the atmosphere (Ridley and Partington, 1988; Rignot et al., 2001; Zebker and Weber Hoen, 2000). The laser altimeter onboard Ice, Cloud, and Land Elevation Satellite-1 (ICESat-1) transmitted 1064 nm laser pulses to measure the distance (range) between the satellite and ice sheet surfaces (Schutz et al., 2005). Photons with wavelength 1064 nm penetrate snowpack no more than a few centimetres (Brandt and Warren, 1993; Järvinen and Leppäranta, 2013). This length scale is smaller than typical laser altimetry surface elevation errors due to ice and snow surface roughness and geolocation uncertainty (Deems et al., 2013). In contrast, the laser altimeter onboard ICESat-2 transmits 532

150 nm laser pulses (Markus et al., 2017). Ice is ~10 times more transparent at 532 nm than at 1064 nm (Warren and Brandt, 2008), and photons at 532 nm may penetrate many tens of centimetres into glacier ice. These subsurface scattered photons may introduce a range bias in ICESat-2 surface elevation retrievals over glacier ice, similar to radar penetration into snow (Brunt et al., 2016; Gardner et al., 2015; Greeley et al., 2017; Smith et al., 2018). To our knowledge no in situ observations of 532 nm optical penetration depth for bare glacier ice exist, precluding field validation of penetration depth obtained from theoretical radiative transfer models.

155 The purpose of this investigation is to provide experimental values for k_{att} obtained from measurements of solar flux attenuation in bare ice in the Greenland Ice Sheet ablation zone, and to compare them with theoretical values for k_{att} obtained from the two-stream analytical solution (c.f. Eq. 26 Bohren, 1987; Schuster, 1905). We benchmark our field estimates against the two-stream solution because of its wide use in surface energy balance models applied to snow and ice. In Sect. 2 we describe the field measurements and the optical theory used to interpret the solar flux attenuation. In Sect. 3 we report values for k_{att} obtained from our measurements, compare them with values obtained from two-stream theory, and propose a simple empirical model that accounts for enhanced near-surface attenuation. In Sect. 4 we discuss how our k_{att} values differ from prior experimental values acquired in sea ice, snowpack, and deep South Pole glacial ice, and the implication of these differences for modelling radiative transfer in bare glacier ice. To demonstrate the broader implications of our study, we suggest how our findings can be used to improve models for subsurface heating of ablating glacier ice.

2 Methods

2.1 Transmittance measurements

170 Ice transmittance was measured on 20 July and 21 July 2018 in the Kangerlussuaq sector of the western Greenland Ice Sheet. The study site is located ~1 km from the ice sheet margin at 840 m a.s.l. (67.15 °N, 50.02 °W). Subsurface (in-ice) spectral irradiance was measured at ~0.35 nm spectral resolution in the wavelength range 350–900 nm with an Ocean Optics® JAZ spectrometer. Light was guided from the ice interior to the spectrometer with a 3 mm diameter Kevlar-sheathed fibre optic cable fitted inside a 2 m long insulated white PVC tube (Fig. 1). The fibre was attached at one end to an irradiance sensor consisting of a 90° collimating lens adapter and a remote cosine receptor (RCR) with a Spectralon™ diffusing element. The RCR lens barrel was wrapped in white PTFE tape and set 2 mm out from the PVC tube exterior to act as a contact horizon between its diffusing element and the ice. The system was operated from a battery-powered computer running the Ocean Optics® OceanView software, placed on a tripod platform oriented 180° away from the sun and 2.5 m horizontal distance from the measurement location.

180 To access the interior of the ice, holes were drilled horizontally into a 2-m high sidewall of a natural ice feature with a battery powered hand drill fitted with a 3 cm diameter Kovacs auger bit. To drill these holes, the auger was advanced into the sidewall

Deleted: calibrated for absolute irradiance

Deleted: affixed

Deleted: diffuser

Deleted: . The computer and spectrometer were

Deleted: Each hole was drilled 2 m deep into the ice. Starting at the lowest hole near the bottom of the sidewall, t

approximately 20 cm, levelled horizontally with a digital spirit level, and the sequence was repeated to 2 m horizontal depth. The PVC tube-fibre optic assembly was then inserted into the hole, RCR facing upward, and a 2 m long ruler was shimmed under the bottom of the PVC tube to ensure the RCR barrel preserved contact with the overlying ice thus minimizing stray light contamination into the RCR field of view. Ice shavings were packed around the drill hole to prevent light reflection into the hole. Spectral irradiance was measured using a 20-scan average with 0.0228 s integration time per scan, yielding 0.46 s total integration time per irradiance measurement. Irradiance measurements were recorded at 1 Hz frequency for thirty seconds yielding thirty irradiance profiles at each depth, after which the tube was removed, the next hole was drilled, and the sequence was repeated, working from the top hole toward the bottom on 20 July, and from the bottom hole toward the top on 21 July.

The measurements were completed between 13:45 and 14:35 local time (UTC -3) on 20 July, and between 13:09 and 14:00 on 21 July, at solar zenith angles of ~48–51°. Solar noon at this time and location is ~13:26.

Deleted: and 44 Hz integration time

Background upwelling and downwelling spectral irradiances were measured continuously at 2 m height above the ice surface ~3 m away from the in-ice measurements with a dual-channel Ocean Optics JAZ spectrometer. These data were recorded at 1 min frequency using a 30-scan average with 0.011 s integration time. Light was guided to the spectrometer via two 3 m fibre optic cables attached to two RCRs mounted in upward-looking and downward-looking orientation on a 2 m long horizontally levelled boom attached to a vertical mast frozen into the ice. The horizontal boom became unstable on 21 July and the upward-looking RCR was moved to the vertical mast; the downward-looking RCR was decommissioned.

Deleted: approximately

Deleted: calibrated for absolute irradiance.

Deleted: 92 Hz

Deleted: drilled

The surface-based spectrometer was calibrated for absolute irradiance in a controlled setting prior to the field experiment using an Ocean Optics HL-3P radiometrically calibrated halogen light source. During the field experiment, the in-ice spectrometer was cross-calibrated to the surface spectrometer by holding it level above the ice surface in an upward-looking orientation ~3 m away from the surface spectrometer. Cross-calibration irradiance profiles were collected on 20 July and 21 July immediately prior to subsequent in-ice measurements. All in-ice irradiances are cross-calibrated to the surface spectrometer as a pre-processing step prior to further analysis.

Dark current spectra were recorded prior to each irradiance measurement as input to the OceanView automated dark current correction module. To measure dark current, the RCR lens barrel was capped with a custom-fit opaque metal cap provided by Ocean Optics. OceanView adjusts these spectra in real-time for changes in integration time and for charge leakage if detected, corrects the nonlinear analogue-to-digital response of the linear silicon charge coupled device, and applies a boxcar smoothing over adjacent pixels to further reduce noise. Following these automated corrections, the opaque cap was left in place and residual dark current (noise) was recorded with the reference spectrometer in its identical setup during the experiment and with the in-ice spectrometer held level above the ice surface in an upward-looking orientation. These residual dark current spectra are treated as systematic errors and are subtracted from all irradiance profiles as a pre-processing step prior to analysis (Fig. 2a).

2.2 Weather conditions

The 20 July experiment was conducted under low, thick cloud cover with light rain and no direct sun, ideal conditions for estimating the attenuation of diffuse light in ice. The 21 July experiment was conducted under higher, thinner cloud cover with no rain and very brief periods of intermittent direct sun (see Fig. 1). The effect of intermittent direct sun was easily identified in the in-ice irradiance measurements as a rapid increase in light intensity, which only occurred during the third measurement on 21 July. This was mitigated by averaging over the first ten in-ice irradiance profiles for that measurement, prior to the rapid increase in light intensity, and discarding the remainder.

2.3 Ice thickness and density

The ice thickness between detector positions was measured to the nearest millimetre with a metre stick and converted to units of solid ice thickness with the relation:

$$\Delta z = \Delta h \frac{\rho}{\rho_{ice}}, \quad (4)$$

where Δh is in-situ ice thickness between detector positions, ρ is in-situ ice density, ρ_{ice} is solid ice density (917 kg m^{-3}), and Δz is solid ice thickness between detector positions. Two separate observers made ten independent measurements of Δh . In addition, one observer made 41 replicate measurements of an ablation stake using the same metre stick, yielding a mean difference and standard error on Δh . The ice density ρ was measured on a 1.2 m ice core extracted at the measurement location with a Kovacs Mark IV corer (www.kovacsicedrillingequipment.com) (Fig. 3). The ice core was split along natural breaks into three segments that were measured to the nearest millimetre with a calliper and weighed to the nearest gram on an Acculab digital scale.

2.4 Experimental asymptotic flux attenuation coefficients and ice surface albedo

Spectral asymptotic flux attenuation coefficients are estimated by fitting a Bouguer-law exponential decay model to the in-ice irradiance profiles (Grenfell and Maykut, 1977):

$$I_z(\lambda) = I_0(\lambda) \exp[-k_{att}(\lambda)(z - z_0)], \quad (5)$$

where $k_{att}(\lambda)$ is the asymptotic flux attenuation coefficient, $I_z(\lambda)$ is in-ice spectral irradiance at depth z , $I_0(\lambda)$ is background downwelling spectral irradiance, z_0 is the ice surface, and:

$$T_z(\lambda) = I_z(\lambda)/I_0(\lambda) \quad (6)$$

is spectral transmittance. The optical depth $\tau_z(\lambda)$ is a dimensionless path length that scales the physical thickness of a layer by its attenuation rate:

$$\tau_z(\lambda) = -\ln T_z(\lambda) = k_{att}(\lambda)(z - z_0). \quad (7)$$

Estimates of $k_{att}(\lambda)$ for each spectral band are obtained by solving a linear equation of the form:

$$\tau_z(\lambda) = \tau_0(\lambda) + k_{att}(\lambda)(\Delta z + \epsilon_{\Delta z}) + \epsilon_{\Delta \tau}, \quad (8)$$

Deleted: The raw 0.35 nm spectra were interpolated to 1 nm using bilinear interpolation and smoothed with a centred moving mean filter with window size 3 nm.

Deleted: 1 nm

260 where τ_0 is a parameter (y-intercept), $\Delta z = z - z_0$ is ice thickness, $\varepsilon_{\Delta z}$ is an error term that represents ice thickness measurement uncertainty, and $\varepsilon_{\Delta\tau}$ is an error term that represents optical path measurement uncertainty. Equation 8 is solved by Maximum Likelihood Estimation (MLE), which gives an unbiased estimate of the slope when measurement errors are present in both the independent and dependent variables (see Sect. 2.9) (York et al., 2004).

265 The attenuation length $l_{\text{att}}(\lambda)$ is the inverse of $k_{\text{att}}(\lambda)$ and is analogous to the photon mean free path or transport length (Ackermann et al., 2006). It is equivalent to the path length in ice required to attenuate irradiance to 37% ($1/e$) of its incident intensity, i.e., the path length at which $T = 1/e$ and $\tau = 1$:

$$l_{\text{att}}(\lambda) = \frac{1}{k_{\text{att}}(\lambda)}. \quad (9)$$

The ice surface spectral albedo is the ratio of the upwelling spectral irradiance to the downwelling spectral irradiance:

270
$$\alpha(\lambda) = \frac{I \uparrow (\lambda)}{I \downarrow (\lambda)}, \quad (10)$$

and the broadband albedo is:

$$\alpha = \int_{\lambda_1}^{\lambda_2} \alpha(\lambda) I_0(\lambda) d\lambda / \int_{\lambda_1}^{\lambda_2} I_0(\lambda) d\lambda. \quad (11)$$

2.5 Asymptotic flux attenuation coefficients

Theoretical $k_{\text{att}}(\lambda)$ values are calculated using the asymptotic solution to the delta-Eddington two-stream radiative transfer approximation (Joseph et al., 1976; Schuster, 1905):

$$k_{\text{att}}(\lambda) = \frac{3}{4} \frac{Q_{\text{ext}}(\lambda)}{r_{\text{eff}}} \sqrt{3(1 - \bar{\omega}(\lambda))(1 - g(\lambda)\bar{\omega}(\lambda))}, \quad (12)$$

where $Q_{\text{ext}}(\lambda)$ is the extinction efficiency, r_{eff} is the effective scattering particle radius (m), $g(\lambda)$ is the average cosine of the scattering angle, also referred to as the asymmetry parameter, and $\bar{\omega}(\lambda)$ is the single-scattering albedo:

$$\bar{\omega}(\lambda) = \frac{\sigma_{\text{sca}}(\lambda)}{\sigma_{\text{att}}(\lambda)}, \quad (13)$$

280 where $\sigma_{\text{att}}(\lambda)$ and $\sigma_{\text{sca}}(\lambda)$ are the single-scattering attenuation coefficient (m^{-1}) and scattering coefficient (m^{-1}), respectively. Equation (12) describes light attenuation by multiple scattering and absorption in a homogeneous plane-parallel slab of absorbing spheres far from any boundaries (Bohren, 1987).

To estimate r_{eff} , Eq. (12) is inverted and solved by iteration for the value of r_{eff} that minimizes the difference between measured and calculated k_{att} at $\lambda = 600$ nm. This method assumes that all absorption at 600 nm is due to ice (Warren et al., 2006). If absorption was influenced by LAPs r_{eff} would be over-estimated. Values for $Q_{\text{ext}}(\lambda)$, $g(\lambda)$, and $\bar{\omega}(\lambda)$ are obtained

Deleted: 2.Theoretical

Deleted: given by

Deleted: ρ_i is ice sample density (kg m^{-3}), ρ_{ice} is pure ice density (917 kg m^{-3}),

Deleted: iteratively solved

from Mie scattering algorithms (Mätzler, 2002) provided as MATLAB® code with input $m(\lambda)$ from Warren and Brandt (2008). The Mie solutions at each wavelength are integrated over a Gaussian size distribution (N=1000) of scattering radii $\mathcal{N}(\mu_r = r_{\text{eff}}, \sigma_r = 0.15r_{\text{eff}})$ to eliminate ripples associated with Bessel function solutions to the Mie equations (Gardner and Sharp, 2010). The optimal r_{eff} value is 3.1 mm, which corresponds to a specific surface area of $1.05 \text{ m}^2 \text{ kg}^{-1}$. These values are used in all subsequent calculations.

2.6 Flux absorption coefficients

Warren et al. (2006) developed a method to estimate k_{abs} for pure ice (i.e. $k_{\text{abs}}^{\text{ice}}$) from measurements of flux attenuation in snow in Antarctica. The method relies on three assumptions: 1) the value of $k_{\text{abs}}^{\text{ice}}$ at the reference wavelength ($\lambda_0 = 600 \text{ nm}$) is known accurately, 2) the value of k_{att} at λ_0 is not affected by LAPs in the measured snow or ice, and 3) $\bar{\omega}(\lambda)$ varies so little as to be effectively independent of wavelength in the spectral range considered (here the near-UV and visible). Warren et al. (2006) verified the validity of these assumptions for the spectral range 350–600 nm and obtained the following relation (Eq. 15 of that paper) between flux attenuation and flux absorption:

$$\left[\frac{k_{\text{att}}(\lambda)}{k_{\text{att}}(\lambda_0)} \right]^2 \approx \left[\frac{k_{\text{abs}}(\lambda)}{k_{\text{abs}}(\lambda_0)} \right]. \quad (14)$$

Equation 14 assumes that k_{abs} is not affected by LAPs at the reference wavelength (600 nm) but the relation can be used to estimate k_{abs} at any other wavelength, including those where absorption is affected by LAPs. At those wavelengths, Eq. (14) will predict values for k_{abs} higher than $k_{\text{abs}}^{\text{ice}}$ if LAPs are present in the measured snow or ice volume, due to the influence of LAPs on k_{att} .

The inferred k_{abs} values can be related to a mass absorption cross-section (MAC) (Doherty et al., 2010):

$$k_{\text{abs}}(\lambda) = k_{\text{abs}}^{\text{ice}}(\lambda) + \beta c \rho_i, \quad (15)$$

where β is the spectral MAC ($\text{m}^2 \text{ kg}^{-1}$) and c is the mass-mixing ratio of LAPs in the ice volume (g LAPs g^{-1} ice). We exploit this to interpret differences between our theoretical and experimental values of k_{att} on the basis of differences between $k_{\text{abs}}^{\text{ice}}$ (Warren et al., 2006) and the k_{abs} values that we obtain for glacier ice from Eq. (14). To provide context, we use representative values of β for black carbon β_{BC} and insoluble mineral dust (hereafter ‘dust’) β_{dust} to estimate corresponding equivalent mass mixing ratios $c_{\text{eq,BC}}$ and $c_{\text{eq,dust}}$ (Di Mauro et al., 2017; Doherty et al., 2010). The ‘equivalent’ mass mixing ratio is the mass-mixing ratio of each LAP species required to explain the difference between $k_{\text{abs}}^{\text{ice}}$ and our inferred k_{abs} values at a reference wavelength, and follows a similar approach used to infer LAP absorption in snowpack (Tuzet et al., 2019). For β_{BC} , we use $6 \text{ m}^2 \text{ g}^{-1}$ as a representative MAC at 550 nm and an absorption Ångstrom exponent range 0.8–1.9 to scale this value to 400 nm (Doherty et al., 2010). For β_{dust} , we use $0.013 \text{ m}^2 \text{ g}^{-1}$ at 550 nm (Di Mauro et al., 2017) and an absorption Ångstrom

Deleted: requires

Deleted: Consequently, although not developed for this purpose, Eq. (X) provides a means to infer the influence of LAPs on measured flux attenuation by comparison with values of $k_{\text{abs,ice}}$ provided by Warren et al. (2006), which are compiled in Warren (2008).

exponent range 2–3 (Doherty et al., 2010). We note that these descriptive estimates provide context for discussion; actual LAP species concentrations were not measured.

2.7 Near surface effects

330 Equations 7 and 12 are applicable at distances far enough from the incident boundary (here the ice surface) that the radiation field is diffuse and k_{att} is constant with depth. Near the ice surface the radiation field is converted via multiple scattering from direct to diffuse flux, and attenuation may be enhanced by direct reflection, enhanced scattering and/or absorption by the granular near-surface ice microstructure, or specular reflection at the ice surface, depending on its roughness (Dadic et al., 2013; Light et al., 2008; Mullen and Warren, 1988). To account for non-diffuse near-surface attenuation, we define a piecewise optical depth:

$$\tau(\lambda) = \int_0^{z'} k'(\lambda) dz + \int_{z'}^z k_{\text{att}}(\lambda) dz, \quad (16)$$

where k' is an effective attenuation coefficient for the near-surface non-diffuse layer and z' is a depth chosen to partition this layer from the interior diffuse region. We estimate k' with a centred finite-difference form of Eq. (7):

$$k'(\lambda) = -\frac{1}{\Delta z'} \ln \left[\frac{I_{z'}(\lambda)}{I_0(\lambda)} \right]. \quad (17)$$

340 Here, $\Delta z' = 12$ cm and $I_{z'}$ is the 12 cm in-ice irradiance measured on 20 July. Accordingly, the asymptotic attenuation length (Eq. 9) is distinguished from an effective penetration depth d_λ to include the effect of near-surface attenuation. The attenuation length is the depth at which $\tau = 1$. Setting $\tau(\lambda) = 1$ in Eq. (16) and solving for z yields:

$$z = \frac{1 - \Delta z' [k'(\lambda) - k_{\text{att}}(\lambda)]}{k_{\text{att}}(\lambda)} = d_\lambda. \quad (18)$$

345 Equation 16 gives estimates of spectral transmittance that account for non-diffuse near-surface attenuation but relies on knowledge of k' , which is sensitive to the spectral composition and directional distribution of I_0 and the structure and composition of the near-surface ice (Grenfell and Maykut, 1977; Light et al., 2008). To generalize the magnitude of near-surface attenuation, we calculate the fraction of downwelling spectral irradiance that transmits the non-diffuse layer weighted by the downwelling spectral irradiance:

$$\chi_0 = \int_{\lambda_1}^{\lambda_2} I_0(\lambda) \exp[-k'_\lambda \Delta z'] d\lambda / \int_{\lambda_1}^{\lambda_2} I_0(\lambda) d\lambda. \quad (19)$$

350 The χ_0 parameter is analogous to the i_0 parameter introduced by Grenfell and Maykut (1977) to partition the fraction of solar irradiance absorbed in the upper 10 cm of sea ice, which they termed the “Surface Scattering Layer” (SSL), and the ice interior, in which radiation is exponentially absorbed at a constant rate:

Deleted: The $k_{\text{att}}(\lambda)$ values calculated using

Deleted: are

Deleted: is

Deleted: transmission of upward reflected light out of the ice volume before re-reflection downward (Briegleb and Light, 2007). Attenuation may also be enhanced by

$$i_0 = \int_{\lambda_1}^{\lambda_2} [1 - \alpha_\lambda] J_0(\lambda) \exp[-k'_\lambda \Delta z'] d\lambda / \int_{\lambda_1}^{\lambda_2} [1 - \alpha_\lambda] J_0(\lambda) d\lambda. \quad (20)$$

360 The i_0 parameter has been widely adopted in energy balance models of glaciers and sea ice to compute [subsurface flux divergence \(heating rates\)](#) when radiation penetration is considered important (Bintanja and Van Den Broeke, 1995; Hoffman et al., 2014; Holland et al., 2012). For example, the sea ice component of the Community Earth System Model (CESM) uses $i_0 = 70\%$ for the visible (200–700 nm) and $i_0 = 0\%$ for the infrared (700–5000 nm) (Briegleb and Light, 2007). The important distinction is that i_0 partitions the [absorbed flux](#) whereas χ_0 partitions the downward flux (Brandt and Warren, 1993). For both χ_0 and i_0 , we set $\Delta z'$ to 10 cm for consistency with prior work (Grenfell and Maykut, 1977; Light et al., 2008; Maykut and Untersteiner, 1971).

Deleted: 3

2.8 Monte Carlo simulations of detector interference

370 We developed a Monte Carlo radiative transfer model to estimate the effect of detector interference on measured irradiances and fitted k_{att} values, following methods developed to simulate light propagation in biological tissue, ocean waters, and sea ice (Leathers et al., 2004; Light et al., 2003; Wang et al., 1995). Photon scattering is specified by a Henyey-Greenstein scattering phase function with single-scattering properties $Q_{\text{ext}}(\lambda)$, $g(\lambda)$, and $\bar{\omega}(\lambda)$ inferred from our optical measurements (Sect. 2.5). A complete technical description is given in Appendix 2, where model accuracy is verified by comparison with benchmark solutions to the radiative transfer equation (van de Hulst, 1980).

375 In the Monte Carlo simulations, photons are launched from an irradiance sensor on a detector rod with dimensions identical to those reported in this study. In the ideal (baseline) simulation, photons originate from an isotropic point source and propagate through ice until they transmit the surface or are terminated by absorption. Detector interference is investigated by repeating the Monte Carlo with an ideal cosine source function describing the angular response to radiance of the RCR, and with a non-ideal (empirical) angular response function (Fig. 2), with and without scattering and absorption interference by the PVC detector rod. The detector rod albedo $\bar{\omega}_{\text{rod}} \approx 0.4$ is calculated from the absorption spectra of polyvinyl chloride (Zhang et al., 2020); scattering by the rod is assumed isotropic. The Monte Carlo is integrated over 10 000 interactions at nine wavelengths in 50 nm increments from 350 nm to 750 nm, allowing us to fit the wavelength dependence of the estimated systematic uncertainty in simulated k_{att} values.

2.9 Uncertainty propagation

385 Unless stated otherwise, all statistical uncertainties reported in this paper are standard errors that correspond to 68% confidence intervals around the mean (Taylor and Kuyatt, 1994). For an individual measurement with standard deviation s_i and sample size $N \geq 30$ the standard error is s_i / \sqrt{N} . For $N < 30$, standard errors are scaled by a critical t-value drawn from the Student's

t-distribution. Standard errors for combined quantities are propagated in quadrature and hereafter referred to as combined uncertainty. The combined uncertainty for spectral irradiance $I(\lambda)$ is:

$$\sigma_I = \sqrt{(\sigma_I^*)^2 + (\sigma_D)^2}, \quad (21)$$

where σ_I^* is the standard deviation of the high-frequency irradiance spectra before dark-noise correction and σ_D is the standard deviation of the high-frequency dark-noise spectra. An analogous procedure is used to estimate the combined uncertainty for calibrated irradiance. The combined uncertainty for spectral transmittance $T(\lambda)$ is:

$$\sigma_T = T \sqrt{\left(\frac{\sigma_{I_z}^{\text{cal}}}{I_z^{\text{cal}}}\right)^2 + \left(\frac{\sigma_{I_0}}{I_0}\right)^2}, \quad (22)$$

where $\sigma_{I_z}^{\text{cal}}$ and σ_{I_0} are the combined uncertainties for calibrated in-ice irradiance and dark-noise corrected surface downwelling irradiance, respectively. The combined uncertainty for optical depth τ_λ is:

$$\sigma_\tau = \frac{\sigma_T}{T}, \quad (23)$$

and the combined uncertainty for k_{att} is:

$$\sigma_k = \sqrt{(\sigma_{\Delta\tau})^2 + (\sigma_{\Delta z})^2}. \quad (24)$$

Equation 24 gives a first order description of σ_k due to statistical propagation of measurement uncertainties, neglecting higher-order interaction terms. A description of the statistical uncertainty in fitted $k_{\text{att}}(\lambda)$ values is given by the MLE estimate of the regression slope of Eq. 8, which can be expressed in terms of an error model as:

$$\tau_z(\lambda) = \tau_0(\lambda) + k_{\text{att}}(\lambda) (\Delta z + \varepsilon_{\Delta z}) + \varepsilon_{\Delta\tau},$$

where τ and Δz are the true but unobserved (due to measurement error) optical depth and ice thickness and $\varepsilon_{\Delta z} \sim \mathcal{N}(0, \sigma_{\Delta z}^2)$ and $\varepsilon_{\Delta\tau} \sim \mathcal{N}(0, \sigma_{\Delta\tau}^2)$ are normally distributed error terms. Unlike Ordinary Least Squares, MLE gives an unbiased estimate of the slope and standard error of a linear functional relationship between two variables measured with error (York et al., 2004). The method has been used in similar studies to infer optical coefficients (Zieger et al., 2011). The MLE standard errors for $k_{\text{att}}(\lambda)$ are adjusted for $N - 2$ degrees of freedom with a two-sided t-statistic (Cantrell, 2008) and combined in quadrature with systematic uncertainty estimated from Monte Carlo simulation to estimate total combined uncertainty for reported $k_{\text{att}}(\lambda)$ values.

3 Results

3.1 Spectral transmittance

Four in-ice irradiance spectra were collected at 12 cm, 36 cm, 58 cm, and 77 cm depth below the ice surface on 20 July (hereafter referred to as Layer A) (Fig. 4a), and at 53 cm, 67 cm, 82 cm, and 124 cm on 21 July (hereafter referred to as Layer B). At all depths, spectral transmittance T is maximum at 430 nm and maintains relatively stable and high values up to ~500 nm in the visible, beyond which T decreases into the red end of the visible spectrum following the well-known exponential

increase in ice absorptivity (Fig. 4c). Maximum T values vary from 78% at 12 cm to 45% at 77 cm. For wavelengths >500 nm, T rapidly decreases both with wavelength and with depth; beyond ~800 nm nearly all incident light is attenuated within 36 cm of the ice surface, although substantial attenuation is apparent in the 12–36 cm depth region (Fig. 4b). The standard deviation of the 1 Hz raw data is $<1 \text{ W m}^{-2} \text{ nm}^{-1}$ at all wavelengths, consistent with field observations of thick cloud cover and diffuse light conditions described in Sect. 2.6. Instrumental noise and high-frequency measurement variations propagate as $\pm 1.6\%$ uncertainty on T for wavelengths between 400–600 nm, ± 1 –8% for wavelengths between 350–400 nm, where instrumental noise is higher, and ± 1 –12% uncertainty for wavelengths between 600–750 nm, where noise is higher and light levels are low.

3.2 Experimental flux attenuation coefficients and albedo

Fitted k_{att} values for Layer A range from $0.98 \pm 0.17 \text{ m}^{-1}$ to $7.86 \pm 0.43 \text{ m}^{-1}$ for wavelengths between 350 and 750 nm (Fig. 5a), with uncertainty bounds that represent statistical and systematic uncertainty summed in quadrature (see Sect. 4 for a discussion of systematic error). These values correspond to attenuation lengths of $1.02 \pm 0.18 \text{ m}$ to $0.13 \pm 0.007 \text{ m}$, respectively. Layer B k_{att} values are ~12% lower than Layer A values at 350–500 nm and within 1% at 650 nm (see inset Fig. 6). For Layer A, the minimum in k_{att} is at 390 nm, blue-shifted relative to the maximum in T at 430 nm. For Layer B, the minimum is at 397 nm. The coefficient of determination (r^2) ranged from 0.96–1.0 ($p < 0.01$), with a median value of 0.98, suggesting the data are described appropriately by the Bouguer-law exponential decay model up to ~700 nm, beyond which measured values of in-ice irradiance at 58 cm and 77 cm depth were too low to reliably fit k_{att} values (see Fig. 4b and Fig. 5c). For Layer B values, low light levels prevented fits beyond ~650 nm.

Albedo spectra correspond closely to patterns in transmittance and attenuation (Fig. 5c). The near-UV and blue wavelengths that efficiently transmit ice mostly re-emerge as reflected light, owing to the extremely low values of ice absorptivity in the wavelength range 350–500 nm where albedo is maximum (Gardner and Sharp, 2010; He and Flanner, 2020; Warren et al., 2006). The maximum measured albedo value is 0.81 ± 0.004 at 452 nm, further red shifted from the minimum in k_{att} and the maximum in T . All three quantities have low variability near the minimum; albedo is 0.79 at 390 nm. The broadband albedo α (Eq. 11) for the 350–900 nm wavelength range is 0.70 ± 0.006 , which is high but not atypical for melting white ice under overcast skies (Bøggild et al., 2010).

3.3 Theoretical flux attenuation coefficients

Asymptotic k_{att} values predicted by two-stream theory for optically-clean bubbly ice are nearly one order of magnitude lower than field estimates for wavelengths <500 nm, where very small concentrations of LAPs in the measured ice volume dominate absorption (compare dotted grey line to solid blue line, Fig. 6) (Warren et al., 2006). In contrast, field-estimates and two-stream theory converge at wavelengths >540 nm, where absorption is dominated by grain-size effects (He et al., 2017; Libois et al.,

Deleted: by ice is higher

Deleted: For $\lambda > 500 \text{ nm}$

Deleted: below

Deleted: , with $T < 2\%$ at 36 cm and $< 0.6\%$ at 58 and 77 cm depth.

Deleted: In contrast, T at 12 cm decreases from 18% at 800 nm to 5% at 900 nm, suggesting

Deleted: value (0.96 m^{-1}) occurs

Deleted: Theoretical

Deleted: the

Deleted: solution

Deleted: This discrepancy can be inferred to relate to the presence of LAPs embedded in the ice matrix that increase the effective absorption of the ice volume. For example, differences between the field-estimate of k_{abs} and $k_{\text{abs,ice}}$ (Fig. 5b) mirror those between the field-estimate and theoretical estimate of k_{att} (Fig. 5a). Namely, k_{abs} is nearly one order of magnitude larger than $k_{\text{abs,ice}}$ in the range 350–500 nm, where even very small concentrations of LAPs in the measured ice volume would dominate absorption (Warren et al., 2006). ...

Deleted: solution and

Deleted: $\lambda > 530 \text{ nm}$

470 2013). The magnitude of inferred absorption enhancement in the visible due to LAPs (the quantity $\beta c \rho_i$ in Eq. 15) varies from
 0.009–0.015 m^{-1} at 350–530 nm. The equivalent black carbon concentration c_{eqBC} inferred at 400 nm is 1–2 ng g^{-1} for both
 Layer A and Layer B, where the range covers uncertainty in both the absorption spectra and the absorption Ångstrom exponent
 (Doherty et al., 2010). The equivalent mineral dust concentration c_{eqdust} is ~344–620 ng g^{-1} for Layer A and 303–545 ng g^{-1}
 for Layer B. Monte Carlo simulations without detector interference replicate both asymptotic theory for clean bubbly ice (i.e.,
 475 when forced with $k_{\text{abs}}^{\text{ice}}$) and field estimates when forced with k_{abs} values inferred from our optical measurements (solid blues
 squares, Fig. 6). Monte Carlo simulations of detector interference are discussed further in Sect. 4.

3.4 Near-surface attenuation and effective penetration depth

Near the ice surface irradiance is not attenuated exponentially and Bouguer’s law does not hold, as indicated by the
 y-intercepts of the straight lines in Fig. 5b at values <100%. Effective k' values (Eq. 17) for the quasi-granular 0–12 cm
 480 layer are ~1.5 times higher than k_{att} values for interior bubbly ice at 12–77 cm depth for wavelengths >570 nm, and are up to
 4 times higher between 400–570 nm (Fig. 7). Owing to higher near-surface attenuation, transmitted irradiance I_z is
 overestimated by 10–60% if Bouguer’s law is applied to the incident downwelling irradiance I_0 using asymptotic k_{att} values,
 with median over-estimation 23% (Fig. 8a). In contrast, the piecewise optical depth (Eq. 16) predicts I_z to within 12% of
 measured values for all wavelengths between 350–700 nm with median error 3%. Integrated over these wavelengths, χ_0 is
 485 0.68 and i_0 is 0.66, suggesting 66% of the total incoming irradiance within this spectral region is absorbed at depths below
 10 cm. If k_{att} is used rather than k' to calculate χ_0 and i_0 , the respective values are 0.81 and 0.79.

Stated in terms of penetration depth, d_{eff} varies from 12–84 cm between 350–700 nm. These values are 13–44% lower than
 attenuation lengths l_{att} inferred from empirical asymptotic k_{att} values. Specifically at 532 nm, d_{eff} is 52 cm, or 10 cm lower
 490 than the 62 cm empirical l_{att} value, and 14 cm lower than the 66 cm theoretical l_{att} value for optically pure bubbly ice. These
 results point to the potential for reduced optical penetration due to enhanced scattering and absorption on or near the ice surface,
 as well as within the ice volume where small LAP concentrations reduce optical backscattering due to enhanced absorption.

For smooth ice surfaces, attenuation may be enhanced by refraction at the ice-air interface (Mullen and Warren, 1988). If
 495 present, a refractive boundary would enhance near-surface attenuation via external specular reflection, and possibly via
 enhanced near-surface absorption of the internally reflected downward flux. Following Briegleb and Light (2007), Eq. 20–24,
 we calculate the external diffuse specular reflectivity for a flat ice surface to be 0.063, meaning specular reflection could
 enhance attenuation by up to 6.3%. This value is smaller than the 18–44% near-surface attenuation implied by the y-intercepts
 in Fig. 5b, suggesting specular reflection alone cannot explain the discrepancy. Instead, we suggest that enhanced scattering
 500 by the granular near-surface ice microstructure, together with absorptive impurities, enhance near-surface light attenuation at

Deleted: confirmed

Deleted: This suggests k_{att} values are higher in the 0–12 cm near-surface region where irradiance measurements were not obtained. Consequently, transmitted irradiance is overestimated by 5–50% if Bouguer’s law is applied to the incident surface irradiance using k_{att} values from the 12–77 cm region, with median over-estimation 16% (Fig. 6a). The value of χ (Eq.) that minimizes the root-mean-squared-difference between measured and predicted transmitted irradiance, weighted equally at all depths and all λ , is 15%. Transmitted irradiance spectra predicted using Eq. () with $\chi = 15\%$ are shown in Fig. 6c. ↓

Expressed in terms of attenuation rate, e

Deleted: region

Deleted: estimated from a finite-difference solution to Eq. 17)

Deleted: the

Deleted: region

Deleted: $\lambda > 570$ nm,

Deleted: attenuation length, l

Deleted: These values are calculated by combining the effective k_{att} values for the 0–12 cm region with the k_{att} values for the 12–77 cm region and therefore correspond to effective penetration depths. Effective penetration depths are smaller than

Deleted: for the 12–77 cm region (i.e., from Eq.)

Deleted: effective penetration depth

Deleted: in the 12–77 cm region

Deleted: light that escapes the ice volume.

Deleted: 10

Deleted: 25

our field site, consistent with observations of the granular and porous surface layer on sea ice (Grenfell and Maykut, 1977; Light et al., 2008).

4 Uncertainty analysis

535 The effect of random and systematic uncertainties on our optical measurements and fitted k_{att} values is evaluated with Monte Carlo simulation and statistical analysis. We considered systematic uncertainties in detector positions, spectrometer sensitivity to dark current, the non-ideal angular response of the irradiance sensor, and attenuation interference by the PVC detector rod.

540 The detector positions are known to within 0.9 ± 0.4 cm from independent measurements of the vertical ice thickness Δh . The in-situ ice density ρ varied from $801\text{--}888$ kg m⁻³ between $4\text{--}124$ cm where irradiances were measured. The variation in ρ was examined by repeating the analysis with Δz values computed with a single depth-weighted average ρ applied to each Δh , and with ρ values estimated for each Δh from linear and cubic interpolation of the vertical density profile. The maximum Δz difference was 0.9 cm. The k_{att} values differed by $<1\%$, and r^2 values were nearly identical. We use the depth-weighted average ρ values to calculate Δz , which are 835 kg m⁻³ and 855 kg m⁻³ for the measurements collected on 20 July and 21 July, respectively.

550 Detector position uncertainty was further assessed by fitting k_{att} values with an ensemble of 10 000 Δz values perturbed with random errors drawn from a normal distribution $\mathcal{N}(\mu = 0.9 \text{ cm}, \sigma = 0.4 \text{ cm})$. At all wavelengths, the chance of obtaining a fitted k_{att} value $>2\%$ from the mean value was $<5\%$. We take 2% as a conservative estimate of systematic uncertainty due to ice thickness measurement bias.

555 As described in Sect. 3, all irradiance spectra are corrected for residual dark noise. The noise may have varied during the experiment, and dark noise measurements with the in-ice spectrometer were made on the surface, rather than within the ice. To assess possible bias, we fit k_{att} values with and without residual dark noise correction. The mean difference was $-0.01 \pm 0.13\%$ averaged over the $350\text{--}700$ nm wavelength range. For a few discrete wavelengths between $350\text{--}400$ nm and $700\text{--}750$ nm, differences approached 2% . These wavelengths are those with the highest dark noise in the reference spectrometer (Fig. 2). At wavelengths between $400\text{--}700$ nm, differences were $<0.5\%$.

560 Monte Carlo simulations indicate a possible $+1\text{--}14\%$ systematic bias due to detector interference for Layer A values, and $+2\text{--}8\%$ for Layer B values (Fig. 9; also see purple stars minus solid squares, Fig. 6). The high end of this range applies to the wavelength region of minimum absorption $\sim 350\text{--}450$ nm. The simulated bias is within statistical uncertainty at wavelengths >450 nm for Layer A and at wavelengths >400 nm for Layer B (Fig. 9). The non-ideal cosine response of the RCR and the presence of the detector rod both tend to increase k_{att} values relative to the ideal case, as expected given the low albedo of the

565 detector rod. However, detector interference is masked somewhat by the presence of LAPs, as indicated by the larger simulated
570 interference for bubbly ice without LAPs (see dotted grey line and associated Monte Carlo values, Fig. 6). Overall, the
combined statistical and systematic uncertainty for the 350–450 nm region is <20% for Layer A values and <14% for Layer B
values, and as low as ~5% for wavelengths >450 nm (Fig. 9).

5 Discussion

5.1 Comparison with attenuation spectra for sea ice, snowpack, and deep glacial ice

570 We report spectral measurements of near-UV and visible light transmission in bare ablating glacier ice. These measurements
are used to estimate irradiance attenuation coefficients k_{att} for the spectral range 350–750 nm. Prior studies quantified k_{att} for
sea ice and snowpack (c.f. Fisher et al., 2005; Frey et al., 2011; Gerland et al., 2000; Grenfell and Maykut, 1977; Järvinen and
Leppäranta, 2013; King and Simpson, 2001; Light et al., 2008; Meirold-Mautner and Lehning, 2004; Pegau and Zaneveld,
2000; Picard et al., 2016; Tuzet et al., 2019; Warren et al., 2006). Scattering and absorption coefficients were quantified for
575 compressed South Pole glacial ice at 800–2350 m depth by the AMANDA (Antarctic Muon and Neutrino Detector Array)
experiment (Ackermann et al., 2006; Askebjerg et al., 1995, 1997). For South Pole ice at 800–1000 m depth, visible and near-UV
light scatters on air bubbles, below which bubbles transition under pressure to clathrates and light scatters on dust grains (Price
and Bergström, 1997b). In the bubbly ice regime studied by AMANDA, k_{sc} values at 532 nm are ~1–3 m⁻¹, comparable to
the 1.6 m⁻¹ value quantified in this study. Light scattering in the dusty-ice regime (>1000 m depth) is not comparable to this
580 study; absorption by dust is discussed below.

Fig. 10 compares our k_{att} spectra for glacier ice to seven previously published spectra for snowpack and sea ice. In general,
glacier ice is the most transparent structure examined with the exception of multi-year and first-year interior sea ice at
wavelengths >540 nm (Grenfell et al., 2006). Light attenuation in sea ice is controlled by its unique vertical composition
585 including brine inclusions, air pockets, solid salts, sea ice algae, dissolved organic matter, water saturation, and radiative
interactions between the ice and underlying ocean (Perovich, 1996). The latter factor, together with differences in
optically-equivalent grain size, may explain the low attenuation at longer wavelengths for sea ice shown here. Relative to
snowpack in Greenland and Antarctica (Järvinen and Leppäranta, 2013; Meirold-Mautner and Lehning, 2004; Warren et al.,
2006), attenuation by glacial ice has similar spectral structure but is lower at all wavelengths, reflecting the higher specific
590 surface area of fine-grained polar snow. Attenuation within the surface scattering layer (SSL) of sea ice is intermediate, with
spectral structure similar to snowpack and glacial ice. Attenuation at 5 cm depth in snow near Summit, Greenland is highest
of all, possibly due to direct light scattering in the near-surface optical boundary layer. The comparison demonstrates that k_{att}
values vary by nearly two orders of magnitude at visible wavelengths due to differences in ice structure and composition.

Deleted: first

Deleted: asymptotic flux

Deleted: 700

Deleted: and for

Deleted: for which optical scattering is not representative of near-surface ablating glacier ice

Deleted: observed in

Deleted: Relative to prior measurements in sea ice (Grenfell et al., 2006; Grenfell and Maykut, 1977), our results suggest light attenuation by glacial ice is lower at blue-green wavelengths and higher at orange-red wavelengths, likely reflecting differences in the absorption spectra of light absorbing material found in sea ice relative to that found in glacier ice.

Deleted: likely due to

Deleted: enhanced scattering from the

Deleted: structure of

Deleted: likely

Deleted: the tremendous variation in

Deleted: caused by

Deleted: 4.2 Relevance to surface energy balance modelling and subsurface meltwater production*

At visible wavelengths between 350–530 nm our field estimates of k_{att} are up to one order of magnitude larger than those obtained from two-stream theory for optically pure bubbly ice, consistent with selective absorption by mineral dust, black carbon, and microorganisms found on glaciers and ice sheet surfaces (Bøggild et al., 2010; Ryan et al., 2018; Stibal et al., 2017; Takeuchi, 2002; Yallop et al., 2012). For context, the absorptivity we document at 400 nm for Layer B can be explained by 1.2–1.8 ng g⁻¹ (ppb) equivalent black carbon concentration. Values for Layer A are 1.4–2.0 ppb. Both estimates are relative to pure ice absorptivity values reported by Warren et al. (2006). These values are within the range 2±2 ppb reported for clean snow near Dye-2 on the interior Greenland Ice Sheet considered representative of pre-industrial fallout rates (Doherty et al., 2010). The equivalent mineral dust concentration is ~344–620 ppb for Layer A and 303–545 ppb for Layer B.

Deleted: clean

Relative to South Pole ice, our absorptivity values broadly agree with AMANDA values within two depth regions corresponding to peaks in atmospheric dust concentration during the Last Glacial Maximum (LGM) and Marine Isotope Stage 4 (MIS-4) glacial periods ~23 000 and ~66 000 years before present (Fig. 11). For these periods in Earth's history, Southern Hemisphere dust concentrations inferred from the Vostok and Dome-C ice cores are ~300–1500 ppb (Muhs, 2013; Petit et al., 1999). Hemispherical dust fluxes are generally synchronous at these timescales; similar peaks at LGM and MIS-4 are observed in Greenland ice cores (Ruth et al., 2003). However, Northern Hemisphere dust concentrations are several times higher (Muhs, 2013; Ruth et al., 2003), meaning correlation with South Pole absorptivity does not map age at our site. Rather, our optical measurements are consistent with the relatively low dust concentrations during Northern Hemisphere warm periods. For the western Greenland ablation zone, alternating bands of visibly dark and bright outcropping ice are associated with periods of higher and lower aeolian activity during both the early Holocene (post-LGM) and late Pleistocene, with a characteristic band of older brighter interglacial ice ~0.7–1 km from the margin where our field site is located (Bøggild et al., 2010; Petrenko et al., 2006; Reeh et al., 2002; Wientjes et al., 2012). Taken together, this suggests the optical properties documented here are representative of Pleistocene interglacial ice with relatively low volumetric LAP concentration and smaller crystal diameters than Holocene ice associated with the 'dark zone' further inland (Gow et al., 1997; Petrenko et al., 2006; Wientjes et al., 2011).

Deleted: Comparison with the spectral absorption coefficient of pure ice (Fig. 6c) suggests the discrepancy we find is likely due to LAPs present in the measured ice volume, which appear to disproportionately enhance energy absorption near the ice surface.

Regarding pure ice absorptivity, our k_{abs} values support the lower bound pure ice estimate from Warren et al. (2006) (Fig. 12). The steeply sloping high values in the near-UV in the laboratory measurements (Grenfell and Perovich, 1981; Perovich and Govoni, 1991) are now understood as signatures of Rayleigh scattering on nanoscale defects in the laboratory-grown ice (Price and Bergström, 1997a). The South Pole values at 1755 m depth and 830 m depth are contaminated by trace dust deposited during the late Pleistocene and early Holocene, respectively (Ackermann et al., 2006) (Fig. 11). The lowest values reported by Warren et al. (2006) (hereafter W06) were obtained by applying Eq. 7 to measurements of transmitted radiance in a single snow layer at ~90–135 cm depth near Dome C in Antarctica contaminated by ~0.6 ppb black carbon (75°S, 123°E, 3230 m). Picard et al. (2016) (hereafter P16) repeated the W06 experiment on 56 transmitted radiance profiles collected in snow near Dome C with variable impurity content. The values shown in Fig. 12 are their best estimate of pure ice absorptivity from radiance profiles collected in snow with low impurity content (see 'clean' subset, Fig. 17 of that paper). P16 was unable

655 to reconcile their values with W06, after considering published values for impurity loadings in the vicinity of Dome C, suggesting the W06 values were unreasonably low. Regardless of that discrepancy, our values were undoubtedly influenced by LAPs but are lower than the P16 values. Treating our Monte Carlo estimate of detector interference as a known systematic error would bring our values closer to the AMANDA and W06 lower bound and further from the P16 values (see dotted line, Fig. 12).

660 Our absorption minimum is at 390 nm for Layer A values and 397 nm for Layer B values, in agreement with W06 and AMANDA. The wavelength shift in the P16 absorption minimum (430 nm) is apparent in all attenuation coefficient spectra shown in Fig. 10 that use the surface as a reference horizon, but is absent in those that use an interior layer as a reference horizon. P16 used the latter method and excluded radiance measurements within 8 cm of the surface based on Monte Carlo simulations of detector interference and visual inspection of homogeneous attenuation rates, but their shifted minimum may indicate that radiance profiles in the near-UV and blue were impacted disproportionately by detector rod interference. The same can be said of our values, and may explain in part the spectral structure in our near-surface (0–12 cm) effective attenuation coefficient profile between 350–400 nm (dotted line Fig. 10). Similar spectral structure is apparent in diffuse attenuation coefficients obtained in snowpack in the French Alps using the same method as P16 (c.f. Fig. 3b, Tuzet et al., 2019). Differences aside, our inferred absorption spectrum provides new insight into the magnitude of this fundamental but uncertain optical property, and supports a conclusion that the minimum is likely $<10^{-2} \text{ m}^{-1}$ and possibly lower (Ackermann et al., 2006; Picard et al., 2016; Warren et al., 2006).

675 In addition to the traditional concept of surface melt, visible light transmission provides an energy source for subsurface heating and internal melting of near-surface glacier ice (Cooper et al., 2018; Hoffman et al., 2014; Liston and Winther, 2005; Schuster, 2001). Prior estimates of subsurface meltwater production in bare ice used two-stream theory forced with values of k_{abs} for pure ice to calculate k_{att} and the absorbed solar flux as a function of depth below the ice surface in both Greenland and Antarctica (van den Broeke et al., 2008; Hoffman et al., 2014; Kuipers Munneke et al., 2009; Liston and Winther, 2005). The influence of LAPs on subsurface meltwater production has not been quantified to our knowledge and is beyond our scope, but our results suggest LAPs enhance subsurface energy absorption in ablating glacier ice, consistent with enhanced surface melt rates caused by LAPs distributed on bare ice surfaces and within snowpack (Boggild et al., 1996; Goelles et al., 2015; Goelles and Boggild, 2017; Tuzet et al., 2019). From a practical perspective, this suggests that k_{abs} values for contaminated ice given here and snowpack given elsewhere (Picard et al., 2016) could provide realistic input for radiative transfer models absent explicit knowledge of realistic LAP concentrations. In contrast, simulations that use the canonical pure ice absorptivity values compiled in Warren and Brandt (2008) will likely underestimate light attenuation and misrepresent the distribution of subsurface absorbed flux unless LAP concentrations are otherwise accounted for.

Deleted: This is important because v

Deleted: in the ablation zones of glaciers and ice sheets

Deleted: To our knowledge, t

690 **5 Conclusion**

We report in-situ spectral measurements of near-UV and visible light transmission in near-surface bare glacial ice, collected at a site in the western Greenland ablation zone during July 2018. In general, our empirical irradiance attenuation coefficients are nearly one order of magnitude larger in the range 350–530 nm than predicted by asymptotic two-stream theory using canonical values for the absorption coefficient of pure ice (Warren and Brandt, 2008). The absorption minimum is 0.013–
695 $0.014 \pm 0.003 \text{ m}^{-1}$ at 390–397 nm implying absorption length scales of 69–77 m. The volumetric scattering coefficient is $1.6 \pm 0.2 \text{ m}^{-1}$ at 532 nm, with an asymptotic attenuation length scale $0.62 \pm 0.08 \text{ m}$. In addition to light scattering on air bubbles, we find that light attenuation is enhanced by a layer of quasi-granular white ice that extends from the surface to ~10 cm depth at our field site. The effective penetration depth, which accounts for reduced optical transmission through this granular layer relative to interior bubbly ice, is $0.52 \pm 0.07 \text{ m}$ at 532 nm. Our co-located measurements of transmittance and albedo suggest
700 that about 34% of cloudy sky downwelling solar irradiance at 350–700 nm is absorbed within 10 cm of the ice surface at this time and location, consistent with observations of the semi-granular surface layer on sea ice. The estimated absorption spectrum suggests equivalent black carbon and mineral dust concentrations consistent with pre-industrial periods in Earth’s history with low Northern Hemisphere aeolian activity, and therefore may provide a reasonable lower bound on volumetric absorption enhancement due to impurities embedded in outcropping glacial ice in the western Greenland ablation zone.

Deleted: first

705 **Data availability**

The data is hosted by the Pangaea open access data repository (<https://doi.pangaea.de/10.1594/PANGAEA.913508>) and is provided as a supplement to this manuscript in Appendix 2.

Author contribution

M.G. Cooper and L.C. Smith designed the experiment. M.G. Cooper, A.K. Rennermalm, M. Tedesco, R. Muthyala, S.Z.
710 Liedman, and S.E. Moustafa collected the field data. M.G. Cooper performed the data analysis and wrote the manuscript. L.C. Smith, A.K. Rennermalm, M. Tedesco, S.Z. Liedman, and J.V. Fayne edited the manuscript. The authors declare they have no conflict of interest.

Acknowledgements

This project was funded by the NASA Cryosphere Program grant 80NSSC19K0942, managed by Dr. Colene Haffke and Dr.
715 Thomas P. Wagner and a graduate fellowship from the NASA Earth and Space Sciences Fellowship Program managed by Dr. Lin Chambers. We thank Polar Field Services for their field support. The authors thank S. G. Warren, R. E. Brandt, and G. Picard for advice on the experimental design.

References

- 720 Ackermann, M., Ahrens, J., Bai, X., Bartelt, M., Barwick, S. W., Bay, R. C., Becka, T., Becker, J. K., Becker, K.-H., Berghaus, P., Bernardini, E., Bertrand, D., Boersma, D. J., Böser, S., Botner, O., Bouchta, A., Bouhali, O., Burgess, C., Burgess, T., Castermans, T., Chirkin, D., Collin, B., Conrad, J., Cooley, J., Cowen, D. F., Davour, A., Clercq, C. D., Heros, C. P. de los, Desiati, P., DeYoung, T., Ekström, P., Feser, T., Gaißer, T. K., Ganugapati, R., Geenen, H., Gerhardt, L., Goldschmidt, A., Groß, A., Hallgren, A., Halzen, F., Hanson, K., Hardtke, D. H., Harenberg, T., Hauschildt, T., Helbing, K., Hellwig, M., Herquet, P., Hill, G. C., Hodges, J., Hubert, D., Hughey, B., Hulth, P. O., Hultqvist, K., Hundertmark, S., Jacobsen, J., Kampert, K. H., Karle, A., Kestel, M., Kohnen, G., Köpke, L., Kowalski, M., Kuehn, K., Lang, R., Leich, H., Leuthold, M., Liubarsky, I., Lundberg, J., Madsen, J., Marciniwski, P., Matis, H. S., McParland, C. P., Messarius, T., Minaeva, Y., Miočinović, P., Morse, R., München, K., Nahnhauser, R., Nam, J. W., Neunhöffer, T., Niessen, P., Nygren, D. R., Olbrechts, P., Pohl, A. C., Porrata, R., Price, P. B., Przybylski, G. T., Rawlins, K., Resconi, E., Rhode, W., Ribordy, M., Richter, S., Martino, J. R., Sander, H.-G., Schlenstedt, S., Schneider, D., Schwarz, R., Silvestri, A., Solarz, M., Spiczak, G. M., et al.: Optical properties of deep glacial ice at the South Pole, *Journal of Geophysical Research: Atmospheres*, 111(D13), <https://doi.org/10.1029/2005JD006687>, 2006.
- Askebjerg, P., Barwick, S. W., Bergström, L., Bouchta, A., Carius, S., Coulthard, A., Engel, K., Erlandsson, B., Goobar, A., Gray, L., Hallgren, A., Halzen, F., Hulth, P. O., Jacobsen, J., Johansson, S., Kandhadai, V., Liubarsky, I., Lowder, D., Miller, T., Mock, P. C., Morse, R., Porrata, R., Price, P. B., Richards, A., Rubinstein, H., Schneider, E., Sun, Q., Tilav, S., Walck, C. and Yodh, G.: Optical Properties of the South Pole Ice at Depths Between 0.8 and 1 Kilometer, *Science*, 267(5201), 1147–1150, <https://doi.org/10.1126/science.267.5201.1147>, 1995.
- 740 Askebjerg, P., Barwick, S. W., Bergström, L., Bouchta, A., Carius, S., Dalberg, E., Erlandsson, B., Goobar, A., Gray, L., Hallgren, A., Halzen, F., Heukenkamp, H., Hulth, P. O., Hundertmark, S., Jacobsen, J., Kandhadai, V., Karle, A., Liubarsky, I., Lowder, D., Miller, T., Mock, P., Morse, R., Porrata, R., Price, P. B., Richards, A., Rubinstein, H., Schneider, E., Spiering, C., Streicher, O., Sun, Q., Thon, T., Tilav, S., Wischniewski, R., Walck, C. and Yodh, G.: UV and optical light transmission properties in deep ice at the South Pole, *Geophysical Research Letters*, 24(11), 1355–1358, <https://doi.org/10.1029/97GL01246>, 1997.
- 745 Bintanja, R. and Van Den Broeke, M. R.: The Surface Energy Balance of Antarctic Snow and Blue Ice, *J. Appl. Meteor.*, 34(4), 902–926, [https://doi.org/10.1175/1520-0450\(1995\)034<0902:TSEBOA>2.0.CO;2](https://doi.org/10.1175/1520-0450(1995)034<0902:TSEBOA>2.0.CO;2), 1995.
- Bøggild, C. E., Oerter, H. and Tukiainen, T.: Increased ablation of Wisconsin ice in eastern north Greenland: observations and modelling, *Annals of Glaciology*, 23, 144–148, <https://doi.org/10.3189/S0260305500013367>, 1996.
- 750 Bøggild, C. E., Brandt, R. E., Brown, K. J. and Warren, S. G.: The ablation zone in northeast Greenland: ice types, albedos and impurities, *Journal of Glaciology*, 56(195), 101–113, <https://doi.org/10.3189/002214310791190776>, 2010.
- Bohren, C. F.: Colors of snow, frozen waterfalls, and icebergs, *J. Opt. Soc. Am.*, JOSA, 73(12), 1646–1652, <https://doi.org/10.1364/JOSA.73.001646>, 1983.

- Bohren, C. F.: Multiple scattering of light and some of its observable consequences, *American Journal of Physics*, 55(6), 524–533, <https://doi.org/10.1119/1.15109>, 1987.
- Bohren, C. F. and Barkstrom, B. R.: Theory of the optical properties of snow, *J. Geophys. Res.*, 79(30), 4527–4535, <https://doi.org/10.1029/JC079i030p04527>, 1974.
- 760 Brandt, R. E. and Warren, S. G.: Solar-heating rates and temperature profiles in Antarctic snow and ice, *Journal of Glaciology*, 39(131), 99–110, <https://doi.org/10.3189/S0022143000015756>, 1993.
- Briegleb, B. P. and Light, B.: A Delta-Eddington Multiple Scattering Parameterization for Solar Radiation in the Sea Ice Component of the Community Climate System Model, Technical Note, National Center for Atmospheric Research, Boulder, Colorado. <http://dx.doi.org/10.5065/D6B27S71>, last access: 18 February 2019, 2007.
- 765 van den Broeke, M. R., Smeets, P., Ettema, J., van der Veen, C., van de Wal, R. and Oerlemans, J.: Partitioning of melt energy and meltwater fluxes in the ablation zone of the west Greenland ice sheet, *The Cryosphere*, 2(2), 179–189, <https://doi.org/10.5194/tc-2-179-2008>, 2008.
- Brunt, K. M., Neumann, T. A., Amundson, J. M., Kavanaugh, J. L., Moussavi, M. S., Walsh, K. M., Cook, W. B. and Markus, T.: MABEL photon-counting laser altimetry data in Alaska for ICESat-2 simulations and
770 development, *The Cryosphere*, 10(4), 1707–1719, <https://doi.org/10.5194/tc-10-1707-2016>, 2016.
- Cantrell, C. A.: Technical Note: Review of methods for linear least-squares fitting of data and application to atmospheric chemistry problems, *Atmos. Chem. Phys.*, 8(17), 5477–5487, <https://doi.org/10.5194/acp-8-5477-2008>, 2008.
- Cooper, M. G., Smith, L. C., Rennermalm, A. K., Miège, C., Pitcher, L. H., Ryan, J. C., Yang, K. and Cooley, S.
775 W.: Meltwater storage in low-density near-surface bare ice in the Greenland ice sheet ablation zone, *The Cryosphere*, 12(3), 955–970, <https://doi.org/10.5194/tc-12-955-2018>, 2018.
- Dadic, R., Mullen, P. C., Schneebeli, M., Brandt, R. E. and Warren, S. G.: Effects of bubbles, cracks, and volcanic tephra on the spectral albedo of bare ice near the Transantarctic Mountains: Implications for sea glaciers on Snowball Earth, *Journal of Geophysical Research: Earth Surface*, 118(3), 1658–1676,
780 <https://doi.org/10.1002/jgrf.20098>, 2013.
- Deems, J. S., Painter, T. H. and Finnegan, D. C.: Lidar measurement of snow depth: a review, *Journal of Glaciology*, 59(215), 467–479, <https://doi.org/10.3189/2013JoG12J154>, 2013.
- Di Mauro, B., Baccolo, G., Garzonio, R., Giardino, C., Massabò, D., Piazzalunga, A., Rossini, M. and Colombo, R.: Impact of impurities and cryoconite on the optical properties of the Morteratsch Glacier (Swiss Alps), *The Cryosphere*, 11(6), 2393–2409, <https://doi.org/10.5194/tc-11-2393-2017>, 2017.
- Doherty, S. J., Warren, S. G., Grenfell, T. C., Clarke, A. D. and Brandt, R. E.: Light-absorbing impurities in Arctic snow, *Atmos. Chem. Phys.*, 10(23), 11647–11680, <https://doi.org/10.5194/acp-10-11647-2010>, 2010.

- 790 Fisher, F. N., King, M. D. and Lee-Taylor, J.: Extinction of UV-visible radiation in wet midlatitude (maritime) snow: Implications for increased NO_x emission, *J. Geophys. Res.*, 110(D21), D21301, <https://doi.org/10.1029/2005JD005963>, 2005.
- France, J. L., King, M. D., Frey, M. M., Erbland, J., Picard, G., Preunkert, S., MacArthur, A. and Savarino, J.: Snow optical properties at Dome C (Concordia), Antarctica; implications for snow emissions and snow chemistry of reactive nitrogen, *Atmospheric Chemistry and Physics*, 11(18), 9787–9801, <https://doi.org/10.5194/acp-11-9787-2011>, 2011.
- 795 Frey, K. E., Perovich, D. K. and Light, B.: The spatial distribution of solar radiation under a melting Arctic sea ice cover, *Geophysical Research Letters*, 38(22), <https://doi.org/10.1029/2011GL049421>, 2011.
- Gardner, A. S. and Sharp, M. J.: A review of snow and ice albedo and the development of a new physically based broadband albedo parameterization, *J. Geophys. Res.*, 115(F1), F01009, <https://doi.org/10.1029/2009JF001444>, 2010.
- 800 Gardner, A. S., Smith, B. E., Brunt, K. M., Harding, D. J., Neumann, T. and Walsh, K.: ICESat2 subsurface-scattering biases estimated based on the 2015 SIMPL/AVRIS campaign, in *AGU Fall Meeting Abstracts*, vol. 41, pp. C41C-0710, <http://adsabs.harvard.edu/abs/2015AGUFM.C41C0710G>, last access: 25 January 2019, 2015.
- Gerland, S., Liston, G. E., Winther, J.-G., Ørbæk, J. B. and Ivanov, B. V.: Attenuation of solar radiation in Arctic snow: field observations and modelling, *Annals of Glaciology*, 31, 364–368, <https://doi.org/10.3189/172756400781820444>, 2000.
- Goelles, T. and Bøggild, C. E.: Albedo reduction of ice caused by dust and black carbon accumulation: a model applied to the K-transect, West Greenland, *Journal of Glaciology*, 63(242), 1063–1076, <https://doi.org/10.1017/jog.2017.74>, 2017.
- 810 Goelles, T., Bøggild, C. E. and Greve, R.: Ice sheet mass loss caused by dust and black carbon accumulation, *The Cryosphere*, 9(5), 1845–1856, <https://doi.org/10.5194/tc-9-1845-2015>, 2015.
- Gow, A. J., Meese, D. A., Alley, R. B., Fitzpatrick, J. J., Anandkrishnan, S., Woods, G. A. and Elder, B. C.: Physical and structural properties of the Greenland Ice Sheet Project 2 ice core: A review, *J. Geophys. Res.*, 102(C12), 26559–26575, <https://doi.org/10.1029/97JC00165>, 1997.
- 815 Greeley, A., Kurtz, N. T., Neumann, T. and Markus, T.: Estimating Surface Elevation Bias Due to Subsurface Scattered Photons from Visible Wavelength Laser Altimeters, in *AGU Fall Meeting Abstracts*, vol. 51, <http://adsabs.harvard.edu/abs/2017AGUFM.C51A0961G>, last access: 25 January 2019, 2017.
- Grenfell, T. C.: The Effects of Ice Thickness on the Exchange of Solar Radiation Over the Polar Oceans, *Journal of Glaciology*, 22(87), 305–320, <https://doi.org/10.3189/S0022143000014295>, 1979.
- 820 Grenfell, T. C.: A radiative transfer model for sea ice with vertical structure variations, *Journal of Geophysical Research: Oceans*, 96(C9), 16991–17001, <https://doi.org/10.1029/91JC01595>, 1991.

- Grenfell, T. C. and Maykut, G. A.: The Optical Properties of Ice and Snow in the Arctic Basin*, *Journal of Glaciology*, 18(80), 445–463, <https://doi.org/10.3189/S0022143000021122>, 1977.
- 825 Grenfell, T. C. and Perovich, D. K.: Radiation absorption coefficients of polycrystalline ice from 400–1400 nm, *J. Geophys. Res.*, 86(C8), 7447–7450, <https://doi.org/10.1029/JC086iC08p07447>, 1981.
- Grenfell, T. C. and Warren, S. G.: Representation of a nonspherical ice particle by a collection of independent spheres for scattering and absorption of radiation, *J. Geophys. Res.*, 104(D24), 31697–31709, <https://doi.org/10.1029/1999JD900496>, 1999.
- 830 Grenfell, T. C., Light, B. and Perovich, D. K.: Spectral transmission and implications for the partitioning of shortwave radiation in arctic sea ice, *Ann. Glaciol.*, 44, 1–6, <https://doi.org/10.3189/172756406781811763>, 2006.
- He, C. and Flanner, M.: Snow Albedo and Radiative Transfer: Theory, Modeling, and Parameterization, in *Springer Series in Light Scattering*, edited by A. Kokhanovsky, pp. 67–133, Springer International Publishing, Cham, https://doi.org/10.1007/978-3-030-38696-2_3, 2020.
- 835 He, C., Takano, Y., Liou, K.-N., Yang, P., Li, Q. and Chen, F.: Impact of Snow Grain Shape and Black Carbon–Snow Internal Mixing on Snow Optical Properties: Parameterizations for Climate Models, *J. Climate*, 30(24), 10019–10036, <https://doi.org/10.1175/JCLI-D-17-0300.1>, 2017.
- Hoffman, M. J., Fountain, A. G. and Liston, G. E.: Near-surface internal melting: a substantial mass loss on Antarctic Dry Valley glaciers, *Journal of Glaciology*, 60(220), 361–374, <https://doi.org/10.3189/2014JoG13J095>, 2014.
- 840 Holland, M. M., Bailey, D. A., Briegleb, B. P., Light, B. and Hunke, E.: Improved Sea Ice Shortwave Radiation Physics in CCSM4: The Impact of Melt Ponds and Aerosols on Arctic Sea Ice*, *J. Climate*, 25(5), 1413–1430, <https://doi.org/10.1175/JCLI-D-11-00078.1>, 2012.
- van de Hulst, H. C.: Multiple light scattering: tables, formulas, and applications, Academic Press, New York., 1980.
- 845 Järvinen, O. and Leppäranta, M.: Solar radiation transfer in the surface snow layer in Dronning Maud Land, Antarctica, *Polar Science*, 7(1), 1–17, <https://doi.org/10.1016/j.polar.2013.03.002>, 2013.
- Joseph, J. H., Wiscombe, W. J. and Weinman, J. A.: The Delta-Eddington Approximation for Radiative Flux Transfer, *J. Atmos. Sci.*, 33(12), 2452–2459, [https://doi.org/10.1175/1520-0469\(1976\)033<2452:TDEAFR>2.0.CO;2](https://doi.org/10.1175/1520-0469(1976)033<2452:TDEAFR>2.0.CO;2), 1976.
- 850 King, M. D. and Simpson, W. R.: Extinction of UV radiation in Arctic snow at Alert, Canada (82°N), *J. Geophys. Res.*, 106(D12), 12499–12507, <https://doi.org/10.1029/2001JD900006>, 2001.
- Kokhanovsky, A. A. and Zege, E. P.: Scattering optics of snow, *Appl. Opt.*, 43(7), 1589, <https://doi.org/10.1364/AO.43.001589>, 2004.

- 855 Kuipers Munneke, P., van den Broeke, M. R., Reijmer, C. H., Helsen, M. M., Boot, W., Schneebeli, M. and Steffen, K.: The role of radiation penetration in the energy budget of the snowpack at Summit, Greenland, *The Cryosphere*, 3(2), 155–165, <https://doi.org/10.5194/tc-3-155-2009>, 2009.
- Leathers, R. A., Downes, T. V., Davis, C. O. and Mobley, C. D.: Monte Carlo Radiative Transfer Simulations for Ocean Optics: A Practical Guide, Memorandum, Naval Research Laboratory, Washington, D.C. https://www.oceanopticsbook.info/packages/iws_12h/conversion/files/Leathersetal_NRL2004.pdf, last access: 11 October 2020, 2004.
- 860 Libois, Q., Picard, G., France, J. L., Arnaud, L., Dumont, M., Carmagnola, C. M. and King, M. D.: Influence of grain shape on light penetration in snow, *The Cryosphere*, 7(6), 1803–1818, <https://doi.org/10.5194/tc-7-1803-2013>, 2013.
- Light, B., Maykut, G. A. and Grenfell, T. C.: A two-dimensional Monte Carlo model of radiative transfer in sea ice, *Journal of Geophysical Research: Oceans*, 108(C7), 3219, <https://doi.org/10.1029/2002JC001513>, 2003.
- Light, B., Maykut, G. A. and Grenfell, T. C.: A temperature-dependent, structural-optical model of first-year sea ice, *J. Geophys. Res.*, 109(C6), C06013, <https://doi.org/10.1029/2003JC002164>, 2004.
- Light, B., Grenfell, T. C. and Perovich, D. K.: Transmission and absorption of solar radiation by Arctic sea ice during the melt season, *J. Geophys. Res.*, 113(C3), C03023, <https://doi.org/10.1029/2006JC003977>, 2008.
- 870 Liston, G. E. and Winther, J.-G.: Antarctic Surface and Subsurface Snow and Ice Melt Fluxes, *J. Climate*, 18(10), 1469–1481, <https://doi.org/10.1175/JCLI3344.1>, 2005.
- Liston, G. E., Bruland, O., Elvehøy, H. and Sand, K.: Below-surface ice melt on the coastal Antarctic ice sheet, *Journal of Glaciology*, 45(150), 273–285, <https://doi.org/10.3189/002214399793377130>, 1999a.
- 875 Liston, G. E., Bruland, O., Winther, J.-G., Elvehøy, H. and Sand, K.: Meltwater production in Antarctic blue-ice areas: sensitivity to changes in atmospheric forcing, *Polar Research*, 18(2), 283–290, <https://doi.org/10.1111/j.1751-8369.1999.tb00305.x>, 1999b.
- Malinka, A., Zege, E., Heygster, G. and Istomina, L.: Reflective properties of white sea ice and snow, *The Cryosphere*, 10(6), 2541–2557, <https://doi.org/10.5194/tc-10-2541-2016>, 2016.
- Malinka, A. V.: Light scattering in porous materials: Geometrical optics and stereological approach, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 141, 14–23, <https://doi.org/10.1016/j.jqsrt.2014.02.022>, 2014.
- 880 Markus, T., Neumann, T., Martino, A., Abdalati, W., Brunt, K., Csatho, B., Farrell, S., Fricker, H., Gardner, A., Harding, D., Jasinski, M., Kwok, R., Magruder, L., Lubin, D., Luthcke, S., Morison, J., Nelson, R., Neuenschwander, A., Palm, S., Popescu, S., Shum, C., Schutz, B. E., Smith, B., Yang, Y. and Zwally, J.: The Ice, Cloud, and land Elevation Satellite-2 (ICESat-2): Science requirements, concept, and implementation, *Remote Sensing of Environment*, 190, 260–273, <https://doi.org/10.1016/j.rse.2016.12.029>, 2017.
- 885 Mätzler, C.: MATLAB Functions for Mie Scattering and Absorption, Version 2, Research Report, Institut für Angewandte Physik, Bern, Switzerland. <https://boris.unibe.ch/146550/>, 2002.

- 890 Maykut, G. A. and Untersteiner, N.: Some results from a time-dependent thermodynamic model of sea ice, *Journal of Geophysical Research* (1896-1977), 76(6), 1550–1575, <https://doi.org/10.1029/JC076i006p01550>, 1971.
- Meirolid-Mautner, I. and Lehning, M.: Measurements and model calculations of the solar shortwave fluxes in snow on Summit, Greenland, *Annals of Glaciology*, 38, 279–284, <https://doi.org/10.3189/172756404781814753>, 2004.
- 895 Muhs, D. R.: The geologic records of dust in the Quaternary, *Aeolian Research*, 9, 3–48, <https://doi.org/10.1016/j.aeolia.2012.08.001>, 2013.
- Mullen, P. C. and Warren, S. G.: Theory of the optical properties of lake ice, *J. Geophys. Res.*, 93(D7), 8403–8414, <https://doi.org/10.1029/JD093iD07p08403>, 1988.
- Pegau, W. S. and Zaneveld, J. R. V.: Field measurements of in-ice radiance, *Cold Regions Science and Technology*, 31(1), 33–46, [https://doi.org/10.1016/S0165-232X\(00\)00004-5](https://doi.org/10.1016/S0165-232X(00)00004-5), 2000.
- 900 Perovich, D. K.: The Optical Properties of Sea Ice., U.S. Army Cold Regions Research and Engineering Laboratory, Hanover, NH. <https://apps.dtic.mil/dtic/tr/fulltext/u2/a310586.pdf>, 1996.
- Perovich, D. K. and Govoni, J. W.: Absorption coefficients of ice from 250 to 400 nm, *Geophys. Res. Lett.*, 18(7), 1233–1235, <https://doi.org/10.1029/91GL01642>, 1991.
- 905 Petit, J. R., Jouzel, J., Raynaud, D., Barkov, N. I., Barnola, J.-M., Basile, I., Bender, M., Chappellaz, J., Davis, M., Delaygue, G., Delmotte, M., Kotlyakov, V. M., Legrand, M., Lipenkov, V. Y., Lorius, C., Pépin, L., Ritz, C., Saltzman, E. and Stievenard, M.: Climate and atmospheric history of the past 420,000 years from the Vostok ice core, Antarctica, *Nature*, 399(6735), 429–436, <https://doi.org/10.1038/20859>, 1999.
- Petrenko, V. V., Severinghaus, J. P., Brook, E. J., Reeh, N. and Schaefer, H.: Gas records from the West Greenland ice margin covering the Last Glacial Termination: a horizontal ice core, *Quaternary Science Reviews*, 910 25(9), 865–875, <https://doi.org/10.1016/j.quascirev.2005.09.005>, 2006.
- Picard, G., Libois, Q. and Arnaud, L.: Refinement of the ice absorption spectrum in the visible using radiance profile measurements in Antarctic snow, *The Cryosphere*, 10(6), 2655–2672, <https://doi.org/10.5194/tc-10-2655-2016>, 2016.
- 915 Price, P. B. and Bergström, L.: Enhanced Rayleigh scattering as a signature of nanoscale defects in highly transparent solids, *null*, 75(5), 1383–1390, <https://doi.org/10.1080/01418619708209861>, 1997a.
- Price, P. B. and Bergström, L.: Optical properties of deep ice at the South Pole: scattering, *Appl. Opt.*, AO, 36(18), 4181–4194, <https://doi.org/10.1364/AO.36.004181>, 1997b.
- Reeh, N., Oerter, H. and Thomsen, H. H.: Comparison between Greenland ice-margin and ice-core oxygen-18 records, *Annals of Glaciology*, 35, 136–144, <https://doi.org/10.3189/172756402781817365>, 2002.

- 920 Ridley, J. K. and Partington, K. C.: A model of satellite radar altimeter return from ice sheets, *International Journal of Remote Sensing*, 9(4), 601–624, <https://doi.org/10.1080/01431168808954881>, 1988.
- Rignot, E., Echelmeyer, K. and Krabill, W.: Penetration depth of interferometric synthetic-aperture radar signals in snow and ice, *Geophysical Research Letters*, 28(18), 3501–3504, <https://doi.org/10.1029/2000GL012484>, 2001.
- 925 Ruth, U., Wagenbach, D., Steffensen, J. P. and Bigler, M.: Continuous record of microparticle concentration and size distribution in the central Greenland NGRIP ice core during the last glacial period, *Journal of Geophysical Research: Atmospheres*, 108(D3), <https://doi.org/10.1029/2002JD002376>, 2003.
- Ryan, J. C., Hubbard, A. L., Stibal, M., Irvine-Fynn, T. D., Cook, J., Smith, L. C., Cameron, K. and Box, J. E.: Dark zone of the Greenland Ice Sheet controlled by distributed biologically-active impurities, *Nature*
- 930 *Communications*, 9(1), 1065, <https://doi.org/10.1038/s41467-018-03353-2>, 2018.
- Schuster, A.: Radiation through a foggy atmosphere, *The Astrophysical Journal*, XX1(1), 1–22, 1905.
- Schuster, C.: Weathering crust processes on melting glacier ice (Alberta, Canada), *Theses and Dissertations (Comprehensive)* <http://scholars.wlu.ca/etd/489>, 2001.
- Schutz, B. E., Zwally, H. J., Shuman, C. A., Hancock, D. and DiMarzio, J. P.: Overview of the ICESat Mission,
- 935 *Geophys. Res. Lett.*, 32(21), L21S01, <https://doi.org/10.1029/2005GL024009>, 2005.
- Smith, B. E., Gardner, A., Schneider, A. and Flanner, M.: Modeling biases in laser-altimetry measurements caused by scattering of green light in snow, *Remote Sensing of Environment*, 215, 398–410, <https://doi.org/10.1016/j.rse.2018.06.012>, 2018.
- Stibal, M., Box, J. E., Cameron, K. A., Langen, P. L., Yallop, M. L., Mottram, R. H., Khan, A. L., Molotch, N.
- 940 P., Christmas, N. A. M., Quaglia, F. C., Remias, D., Smeets, C. J. P. P., Broeke, M. R. van den, Ryan, J. C., Hubbard, A., Tranter, M., As, D. van and Ahlström, A. P.: Algae Drive Enhanced Darkening of Bare Ice on the Greenland Ice Sheet, *Geophysical Research Letters*, 44(22), 11,463–11,471, <https://doi.org/10.1002/2017GL075958>, 2017.
- Takeuchi, N.: Optical characteristics of cryoconite (surface dust) on glaciers: the relationship between light absorbcency and the property of organic matter contained in the cryoconite, *Annals of Glaciology*, 34(1), 409–414, <https://doi.org/10.3189/172756402781817743>, 2002.
- Taylor, B. N. and Kuyatt, C. E.: Guidelines for Evaluating and Expressing the Uncertainty of NIST Measurement Results (NIST Technical Note vol 1297), National Institute of Standards and Technology, Gaithersburg, MD. <http://physics.nist.gov/Pubs/guidelines/TN1297/tn1297s.pdf>, 1994.
- 950 Tuzet, F., Dumont, M., Arnaud, L., Voisin, D., Lamare, M., Larue, F., Revuelto, J. and Picard, G.: Influence of light-absorbing particles on snow spectral irradiance profiles, *The Cryosphere*, 13(8), 2169–2187, <https://doi.org/10.5194/tc-13-2169-2019>, 2019.

- Wang, L., Jacques, S. L. and Zheng, L.: MCML—Monte Carlo modeling of light transport in multi-layered tissues, *Computer Methods and Programs in Biomedicine*, 47(2), 131–146, [https://doi.org/10.1016/0169-2607\(95\)01640-F](https://doi.org/10.1016/0169-2607(95)01640-F), 1995.
- 955 Warren, S. G.: Optical properties of snow, *Rev. Geophys.*, 20(1), 67–89, <https://doi.org/10.1029/RG020i001p00067>, 1982.
- Warren, S. G.: Optical constants of ice from the ultraviolet to the microwave, *Appl. Opt.*, AO, 23(8), 1206–1225, <https://doi.org/10.1364/AO.23.001206>, 1984.
- 960 Warren, S. G. and Brandt, R. E.: Optical constants of ice from the ultraviolet to the microwave: A revised compilation, *J. Geophys. Res.*, 113(D14), D14220, <https://doi.org/10.1029/2007JD009744>, 2008.
- Warren, S. G., Brandt, R. E., Grenfell, T. C. and McKay, C. P.: Snowball Earth: Ice thickness on the tropical ocean, *Journal of Geophysical Research: Oceans*, 107(C10), 31-1-31–18, <https://doi.org/10.1029/2001JC001123>, 2002.
- 965 Warren, S. G., Brandt, R. E. and Grenfell, T. C.: Visible and near-ultraviolet absorption spectrum of ice from transmission of solar radiation into snow, *Appl. Opt.*, AO, 45(21), 5320–5334, <https://doi.org/10.1364/AO.45.005320>, 2006.
- Wientjes, I. G. M., Van de Wal, R. S. W., Reichart, G. J., Sluijs, A. and Oerlemans, J.: Dust from the dark region in the western ablation zone of the Greenland ice sheet, *The Cryosphere*, 5(3), 589–601, <https://doi.org/10.5194/tc-5-589-2011>, 2011.
- 970 Wientjes, I. G. M., De Van Wal, R. S. W., Schwikowski, M., Zapf, A., Fahrni, S. and Wacker, L.: Carbonaceous particles reveal that Late Holocene dust causes the dark region in the western ablation zone of the Greenland ice sheet, *Journal of Glaciology*, 58(210), 787–794, <https://doi.org/10.3189/2012JoG11J165>, 2012.
- Wiscombe, W. J. and Warren, S. G.: A Model for the Spectral Albedo of Snow. I: Pure Snow, *J. Atmos. Sci.*, 37(12), 2712–2733, [https://doi.org/10.1175/1520-0469\(1980\)037<2712:AMFTSA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1980)037<2712:AMFTSA>2.0.CO;2), 1980.
- 975 Woschnagg, K. and Price, P. B.: Temperature dependence of absorption in ice at 532 nm, *Appl. Opt.*, 40(15), 2496–2500, <https://doi.org/10.1364/AO.40.002496>, 2001.
- Yallop, M. L., Anesio, A. M., Perkins, R. G., Cook, J., Telling, J., Fagan, D., MacFarlane, J., Stibal, M., Barker, G., Bellas, C., Hodson, A., Tranter, M., Wadham, J. and Roberts, N. W.: Photophysiology and albedo-changing potential of the ice algal community on the surface of the Greenland ice sheet, *ISME J*, 6(12), 2302–2313, <https://doi.org/10.1038/ismej.2012.107>, 2012.
- 980 Yang, Y., Marshak, A., Han, M., Palm, S. P. and Harding, D. J.: Snow grain size retrieval over the polar ice sheets with the Ice, Cloud, and land Elevation Satellite (ICESat) observations, *Journal of Quantitative Spectroscopy and Radiative Transfer*, 188, 159–164, <https://doi.org/10.1016/j.jqsrt.2016.03.033>, 2017.

985 York, D., Evensen, N. M., Martínez, M. L. and De Basabe Delgado, J.: Unified equations for the slope, intercept, and standard errors of the best straight line, *American Journal of Physics*, 72(3), 367–375, <https://doi.org/10.1119/1.1632486>, 2004.

Zebker, H. A. and Weber Hoen, E.: Penetration depths inferred from interferometric volume decorrelation observed over the Greenland Ice Sheet, *IEEE Transactions on Geoscience and Remote Sensing*, 38(6), 2571–2583, <https://doi.org/10.1109/36.885204>, 2000.

Zhang, X., Qiu, J., Li, X., Zhao, J. and Liu, L.: Complex refractive indices measurements of polymers in visible and near-infrared bands, *Appl. Opt.*, 59(8), 2337, <https://doi.org/10.1364/AO.383831>, 2020.

Zieger, P., Weingartner, E., Henzing, J., Moerman, M., de Leeuw, G., Mikkilä, J., Ehn, M., Petäjä, T., Clémer, K., van Roozendaal, M., Yilmaz, S., Frieß, U., Irie, H., Wagner, T., Shaiganfar, R., Beirle, S., Apituley, A.,
995 Wilson, K. and Baltensperger, U.: Comparison of ambient aerosol extinction coefficients obtained from in-situ, MAX-DOAS and LIDAR measurements at Cabauw, *Atmos. Chem. Phys.*, 11(6), 2603–2624, <https://doi.org/10.5194/acp-11-2603-2011>, 2011.

1000

Figures




Fig. 1: The irradiance sensor is comprised of a remote cosine receptor and fibre optic light guide fitted inside an insulated white PVC tube of 2 m length. Holes are drilled level and horizontal into the ice, the tube is inserted, and drill shavings are packed around the hole to prevent stray reflections. The cosine receptor collects the downwelling light, guides it to the fibre optic cable that transmits the light to an Ocean Optics® JAZ spectrometer, and a computer running the Ocean Optics® Ocean View software records the spectra. Background downwelling surface spectra are recorded on a 2 m mast drilled into the ice approximately 3 m to the northwest of the in-ice measurement location (see photo background). This photograph was taken on 21 July 2018 at ~13:22 local time (UTC-3).

- Deleted: Experimental setup schematic.
- Deleted: horizontal cylindrical rod is
- Deleted: , working from the bottom toward the top
- Deleted: Inside the tube is a fibre optic cable attached to a remote cosine receptor with a Spectralon diffusing element oriented parallel to the rod (normal to the vertical), set approximately 2 mm out from the tube exterior and in contact with the overlying ice.
- Deleted: After all measurements are complete, a 2 m ruler is inserted into each hole, two at a time, and the distance between the holes is recorded to determine their depth relative to each other and to the surface.




Fig. 2: (a) Residual dark-current spectra for the surface-based reference spectrometer and the in-ice spectrometer. Dark current spectra are recorded prior to each absolute irradiance measurement as input to the OceanView software dark-current correction module. Shown here are residual dark-current spectra after automated software correction, which are treated as systematic errors and subtracted from irradiance profiles prior to fitting experimental k_{att} values. (b) Ideal angular response function (ideal cosine) and empirical angular response function provided by Ocean Optics from laboratory measurements on the same type of irradiance sensor used in this study. The dashed red line in (b) is used as an empirical probability density function for the angular response of the cosine receptor in our Monte Carlo simulations of detector interference.

1025

1030





Fig. 3: Photographs of an ice core collected at the field site. (a) The upper few centimetres of ice is semi-granular, with ~4 cm of unrecovered granular ice not shown. (b) The 122 cm ice core was broken into three segments corresponding to depths of 4–45 cm, 45–74 cm, and 74–122 cm below the ice surface (the far right of the image in (b) is at 74 cm). The density of these segments is 801 kg m^{-3} , 884 kg m^{-3} , and 888 kg m^{-3} , respectively. Below ~10 cm, the bubbly ice appears foliated, reflecting variations in bubble density and size distribution that affect scattering. Black box in (b) is approximately the image area in (a).



040 Fig. 4: (a) Field spectra of surface **downwelling** ($z = z_0$) and in-ice irradiance at four depths below the ice surface collected on 20
 045 July 2018 between 13:45 and 14:35 local time in the western Greenland ablation zone (67.15 °N, 50.02 °W). Raw data were recorded
 at 1 Hz frequency for 30 seconds, yielding 30 irradiance profiles at each depth. Shown here are 30-second averages at ~0.35 nm
 spectral resolution for each depth (black dots), and 1-nm interpolated values smoothed with a 3-nm centred moving mean filter for
 clarity (continuous lines). (b) **Transmittance** at each depth, with 30-second averages (black dots), 1-nm interpolated values
 (continuous lines), and shaded bounds ($\pm 2\sigma$) representing propagated measurement uncertainty deduced from the standard
 deviations of the 1 Hz raw data ($N=30$ for each value).

Deleted: Relative irradiance

Deleted: is $<1 \text{ W m}^{-2} \text{ nm}^{-1}$ at all wavelengths and measured depths...






Fig. 5: (a) Sample linear regressions between measured transmittance (indicated by rectangles with width and height proportional to measurement uncertainty in both variables) and depth for five representative wavelengths spanning the measured spectral range. The slope of each line is the attenuation coefficient k_{att} . Shaded bounds are $\pm 95\%$ confidence intervals from a two-sided t-distribution. (b) Red box inset in (a) shows the y-axis intercept of each regression is less than 100%, indicating the magnitude of deviation from Bouguer's law near the surface. (c) Spectral k_{att} (blue dots with shaded uncertainty; left axis) and spectral albedo (red dots with shaded uncertainty; right axis). Beyond ~ 700 nm, in-ice transmitted irradiance is too low to reliably estimate k_{att} (see Fig. 4a and Fig. 2c), as indicated by the increased scatter in k_{att} values.

1060



1065 Fig. 6: Visible and near-UV attenuation coefficient k_{att} spectra from measurements of light transmission in bare glacier ice collected
 1070 on 20 July 2018 at 12–77 cm depth below the ice sheet surface. Field estimates are compared with asymptotic two-stream theory for
 optically clean bubbly ice (continuous dotted line) and with values at nine wavelengths from four simulations with a 3-dimensional
 Monte Carlo radiative transfer model (solid symbols). Monte Carlo values for clean bubbly ice are shown for 350, 400, and 450 nm
 to demonstrate detector interference at these wavelengths; values at wavelengths >550 nm converge with field spectra and are
 omitted for clarity. Two measures of uncertainty are shown: 1) statistical linear model uncertainty ε_{LM} (shaded uncertainty bounds;
 ± 1 standard error in the linear regression) and, 2) ε_{LM} combined with systematic uncertainty ε_{MC} due to detector interference
 estimated with Monte Carlo (error bars; $\mu \pm \varepsilon$). The same comparison for the 21 July experiment (inset) suggests detector
 interference is lower and within statistical uncertainty at wavelengths >400 nm.



075 Fig. 7: (a) Effective attenuation coefficient k_{att} for the near-surface 0–12 cm region compared to k_{att} values estimated for the interior 12–77 cm region. (b) Effective k_{att} values are ~ 1.6 times larger at wavelengths beyond about 600 nm but are ~ 4 times larger between 400–600 nm. The spectral dependence suggests higher influence of absorptive impurities on attenuation enhancement near the ice surface than in the ice interior. The shaded bounds represent propagated ± 1.2 cm vertical measurement uncertainty.

Deleted: estimated with a finite-difference solution to Eq. ()

Deleted: The shaded bounds represent ± 1.2 cm vertical measurement uncertainty.






Fig. 8: Measured in-ice irradiance compared with: (a) Bouguer's law (Eq. 7) with no modification, and (b) the **piece-wise** Bouguer law (Eq. 16). The error structure (c–d) provides insight into the near-surface attenuation processes: relative errors (%) are positive (model under-predicts attenuation) at all wavelengths but are highest in the near-UV, lowest in the blue, and increase monotonically into the red end of the visible spectrum. (d) errors are small, and generally decrease monotonically with increasing wavelength. Taken together, near-surface attenuation enhancement is on the order 10–60%.

- Deleted: modified
- Deleted: with $\chi = 15\%$.
- Deleted: c
- Deleted: d




095
100
Fig. 9: Uncertainty budget for reported asymptotic attenuation coefficient k_{att} values for Layer A (12–77 cm) (20 July experiment). Systematic uncertainties examined include spectrometer sensitivity to dark noise, ice thickness (detector position) measurement uncertainty, the non-ideal angular response of the irradiance sensor, and scattering and absorption interference by the polyvinyl chloride detector rod (estimated with Monte Carlo simulation). These systematics are summed in quadrature with statistical uncertainty represented by one standard error in the Maximum Likelihood Estimate (MLE) linear regression fits. Values for Layer B (53–124 cm) (not shown) are qualitatively similar but <14% in the region of maximum uncertainty.




1105 Fig. 10: Attenuation coefficient spectra for seven distinct ice structures (from lower left clockwise): interior of clean, dry snowpack ($r_{\text{eff}} \approx 10^{-5}$ m) near Dome-C in Antarctica for two depth regions (90–135 cm and 40–90 cm) (Warren et al., 2006), near-surface (0–10 cm) dry snowpack ($r_{\text{eff}} \approx 10^{-3}$ m) in Dronning Maud Land, Antarctica (Järvinen and Leppäranta, 2013), near-surface (0–5 cm) and interior (0–18 cm) dry snowpack ($r_{\text{eff}} \approx 10^{-4}$ m) near Summit, Greenland (Meirolid-Mautner and Lehning, 2004), surface scattering layer (SSL; 0–5 cm) of multi-year sea ice in the Arctic Ocean (Grenfell and Maykut, 1977), interior of ablating glacier ice in Greenland (this study) (12–77 cm in solid line; 0–12 cm in dotted line), interior of first-year sea ice in the coastal zone near Barrow, Alaska (Grenfell et al., 2006), and interior of multi-year sea ice in the Arctic Ocean (Grenfell et al., 2006). Differences in attenuation magnitude at each wavelength are mostly controlled by structural differences that control scattering, whereas spectral differences are mostly controlled by differences in type and concentration of absorbing impurities.

1110



1115 Fig. 11: Depth profiles of South Pole ice absorptivity at four wavelengths obtained from Eq. 26 and Table 3 of Ackermann et al. (2006). The 1% K^{-1} temperature dependence of pure ice absorptivity (Woschnagg and Price, 2001) is removed for comparison with this study's lower (Layer B) and upper (Layer A) absorption coefficient estimate at each wavelength (dashed lines). Values reported in this study are consistent with South Pole values at depths corresponding to the Last Glacial Maximum and Marine Isotope Stage 4 when atmospheric dust concentrations peaked in both hemispheres. Note that South Pole age does not map to ice near the Greenland Ice Sheet margin. Rather, Southern Hemisphere dust concentrations during these periods are consistent with Northern Hemisphere dust concentrations during warm interglacial periods and/or periods with low aeolian activity (Muhs, 2013; Reeh et al., 2002).

1120



1125 Fig. 12: Estimates of ice absorption coefficient k_{abs} , obtained from five distinct sources: laboratory-grown pure ice (Grenfell and
 Perovich, 1981; Perovich and Govoni, 1991), as compiled in Warren (1984), snow in Antarctica, contaminated by trace
 concentrations of light absorbing particles (LAPs) (Picard et al., 2016), glacial ice in Greenland with unknown concentration of
 LAPs (this study), compressed glacial ice at 1755 m depth and 830 m depth in the Antarctic Ice Sheet contaminated by dust deposited
 1130 during the late Pleistocene and Early Holocene, respectively (Ackermann et al., 2006), and snow in Antarctica with the effect of
 LAPs removed (pure ice estimate) (Warren et al., 2006). Values from this study with the detector interference subtracted are shown
 as dotted blue line.

Appendix 1: Monte Carlo radiative transfer model

See attached document.




Table A1: Estimates of attenuation coefficient and absorption coefficient obtained from solar irradiance transmission measurements in glacier ice, one standard error in the linear regression, and coefficient of determination.

wavelength (nm)	k_{att} (m^{-1})	standard error (k_{att})	r^2	k_{abs} (m^{-1})	standard error (k_{abs})
350	0.975	0.021	0.999	0.0162	0.00035
351	0.970	0.022	0.999	0.0160	0.00037
352	0.965	0.023	0.999	0.0159	0.00038
353	0.960	0.024	0.999	0.0157	0.00039
354	0.955	0.025	0.999	0.0155	0.00040
355	0.951	0.025	0.997	0.0154	0.00041
356	0.947	0.026	0.998	0.0153	0.00042
357	0.943	0.027	0.999	0.0152	0.00043
358	0.940	0.026	0.999	0.0150	0.00041
359	0.936	0.026	0.999	0.0149	0.00041
360	0.935	0.026	0.997	0.0149	0.00041
361	0.933	0.027	0.998	0.0148	0.00042
362	0.931	0.029	0.999	0.0148	0.00046
363	0.931	0.030	0.997	0.0148	0.00048
364	0.931	0.031	0.997	0.0148	0.00049
365	0.927	0.031	0.998	0.0147	0.00049
366	0.923	0.031	0.997	0.0145	0.00048
367	0.919	0.032	0.998	0.0144	0.00050
368	0.915	0.031	0.997	0.0143	0.00049
369	0.912	0.030	0.998	0.0142	0.00046
370	0.909	0.029	0.998	0.0141	0.00045
371	0.908	0.029	0.998	0.0140	0.00045
372	0.905	0.030	0.997	0.0140	0.00047
373	0.901	0.031	0.999	0.0138	0.00048
374	0.897	0.033	0.998	0.0137	0.00050
375	0.894	0.034	0.996	0.0136	0.00052
376	0.892	0.035	0.996	0.0136	0.00053

377	0.892	0.035	0.997	0.0136	0.00053
378	0.892	0.038	0.996	0.0136	0.00057
379	0.893	0.039	0.996	0.0136	0.00060
380	0.892	0.040	0.997	0.0135	0.00061
381	0.892	0.042	0.997	0.0136	0.00064
382	0.890	0.044	0.994	0.0135	0.00067
383	0.886	0.045	0.993	0.0134	0.00069
384	0.883	0.046	0.995	0.0133	0.00070
385	0.880	0.047	0.994	0.0132	0.00070
386	0.878	0.046	0.995	0.0131	0.00069
387	0.876	0.045	0.995	0.0131	0.00067
388	0.876	0.044	0.995	0.0131	0.00066
389	0.879	0.046	0.995	0.0132	0.00069
390	0.881	0.047	0.995	0.0132	0.00071
391	0.880	0.048	0.994	0.0132	0.00073
392	0.881	0.050	0.993	0.0132	0.00076
393	0.881	0.052	0.992	0.0132	0.00078
394	0.880	0.054	0.993	0.0132	0.00081
395	0.878	0.055	0.992	0.0131	0.00082
396	0.877	0.056	0.991	0.0131	0.00083
397	0.875	0.056	0.991	0.0131	0.00084
398	0.875	0.057	0.992	0.0130	0.00085
399	0.877	0.059	0.992	0.0131	0.00088
400	0.881	0.060	0.991	0.0132	0.00091
401	0.883	0.062	0.990	0.0133	0.00093
402	0.886	0.064	0.990	0.0134	0.00096
403	0.890	0.066	0.988	0.0135	0.00100
404	0.892	0.067	0.988	0.0136	0.00102
405	0.892	0.068	0.988	0.0136	0.00104
406	0.893	0.069	0.988	0.0136	0.00105
407	0.894	0.070	0.989	0.0136	0.00107
408	0.894	0.071	0.988	0.0136	0.00108
409	0.893	0.072	0.987	0.0136	0.00110
410	0.894	0.074	0.987	0.0136	0.00112
411	0.895	0.075	0.986	0.0136	0.00114
412	0.896	0.075	0.986	0.0137	0.00115

413	0.897	0.076	0.985	0.0137	0.00116
414	0.897	0.076	0.985	0.0137	0.00117
415	0.897	0.077	0.987	0.0137	0.00118
416	0.896	0.077	0.987	0.0137	0.00118
417	0.896	0.078	0.984	0.0137	0.00119
418	0.896	0.079	0.983	0.0137	0.00120
419	0.895	0.080	0.985	0.0137	0.00122
420	0.896	0.081	0.984	0.0137	0.00124
421	0.896	0.082	0.984	0.0137	0.00125
422	0.897	0.082	0.983	0.0137	0.00125
423	0.897	0.082	0.983	0.0137	0.00126
424	0.897	0.083	0.983	0.0137	0.00127
425	0.896	0.084	0.984	0.0137	0.00128
426	0.896	0.084	0.982	0.0137	0.00129
427	0.895	0.085	0.982	0.0137	0.00130
428	0.894	0.085	0.983	0.0136	0.00130
429	0.893	0.086	0.982	0.0136	0.00131
430	0.894	0.088	0.980	0.0136	0.00133
431	0.895	0.088	0.982	0.0137	0.00134
432	0.896	0.089	0.981	0.0137	0.00135
433	0.897	0.090	0.979	0.0137	0.00137
434	0.898	0.091	0.980	0.0137	0.00139
435	0.899	0.091	0.981	0.0138	0.00140
436	0.900	0.092	0.980	0.0138	0.00141
437	0.900	0.093	0.978	0.0138	0.00142
438	0.899	0.093	0.979	0.0138	0.00142
439	0.899	0.093	0.979	0.0138	0.00142
440	0.899	0.093	0.979	0.0138	0.00143
441	0.900	0.093	0.980	0.0138	0.00142
442	0.901	0.093	0.979	0.0138	0.00143
443	0.904	0.093	0.979	0.0139	0.00144
444	0.907	0.094	0.979	0.0140	0.00146
445	0.910	0.095	0.979	0.0141	0.00148
446	0.913	0.096	0.978	0.0142	0.00150
447	0.916	0.097	0.978	0.0143	0.00152
448	0.918	0.098	0.978	0.0144	0.00153

449	0.920	0.099	0.977	0.0144	0.00155
450	0.922	0.099	0.977	0.0145	0.00155
451	0.922	0.099	0.978	0.0145	0.00155
452	0.923	0.099	0.978	0.0145	0.00156
453	0.925	0.099	0.977	0.0146	0.00157
454	0.926	0.100	0.978	0.0146	0.00157
455	0.927	0.100	0.977	0.0146	0.00158
456	0.929	0.100	0.976	0.0147	0.00158
457	0.931	0.100	0.977	0.0148	0.00158
458	0.933	0.100	0.978	0.0148	0.00159
459	0.936	0.101	0.978	0.0149	0.00161
460	0.939	0.101	0.978	0.0150	0.00162
461	0.942	0.102	0.977	0.0151	0.00164
462	0.946	0.103	0.975	0.0153	0.00166
463	0.951	0.104	0.976	0.0154	0.00169
464	0.955	0.105	0.976	0.0155	0.00171
465	0.960	0.105	0.977	0.0157	0.00172
466	0.964	0.105	0.977	0.0158	0.00172
467	0.968	0.105	0.977	0.0160	0.00173
468	0.972	0.105	0.977	0.0161	0.00174
469	0.975	0.105	0.977	0.0162	0.00175
470	0.979	0.106	0.977	0.0163	0.00177
471	0.981	0.107	0.977	0.0164	0.00178
472	0.984	0.107	0.977	0.0165	0.00180
473	0.987	0.107	0.976	0.0166	0.00181
474	0.991	0.107	0.976	0.0167	0.00181
475	0.994	0.107	0.977	0.0168	0.00182
476	0.997	0.107	0.978	0.0169	0.00182
477	1.001	0.107	0.978	0.0171	0.00183
478	1.004	0.107	0.977	0.0172	0.00183
479	1.008	0.107	0.978	0.0173	0.00184
480	1.012	0.107	0.978	0.0175	0.00185
481	1.016	0.107	0.978	0.0176	0.00186
482	1.021	0.107	0.978	0.0178	0.00187
483	1.026	0.108	0.978	0.0179	0.00188
484	1.031	0.108	0.979	0.0181	0.00189

485	1.037	0.108	0.979	0.0183	0.00190
486	1.042	0.107	0.979	0.0185	0.00191
487	1.048	0.107	0.979	0.0187	0.00192
488	1.053	0.107	0.980	0.0189	0.00192
489	1.058	0.107	0.980	0.0191	0.00193
490	1.064	0.107	0.980	0.0193	0.00194
491	1.069	0.107	0.980	0.0195	0.00195
492	1.075	0.107	0.981	0.0197	0.00196
493	1.081	0.107	0.980	0.0199	0.00198
494	1.087	0.107	0.981	0.0202	0.00199
495	1.094	0.108	0.981	0.0204	0.00201
496	1.101	0.108	0.981	0.0207	0.00203
497	1.108	0.108	0.982	0.0209	0.00204
498	1.115	0.108	0.981	0.0212	0.00205
499	1.123	0.108	0.981	0.0215	0.00207
500	1.130	0.108	0.982	0.0218	0.00208
501	1.138	0.108	0.983	0.0221	0.00209
502	1.145	0.108	0.982	0.0223	0.00210
503	1.152	0.107	0.983	0.0226	0.00211
504	1.159	0.107	0.983	0.0229	0.00212
505	1.167	0.108	0.983	0.0232	0.00214
506	1.174	0.108	0.983	0.0235	0.00216
507	1.181	0.108	0.983	0.0238	0.00218
508	1.188	0.108	0.984	0.0240	0.00220
509	1.195	0.109	0.984	0.0243	0.00222
510	1.202	0.109	0.983	0.0246	0.00223
511	1.209	0.109	0.984	0.0249	0.00224
512	1.216	0.109	0.984	0.0252	0.00225
513	1.226	0.109	0.984	0.0256	0.00227
514	1.237	0.108	0.985	0.0261	0.00228
515	1.248	0.108	0.985	0.0265	0.00229
516	1.259	0.108	0.986	0.0270	0.00231
517	1.269	0.108	0.986	0.0274	0.00233
518	1.279	0.108	0.986	0.0279	0.00234
519	1.290	0.108	0.986	0.0283	0.00236
520	1.300	0.108	0.986	0.0288	0.00238

521	1.310	0.107	0.987	0.0292	0.00240
522	1.320	0.107	0.987	0.0297	0.00241
523	1.333	0.107	0.987	0.0303	0.00242
524	1.345	0.106	0.988	0.0308	0.00243
525	1.358	0.105	0.988	0.0314	0.00244
526	1.370	0.105	0.988	0.0320	0.00245
527	1.382	0.104	0.989	0.0326	0.00246
528	1.394	0.104	0.989	0.0331	0.00247
529	1.407	0.104	0.989	0.0337	0.00248
530	1.419	0.104	0.989	0.0343	0.00250
531	1.432	0.103	0.990	0.0349	0.00252
532	1.445	0.104	0.990	0.0356	0.00255
533	1.458	0.104	0.990	0.0362	0.00258
534	1.471	0.103	0.990	0.0369	0.00259
535	1.483	0.103	0.990	0.0375	0.00260
536	1.497	0.103	0.990	0.0382	0.00262
537	1.511	0.102	0.991	0.0389	0.00264
538	1.525	0.102	0.991	0.0396	0.00266
539	1.539	0.102	0.991	0.0404	0.00268
540	1.553	0.102	0.991	0.0411	0.00270
541	1.567	0.102	0.991	0.0419	0.00273
542	1.581	0.102	0.992	0.0426	0.00275
543	1.595	0.102	0.992	0.0433	0.00278
544	1.608	0.102	0.992	0.0440	0.00280
545	1.621	0.102	0.992	0.0448	0.00282
546	1.634	0.103	0.992	0.0455	0.00286
547	1.648	0.103	0.992	0.0463	0.00290
548	1.662	0.104	0.992	0.0471	0.00294
549	1.677	0.104	0.992	0.0479	0.00296
550	1.690	0.103	0.992	0.0487	0.00298
551	1.704	0.103	0.993	0.0495	0.00299
552	1.718	0.103	0.993	0.0503	0.00300
553	1.732	0.102	0.993	0.0511	0.00301
554	1.747	0.102	0.993	0.0520	0.00303
555	1.763	0.102	0.993	0.0530	0.00306
556	1.780	0.102	0.993	0.0540	0.00310

557	1.798	0.102	0.993	0.0551	0.00313
558	1.817	0.102	0.994	0.0562	0.00316
559	1.835	0.102	0.994	0.0574	0.00319
560	1.853	0.102	0.994	0.0585	0.00321
561	1.870	0.101	0.994	0.0596	0.00322
562	1.888	0.101	0.994	0.0607	0.00324
563	1.906	0.100	0.995	0.0619	0.00326
564	1.925	0.100	0.995	0.0631	0.00328
565	1.944	0.099	0.995	0.0644	0.00328
566	1.964	0.099	0.995	0.0657	0.00331
567	1.983	0.098	0.995	0.0670	0.00333
568	2.001	0.098	0.995	0.0683	0.00335
569	2.019	0.098	0.995	0.0695	0.00337
570	2.037	0.098	0.995	0.0707	0.00338
571	2.055	0.097	0.995	0.0720	0.00341
572	2.072	0.097	0.996	0.0732	0.00343
573	2.090	0.097	0.996	0.0744	0.00347
574	2.108	0.098	0.996	0.0757	0.00350
575	2.126	0.097	0.996	0.0770	0.00353
576	2.143	0.098	0.996	0.0782	0.00357
577	2.160	0.098	0.996	0.0795	0.00360
578	2.178	0.097	0.996	0.0808	0.00362
579	2.197	0.097	0.996	0.0822	0.00364
580	2.216	0.097	0.996	0.0837	0.00366
581	2.237	0.097	0.996	0.0853	0.00370
582	2.257	0.097	0.996	0.0868	0.00373
583	2.276	0.097	0.996	0.0882	0.00376
584	2.296	0.097	0.996	0.0898	0.00379
585	2.317	0.097	0.996	0.0914	0.00382
586	2.337	0.097	0.997	0.0930	0.00388
587	2.358	0.097	0.997	0.0947	0.00391
588	2.380	0.097	0.997	0.0965	0.00393
589	2.402	0.097	0.997	0.0983	0.00395
590	2.424	0.096	0.997	0.1001	0.00398
591	2.446	0.096	0.997	0.1020	0.00400
592	2.468	0.096	0.997	0.1038	0.00402

593	2.489	0.096	0.997	0.1055	0.00405
594	2.512	0.095	0.997	0.1076	0.00407
595	2.536	0.095	0.997	0.1096	0.00412
596	2.560	0.096	0.997	0.1116	0.00417
597	2.583	0.096	0.997	0.1137	0.00421
598	2.606	0.096	0.997	0.1158	0.00426
599	2.630	0.096	0.997	0.1179	0.00431
600	2.653	0.096	0.997	0.1200	0.00435