# Tracing devastating fires in Portugal to a snow archive in the Swiss Alps: a case study

Dimitri Osmont<sup>1,2,3,a</sup>, Sandra Brugger<sup>3,4,\*</sup>, Anina Gilgen<sup>5,\*</sup>, Helga Weber<sup>3,6,\*</sup>, Michael Sigl<sup>1,3</sup>, Robin L. Modini<sup>7</sup>, Christoph Schwörer<sup>3,4</sup>, Willy Tinner<sup>3,4</sup>, Stefan Wunderle<sup>3,6</sup>, Margit Schwikowski<sup>1,2,3</sup>

- <sup>1</sup>Laboratory of Environmental Chemistry, Paul Scherrer Institut, 5232 Villigen, Switzerland
   <sup>2</sup>Department of Chemistry and Biochemistry, University of Bern, 3012 Bern, Switzerland
   <sup>3</sup>Oeschger Centre for Climate Change Research, University of Bern, 3012 Bern, Switzerland
   <sup>4</sup>Institute of Plant Sciences, University of Bern, 3012 Bern, Switzerland
   <sup>5</sup>Institute for Atmospheric and Climate Science, ETH Zürich, 8092 Zürich, Switzerland
   <sup>6</sup>Institute of Geography, University of Bern, 3012 Bern, Switzerland
   <sup>7</sup>Laboratory of Atmospheric Chemistry, Paul Scherrer Institut, 5232 Villigen, Switzerland
  - \*These authors equally contributed to this work. \*Now at Institut des Géosciences de l'Environnement, Université Grenoble-Alpes, 38400 Saint Martin d'Hères, France

Correspondence to: Margit Schwikowski (margit.schwikowski@psi.ch)

- 15 Abstract. Recent large wildfires, such as those in Portugal in 2017, have devastating impacts on societies, economy, ecosystems and environments. However, wildfires are a natural phenomenon, which has been exacerbated by land use during the past millennia. Ice cores are one of the archives preserving information on fire occurrences over these timescales. A difficulty is that emission sensitivity of ice cores is often unknown, which constitutes a source of uncertainty in the interpretation of such archives. Information from specific and well-documented case studies is therefore useful to better
- 20 understand the spatial representation of ice-core burning records. The wildfires near Pedrógão Grande in Central Portugal in 2017 provided a test bed to link a fire event to its footprint left in a high-alpine snowpack considered a surrogate for highalpine ice-core sites. Here, we (1) analyzed black carbon (BC) and microscopic charcoal particles deposited in the snowpack close to the high-alpine research station Jungfraujoch in the Swiss Alps, (2) calculated backward trajectories based on ERA-Interim reanalysis data and simulated the transport of these carbonaceous particles using a global aerosol-climate model, and
- (3) analyzed the fire spread, its spatial and temporal extent, as well as its intensity, with remote sensing (e.g. MODIS) active fire and burned area products. According to modelled emissions of the FINN v1.6 database, the fire emitted a total amount of 203.5 tons BC from a total burned area of 501 km<sup>2</sup> as observed on the basis of satellite fire products. Backward trajectories unambiguously linked a peak of atmospheric equivalent BC observed at the Jungfraujoch research station on 22<sup>nd</sup> June, with elevated levels until the 25<sup>th</sup> June, with the highly intensive fires in Portugal. The atmospheric signal is in correspondence with
- 30 an outstanding peak in microscopic charcoal observed in the snow layer, depositing nearly as many charcoal particles as during an average year in other ice archives. In contrast to charcoal, the amount of atmospheric BC deposited during the fire episode was minor due to a lack of precipitation. Simulations with a global aerosol climate model supported that the observed microscopic charcoal particles originated from the fires in Portugal and that their contribution to the BC signal in snow was negligible. Our study revealed that microscopic charcoal can be transported over long distances (1500 km), and that snow and

ice archives are much more sensitive to distant events than sedimentary archives, for which the signal is dominated by local 35 fires. The findings are important for future ice-core studies, as they document that for BC as fire tracer the signal preservation depends on precipitation. Single events, like this example, might not be preserved due to unfavorable meteorological conditions.

#### **1** Introduction

- 40 Fires are an important component of terrestrial ecosystems as they substantially control vegetation cover and contribute to the global carbon budget (e.g. Bond et al., 2005; Hantson et al., 2015). Global CO<sub>2</sub> emissions from fires, including landscape and biomass (i.e. biomass combustion from domestic and industrial uses), represent around 50% of those produced by fossil fuel burning (Bowmann et al., 2009). Fires exert an influence on the climate system as they emit greenhouse gases and aerosols (Andreae and Merlet, 2001) and change surface albedo (Randerson et al., 2006), as well as on vegetation and soil carbon (Page
- 45 et al., 2002). In the last two decades, the occurrence of devastating wildfires has increased in many regions of the world, leading to substantial socioeconomic and environmental consequences (Moritz et al., 2014). In the context of global warming, fire risk (Pechony et al., 2010), frequency (Keywood et al., 2013), and season severity (Flannigan et al., 2013) are potentially increasing with significant feedbacks on terrestrial and atmospheric systems (Bowman et al., 2011). To understand these future impacts, paleofire reconstructions provide an important tool for assessing long-term changes in past fire activity and can help
- 50 to disentangle the influence of climate and humans on biomass burning. Most of the currently available sedimentary records have large chronological uncertainties in the youngest part of the records (Marlon et al., 2016) and reflect small local to regional catchments (Adolf et al., 2018). To address global fire activity trends, ice cores from polar (see e.g. Arienzo et al., 2017; Fischer et al., 2015; Keegan et al., 2014; Legrand et al., 1992; Legrand and De Angelis, 1996, Legrand et al., 2016; Whitlow et al., 1994; Zennaro et al., 2014) and high-altitude glaciers (see e.g. Brugger et al., 2018a, 2019a; Eichler et al., 2011; Osmont
- 55 et al., 2018, 2019; Yalcin et al., 2006) have a high potential since they record fire activity from regional to continental scales and usually provide well-constrained chronologies (see e.g. Herren et al., 2013; Konrad et al., 2013; Uglietti et al., 2016). Fires emit a wide range of chemical compounds and particles to the atmosphere, such as black carbon (BC) and charcoal. If transported over long distances, these particles can be deposited with precipitation or by gravitational settling on the snowpack, where they will be archived, and can be subsequently retrieved by ice-core drilling. However, ice-core catchment areas are not
- often precisely known due to a lack of information about emission, transport and deposition processes. Case studies are 60 essential to understand these processes including fire extent, fuel load, plume transport, biomass burning tracer deposition and preservation in the snowpack (e.g. Kaspari et al., 2015). Modelling of fire tracers (e.g. BC or charcoal) from the fire source to the deposition site provides a direct link and quantification of the archived biomass burning tracers. Detailed information to quantify the amount, frequency and intensity of biomass burning emissions relies mainly on satellite observations (e.g. 65 occurrence of active fires, fire radiative power (i.e. intensity; FRP), burned area and vegetation cover). A limiting factor is the

availability of long-term satellite products with sufficient temporal and spatial resolution. Recently, an active fire product was

developed for Europe reaching back until 1985 (see Weber and Wunderle, 2019). Therefore, a direct linkage between specific fire events and ice core observations is very challenging, especially for biomass burning events that occurred some decades or centuries ago. Thus, the fire footprint of ice-core sites and the preservation of single biomass burning events in these archives

- 70 remain largely unknown. Pioneer case studies have shown that elevated concentrations of ammonium, potassium and formate in Greenland's atmosphere and snow could be directly linked to forest fires in Canada (Dibb et al., 1996). More recent case studies confirmed that BC emissions from fires can be preserved in snow and ice archives at sub-continental scales. For instance, BC peaks in Greenland snowpits were associated with specific biomass burning events in Canada with the help of remote sensing and modelling tools (Thomas et al., 2017). BC emissions from the oil well fires in Kuwait in 1991 during the
- 75 Gulf War were detected unambiguously in an ice core from Muztagh Ata, Northern Tibet (Zhou et al., 2018). However, for Europe, characterized by highly fragmented landscapes and smaller mean burned area compared to other continents (Mouillot and Field, 2005), specific case studies are missing.

Here, we focus on an outstanding wildfire event starting on 17<sup>th</sup> June 2017 in Portugal. Portugal was affected by a severe heat wave with temperatures above 40 °C in June 2017 and during a dry thunderstorm, lightning ignited the forests dominated by

- 80 non-native *Eucalyptus* plantations near Pedrógão Grande in Central Portugal (Fig. 1). The highly flammable vegetation resulted in a rapid and uncontrolled spread of the fire with devastating impacts, leading to the worst death toll Portugal ever experienced for a fire. 64 people lost their life and 254 were injured (Gómez-González et al., 2018) until the fires were finally extinguished on 24<sup>th</sup> June. A large plume of smoke was emitted and transported to the northeast from the burning site. On 21<sup>st</sup> June, the automatic lidar operated by MeteoSwiss (Federal Office of Meteorology and Climatology, Switzerland) in Payerne,
- Switzerland, detected a layer of smoke between 3000 and 5000 m a.s.l. corresponding to the arrival of the plume. The signal intensified on 22<sup>nd</sup> June and its Portuguese origin was confirmed by atmospheric backward trajectory analysis (MeteoSwiss, 2017). The smoke layer became visible in the morning hours of 22<sup>nd</sup> June on the Jungfraujoch (JFJ) research station webcam, located at 3580 m a.s.l. in the Bernese Alps. At the same time, a peak in atmospheric equivalent black carbon (eBC) was detected by the Multi-Angle Absorption Photometer (MAAP, optical method for BC quantification, Petzold and Schönlinner,
- 90 2004) installed at the research station (Fig. 2a). The term eBC is used for black carbon data derived from optical absorption methods, together with a suitable mass absorption cross-section (MAC) for the conversion of light absorption coefficient into mass concentration (Petzold et al., 2013). Elevated eBC values lasted until 25<sup>th</sup> June when the first snowfall after the event occurred.

The exceptional conditions of this Portuguese fire event producing a plume that was recorded unambiguously at the high-

95 alpine research station JFJ in Switzerland, located close to perennial snow archives, provided an ideal situation to analyze in detail the processes related to emission, transport and deposition of fire tracers in a high-mountain snowpack.

# 2 Methods

## 2.1 Study site and meteorological conditions

The high-altitude research station Jungfraujoch (46°32′ N 7°59′ E, Fig. 1) is located on the pass between the summits of Jungfrau and Mönch in the Bernese Alps. Built at an altitude of 3580 m a.s.l., the surrounding high-alpine environment is characterized by glaciers and cliffs. The north side, facing the Swiss Plateau, consists of a steep ice fall while the relatively flat south slopes host the large Jungfraufirn glacier feeding the Aletsch glacier. The JFJ site lies only partially within the free troposphere, being frequently influenced by uplifted air from the planetary boundary layer (e.g. Baltensperger et al., 1997; Bukowiecki et al., 2016). Aerosol monitoring at JFJ provides continuous long-term atmospheric records of many parameters

- 105 including eBC concentration. In addition, a weather station provides meteorological data, but precipitation is not monitored at JFJ due to the difficulty of obtaining accurate data as most precipitation occurs in form of snow in frequent association with strong winds. The closest precipitation data available is from the automatic weather station Itramen (2162 m a.s.l.), operated by the Institute for Snow and Avalanche Research (WSL-SLF) and located 9 km to the northwest of JFJ. Following the Portugal fires, precipitation (rainfall) was recorded at Itramen on 25<sup>th</sup>, 26<sup>th</sup>, 27<sup>th</sup>, 28<sup>th</sup> and 29<sup>th</sup> June, bringing 32, 42.8, 2.0, 49.6 and 35.2
- 110 mm of water, respectively (data from SLF © 2019, SLF). We used the JFJ webcam to determine whether snowfall occurred at the same time at JFJ, which was the case except for 27<sup>th</sup> June.

## 2.2 Snowpit and sampling

A snowpit was collected on 30<sup>th</sup> June 2017, around 20 m off the prepared trail between JFJ and Mönchsjochhütte and 400 m east from the Jungfraujoch tunnel exit onto the glacier, at an altitude of 3460 m a.s.l. A massive ice layer prevented us from

115 digging deeper than 1.10 m. Density was measured on the spot for each layer, by weighing a stainless steel cylinder of known volume filled with snow. Following the stratigraphic study, 20 samples were collected at about 5 cm resolution by pushing pre-cleaned 50-mL polypropylene vials vertically in the snow wall. Ice layers were sampled specifically (if thick enough) to check the potential impact of melting on the chemical composition of the snowpack. 20 replicate samples were retrieved at the same resolution to test the reproducibility of the experiment. In addition, 6 pre-cleaned 1-L PETG jars were filled with 0.2 to 0.5 kg snow for microscopic charcoal analysis, at 10 cm resolution between 20 and 80 cm depth.

# 2.3 Analytical methods

Samples were stored in frozen state and were melted just before analysis of refractory BC (rBC) and major ions at the Paul Scherrer Institute. rBC stands for black carbon measured by incandescence methods (Petzold et al., 2013) and analysis followed the method described by Wendl et al. (2014): after sample melting at room temperature and a 25-min sonication in a ultrasonic

125 bath, rBC was quantified using a Single Particle Soot Photometer (SP2, Droplet Measurement Technologies, USA, Schwarz et al., 2006; Stephens et al., 2003) coupled to an APEX-Q jet nebulizer (Elemental Scientific Inc., USA). Further analytical details regarding calibration, reproducibility and autosampling method can be found in Osmont et al. (2018). The samples were subsequently analyzed for 13 ions (5 cations: ammonium  $NH_4^+$ , calcium  $Ca^{2+}$ , magnesium  $Mg^{2+}$ , potassium  $K^+$  and sodium  $Na^+$ ; and 8 anions: acetate  $CH_3COO^-$ , chloride  $Cl^-$ , fluoride  $F^-$ , formate  $HCOO^-$ , methanesulfonate (MSA)  $CH_3SO_3^-$ , nitrate  $NO_3^-$ , oxalate  $C_2O_4^{2-}$  and sulfate  $SO_4^{2-}$ ) by ion chromatography (850 Professional IC, Metrohm, Switzerland).

- To estimate microscopic charcoal concentrations, we added a known amount of *Lycopodium* marker spores to the six snow samples dedicated to palynological analyses, and prepared the samples following the evaporation-based protocol for ice samples developed by Brugger et al. (2018b). Microscopic charcoal particles were identified as completely opaque particles with a major axis >10 µm and with angular shape following Tinner and Hu (2003). We counted to a combined sum of 200
- 135 items (microscopic charcoal + *Lycopodium* spores; Finsinger and Tinner, 2005).

# 2.4 Air mass trajectories and transport simulations

130

140

To link the observations in the snowpack with the fire emissions in Portugal, we calculated 5-day backward trajectories at a 6-hourly resolution with the Lagrangian analysis tool LAGRANTO (Sprenger and Wernli, 2015) based on ERA-Interim reanalysis data (Dee et al., 2011). The trajectories started at the coordinates of JFJ, at 20 equidistant levels in pressure coordinates between 700 hPa and 500 hPa, in accordance with the detected smoke plume height.

- We used the global aerosol-climate model ECHAM6.3-HAM2.3 to simulate the transport of BC and microscopic charcoal (Gilgen et al., 2018; Stier et al., 2005; Zhang et al., 2012). To the best of our knowledge, this is the only model with a charcoal module implemented. The large-scale wind velocities of the simulations were nudged towards ERA-Interim reanalysis data. Two simulations were conducted: i) a simulation without any fire emissions and ii) a simulation including the fire emissions
- 145 that occurred in Central Portugal (-8.45° to -7.85° E; 39.75° to 40.25° N) in June. Daily fire emission fluxes were calculated from the Fire Inventory from NCAR (FINN version 1.6.; Wiedinmyer et al., 2011) and interpolated to the model grid. For anthropogenic aerosol emissions, we used ACCMIP (Atmospheric Chemistry and Climate Model Intercomparison Project) interpolated emissions for the year 2008 (Lamarque et al., 2010), being constant over the year. A spatial resolution of T127L95 was applied (≈100 km at the equator). To simulate the microscopic charcoal particles, we chose a density of 0.6 g cm<sup>-3</sup>, a
- 150 geometric mean radius upon emission of 5 μm, a threshold radius of 4.9 μm and a scaling factor of BC mass emissions of 250. These parameters were extensively tested, compared and validated by Gilgen et al. (2018).

#### 2.5 Satellite observations

To analyze the fire source itself, i.e. the spatial and temporal extent as well as the intensity of the burning near Pedrógão Grande, we used different fire products retrieved from satellite imagery over the study area. The fires were detected by several

155 satellite sensors and during multiple overpasses (e.g. Visible Infrared Imaging Radiometer Suite (VIIRS), Moderate Resolution Imaging Spectroradiometer (MODIS) or Advanced Very High Resolution Radiometer (AVHRR)), which allows to study the temporal evolution (i.e. progression) of the fires. In consistency with FINN v1.6, we chose the Thermal Anomalies & Fire Daily L3 Global Product, which is also utilized for the emission model. In addition, we obtained the Burned Area Monthly L3 Global Product of MODIS.

- 160 The MODIS sensors on board NASA's Earth Observing System Terra and Aqua satellites have a return period of one to two days with a daytime equator crossing time at 10:30 am (1:30 pm) for Terra (Aqua). The areal extent of the study site was defined as -8.45° to -7.85° longitude and 39.75° to 40.25° latitude according to the location of the forest fires and also used for the modelling (see Section 2.4). For June 2017, we downloaded the respective tiles over the study area of both products: i) the 1 km daily thermal anomaly and fires MOD14A1 (MODIS/Terra) and MYD14A1 (MODIS/Aqua: both V006; Giglio et
- 165 al., 2015a) products and ii) the monthly 500m Burned Area Product MCD64A1 (V006; Giglio et al., 2015b). The latter is a combined product of the burned area detected by the MODIS sensors onboard the Terra and Aqua satellites. It was resampled to a spatial resolution of 1 km. The tiles of the MOD14A1/MYD14A1 products were mosaicked into one image. We lastly reprojected the scenes of both products accordingly to the FINN emission data (version 1.6; Wiedinmyer et al., 2011). Both products were analyzed in terms of active fire progression, fire radiative power (FRP), burned area date and extent, and cloud
- 170 coverage. Moreover, we compared them with the spatial distribution of emission species, like BC,  $NO_x$  and  $PM_{2.5}$ , derived from the FINN database.

To detect the fire plume, we analyzed different daily Aerosol Optical Depth (AOD) products from OMAEROe v003 at 0.25° (smallest spatial resolution in the GIOVANNI database), MODIS-Terra MOD08\_D3 v6.1. at 1° spatial resolution, and from the Modis sensor flying onboard of the satellites Terra and Aqua at a spatial resolution of 10 km at nadir and their area-averaged time series for 17<sup>th</sup> until 26<sup>th</sup> of June.

175 time series for 17<sup>th</sup> until 26<sup>th</sup> of Jur

#### **3 Results and discussions**

### 3.1 Snowpit profile

The snowpit profile (Fig. 2b) showed five layers with different grain size and density (Fig. 2c). Layer A, from 0 to 10 cm depth (d = 0.2 g cm<sup>-3</sup>), was composed of fresh and very light powder snow corresponding to the snowfall on 29<sup>th</sup> June. Layer B, from 10 to 22 cm (d = 0.37, g cm<sup>-3</sup>) was made of light snow probably originating from the snowfall on 28<sup>th</sup> June. Snow in layer C, from 22 to 50 cm (d = 0.54 g cm<sup>-3</sup>), already experienced some transformation as bigger (2–3 mm) round-shaped grains were observed that could relate to the snowfalls that occurred on 25<sup>th</sup> and 26<sup>th</sup> June. For those days, a good agreement was obtained between the precipitation amount from Itramen and the snow layer height corrected for density (Fig. 2a), confirming our time attribution. Below 50 cm depth, accurate dating becomes impossible, but the frequent presence of ice layers and more compact

185 snow indicates melting due to warmer temperatures. This is in line with the June 2017 heat wave in Switzerland that lasted from 19<sup>th</sup> to 24<sup>th</sup> June (MétéoSuisse, 2017). Layer D, from 50 to 76 cm depth, was composed of denser and compact snow (d = 0.55 g cm<sup>-3</sup>). Lastly, below 78 cm depth, layer E seemed even more compact, although the density did not significantly change (d = 0.52 g cm<sup>-3</sup>).

## 3.2 Fire tracers: rBC, microscopic charcoal and major ions

- 190 A peak with concentrations up to 9.8 ng g<sup>-1</sup> is visible in the rBC profile (Fig. 2d) from 22 to 37 cm depth (samples 4 to 6; layer C), corresponding to the first snowfalls recorded after the event. This suggests that atmospheric BC was probably scavenged by snow on 25<sup>th</sup> and 26<sup>th</sup> June, which is in agreement with a drop in atmospheric eBC concentration observed simultaneously (Fig. 2a). Wet deposition seems to be the preferential pathway as the rBC peak is spread over the whole accumulated snow layer while dry deposition would rather create a thin and highly concentrated layer. Several studies indicate that rBC is mainly
- 195 scavenged from the atmosphere via wet deposition processes (Cape et al., 2012; Ruppel et al., 2017; Sinha et al., 2018). The uppermost two layers (A–B, samples 1 to 3) show very low rBC concentrations (average: 0.21 ng g<sup>-1</sup>), in agreement with the clean atmospheric conditions that prevailed on 28<sup>th</sup> and 29<sup>th</sup> June with eBC concentrations below 10 ng m<sup>-3</sup>. Below the rBC peak, a secondary rBC maximum with 4.5 ng g<sup>-1</sup> was observed between 60 and 70 cm in one replicate series, but otherwise average concentrations are low (2.0 ng g<sup>-1</sup>). Except for this secondary maximum, a very good agreement is obtained between
- 200 the two series of replicate rBC samples (r = 0.90, all samples).

The microscopic charcoal record (Fig. 2e) shows a smaller peak at 60–70 cm depth (4000 fragments L<sup>-1</sup>), and the main maximum at 30–40 cm depth, reaching a factor of 10 higher concentrations (20 000 fragments L<sup>-1</sup>) compared to the average of all six samples. We relate the main peak to the fire event in Portugal. Compared to the rBC peak, the narrower microscopic charcoal peak could indicate a more important contribution of dry deposition versus wet deposition, consistent with its larger

205 particle size (> 10 μm major axis for microscopic charcoal vs. < 1 μm diameter for rBC). Microscopic charcoal is a specific proxy for biomass burning, whereas rBC is less specific and can also originate from fossil fuel combustion. No ice layer was observed in the uppermost 50 cm of the snowpit, which indicates that both rBC and microscopic charcoal profiles were not affected by melting processes.

The mass size distribution for the broadband low gain (BBLG) channel of the SP2 remained similar throughout the profile

- 210 with 306 nm and 291 nm for the two series of replicates, respectively (Fig. 2d). This indicates that rBC size distributions in the falling snow are rather stable, as already observed by Sinha et al. (2018). Only sample 13 shows a bigger fraction of large rBC particles, in association with a small peak in rBC and charcoal concentration. On the contrary, no shift in the rBC size distribution is visible for samples 4 to 6 during the peak in rBC and charcoal concentrations.
- Continental-scale calibrations comparing satellite-based fire incidence and charcoal influx into lake sediment or peat bogs suggest that microscopic charcoal particles > 10 µm mainly originate from regional sources with an average radius of ca. 40 km (Adolf et al., 2018). However, previous studies of charcoal in ice cores indicate that charcoal can be transported over distances larger than 500 km (e.g. Brugger et al., 2018a; Eichler et al., 2011; Reese et al., 2013, Hicks and Isaksson, 2006). These differences can be explained by dissimilar environmental conditions, as lake sediment and peat bog studies are usually performed at low-elevation sites surrounded by potential burning sources while snowpits and ice cores originate from high-
- 220 latitude or high-altitude sites remote from vegetation (Brugger et al., 2018a, 2019b). These remote sites frequently lie within the free troposphere: for JFJ, free troposphere background conditions are observed about 39% of the time, with a maximum of

60% in winter and a minimum of 20% in summer (Bukowiecki et al., 2016). However, a travel distance from Portugal to JFJ of around 1500 km seems exceptional. Given the strong intensity of this fire (see Section 3.5), we hypothesize that convection lifted a large amount of microscopic charcoal particles high up in the atmosphere enabling the long-distance travel. The

- 225 microscopic charcoal peak concentrations observed in this study are outstandingly high and only comparable to exceptional peak events in other ice cores (Brugger et al., 2018a; Eichler et al., 2011; Reese et al., 2013), which might be due partly to the highly resolved sampling. This is in contrast to rBC, for which such peak concentrations have been observed in the preindustrial part of European high-altitude ice cores, when rBC mainly originated from biomass burning (Lim et al., 2017; Sigl et al., 2018).
- In addition to rBC and charcoal, ions such as ammonium ( $NH_4^+$ ), formate ( $HCOO^-$ ), acetate ( $CH_3COO^-$ ) and nitrate ( $NO_3^-$ ) have been used as fire tracers (Arienzo et al., 2017; Fischer et al., 2015; Legrand et al., 2016; Savarino and Legrand, 1998), although other emission sources exist, such as direct biogenic emissions (ammonium, formate), oxidation from volatile organic compounds (formate, acetate) and anthropogenic sources (agriculture for ammonium, traffic and agriculture for nitrate). Except for a few values, a good agreement was obtained between the two sets of replicates of the major ion profiles, particularly for
- 235 ammonium and nitrate (Fig. 3). None of the ionic fire tracers shows increased concentrations at the depth of the rBC peak (22-37 cm).

We hypothesize that charcoal originating from the Portugal fires was deposited by dry deposition during the period 22<sup>nd</sup> to 24<sup>th</sup> June, when the fire plume arrived at the JFJ as detected by elevated atmospheric eBC concentrations. Since no snowfall occurred during that period, the majority of BC was not deposited. Dry deposition most likely resulted in a confined charcoal

240 layer, which was not separated from the rBC peak in the snowpit due to the coarse sampling resolution of 10 cm. With the beginning of snowfall on 25<sup>th</sup> June, air mass transport changed, ending the advection of the fire plume to the JFJ as indicated by backward trajectories (see below and Fig. 4). Instead more regional, polluted air masses were scavenged, which explains the absence of ionic fire tracers and of a shift in the rBC size distribution.

# 3.3 rBC scavenging ratios

- rBC scavenging ratios (W) were calculated to determine the total scavenging of rBC from air to snow by snowfall, based on the following formula:  $W = \rho C_s/C_a$ , with  $\rho$  the air density in g m<sup>-3</sup>, C<sub>s</sub> the concentration of rBC in snow in ng g<sup>-1</sup> and C<sub>a</sub> the concentration of rBC in air in ng m<sup>-3</sup> (Schwikowski et al., 1995). For C<sub>a</sub>, we considered the daily average eBC concentration on 24<sup>th</sup> June, just before the precipitation starts, while for C<sub>s</sub>, we used the average of samples 4–6 of the snowpit, corresponding to the days 25<sup>th</sup> and 26<sup>th</sup> June when peak values are found. The air density at JFJ of 842 g m<sup>-3</sup> was calculated with the ideal gas law using a temperature of 273.15 K and a pressure of 66000 Pa obtained from MeteoSwiss for the JFJ weather station.
- In our case, the calculation of rBC scavenging ratios (Table 1) is highly dependent on the choice of a mass absorption coefficient (MAC) for converting the light absorption intensity given by the MAAP into an eBC concentration. The objective is that the eBC mass concentrations from the MAAP precisely match the rBC mass concentrations from the SP2, in order not to introduce a methodological bias as two different quantification methods are used. Previous studies suggested a median MAC

- of  $10.2 \pm 3.2 \text{ m}^2 \text{ g}^{-1}$  for JFJ (Liu et al., 2010) or  $11.1 \pm 0.2 \text{ m}^2 \text{ g}^{-1}$  in summer (Cozic et al., 2008). Higher values ( $13.3 \pm 3.0 \text{ m}^2 \text{ g}^{-1}$ ) have also been reported in the case of eBC from a purely biomass burning origin (Schwarz et al., 2008). An upper limit estimate of 20 m<sup>2</sup> g<sup>-1</sup> for the MAC was obtained during the CLACE 2016 summer campaign at JFJ (Motos et al., 2019). This value is of particular interest as both a MAAP and a SP2 were simultaneously sampling the air at JFJ during this campaign. By applying a MAC of 20 m<sup>2</sup> g<sup>-1</sup>, a perfect agreement could be obtained between the BC mass concentrations given by the
- 260 MAAP and those given by SP2 (measurements not shown). Therefore, MACs of 10 and 20 m<sup>2</sup> g<sup>-1</sup> were chosen here (λ = 637 nm) to test the lower and upper limits, leading to a range of scavenging ratios from 41 to 91 (Table 1). Few BC air-to-snow scavenging ratio values are available in the literature, usually ranging from 100 to 150 (Table 2). By comparing our scavenging ratios obtained at JFJ with values from previous studies, it appears that the choice of a MAC value

of 20 m<sup>2</sup> g<sup>-1</sup> seems preferable. Discrepancies can arise from the different locations implying various climatic conditions, from

the use of different methods to quantify BC (light absorption, thermal-optical or incandescence), and from the different relative contribution of wet and dry deposition processes, depending on the location and elevation, with higher scavenging ratios at higher altitudes (Gogoi et al., 2018).

#### 3.4 Atmospheric transport

The 5-day air mass backward trajectory analyses suggest Portugal as a very likely source for the atmospheric eBC (Fig. 4).

- For all the days between the 22<sup>nd</sup> and the 24<sup>th</sup> June, when atmospheric eBC concentrations were high, part of the air masses originated from Portugal, which is also supported by simulations from MeteoSwiss (MeteoSwiss, 2017). The back trajectories with arrival at JFJ on 25<sup>th</sup> June at 18 UTC indicate a major change in the synoptic situation to more north-westerly flow directions. We can only speculate that this happened concomitant with the onset of precipitation, since the timing of the latter is not precisely known. AOD data at 10 km resolution support that between 20<sup>th</sup> and 22<sup>nd</sup> of June the JFJ site received air
- 275 masses with elevated AOD from Portugal. The AOD levels increased over Switzerland mainly north of the Alps during that time, in agreement with the backward trajectories (Fig. 4). From 23<sup>rd</sup> to 25<sup>th</sup> of June the area was cloud covered, so AOD could not be retrieved. Simulations with ECHAM-HAM show no clear difference for BC when we compare the simulation with fires to the simulation without fires (Fig. 5). In our simulations, BC is preferentially removed by wet scavenging. This suggests that the simulated BC deposition peak observed from 26<sup>th</sup> to 28<sup>th</sup> June is mainly reflecting the precipitation pattern. Furthermore,
- 280 other BC sources than fires seem to dominate in our simulations, which is in agreement with our hypothesis that during fire plume arrival at JFJ the majority of fire-related BC was not deposited due to the lack of snowfall. With the beginning of snowfall at JFJ, air mass transport changed and more regional polluted air masses with minor or without fire contribution were scavenged. In addition, the relatively low spatial resolution of the model smooths the topography, resulting in a more efficient vertical mixing in the lower troposphere. As a consequence, overestimated levels of BC from regional anthropogenic sources
- 285 can probably reach the location of JFJ in the model compared to reality. Microscopic charcoal has no other sources than biomass burning in our model. We observe a clear peak in the deposited microscopic charcoal fluxes on the 23<sup>rd</sup> June (Fig. 5), which is predominantly caused by dry removal processes (i.e. dry

deposition and gravitational settling) and near the observed peak on the  $22^{nd}$  June for atmospheric eBC (Fig. 2a). The simulated deposition fluxes are still rather high on the  $25^{th}$  of June, suggesting that dry deposition continued for the entire fire episode at

290 JFJ. An exact temporal match cannot be expected due to the daily resolution of the fire emissions and the rather low spatial resolution of the model together with the dating uncertainties of the snow pit. Nevertheless, this result qualitatively confirms the hypothesis that the microscopic charcoal particles observed at JFJ originated from Portugal.

#### 3.5 Observations by satellites

- Quantification of the occurrence (active fires), intensity (maximum FRP) and extent (burned area) provided by satellite products allows the modelling of biomass burning emissions. Here, we analyzed the temporal and spatial extent as well as the FRP of the wildfires with the MODIS thermal anomaly and fire (MOD14A1/MYD14A1) and the MCD64A1 burned area products together with emission species of the FINN v1.6. database. The fires evolved on the 17<sup>th</sup> June 2017 in the afternoon (Fig. 6a) in a central location between the fire clusters. An additional comparison of the Aqua/MODIS and VIIRS/NPP Active Fire Product (375 m spatial resolution; Schroeder at el., 2014) showed, that the fire was early detected by the VIIRS/NPP Fire
- 300 Product at ~13:46 UTC. The fire was located few kilometres away from a thick cumulonimbus cloud. Though, the overpass time of Aqua/MODIS was almost simultaneous (i.e. 13:45-13:50 UTC), the fire was too small to be detected by the coarser spatial resolution sensor (1 km). This explains why FINN v1.6 has no emission entries for 17<sup>th</sup> June. Later during the day, several fires were already burning according to the burned area product, which indicates a fast spread in south- and northward direction. Clouds obscured observation by the MODIS sensor on the 18<sup>th</sup> June. Therefore, fewer active fires and lower
- 305 maximum FRP were detected on this day, resulting in gaps in the MOD14A1/MYD14A1 product (Fig. 6a). However, the MCD64A1 burned area product shows the day by day evolution of the fires, forming two big fires clusters. The maximum fire activity and vegetation consumption was observed on the 19<sup>th</sup> and 20<sup>th</sup>, as indicated by the burned area and the high maximum FRP values. Less powerful fires on the south and mainly north edges of the two fire clusters burned from the 20<sup>th</sup> to 22<sup>nd</sup> until the fires were completely extinguished on the 24<sup>th</sup> of June.
- The total area burned over these days accumulates to 501 km<sup>2</sup> according to the burned area product. The total area might be underestimated by this product compared to the maximum FRP, which indicates a larger spatial extent of burning. The detection signals of the burned area were probably too low to detect the burned area along the outer edges of the fires in the south and north (Fig. 6a). Nevertheless, this area agrees well with the actual burned area of 470 km<sup>2</sup> for the two major fire outbreaks occurring in the municipalities of Pedrógão Grande and Góis, based on ground observations (CTI, 2017). The
- 315 modelled BC emissions (Fig. 6b), based on the FINN v1.6. database, range from less than 500 kg day<sup>-1</sup> pixel<sup>-1</sup> to over 2250 kg day<sup>-1</sup> pixel<sup>-1</sup>. A total of about 203.5 tons of BC was emitted by this exceptional fire event (Table 3). Due to cloud obscuration and masking on 18<sup>th</sup> June, this value is probably underestimated.

# **3.6 Deposition fluxes**

Deposition fluxes were calculated for rBC and microscopic charcoal by multiplying the concentration of the respective

320 compounds by the snow accumulation corrected for density. Even if these values remain uncertain due to dating limitations of the snowpit and to the lack of detailed snowfall monitoring at JFJ, preventing us from knowing the exact duration of a snowfall event, they constitute the first step towards a quantitative transfer function.

For microscopic charcoal, we observed a total deposition flux of 104 fragments cm<sup>-2</sup> in the snowpit, around 8 times more than the modelled flux of 13.8 fragments cm<sup>-2</sup> (integral over 23<sup>rd</sup> and 24<sup>th</sup> June, Fig. 5). Compared to yearly average fluxes from

325 high-alpine ice archives (Table 4), the estimated influx at JFJ during the event is exceptional and cannot be explained by the somewhat lower altitude of the JFJ site compared to the other alpine ice-core locations. The comparison suggests this single outstanding event deposited nearly as many charcoal particles as during an average year in other ice archives (e.g. Brugger et al. 2018a).

For rBC, the total deposition flux for samples 4, 5, and 6 was 62 ng cm<sup>-2</sup> (average of replicate samples). The integral flux retrieved from the model for 26<sup>th</sup>, 27<sup>th</sup>, and 28<sup>th</sup> June is 195 ng cm<sup>-2</sup> (Fig. 5); a factor of three higher than the observation. Interestingly, the model underestimates the charcoal and overestimates the BC flux. When the model performance was evaluated previously with longer-term charcoal deposition fluxes, the opposite was observed (Gilgen et al., 2018). It was hypothesized that the model overestimates the fluxes at ice core sites because of their high location within complex topography. The model is not able to simulate these high locations correctly, since the surface altitude is constant over the whole grid box;

- i.e. the topography is smoothed. In addition, ice core sites are often located above the top plume height of most fires (Rémy et al., 2017), which may prevent transport of charcoal particles to them. Obviously the Pedrógão Grande case was exceptional, since the plume was transported at elevations between 3000 and 5000 m a.s.l.. In addition, an underestimation of the fire emissions might have played a role for the charcoal flux. Contrary to the effect on charcoal, the smoothed topography in the model and the corresponding more efficient vertical mixing might have resulted in overestimated levels of BC from regional
- anthropogenic sources, explaining the overestimation of the BC flux. Large discrepancies between modelled and actual deposited rBC fluxes were already pointed out by Thomas et al. (2017), who found a factor of 2–100 in a case study from Greenland and advocated for a better description of precipitation scavenging and fire emissions by the models.

# Conclusions

345

In this case study, biomass burning emissions from an outstanding fire event in Portugal in June 2017 were observed at the high-alpine site Jungfraujoch, Swiss Alps, in both the atmosphere and the snowpack. According to satellite observations, the fire burned a total area of 501 km<sup>2</sup> from 17<sup>th</sup> to 24<sup>th</sup> June, in close agreement with ground observations. At least about 203.5 tons of BC were emitted during this event. Atmospheric backward trajectory analyses showed that the resulting plume of smoke traveled three days before reaching Switzerland, leading to a peak in atmospheric eBC at JFJ on 22<sup>nd</sup> June, lasting until 25<sup>th</sup> June when snowfall occurred. Dry deposition of microscopic charcoal resulted in an outstanding peak in the snowpack.

- 350 This event deposited nearly as many charcoal particles as during an average year in other ice archives. Our study highlights that, for microscopic charcoal, snow and ice archives are more sensitive to distant fires than sedimentary archives, due to the special settings at high elevation. For snow and ice archives, it also reveals that a much longer traveling distance (≈1500 km) than previously thought can be reached, with outstandingly high concentrations in the case of events with optimal weather and transport conditions, thus making microscopic charcoal an excellent biomass burning tracer in ice archives. For rBC, in
- 355 contrast, concentrations in the snow were not exceptionally high. In combination with the absence of a peak in ionic fire tracers such as ammonium, this suggest that the majority of atmospheric BC was not deposited during the fire episode due to a lack of precipitation. Instead the observed rBC peak was mostly likely caused by scavenging of air masses containing regional pollution with the beginning of snowfall on 25<sup>th</sup> June, which ended the advection of the fire plume to the JFJ. rBC scavenging ratios were in line with previous studies, giving additional evidence that rBC was predominantly scavenged by wet deposition.
- 360 Simulations with a global aerosol climate model supported that the observed microscopic charcoal particles originated from the fires in Portugal, whereas their contribution to the BC signal in snow was minor. The findings of our case study are important for future ice-core studies, as they document that for BC as fire tracer the signal preservation depends on precipitation and wet deposition. Single events, like this example, might not be preserved due to unfavorable meteorological conditions. Nevertheless, an exhaustive quantification of the process remains challenging due to the intrinsic uncertainties of each

365 parameter, which requires further collaboration between the different disciplines involved.

#### Data availability

The data underlying Figs. 2 and 3 are available in the Supplement.

#### 370 Author contributions

DO designed the project, carried out sampling, performed rBC analyses and wrote the paper. SB performed microscopic charcoal analyses and contributed to the manuscript writing. AG made atmospheric transport simulations and contributed to the manuscript writing. HW retrieved and analyzed satellite data and contributed to the manuscript writing. MiS organized and led the snowpit study, and commented the manuscript. RLM provided the eBC data from JFJ and advice to interpret them, and

375

5 commented the manuscript. CS, WT and SW contributed to the manuscript writing. MaS led the project, contributed to and supervised writing of the manuscript.

# **Competing interests**

The authors declare that they have no conflict of interest.

# Acknowledgements

- 380 We thank the Swiss National Science Foundation (SNF) for granting the Sinergia project "Paleo fires from high-alpine ice cores", which funded this research (CRSII2\_154450). Furthermore, we thank Nicolas Bukowiecki for providing the atmospheric eBC data from Jungfraujoch, the Institute for Snow and Avalanche Research (SLF) for the precipitation data and Sabina Brütsch for ion chromatography analyses. The online eBC measurements at Jungfraujoch were conducted with financial support from MeteoSwiss (GAW-CH aerosol monitoring program) and from the European Union as well as the Swiss State
- 385 Secretariat for Education, Research and Innovation (SERI) for the European Research Infrastructure for the observation of Aerosol, Clouds and Trace Gases (ACTRIS). The International Foundation High Altitude Research Station Jungfraujoch and Gornergrat (HSFJG) is acknowledged for hosting the online aerosol measurements and for giving access for snow sampling. The authors gratefully acknowledge the personal communication and data provided by C. Wiedinmyer, Cooperative Institute for Research and Environmental Sciences (CIRES), University of Colorado Boulder, Boulder, USA. The modelling was
- 390 supported by a grant from the Swiss National Supercomputing Centre (CSCS) under project ID s652. The ECHAM-HAMMOZ model is developed by a consortium composed of ETH Zürich, Max Planck Institut für Meteorologie, Forschungszentrum Jülich, the University of Oxford, the Finnish Meteorological Institute, and the Leibniz Institute for Tropospheric Research, and managed by the Center for Climate Systems Modeling (C2SM) at ETH Zürich. The MOD14A1/MYD14A1 and MCD64A1 data products were retrieved from the online Data Pool, courtesy of the NASA Land Processes Distributed Active
- 395 Archive Center (LP DAAC), USGS/Earth Resources Observation and Science (EROS) Center, Sioux Falls, South Dakota, https://lpdaac.usgs.gov/data\_access/data\_pool.

# References

400

Adolf, C., Wunderle, S., Colombaroli, D., Weber, H., Gobet, E., Heiri, O., van Leeuwen, J. F. N., Bigler, C., Connor, S. E., Gałka, M., La Mantia, T., Makhortykh, S., Svitavská-Svobodová, H., Vannière, B., and Tinner, W.: The sedimentary and remote-sensing reflection of biomass burning in Europe, Global Ecol. Biogeogr., 27, 199–212, 2018.

Andreae, M. O. and Merlet, P.: Emission of trace gases and aerosols from biomass burning, Global Biogeochem. Cy., 15(4), 955–966, 2001.

Arienzo, M. M., McConnell, J. R., Murphy, L. N., Chellman, N., Das, S., Kipfstuhl, S., and Mulvaney, R.: Holocene black carbon in Antarctica paralleled Southern Hemisphere climate, J. Geophys. Res.-Atmos., 122, 2017.

- Baltensperger, U., Gäggeler, H. W., Jost, D. T., Lugauer, M., Schwikowski, M., and Weingartner, E.: Aerosol climatology at the high-alpine site Jungfraujoch, Switzerland, J. Geophys. Res., 102(D16), 19707–19715, 1997.
  Bond, W. J., Woodward, F. I., and Midgley, G. F.: The global distribution of ecosystems in a world without fire, New phytol., 165(2), 525–537, 2005.
  Bowman, D. M. J. S., Balch, J. K., Artaxo, P., Bond, W. J., Carlson, J. M., Cochrane, M. A., D'Antonio, C. M., DeFries, R.
- 410 S., Doyle, J. C., Harrison, S. P., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Marston, J. B., Moritz, M. A.,

Prentice, I. C., Roos, C. I., Scott, A. C., Swetnam, T. W., van der Werf, G. R., and Pyne, S. J.: Fire in the Earth System, Science, 324(5926), 481–484, 2009.

Bowman, D. M. J. S., Balch, J. K., Artaxo, P., Bond, W. J., Cochrane, M. A., D'Antonio, C. M., DeFries, R. S., Johnston, F. H., Keeley, J. E., Krawchuk, M. A., Kull, C. A., Mack, M., Moritz, M. A., Pyne, S. J., Roos, C. I., Scott, A. C., Sodhi, N. S.,

415 and Swetnam, T. W.: The human dimension of fire regimes on Earth, J. Biogeogr., 38(12), 2223–2236, 2011. Brugger, S. O., Gobet, E., Sigl, M., Osmont, D., Papina, T., Rudaya, N., Schwikowski, M., and Tinner, W.: Ice records provide new insights into climatic vulnerability of Central Asian forest and steppe communities, Global Planet. Change, 169, 188–201, 2018a.

Brugger, S. O., Gobet, E., Schanz, F. R., Heiri, O., Schwörer, C., Sigl, M., Schwikowski, M., and Tinner, W.: A quantitative comparison of microfossil extraction methods from ice cores, J. Glaciol., 64(245), 432–442, 2018b.

Brugger, S. O., Gobet, E., Osmont, D., Behling, H., Fontana, S. L., Hooghiemstra, H., Morales-Molino, C., Sigl, M., Schwikowski, M., and Tinner, W.: Tropical Andean glacier reveals colonial legacy in modern mountain ecosystems, Quat. Sci. Rev., 2019a.

Brugger, S.O., Gobet, E., Blunier, T., Morales-Molino, C., Lotter, A. F., Fischer, H., Schwikowski, M., and Tinner, W.:

Palynological insights into global change impacts on Arctic vegetation, fire, and pollution recorded in Central Greenland ice,
 Holocene, 29(7), 1189–1197, 2019b.

Bukowiecki, N., Weingartner, E., Gysel, M., Collaud Coen, M., Zieger, P., Herrmann, E., Steinbacher, M., Gäggeler, H. W., and Baltensperger, U.: A review of more than 20 years of aerosol observation at the high altitude research station Jungfraujoch, Switzerland (3580 m asl), Aerosol Air Qual. Res, 16, 764–788, 2016.

Cape, J. N., Coyle, M., and Dumitrean, P.: The atmospheric lifetime of black carbon, Atmos. Environ., 59, 256–263, 2012.
Clarke, A. D. and Noone, K. J.: Soot in the Arctic snowpack: a cause for perturbations in radiative transfer, Atmos. Environ., 19(12), 2045–2053, 1985.

Cozic, J., Verheggen, B., Weingartner, E., Crosier, J., Bower, K. N., Flynn, M., Coe, H., Henning, S., Steinbacher, M., Henne,
S., Collaud Coen, M., Petzold, A., and Baltensperger, U.: Chemical composition of free tropospheric aerosol for PM1 and
coarse mode at the high alpine site Jungfraujoch, Atmos, Chem. Phys., 8, 407–423, 2008.

CTI – Comissão Técnica Independente, Análise e apuramento dos factos relativos aos incêndios que ocorreram em Pedrógão Grande, Castanheira de Pera, Ansião, Alvaiázere, Figueiró dos Vinhos, Arganil, Góis, Penela, Pampilhosa da Serra, Oleiros e Sertã, entre 17 e 24 de junho de 2017, *Relatório*, Assembleia da República, 296 p., 2017.

435

Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S., Andrae, U., Balmaseda, M. A., Balsamo, G.,
Bauer, P., Bechtold, P., Beljaars, A. C., van de Berg, L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer,
A. J., Haimberger, L., Healy, S. B., Hersbach, H., Hólm, E. V., Isaksen, L., Kållberg, P., Köhler, M., Matricardi, M., McNally,
A. P., Monge-Sanz, B. M., Morcrette, J., Park, B., Peubey, C., de Rosnay, P., Tavolato, C., Thépaut, J., and Vitart, F.: The
ERA-Interim reanalysis: configuration and performance of the data assimilation system, Q.J.R. Meteorol. Soc., 137, 553–597, 2011.

445 Dibb, J. E., Talbot, R. W., Whitlow, S. I., Shipham, M. C., Winterle, J., McConnell, J., and Bales, R.: Biomass burning signatures in the atmosphere and snow at Summit, Greenland: An event on 5 August 1994, Atmos. Environ., 30(4), 553–561, 1996.

Eichler, A., Tinner, W., Brütsch, S., Olivier, S., Papina, T., and Schwikowski, M.: An ice-core based history of Siberian forest fires since AD 1250, Quaternary Sci. Rev., 30, 1027–1034, 2011.

450 Finsinger, W. and Tinner, W.: Minimum count sums for charcoal concentration estimates in pollen slides: accuracy and potential errors, Holocene, 15(2), 293–297, 2005.
Fischer, H., Schüpbach, S., Gfeller, G., Bigler, M., Rothlisberger, R., Erhardt, T., Stocker, T. F., Mulvaney, R., and Wolff, E.: Millennial changes in North American wildfire and soil activity over the last glacial cycle, Nat. Geosci., 8, 723–728, 2015.

Flannigan, M., Cantin, A. S., de Groot, W. J., Wotton, M., Nwebery, A., and Gowman, L. M.: Global wildland fire season severity in the 21<sup>st</sup> century, Forest Ecol. Manag., 294, 54–61, 2013.

- Giglio, L., and Justice, C.: MOD14A1 MODIS/Terra Thermal Anomalies/Fire Daily L3 Global 1km SIN Grid V006 [Data set], NASA EOSDIS LP DAAC, doi: 10.5067/MODIS/MOD14A1.006, 2015a.
  Giglio, L., Justice, C., Boschetti, L., and Roy, D.: MCD64A1 MODIS/Terra+Aqua Burned Area Monthly L3 Global 500m SIN Grid V006 [Data set], NASA EOSDIS Land Processes DAAC, doi: 10.5067/MODIS/MCD64A1.006, 2015b.
- 460 Gilgen, A., Adolf, C., Brugger, S. O., Ickes, L., Schwikowski, M., van Leeuwen, J. F. N., Tinner, W., and Lohmann, U.: Implementing microscopic charcoal particles into a global aerosol-climate model, Atmos. Chem. Phys., 18, 11813–11829, 2018.

Gogoi, M. M., Babu, S. S., Moorthy, K. K., Thakur, R. C., Chaubey, J. P., and Nair, V. S.: Aerosol black carbon over Svalbard regions of Arctic, Polar Sci., 10, 60–70, 2016.

Gogoi, M. M., Babu, S. S., Pandey, S. K., Nair, V. S., Vaishya, A., Girach, I. A., and Koushik, N.: Scavenging ratio of black carbon in the Arctic and the Antarctic, Polar Sci., 16, 10–22, 2018.
Gómez-González, S., Ojeda, F., and Fernandes, P. M.: Portugal and Chile: Longing for sustainable forestry while rising from the ashes, Environ. Sci. Policy, 81, 104–107, 2018.
Hantson, S., Pueyo, S., and Chuvieco, E.: Global fire size distribution is driven by human impact and climate, Global Ecol.

470 Biogeogr., 24, 77–86, 2015.
Herren, P.-A., Eichler, A., Machguth, H., Papina, T., Tobler, L., Zapf, A., and Schwikowski, M.: The onset of Neoglaciation

6000 years ago in western Mongolia revealed by an ice core from the Tsambagarav mountain range, Quaternary Sci. Rev., 69, 59–69, 2013.

Hicks, S. and Isaksson, E.: Assessing source areas of pollutants from studies of fly ash, charcoal and pollen from Svalbard snow and ice, J. Geophys. Res., 111, D02113, 2006.

Jacobson, J.Z.: Climate response of fossil fuel and biofuel soot, accounting for soot's feedback to snow and sea ice albedo and emissivity, J. Geophys. Res., 109, D21201, 2004.

Jenk, T. M., Szidat, S., Schwikowski, M., Gaggeler, H. W., Brütsch, S., Wacker, L., Synal, H. A., and Saurer, M.: Radiocarbon analysis in an Alpine ice core: record of anthropogenic and biogenic contributions to carbonaceous aerosols in the past (1650-

Kaspari, S., Skiles, S. M., Delaney, I., Dixon, D., and Painter, T. H.: Accelerated glacier melt on Snow Dome, Mount Olympus, Washington, USA, due to deposition of black carbon and mineral dust from wildfire, J. Geophys. Res.-Atmos., 120(7), 2793–2807, 2015.

Keegan, K. M., Albert, M. R., McConnell, J. R., and Baker, I.: Climate change and forest fires synergistically drive widespread
melt events of the Greenland Ice Sheet, P. Natl. Acad. Sci. USA, 111, 7964–7967, 2014.

Keywood, M., Kanakidou, M., Stohl, A., Dentener, F., Grassi, G., Meyer, C. P., Torseth, K., Edwards, D., Thompson, A. M., Lohmann, U., and Burrows, J.: Fire in the air: Biomass burning impacts in a changing climate, Crit. Rev. Env. Sci. Tech., 43(1), 40–83, 2013.

Konrad, H., Bohleber, P., Wagenbach, D., Vincent, C., and Eisen, O.: Determining the age distribution of Colle Gnifetti, Monte Rosa, Swiss Alps, by combining ice cores, ground-penetrating radar and a simple flow model, J. Glaciol., 59(213), 179–189,

490 Rosa, Swiss Alps, by combining ice cores, ground-penetrating radar and a simple flow model, J. Glaciol., 59(213), 179–189, 2013.

Lamarque, J.-F., Bond, T. C., Eyring, V., Granier, C., Heil, A., Klimont, Z., Lee, D., Liousse, C., Mieville, A., Owen, B., Schultz, M. G., Shindell, D., Smith, S. J., Stehfest, E., Van Aardenne, J., Cooper, O. R., Kainuma, M., Mahowald, N., McConnell, J. R., Naik, V., Riahi, K., and van Vuuren, D. P.: Historical (1850–2000) gridded anthropogenic and biomass

burning emissions of reactive gases and aerosols: methodology and application, Atmos. Chem. Phys., 10, 7017–7039, 2010.
 Legrand, M., McConnell, J., Fischer, H., Wolff, E. W., Preunkert, S., Arienzo, M., Chellman, N., Leuenberger, D., Maselli, O., Place, P., Sigl, M., Schüpbach, S., and Flannigan, M.: Boreal fire records in Northern Hemisphere ice cores: a review, Clim. Past, 12, 2033–2059, 2016.

Legrand, M., and DeAngelis, M.: Light carboxylic acids in Greenland ice: A record of past forest fires and vegetation emissions from the boreal zone, J. Geophys. Res.-Atmos, 101(D2), 4129-4145, 1996.

Legrand, M., De Angelis, M., Staffelbach, T., Neftel, A., and Stauffer, B.: Large perturbations of ammonium and organic acids content in the Summit-Greenland ice core: Fingerprint from forest fires?, Geophys. Res. Lett., 19(5), 473–475, 1992. Lim, S., Faïn, X., Ginot, P., Mikhalenko, V., Kutuzov, S., Paris, J.-D., Kozachek, A., and Laj, P.: Black carbon variability since preindustrial times in the eastern part of Europe reconstructed from Mt. Elbrus, Caucasus, ice cores, Atmos. Chem. Phys.,

505 17, 3489–3505, 2017.

500

480

1940), Atmos. Chem. Phys., 6, 5381–5390, 2006.

- Liu, D., Flynn, M., Gysel, M., Targino, A., Crawford, I., Bower, K., Choularton, T., Jurányi, Z., Steinbacher, M., Hueglin, C.,
  Curtius, J., Kampus, M., Petzold, A., Weingartner, E., Baltensperger, U., and Coe, H.: Single Particle Characterization of
  Black Carbon Aerosols at a Tropospheric Alpine Site in Switzerland, Atmos. Chem. Phys., 10, 7389–7407, 2010.
  Marlon, J. R., Kelly, R., Daniau, A.-L., Vannière, B., Power, M. J., Bartlein, P., Higuera, P., Blarquez, O., Brewer, S., Brücher,
- 510 T., Feurdean, A., Romera, G. G., Iglesias, V., Maezumi, S. Y., Magi, B., Courtney Mustaphi, C. J., and Zhihai, T.:

Reconstructions of biomass burning from sediment-charcoal records to improve data-model comparisons, Biogeosciences, 13, 3225–3244, 2016.

MétéoSuisse, Bulletin Climatologique Juin 2017, Genève, 2017. Available at: https://www. meteosuisse.admin.ch/home/service-et-publications/publications.subpage.html/fr/data/publications/2017/7/

 515 bulletin-climatologique-juin-2017.html?pageIndex=5&tab=search\_tab, last accessed 25.04.2019.
 MeteoSwiss, Cendres des incendies portugais détectées en Suisse, https://www.meteosuisse.admin.ch/home.subpage.html/fr/data/blogs/2017/6/incendies-du-portugal-mesures-en-suisse.html, 2017, last accessed 08.10.2018.

Moritz, M. A., Batllori, E., Bradstock, R. A., Malcolm Gill, A., Handmer, J., Hessburg, P. F., Leonard, J., McCaffrey, S., 520 Odion, D. C., Schoennagel, T., and Syphard, A. D.: Learning to coexist with wildfire, Nature, 515, 58–66, 2014.

Motos, G., Schmale, J., Corbin, J. C., Modini, R. L., Karlen, N., Bertò, M., Baltensperger, U., and Gysel-Beer, M.: Cloud droplet activation properties and scavenged fraction of black carbon in liquid-phase clouds at the high-alpine research station Jungfraujoch (3580m a.s.l.), Atmos. Chem. Phys., 19, 3833–3855, 2019. Mouillot, F. and Field, C.: Fire history and the global carbon budget: a 1° × 1° fire history reconstruction for the 20th century,

Global Change Biol., 11, 398–420, 2005.
Noone, K. J. and Clarke, A. D.: Soot scavenging measurements in Arctic snowfall, Atmos. Environ., 22(12), 2773–2778, 1988.
Osmont, D., Wendl, I. A., Schmidely, L., Sigl, M., Vega, C. P., Isaksson, E., and Schwikowski, M.: An 800-year high-resolution black carbon ice core record from Lomonosovfonna, Svalbard, Atmos. Chem. Phys., 18, 12777–12795, 2018.

- Osmont, D., Sigl, M., Eichler, A., Jenk, T., and Schwikowski, M.: A Holocene black carbon ice-core record of biomass burning 530 in the Amazon Basin from Illimani, Bolivia, Clim. Past, 15, 579–592, 2019.
- Page, S. E., Siegert, F., Rieley, J. O., Boehm, H.-D. V., Jaya, A., and Limin, S.: The amount of carbon released from peat and forest fires in Indonesia during 1997, Nature, 420, 61–65, 2002.
  Pechony, O. and Shindell, D. T.: Driving forces of global wildfires over the past millennium and the forthcoming century, P. Natl. Acad. Sci. USA, 107(45), 19167–19170, 2010.
- Petzold, A. and Schönlinner, M.: Multi-angle absorption photometry a new method for the measurement of aerosol light absorption and atmospheric black carbon, J. Aerosol Sci., 35, 421–441, 2004.
  Petzold, A., Ogren, J. A., Fiebig, M., Laj, P., Li, S. M., Baltensperger, U., Holzer-Popp, T., Kinne, S., Pappalardo, G., Sugimoto, N., Wehrli, C., Wiedensohler, A., and Zhang, X. Y.: Recommendations for reporting "black carbon" measurements, Atmos. Chem. Phys., 13, 8365–8379, 2013.
- 540 Randerson, J. T., Liu, H., Flanner, M. G., Chambers, S. D., Jin, Y., Hess, P. G., Pfister, G., Mack, M. C., Treseder, K. K., Welp, L. R., Chapin, F. S., Harden, J. W., Goulden, M. L., Lyons, E., Neff, J. C., Schuur, E. A. G., and Zender, C. S.: The impact of boreal forest fire on climate warming, Science, 314, 1130–1132, 2006.

Reese, C. A., Liu, K. B., and Thompson, L. G.: An ice-core pollen record showing vegetation response to Late-glacial and Holocene climate changes at Nevado Sajama, Bolivia, Ann. Glaciol., 54(63), 183–190, 2013.

Rémy, S., Veira, A., Paugam, R., Sofiev, M., Kaiser, J.W., Marenco, F., Burton, S. P., Benedetti, A., Engelen, R. J., Ferrare, R., and Hair, J. W.: Two global data sets of daily fire emission injection heights since 2003, Atmos. Chem. Phys., 17, 2921–2942, 2017.

Ruppel, M. M., Soares, J., Gallet, J.-C., Isaksson, E., Martma, T., Svensson, J., Kohler, J., Pedersen, C. A., Manninen, S., Korhola, A., and Ström, J.: Do contemporary (1980–2015) emissions determine the elemental carbon deposition trend at

- 550 Holtedahlfonna glacier, Svalbard?, Atmos. Chem. Phys., 17, 12779–12795, 2017.
  Savarino, J. and Legrand, M.: High northern latitude forest fires and vegetation emissions over the last millennium inferred from the chemistry of a central Greenland ice core, J. Geophys. Res.-Atmos., 103(D7), 8267–8279, 1998.
  Schwarz, J. P., Gao, R. S., Fahey, D. W., Thomson, D. S., Watts, L. A., Wilson, J. C., Reeves, J. M., Darbeheshti, M., Baumgardner, D. G., Kok, G. L., Chung, S. H., Schulz, M., Hendricks, J., Lauer, A., Karcher, B., Slowik, J. G., Rosenlof, K.
- 555 H., Thompson, T. L., Langford, A. O., Loewenstein, M., and Aikin, K. C.: Single-particle measurements of midlatitude black carbon and light-scattering aerosols from the boundary layer to the lower stratosphere, J. Geophys. Res., 111, D16207, 2006. Schroeder, W., Oliva, P., Giglio, L., and Csiszar, I. A.: The New VIIRS 375 m active fire detection data product: algorithm description and initial assessment. Remote Sens. Environ. 143, 85-96. doi: 10.1016/j.rse.2013.12.008, 2014.
- Schwarz, J. P., Gao, R. S., Spackman, J. R., Watts, L. A., Thomson, D. S., Fahey, D. W., Ryerson, T. B., Peischl, J., Holloway,
  J. S., Trainer, M., Frost, G. J., Baynard, T., Lack, D. A., de Gouw, J. A., Warneke, C., and Del Negro, L. A.: Measurement of the mixing state, mass, and optical size of individual black carbon particles in urban and biomass burning emissions, Geophys. Res. Lett., 35, L13810, 2008.

Schwikowski, M., Seibert, P., Baltensperger, U., and Gäggeler, H. W.: A study of an outstanding Saharan dust event at the high-alpine site Jungfraujoch, Switzerland, Atmos. Environ., 29(15), 1829–1842, 1995.

- 565 Sigl, M., Abram, N. J., Gabrieli, J., Jenk, T. M., Osmont, D., and Schwikowski, M.: 19th century glacier retreat in the Alps preceded the emergence of industrial black carbon deposition on high-alpine glaciers, Cryosphere, 12, 3311-3331, 2018. Sinha, P. R., Kondo, Y., Goto-Azuma, K., Tsukagawa, Y., Fukuda, K., Koike, M., Ohata, S., Moteki, N., Mori, T., Oshima, N., Førland, E. J., Irwin, M., Gallet, J.-C., and Pedersen, C. A.: Seasonal Progression of the Deposition of Black Carbon by Snowfall at Ny-Ålesund, Spitsbergen, J. Geophys. Res.-Atmos., 123, 997–1016, 2018.
- 570 Sprenger, M. and Wernli, H.: The LAGRANTO Lagrangian analysis tool version 2.0, Geosci. Model Dev., 8, 2569–2586, 2015.

Stephens, M., Turner, N., and Sandberg, J.: Particle identification by laser-induced incandescence in a solid-state laser cavity, Appl. Optics, 42, 3726–3736, 2003.

Stier, P., Feichter, J., Kinne, S., Kloster, S., Vignati, E., Wilson, J., Ganzeveld, L., Tegen, I., Werner, M., Balkanski, Y.,

575 Schulz, M., Boucher, O., Minikin, A., and Petzold, A.: The aerosol-climate model ECHAM5-HAM, Atmos. Chem. Phys., 5, 1125–1156, 2005.

Thomas, J. L., Polashenski, C. M., Soja, A. J., Marelle, L., Casey, K. A., Choi, H. D., Raut, J.-C., Wiedinmyer, C., Emmons, L. K., Fast, J. D., Pelon, J., Law, K. S., Flanner, M. G., and Dibb, J. E.:: Quantifying black carbon deposition over the Greenland ice sheet from forest fires in Canada, Geophys. Res. Lett., 44, 7965–7974, 2017.

580 Tinner, W. and Hu, F. S.: Size parameters, size-class distribution and area-number relationship of microscopic charcoal: relevance for fire reconstruction, Holocene, 13(4), 499–505, 2003.

Uglietti, C., Zapf, A., Jenk, T. M., Sigl, M., Szidat, S., Salazar, G., and Schwikowski, M.: Radiocarbon dating of glacier ice: overview, optimisation, validation and potential, Cryosphere, 10, 3091–3105, 2016.

Wang, Z. W., Gallet, J. C., Pedersen, C. A., Zhang, X. S., Ström, J., and Ci, Z. J.: Elemental carbon in snow at Changbai
mountain, northeastern China: concentrations, scavenging ratios, and dry deposition velocities, Atmos. Chem. Phys., 14, 629–640, 2014.

Warren, S. G. and Clarke, A. D.: Soot in the atmosphere and snow surface of Antarctica, J. Geophys. Res., 95(D2), 1811–1816, 1990.

Weber and Wunderle, Drifting Effects of NOAA Satellites on Long-Term Active Fire Records of Europe, Remote Sens., 11(4), 467, 2019.

590

Wendl, I. A., Menking, J. A., Färber, R., Gysel, M., Kaspari, S. D., Laborde, M. J. G., and Schwikowski, M.: Optimized method for black carbon analysis in ice and snow using the Single Particle Soot Photometer, Atmos. Meas. Tech., 7, 2667–2681, 2014.

Wiedinmyer, C., Akagi, S. K., Yokelson, R. J., Emmons, L. K., Al-Saadi, J. A., Orlando, J. J., and Soja, A. J.: The Fire

595 Inventory from NCAR (FINN): A High Resolution Global Model to Estimate the Emissions from Open Burning, Geosci. Model Dev., 4, 625–641, 2011.

Whitlow, S., Mayewski, P., Dibb, J., Holdsworth, G., and Twickler, M.: An ice-core-based record of biomass burning in the Arctic and sub-Arctic, 1750-1980, Tellus Series B-Chemical and Physical Meteorology, 46(3), 234-242, 1994.

Yalcin, K., Wake, C. R., Kreutz, K. J., and Whitlow, S. I.: A 1000-yr record of forest fire activity from Eclipse Icefield, Yukon, 600 Canada, Holocene, 16(2), 200–209, 2006.

Zennaro, P., Kehrwald, N., McConnell, J. R., Schüpbach, S., Maselli, O. J., Marlon, J., Vallelonga, P., Leuenberger, D., Zangrando, R., Spolaor, A., Borrotti, M., Barbaro, E., Gambaro, A., and Barbante, C.: Fire in ice: two millennia of boreal forest fire history from the Greenland NEEM ice core, Clim. Past, 10, 1905–1924, 2014.

Zhang, K., O'Donnell, D., Kazil, J., Stier, P., Kinne, S., Lohmann, U., Ferrachat, S., Croft, B., Quaas, J., Wan, H., Rast, S.,

605 and Feichter, J.: The global aerosol-climate model ECHAM-HAM, version 2: sensitivity to improvements in process representations, Atmos. Chem. Phys., 12, 8911–8949, 2012.

Zhou, J., Tie, X., Xu, B., Zhao, S., Wang, M., Li, G., Zhang, T., Zhao, Z., Liu, S., Yang, S., Chang, L., and Cao, J.: Black carbon (BC) in a northern Tibetan mountain: effect of Kuwait fires on glaciers, Atmos. Chem. Phys., 18, 13673–13685, 2018.

Table 1: Values used for the calculation of rBC scavenging ratios (W) at Jungfraujoch, Switzerland. For air concentration (Ca), the610eBC average value of June 24<sup>th</sup> was used, under two different MACs. For snow concentration (Cs), concentrations from the two sets<br/>of replicates were used.

-	MAC $(m^2 g^{-1})$	C <sub>a</sub> (ng m <sup>-3</sup> )	C <sub>s</sub> (ng g <sup>-1</sup> )	W	
_	10	147.5	7.14-8.01	41–46	
	20	73.8	7.14-8.01	81–91	

# Table 2: Examples of BC scavenging ratios available in the literature.

Location	W	BC measurement quantity	Authors
Jungfraujoch (JFJ)	81–91	rBC/eBC	This study
Arctic	160	eBC	Clarke and Noone, 1985
Abisko, Sweden	$97 \pm 34$	eBC	Noone and Clarke, 1988
Antarctica	150	eBC	Warren and Clarke, 1990
N-E China	$140\pm100$	EC	Wang et al., 2014
Svalbard	$98\pm46$	eBC	Gogoi et al., 2016
Antarctic	$120\pm23$	eBC	Gogoi et al., 2018
Global	125	Modelled	Jacobson, 2004

615 Table 3: Burned area and BC emissions per day for the June 2017 forest fire in Portugal (\*cloud covered).

Day	Burned area (km <sup>2</sup> )	BC emissions (tons)
17 <sup>th</sup> June	13	-
18 <sup>th</sup> June	148*	53.7
19 <sup>th</sup> June	243	99.8
20 <sup>th</sup> June	45	25.5
21 <sup>st</sup> June	30	24.5
22 <sup>nd</sup> June	19	-
23 <sup>rd</sup> June	3	-
Total	501	203.5

Table 4: Comparison of microscopic charcoal influx at Jungfraujoch with the yearly average influx at selected glacier sites based on identical laboratory preparation and analytical methods.

Site	Altitude	Microscopic charcoal influx	Deference	
Sile	(m asl)	(fragments cm <sup>-2</sup> year <sup>-1</sup> )	Reference	
Jungfraujoch	3560	104*	This study	
Tsambagarav (Mongolian Altai)	4100	200	Brugger et al. (2018a)	
Colle Gnifetti (Swiss Alps)	4500	390	Gilgen et al. (2018)	
Summit (Central Greenland)	3200	9	Brugger et al. (2019b)	
Illimani (Bolivian Andes)	6300	130	Brugger et al. (2019a)	

620 \*This event only in fragments cm<sup>-2</sup>



Figure 1: Source and deposition sites. a) Map of South-Western Europe with the area of Pedrógão Grande, in Central Portugal, where the fires burned (red box) and the area of the zoom in on the right panel (grey box). b) Map of Switzerland with the sites of interest mentioned in the study. Triangles indicate high-altitude ice-core and snow study sites, circles stand for weather stations.



Figure 2: a) Atmospheric equivalent black carbon (eBC) concentrations at Jungfraujoch showing the peak on  $22^{nd}$  June when the plume of smoke reached the site. A mass absorption coefficient of 10 m<sup>2</sup> g<sup>-1</sup> was assumed. Blue bars indicate days with significant snowfall at JFJ, with a comparison between the daily precipitation amount measured at Itramen (values in blue, in mm, data from

630 SLF © 2019, SLF) and the daily snowfall height inferred from the snowpit at JFJ, corrected for density (values in orange, in mm water equivalent (weq)). b) Snowpit stratigraphy with ice lenses in dark blue. c) Density profile of the snowpit. d) rBC concentration profile (top scale) with the two series of replicates (black and grey) and sample number associated. Mode of the rBC mass size distribution (bottom scale) with the two series of replicates (dark red and red). Blue bars are ice lenses. e) Microscopic charcoal concentration record.



Figure 3: Ionic records from the Jungfraujoch snowpit with the two sets of replicates (darker/lighter lines). The orange bar indicates the depth at which rBC and microscopic charcoal peaks are observed. Blue bars represent ice layers. Sample numbers are specified for calcium and are similar for the other ions. Potential sources are indicated above each graph.



Figure 4: 5-day air mass backward trajectories starting from the Jungfraujoch site at 00 UTC on 22<sup>nd</sup> to 24<sup>th</sup> June 2017. For 25<sup>th</sup> June starting time is 18 UTC.



Figure 5: Simulated deposition fluxes of BC and microscopic charcoal at Jungfraujoch. In line with the observations, the deposition (dep) fluxes for BC are given as mass fluxes, whereas the deposition fluxes of microscopic charcoal are given as the number fluxes of particles larger than a certain threshold (10 µm major axis). For BC, a simulation without fire emissions (dotted gray line) and a simulation including the fire emissions near Pedrógão Grande in Central Portugal (solid gray line) are shown. For microscopic charcoal, only the simulation including fire emissions (solid orange line) is shown since biomass burning is the only source of microscopic charcoal.



655 Figure 6: Spatial and temporal observed wild fires and their BC emissions near Pedrógão Grande, Portugal, for the 17<sup>th</sup> to 23<sup>th</sup> of June at 1km spatial resolution. a) Date of the area burned (MCD64A1) and maximum Fire Radiative Power (FRP) values of the active fires according to MOD14A1/MYD14A1 showing the evolved two fire clusters. b) Corresponding BC emission values based on the FINN v1.6 database with high values at the outer edges.