

Interactive comment on “Impact of coastal East Antarctic ice rises on surface mass balance: insights from observations and modeling” by Thore Kausch et al.

Thore Kausch et al.

t.kausch@tudelft.nl

Received and published: 23 June 2020

TC Response to referee 2

C1

Dear referee,

First of all thank you for taking your time read through our manuscript as well as for your positive review and constructive criticism. The points you are raising are all very much justified and we tried our best to address them as good as possible.

Below you can find your original comments with our answers in [blue](#) as well as updated sections from the text in *italic*.

Yours sincerely,
Thore Kausch and co-authors

1 Referee 2 comments:

Interactive comment on “Impact of coastal East Antarctic ice rises on surface mass balance: insights from observations and modeling” by Thore Kausch et al.

1.1 General

This paper presents data and model results for surface mass balance in the region of two East Antarctic ice rises. The authors show using ground-penetrating-radar-based estimates of SMB and results from RACMO2 and SnowMod el that SMB varies on a local scale, due to erosion and redeposition by wind, and on a regional scale, due to orographic lifting. Neither of these results are surprising, but the paper argues the implications for ice-core records recovered on ice rises.

C2

1.2 Main issues

In section 2.5 a number of parameters are described about the SnowModel, and it's not that clear if the 'knob-twiddling' was to match observations or actually based on a-priori information and assumptions. It would be good to make this more clear.

For the roughness length and the threshold shear velocity we indeed used forward modelling to try several different values and in the end chose the ones, which would fit the observations best. This was not clear in the text before. We added a few sentence to clarify this in the text.

L182-184: For both, the roughness length and the threshold shear velocity we tried different values using forward modelling. We varied the roughness length between 0.05 and 0.00005 m and the threshold shear velocity between 0.3 and 0.8 m/s, and chose the values which qualitatively fit the reconstructed SMB values from the GPR best. In the end we chose a roughness length of 0.005 m and 0.6 m/s for the threshold shear velocity.

It is not that clear how much the analysis from the second ice rise, TIR, really adds to the result. The SMB estimates from TIR can only be considered in a relative sense, and the GPR transects from TIR do not really capture the features that are prominently discussed for the FKIR.

TIR mostly adds the information that the features we see are wind driven SMB features not internal deformation. This becomes evident by the fact that we see similar features to the ones we are prominently discussing for the FKIR, namely enhanced SMB on the windward side of the ice rise and erosion on the local ice divide, only in one of the three profiles. The one profile which also shows these features is the one parallel to the dominant wind direction. This is also additional evidence for the idea, that both processes generally occur on ice rises with ridges perpendicular to the predominant wind direction. We added a sentence to clarify this.

C3

L262-266: The absence of the local minima at the ice divide on profile 3 and 4 across the TIR (Fig. 3) provides further evidence that this is a wind driven accumulation feature and not a result of internal deformation. All three profiles are crossing one of the ridges of the ice rise at a similar elevation, however only the one parallel to the dominant wind direction (profile 2) shows the local SMB minimum and maximum near the peak of the profile (Fig. 3). Therefore, it seems that positive curvature of the topography, parallel to the dominant wind direction, is necessary to create this erosion and deposition feature.

The implications of the work should be made more prominent. This work has conclusions that impact the interpretation of ice core records from ice rises, but these implications are only given a few sentences near the end of the discussion, and only just mentioned in the abstract and conclusions.

We agree with the referee that the implications are not prominent enough in the discussion. We changed lines 314-315 in the discussion and added a few sentences to highlight the implications. We also modified figure 7 to highlight the differences in SMB evolution between windward and leeward side. See below.

L314-323: This becomes evident when looking at the temporal SMB evolution on the windward side of the ice rise compared to the SMB evolution at the peak of the ice rise or the leeward side (Fig. 7). While the SMB at the peak of the ice rise and on the leeward side decreases with time, the SMB on the windward side decreases only until 2002, but then increases in the latest time period from 2002 to 2018. This shows a disconnection between the SMB evolution at the peak of the ice rise (where the ice core is recovered) and the windward side of the ice rise, where snowfall is highest. Now since we observe high SMB on the windward side of an ice rise and erosion at the peak of an ice rise on other ice rises too, it is not unlikely that this disconnection between SMB evolution at the peak and on the windward side is also a generic ice rise feature. A consequence of this would be that an SMB record from an ice core at the peak of an ice rise alone would only be sufficient to derive the SMB at the very peak

C4

of the ice rise and the leeward side, but would fail to capture the SMB evolution on the windward side of the same ice rise.

1.3 Small, line-by-line stuff

Figure 1: wind-rose plot is difficult to read at magnifications under 200- suggest reworking that part of the plot if it's important, or eliminating it and replacing with a 'prevalent wind direction' arrow. The black dot at the summit of FKIR is not called out explicitly- presumably it is the ice core location but not clear in the caption- add it in. [Removed the wind rose and added a 'prevalent wind direction' arrow. Also added a note marking the black dot as the location of the ice core.](#)

Line 27: hyphenate "inter-annual"
[Changed to interannual by request of the other referee.](#)

Line 52, set of references: this actually goes back to Black and Budd 1964 (who King et al cite)- might be good to cite them too! Black, H. P., and W. Budd (1964), Accumulation in the region of Wilkes, Wilkes Land, Antarctica, *J. Glaciol.*, 5(37), 3–15.
[Added the citation.](#)

Line 62: Might as well name the ice rises before shortening them to abbreviations (I'm assuming the abbreviations are for "T" ice rise and "FK" ice rise).
[FKIR and TIR were project internal names, named after leading scientist in the Mass2Ant field expedition to the ice rises. However since there are official norwegian names for both ice rises we decided to rename them. Therefore we changed FKIR to Lokeryggen ice rise \(LIR\) and TIR to Hammaryggen ice rise \(HIR\). But we kept the](#)

C5

[name Derwael ice rise \(DIR\) since it was already named like this in earlier publications we are citing. We added a note though mentioning the norwegian name of DIR \(Kilekollen\).](#)

Line 70: delete comma after "rises"
[Done](#)

Lines 76, 77: including make and model of the (probably commercial) GPR would be of interest to GPR folks
[Included the model of the GPR \(GSSI:SIR 3000\)](#)

Figure 2: Fonts too small to read even at 135 zoom. enlarge. Also, the IRHs are challenging to see without magnification of close to 300. Can the contrast be improved, or a different color scheme be employed?
[Increased font size and changed the color of the IRHs. See below](#)

Line 102: Best not to start a sentence with an acronym, even if it's a common one like GPR.
[Changed the sentence to:](#)
L102-103: To study the spatial SMB distribution across the ice rise we used a GPR. The GPR emits radar beams into the ground and records the time it takes for the signal to return, after being reflected on internal layers within the snowpack.

Line 104: Neither of these references establishes that the IRH is an isochron, and you really should include one (Callens in particular uses the vague "generally accepted" language that should be avoided. A good early reference for this is Spikes et. al., 2004 (see figure 2): Spikes, Vandy B., et al. "Variability in accumulation rates from GPR

C6

profiling on the West Antarctic plateau.” Annals of Glaciology 39 (2004): 238-244.
[Added the citation.](#)

Line 106: In section 2.1 you said you used 400 Mhz. Close enough that the science doesn't change, but choose one (preferably the accurate one) and be consistent (unless of course you used both!).
[Corrected to 400 Mhz](#)

Line 106: to clarify the language, suggest changing “the first 50 m of the snow- pack in the vertical direction” to “the shallowest 50 m of the snowpack”
[Done](#)

Line 121: delete “do” from the phrase “in order to do get the density”
[Done](#)

Line 133: Not clear if the initial layer depth mentioned here is part of an iterative procedure or simply part of an initial data analysis step. If the former, describe more fully, if the latter, delete as it's not necessary!
[Removed the unnecessary part of the sentence](#)
L132-133: This layer depth was calculated from the two-way travel time using a radar velocity which increases with density following the mixing formula of Looyenga

Line 135: with such detail in the procedure the omission of an actual equation describing mathematically how SMB is derived seems glaring. Suggest adding here.
[We added an equation and some explanatory sentences.](#)
Finally we combined the density model, one layer depth value for every hundred meter along GPR IRHs and the age dating from the ice core to calculate 220 SMB values for

C7

each time interval, along the profile (Fig. 3 B). Each SMB was calculated by summing up the density between two IRH and dividing by their age difference.

$$SMB = \frac{\sum_{z=d_i}^{d_{i+1}} \rho(z) * V_e}{A_{i+1} - A_i} \quad (1)$$

Where $\rho(z)$ is the density at depth z and $V_e = 1$ is a volume element in our case 1 m times 1 m times 1 m, d_i is the depth of an IRH and A_i is the age of that IRH. The error bars around the SMB profile represent the uncertainty due to the age measurement and the density model.

Lines 138-148: This approach is fine, and using relative magnitudes of SMB is ok, but I don't think it's appropriate in that case to use absolute values on the y axes of Figures 3 D-F. instead, use some scaled value and leave units off. On line 146 you state that the absolute values should be disregarded- thus leaving them on the figure almost invites misuse of the result.
[Removed the units from the y-axis](#)

Line 161: “it consist of” -> “it consists of”
[Done](#)

Line 165: “for a, in our case, single. . . ” awkward. Rewrite.
[Changed it to:](#)
L165-166: SnowPack simulates the changes of snow depth for multiple snow layers, but does not consider any snow micro structure. Here we ran SnowPack with a single snow layer for simplicity reasons.

C8

Line 191: mention Figure 3 first, and then Figure 4.

Done

Figure 3: Again, because the SMB estimates in panels D-F are only relative, leave units off the y-axis.

Done. See below.

Figure 4: make the box in panel A (denoting the location of panel B) more prominent.

Done. See below.

Lines 203-204: start referring to figure 5 here; will be useful.

Done

Figure 5: Needs A and B labels. Also, either repeat legend in panel B (there's plenty of room), or move it outside both panels. Finally, since SMB in each model is what is being directly compared to the GPR SMB, make that curve bold.

Done. See below.

Figure 6: If a figure needs to be removed to save space, this one is a good candidate.

Fig. 6 is indeed the least important figure of all of them, but since we are claiming in the text that the arch amplitudes of the IRHs are increasing linearly with depth, we thought it might still be important to show this.

Lines 238 and 265: I don't see the ice rise DIR actually named anywhere.

Changed the text, DIR is now named in line 195 (Derwael ice rise)

C9

Line 327: "For this ice rise the erosion at the peak. . ." This sentence is unclear- in particular the expression "ice rise wide temporal variations". I understand the idea but it is not well expressed here- suggest split to a few sentences. . .

Changed the sentence.

L339-343: For this ice rise the erosion at the peak of the ice rise locally evens out the higher snowfall values on the windward side of the ice rise. Not only does the erosion at the peak reduce the higher snowfall values due to orographic uplift on the windward side, it also compensates for the higher temporal variability in snowfall on the windward side, as the erosion values are higher when more freshly fallen snow is available.

Interactive comment on The Cryosphere Discuss., <https://doi.org/10.5194/tc-2020-66>, 2020.

C10




Fig. 1.

C11




Fig. 2.

C12




Fig. 3.

C13




Fig. 4.

C14




Fig. 5.

C15




Fig. 6.

C16