

Subglacial lakes and hydrology across the Ellsworth Subglacial Highlands, West Antarctica

Felipe Napoleoni¹, Stewart S. R. Jamieson¹, Neil Ross², Michael J. Bentley¹, Andrés Rivera^{3,4}, Andrew M. Smith⁵, Martin J. Siegert⁶, Guy J. G. Paxman^{1,7}, Guisella Gacitúa⁸, José A. Uribe⁹, Rodrigo Zamora⁹, Alex M. Brisbourne⁵, and David G. Vaughan⁵

¹Department of Geography, Durham University, Durham, UK

²School of Geography, Politics and Sociology, Newcastle University, Newcastle upon Tyne, UK

³Departamento de Geografía, Universidad de Chile, Santiago, Chile

⁴Instituto de Conservación, Biodiversidad y Territorio, Facultad de Ciencias Forestales y Recursos Naturales, Universidad Austral de Chile, Valdivia, Chile.

⁵British Antarctic Survey, Cambridge, UK

⁶Grantham Institute and Department of Earth Science and Engineering, Imperial College London, London, UK

⁷Lamont-Doherty Earth Observatory, Columbia University, Palisades, New York, USA

⁸Instituto de Ciencias Físicas y Matemáticas, Universidad Austral de Chile, Valdivia, Chile

⁹Centro de Estudios Científicos, Arturo Prat 514, Valdivia, Chile

Correspondence: Felipe Napoleoni (felipe.a.napoleoni@durham.ac.uk)

Abstract. Subglacial water plays an important role in ice sheet dynamics and stability. Subglacial lakes are often located at the onset of ice streams and have been hypothesised to enhance ice flow downstream by lubricating the ice-bed interface. The most recent subglacial lake inventory of Antarctica mapped nearly 400 lakes, of which ~14% are found in West Antarctica. Despite the potential importance of subglacial water for ice dynamics, there is a lack of detailed subglacial water characterization in West Antarctica. Using radio-echo sounding data, we analyse the ice-bed interface to detect subglacial lakes. We report 33 previously uncharted subglacial lakes and present a systematic analysis of their physical properties. This represents a ~40% increase in subglacial lakes in West Antarctica. Additionally, a new digital elevation model of basal topography of the Ellsworth Subglacial Highlands was built and used to create a hydropotential model to simulate the subglacial hydrological network. This allows us to characterize basal hydrology, determine subglacial water catchments and assess their connectivity. We show that the simulated subglacial hydrological catchments of Rutford Ice Stream, Pine Island Glacier and Thwaites Glacier do not correspond to their ice surface catchments.

1 Introduction

Subglacial water is important for ice sheet flow, with the potential to control the location of ice stream onset (e.g., Siegert and Bamber, 2000; Vaughan et al., 2007; Winsborrow et al., 2010; Wright and Siegert, 2012) by lubricating the ice base and reducing basal friction (Bell et al., 2011; Pattyn, 2010; Pattyn et al., 2016; Gudlaugsson et al., 2017). Some studies have reported acceleration of ice velocity in response to varying of basal hydrologic conditions (e.g., Stearns et al., 2008). Subglacial water piracy has been invoked to explain the on and off switching of ice streams (e.g., Vaughan et al., 2008; Anandakrishnan

and Alley, 1997; Diez et al., 2018). Other studies have demonstrated the potential variability of subglacial flow routing and that many subglacial lakes form part of a dynamic drainage network (e.g., Siegert, 2000; Fricker et al., 2014; Pattyn et al., 2016).
20 For example, there is evidence of a well-organized and dynamic subglacial hydrological system which formed paleo-channels and basins underneath the present Amundsen Sea Embayment (ASE) (Kirkham et al., 2019). This subglacial hydrological system was hypothesized to be caused by episodic releases of meltwater trapped in upstream subglacial lakes (Kirkham et al., 2019). Additionally, small changes in the ice sheet surface or ice thickness can lead to large-scale changes in basal hydrology; causing water flow to change direction (Wright et al., 2008). Significant glaciological change is known to have taken place
25 in West Antarctica since the Last Glacial Maximum (LGM) (Siegert et al., 2004b, 2019). However, changes to subglacial hydrology associated with post-LGM ice surface elevation changes have not been identified across the Ellsworth Subglacial Highlands (ESH), which are located within the Ellsworth-Whitmore Mountain (EWM) block (Figure 1). Ross et al. (2011) demonstrated that the ice divide and ice flow across ESH have been stable for more than 10 kyr. At present our understanding of the subglacial hydrology in the ESH is relatively limited. Understanding the current hydrological network, and assessing its
30 evolution and sensitivity through time, is important for an improved understanding of Antarctic ice-sheet dynamics. A better understanding of the relationship between subglacial hydrology and ice dynamics is important for studies of ice sheet mass balance and supplies of water to the ocean, where meltwater can affect circulation and nutrient productivity (Ashmore and Bingham, 2014).

The most recent inventory identified ~400 subglacial lakes across Antarctica (Wright and Siegert, 2012, Figure 1), ~14%
35 of which are located beneath the West Antarctic Ice Sheet (WAIS). Some of these subglacial lakes are connected (Wingham et al., 2006; Fricker et al., 2014) and drain and refill dynamically (e.g., Fricker et al., 2007, 2014). Active subglacial lakes have been identified using a range of techniques including satellite measurements of ice surface elevation changes (e.g., Wingham et al., 2006; Smith et al., 2009). Stable deep water subglacial lakes have been identified using airborne radio echo sounding (RES) (e.g., Robin et al., 1970; Popov and Masolov, 2003) and/or ground-based RES (e.g., Rivera et al., 2015).

40 Previous work in the ESH area identified Subglacial Lake Ellsworth (SLE) and Subglacial Lake CECs (SLC) (Figure 1) by interpreting specular basal reflections in RES data as an indicator of deep (>10 m) subglacial water (e.g., Siegert et al., 2004a; Rivera et al., 2015) and although Vaughan et al. (2007) identified other potential subglacial lakes near SLE, none of these candidates were quantitatively confirmed and they were not included in the last subglacial lake inventory (Wright and Siegert, 2012). Subglacial Lake Ellsworth's water depth, geometry and lake floor sediments were characterised using seismic reflection
45 surveys (Woodward et al., 2010; Smith et al., 2018). SLE and SLC are components of a subglacial hydrological network in the upper reaches of multiple West Antarctic ice streams (e.g., Vaughan et al., 2007). However, despite the evidence of subglacial water, and a potential subglacial network connecting multiple subglacial water bodies, many hypotheses remain untested in terms of subglacial hydrological dynamics (e.g., channelized and/or distributed subglacial drainage). Given the fact that this region is located up-ice of the fastest-changing ice streams in the world (i.e. Pine Island Glacier and Thwaites Glacier), and that
50 these are some of the most vulnerable glaciers to ongoing climate change (e.g., Rignot et al., 2014, 2019; Joughin et al., 2014), a more detailed study of the subglacial hydrological system using RES data is justified. Our aim is to produce an inventory of subglacial lakes for the ESH, and to model the modern subglacial hydrology in the ESH draining towards the ASE. We then




Figure 1. The distribution of West Antarctic subglacial lakes between 60°W and 120°W (Wright and Siegert, 2012). EWM: Ellsworth-Whitmore Mountain (in red, polygon from Jordan et al. (2013)); ESH: Ellsworth Subglacial Highlands (in white); Black lines: catchment boundaries produced using data from the SCAR Antarctic Digital Database (<https://www.add.scar.org/>); TG: Thwaites Glacier; PIG: Pine Island Glacier; RIS: Rutford Ice Stream; IIS: Institute Ice Stream. In the background is BedMachine Antarctica v1 elevation model (500 m) (Morlighem et al., 2019), with contour lines every 500 m. Projection: Antarctic Polar Stereographic (EPSG 3031)

assess the connectivity of these new subglacial lakes and the potential flowpaths of basal water from the ice sheet interior to the edge of the grounded ice sheet.

55 2 Methods

During the 2004/2005 austral summer the British Antarctic Survey collected $\sim 35,000$ km of airborne RES data (Vaughan et al., 2006), mostly over the catchment of Pine Island Glacier (PIG) (Figure 5), during the Basin Balance and Synthesis (BBAS) aerogeophysical survey (Vaughan et al., 2006). The survey aircraft was equipped with dual-frequency carrier-phase GPS for navigation, a radar altimeter for surface mapping, magnetometers and a gravimeter for potential field measurements and the
60 Polarimetric radar Airborne Science INstrument (PASIN) ice-sounding radar system (Vaughan et al., 2006, 2007; Corr et al., 2007). The radar system was configured to operate with a transmit power of 4 kW around a central frequency of 150 MHz. A 10-MHz chirp pulse was used to successfully obtain bed-echoes through ice more than 4200 m thick (Vaughan et al., 2006). Here, we use the radar data processed as a combination of coherent and incoherent summation without SAR processing to obtain ice-bed interface information (Vaughan et al., 2006). We analyse the available BBAS data from 2004/2005 to characterize the
65 bed conditions of the northern margin of the ESH. We focus on three main tasks: first, identifying subglacial water at the ice base; second, defining and characterizing the modern subglacial hydrological network; and third, simulating the subglacial flow routing. We identify subglacial lakes by analysing the power of the reflected energy from the ice-bed interface (Gades et al., 2000), i.e. Bed Reflection Power (BRP), using four steps:

1. We use the radar data to identify bright reflections underneath the ice (Section 2.1).
- 70 2. We correct the ice attenuation of the radar power to obtain absolute reflections (Section 2.2).
3. We classify the degree of confidence we have in the identification of each potential water body (Section 2.4) according to the methodology of Carter et al. (2007).
4. Lastly, we analyse the hydraulic potential and ice surface slope of each bright reflection (Section 2.5).

2.1 Identification of subglacial lakes

75 Nearly 7500 km of the BBAS airborne radar (i.e. flight lines: B01, B02, B03, B05, B08, B09, B22, T04) data were analysed to identify high amplitude basal reflections potentially associated with a subglacial lake signature. A preliminary qualitative approach allowed us to identify a variety of bright surfaces at the base of the ice sheet. Following previous methods (e.g., Robin et al., 1970; Oswald and Robin, 1973; Popov and Masolov, 2003; Siegert, 2005; Rivera et al., 2015), these reflections were then quantitatively analysed to determine the BRP in order to classify them as potential subglacial lake candidates, saturated
80 sediments and/or smooth surfaces. We looked for lake candidates that satisfy the following five criteria:

1. Ice surface above subglacial lake candidate must be smooth and planar (Kapitsa et al., 1996; Siegert, 2005).
2. The subglacial lake candidate reflection should have a slope ~ 11 times the magnitude of, and in the opposite direction to, the ice surface slope (Oswald and Robin, 1973).
3. There should be a flat hydraulic potential along the length of the lake (Vaughan et al., 2007).

- 85 4. The lake should have BRP values that are significantly higher than the surrounding surface return (15-20 dB) Siebert (2000).
5. There should be low amplitude strength variation (specularity) of the ice-water interface (i.e. $< 3dB\sigma$ BRP). We followed previous approaches (e.g., Carter et al., 2007) to obtain a specularity proxy using the dispersion of the Bed Reflected Power measured (BRPm). We then used a threshold of 3 dB σ to define a specular surface over all the subglacial lake
90 candidates.

2.2 Bed Reflection Power (BRP)

The returned power measured, $BRPm$, from the ice/bed interface is computed within a defined sampling window using eq.1.

$$BRPm = \frac{1}{n_2 - n_1 + 1} \sum_{n=n_1}^{n_2} s_n^2 \quad (1)$$

where s_n is the amplitude (V) of the signal, and n_1 and n_2 defines the sample numbers in the calculation window.

95

To distinguish the potential lakes from their surroundings, we use a normalized depth-corrected BRP over each candidate lake. We calculate the BRP following previous studies (e.g., Gades et al., 2000; Gacitúa et al., 2015), as the ratio of the locally measured power of the basal echo (BRP_m) to the energy corrected by geometrical spreading and ice attenuation loss (BRP_e):

$$100 \quad BRP = \frac{BRP_m}{BRP_e} \quad (2)$$

In this study, we used as a general expression for the estimated power, BRP_e (e.g., Gacitúa et al., 2015):

$$BRP_e = P_{tx} G L_g T_{ai}^2 L_i^2 R_{ib} \quad (3)$$

where P_{tx} is the transmitted power; G , is the system gain; and L_g , geometrical spreading losses. Additionally, the transmission coefficient at the air/ice interface is defined as T_{ai} ; and the reflection coefficient of the ice/bedrock interface as R_{ib} . Lastly, L_i
105 represents the ice attenuation loss.

We then calculated the geometrical spreading loss (L_g) modifying the original approach by Bentley et al. (1998) as:

$$L_g = \frac{(G_{ant}\lambda)^2}{[8\pi(h_a + \frac{h_i}{\sqrt{\epsilon_i}})]^2} \quad (4)$$

where G_{ant} is the antenna gain (11 dBi), λ is the wavelength of the radar signal at the air medium (1.935 m), and h_a is the
110 height of the antenna above the ice surface (values taken from Corr et al. (2007)). h_i is the ice thickness, derived from BBAS ice thickness picks, and ϵ_i is the relative dielectric permittivity of ice (3.188) (Glen and Paren, 1975).




Figure 2. Example of the ice-attenuation correction for a window of PASIN radar data (BBAS line B01: SL2 in Fig.7). The red best fit line is a log of normalized power representing the BRP dependency to ice thickness.

For each section of the radar profile with a subglacial lake candidate, we quantify the ice attenuation loss (L_i) following previous studies (e.g., Winebrenner et al., 2003; MacGregor et al., 2007; Matsuoka et al., 2012; Schroeder et al., 2016) on a 'section by section' basis. We first use an empirical relationship between the ice thickness (Z_s) and BRP; and we then apply a depth-averaged attenuation rate to correct for the power loss (Figure 2). To calculate the Bed Reflection Power estimated (BRPe) we substitute the L_g calculation and the L_i estimation into eq. 3.

2.3 Specularity calculation

In this work we use the standard deviation (σ) of the BRP as a proxy to infer the surface specularity of each potential water body is (Carter et al., 2007). This is used as a threshold to determine whether the surface of each subglacial lake candidate is smooth or not (Peters et al., 2005). σ was calculated 20 samples either side of a particular point, i.e. ~ 400 m each side, to ensure a representative number of radar observations are included. A low value of σ indicates a smooth surface is present at the base of the ice as would be expected for the surface of a water body (e.g., Peters et al., 2005; Carter et al., 2007).

125 2.4 Subglacial Lake classifications

Having identified the candidate subglacial lakes, we determine the degree of confidence in our identification by ranking them from the most to least likely to be a subglacial lake. To achieve this we initially use the BRP values for each lake candidate following previous work (Carter et al., 2007), comparing these values to known subglacial water bodies such as SLE, SLC and Lake Vostok. This leads to a suite of four physically-based categories to which each potential subglacial lake is assigned, ranging from definite to indistinct as follows:

1. **Definite subglacial lakes.** This category has an absolute reflectivity higher than the surroundings (BRP at least 15db higher) and displays low variation in BRP (high specularity: $< 3 \text{ dB}\sigma$ BRP). Therefore, this category is defined by a high absolute reflection power and a low standard deviation (σ) across the candidate subglacial lake surface.
2. **Dim subglacial lakes.** These candidates have a high relative reflection strength and surface specularity, but lack the absolute reflectivity values of definite lakes in that their BRP is no more than 10db higher than the surroundings.
3. **Fuzzy subglacial lakes.** These candidates show higher relative and absolute reflection coefficients than the surroundings, but are not specular along their surfaces (i.e. $> 3\sigma$ BRP). We note that a challenge with such candidates is that they could also potentially be saturated basal sediments. (Dowdeswell and Siegert, 2003; Peters et al., 2005; Siegert, 2000). In addition, if the water is less than 8 m deep, reflections from the water-lake bottom interface may interfere with the signal from the ice-water interface (Gorman and Siegert, 1999). Furthermore, exceptionally smooth surfaces with no water present could also have similar signal characteristics to these fuzzy subglacial lakes (Carter et al., 2007).
4. **Indistinct subglacial lakes.** This category is composed of lake candidates that are specular but their reflectivity is difficult to distinguish from the surroundings. Although these could still represent subglacial lakes, such characteristics are also common to transient water systems or to exceptionally smooth beds with fine grained sediments surrounded by rougher saturated sediments (Carter et al., 2007).

2.5 Hydropotential (Φ) calculation

To determine the subglacial hydrological characteristics of the EWM region we produced a new bed Digital Elevation Model (DEM). We use existing gridded bed elevation data from: Bedmap2 (Fretwell et al., 2013); ice thickness from DELORES (2007-2009) and CECs (2005/2006) (Siegert et al., 2012); along-track ice thickness measurements from the 2014 CECs RES campaign (Rivera et al., 2015); and new unpublished radar measurements from the CECs 2017 RES field campaign in ESH region (Zamora et al., 2019; Uribe et al., 2019). In January 2014 ~ 1100 km of RES data collected by CECs over an area of $\sim 7000 \text{ km}^2$ were acquired with a line spacing of ~ 8 km. The CECs survey during December of the same year was collected over a nested grid around SLC and along the host subglacial trough further north towards RIS, surveying a total of ~ 1050 km. During December 2017 the CECs radar survey acquired a total of ~ 700 km of RES data along the same trough (Figure 3).

We used a 2 km grid mesh with a continuous curvature tension spline algorithm (Paxman et al., 2017; Wessel et al., 2013) to grid the data. We masked the grid to remove interpolated values more than 5 km from the nearest measured data point, and replaced them with Bedmap2 bed elevation values. As Bedmap2 has a resolution of 1 km, we used the nearest neighbor algorithm to downsample the DEM cell size to 2 km. We also masked the ice shelves from the radar measurements and replaced these values with those from Bedmap2 to obtain offshore bathymetry. The 1 km resolution CryoSat2 ice surface DEM (Slater et al., 2018) was down-sampled to 2 km to match the new bed DEM using the nearest neighbor routine. This enabled us to use the ice surface elevation and subglacial bed to determine the hydrological head, Φ , following Shreve (1972).

At the ice sheet bed, water flows in the direction of steepest descent of the hydraulic potential. Hydropotential (Φ_h) is the sum of the water pressure, P_w (Pa), and water density, ρ_w (1000 and 1020 kg m^{-3} for fresh and sea water, respectively),




Figure 3. Coverage of data sets used in the construction of the Bed Digital Elevation Model. Blue lines show data from Bedmap2 and SLE DELORES surveys; Black lines: catchment boundaries produced using data from the SCAR Antarctic Digital Database v7.1 (<https://www.add.scar.org/>). Red lines show new radar lines from CECs surveys. Projection: Antarctic Polar Stereographic (EPSG 3031).

normalized by the gravitational acceleration, g (9.8 m s^{-2}) and the water system bed elevation, Z (m) (e.g., Shreve, 1972; Cuffey and Paterson, 2010; Livingstone et al., 2013) as follows:

$$\Phi = P_w + \rho_w g Z \quad (5)$$

Equation 5 can be rewritten into an alternative equation in terms of subglacial bed and ice surface elevation (Shreve, 1972).

$$\Phi = \rho_w g Z + k \rho_i g H \quad (6)$$

where ρ_i is the density of ice (917 kg m^{-3}), H is the ice thickness (m) and k is a dimensionless factor, representing the influence of ice overburden pressure on the local subglacial water pressure.

We assumed the water pressure is close to the overburden ice pressure as suggested for the ESH in previous studies (e.g., Vaughan et al., 2007; Rivera et al., 2015), so equation 6 can be rearranged as follows:

$$\Phi = \rho_w g Z + \rho_i g H \quad (7)$$

This approach has been widely used across Antarctica to model the subglacial hydrological drainage (e.g., Livingstone et al., 2013; Carter et al., 2017; Kirkham et al., 2019).

2.6 Subglacial water flow routing

Because subglacial water moves from areas of high to low subglacial water pressure, following the hydropotential (Φ) gradient (Shreve, 1972), we can determine present-day large-scale subglacial flow routing and identify whether the candidate subglacial lakes connect into this subglacial hydrological network. We modelled the subglacial flow routing using the hydropotential (Φ) and followed Schwanghart and Scherler (2014) to calculate the flow routing using the following steps:

1. Lows (sinks) in hydropotential (Φ) were filled to their lowest pour point.
2. The channelized network was then determined using a multiple flow direction (MFD) algorithm (Schwanghart and Scherler, 2014). The subglacial hydraulic drainage basin was then delineated, and we computed the flow accumulation in order to understand the upstream contributing area above the lakes.
3. The stream network was then defined using an up-slope area threshold for channel initiation. We set an arbitrary threshold of 25 connected cells (50 km²) as defining a channel.

3 Results

3.1 New bed elevation model

The new bed DEM of the ESH provides new detail on two major subglacial troughs and shows that they are much deeper than shown in existing DEMs (e.g. Bedmap2 and BedMachine Antarctica v1): Ellsworth Trough (ET) and a parallel trough east of ET, informally referred to here as CECs Trough (CT) (Figure 4). These troughs extend parallel to the Ellsworth Mountains and are extensive linear features that appear to connect the interior of the ESH to the deep basins that lie beneath the WAIS. These are a common feature within the ESH and likely reflects a topography which has evolved under conditions of alpine glacial erosion, fluvial erosion and tectonic processes prior to glacial inception (Jamieson et al., 2014; Sugden et al., 2017; Vaughan et al., 2007; Ross et al., 2014). This area may have been subject to a mix of areal scour and selective linear erosion beneath the WAIS (e.g., Jamieson et al., 2014; Sugden et al., 2017; Paxman et al., 2019). The main difference from Bedmap2 and BedMachine Antarctica v1 are the two parallel subglacial troughs that run from IIS to RIS and PIG, and a subglacial mountain range between the new troughs with a perpendicular transection valley near SLE (Figure 4). In Figure 4, we compare these new features in our DEM against Bedmap2 and BedMachine Antarctica v1.

3.2 Subglacial lakes

Using qualitative visual analysis of the BBAS radar data, we identified 107 bright reflections potentially caused by subglacial lakes within the BBAS dataset (Figure 5). These reflections were further analysed by comparing the characteristics of these features with other Antarctic subglacial lakes (Siegert, 2005; Carter et al., 2007) in order to either confirm or reject each feature as a subglacial lake (Figure 6).




Figure 4. Comparison between three different bed Digital Elevation Models. a) New DEM gridded in this work; b) Bedmap2 DEM; c) BedMachine Antarctica v1. White lines show the catchment boundaries produced using data from the SCAR Antarctic Digital Database (<https://www.add.scar.org/>). ET: Ellsworth Trough; CT: CECs Trough; RIS: Rutford Ice Stream; PIG: Pine Island Glacier; IIS: Institute Ice Stream. Projection: Antarctic Polar Stereographic (EPSG 3031).

The lakes are largely distributed within a series of subglacial valleys that emerge from the ESH into the Bentley Subglacial Trench near the ice divide between PIG, RIS and IIS at the northern edge of the ESH (Figure 7). We observe two clusters of subglacial lakes and one potential chain of subglacial lakes in the ESH region. The first cluster is near SLE, <100 km from the ice divide between IIS, RIS and PIG (Figure 8a). Most of the subglacial lakes in this first cluster are located in ET, or are
 210 connected to the same trough system. However, in some cases the drainage of subglacial water in this cluster may be in two distinct directions, i.e. towards Weddell Sea Embayment (WSE) or ASE (Figure 8c). The second cluster is located in a valley in the upper PIG catchment near the water divide between ASE and WSE (Figure 8b). Hydraulic modelling suggests that this cluster drains towards the WSE (Figure 8d). Furthermore, the subglacial hydrological simulation also shows that part of the drainage flowing beneath PIG is diverted to flow beneath Thwaites Glacier (TG), collecting the water of these ice catchments
 215 and draining to the ASE.

Using the quantitative analysis of the bed reflectivity at 107 locations, and classifying accordingly, we confirm the presence of 33 previously unrecognised subglacial lakes (Supplementary information: Table 1), which is a ~40% increase in the total number of subglacial lakes known to exist beneath the WAIS (Wright and Siegert, 2012). Although a small number of these subglacial lakes were hypothesised or identified by other studies (e.g., Livingstone et al., 2013; Vaughan et al., 2007), none
 220 of these water bodies were included in the most recent subglacial lake inventory of Wright and Siegert (2012). Using the categories described above, we categorise these subglacial lakes into 4 groups with different confidence levels (Figure 7). We identify 7 definite subglacial lakes (very high confidence), 5 dim subglacial lakes (high confidence), 20 fuzzy subglacial lakes (medium confidence) and 1 indistinct subglacial lake (low confidence).




Figure 5. Qualitative analysis of high amplitude reflection (red dots) identified using eight flight lines of the BBAS radar set. The black and red stars show the position of Subglacial Lake CECS (SLC) and Subglacial Lake Ellsworth (SLE), respectively. The red star also marks the location of the radar section shown in Figure 6. Data from the BedMachine Antarctica v1 Elevation model (500 m) (Morlighem et al., 2019) is shown in the background. White lines show the 7500 km of analysed BBAS radar lines. Black lines: catchment boundaries produced using data from the SCAR Antarctic Digital Database (<https://www.add.scar.org/>). ET: Ellsworth Trough; CT: CECs Trough; RIS: Rutford Ice Stream; PIG: Pine Island Glacier; TG: Thwaites Glacier; BST: Bentley Subglacial Trench; IIS: Institute Ice Stream. Projection: Antarctic Polar Stereographic (EPSG 3031).

3.3 Distribution of subglacial lakes

225 The ESH hosts 24 of the new subglacial lakes, 7 others are in the Bentley Subglacial Trench, and 2 lakes are located in the region of relatively high topography between tributaries 3 and 5 of PIG, between Byrd Subglacial Basin and the outlet of PIG (SL21 and SL22 in Figure 7). The majority of these subglacial lakes (30) are located less than 50 km from the ice divide and thirteen are located within 20 km from the ice divide between IIS and RIS (Figure 9a). Sixteen small subglacial lakes (less than




Figure 6. Example of subglacial lake identification in a radargram from the BBAS survey (Flight line B05) and its location in WAIS. a) Shows the location of the flight line (B05, white-line) and the position of the radargram (red star) in the line overlain on the new subglacial DEM (2 km). Ice catchment boundaries produced using data from the SCAR Antarctic Digital Database v7.1 (<https://www.add.scar.org/>) are shown in black. The inset on 'a' shows SLE bed topography contour, and the portion in the profile above SLE between x and x' represents the subglacial lake in 'b'. ET: Ellsworth Trough; CT: CECs Trough; RIS: Rutford Ice Stream; IIS: Institute Ice Stream; PIG: Pine Island Glacier. Projection: Antarctic Polar Stereographic (EPSG 3031). b) Shows the bright reflection (red line) classified as a subglacial lake. The radargram shows both the time (ns) for the returned echo from the ice/bed interface (Y1) and the ice thickness (m) calculated using a radio wave propagation through the ice of 0.168 m ns^{-1} (Y2). c) Bed Reflection Power (dBm). d) Specularity (σBRP). e) Hydropotential considering fresh (blue) and salt (red) water densities. f) Ice surface elevation (blue line) and bed elevation (red line).

3 km length) lie within 30 km of an ice divide, whereas the two largest subglacial lakes (SL6 and SL23) are all located within
 230 25 km of the PIG ice divide and ~ 3 km, respectively. The ice thickness over the small subglacial lakes is variable ranging
 from 1600 m to 4000 m with an average thickness of ~ 2600 m (Figure 9b). Five subglacial lakes are covered by ice thicker
 than 3000 m thick and ten lie underneath ice less than 2000 m thick. Most subglacial lakes associated with thinner ice columns
 are located south of the IIS-PIG ice divide, while those underneath thicker ice columns are distributed at the head of PIG. The
 majority of subglacial lakes with an overlying ice thickness of between 2000 and 3000 m are situated close to the triple ice
 235 divide between IIS-RIS and PIG and along the border between PIG and RIS, where the ice surface slope is near zero (Figure
 9d) and, hence, the subglacial hydraulic gradient is also close to zero. Those subglacial lakes located near the PIG-RIS-IIS
 ice divide lie beneath very slow flowing ice (Figure 9c and Figure 10). However, the ice surface velocity is higher (between
 50 myr^{-1} and 60 myr^{-1}) above those subglacial lakes located at lower elevation (Figure 10). The length of subglacial lakes




Figure 7. Location of subglacial lakes within the Ellsworth Subglacial Highlands. Inset: Location of ESH in Antarctica. The subglacial lakes are represented by different shapes and colours according to the classification of their BRP. The subglacial lakes are classified from a greater to a lesser degree of confidence, as follows: red dots for definite; yellow squares for dim; pink diamonds for fuzzy; and cyan diamonds for indistinct subglacial lakes. The black and red stars represent SLC and SLE, respectively. In the background the new subglacial DEM and the PIG tributaries are numbered according to the scheme given by Stenoien and Bentley (2000). Contours are every 500 m. The red rectangle shows the location of panel a and b (also c and d) in Figure 8. Catchment boundaries produced using data from the SCAR Antarctic Digital Database v7.1 (<https://www.add.scar.org/>) are shown in thick black lines. ASE: Amundsen Sea Embayment; BST: Bentley Subglacial Trench; BSB: Byrd Subglacial Basin. Projection: Antarctic Polar Stereographic (EPSG 3031).

is variable, ranging from a minimum of ~ 0.35 km to a maximum of ~ 8 km, but most (18) are less than 2 km in length (Figure 9e). Some of the subglacial lakes (13) are larger than 2 km and two (SL6 and SL23), near PIG ice divide (Figure 7), are relatively large lakes (~ 8 km). In addition, the new DEM reveals that subglacial lakes are located within two different subglacial topographic contexts: within linear subglacial troughs with steep side walls, and within lowland terrain constrained




Figure 8. Two main subglacial lakes clusters: Ellsworth Trough and western ESH. The location of this cluster is delimited by red squares in Figure 7. a) First cluster of subglacial lakes in Ellsworth Trough. b) second cluster of subglacial lakes identified in upstream Institute area. The new subglacial DEM is shown in the background. In a) and b) white lines are airborne RES survey tracks. Elevation contours lines at 500 m intervals. c) and d) shows the subglacial water hydrological network, for the same areas of a) and b), plotted in different colours depending the flow direction. The red lines show modelled subglacial drainage towards Amundsen Sea Embayment (ASE); the yellow lines, drainage to the Weddell Sea Embayment (WSE). In c) and d) the white lines shows the subglacial water catchment boundaries and the thick black lines the ice sheet boundaries produced using data from the SCAR Antarctic Digital Database (<https://www.add.scar.org/>). Projection: Antarctic Polar Stereographic (EPSG 3031).

by subglacial hills (Figure 4).




Figure 9. Frequency-distribution histograms of subglacial lakes identified in this study. a) Distance from major ice divides. b) Average ice thickness, obtained from BBAS ice thickness measurements.c) Ice surface velocity (Mouginot et al., 2019). d) Ice surface slope, obtained using Cryosat-2 (Slater et al., 2018). e) Minimum length of identified subglacial lake, calculated by measuring the horizontal extent of the lake reflection. Subglacial lakes bed elevation (new DEM).

245 3.4 Subglacial flow routing

The modern configuration of the ice sheet and the improved bed DEM allowed us to calculate a new hydropotential map around ESH. This hydropotential model suggest that the subglacial water catchment of WAIS differ in shape to those previously calculated. The new hydraulic potential model suggests an increase of $\sim 1500 \text{ km}^2$ in the water catchment area of TG (in comparison with Bedmap2), mainly because the contribution of the newly discovered CT (Figure 4). The water flow routing in the hydrological network initiates near the major ice divides in the region (i.e. RIS-IIS-PIG) and flows to the margin of the continent (Figure 7d and e). The hydraulic model assumes a wet bed throughout the ESH and the geometry of the subglacial hydrological network is classically dendritic. It extends almost from the major ice divide (RIS-IIS-PIG) to the edge of the ice sheet, connecting the majority (i.e. 30 subglacial lakes) of the subglacial lakes identified in this work into a wider subglacial hydrological network. We identify 2 main drainage systems: WSE and ASE. There are 15 subglacial lakes draining towards the WSE and 18 draining towards the ASE. In ASE, most of the subglacial lakes are connected into a single drainage network flowing underneath PIG and TG, and partially beneath RIS. Only two subglacial lakes are disconnected from the main ASE

250

255

drainage system: they originate near the border between RIS and PIG over the subglacial highland (SL28) and within a valley sub-parallel to ET (SL31). Only one subglacial lake, located in the upper catchment of IIS (SL10), is disconnected from WSE drainage system. Significantly, we note that the surface flow patterns of ice and the flow patterns of subglacial water are not always co-incident. The most evident difference between hydrological and ice flow catchments are observed underneath the PIG-TG catchments (Figure 8c and 8d). Here, the ice flows from the PIG-IIS ice divide into the PIG catchment, however the subglacial drainage catchment flows first into the head of the PIG catchment and then diverts into the TG catchment. We find that the area of the subglacial hydrological catchments of PIG and TG are $\sim 1.35 \times 10^5 \text{ km}^2$ and $\sim 2.83 \times 10^5 \text{ km}^2$ respectively (if calculated using BedMachine Antarctica v1 they are $\sim 1.25 \times 10^5 \text{ km}^2$ and $\sim 2.88 \times 10^5 \text{ km}^2$, respectively). These are markedly different to the ice surface catchments of PIG and TG, which have areas of $\sim 1.76 \times 10^5$ and $\sim 1.86 \times 10^5 \text{ km}^2$, respectively (Mouginot et al., 2017).

We find that the majority of the known subglacial lakes in the region coincide with channels delineated in our subglacial hydraulic modelling. This gives additional confidence to the assessment of the lakes because it provides an indication that there are upstream areas which might capture enough water to fill a lake, but also that the routing would be appropriately directed to deliver that water into the lake locations. A challenge with the subglacial flow routing is that the bed DEM is based on sparse data due to the reconnaissance nature of the primary airborne RES data-sets across the study area. As a consequence, the interpolation across gaps in the DEM means that the routing is subject to uncertainty associated with overly smooth data in some areas, and potentially noisy data in others. Because we use tension spline interpolation we believe that noise is minimised.

275 4 Discussion

4.1 Subglacial lakes and the production of basal water

We have identified several bright basal reflections, which we interpret as subglacial lakes. Many of these are located at the head of the catchments of PIG and TG and many are very close to the ice divide between IIS, PIG and RIS (Figure 7). Because some of these were classified as 'definite' lakes, we have confidence that they may be up to as much as 100 m deep like SLC and SLE. The subglacial water may be produced by a combination of basal melting directly over subglacial lakes and from the input of water produced elsewhere in the subglacial catchment (e.g., Ross and Siegert, 2020).

The spatial distribution of subglacial lakes in terms of size does not have a evident pattern to its distribution. The RES data shows that most of the subglacial lakes ($\sim 60\%$) are smaller than 2 km and only $\sim 10\%$ are bigger than 5 km. Only $\sim 30\%$ of the subglacial lakes that are between 2 km and 5 km in length and distributed within a region of steep subglacial topographic relief (e.g. the ESH), while the smaller subglacial lakes are found in much lower subglacial terrain. However, the largest new subglacial lakes are also located in a low elevation area, underneath RIS. The distance from the lakes previously inventoried to the ice divide is similar to our new lakes, confirming the tendency of subglacial lakes to be located close to the thick, flat ice and rough basal topography associated with ice divides (Dowdeswell and Siegert, 1999). While Wright and Siegert (2011)




Figure 10. Mean annual ice surface velocity (Mouginot et al., 2019) of Pine Island Glacier, Rutford Ice Stream, Institute Ice Stream and Thwaites Glacier. The red lines show the subglacial water towards Amundsen Sea Embayment (ASE), while the yellow lines indicate the drainage towards Weddell Sea Embayment (WSE). The inset in the upper corner (black square in main panel), shows details in the water catchment boundary between ASE & WSE. ET: Ellsworth Trough; CT: CECs Trough. The white line indicates the boundary of the water catchment (this work) and the blue line indicates the previous boundary (Bedmap2). Ice surface velocity (Mouginot et al., 2019) and the new DEM are shown in the background, with elevation contour every 500 m (thin black lines). Velocities lower than 50 m yr^{-1} are not displayed. BST: Bentley Subglacial Trench. Projection: Antarctic Polar Stereographic (EPSG 3031).

290 mapped subglacial lakes as large as $\sim 10 \text{ km}$, the subglacial lakes identified in this work are all less than 8 km in length. Despite this, the predominance of lakes with lengths less than 3 km is similar to the distribution noted by Wright and Siegert (2011).

Subglacial lakes are likely to form beneath areas suitable for the existence of basal meltwater, for example: areas of thick ice (i.e. $> 2.5 \text{ km}$) where the pressure-melting temperature is enough to maintain liquid water (Dowdeswell and Siegert, 1999; 295 Gorman and Siegert, 1999), in areas of internal ice deformation and sliding that contribute heat to produce melting (Siegert

et al., 1996), or in areas of elevated geothermal heatflux (GHF). Observations of GHF are absent in the ESH, and modelled GHF estimates (e.g., Shapiro and Ritzwoller, 2004; Maule et al., 2005; An et al., 2015; Martos et al., 2017) vary depending on the technique used and may not show localized highs in the GHF. This allows for the possibility of localized elevated GHF, and therefore enhanced production of basal water in the ESH. One potential localised source of elevated basal heat is enhanced radiogenic heat flux from granite intrusions known to exist within the EWM block (Burton-Johnson et al., 2017; Leat et al., 2018). In addition, the high relief basal morphology of ESH (e.g., Vaughan et al., 2007; Ross et al., 2014; Winter et al., 2015) with its narrow and deep subglacial troughs, will enable a localised intensification of GHF via topographic focusing (van der Veen et al., 2007). Continent-wide models of the basal thermal regime (e.g., Pattyn, 2010) suggest that the ESH are warm-based throughout, although given the thin ice located between the deep subglacial troughs, it is more likely to have a patchwork basal thermal regime, with warm-based ice within the deep troughs and cold-based ice on the subglacial interfluves. Beyond the ESH (e.g. PIG, RIS, TG) it is likely that the bed is predominantly warm-based in line with continent-scale models (e.g., Pattyn, 2010). The discrepancy between the Pattyn model and actual basal conditions in the ESH is likely to be due to the coarse resolution of Bedmap (Lythe and Vaughan, 2001) used by Pattyn (2010). Future numerical modelling of basal thermal regime using our new high-resolution DEM and newly identified subglacial lakes would therefore be an important aspect of improving assessments of the basal thermal regime in this region.

4.2 Large-scale subglacial drainage network and lake connectivity

In the ESH, the subglacial water network is mainly controlled by subglacial topography and pre-existing troughs and deep valleys (e.g., ET and CT) (Siegert et al., 2012). Flow routing into Bentley Subglacial Trench, RIS, IIS and TG is determined by the combination of the steep-sided trough walls and the overall form of the ice sheet surface, which partially aligns with ice flow. The new hydropotential map suggests that the subglacial water in the ESH is flowing along the axis of the new deep subglacial troughs (i.e. CT and ET) and is likely connected with the northern edge of the ESH. This implies that more water than previously expected is draining through the TG subglacial water catchment, potentially lubricating its bed.

It is possible that the subglacial lakes identified in the Ellsworth Trough (Figure 8c) are connected in a very well-defined local drainage system (Siegert et al., 2012; Ross and Siegert, 2020). The hydropotential modelling also shows that the subglacial hydrological divide between water flowing into the WSE and ASE is less than 20 km upstream of SLE and most of the seven subglacial lakes in this system are connected or very close to a subglacial water path. This raises the possibility that some episodic events could link these subglacial lakes by draining subglacial water from one hydropotential sink to another, from areas with high hydraulic potential (i.e. modern ice divide between PIG and IIS) to areas with lower hydraulic potential (i.e. the Bentley Subglacial Trench), forming a cascade-type system (e.g., Smith et al., 2017). However, these bridging events may not be large enough to displace the necessary subglacial water along this flow path for detection by remote sensing methods. Another possibility is that these episodes occur on the timescale of tens of years and therefore it may not be noticed from the surface. We also identify a dim subglacial lake (SL16) in the CT, downstream of SLC toward the RIS (Figure 7). Although the

330 current hydraulic modelling does not show a clear connection between this new lake and SLC, it is possible that under different ice sheet configurations both subglacial lakes were connected hydrologically. Alternatively, this lack of connection could be related to uncertainty in topography; or because the model we used to simulate the drainage system does not accurately represent the current hydrological regime.

335 **4.3 The subglacial hydrological catchments of Pine Island and Thwaites Glaciers**

We observe that most of the subglacial water draining towards ASE is routed through the Bentley Subglacial Trench in the upper part of the hydrological catchment and driven through the Byrd Subglacial Basin towards the trunk of TG. The high topography in the centre of PIG catchment (Vaughan et al., 2006) means that the hydrological drainage system does not link to the faster flowing trunk of PIG (Figure 10) and instead the basal hydrological system is captured by TG. This drainage
340 pattern has two main implications. Firstly, the subglacial hydrological catchments of PIG and TG do not correspond to the ice catchments; they do not coincide either in position or size. Secondly, the hydrological system of TG trunk (Schroeder et al., 2013) may be fed by water sourced in the upper glaciological catchment of PIG, within the ESH.

Following from this, any change in the water catchment of the TG, at the head of PIG, could therefore have important glacio-
345 logical consequences for the ice dynamics of TG and the wider ASE. This is particularly critical because the subglacial water drainage area of TG is larger than previously thought and recent investigations (e.g., Smith et al., 2017) have demonstrated the presence of active subglacial lakes, in a cascade system-type, beneath the trunk of TG. Any water accumulation/drainage (e.g., chain of active subglacial lakes) in this area may therefore affect the basal friction of the ice and therefore the ice flow velocity. Conversely, this pattern may have a reduced importance for PIG in terms of magnitude or timing due to the topographic barrier
350 disconnecting the drainage upstream with the lower/marginal section of PIG. If we are to clearly understand the potential role of subglacial water on the ice dynamics of the PIG and TG systems, then more detailed investigations of the subglacial and hydrological conditions are required.

4.4 The spatial relationship between subglacial lakes and ice flow in West Antarctica

355 Almost 90% of the newly identified subglacial lakes in West Antarctica are located in areas of slow ice flow velocity (<20 m/yr) (Mouginot et al., 2019). There are two likely reasons for this: Firstly, the ice surface slope, which is a crucial driver of basal hydraulic conditions, is typically low in these regions (Supplementary information: Table 1), thus enabling ponding to occur even in relatively low magnitude topographic depressions. Secondly, given the ice flow is slow in this area, we can infer that the subglacial network may be an efficiently draining system that does not enable the pressurization of a deforming bed,
360 but instead may allow efficient water transport.

4.5 Limitations of the RES data, the bed DEM and hydropotential modelling

The BBAS radar data were processed as un-focused SAR images and were based on 1-D reflection (Helie et al., 2007). In some areas (e.g., subglacial troughs), this provided a poor constraint on the subglacial bed because of the presence of diffracted hyperbolae caused by an unfocused return of the energy. This means the ice-bed interface is not clear enough to identify in some of the radar lines, and it is especially difficult to detect the bed topography in areas where the strength of the radar return is low. This way of processing could have influenced the BRP calculation, misleading some subglacial lake classification, or underestimating the size of the recognized subglacial lakes due to an uncertainty as to which axis (longitudinal or transversal) had been identified.

370

The BRP calculation was focused in single portions with bright reflectors on each radar profile; and we assume a constant ice attenuation rate by considering the attenuation within these portions as proportional to the ice thickness. Although this attenuation is proportional to the dielectric attenuation, and therefore to the ice temperature and to solute content in the ice (Gudmandsen, 1971; Corr et al., 1993), we did not include any temperature model, which resulted in limited capacity to distinguish differences in reflectance.

375

We categorized every subglacial lake based on their physical characteristics using the classification method proposed by Carter et al. (2007) as a guideline. However, specularity criteria could be modified increasing the threshold to 6 dB sigma as suggested in previous studies (e.g., Gudmandsen, 1971; Peters et al., 2005) allowing some roughness in the ice/bed interface and considering the values for the same criteria in SLE and SLC. In this case, the number of definite subglacial lakes would rise to 17; while the number of fuzzy lakes would decrease to 10 subglacial lakes (Supplementary information: Table 2).

380

Our new subglacial DEM improves our previous knowledge of subglacial bed conditions in the ESH (Bedmap2 (Fretwell et al., 2013), and BedMachine Antarctica v1 (Morlighem et al., 2019)). Despite its relatively coarse resolution, it does account for topographic features which connect the WSE with the ASE that are not yet present in these continental DEMs (Figure 4). We note that our hydropotential model was derived from this new gridded DEM at a resolution of 2 km, and therefore it may not detect some smaller (< 2km) subglacial flow pathways. Although new and/or more detailed subglacial water or drainage systems could be identified in future RES campaigns, the main hydrological outlets would not be substantially different to those which we have identified under the modern ice sheet configuration.

390

Additional targeted surveys across the newly identified lake reflections would better constrain the size and area of the subglacial lakes and therefore improve our estimations of the potential volume of subglacial lakes within this region of WAIS. Moreover, for the parts of the survey closest to the ESH the BBAS survey collected airborne RES using a regular 30-km grid flown at constant altitude (Vaughan et al., 2006), which could have missed some features and/or subglacial lakes hosted in areas between each flight line. Therefore, the number of subglacial lakes discovered in this work could increase with more

395

RES data acquired using a grid optimized to characterize the subglacial interface, considering the geometry of the subglacial topography. Ideally, this grid would have survey lines aligned perpendicular to topographical features (e.g., across subglacial troughs) as opposed to diagonally.

400 5 Conclusions

1. We used RES data from the 2004/5 BBAS survey to locate subglacial lakes within the ESH region of West Antarctica. This analysis allowed us to identify 33 new subglacial lakes to add to the existing inventory (Wright and Siegert, 2012). We then classified these subglacial lakes according to how confident we are in their detection. Using this classification, we identify 7 subglacial lakes with a very high degree of confidence. A further 5 dim subglacial lakes, 20 fuzzy lakes
405 and 1 indistinct subglacial lake were also identified.
 2. We observed that the majority (28) of the subglacial lakes are situated underneath or close (< 40 km) to the modern ice divide between Institute Ice Stream and Rutford Ice Stream, Pine Island Glacier and Thwaites Glacier. Furthermore, we also detected that slow ice flow is associated with these lakes and that there are always low gradient ice surfaces above them.
 - 410 3. We developed a new bed DEM based on recently collected radar survey data that was not previously incorporated in Antarctic topographic models. This allowed us to recognize new topographic features such as the long and linear subglacial trough systems (i.e. CT and ET) which connect to multiple sub-catchments and which therefore may play an important role in the basal hydrology and dynamics of the West Antarctic Ice Sheet.
 4. Using the new DEM and the up to date surface elevation model from Cryosat-2 (Slater et al., 2018) we analysed the
415 subglacial hydraulic network. We identified the potential for connection between the subglacial lakes and the wider subglacial hydrological system, thus providing a mechanism for cascading and active lake drainage. Most importantly, however, we show that the hydrological catchments of RIS, PIG and TG do not correspond precisely with glaciological catchments. Indeed, TG's hydrological catchment appears larger than previously thought, capturing basal water from the upper region of PIG.
- 420 *Author contributions.* The study was conceived by FN, MB, SJ, NR and AS. BBAS RES data was originally collected and provided by DV, NR, MS. CECs RES data was collected and provided by AR, RZ and JAU. RES processing was undertaken by FN, NR, GG and JAU. DEM was created by FN and GP. RES analysis was undertaken by FN, NR, SJ and MB. The manuscript was written by FN with input from all authors.

Competing interests. The authors declare that they have no conflict of interest.

425 *Acknowledgements.* We acknowledge the British Antarctic Survey (BAS) and Centro de Estudios Científicos (CECs) for providing their radar data for analysis. This work is supported by Natural Environment Research Council (NERC) UK grant NE/J008176/1. FN acknowledges support from the Agencia Nacional de Investigación y Desarrollo (ANID) Programa Becas de Doctorado en el Extranjero, Beca Chile for the doctoral scholarship. The subglacial lakes information is available in supplementary information or from corresponding author.

References

- 430 An, M., Wiens, D. A., Zhao, Y., Feng, M., Nyblade, A., Kanao, M., Li, Y., Maggi, A., and L ev eque, J.-J.: Temperature, lithosphere-asthenosphere boundary, and heat flux beneath the Antarctic Plate inferred from seismic velocities, *Journal of Geophysical Research: Solid Earth*, 120, 8720–8742, <https://doi.org/10.1002/2015JB011917>, 2015.
- Anandakrishnan, S. and Alley, R. B.: Stagnation of ice stream C, West Antarctica by water piracy, *Geophysical Research Letters*, 24, 265–268, 1997.
- 435 Ashmore, D. W. and Bingham, R. G.: Antarctic subglacial hydrology: current knowledge and future challenges, *Antarctic Science*, 26, 758–773, <https://doi.org/10.1017/S0954102014000546>, 2014.
- Bell, R. E., Ferraccioli, F., Creyts, T. T., Braaten, D., Corr, H., Das, I., Damaske, D., Frearson, N., Jordan, T., Rose, K., Studinger, M., and Wolovick, M.: Widespread Persistent Thickening of the East Antarctic Ice Sheet by Freezing from the Base, *Science*, 331, 1592–1595, <https://doi.org/10.1126/science.1200109>, <https://science.sciencemag.org/content/331/6024/1592>, 2011.
- 440 Bentley, C. R., Lord, N., and Liu, C.: Radar reflections reveal a wet bed beneath stagnant Ice Stream C and a frozen bed beneath ridge BC, West Antarctica, *Journal of Glaciology*, 44, 149–156, <https://doi.org/10.3189/S0022143000002434>, 1998.
- Burton-Johnson, A., Halpin, J. A., Whittaker, J. M., Graham, F. S., and Watson, S. J.: A new heat flux model for the Antarctic Peninsula incorporating spatially variable upper crustal radiogenic heat production, *Geophysical Research Letters*, 44, 5436–5446, <https://doi.org/10.1002/2017GL073596>, 2017.
- 445 Carter, S. P., Blankenship, D. D., Peters, M. E., Young, D. A., Holt, J. W., and Morse, D. L.: Radar-based subglacial lake classification in Antarctica, *Geochemistry, Geophysics, Geosystems*, 8, <https://doi.org/10.1029/2006GC001408>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2006GC001408>, 2007.
- Carter, S. P., Fricker, H. A., and Siegfried, M. R.: Antarctic subglacial lakes drain through sediment-floored canals: theory and model testing on real and idealized domains, *The Cryosphere*, 11, 381–405, <https://doi.org/10.5194/tc-11-381-2017>, <https://www.the-cryosphere.net/11/381/2017/>, 2017.
- 450 Corr, H., Moore, J. C., and Nicholls, K. W.: Radar absorption due to impurities in Antarctic ice, *Geophysical research letters*, 20, 1071–1074, 1993.
- Corr, H. F., Ferraccioli, F., Frearson, N., Jordan, T., Robinson, C., Armadillo, E., Caneva, G., Bozzo, E., and Tabacco, I.: Airborne radio-echo sounding of the Wilkes Subglacial Basin, the Transantarctic Mountains and the Dome C region, *Terra Antarctica Reports*, 13, 55–63, 2007.
- 455 Cuffey, K. M. and Paterson, W. S. B.: *The physics of glaciers*, Academic Press, 2010.
- Diez, A., Matsuoka, K., Ferraccioli, F., Jordan, T. A., Corr, H. F., Kohler, J., Olesen, A. V., and Forsberg, R.: Basal Settings Control Fast Ice Flow in the Recovery/Slessor/Bailey Region, East Antarctica, *Geophysical Research Letters*, 45, 2706–2715, <https://doi.org/10.1002/2017GL076601>, 2018.
- Dowdeswell, J. A. and Siegert, M. J.: The dimensions and topographic setting of Antarctic subglacial lakes and implications for large-scale water storage beneath continental ice sheets, *Geological Society of America Bulletin*, 111, 254–263, 1999.
- 460 Dowdeswell, J. A. and Siegert, M. J.: The physiography of modern Antarctic subglacial lakes, *Global and Planetary Change*, 35, 221 – 236, [https://doi.org/https://doi.org/10.1016/S0921-8181\(02\)00128-5](https://doi.org/https://doi.org/10.1016/S0921-8181(02)00128-5), <http://www.sciencedirect.com/science/article/pii/S0921818102001285>, subglacial Lakes: A Planetary Perspective, 2003.
- Fretwell, P., Pritchard, H. D., Vaughan, D. G., Bamber, J. L., Barrand, N. E., Bell, R., Bianchi, C., Bingham, R. G., Blankenship, D. D., 465 Casassa, G., Catania, G., Callens, D., Conway, H., Cook, A. J., Corr, H. F. J., Damaske, D., Damm, V., Ferraccioli, F., Forsberg, R., Fujita,

- S., Gim, Y., Gogineni, P., Griggs, J. A., Hindmarsh, R. C. A., Holmlund, P., Holt, J. W., Jacobel, R. W., Jenkins, A., Jokat, W., Jordan, T., King, E. C., Kohler, J., Krabill, W., Riger-Kusk, M., Langley, K. A., Leitchenkov, G., Leuschen, C., Luyendyk, B. P., Matsuoka, K., Mouginot, J., Nitsche, F. O., Nogi, Y., Nost, O. A., Popov, S. V., Rignot, E., Rippin, D. M., Rivera, A., Roberts, J., Ross, N., Siegert, M. J., Smith, A. M., Steinhage, D., Studinger, M., Sun, B., Tinto, B. K., Welch, B. C., Wilson, D., Young, D. A., Xiangbin, C., and Zirizzotti, A.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica, *The Cryosphere*, 7, 375–393, <https://doi.org/10.5194/tc-7-375-2013>, 2013.
- 470 Fricker, H. A., Scambos, T., Bindschadler, R., and Padman, L.: An Active Subglacial Water System in West Antarctica Mapped from Space, *Science*, 315, 1544–1548, <https://doi.org/10.1126/science.1136897>, 2007.
- Fricker, H. A., Carter, S. P., Bell, R. E., and Scambos, T.: Active lakes of Recovery Ice Stream, East Antarctica: a bedrock-controlled subglacial hydrological system, *Journal of Glaciology*, 60, 1015–1030, <https://doi.org/10.3189/2014JG14J063>, 2014.
- 475 Gacitúa, G., Uribe, J. A., Wilson, R., Loriaux, T., Hernández, J., and Rivera, A.: 50 MHz helicopter-borne radar data for determination of glacier thermal regime in the central Chilean Andes, *Annals of Glaciology*, 56, 193–201, <https://doi.org/10.3189/2015AoG70A953>, 2015.
- Gades, A. M., Raymond, C. F., Conway, H., and Jagobel, R. W.: Bed properties of Siple Dome and adjacent ice streams, West Antarctica, inferred from radio-echo sounding measurements, *Journal of Glaciology*, 46, 88–94, <https://doi.org/10.3189/172756500781833467>, 2000.
- 480 Glen, J. W. and Paren, J. G.: The Electrical Properties of Snow and Ice, *Journal of Glaciology*, 15, 15–38, <https://doi.org/10.3189/S0022143000034249>, 1975.
- Gorman, M. R. and Siegert, M. J.: Penetration of Antarctic subglacial lakes by VHF electromagnetic pulses: Information on the depth and electrical conductivity of basal water bodies, *Journal of Geophysical Research: Solid Earth*, 104, 29 311–29 320, <https://doi.org/10.1029/1999JB900271>, 1999.
- 485 Gudlaugsson, E., Humbert, A., Andreassen, K., Clason, C. C., Kleiner, T., and Beyer, S.: Eurasian ice-sheet dynamics and sensitivity to subglacial hydrology, *Journal of Glaciology*, 63, 556–564, 2017.
- Gudmandsen, P.: Electromagnetic Probing of Ice, in: *Electromagnetic Probing in Geophysics*, edited by Wait, J. R., vol. 79-147274, p. 321, 1971.
- Heliere, F., Lin, C., Corr, H., and Vaughan, D.: Radio Echo Sounding of Pine Island Glacier, West Antarctica: Aperture Synthesis Processing and Analysis of Feasibility From Space, *IEEE Transactions on Geoscience and Remote Sensing*, 45, 2573–2582, <https://doi.org/10.1109/TGRS.2007.897433>, 2007.
- 490 Jamieson, S. S., Stokes, C. R., Ross, N., Rippin, D. M., Bingham, R. G., Wilson, D. S., Margold, M., and Bentley, M. J.: The glacial geomorphology of the Antarctic ice sheet bed, *Antarctic Science*, 26, 724–741, <https://doi.org/10.1017/S0954102014000212>, 2014.
- Jordan, T. A., Ferraccioli, F., Ross, N., Corr, H. F., Leat, P. T., Bingham, R. G., Rippin, D. M., le Brocq, A., and Siegert, M. J.: Inland extent of the Weddell Sea Rift imaged by new aerogeophysical data, *Tectonophysics*, 585, 137–160, 2013.
- 495 Joughin, I., Smith, B. E., and Medley, B.: Marine Ice Sheet Collapse Potentially Under Way for the Thwaites Glacier Basin, West Antarctica, *Science*, 344, 735–738, <https://doi.org/10.1126/science.1249055>, <https://science.sciencemag.org/content/344/6185/735>, 2014.
- Kapitsa, A. P., Ridley, J. K., de Q. Robin, G., Siegert, M. J., and Zotikov, I. A.: A large deep freshwater lake beneath the ice of central East Antarctica, *Nature*, 381, 684–686, <https://doi.org/10.1038/381684a0>, 1996.
- 500 Kirkham, J. D., Hogan, K. A., Larter, R. D., Arnold, N. S., Nitsche, F. O., Gолledge, N. R., and Dowdeswell, J. A.: Past water flow beneath Pine Island and Thwaites glaciers, West Antarctica, *The Cryosphere*, 13, 1959–1981, <https://doi.org/10.5194/tc-13-1959-2019>, <https://www.the-cryosphere.net/13/1959/2019/>, 2019.

- Leat, P. T., Jordan, T. A., Flowerdew, M. J., Riley, T. R., Ferraccioli, F., and Whitehouse, M. J.: Jurassic high heat production granites associated with the Weddell Sea rift system, Antarctica, *Tectonophysics*, 722, 249–264, 2018.
- 505 Livingstone, S. J., Clark, C. D., Woodward, J., and Kingslake, J.: Potential subglacial lake locations and meltwater drainage pathways beneath the Antarctic and Greenland ice sheets, *The Cryosphere*, 7, 1721–1740, <https://doi.org/10.5194/tc-7-1721-2013>, 2013.
- Lythe, M. B. and Vaughan, D. G.: BEDMAP: A new ice thickness and subglacial topographic model of Antarctica, *Journal of Geophysical Research: Solid Earth*, 106, 11 335–11 351, 2001.
- MacGregor, J. A., Winebrenner, D. P., Conway, H., Matsuoka, K., Mayewski, P. A., and Clow, G. D.: Modeling englacial radar attenuation at Siple Dome, West Antarctica, using ice chemistry and temperature data, *Journal of Geophysical Research: Earth Surface*, 112, <https://doi.org/10.1029/2006JF000717>, 2007.
- 510 Martos, Y. M., Catalán, M., Jordan, T. A., Golynsky, A., Golynsky, D., Eagles, G., and Vaughan, D. G.: Heat Flux Distribution of Antarctica Unveiled, *Geophysical Research Letters*, 44, 11,417–11,426, <https://doi.org/10.1002/2017GL075609>, 2017.
- Matsuoka, K., MacGregor, J. A., and Pattyn, F.: Predicting radar attenuation within the Antarctic ice sheet, *Earth and Planetary Science Letters*, 359, 173–183, 2012.
- 515 Maule, C. F., Purucker, M. E., Olsen, N., and Mosegaard, K.: Heat flux anomalies in Antarctica revealed by satellite magnetic data, *Science*, 309, 464–467, 2005.
- Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., Fretwell, P., et al.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, pp. 1–6, 2019.
- 520 Mouginot, J., Scheuchl, B., and Rignot, E.: MEaSURES Antarctic boundaries for IPY 2007–2009 from satellite radar, version 2, Boulder, CO: NASA National Snow and Ice Data Center Distributed Active Archive Center. <https://doi.org/10.5067/AXE4121732AD>, 2017.
- Mouginot, J., Rignot, E., and Scheuchl, B.: Continent-Wide, Interferometric SAR Phase, Mapping of Antarctic Ice Velocity, *Geophysical Research Letters*, 0, <https://doi.org/10.1029/2019GL083826>, 2019.
- Oswald, G. and Robin, G.: Lakes Beneath the Antarctic Ice Sheet, *Nature*, 245, 251–254, <https://doi.org/10.1038/245251a0>, 1973.
- 525 Pattyn, F.: Antarctic subglacial conditions inferred from a hybrid ice sheet/ice stream model, *Earth and Planetary Science Letters*, 295, 451–461, <https://doi.org/https://doi.org/10.1016/j.epsl.2010.04.025>, 2010.
- Pattyn, F., Carter, S. P., and Thoma, M.: Advances in modelling subglacial lakes and their interaction with the Antarctic ice sheet, *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 374, 20140296, <https://doi.org/10.1098/rsta.2014.0296>, 2016.
- 530 Paxman, G., Jamieson, S., Ferraccioli, F., Bentley, M., Forsberg, R., Ross, N., Watts, A., F.J. Corr, H., and Jordan, T.: Uplift and tilting of the Shackleton Range in East Antarctica driven by glacial erosion and normal faulting: Flexural Uplift of the Shackleton Range, *Journal of Geophysical Research: Solid Earth*, 122, <https://doi.org/10.1002/2016JB013841>, 2017.
- Paxman, G. J., Jamieson, S. S., Hochmuth, K., Gohl, K., Bentley, M. J., Leitchenkov, G., and Ferraccioli, F.: Reconstructions of Antarctic topography since the Eocene–Oligocene boundary, *Palaeogeography, Palaeoclimatology, Palaeoecology*, 535, 109–136, <https://doi.org/https://doi.org/10.1016/j.palaeo.2019.109346>, 2019.
- 535 Peters, M. E., Blankenship, D. D., and Morse, D. L.: Analysis techniques for coherent airborne radar sounding: Application to West Antarctic ice streams, *Journal of Geophysical Research: Solid Earth*, 110, <https://doi.org/10.1029/2004JB003222>, 2005.
- Popov, S. and Masolov, V.: Novye dannye o podlednikovih ozerah tsentral'noy chasty Vostochnoy Antarktity [New data on subglacial lakes in central part of Eastern Antarctica], *Materialy Glatsiologicheskikh Issledovaniy*, 95, 161–167, 2003.

- 540 Rignot, E., Mouginot, J., Morlighem, M., Seroussi, H., and Scheuchl, B.: Widespread, rapid grounding line retreat of Pine Island, Thwaites, Smith, and Kohler glaciers, West Antarctica, from 1992 to 2011, *Geophysical Research Letters*, 41, 3502–3509, <https://doi.org/10.1002/2014GL060140>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2014GL060140>, 2014.
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proceedings of the National Academy of Sciences*, 116, 1095–1103, <https://doi.org/10.1073/pnas.1812883116>,
545 <https://www.pnas.org/content/116/4/1095>, 2019.
- Rivera, A., Uribe, J., Zamora, R., and Oberreuter, J.: Subglacial Lake CECs: Discovery and in situ survey of a privileged research site in West Antarctica, *Geophysical Research Letters*, 42, 3944–3953, <https://doi.org/10.1002/2015GL063390>, 2015.
- Robin, G. d. Q., Swithinbank, C., Smith, B., et al.: Radio echo exploration of the Antarctic ice sheet, *International Association of Scientific Hydrology Publication*, 86, 97–115, 1970.
- 550 Ross, N. and Siegert, M.: Basal melting over Subglacial Lake Ellsworth and its catchment: insights from englacial layering, *Annals of Glaciology*, p. 1–8, <https://doi.org/10.1017/aog.2020.50>, 2020.
- Ross, N., Siegert, M., Woodward, J., Smith, A., Corr, H., Bentley, M., Hindmarsh, R., King, E., and Rivera, A.: Holocene stability of the Amundsen-Weddell ice divide, West Antarctica, *Geology*, 39, 935–938, <https://doi.org/10.1130/G31920.1>, 2011.
- Ross, N., Jordan, T. A., Bingham, R. G., Corr, H. F., Ferraccioli, F., Le Brocq, A., Rippin, D. M., Wright, A. P., and Siegert, M. J.: The
555 Ellsworth subglacial highlands: inception and retreat of the West Antarctic Ice Sheet, *Bulletin*, 126, 3–15, 2014.
- Schroeder, D. M., Blankenship, D. D., and Young, D. A.: Evidence for a water system transition beneath Thwaites Glacier, West Antarctica, *Proceedings of the National Academy of Sciences*, 110, 12 225–12 228, 2013.
- Schroeder, D. M., Seroussi, H., Chu, W., and Young, D. A.: Adaptively constraining radar attenuation and temperature across the Thwaites Glacier catchment using bed echoes, *Journal of Glaciology*, 62, 1075–1082, 2016.
- 560 Schwanghart, W. and Scherler, D.: Short Communication: TopoToolbox 2 “MATLAB-based software for topographic analysis and modeling in Earth surface sciences, *Earth Surface Dynamics*, 2, 1–7, <https://doi.org/10.5194/esurf-2-1-2014>, <https://www.earth-surf-dynam.net/2/1/2014/>, 2014.
- Shapiro, N. M. and Ritzwoller, M. H.: Inferring surface heat flux distributions guided by a global seismic model: particular application to Antarctica, *Earth and Planetary Science Letters*, 223, 213–224, 2004.
- 565 Shreve, R. L.: Movement of Water in Glaciers, *Journal of Glaciology*, 11, 205–214, <https://doi.org/10.3189/S002214300002219X>, 1972.
- Siegert, M., Dowdeswell, J., Gorman, M., and McIntyre, N.: An inventory of Antarctic sub-glacial lakes, *Antarctic Science*, 8, 281 – 286, <https://doi.org/10.1017/S0954102096000405>, 1996.
- Siegert, M. J.: Radar evidence of water-saturated sediments beneath the East Antarctic Ice Sheet, *Geological Society, London, Special Publications*, 176, 217–229, <https://doi.org/10.1144/GSL.SP.2000.176.01.17>, <https://sp.lyellcollection.org/content/176/1/217>, 2000.
- 570 Siegert, M. J.: Lakes Beneath the Ice Sheet: The Occurrence, Analysis, and Future Exploration of Lake Vostok and Other Antarctic Subglacial Lakes, *Annual Review of Earth and Planetary Sciences*, 33, 215–245, <https://doi.org/10.1146/annurev.earth.33.092203.122725>, 2005.
- Siegert, M. J. and Bamber, J. L.: Subglacial water at the heads of Antarctic ice-stream tributaries, *Journal of Glaciology*, 46, 702–703, 2000.
- Siegert, M. J., Hindmarsh, R., Corr, H., Smith, A., Woodward, J., King, E. C., Payne, A. J., and Joughin, I.: Subglacial Lake Ellsworth: A candidate for in situ exploration in West Antarctica, *Geophysical Research Letters*, 31, <https://doi.org/10.1029/2004GL021477>, 2004a.
- 575 Siegert, M. J., Welch, B., Morse, D., Vieli, A., Blankenship, D. D., Joughin, I., King, E. C., Vieli, G. J.-M. C. L., Payne, A. J., and Jacobel, R.: Ice Flow Direction Change in Interior West Antarctica, *Science*, 305, 1948–1951, <https://doi.org/10.1126/science.1101072>, 2004b.

- Siegert, M. J., Clarke, R. J., Mowlem, M., Ross, N., Hill, C. S., Tait, A., Hodgson, D., Parnell, J., Tranter, M., Pearce, D., Bentley, M. J., Cockell, C., Tsaloglou, M.-N., Smith, A., Woodward, J., Brito, M. P., and Waugh, E.: Clean access, measurement, and sampling of Ellsworth Subglacial Lake: A method for exploring deep Antarctic subglacial lake environments, *Reviews of Geophysics*, 50, 580 <https://doi.org/10.1029/2011RG000361>, 2012.
- Siegert, M. J., Kingslake, J., Ross, N., Whitehouse, P. L., Woodward, J., Jamieson, S. S. R., Bentley, M. J., Winter, K., Wearing, M., Hein, A. S., Jeofry, H., and Sugden, D. E.: Major Ice Sheet Change in the Weddell Sea Sector of West Antarctica Over the Last 5,000 Years, *Reviews of Geophysics*, n/a, <https://doi.org/10.1029/2019RG000651>, 2019.
- Slater, T., Shepherd, A., McMillan, M., Muir, A., Gilbert, L., Hogg, A. E., Konrad, H., and Parrinello, T.: A new digital elevation model of Antarctica derived from CryoSat-2 altimetry, *The Cryosphere*, 12, 1551–1562, <https://doi.org/10.5194/tc-12-1551-2018>, <https://www.the-cryosphere.net/12/1551/2018/>, 2018.
- Smith, A. M., Woodward, J., Ross, N., Bentley, M. J., Hodgson, D. A., Siegert, M. J., and King, E. C.: Evidence for the long-term sedimentary environment in an Antarctic subglacial lake, *Earth and Planetary Science Letters*, 504, 139 – 151, <https://doi.org/10.1016/j.epsl.2018.10.011>, 2018.
- 590 Smith, B. E., Fricker, H. A., Joughin, I. R., and Tulaczyk, S.: An inventory of active subglacial lakes in Antarctica detected by ICESat (2003–2008), *Journal of Glaciology*, 55, 573–595, <https://doi.org/10.3189/002214309789470879>, 2009.
- Smith, B. E., Gourmelen, N., Huth, A., and Joughin, I.: Connected subglacial lake drainage beneath Thwaites Glacier, West Antarctica, *The Cryosphere*, 11, 451–467, <https://doi.org/10.5194/tc-11-451-2017>, 2017.
- Stearns, L. A., Smith, B. E., and Hamilton, G. S.: Increased flow speed on a large East Antarctic outlet glacier caused by subglacial floods, 595 *Nature Geoscience*, 1, 827, 2008.
- Stenoien, M. D. and Bentley, C. R.: Pine Island Glacier, Antarctica: A study of the catchment using interferometric synthetic aperture radar measurements and radar altimetry, *Journal of Geophysical Research: Solid Earth*, 105, 21 761–21 779, 2000.
- Sugden, D. E., Hein, A. S., Woodward, J., Marrero, S. M., Ángel Rodés, Dunning, S. A., Stuart, F. M., Freeman, S. P., Winter, K., and Westoby, M. J.: The million-year evolution of the glacial trimline in the southernmost Ellsworth Mountains, Antarctica, *Earth and Planetary Science Letters*, 469, 42 – 52, <https://doi.org/https://doi.org/10.1016/j.epsl.2017.04.006>, 2017.
- 600 Uribe, J., Zamora, R., Pulgar, S., Oberreuter, J., and Rivera, A.: Overview of the low-frequency ice penetrating radar system survey conducted to Subglacial Lake CECs, West Antarctica, https://www.igsoc.org/symposia/2019/stanford/proceedings/procsfiles/procabstracts_75.html#A2948, 2019.
- van der Veen, C. J., Leftwich, T., von Frese, R., Csatho, B. M., and Li, J.: Subglacial topography and geothermal heat flux: Potential interactions with drainage of the Greenland ice sheet, *Geophysical Research Letters*, 34, <https://doi.org/10.1029/2007GL030046>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2007GL030046>, 2007.
- 605 Vaughan, D. G., Corr, H. F. J., Ferraccioli, F., Frearson, N., O’Hare, A., Mach, D., Holt, J. W., Blankenship, D. D., Morse, D. L., and Young, D. A.: New boundary conditions for the West Antarctic ice sheet: Subglacial topography beneath Pine Island Glacier, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2005GL025588>, 2006.
- 610 Vaughan, D. G., Rivera, A., Woodward, J., Corr, H. F. J., Wendt, J., and Zamora, R.: Topographic and hydrological controls on Subglacial Lake Ellsworth, West Antarctica, *Geophysical Research Letters*, 34, L18 501, <https://doi.org/10.1029/2007GL030769>, 2007.
- Vaughan, D. G., Corr, H. F., Smith, A. M., Pritchard, H. D., and Shepherd, A.: Flow-switching and water piracy between Rutford ice stream and Carlson inlet, West Antarctica, *Journal of Glaciology*, 54, 41–48, 2008.

- Wessel, P., Smith, W. H. F., Scharroo, R., Luis, J., and Wobbe, F.: Generic Mapping Tools: Improved Version Released, *Eos, Transactions American Geophysical Union*, 94, 409–410, <https://doi.org/10.1002/2013EO450001>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2013EO450001>, 2013.
- Winebrenner, D. P., Smith, B. E., Catania, G. A., Conway, H. B., and Raymond, C. F.: Radio-frequency attenuation beneath Siple Dome, West Antarctica, from wide-angle and profiling radar observations, *Annals of Glaciology*, 37, 226–232, 2003.
- Wingham, D. J., Siegert, M. J., Shepherd, A., and Muir, A. S.: Rapid discharge connects Antarctic subglacial lakes, *Nature*, 440, <https://doi.org/10.1038/nature04660>, 2006.
- Winsborrow, M. C., Clark, C. D., and Stokes, C. R.: What controls the location of ice streams?, *Earth-Science Reviews*, 103, 45–59, 2010.
- Winter, K., Woodward, J., Ross, N., Dunning, S. A., Bingham, R. G., Corr, H. F., and Siegert, M. J.: Airborne radar evidence for tributary flow switching in Institute Ice Stream, West Antarctica: Implications for ice sheet configuration and dynamics, *Journal of Geophysical Research: Earth Surface*, 120, 1611–1625, 2015.
- Woodward, J., Smith, A. M., Ross, N., Thoma, M., Corr, H. F. J., King, E. C., King, M. A., Grosfeld, K., Tranter, M., and Siegert, M. J.: Location for direct access to subglacial Lake Ellsworth: An assessment of geophysical data and modeling, *Geophysical Research Letters*, 37, <https://doi.org/10.1029/2010GL042884>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2010GL042884>, 2010.
- Wright, A. and Siegert, M.: A fourth inventory of Antarctic subglacial lakes, *Antarctic Science*, 24, 659–664, <https://doi.org/10.1017/S095410201200048X>, 2012.
- Wright, A. and Siegert, M. J.: The Identification and Physiographical Setting of Antarctic Subglacial Lakes: An Update Based on Recent Discoveries, pp. 9–26, American Geophysical Union (AGU), <https://doi.org/10.1002/9781118670354.ch2>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/9781118670354.ch2>, 2011.
- Wright, A. P., Siegert, M. J., Le Brocq, A. M., and Gore, D. B.: High sensitivity of subglacial hydrological pathways in Antarctica to small ice-sheet changes, *Geophysical Research Letters*, 35, <https://doi.org/10.1029/2008GL034937>, <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1029/2008GL034937>, 2008.
- Zamora, R., Uribe, J., Pulgar, S., Oberreuter, J., and Rivera, A.: Ground penetrating radar system for measuring deep ice in Antarctica using software-defined radio approach, https://www.igsoc.org/symposia/2019/stanford/proceedings/procsfiles/procabstracts_75.html#A2968, 2019.