

Response to Editor Comments

Journal: The Cryosphere

Title: **Sensitivity of ice flow to uncertainty in flow law parameters in an idealized one-dimensional geometry**

Author(s): Maria Zeitz, Anders Levermann, Ricarda Winkelmann

MS No.: tc-2020-79

MS Type: Research Article

We would like to thank the editor Alexander Robinson for his helpful comments! We provide detailed answers to all comments below. In particular, we added a new figure, which addresses the main question brought up by the editor, to the supplemental material. We are just as happy to move this figure to the main document if you see fit.

The editor's comments are given in black and the authors' in blue. The changes made to the manuscript can be found at the end of this document (created with latexdiff).

Editor Decision: Publish subject to minor revisions (review by editor) (16 Aug 2020) by Alexander Robinson

Comments to the Author:

The manuscript is in very good condition, and near-ready for publication, except for potentially one critical point that I believe has been overlooked until now.

I apologize that I didn't notice it until now, but I am somewhat concerned by the experimental setup described in Section 3.2, in which the initial volume is kept constant by tuning accumulation. This procedure may introduce an unwanted change in the temperature of the ice sheet, since the accumulation rate directly controls the vertical velocity profile, which enters the thermodynamic equation. By e.g., increasing accumulation, one can generally expect a colder ice sheet and a lower rate factor. Accumulation rate differences of between 50% and 300% seem large enough to have a strong impact on internal temperature. Since flow differences due to Q and n are precisely what is studied here, I am worried that the interpretation of the results is contaminated by the systematic changes in accumulation.

The best solution at this stage would be to be able to confirm that this is not the case – i.e., that despite the influence of accumulation on internal temperature, its impact is much smaller than that of the rate factor parameters themselves. Maybe this could be done with a fixed ice thickness profile: then you could diagnose changes in the rate factor and/or ice velocity as a function of the rate-factor parameters and accumulation and see the magnitude of their impacts? If changes in accumulation do play a large role, perhaps you could consider simply allowing the initial volumes to differ – since you report volume losses in percent, the difference in absolute volume should not have a large impact given the idealized configuration of the experiment.

Thank you very much for bringing this up. In order to keep results most comparable we chose to fix the equilibrium volume and the geometry of the ice sheet though adapted accumulation rates. Indeed, the temperature distribution in the ice changes with variations in the accumulation rate, as we show in

Supplementary Figure S4 (a) of the revised supplementary material. Temperatures decrease with increased accumulation rates, which in this setup are associated with high activation energies, and vice versa. The temperature change due to adapted accumulation rates is most prominent at the top of the ice sheet.

The softness (or rate factor) of the ice depends on pressure adjusted temperature and flow parameters. In this setup high activation energies are associated with high softness and increased accumulation rates and low activation energies with low softness and low accumulation rates. The temperature decrease from the increased accumulation rates thus counteracts the increase in softness due to the change in parameters.

The equilibrium simulations with adapted accumulation rates show distinct variations in softness, in particular close to the base of the ice sheet (see Figure S4 b). We disentangle the influences of the change in flow parameters (i.e. activation energies) and the change in the temperature distribution: Replacing the flow parameters by the standard values leads to very similar softness profiles, despite changes in the ice-sheet temperature (see S4 c). On the other hand, fixing the ice temperatures and varying only the flow parameters reproduces the softness profiles which are observed in equilibrium (see S4 d).

We included a short discussion with a reference to the supplementary figure in the main text.

In either case, I think it is important to address this point before moving forward with publication.

Please find additional minor points below:

L42: I suggest separating discussion and references about the parameter n and the parameter Q . Therefore, I would move the sentences of L52-56 concerning the parameter n to just after “which can be between 2 and 4” on L42, to keep the discussion of n together. Then proceed with discussion of Q . I.e., “..., which can be between 2 and 4 [REFS]. [Sentences from L52-56]. The activation energies Q ... can also vary by a factor of two [REFS].

done

L43: 50+ citations to support the statement that n shows a large spread and that the activation energies “can vary by a factor of two” seems a bit excessive. I support the notion of such an extensive review of older and perhaps now forgotten literature, especially if nothing new has been done since beyond it. However, for that I would expect some specific information that came from each paper with separated citations (essentially, a more detailed review paragraph). If you are only interested in supporting the above statement, I would suggest reducing the number of references to the most relevant and ground-breaking ones.

done

L52: Merge “(Zeit et al. submitted)” into the other citations here, or specify what is different about the reference, aside from only having been submitted. As it is now, it’s not clear why it stands apart.

L75: It may depend, among others, on \Rightarrow It may depend on water content ..., among other things.

done

L79: temperature difference \Rightarrow temperature relative

done

L95: independently \Rightarrow independent

done

L115: the typical standard value => a typical reference value

done

L116: the standard value => a typical reference value

done

L118: The standard values serve as a reference point and correspond => The reference values above correspond

done

L130: warm or cold ice respectively => cold or warm ice, respectively,

done

L134: in the same order => on the same order

done

L146: I would expect to see the basal boundary condition here as well, ie, geothermal heat flux.

done

L196: the volume calculated => the ice profile

done

L210: , which matches => that matches

done

Fig. 6: Remove grey lines connecting the zoom of panel (b) to that of panel (a). They do not help comprehension, and the caption and axes are clear enough.

done

Fig. 8: Panel (b) should include a y-axis label as well (despite having the same units, I assume), to avoid ambiguity.

done

L259-261: I'm not able to follow this logic – it sounds like something contradictory is stated. If A_0 reduces with a lower n , or rather if A_0 increases with a higher n , then the rate factor should increase with a higher n . This would lead to faster flow too. Perhaps another sentence or two would help with clarity of the message here. Or simply delete the sentence “This might be linked ...”. The last sentence of the paragraph is clear and seems warranted by the results.

done

L270: shallow ice approximation => shallow shelf approximation [?]

done

L290: allows to => allows us to

done

L296: allows to => allows us to
done

L328: by e.g => by e.g.
done

L330: to assess => to assessing
done

L331: allows informed => helps to inform
done

L332: the Greenland or Antarctic Ice Sheet => continental-scale ice sheets [since these insights could apply to any ice sheet, right?]
done

Sensitivity of ice loss to uncertainty in flow law parameters in an idealized one-dimensional geometry

Maria Zeitz^{1,2}, Anders Levermann^{1,2,3}, and Ricarda Winkelmann^{1,2}

¹Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, P.O. Box 60 12 03, 14412 Potsdam, Germany

²University of Potsdam, Institute of Physics and Astronomy, Karl-Liebknecht-Str. 24-25, 14476 Potsdam, Germany

³LDEO, Columbia University, New York, USA.

Correspondence: Maria Zeitz (maria.zeitz@pik-potsdam.de), Ricarda Winkelmann (ricarda.winkelmann@pik-potsdam.de)

Abstract. Acceleration of the flow of ice drives mass losses in both the Antarctic and the Greenland Ice Sheet. The projections of possible future sea-level rise re<https://www.overleaf.com/projectly> on numerical ice-sheet models, which solve the physics of ice flow, melt, and calving. While major advancements have been made by the ice-sheet modeling community in addressing several of the related uncertainties, the flow law, which is at the center of most process-based ice-sheet models, is not in the focus of the current scientific debate. However, recent studies show that the flow law parameters are highly uncertain and might be different from the widely accepted standard values. Here, we use an idealized flowline setup to investigate how these uncertainties in the flow law translate into uncertainties in flow-driven mass loss. In order to disentangle the effect of future warming on the ice flow from other effects, we perform a suite of experiments with the Parallel Ice Sheet Model (PISM), deliberately excluding changes in the surface mass balance. We find that changes in the flow parameters within the observed range can lead up to a doubling of the flow-driven mass loss within the first centuries of warming, compared to standard parameters. The spread of ice loss due to the uncertainty in flow parameters is of the same order of magnitude as the increase in mass loss due to surface warming. While this study focuses on an idealized flowline geometry, it is likely that this uncertainty carries over to realistic three-dimensional simulations of Greenland and Antarctica.

Copyright statement. TEXT

15 1 Introduction

Current and future sea-level rise is one of the most iconic impacts of a warming climate and affects shorelines worldwide (Hinkel et al., 2014; Strauss et al., 2015). The contribution of the large ice sheets in Greenland and Antarctica to sea-level rise sums up to 13.7 + 14.0 mm over the last four decades (Mouginot et al., 2019; Rignot et al., 2019). It has been accelerating in recent years and is expected to further increase with sustained warming (Levermann et al., 2014, 2020; Mengel et al., 2016; Seroussi et al., 2020; Goelzer et al., 2020; Aschwanden et al., 2019; Bamber et al., 2019). Although some convergence can be observed in the projections of the median contribution of ice loss from Antarctica and Greenland, large uncertainties remain,

and coastal protection cannot rely on the median estimate since there is a 50% likelihood that it will be exceeded. Rather, an estimate of the upper uncertainty range is crucial. The most recent IPCC Special Report on the Ocean and Cryosphere in a Changing Climate provides projections of sea-level rise for the year 2100 of 0.43 m (0.29 – 0.59 m) and 0.84 m (0.61 – 1.10 m) for RCP2.6 and RCP8.5 scenarios, respectively (Pörtner et al., 2019). Other studies find slightly different (Goelzer et al., 2011, 2016; Huybrechts et al., 2011) and partly wider ranges (Levermann et al., 2020). Such projections are typically performed with process-based ice-sheet models which represent the physics in the interior and the processes at the boundaries of the ice-sheet.

In contrast to these processes at the boundaries of the ice sheet, many rheological parameters of the ice are typically not represented as an uncertainty in sea-level projections. The theoretical basis of ice flow, as implemented in ice-sheet models, has been studied in the lab and by field observations for more than half a century and is perceived as well established (Glen, 1958; Paterson and Budd, 1982; Budd and Jacka, 1989; Greve and Blatter, 2009; Cuffey and Paterson, 2010; Schulson and Duval, 2009; Duval et al., 2010). *Glen's flow law*, which relates stress and strain rate in a power law, is most widely used in ice-flow models. It is described in more detail in section 2.1. Some alternatives to the mathematical form of the flow law have been proposed: multi-term power laws like the *Goldsby-Kohlstedt* law or similar (Peltier et al., 2000; Pettit and Waddington, 2003; Ma et al., 2010; Quiquet et al., 2018) and anisotropic flow laws (Ma et al., 2010; Gagliardini et al., 2013) might be better suited to describe ice flow over a wide range of stress regimes. However, they have not been picked up by the ice-modeling community widely, possibly because this would require introducing another set of parameters which are not very well constrained.

Of all flow parameters, the enhancement factor is varied most routinely and its influence on ice dynamics is well understood (Quiquet et al., 2018; Ritz et al., 1997; Aschwanden et al., 2016). However, recent developments suggest that also the other parameters of the flow law are less certain than typically acknowledged in modelling approaches: A review of the original literature on experiments and field observations shows a large spread in the flow exponent n (which describes the nonlinear response in deformation rate to a given stress), which can be between 2 and 4, and the activation energies Q in the Arrhenius law (which describe the dependence of the deformation rate on temperature), which also can vary by a factor of two (Jellinek and Brill, 1956; Butkovich and Landauer, 1960; Mellor and Smith, 1967; Weertman, 1968; Mellor and Testa, 1969; Muguruma, Zeitz et al. submitted). New experimental approaches suggest a flow exponent larger than $n = 3$, which has been the most accepted value so far (Qi et al., 2017). Further, via an analysis of the thickness, surface slope and velocities of the Greenland Ice Sheet from remote sensing data, Bons et al. (2018) relate the driving stress to the ice velocities in regions where sliding is negligible, and can thus infer a flow exponent $n = 4$ under more realistic conditions. The activation energies Q in the Arrhenius law (which describe the dependence of the deformation rate on temperature), which also can vary by a factor of two (Glen, 1955; Nye, 1953; Mellor and Testa, 1969; Barnes et al., 1971; Weertman, 1973; Paterson, 1977; Goldsby and Kohlstedt, 2001; Trevisan et al., 2018).

Here we assess the implications of this uncertainty in simulations with the thermomechanically coupled Parallel Ice Sheet Model (the PISM authors, 2018; Bueler and Brown, 2009; Winkelmann et al., 2011), showing that variations in flow parameters have an important influence on flow-driven ice loss in an idealized flowline scenario.

This paper is structured as follows: in Section 2 we recapitulate the theoretical background of ice flow physics and describe the simulation methods used. The results of the equilibrium and warming experiments in a flow-line setup with different flow

parameters are presented in Section 3. Section 4 discusses the results and the limitations of the experimental approach and draws conclusions and suggests possible implications of these results.

2 Methods

60 2.1 Theoretical background of ice flow physics

The flow of ice cannot be described by the equations of fluid dynamics alone, but needs to be complemented by a material-dependent constitutive equation which relates the internal forces (stress) to the deformation rate (strain rate). Numerous laboratory experiments and field measurements show that the ice deformation rate responds to stress in a nonlinear way. Under the assumptions of isotropy, incompressibility and uni-axial stress this observation is reflected in Glen's flow law, which gives the
65 constitutive equation for ice,

$$\dot{\epsilon} = A\tau^n, \quad (1)$$

where $\dot{\epsilon}$ is the strain rate, τ the dominant shear stress, n the flow exponent and A the softness of ice (Glen, 1958).

Both, the flow exponent and the softness are important parameters which determine the flow of ice. Usually, the exponent n is assumed to be constant through space and time. Until today, there is no comprehensive understanding of all the physical
70 processes determining the softness A . It may depend ~~-,among others,-~~ on water content, impurities, grain size and anisotropy as well as temperature of the ice, among other things. Within the scope of ice-sheet modeling A is typically expressed as a function of temperature alone

$$A = A_0 \exp\left(-\frac{Q}{RT'}\right), \quad (2)$$

where A_0 is a constant factor, Q is an activation energy, R is the universal gas constant and T' the temperature **difference**
75 relative to the pressure melting point (Greve and Blatter, 2009; Cuffey and Paterson, 2010).

Due to pre-melt processes the softness responds more strongly to warming at temperatures close to the pressure melting point, which is often described by a piece-wise adaption of the activation energy Q (Barnes et al., 1971; Paterson, 1991), with a larger value of Q at temperatures $T' > -10^\circ\text{C}$. When using these piece-wise defined values for Q for warm and for cold ice in the functional form of the flow law, the respective factors A_0 ensure that the function is continuous at $T' = -10^\circ\text{C}$. A_0 is
80 therefore dependent on the values of the flow exponent n and both values of Q for cold and for warm ice.

The scalar form of Glen's flow law (Equation (1)) is only valid for uni-axial stresses, acting in only one direction. For a complete picture the stress is described as a tensor of order two. The generalized flow law reads

$$\dot{\epsilon}_{jk} = A(T')\tau_e^{n-1}\tau_{jk}, \quad (3)$$

where $\dot{\epsilon}_{jk}$ are the components of the strain rate tensor and τ_{jk} are the components of the stress deviator, τ_e is the effective
85 stress, which is closely related to the second invariant of the deviatoric stress tensor:

$$\tau_e^2 = \frac{1}{2} [\tau_{xx}^2 + \tau_{yy}^2 + \tau_{zz}^2] + \tau_{xy}^2 + \tau_{xz}^2 + \tau_{yz}^2. \quad (4)$$

Each component of the strain rate tensor depends on all the components of the deviatoric stress tensor through the effective stress τ_e .

90 ~~independently-independent~~ of the other approximations they might use (the PISM authors, 2018; Winkelmann et al., 2011; Greve, 1997; Pattyn, 2017; Larour et al., 2012; de Boer et al., 2013; Fürst et al., 2011; Lipscomb et al., 2018).

2.2 Ice flow model PISM

The simulations in this study were performed with the Parallel Ice Sheet Model (PISM) release stable v1.1. PISM uses shallow approximations for the discretized, physical equations: The shallow-ice approximation (SIA) (Hutter, 1983) and the shallow-shelf approximation (SSA) (Weis et al., 1999) are solved in parallel within the entire simulation domain. The shallow ice approximation is typically dominant in regions with high bottom friction, such that the vertical shear stresses dominate over horizontal shear stresses and longitudinal stresses. The shallow shelf approximation is typically dominant for ice shelves, with zero traction at the base of the ice, and for the fast flow regime in ice streams (Winkelmann et al., 2011). PISM assumes a non-sliding SIA flow and uses the results of the SSA approximations for fast flowing and sliding ice. In PISM, the flow law enters both the SIA and the SSA part of the velocities, as detailed in Winkelmann et al. (2011). It is possible to choose different flow exponents n for the SSA and the SIA, but the softness is the same for both approximations.

105 The simulations performed here use mostly the SIA mode: The geometry of a two dimensional ice sheet sitting on a flat bed and the SIA mode serve to study the effects of changes in flow parameters onto internal deformation and to separate those effects from changes in sliding, etc. Including the shallow shelf approximation reproduces and even enhances the effect of changes in the activation energies Q (see section 3.5).

2.3 Uncertainty in flow exponent and activation energies

The flow exponent n and the activation energies for warm and for cold ice, Q_w and Q_c , determine the deformation of the ice as a response to stress or temperature. A recent review (Zeitzi et al. submitted, see also literature in the introduction above) reveals a broad range of potential flow parameters n , Q_w and Q_c . The activation energy for cold ice Q_c is varied between 42 kJ/mol to 85 kJ/mol (~~the typical standard a typical reference~~ value is $Q_c = 60$ kJ/mol). The activation energy for warm ice Q_w is varied between 120 kJ/mol to 200 kJ/mol (~~the standard a reference~~ value is $Q_w = 139$ kJ/mol). For the flow exponent n , values as low as 1 have been reported, but since many experiments and observations confirm a nonlinear flow of ice, n has been varied between 2 and 4, with ~~the standard a reference~~ value of $n = 3$. The ~~standard values serve as a reference point and reference values above~~ correspond to the default values in many ice-sheet models (the PISM authors, 2018; Greve, 1997; Pattyn, 2017; Larour et al., 2012; de Boer et al., 2013; Fürst et al., 2011; Lipscomb et al., 2018).

2.4 Adaption of the flow factor A_0

The flow factor A_0 in the flow law must be adapted to fulfill the following conditions: First, the continuity of the piece-wise defined softness $A(T')$ must be ensured for all combinations of Q_w , Q_c and n . Secondly, a reference deformation rate $\dot{\epsilon}$ at a reference driving stress τ_0 and a reference temperature T'_0 (the PISM authors, 2018) should be maintained regardless of the parameters. This is because the coefficient and the power are non-trivially linked when a power law is fitted to experimental data. These conditions give:

$$A_{0,\text{old}} \cdot \exp\left(-\frac{Q_{\text{old}}}{RT'_0}\right) \cdot \tau_0^{n_{\text{old}}} = A_{0,\text{new}} \cdot \exp\left(-\frac{Q_{\text{new}}}{RT'_0}\right) \cdot \tau_0^{n_{\text{new}}}, \quad (5)$$

$$A_{0,\text{new}} = A_{0,\text{old}} \cdot \exp\left(-\frac{Q_{\text{old}} - Q_{\text{new}}}{RT'_0}\right) \cdot \tau_0^{n_{\text{old}} - n_{\text{new}}}. \quad (6)$$

If the reference temperature is $T'_0 < -10^\circ\text{C}$ the values for cold ice $A_{0,c}$ and Q_c are used in the equation above, or else $A_{0,w}$ and Q_w are used. The corresponding $A_{0,\text{new}}$ for ~~warm or cold~~ cold and warm ice respectively is calculated from the continuity condition at $T' = -10^\circ\text{C}$. For e.g. $T'_0 < -10^\circ\text{C}$ it follows

$$A_{0,c,\text{new}} = A_{0,c,\text{old}} \cdot \exp\left(-\frac{Q_{c,\text{old}} - Q_{c,\text{new}}}{RT'_0}\right) \cdot \tau_0^{n_{\text{old}} - n_{\text{new}}} \quad \text{and} \quad (7)$$

$$A_{0,w,\text{new}} = A_{0,c,\text{new}} \cdot \exp\left(-\frac{(Q_{c,\text{new}} - Q_{w,\text{new}})}{R \cdot 263.15\text{K}}\right). \quad (8)$$

Here we choose $\tau_0 = 80\text{kPa}$ as a typical stress in a glacier and $T'_0 = -20^\circ\text{C}$. Choosing another τ_0 ~~in~~ on the same order of magnitude has only little effect on the differences in dynamic ice loss. Choosing another T'_0 on the other hand influences how the softness changes with the activation energy Q , see Supplemental Figure S1. With T'_0 closer to the melting temperature, the difference in softness at the pressure melting point decreases thus the ice loss is less sensitive to changes in the activation energy Q .

2.5 Experimental design

The study is performed in a flow-line setup, similar to Pattyn et al. (2012), where the computational domain has an extent of 1000km in x -direction and 3km in y -direction (with a periodic boundary condition). The spatial horizontal resolution is 1 km. The ice rests on a flat bed of length $L = 900\text{ km}$ with a fixed calving front at the edge of the bed, such that no ice shelves can form (Figure 1). In contrast to Pattyn et al. (2012), the temperature and the enthalpy of ice sheet are allowed to evolve freely.

The model is initialized with a spatially constant ice thickness and is run into equilibrium for different combinations of flow parameters Q_c , Q_w and n . The ice surface temperature is altitude dependent, $T_s = -6^\circ\text{C}/\text{km} \cdot z - 2^\circ\text{C}$, where z is the surface elevation in km. The accumulation rate is constant in space and time for each simulation. Similar to the MISMIP setup, there is no geothermal heat flux prescribed. In the warming experiments, for each ensemble member an instantaneous temperature increase of $\Delta T \in [1, 2, 3, 4, 5, 6]^\circ\text{C}$ is applied to the ice surface for the duration of 15,000 years (until a new equilibrium is reached), while the climatic mass balance remains unchanged. That means, the temperature increase can lead to




Figure 1. Sketch of the flow-line setup The ice is sitting on a flat bed, the fixed calving front does not allow ice-shelves. The accumulation rate is constant throughout the simulation domain and the temperature is altitude dependent.

145 an acceleration of ice flow, but is prohibited from inducing additional melt. This idealized forcing allows us to disentangle the effect of warming on the ice flow from climatic drivers of ice loss.

The thickness profile of the equilibrium state is similar to the *Vialov profile* (see e.g. Cuffey and Paterson (2010); Greve and Blatter (2009)). However, in contrast to the isothermal Vialov profile, here the temperature of the ice is allowed to evolve freely, leading to a non-uniform softness of the ice (the PISM authors, 2018). The extent in x -direction is given by the geometry of
 150 the setup, a flat bed with a calving boundary condition at the margin, and the height and shape of the ice sheet depend on the flow parameters n , Q_w and Q_c and the accumulation rate a .

3 Results

3.1 Effect of activation energies in model simulations compared to analytical solution

In order to gain a deeper understanding of the influences of Q_c and Q_w on the equilibrium shape of ice-sheets, we here compare
 155 the simulated results to analytical considerations based on the Vialov profile.

At a fixed accumulation rate of $a = 0.5\text{m/yr}$, each flow parameter combination leads to an equilibrium state with a thickness profile similar to the Vialov profile but differences in maximal thickness and volume (Figure 2 a). Overall, high activation energies increase ice-flow velocities and reduce the ice-sheet volume. The activation energy for warm ice, Q_w , affects the volume and the velocities more strongly than the activation energy for cold ice, Q_c . A high Q_w leads to softer ice close to the
 160 pressure melting point (supplemental Figure 1) and at the base of the ice sheet, which leads to higher velocities and a lower equilibrium volume of the ice sheet while a low Q_w leads to stiffer ice close to the pressure melting point and at the base of the ice sheet and in consequence the velocities decrease and the volume increases (Figure 2, b and c). For a fixed Q_w , the volume appears to decrease linearly with increasing Q_c and the velocity appears to increase linearly with increasing Q_c .




Figure 2. Effect of activation energies on equilibrium volume and velocities with fixed accumulation rate $a = 0.5\text{m/yr}$ and flow exponent $n = 3$. Thickness profiles of equilibrium states for different combinations of activation energies Q_w and Q_c (a). Relative difference of average equilibrium volumes (b) and velocities (c) compared to the reference state with standard parameters for parameter combinations of Q_w and Q_c . Q_c is shown on the x-axis and Q_w is given through the color of the the markers (Blue: $Q_w = 120$ kJ/mol, orange: $Q_w = 139$ kJ/mol, red: Blue: $Q_w = 200$ kJ/mol)




Figure 3. Comparison of simulated equilibrium thickness with analytical results: (a) Dots: Maximal thickness h_m of the simulated polythermal ice sheet versus the analytical solution for the maximal thickness h_m of a temperate Vialov profile with the same flow parameters and accumulation rate. Colors indicate the flow-parameter combination. The grey line indicates identity. Short, dashed lines indicate the analytical h_m with a temperature lower than the pressure melting point versus h_m at the pressure melting point with the same flow parameters and accumulation rate. The temperature, which fits the simulated results best, is indicated in the legend. **(b)** Ratio of the simulated h_m to the analytic h_m (assuming a temperate ice sheet) versus Q_c for different parameter combinations Q_c, Q_w . The value of Q_w is indicated by the color.

The maximal thickness of an isothermal ice sheet can be estimated with the Vialov profile

$$165 \quad h_m = 2^{n/(2n+2)} L^{1/2} \left(\frac{(n+2)a}{2A(T')(\rho g)^n} \right) \quad (9)$$

with the Glen exponent n , the ice sheet extent $2L$, the pressure adjusted temperature T' , the gravity g , and the ice density ρ (Greve and Blatter, 2009). The Vialov thickness of a temperate ice sheet (isothermal at the pressure melting point), where the softness is evaluated at the pressure melting point depending on the activation energies Q_c and Q_w (see Equation (2)) gives a lower bound to the thickness, given the same geometry and flow parameters. The simulated maximal thickness is larger than the lower bound for all parameter combinations (Figure 3 a, lower bound indicated by grey line) and the ratio between the maximal thickness h_m from the PISM simulation to the lower bound from the Vialov profile depends on both, Q_w and Q_c . The ratio increases with higher Q_w and decreases with higher Q_c (Figure 3 b). The ice-sheet thickness of the polythermal ice sheet, as simulated with PISM, matches well the Vialov thickness calculated with Equation (9), if an effective temperature $T'_{\text{eff}} < 0^{\circ}\text{C}$ is assumed. The effective temperature T'_{eff} , which matches simulations best, varies for different Q_w . For $Q_w =$ 175 120 kJ/mol, an effective temperature of $T'_{\text{eff}} = -5^{\circ}\text{C}$ matches well the equilibrium thickness of the polythermal ice sheets. For

$Q_w = 200 \text{ kJ/mol}$, an effective temperature of $T'_{\text{eff}} = -3.3^\circ\text{C}$ matches well the equilibrium thickness of the polythermal ice sheets. These differences can be partly explained by the altitude-dependent surface temperature: The maximal thickness of the ice sheets varies by approximately 800 m, which leads to a difference in ice surface temperature of approximately 4.8°C between the thickest and the thinnest ice and thus influences the temperature within the ice sheet.

180 The relative difference of average velocities $d_v = (\bar{v} - \bar{v}_0) / \bar{v}_0$ spans from $d_v = -7\%$ (with a corresponding relative difference in ice sheet volume of $d_{vol} = +10\%$) for the lowest combination of activation energies to $d_v = +18\%$ with a difference in volume of $d_{vol} = -15\%$ for the highest combination of values for Q_c and Q_w (Figure 2 b).

3.2 Ice-sheet initial states

In order to keep the initial ice volume largely fixed (with variations of less than one percent) in the warming experiments, we
185 adapt the accumulation rate for each parameter combination of Q_c and Q_w .

Since simulations with high activation energies Q_w have a smaller equilibrium volume at the same accumulation rate than simulations with standard activation energies, the accumulation rate a is increased to maintain an equilibrium volume close to the reference value. Simulations with low activation energies Q_c have a higher volume at the same accumulation rate, so the accumulation rate a is decreased. In the case of an isothermal ice sheet the maximal thickness and the volume can be computed
190 analytically as shown above in Equation (9). In our model simulations, however, the temperature distribution within the ice can evolve freely, thus the softness is not uniform and an analytical solution cannot be found.

In order to find the right adaptation for the accumulation rates, we start from the ~~volume-calculated-ice profile~~ from the isothermal approximation as a first guess and run the model into equilibrium. If the relative difference between the new equilibrium volume and the standard equilibrium volume exceeds 1%, we further change the accumulation rate and repeat the
195 equilibrium simulation, always starting from the same initial state. The final equilibrium states found via this iterative approach differ by max. 0.8% in ice volume (supplemental Figure S2) and the difference in maximal thickness is less than 100m (Figure 4, a and b).

For the combination of high activation energies Q_w and Q_c , the relative differences $d_x = (x - x_0) / x_0$ of both, adapted accumulation rates a and mean surface velocities v , increase by more than 300% (Fig. 4, c and d) and for the combination of
200 low activation energies Q_c and Q_w both, adapted accumulation rates a and surface velocities v are approximately 50% lower compared to the case with standard parameters. Both, the accumulation rate and the velocities, change in the same way since they balance each other in equilibrium. A change in accumulation rates controls the vertical velocity profile and thus influence the thermodynamics in the ice, which leads to differences in the temperatures of the ice sheet (pressure adjusted temperature distributions shown in Supplemental Figure S4 a). The change in temperature is most prominent at the top of the ice sheet, where higher accumulation rates (associated with high activation energies) lead to lower temperatures and vice versa. Thus the temperature change introduced from increased accumulation counteracts the effect of increased softness. In order to estimate how changed temperature on the one hand and changed flow parameters on the other hand impacts the resulting ice softness, either one was kept fixed. The effect of the temperature changes on the ice softness is negligible, compared to parameter changes (see supplemental Figure S4 b,c and d).




Figure 4. Effect of flow parameters on equilibrium state without warming with adapted accumulation rates and flow exponent $n = 3$. (a) Thickness profile of equilibrium states for parameter combinations of Q_w and Q_c with a zoom on the ice divide (b). Relative difference of accumulation rates (c) needed to keep the ice sheet volume in equilibrium close to the reference simulations with standard flow parameters and relative difference in average surface velocities (d) versus Q_c . The value of Q_w is given by the color.




Figure 5. Comparison of simulated equilibrium thickness with analytical results: (a) Dots: Maximal thickness h_m of the simulated polythermal ice sheet versus the analytical solution for the maximal thickness h_m of a temperate Vialov profile with the same flow parameters and accumulation rate. Colors indicate the parameter combination. The grey line indicates identity. Short, dashed lines indicate the analytical h_m with a temperature lower than the pressure melting point versus h_m at the pressure melting point with the same flow parameters and accumulation rate. The temperature, which fits the simulated results best, is indicated in the legend. (b) Ratio of the simulated h_m to the analytic h_m (assuming a temperate ice sheet) versus Q_c for different parameter combinations Q_c, Q_w . The value of Q_w is indicated by the color.

210 The maximal thickness of the polythermal simulated ice sheet is approximately 13-16% larger than the lower bound estimated with a temperate ice sheet (Figure 5, a and b) with the same flow parameters and accumulation rates. Similar to the case with fixed accumulation rates, the simulated thickness matches the Vialov thickness well, if an effective temperature $T'_{\text{eff}} < 0^\circ\text{C}$ is assumed. The effective temperature, ~~which matches that matches~~ simulations best, varies for different Q_w , from -5°C for $Q_w = 120$ kJ/mol to -3.6°C for $Q_w = 200$ kJ/mol. This difference can not be sufficiently explained by variations in surface

215 temperature due to the difference in ice-sheet thickness. Rather the higher effective temperatures are linked to increased flow velocities of the ice, which in turn might lead to strain heating. In simulations with a high Q_w the simulated thickness has a higher discrepancy to the estimated lower bound (assuming a temperate ice sheet) than simulations with a low Q_w . In contrast to the case with fixed accumulation rate (Figure 3) the ratio between the estimated and the simulated thickness depends only very little on Q_c .

Disentangling the purely flow-driven ice losses from the influences of melting, different initial temperature profiles and variations in sliding requires several conditions: 1) The initial volume is fixed, which is here attained through adjustment of the accumulation rate for the different flow parameter combinations as explained in section 3.2. 2) The surface mass balance is fixed, i.e., we do not allow for additional melt, and the accumulation rate does not change with warming. 3) Sliding is effectively inhibited (which is here ensured by applying an SIA-only condition).

The effect of the temperature increase is limited to warming at the ice surface which can propagate into the interior of the ice sheet though diffusion and advection. Warming makes the ice softer thus accelerates the flow and ice discharge. Since temperature diffusion in an ice sheet is a very slow process, we apply the temperature anomaly for a total duration of 15,000 years. The total mass balance is evaluated and compared to the standard parameter simulation after 100, 1000 and 10,000 years of warming. A new equilibrium state is reached after 10,000 years for all parameter combinations (see longer time-series in Supplemental Figure S3).




Figure 6. Time series for flow driven ice discharge under 2°C warming: (a) Time evolution of ice loss with different activation energies Q_c and Q_w and the flow exponent $n = 3$, subject to a temperature anomaly forcing of $\Delta T = 2^\circ\text{C}$. (b) Zoom on the first 100 years.

In the experiments, the ice sheet loses mass for all warming levels and all parameter combinations. However, the amount and rate of the ice loss is dependent on the flow parameters. Figure 6 shows the ice-sheet response to a warming of 2°C. For a fixed flow exponent of $n = 3$ the fastest ice loss is observed for the flow parameter combination of $Q_c = 85\text{kJ/mol}$ and $Q_w = 200\text{kJ/mol}$ and the slowest ice loss for $Q_c = 42\text{kJ/mol}$ and $Q_w = 120\text{kJ/mol}$. Simulations with $Q_w = 200\text{kJ/mol}$ reach a new, temperature adapted equilibrium already after 2,000 yrs, while simulations with lower Q_w continue to lose mass.




Figure 7. Effect of activation energy on flow driven ice discharge under 2°C warming: Relative difference of flow-driven ice loss after 100 (a), 1000 (b) and 10,000 (c) years versus Q_c . The value of Q_w is given by the color. The simulations have reached a new equilibrium after 10,000 years.

The sensitivity to variations in flow parameters is measured via the relative differences for flow-driven ice loss $d_m = (\Delta m - \Delta m_0) / \Delta m_0$, where the reference Δm_0 is always given by the simulation with standard parameters under the same temperature increase (Figure 7). While the long term response to warming, after 10,000 years, is not very sensitive to the particular choice of flow parameters, the rate of flow-driven ice loss is. The largest relative differences in ice loss is found in the first century after the temperature increase (Figure 7, a), indicating that high Q_w speed up the flow-driven ice loss. Under 2°C of warming, ice loss after 100 years is enhanced more than two-fold (i.e. increased by up to 118%) in simulations with $Q_w = 200$ kJ/mol, while low Q_w reduces the relative ice loss by up to 37%.




Figure 8. Effect of temperature forcing and activation energy on flow-driven ice (a) Time evolution of ice loss under warming of 1°C and 6°C . For warming of $\Delta T = 1^{\circ}\text{C}$ the conceptual ice sheet loses 0.35% of ice after 1000 yr for standard parameters (solid, yellow line). Variations in the activation energies Q lead to variations in ice loss on the same order of magnitude (shaded area). For a warming of $\Delta T = 6^{\circ}\text{C}$ the conceptual ice sheet loses 1.89% of ice after 1000 yr for standard parameters (solid, red line). The variations in ice loss due to different parameters for the activation energy Q (shaded area) are strongly asymmetrical and, in particular during the first 300 years, high compared to the total amount of ice loss. (b) Uncertainty in flow induced ice loss after 100 years of simulation time over all combinations of Q_w , Q_c and temperature anomalies ΔT . The flow exponent $n = 3$ is kept fixed for all simulations.

The effect of the flow parameters on flow-driven ice loss upon warming is robust for different temperature increases. Ice losses as well as the spread in flow-driven ice loss both increase for higher warming levels (see Figure 8). For a warming of $\Delta T = 1^{\circ}\text{C}$ the idealized ice sheet loses 0.09% after 100 yr and 0.35% of ice after 1000 yr for standard parameters. For a warming of $\Delta T = 6^{\circ}\text{C}$ the ice sheet loses 0.46% after 100 yr and 1.89% of ice after 1000 yr for standard parameters (solid, red line). For comparison, the Greenland Ice Sheet has lost approximately 0.18% of its mass in the period between 1972 and 2018 (Mouginot et al., 2019), which includes all processes: increase in flow, melting, and sliding.

The effect of flow parameter changes onto the purely flow-driven ice loss after 100 years is of the same order of magnitude as the effect of surface warming by several degrees. In particular the uncertainty ranges of ice loss for warming of 2°C and warming of 6°C overlap (Figure 8 b), when solely considering the ice loss is driven by changes in flow and excluding surface mass balance changes.

3.4 Influence of the flow exponent n




Figure 9. Effect of the flow exponent and activation energies on flow-driven ice loss after 100 years under 2°C of warming: Relative difference in flow-driven ice discharge for $n = 2$ (a), $n = 3$ (b) and $n = 4$ (c) for different combinations of the flow exponent n and activation energies Q_c and Q_w . The reference is always a simulation performed with standard parameters $n = 3$, $Q_c = 60$ kJ/mol and $Q_w = 139$ kJ/mol.




Figure 10. Effect of the flow exponent and activation energies on mean velocity change after 100 years under 2°C of warming: Relative difference in average surface velocity f or $n = 2$ (a), $n = 3$ (b) and $n = 4$ (c) for different combinations of the flow exponent n and activation energies Q_c and Q_w . The reference is always a simulation performed with standard parameters $n = 3$, $Q_c = 60$ kJ/mol and $Q_w = 139$ kJ/mol. Variations in the flow exponent n do not significantly influence the relative difference of mean velocities after 100 years.

255 Variations in the flow exponent n do not change the qualitative effect of variations in activation energies Q on the ice loss. After 100 years for a temperature anomaly of $\Delta T = 2^\circ\text{C}$ a higher n seems to mitigate the effect of the activation energy on differences in ice loss, while a lower n seems to enhance this effect (Figure 9). However, the effect of variations in activation energy on the average surface velocity is almost independent of the choice for the flow exponent n (Figure 10).

The influence of the activation energies Q_c and Q_w on ice flow is similar even with different flow exponents n . This is
 260 robust for different warming scenarios from $+1$ to $+6^\circ\text{C}$. A higher flow exponent n , which leads to a more pronounced nonlinearity in ice flow, does not enhance but reduce variations in dynamic ice loss. ~~This might be linked to the details of the parametrization in the flow law: As seen in both equations and, the factor A_0 decreases if the flow exponent n is reduced and $n_{\text{new}} < n_{\text{old}}$.~~ Compared to the nonlinear stress dependency τ^n in the flow law the temperature dependent softness $A(T') = A_0 \cdot \exp(-Q/RT')$ becomes less important with increasing flow exponent n .

265 3.5 Robustness of results to changes in accumulation and sliding

The overall effect of uncertainties in the activation energies Q remains robust, even if an additional driver of ice loss is taken into account. In a simulation where in addition to warming of 2°C we also reduce the accumulation rate by 50%, the ice losses remain dependent on the flow parameters Q_c and Q_w (Figure 11, lines indicate results without a change in accumulation rate, analogous to Figure 7 and squares indicate results with an additional 50% decrease in accumulation rate). After 100 years of
 270 forcing the relative spread of ice loss is slightly larger if accumulation changes are included. In particular, for $Q_w = 200$ kJ/mol,




Figure 11. Effect of the flow exponent and activation energies on flow-driven ice loss under 2°C of ice warming in combination with a 50% reduction in accumulation rates: Relative difference in flow-driven ice discharge after 100 (a), 1000 (b) and 10,000 (c) years. The ice sheet has reached a new equilibrium after 10,000 years. Relative difference for 2°C warming with an additional reduction of the accumulation rate of 50% (squares) are compared to the results without changes in the accumulation rate (lines, also see Figure 7).

the relative increase of mass loss mounts from 118% to 190%. On longer time scales, the spread in ice loss is reduced (after 10,000 years of forcing, when the ice sheet has reached a new equilibrium, the relative spread is below $\pm 10\%$).

When sliding is taken into account via the shallow [ice-shelf](#) approximation for sliding ice (see the PISM authors (2018)) the uncertainty in flow parameters leads to relative changes in ice loss from -30% to +470% after 100 years, which is a considerably larger spread than without sliding. The relative differences decrease with time, but remain ranger than without sliding. After 1000 years the ensemble member with low activation energies lost 40% less ice than the standard parametrization and high activation energies almost double the ice loss (+90%). After 10,000 years, when the ice sheets have reached a new equilibrium, the relative differences still range from -16% to +40% (see Figure 12).

4 Discussion and Conclusion

In this study we present a first attempt to disentangle and quantify the effect of uncertainties in the flow law parameters, in particular the activation energies Q and the flow exponent n , onto ice dynamics.

The effect of ice rheology in ice-sheet models has been adressed in several studies with different experimental setups and different time frames. In particular the effect of the enhancement factors, which are often used to approximate the change in ice flow due to anisotropy, has been explored (Ritz et al., 1997; Ma et al., 2010; Humbert et al., 2005; Quiquet et al., 2018). In addition, the effect of the initial conditions (Seroussi et al., 2013; Nias et al., 2016; Humbert et al., 2005) and the effect




Figure 12. Effect of activation energy on flow driven ice discharge under 2°C warming, including sliding: Relative difference of flow and sliding driven ice loss after 100 (a), 1000 (b) and 10,000 (c) years. The simulations have reached a new equilibrium after 10,000 years.

of the mathematical form of the flow law itself (Quiquet et al., 2018; Peltier et al., 2000; Pettit and Waddington, 2003) have been studied. These studies have been crucial for the understanding of different enhancement factors in the shallow ice and the shallow shelf approximation (Ma et al., 2010), for the reconciliation of the aspect ratios of the Greenland Ice Sheet and the Laurentide Ice Sheet during the last glacial maximum (Peltier et al., 2000) and the ice flow in Antarctica and the Greenland Ice Sheet (Ritz et al., 1997; Seroussi et al., 2013; Quiquet et al., 2018; Nias et al., 2016; Humbert et al., 2005).

However, the approach presented in this manuscript is different in two important aspects: Firstly, the systematic study of not only the flow exponent n but also the activation energies Q has not been performed so far. Secondly, the idealized experimental setup, as presented in this study, allows us to disentangle the effects of the flow itself from other drivers and other sources of uncertainty. Several conditions need to hold to this end: The ice sheet is sitting on a flat bed and its maximal extent is determined by a calving front at the borders of the bed, thus no ice-ocean interactions or impacts of the bed geography influence the ice flow. Sliding is generally inhibited (the ice dynamics is described by the shallow ice approximation, with zero basal velocity), no changes in sliding velocity influence the ice flow. The accumulation rate is fixed and independent of the temperature change, so that the ice loss is only driven by changes in flow and not by melting. These idealizations allow a clear understanding of the impact of the flow exponent and the activation energies on ice flow. In addition, they allow us to compare the simulations of the polythermal ice sheet to the analytically solvable limit of an isothermal ice sheet by using the Vialov approximation.

In this setup the largest effect of the uncertainties in the flow parameters is observed in the first century after warming, while the effect of the uncertainties on ice loss becomes less important as the ice approaches a new equilibrium. Uncertainties in the activation energies alone account for up to a doubling in ice loss during the first 100 years of warming and are on the

same order of magnitude as the effects of increased temperature forcing, under fixed surface mass balance. This effect remains
305 robust, even if changes in the surface mass balance are taken into account. Reducing the surface mass balance by 50%, which
is comparable to the changes in total surface mass balance of the Greenland Ice Sheet from 1972 to 2012 (Mouginot et al.,
2019), increases the effect of the flow parameters on a timescale of 100 years and remains comparable on a timescale of 1000
years. Only as the ice sheet approaches its new equilibrium, the effect of the flow parameters becomes negligible. Allowing for
not only flow but sliding while keeping all other conditions equal increases the effect of flow parameters substantially, leading
310 to up to a five-fold increase in ice loss after 100 years compared with standard parameters.

Acknowledging the uncertainty in flow parameters might slightly shift the interpretation of previous studies. For instance,
the effect of the initial thermal regime, as studied by Seroussi et al. (2013) could be enhanced if the the activation energies were
higher than assumed, by making the ice softness more sensitive to changes in temperature. The crossover stress in the multi-
term flow law presented by Pettit and Waddington (2003), at which the linear and the cubic term are of the same importance, is
315 highly sensitive to the values of the activation energies. The positive feedback through shear heating, as studied for example by
Minchew et al. (2018), could also be enhanced if activation energies were higher than usually assumed. The uncertainty in the
flow-law parameters may further provoke a re-evaluation of other parameters, e.g. concerning melting and basal conditions. In
particular, Bons et al. (2018) thorough analysis of observational data of the Greenland Ice Sheet supports a flow exponent of
 $n = 4$, not the standard value of $n = 3$, which is in line with recent laboratory experiments which also find $n > 3$ (Qi et al.,
320 2017). Assuming a higher flow exponent $n = 4$ has shown to significantly reduce the previously assumed area where sliding is
possible (Bons et al., 2018; MacGregor et al., 2016). Moreover, both the flow exponent n and the activation energies Q feed
into the grounding line flux formula (Schoof, 2007). In several ice-sheet models, this formula is used to determine the position
and the flux over the grounding line in transient simulations (Reese et al., 2018). A change in the flow parameters n and Q has
thus implications for the advance and retreat of grounding lines in simulations of the Antarctic Ice Sheet and possibly the onset
325 of the marine ice sheet instability, a particularly relevant process for the long-term stability of the Antarctic Ice Sheet. On the
Greenland Ice Sheet increased ice flow might drives ice masses into ablation regions, where the ice melts. A possible effect of
uncertainty in flow parameters on this particular feedback remains to be explored. Aschwanden et al. (2019) have found that
uncertainty in ice dynamics plays a major role for mass loss uncertainty during the first 100 years of warming. While their
study attributes the uncertainty mostly to large uncertainties in basal motion and only to a lesser extent to the flow via the
330 enhancement factor, the uncertainties of the flow law and of the basal motion are not independent, as suggested by e.g. Bons
et al. (2018).

While the conclusions from the idealized experiments presented here cannot be transferred directly to ~~assess~~assessing
uncertainty in sea-level rise projections, they are an important first step which ~~allows informed~~helps to inform choices about
parameter variations in more realistic simulations of ~~the Greenland or Antaretic Ice Sheet~~continental-scale ice sheets.

335 *Code and data availability.* Data and code are available from the authors upon request.

Author contributions. R.W. and A.L. conceived the study. M.Z., A.L. and R.W. designed the research and contributed to the analysis. M.Z. carried out the literature review and the analysis. M.Z., R.W. and A.L. wrote the manuscript.

Competing interests. No competing interest.

Acknowledgements. M.Z. and R.W. are supported by the Leibniz Association (project DOMINOES). R.W. is grateful for support by the
340 Deutsche Forschungsgemeinschaft (DFG) through grants WI4556/3-1 and WI4556/5-1 and by the PalMod project (FKZ: 01LP1925D),
supported by the German Federal Ministry of Education and Research (BMBF) as a Research for Sustainability initiative (FONA). This
research was further supported by the European Union's Horizon 2020 research and innovation programme under grant agreement no.
820575 (TiPACCs). Development of PISM is supported by NASA grant NNX17AG65G and NSF grants PLR-1603799 and PLR-1644277.
The authors gratefully acknowledge the European Regional Development Fund (ERDF), the German Federal Ministry of Education and
345 Research and the Land Brandenburg for supporting this project by providing resources on the high performance computer system at the
Potsdam Institute for Climate Impact Research. We thank Hilmar Gudmundsson, David Prior and Thomas Kleiner for insightful discussions.

References

- Aschwanden, A., Fahnestock, M. A., and Truffer, M.: Complex Greenland outlet glacier flow captured, *Nature Communications*, 7, 10524, <https://doi.org/10.1038/ncomms10524>, <http://dx.doi.org/10.1038/ncomms10524>, 2016.
- 350 Aschwanden, A., Fahnestock, M. A., Truffer, M., Brinkerhoff, D. J., Hock, R., Khroulev, C., Mottram, R., and Khan, S. A.: Contribution of the Greenland Ice Sheet to sea level over the next millennium, *Science Advances*, 5, eaav9396, <https://doi.org/10.1126/sciadv.aav9396>, <http://advances.sciencemag.org/lookup/doi/10.1126/sciadv.aav9396>, 2019.
- Bamber, J. L., Oppenheimer, M., Kopp, R. E., Aspinall, W. P., and Cooke, R. M.: Ice sheet contributions to future sea-level rise from structured expert judgment, *Proceedings of the National Academy of Sciences*, 116, 11 195–11 200, <https://doi.org/10.1073/pnas.1817205116>, <https://www.pnas.org/content/116/23/11195>, 2019.
- 355 Barnes, P., Tabor, D., and Walker, J. C. F.: The friction and creep of polycrystalline ice, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 324, 127–155, 1971.
- Blinov, K. and Dmitriev, D.: Otsenka reologicheskikh parametrov Ida po rezultatam mnogoletnih nabludeniy v skvazhinah na st. Vostok v Antarktide [Rheological ice parameters estimations on the basis of measurements in bore-holes at Vostok Station, Antarctica], *Antarctica*, 26, 95–106, 1987.
- 360 Bons, P. D., Kleiner, T., Llorens, M.-G., Prior, D. J., Sachau, T., Weikusat, I., and Jansen, D.: Greenland Ice Sheet: Higher Nonlinearity of Ice Flow Significantly Reduces Estimated Basal Motion, *Geophysical Research Letters*, 45, 6542–6548, <https://doi.org/10.1029/2018GL078356>, <http://doi.wiley.com/10.1029/2018GL078356>, 2018.
- Bowden, F. P. and Tabor, D.: *The friction and lubrication of solids*. Vol. 2, OUP, 1964.
- 365 Bromer, D. J. and Kingery, W. D.: Flow of polycrystalline ice at low stresses and small strains, *Journal of Applied Physics*, 39, 1688–1691, <https://doi.org/10.1063/1.1656416>, 1968.
- Budd, W. F. and Jacka, T. H.: A review of ice rheology for ice sheet modelling, *Cold Regions Science and Technology*, 16, 107–144, 1989.
- Bueler, E. and Brown, J.: Shallow shelf approximation as a "sliding law" in a thermomechanically coupled ice sheet model, *Journal of Geophysical Research: Solid Earth*, 114, 1–21, <https://doi.org/10.1029/2008JF001179>, 2009.
- 370 Butkovich, T. and J.K, L.: *The Flow Law For Ice*, <http://hydrologie.org/redbooks/a047/04734.pdf>, 1957.
- Butkovich, T. R. and Landauer, J. K.: Creep of Ice at Low Stresses, <https://doi.org/10.1038/232222b0>, 1960.
- Craw, L., Qi, C., Prior, D. J., Goldsby, D. L., and Kim, D.: Mechanics and microstructure of deformed natural anisotropic ice, *Journal of Structural Geology*, 115, 152–166, <https://doi.org/10.1016/j.jsg.2018.07.014>, <https://doi.org/10.1016/j.jsg.2018.07.014>, 2018.
- Cuffey, K. M. and Paterson, W. S. B.: *The Physics of glaciers*, Elsevier Inc., 4 edn., <http://linkinghub.elsevier.com/retrieve/pii/0016718571900868>, 2010.
- 375 Cyprych, D., Piazzolo, S., Wilson, C. J., Luzin, V., and Prior, D. J.: Rheology, microstructure and crystallographic preferred orientation of matrix containing a dispersed second phase: Insight from experimentally deformed ice, *Earth and Planetary Science Letters*, 449, 272–281, <https://doi.org/10.1016/j.epsl.2016.06.010>, <http://dx.doi.org/10.1016/j.epsl.2016.06.010>, 2016.
- de Boer, B., van de Wal, R. S. W., Lourens, L. J., Bintanja, R., and Reerink, T. J.: A continuous simulation of global ice volume over the past 1 million years with 3-D ice-sheet models, *Climate Dynamics*, 41, 1365–1384, <https://doi.org/10.1007/s00382-012-1562-2>, <http://link.springer.com/10.1007/s00382-012-1562-2>, 2013.
- 380 Doake, C. S. M. and Wolff, E. W.: Flow law for ice in polar ice sheets, *Nature*, 314, 255–257, <https://doi.org/10.1038/315279a0>, 1985.
- Duval, P.: Fluage de la glace polycrystalline, *Rheologica Acta*, 13, 562–566, <https://doi.org/10.1007/BF01521756>, 1974.

- Duval, P.: Lois du fluage transitoire ou permanent de la glace polycristalline pour divers états de contrainte, *Annals of Geophysics*, 32, 385 335–350, 1976.
- Duval, P. and Gac, H. L.: Mechanical behaviour of antarctic ice, *Annals of Glaciology*, 3, 92–95, 1982.
- Duval, P., Montagnat, M., Grennerat, F., Weiss, J., Meyssonier, J., and Philip, A.: Creep and plasticity of glacier ice: a material science perspective, *Journal of Glaciology*, 56, 1059–1068, <https://doi.org/10.3189/002214311796406185>, https://www.cambridge.org/core/product/identifier/S0022143000213269/type/journal_article, 2010.
- 390 Fletcher, N.: *The chemical physics of ice*. Cambridge, Cambridge University Press, Cambridge, 1970.
- Fürst, J. J., Rybak, O., Goelzer, H., De Smedt, B., De Groen, P., and Huybrechts, P.: Improved convergence and stability properties in a three-dimensional higher-order ice sheet model, *Geoscientific Model Development*, 4, 1133–1149, <https://doi.org/10.5194/gmd-4-1133-2011>, 2011.
- Gagliardini, O., Zwinger, T., Gillet-Chaulet, F., Durand, G., Favier, L., de Fleurian, B., Greve, R., Malinen, M., Martín, C., Råback, P., 395 Ruokolainen, J., Sacchetti, M., Schäfer, M., Seddik, H., and Thies, J.: Capabilities and performance of Elmer/Ice, a new-generation ice sheet model, *Geoscientific Model Development*, 6, 1299–1318, <https://doi.org/10.5194/gmd-6-1299-2013>, <https://gmd.copernicus.org/articles/6/1299/2013/>, 2013.
- Glen, J. W.: Experiments on the Deformation of Ice, *Journal of Glaciology*, 2, 111–114, <https://doi.org/10.1017/S0022143000034067>, https://www.cambridge.org/core/product/identifier/S0022143000034067/type/journal_article, 1952.
- 400 Glen, J. W.: Rate of Flow of Polycrystalline Ice, *Nature*, 172, 721–722, <https://doi.org/10.1038/172721a0>, 1953.
- Glen, J. W.: The Creep of Polycrystalline Ice, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 228, 519–538, <https://doi.org/10.1098/rspa.1955.0066>, <http://linkinghub.elsevier.com/retrieve/pii/0165232X85900527><http://rspa.royalsocietypublishing.org/cgi/doi/10.1098/rspa.1955.0066>, 1955.
- Glen, J. W.: The mechanical properties of ice I. The plastic properties of ice, *Advances in Physics*, 7, 254–265, 405 <https://doi.org/10.1080/00018735800101257>, 1958.
- Goelzer, H., Huybrechts, P., Loutre, M.-F., Goosse, H., Fichet, T., and Mouchet, A.: Impact of Greenland and Antarctic ice sheet interactions on climate sensitivity, *Climate dynamics*, 37, 1005–1018, 2011.
- Goelzer, H., Huybrechts, P., Loutre, M.-F., and Fichet, T.: Last Interglacial climate and sea-level evolution from a coupled ice sheet–climate model, *Climate of the Past*, 12, 2195–2213, <https://doi.org/10.5194/cp-12-2195-2016>, <https://www.clim-past.net/12/2195/2016/>, 2016.
- 410 Goelzer, H., Nowicki, S., Payne, A., Larour, E., Seroussi, H., William, H., Gregory, J., Abe-ouchi, A., Shepherd, A., Simon, E., and Agosta, C.: The future sea-level contribution of the Greenland ice sheet : a multi- model ensemble study of ISMIP6, *The Cryosphere Discussions*, pp. 1–43, <https://doi.org/10.5194/tc-2019-319>, 2020.
- Goldsby, D. L. and Kohlstedt, D. L.: Grain boundary sliding in fine-grained ice I, *Scripta Materialia*, 37, 1399–1406, [https://doi.org/10.1016/S1359-6462\(97\)00246-7](https://doi.org/10.1016/S1359-6462(97)00246-7), 1997.
- 415 Goldsby, D. L. and Kohlstedt, D. L.: Superplastic deformation of ice: Experimental observations, *Journal of Geophysical Research*, 106, 11 017–11 030, <https://doi.org/10.1029/2000JB900336>, <http://doi.wiley.com/10.1029/2000JB900336>, 2001.
- Gow, A. J.: Results of measurements in the 309 meter bore hole at byrd station, antarctica, *Journal of Glaciology*, 4, 1963.
- Greve, R.: Application of a Polythermal Three-Dimensional Ice Sheet Model to the Greenland Ice Sheet: Response to Steady-State and Transient Climate Scenarios, *Journal of Climate*, 10, 901–918, [https://doi.org/10.1175/1520-0442\(1997\)010<0901:AOAPTD>2.0.CO;2](https://doi.org/10.1175/1520-0442(1997)010<0901:AOAPTD>2.0.CO;2), [http://journals.ametsoc.org/doi/abs/10.1175/1520-0442\(1997\)010<0901:AOAPTD>2.0.CO;2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0442(1997)010<0901:AOAPTD>2.0.CO;2), 420 [http://journals.ametsoc.org/doi/abs/10.1175/1520-0442\(1997\)010<0901:AOAPTD>2.0.CO;2](http://journals.ametsoc.org/doi/abs/10.1175/1520-0442(1997)010<0901:AOAPTD>2.0.CO;2), 1997.

- Greve, R. and Blatter, H.: Dynamics of Ice Sheets and Glaciers, *Advances in Geophysical and Environmental Mechanics and Mathematics*, Springer Berlin Heidelberg, Berlin, Heidelberg, <https://doi.org/10.1007/978-3-642-03415-2>, <http://link.springer.com/10.1007/978-3-642-03415-2>, 2009.
- Hansen, B. and Landauer, J. K.: Some results of ice cap drill hole measurements, *IASH Publ.*, 47, 313–317, 1958.
- 425 Hinkel, J., Lincke, D., Vafeidis, A. T., Perrette, M., Nicholls, R. J., Tol, R. S., Marzeion, B., Fettweis, X., Ionescu, C., and Levermann, A.: Coastal flood damage and adaptation costs under 21st century sea-level rise, *Proceedings of the National Academy of Sciences*, 111, 3292–3297, 2014.
- Hobbs, P.: *Ice Physics*, Clarendon Press, Oxford, 1974.
- Holdsworth, G. and Bull, C.: The flow law of cold ice: investigations on Meserve Glacier, Antarctica, *International Symposium on Antarctic Glaciologic Exploration*, 86, 204–216, [https://doi.org/\(Symposium at Hanover 1968 — Antarctic Glaciological Exploration \(ISAGE\)\)](https://doi.org/(Symposium at Hanover 1968 — Antarctic Glaciological Exploration (ISAGE))), 1970.
- 430 Humbert, A., Greve, R., and Hutter, K.: Parameter sensitivity studies for the ice flow of the Ross Ice Shelf, Antarctica, *Journal of Geophysical Research: Earth Surface*, 110, 1–13, <https://doi.org/10.1029/2004JF000170>, 2005.
- Hutter, K.: *Theoretical glaciology material science of ice and the mechanics of glaciers and ice sheets*, 1983.
- 435 Huybrechts, P., Goelzer, H., Janssens, I., Driesschaert, E., Fichet, T., Goosse, H., and Loutre, M.-F.: Response of the Greenland and Antarctic ice sheets to multi-millennial greenhouse warming in the Earth system model of intermediate complexity LOVECLIM, *Surveys in Geophysics*, 32, 397–416, 2011.
- Jellinek, H. H. G. and Brill, R.: Viscoelastic Properties of Ice, *Journal of Applied Physics*, 27, 1198–1209, <https://doi.org/10.1063/1.1722231>, <http://aip.scitation.org/doi/10.1063/1.1722231>, 1956.
- 440 Larour, E. Y., Seroussi, H., Morlighem, M., and Rignot, E.: Continental scale, high order, high spatial resolution, ice sheet modeling using the Ice Sheet System Model (ISSM), *Journal of Geophysical Research: Earth Surface*, 117, <https://doi.org/10.1029/2011JF002140>, <http://doi.wiley.com/10.1029/2011JF002140>, 2012.
- Levermann, A., Winkelmann, R., Nowicki, S., Fastook, J. L., Frieler, K., Greve, R., Hellmer, H. H., Martin, M. A., Meinshausen, M., Mengel, M., et al.: Projecting Antarctic ice discharge using response functions from SeaRISE ice-sheet models, 2014.
- 445 Levermann, A., Winkelmann, R., Albrecht, T., Goelzer, H., Golledge, N. R., Greve, R., Huybrechts, P., Jordan, J., Leguy, G., Martin, D., et al.: Projecting Antarctica’s contribution to future sea level rise from basal ice shelf melt using linear response functions of 16 ice sheet models (LARMIP-2), *Earth System Dynamics*, 11, 35–76, 2020.
- Lipscomb, W. H., Price, S. F., Hoffman, M. J., Leguy, G. R., Bennett, A. R., Bradley, S. L., Evans, K. J., Fyke, J. G., Kennedy, J. H., Perego, M., Ranken, D. M., Sacks, W. J., Salinger, A. G., Vargo, L. J., and Worley, P. H.: Description and Evaluation of the Community
- 450 Ice Sheet Model (CISM) v2.1, *Geoscientific Model Development Discussions*, pp. 1–65, <https://doi.org/10.5194/gmd-2018-151>, <https://www.geosci-model-dev-discuss.net/gmd-2018-151/>, 2018.
- Ma, Y., Gagliardini, O., Ritz, C., Gillet-Chaulet, F., Durand, G., and Montagnat, M.: Enhancement factors for grounded ice and ice-shelf both inferred from an anisotropic ice flow model, *Journal of Glaciology*, 56, 805–812, <https://doi.org/10.3189/002214310794457209>, <http://www.agu.org/pubs/crossref/2011/2011GL048892.shtml>, 2010.
- 455 MacGregor, J. A., Colgan, W. T., Fahnestock, M. A., Morlighem, M., Catania, G. A., Paden, J. D., and Gogineni, S. P.: Holocene deceleration of the Greenland Ice Sheet, *Science*, 351, 590–593, <https://doi.org/10.1126/science.aab1702>, <http://doi.wiley.com/10.1002/2015JF003803><http://www.sciencemag.org/cgi/doi/10.1126/science.aab1702>, 2016.

- Martin, P. J. and Sanderson, T. J. O.: Morphology and Dynamics of Ice Rises, *Journal of Glaciology*, 25, 33–46, <https://doi.org/10.3189/S0022143000010261>, https://www.cambridge.org/core/product/identifier/S0022143000010261/type/journal_article, 1980.
- 460 Mellor, M.: Creep tests on antarctic glacier ice, *Nature*, 184, 1959.
- Mellor, M. and Smith, J. S.: Creep of Snow and Ice, US Army Cold Regions Research and Engineering Laboratory, pp. 843–855, 1967.
- Mellor, M. and Testa, R.: Effect of temperature on the creep of ice, *Journal of Glaciology*, 8, 131–145, 1969.
- Mengel, M., Levermann, A., Frieler, K., Robinson, A., Marzeion, B., and Winkelmann, R.: Future sea level rise constrained by observations and long-term commitment, *Proceedings of the National Academy of Sciences of the United States of America*, 113, 2597–2602, <https://doi.org/10.1073/pnas.1500515113>, 2016.
- 465 Minchew, B. M., Meyer, C. R., Robel, A. A., Gudmundsson, G. H., and Simons, M.: Processes controlling the downstream evolution of ice rheology in glacier shear margins : case study on Rutford Ice Stream , West Antarctica, pp. 1–12, <https://doi.org/10.1017/jog.2018.47>, 2018.
- 470 Mouginit, J., Rignot, E., Björk, A. A., van den Broeke, M., Millan, R., Morlighem, M., Noël, B., Scheuchl, B., Wood, M., and Morlighem, M.: Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, *Proceedings of the National Academy of Sciences*, 116, 201904242, <https://doi.org/10.1073/pnas.1904242116>, <http://www.pnas.org/lookup/doi/10.1073/pnas.1904242116>, 2019.
- Muguruma, J.: Effects of surface condition on the mechanical properties of ice crystals, *J. Phys. D: Appl. Phys.*, 2, 1517, 1969.
- Naruse, R., Okuhira, F., Ohmae, H., Kawada, K., and Nakawo, M.: Closure Rate of a 700 m Deep Bore Hole at Mizuho Station, East Antarctica, *Annals of Glaciology*, 11, 100–103, <https://doi.org/10.3189/S0260305500006406>, https://www.cambridge.org/core/product/identifier/S0260305500006406/type/journal_article, 1988.
- 475 Nias, I. J., Cornford, S. L., and Payne, A. J.: Contrasting the Modelled sensitivity of the Amundsen Sea Embayment ice streams, *Journal of Glaciology*, 62, 552–562, <https://doi.org/10.1017/jog.2016.40>, 2016.
- Nye, J. F.: The Flow Law of Ice from Measurements in Glacier Tunnels, Laboratory Experiments and the Jungfrau-firn Borehole Experiment, *Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 219, 477–489, <https://doi.org/10.1098/rspa.1953.0161>, <http://rspa.royalsocietypublishing.org/cgi/doi/10.1098/rspa.1974.0120http://rspa.royalsocietypublishing.org/cgi/doi/10.1098/rspa.1953.0161>, 1953.
- Nye, J. F.: The distribution of stress and velocity in glaciers and ice-sheets, *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 239, 113–133, <https://doi.org/10.1098/rspa.1957.0026>, <http://www.royalsocietypublishing.org/doi/10.1098/rspa.1957.0026>, 1957.
- 485 Paterson, W. S. B.: Observations on Athabaska Glacier and their relation to the theory of glacier flow, Ph.D. thesis, The University of British Columbia, <https://doi.org/10.14288/1.0085872>, 1962.
- Paterson, W. S. B.: Secondary and Tertiary Creep of Glacier Ice as Measured by Borehole Closure Rates, *Reviews of Geophysics*, 15, 47–55, <https://doi.org/10.1029/RG015i001p00047>, 1977.
- 490 Paterson, W. S. B.: Why ice-age ice is sometimes "soft", *Cold Regions Science and Technology*, 20, 75–98, [https://doi.org/10.1016/0165-232X\(91\)90058-O](https://doi.org/10.1016/0165-232X(91)90058-O), 1991.
- Paterson, W. S. B. and Budd, W. F.: Flow parameters for ice sheet modeling, *Cold Regions Science and Technology*, 6, 175–177, [https://doi.org/10.1016/0165-232X\(82\)90010-6](https://doi.org/10.1016/0165-232X(82)90010-6), 1982.
- Paterson, W. S. B. and Savage, J. C.: Measurements on Athabasca Glacier relating to the flow law of ice, *Journal of Geophysical Research*, 68, 4537–4543, <https://doi.org/10.1029/jz068i015p04537>, 1963.
- 495

- Pattyn, F.: Sea-level response to melting of Antarctic ice shelves on multi-centennial timescales with the fast Elementary Thermomechanical Ice Sheet model (f.ETISH v1.0), *The Cryosphere*, 11, 1851–1878, <https://doi.org/10.5194/tc-11-1851-2017>, <https://www.the-cryosphere.net/11/1851/2017/>, 2017.
- Pattyn, F., Schoof, C., Perichon, L., Hindmarsh, R. C. A., Bueler, E., De Fleurian, B., Durand, G., Gagliardini, O., Gladstone, R., Goldberg, D., Gudmundsson, G. H., Huybrechts, P., Lee, V., Nick, F. M., Payne, A. J., Pollard, D., Rybak, O., Saito, F., and Vieli, A.: Results of the marine ice sheet model intercomparison project, MISMIP, *Cryosphere*, 6, 573–588, <https://doi.org/10.5194/tc-6-573-2012>, 2012.
- Peltier, R., Goldsby, D. L., Kohlstedt, D. L., and Tarasov, L.: Ice-age ice-sheet rheology: Constraints from the Last Glacial Maximum form of the Laurentide ice sheet, *Annals of Glaciology*, 30, 163–176, <https://doi.org/10.3189/172756400781820859>, 2000.
- Pettit, E. C. and Waddington, E. D.: Ice flow at low deviatoric stress, *Journal of Glaciology*, 49, 359–369, <https://doi.org/10.3189/172756503781830584>, 2003.
- Pörtner, H.-O., Roberts, D., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska, E., Mintenbeck, K., Nicolai, M., Okem, A., Petzold, J., Rama, B., and Weyer, N.: The IPCC Special Report on the Ocean and Cryosphere, Tech. rep., 2019.
- Qi, C., Goldsby, D. L., and Prior, D. J.: The down-stress transition from cluster to cone fabrics in experimentally deformed ice, *Earth and Planetary Science Letters*, 471, 136–147, <https://doi.org/10.1016/j.epsl.2017.05.008>, <http://dx.doi.org/10.1016/j.epsl.2017.05.008http://linkinghub.elsevier.com/retrieve/pii/S0012821X17302625>, 2017.
- Quiquet, A., Dumas, C., Ritz, C., Peyaud, V., and Roche, D. M.: The GRISLI ice sheet model (version 2.0): Calibration and validation for multi-millennial changes of the Antarctic ice sheet, *Geoscientific Model Development*, 11, 5003–5025, <https://doi.org/10.5194/gmd-11-5003-2018>, 2018.
- Raraty, L. E. and Tabor, D.: The adhesion and strength properties of ice, *Proceedings of the Royal Society of London. Series A. Mathematical and Physical Sciences*, 245, 184–201, <https://doi.org/10.1098/rspa.1958.0076>, 1958.
- Raymond, C. F.: Inversion of flow measurements for stress and rheological parameters in a valley glacier, *Journal of Glaciology*, 12, 1973.
- Reese, R., Winkelmann, R., and Hilmar Gudmundsson, G.: Grounding-line flux formula applied as a flux condition in numerical simulations fails for buttressed Antarctic ice streams, *Cryosphere*, 12, 3229–3242, <https://doi.org/10.5194/tc-12-3229-2018>, 2018.
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979–2017, *Proceedings of the National Academy of Sciences*, 116, 1095–1103, <https://doi.org/10.1073/pnas.1812883116>, <http://www.pnas.org/lookup/doi/10.1073/pnas.1812883116>, 2019.
- Ritz, C., Fabre, A., and Letréguilly, A.: Sensitivity of a Greenland ice sheet model to ice flow and ablation parameters: Consequences for the evolution through the last climatic cycle, *Climate Dynamics*, 13, 11–24, <https://doi.org/10.1007/s003820050149>, 1997.
- Schoof, C.: Ice sheet grounding line dynamics: Steady states, stability, and hysteresis, *Journal of Geophysical Research: Earth Surface*, 112, 1–19, <https://doi.org/10.1029/2006JF000664>, 2007.
- Schulson, E. M. and Duval, P.: Creep and fracture of ice, 2009.
- Seroussi, H., Morlighem, M., Rignot, E., Khazendar, A., Larour, E., and Mouginot, J.: Dependence of century-scale projections of the Greenland ice sheet on its thermal regime, *Journal of Glaciology*, 59, 1024–1034, <https://doi.org/10.3189/2013JoG13J054>, 2013.
- Seroussi, H., Nowicki, S. M. J., Payne, A. J., Goelzer, H., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Albrecht, T., Asay-Davis, X., Barthel, A., Calov, R., Cullather, R. I., Dumas, C., Gladstone, R., Golledge, N. R., Gregory, J. M., Greve, R., Hattermann, T., Hoffman, M. J., Humbert, A., Huybrechts, P., Jourdain, N. C., Kleiner, T., Larour, E. Y., Leguy, G. R., Lowry, D. P., Little, C. M., Morlighem, M., Pattyn, F., Pelle, T., Price, S. F., Quiquet, A., Reese, R., Schlegel, N. J., Shepherd, A., Simon, E., Smith, R. S., Straneo, F., Sun, S., Trusel, L. D., Van Breedam, J., van de Wal, R. S. W., Winkelmann, R., Zhao, C., Zhang, T., and Zwinger, T.: ISMIP6 Antarctica: a multi-model

- ensemble of the Antarctic ice sheet evolution over the 21st century, *The Cryosphere Discussions*, pp. 1–54, <https://doi.org/10.5194/tc-2019-324>, <https://www.the-cryosphere-discuss.net/tc-2019-324/>, 2020.
- 535 Steinemann, S.: Resultats experimentaux sur la dynamique de la glace et leurs correlations avec le mouvement et la petrographie des glaciers, 1958.
- Strauss, B., Tebaldi, C., Kulp, S., Cutter, S., Emrich, C., Rizza, D., and Yawitz, D.: Maryland and the surging sea: a vulnerability assessment with projections for sea level rise and coastal flood risk., 2015.
- 540 Talalay, P. and Hooke, R. L.: Closure of deep boreholes in ice sheets: a discussion, *Annals of Glaciology*, 47, 125–133, <https://doi.org/10.3189/172756407786857794>, https://www.cambridge.org/core/product/identifier/S0260305500253457/type/journal_article, 2007.
- the PISM authors: PISM, a Parallel Ice Sheet Model, <http://www.pism-docs.org>, 2018.
- Thomas, R. H.: The creep of ice shelves: interpretation of observed behaviour, *Journal of Glaciology*, 12, 55–70, 1973.
- 545 Treverrow, A., Budd, W. F., Jacka, T. H., and Warner, R. C.: The tertiary creep of polycrystalline ice: experimental evidence for stress-dependent levels of strain-rate enhancement, *Journal of Glaciology*, 58, 301–314, <https://doi.org/10.3189/2012JoG11J149>, 2012.
- Weertman, J.: Dislocation Climb Theory of Steady-State Creep, *Transactions of the ASM*, 61, 681–694, 1968.
- Weertman, J.: Creep of ice, in: *Physics and Chemistry of Ice*, edited by Whalley, E., Jones, S. J., and Gold, L., pp. 320–337, Royal Society of Canada, Ottawa, 1973.
- 550 Weis, M., Greve, R., and Hutter, K.: Theory of shallow ice shelves, *Continuum Mechanics and Thermodynamics*, 11, 15–50, <https://doi.org/10.1007/s001610050102>, 1999.
- Winkelmann, R., Martin, M. A., Haseloff, M., Albrecht, T., Bueler, E., Khroulev, C., and Levermann, A.: The Potsdam Parallel Ice Sheet Model (PISM-PIK) – Part 1: Model description, *The Cryosphere*, 5, 715–726, <https://doi.org/10.5194/tc-5-715-2011>, <https://www.the-cryosphere.net/5/715/2011/>, 2011.

Sensitivity of ice flow to uncertainty in flow law parameters in an idealized one-dimensional geometry

Supplementary material

Maria Zeitz^{1,2}, Anders Levermann^{1,2,3}, and Ricarda Winkelmann^{1,2}

¹Potsdam Institute for Climate Impact Research (PIK), Member of the Leibniz Association, P.O. Box 60 12 03, 14412 Potsdam, Germany

²University of Potsdam, Institute of Physics and Astronomy, Karl-Liebknecht-Str. 24-25, 14476 Potsdam, Germany

³LDEO, Columbia University, New York, USA.

Correspondence: Maria Zeitz (maria.zeitz@pik-potsdam.de), Ricarda Winkelmann (ricarda.winkelmann@pik-potsdam.de)




Figure S1. Effect of activation energy parameters on the temperature dependence of the softness A . The temperature dependence of the ice softness A is usually shown in an *Arrhenius plot*, where the softness is shown on a semi-log scale over the inverse temperature. Two parameters for the activation energy Q_c and Q_w for $T \leq -10^\circ\text{C}$ and $T > -10^\circ\text{C}$ parametrize the relationship of ice softness to pressure adjusted temperature. Here the softness is fixed at a reference temperature of $T = -20^\circ\text{C}$. The softness at cold temperatures depends only on the choice of Q_c . Softness at pressure melting point is most sensitive for variations in the Q_w , but varies slightly with Q_c . At pressure melting point the softness increases 8-fold between the limits of parameter combinations, from -47% to + 335% compared to standard parameters. For comparison we show the textbook values of the softness for different temperatures (black crosses) (Cuffey and Paterson, 2010).

References

Cuffey, K. M. and Paterson, W. S. B.: The Physics of glaciers, Elsevier Inc., 4 edn., <http://linkinghub.elsevier.com/retrieve/pii/0016718571900868>, 2010.

Relative difference in equilibrium volume with adapted accumulation




Figure S2. Relative difference in equilibrium volume with adapted accumulation rates compared to the equilibrium state of the simulation with standard parameters and an accumulation rate $a = 0.5$ m/yr.

Step temperature increase $\Delta T = 2^\circ C$, $n = 3$




Figure S3. Effect of activation energy and flow exponent on flow-driven ice discharge : Ice loss in a conceptual flowline setup subject to a temperature anomaly forcing of $\Delta T = 2^\circ C$ and $n = 3$ in percent of the initial volume.




Figure S4. Effect of differences in temperature and different activation energies on ice softness in equilibrium: (a) Pressure adjusted temperature of the ice sheet in the equilibrium state for different combinations of activation energies Q_c, Q_w (see title of each tile) with a flow exponent $n = 3$. Note that the accumulation rates are adjusted to keep the equilibrium volume fixed, with decreased accumulation rates for low activation energies and increased accumulation rates for high accumulation energies. The increase in accumulation leads to cooler temperatures at the top of the ice sheet. (b) Ice softness A in equilibrium for different combinations of activation energies Q_c and Q_w . A clear increase in softness, in particular at the base of the ice sheet is observed for high activation Q_c and Q_w . (c) Ice softness calculated from the pressure adjusted temperatures shown in (a) but fixing Q_w and Q_c at the reference values. The changes in temperature distribution alone are not sufficient to reproduce the softness shown in panel (b). (d) Ice softness calculated from a fixed pressure adjusted ice temperature and different Q_c and Q_w parameters. The change in activation energies alone can reproduce the softness shown in panel (b). Here we show equilibrium values without a warming anomaly. A very similar behavior is observed for the equilibrium states with increased temperatures.