

We are grateful to the anonymous referee, to Alex Robinson and Xavier Fettweis for their thoughtful and constructive comments. We hope that we have improved the paper by addressing their concerns, as outlined below, beginning with major points made by more than one referee.

Threshold

Referee 1 (2) *The authors found no evidence for “warming threshold beyond which the ice sheet would be eliminated”. However, one could argue that this depends on how this “threshold” and “eliminated” are defined. The authors based the claim on that “the ice sheet endures, albeit in a much reduced state”, others could call this elimination. They use also use the claim that there is a large spread on the final ice sheet mass for the same global temperature change, coming from the albedo and boundary (ocean) conditions. However, if I look at figure 2c, I could draw two perpendicular lines intersecting at (2.6 K, 3 m SLE) and claim that all simulations with warming of more than ~ 2.6 K would result in very small ice sheets of less than 3 m sea level equivalent, and that no simulation with less than this warming results in final states of less than 3 m sea level equivalent. Yes, for 2 K of warming the spread in the final volume is very large (from almost no change to more than 12 of the original size), but the spread does not include very small ice sheets. Beyond the 2.6 K “threshold” the spread does not include high volume final ice sheets.*

Referee 2 (1) *Threshold or no threshold. The statements in this paper regarding the existence of a threshold for large-scale melting are framed as rather strongly, in a way that greatly contrasts with the results of Robinson et al. (2012) and previous work. However, I feel that this rather binary framing is not really warranted, nor does it help the community gain clarity on the issue.*

- *I would argue that the experimental setup here simply does not allow such a definitive conclusion to be made. Only 47 experiments are performed with rather arbitrary levels of SST warming applied based on available GCM experiments. This leaves some conclusions open to interpretation. For example, in Fig. 2c, it could be argued that the low-albedo model (red points) shows a roughly linear reduction in equilibrium volume as a function of temperature anomaly, while the high-albedo model (blue points) shows a threshold at $\sim 2^\circ\text{C}$.*
- *[A related later comment] While above 2°C only rather low-volume states appear to be accessible, from ~ 0.5 – 2.0°C , a wide range of intermediate states are accessible. It appears that the low- and medium-albedo model versions particularly allow access to volume distributions between 3–6 m sle. In contrast, the high-albedo model version mainly shows states with a large volume or a much smaller volume. Is there a reason that the high-albedo model may exhibit more threshold-like behavior than the low-albedo model? I think a discussion around this point would be a valuable addition to understanding the physics of the system.*
- *It is clear from this and previous work that strong, positive feedbacks do exist that give the system the potential for self-sustained melting of the ice sheet (albedo, elevation feedbacks). This work shows that there are additional important negative feedbacks (circulation changes leading to increased cloudiness and precipitation) that can serve to counterbalance the positive ones. Given this, I think the binary framing of “threshold or no threshold” is rather misleading.*
- *This work will clearly feed into the upcoming IPCC report. Simply including the headline statement “There is no threshold” implies that these results run completely counter to previous work. But one could also make the following statement: “Above 2°C , all simulations show retreat of the ice sheet to less than half of its current size.” This statement is actually quite consistent with previous results, with the difference being how far the ice sheet retreats.*

For the reasons above, I would suggest a general reframing of the discussion of these results in relation to previous work to highlight the continuity in our growing understanding of this complex system.

We appreciate the points that the referees are making. We agree on the need to be clear that our results are not revolutionary. We find that with small enough warming the ice-sheet is little reduced, while with large enough warming little of it remains, and those statements agree with previous results. However, the temperature interval between “small” and “large” is wider in our results, or alternatively the mass interval between “small” and “large” is narrower. Because of this, we think that our results give a qualitatively different impression from those of Levermann *et al.* (2013, using the model of Robinson *et al.*)—see Figure A below. That is what we were intending to convey by the statement that, “There is no threshold.” We think this is an important point to get across.




Figure A: (i) is Figure 1C of Levermann *et al.* (2013), showing steady-state global-mean sea-level rise due to mass-loss from the Greenland ice-sheet as a function of global-mean surface air temperature change. Each thin green line shows the results from the model for a particular choice of parameters. (ii) is Figure 2c of our manuscript.

Levermann *et al.* describe their results as follows.

The contribution to sea-level commitment from the Greenland Ice Sheet is relatively weak (on average $0.18 \text{ m } ^\circ\text{C}^{-1}$ up to 1°C and $0.34 \text{ m } ^\circ\text{C}^{-1}$ between 2 and 4°C) apart from the abrupt threshold of ice loss between 0.8 and 2.2°C above preindustrial (90% credible interval) (Fig. 1C). This corresponds to a transition from a fully ice-covered Greenland to an essentially ice-free state (*i.e.*, a reduction in ice volume to approximately 10% of the present-day volume, corresponding to a sea-level contribution of more than 6 m).

(Note: their text says “of approximately 10%” in the last sentence—we think they meant “to approximately 10%”, or equivalently, “by approximately 90%”.) It can be seen in Figure A(i) that any given version of their model the temperature interval of the transition is very small—the lines look vertical. By comparison, for the low- and medium-albedo versions of our model, a similar transition between ice-sheet masses of about 6 m and 1 m SLE occurs over about a 2°C temperature interval. As referee 2 says, it might be that the high-albedo version of our model has a transition over less than 0.5°C between about 6 m and 3 m. More experiments are

needed to probe that range. Even if so, that is less of a jump than in the model of Levermann *et al.*

We hope that the referees agree with the above assessment. Accordingly, we have removed the bald statement, “There is no threshold”, and have revised the text of the conclusions, abstract and elsewhere describing the behaviour of our model and its differences from previously published ones. Both referees suggest a statement of the kind, “With warming exceeding X , the steady-state ice-sheet mass is smaller than Y ,” and we have included such a statement in the text and abstract. In view of the comment from referee 1, we now say “eliminated” only when we mean *completely* eliminated.

We have remarked on the need implied by referee 2 to explore the transition more thoroughly with further experiments, on the roles of both positive and negative feedbacks (see also response below concerning his point 2), and on the differences in behaviour between the versions of our model. We suggest that, if indeed the high-albedo version has a more abrupt threshold, it could be because the small ice-cap instability is intensified by positive albedo feedback, which is strongest in this version, but we have not included this speculation in the text.

Low resolution

Referee 1, L479–481 *What about the coarse resolution of the atmospheric model, would you include it as a limitation of this study?*

Referee 2 (2) *The FAMOUS atmosphere is necessarily low-resolution for computational speed, but $7.5^\circ\text{lon} \times 5.5^\circ\text{lat}$ corresponds to roughly 7 grid points east-west and 5 grid points north-south over Greenland. Given that this study highlights the importance of atmospheric circulation changes impacting Greenland stability, such a low atmospheric resolution here seems problematic. Have the authors considered running a short experiment with a higher resolution equivalent of the AGCM with the same boundary forcing and a reduced ice-sheet configuration, but no active ice-sheet model, to see if the atmospheric state is similar to that predicted by the very low-resolution version? Such an experiment, if possible, would go very far towards understanding the possible uncertainties related to these non-linear feedbacks with the atmosphere.*

Referee 3, (3), Lines 314–317: *One of the more important results of this study is the necessity of a full coupling with atmosphere to evaluate tipping point of the Greenland as changes in topography impacts on precipitation and cloudiness as negative feedback. I fully agree with this statement (as we have found the same when MAR has been coupled with GRISLI in Le clec’h *et al.*) but I have some reserve about the robustness of these simulated atmospheric changes in view of the spatial resolution ($7.5^\circ \times 5^\circ$) used by the FAMOUS AOCGM. I understand the use of such a huge resolution in this study but an evaluation of these fields over current climate (by comparison with ERA-Interim) will be very useful to evaluate the ability of FAMOUS to simulate precipitation and cloudiness. At such resolution, how many pixels are there over the ice sheet and what is the topography over current climate. Finally, do changes in the Greenland topography impact on only the local climate over Greenland or the climate at a larger scale? This issue linked to the very low resolution used should at least be mentioned in the conclusion (Lines 478–486).*

As all the referees suggest, we have now remarked in the conclusions that the low resolution of the GCM is a limitation of the model. As referee 2 says, Greenland spans only 7 by 5 grid-boxes in the free atmosphere of FAMOUS, and we have now remarked on this when the model is introduced. We have now also drawn attention to the comparison shown by Smith *et al.* (*Geosci. Model Develop. Discuss.*, <https://doi.org/10.5194/gmd-2020-207>) of the

cumulative distributions of area as a function of altitude (the hypsometry) from FAMOUS and the subgridscale scheme. The SMB simulated with the subgridscale scheme has reasonably realistic features, as discussed in Section 3.3 (now 2.3), where we note that the precipitation gradients are probably too weak because of the low GCM resolution.

The evaluation of SMB and other quantities relevant to SMB for late 20th-century climate (as simulated by MIROC5) gives us some confidence in the simulations for climate change. Smith *et al.* present more information about this comparison. We have not carried out a simulation forced with ERA-interim because our experimental design uses surface BCs directly from the AOGCMs for the late 20th and 21st centuries (rather than, for instance, adding anomalies to observational BCs).

The SMB and AGCM–ice-sheet coupling schemes are technically complicated model developments, and it is not trivial to incorporate them in a different AGCM. Over the past few years, at the same time as the work described in this paper and as part of a larger project involving a team of collaborators, we have implemented these schemes, including downscaling to 1.2 km for the Bisicles ice-dynamical model, in the present version of the UK Earth system model, whose atmosphere resolution is 1.875° longitude \times 1.25° latitude (N96). (This model is computationally about 10,000 times more demanding than FAMOUS–ice.) Our climate-change experiments with UKESM–ice are at an early stage, and unfortunately it would be premature to include any results in the present paper. In a preliminary experiment of 1500 ice-sheet years under abrupt4xCO2 forcing, the Greenland ice-sheet contracts towards a shape like Figure 3c2 of the present paper, and the SMB increases in the central part of it. This is due at least partly to an increase in snowfall of a few 0.1 myr^{-1} LWE because of the altered topography, by comparison (as the referee suggests) with another experiment in which the ice-sheet is not dynamic. We are unsure yet about the effects of cloud changes on SMB in this experiment. Because of the systematic uncertainty in any model investigation, it would clearly be valuable if similar experiments to ours were carried out with other coupled ice-sheet–climate models, as we say in the last paragraph of the paper.

The question of referee 3 about the remote climate effect of changes in Greenland is scientifically interesting. However, we feel it is not within the scope of this paper. We note that Ridley *et al.* (2005, 10.1175/JCLI3482.1) found the remote effects to be small, as simulated by HadCM3. We are grateful to Referee 3 for reminding us of Le clec’h *et al.* (2019), now cited, which is closely relevant to our work.

Referee 1

(1) *I find difficult to navigate the results, e.g., figures (with much dependencies and going forth and back in the text) or trace individually the outcome of the 47 simulations. To this point, I have included some comments regarding figure legends and keys, but if the readers can find further ways to help paper navigation, that would be helpful.*

We have followed the referee’s suggestions, thank you. In revising the text we have added some more cross-references as well, which may help.

(2) *See **Threshold**, above*

(3) *The authors label the negative feedbacks found (cloud, precipitation) as “area feedbacks”, as opposed to positive “thickness feedbacks” (elevation), but I am not very clear about this separation. For instance, the present-day distribution of solar radiation and precipitation is related not only to the margin position but to the surface topography (or thickness, Ettema *et al.*, 2010; also see Figure 3c3,d3 and Figure 4c4,d4 where solar radiation and snowfall change*

within the common area as well). Likewise, the albedo feedback is a positive feedback and the albedo effect from area retreat may be included in its definition (besides the melt effects over the remaining ice sheet).

Yes, that is a good point. We have changed the text to refer to the elevation–SMB feedback included in the interpolation from FAMOUS to Glimmer (and present in the uncoupled model) as the “local lapse-rate feedbacks”, and all the others (due to the response of the FAMOUS climate to the evolving ice-sheet) as “regional climate feedbacks”.

(4) Some of the conclusions . . . come from a empirical relationships inferred from this study’s simulations, and not directly from the simulations. The distinction from inferred conclusions and direct conclusions should be made. For instance, when these relationships are applied to present-day rates (as there are actual simulations under present-day climate, but these are “stable” spin-ups) or to results from AR5 (e.g. projections for RCP8.5).

We think that these concerns arise from our rather inconsistent use of “recent” and “present-day”. In revising the text, we have been consistent in referring to 1980–1999 (whose simulated climate is used for spin-up to steady state) as “late 20th century”, and observations of SMB and ice-sheet mass loss from the last couple of decades as “recent”, while “present-day” is used only for the ice-sheet mass and topography, which changes comparatively slowly. The referee has made some specific comments about this general point at line 466, to which we respond below.

(5) The use of a surface energy balance calculation for melt (as opposed to empirical calculations, e.g. PDDs) should be highlighted more, for instance in the abstract. More detail on such calculation (e.g. energy fluxes, snowpack model, refreezing calculation, albedo calculation) should be given here, instead of only referring to a paper in preparation.

We have inserted remarks comparing with empirical SMB schemes in Section 2 preamble and Section 2.1 and mentioned our approach in the abstract—thanks for the suggestions. We would prefer not to include more detail of the schemes in this paper. Please note that Smith *et al.* is presently under open review at <https://doi.org/10.5194/gmd-2020-207> in *Geosci. Model Develop. Discuss.* It contains information on all the matters of interest, and the referee is welcome to comment if the information given there is insufficient.

(6) Very little attention is given to refreezing in the manuscript. How is refreezing evolving within the simulations? Can you explain how refreezing is calculated? Text in general lacks discussion of effects of refreezing (both in introduction and discussion).

Refreezing is described by Smith *et al.* (see previous comment).

(7) Which kind of vegetation cover is simulated under the retreating ice sheet? How different are the properties of this land cover with respect to a glaciated surface?

We have inserted, “When the ice-sheet retreats, the newly exposed land is assigned the properties of bare soil, including a low snow-free albedo; its properties do not subsequently change because vegetation dynamics are not included in the model.”

(8) The spinup is made under 1980-1999 MIROC (ocean) boundary conditions, but scenario climates from other three models are applied without going through the historical period of the corresponding model. For instance, in Figure 2a, the change in SMB is referred to historical MIROC, but in the text the historical value for a different model is given for this change in SMB. In Figure 3 historical HadCM3 SMB is depicted in column a, while the text in Section 6 indicates that this climate is not used.

We think the comment about Figure 2a refers to the mistake we have corrected following the referee’s comment on line 215. The caption for Figure 3a has been corrected to read “initial

state with HadGEM2-ES historical climate”; the referee is correct that this is not a spun-up steady state of the ice-sheet, although not far from being so. Apologies—we do not understand the comment about Section 6.

Abstract *The abstract is difficult to understand without having read the body text first. Much context is missing on the numerous complex statements. I suggest to give more context and/or reduce/generalize the conclusions.*

As suggested in following comments, we have deleted some sentences which were evidently too complex or needed too much explanation.

L7 *clarify meaning of “initially”, and of “warming”.*

We have deleted the phrase containing “initial”. We think “global warming” is a well-understood phrase.

L7 *A bit of introduction on the simulation design (e.g. “steady state”, use of BC from four different models) could be helpful.*

We believe that “steady state” is a familiar concept. We don’t think there is enough space to say more about the experimental design.

L8 *“for all RCP8.5 climates”—clarify that this means for a steady state climate corresponding to 2081-2100, and not RCP/ECP8.5 up to 2300 as e.g. in Aschwanden et al., 2019, Vizcaino et al., 2015*

We have deleted this sentence, for brevity and because we have inserted, “For warming exceeding 3 K, the contribution to GMSLR exceeds 5 m”, in response to comments by referees 1 and 2.

L9 *“if recent climate were maintained”: this is a conclusion not from the simulations, but from empirical relationships obtained from the simulations.*

Yes, it is an estimate obtained from a fairly good linear fit to the results, so it is in effect an interpolation. We don’t think that is essential to include in the abstract. Also note that we have corrected the range from “1.5–2.5” to “0.5–2.5”.

L11,L12 *“The dominant effect is reduction of area”: effect on what? The statement is very cryptic, why is the area important? “The geographical variation of SMB must be taken into account”. This seems to imply that this is not taken into account in previous work, but one could argue that simulation of ablation and accumulation area is a geographical variation already Please specify further “geographical variation of SMB”*

We have deleted the sentence; these comments indicate that it’s too complicated for the abstract.

L14 *“if late twentieth-century climate is restored . . . the ice sheet will not regrow to its present extent”—this would be straightforward after reading line 9, but this line 14 is based directly on the model simulations, and not on the empirical relationships obtained . . . the mixing of direct and indirect results is a bit confusing.*

We don’t share the referee’s concern about this. As stated above, the estimate of the final mass for present SMB is an interpolation of the model results, which is not really “indirect” in our opinion. The other statements about the final mass simply describe model results.

L15 *“owing to such effects: which effects? The cloud and snowfall effects were just said to increase the SMB, so they would help to re-grow the ice sheet?”*

Thanks for pointing to this possible confusion. The “effects” we meant were regional climate change, but in this case they have the opposite consequence, as the referee says. We have rewritten the sentence, also following a similar comment by referee 2.

L71 “Because of the elevation and albedo feedbacks . . . the present ice-sheet could not be regenerated”—Is a decrease of albedo “irreversible”?

The loss of the ice-sheet might be (at least partly) irreversible, because the albedo and elevation are lower than in the present state, making the SMB more negative. The albedo change itself could be reversed if the ice-sheet readvanced, of course. We have rewritten this sentence as follows and hope this avoids the misunderstanding: “Even after CO₂ fell and global climate returned to pre-industrial, it might not be possible to regenerate the ice-sheet, because of greater ablation or reduced snowfall due to lower elevation and albedo in deglaciated regions.”

L123–124 Muntjwerf et al. (*GRL*, 2020) also use this approach with same number of elevation classes for CESM2.

Thank you for this reference, which we have inserted, along with Lipscomb *et al.* (2013).

L126 Sellevold et al. (*TC*, 2019) discusses the sensitivity of the “tiles” downscaling method to lapse rate choice.

Thanks for this reference, which we have inserted at the point just below where we discuss the lapse rate, noting that 6 K km⁻¹ gives the best SMB gradient in their study.

L129–L120 Which gradients are these, can you specify?

The gradient of downwelling longwave radiation is 3.6 W m⁻² K⁻¹ km⁻¹. The specific humidity gradient is not a constant because of the strong dependence of saturation specific humidity on air temperature. Since these are described by Smith *et al.*, we think that they are adequately described here by our phrase “consistent with the prescribed lapse rate”.

L134, L137 “There is an uncertain parameter . . .”, please specify. Since Smith et al. is in preparation, could you give here more detail on the albedo modeling?

We would rather not repeat too much, and think that for the purpose of this paper the summary here is adequate, since Smith *et al.* is now available online and gives further information.

L151–L157 How is sliding parameterized in the ice sheet model?

We have inserted a comment that we run the model with no sliding.

L163 “we run 10 years”, do you mean, each atmospheric year is used 10 times, then next atmospheric year is run, etc?

Yes, that’s right. We have rewritten this sentence in the hope of making it clearer, thus: “Therefore, after each AGCM year, the ice-sheet model runs for ten years with the resulting SMB field, depending on the assumption that the elevation–SMB feedback will be negligible for changes in topography that occur within that decade, before the AGCM runs again.”

L165 “We have verified”: would it be possible to show some proof of this, e.g. a figure in the supplementary information?

We have added another appendix (now Appendix A) to show both this and the effect of using monthly BCs including interannual variability instead of climatological monthly means.

L176 Title “3.3” can be more precise, maybe specify “(Evaluation of) Simulated 1980–1999 surface mass balance”?

We have made it “Simulated surface mass balance for recent climate”, to be more informative, and to contrast with the following sections on warmer climates. It’s correct that the climate is nominally 1980–1999, but the climate data is from AOGCM simulations, which do not simulate real-world unforced interannual variability, and the FAMOUS–ice SMB is compared with RCM SMB—observational data is not involved.

L193 “Similar” is perhaps subjective. Please discuss similarities/differences.

On further analysis, we found that ELA is generally greater in FAMOUS-ice. Smith *et al.* have a paragraph about this comparison (their p13). We have replaced this sentence with their summary and a reference “The equilibrium line (black contour) is generally a little higher and further inland in FAMOUS-ice (see Smith *et al.* for details).”

Table 1 Last sentence is unclear (20, 30 or 100 years, “second group”?)

We have rewritten this to avoid the unclear mention of “groups” and provide further information, thus, “The RCM time-means use 20 years of data, while we use 100 years for the FAMOUS-ice MIROC5 1980–1999 simulations, which supply our initial steady states, and 30 for other FAMOUS-ice simulations, which are transient states.”

Table 1 The SMB for RCP2.6 is very similar to historical. This makes sense as only the 2080–2099 (steady-state) climate is used, as compared to other SMB estimates with evolving ice sheet topography where the full 21st century climate for RCP2.6 is applied. Maybe good to add some cautionary text to avoid misleading comparisons?

The SMB change under RCP2.6 is quite small because the climate change under that scenario is fairly small. The MIROC5 RCP2.6 results in the table are from the experiments marked as squares near 1.0 K and between -100 and 0 Gt yr^{-1} in Figure 2a; CanESM2 and HadGEM2-ES have similar temperature change for RCP2.6, while NorESM1-M, shown as crosses, has about 0.5 K. For RCP2.6, MIROC5 gives smaller SMB change than the other three AOGCMs, but it is within or just outside the AR5 uncertainty indicated with dashed lines. For RCP8.5, MIROC5 is in the middle of the range (the squares at around 3.5 K).

We chose to report the MIROC5 medium-albedo results for comparison with the 1980–1999 climates in the table, but we agree with the comment that this choice could lead to an inaccurate impression of the effect of climate change, especially for RCP2.6, where the systematic uncertainty is proportionately large. Therefore we have included in the table the average of the results of the four AOGCMs for the medium albedo under each scenario, and stated the AOGCM-average SMB change in the text.

We think that the neglect of topography change during the 21st century is comparatively unimportant. At the end of Section 4.1, we report that the elevation feedback is about 20% of the SMB change in the second century for initial perturbations more negative than -100 Gt yr^{-1} ; for the first century with smaller initial perturbations it is not distinguishable from statistical uncertainty in our results. Edwards *et al.* (2014) give a best estimate of 4.3% by 2100 under larger climate change than RCP2.6. We have remarked on this in the text.

Table 2 Why 402 ppmv for the historical period, isnt it too high? The mean concentration was lower than that in 1980-1999.

The CO_2 concentration is “equivalent CO_2 ”, used to represent all forcings. We treated 1980–1999 as “present day”. The AR5 median assessment of the net anthropogenic ERF in 2011 was 2.3 W m^{-2} , with a likely range of 1.1 – 3.3 W m^{-2} . Because the difference between this and the nominal forcing of 2.6 W m^{-2} under RCP2.6 at 2100 is small compared with the large systematic uncertainty in present-day forcing, because the forcing is anyway much less important than the SST boundary conditions, and because our simulations are intended more as indicative than as realistic scenarios, we decided for simplicity to use the same concentration for historical and RCP2.6 simulations. We have added further comments in the caption of Table 2 about this.

L215 In the legend of Figure 2 it says delta_SMB is referred to steady state under historical MIROC5, but here the HadGEM2-ES value (+307) is given, which one is correct?

Thanks for noticing this. We have modified the text to give the correct ΔSMB of -1066 Gt yr^{-1} (rather than -1063) relative to the MIROC5 historical climate with low albedo (rather than

the HadGEM2-ES historical climate), and removed the SMB of $+307 \text{ Gt yr}^{-1}$ from the text to avoid confusion.

L220 *10% larger—I get 14%, am I missing something? (0.67*1.50/0.88)*

To be precise, it is 13% ($= 797/708$, the numbers shown in Figs 3a4 and 3b4, which we refer to the text), or 10% to one significant figure. We have inserted \sim to indicate the rounding, and also with the 50% in the text sentence (which is 49%, to be more precise), and we have rounded the 88% and 67% for consistency.

L238 *“becomes gradually more positive” → increases*

We think “increases” is ambiguous when discussing a negative number, as it might mean “increases in magnitude”, which is the opposite of “becomes more positive”.

L269 *“there are such states” : Could you add detail on those? It is difficult to map them from Figure 2d to Figure 2b to follow the SMB evolution.*

This comment suggests a simpler way to make the point, which is that all the final steady states have positive SMB and non-zero M , even though many of the trajectories start with negative SMB in Figure 2b. We have changed the text accordingly.

L292 *smaller → lower*

We have inserted “in magnitude” to clarify the meaning.

L299 *“more negative”, “4 times more”—confusing, please give values*

The actual values are all negative and the point is mainly qualitative; we think the confusion is about what a larger negative number means. We have inserted the numbers in parenthesis with the Figure references to clarify the comparisons being made.

L307–308 *precipitation contours are difficult to read or labels are absent*

We have now distinguished the 0.5 and 1.0 contour lines by linestyle.

L368, L371 *“All but one”: are you explaining this “one” somewhere else?*

Yes. These exceptions are discussed in Section 6.3. We have inserted a comment in parenthesis.

L398 *“we suppose this dome might regrow in time”: it seems it does not regrow in 20,000 years, when do you expect it will start to regrow?*

Fair comment. We can't rule it out, and maybe some unforced variability might stimulate it, but we have no evidence that it will regrow, so we have deleted this remark. We have also deleted the similar speculation about the regrowth of the southern dome in the EWN state with low albedo.

L411 *“difference” in what? “infinitesimal”—what does this mean? Please quantify.*

We have rewritten this sentence to make it less abstract.

L435–439 *Why 2,500 years at 0.7 mm/yr and 1,700 at more than double the rate (2 mm/yr)? How have you done these calculations? Which GrIS mass are you considering as present-day mass and as NON steady-state mass?*

We have now stated all the numbers to the nearest 100 years (previously we had rounded the first two to the nearest 500 years). They are $(7.4 - 4.0)/0.7 = 4857 \simeq 4900$, $(7.4 - 5.5)/0.7 = 2714 \simeq 2700$, $(7.4 - 4.0)/2.0 = 1700$ years and $(7.4 - 4.0)/6 = 567 \simeq 600$ years. The north (NON) steady-state mass of 5.5 m SLE is stated in the previous paragraph. The GrIS present-day mass of 7.4 m SLE is stated in Section 1.2.

L452 *“outweighed”: in which sense?*

We don't understand this comment. You could say that the positive feedbacks are overwhelmed by the negative feedbacks. That is the sense, but "overwhelmed" sounds too strong.

L453 "Snowfall": do you mean ice-sheet-integrated?

Yes—clarified.

L466 "If a climate (...)" . There is a jump here. The results now are based on the empirical relationships, but not directly on the simulations here as the ice sheet is relatively stable under 1980–1999 forcing.

Here we refer to the recently observed imbalance, rather than the 1980–1999 steady state. We have rephrased it.

L479–481 See **Low resolution**, above

L485 "sketchy" has a negative meaning in informal American English, maybe replace by other adjective ("gross")

We have deleted the adjective.

Figure 1 White contour in b), is this modelled or observed "ice margin"?

The model ice margin coincides with the observed one; we have inserted an explanation of this in the text (in the first paragraph of Section 2.2, formerly 3.2).

Figure 1 The ELA contour of c) makes a strange shape in the NW, any idea why?

The horizontal SMB gradient is small in this region and the bedrock topography has an inlet, but we do have a clear explanation.

Figure 2 It is difficult to read precipitation from the last row of figures. Only one contour line is labeled, there are very few others, and no line interval is given.

We believe that this comment refers to Figure 3. It is similar to the referee's comment on L307–308. We have now distinguished the 0.5 and 1.0 contour lines by linestyle.

Figure 4 Legend text and keys are confusing. My understanding is that the line color indicates albedo choice. "[Timeseries of Greenland ice-sheet mass with constant climates and FAMOUS-ice albedo] indicated by the line colors according to the line key of (c)" would hint to the colors indicating climates, but it is actually albedo?

This is a misinterpretation of what we intended the sentence to mean. We have broken it into two sentences to avoid the misinterpretation, thus: "Timeseries of Greenland ice-sheet mass with constant climates. The FAMOUS-ice albedo is indicated by the line colors ...".

Figure 4 I would take the line key from (c) and put it in a common space as it applies to all panels. Same for the symbols in (c).

Thanks for the suggestion, now implemented.

Figure 4 The orange dotted line corresponds to offline simulation with low albedo, shouldn't it be red dotted for consistency? Otherwise, add it to the key for lines, with a "Low albedo, offline run"

Yes, it was orange by mistake, and is now red.

Figure 4 Panel b legend: "the circles indicate transient and final sites" is not clear. Having circles in the key is confusing, as only the colors are used, and those correspond to the symbols in the key of (c).

Yes, good point. We have changed the key to the final symbol colours in (b) and hope the new version is clearer.

Referee 2 (Alexander Robinson)

Threshold or no threshold. See *Threshold*, above

Section 2 on “Conceptual basis for the existence of a threshold warming.” *I don’t quite see how this section adds value to the manuscript, as it is currently framed, especially since later it is stated that the “the conceptual basis for its existence is incorrect”. It feels somewhat like a straw-man argument. The simple equations described in Section 2 are useful for conceptualizing the possibility of a runaway feedback leading to the complete melting of an ice sheet. But I think it is by now clear to the community that an ice sheet like Greenland is a large, complex system with processes coupled to atmospheric circulation and a wide range of acting timescales. It is clear, for example, that Δs , A and $f(\Delta M)$ will all change over time. Therefore, I would suggest to the authors that, rather than framing this as the current paradigm that should be rejected, it would be more valuable to highlight, conceptually, what could happen when some of those terms vary (i.e., when A becomes smaller, but Δs increases), or to expand $f(\Delta M)$ into the multiple contributions that may exist (like f_{albedo} , which is most often positive in a warming climate, and $f_{\text{cloudiness}}$, which is found here to be an important negative feedback). Because I think the authors would agree that, ignoring possible climate feedbacks like cloudiness for a moment, as done in the uncoupled experiment, the theoretical basis for a threshold for ice sheet retreat (i.e, the small ice cap instability) still applies here. It is just mitigated by additional feedbacks/factors that are not accounted for in this simple equation. Along the same lines, I don’t think it makes sense to summarize the study of Robinson et al. (2012) at the end of this section as estimating ΔM . In that study, as in this one, a fully coupled climate–ice-sheet system is simulated with a dynamically evolving ice sheet and topography. The results of such experiments allow later comparison with expectations from this conceptual framework, but this equation is not used at all for any quantitative analysis.*

The idea of an abrupt threshold is familiar because it has been demonstrated in previous literature, but it is often not explained how it comes about. The intention of this section is simply to do that. It isn’t intended to be a “straw man” in the sense of misleading anyone, but it is indeed an simplification. We have demoted it to become a subsection (1.3), now entitled “Discussion of the threshold warming”, just after the idea of a threshold is introduced, where the explanation may help most. In revising the text, we have tried to make its intention clear, and have rewritten much of it. We hope that referee 2 finds that the study of Robinson *et al.* is now better represented. We have taken up the implied suggestion to use the same framework in the later discussion (formerly Section 5, now Section 3.5) to interpret the results.

(1) *I am very surprised to see an SIA model applied here, and at only 20km resolution. At a minimum, some justification of this relatively low grid resolution should be made (I would not expect computational cost to be an issue for such a model in this framework). More importantly, the authors should acknowledge and discuss the possible impact of a lack of fast ice dynamics in their simulations. For example, is basal sliding parameterized in some way, or is no basal sliding allowed? If it is not, the model is likely underestimating the dynamic adjustment of the ice sheet to the area retreat, which is an important positive feedback on ice decline on these timescales.*

Computational cost is actually a consideration. The ice-sheet model is 20% is the cost of the coupled model. If we doubled the resolution of Glimmer, that would presumably increase its cost fourfold and double the cost of the coupled model and therefore the wallclock time of these experiments, which took many months to run on the resources available. The model is fast, but these are very long experiments!

There is no basal sliding in the model. We are not simulating ice-streams or rapid ice dynamics. We have inserted comments to this effect in the description of the model. To simulate these

phenomena properly would require much higher resolution in some regions as well as higher-order dynamics, with much greater cost. We agree that omitting rapid ice-sheet dynamics means that the rate of ice-loss will be underestimated, but the aim of our experiments, with their simple scenarios of constant late 21st-century climate, is to investigate the steady state. We have added this explanation in the model description and as a caveat in the conclusions.

(2) See **Low resolution**, above

(3) *It would also be valuable to see a figure showing the forcing applied to the model. For example, how do the present-day SST fields compare to reanalysis or observed SSTs? What are the future patterns of warming? Also, it generally seems that the simulations forced by NorESM1-M stand apart from the other two with lower ice volumes predicted for the same level of global warming. Is this reflected in the SST warming patterns in some way?*

We chose these four AOGCMs because of their previously having been assessed as relatively satisfactory for simulation of Greenland regional climate for our reference period (1980–1999) by Fettweis *et al.* (2013) and van Angelen *et al.* (2013). This rationale and these references are given in the manuscript; we have now drawn attention to them for information about the climate evaluation. We have included further plots of global-mean and Greenland regional mean surface air temperature range (in Appendix C) showing that NorESM1-M has the smallest warming in general.

(4) *Finally, in terms of style, I find that the use of abbreviations for different variables throughout the text makes the manuscript harder to follow. For example, on L30, the phrase “The increase in D is probably the ice-dynamical response ...” would be more straightforward replacing “ D ” with “discharge”. Perhaps the authors could consider only using the variable abbreviations (P , D , R , M , etc.) when the text is related to specific equations that use them, and otherwise use the actual names in sentences. Some other abbreviations could be avoided all together (BCs, GMSLR, etc.).*

Thanks for drawing attention to this. We had used the symbols for terms in the mass balance especially and unnecessarily in the introduction and the two conceptual discussions. We have now replaced them with words except where they are needed in equations, as suggested. However, we have kept M because it is widespread throughout the text (about 30 occurrences), and quite a lot shorter than “[Greenland] ice-sheet mass”, and ΔSAT for change in global-mean surface air temperature. We have also kept “SMB” (over 100 occurrences), “BC” (19 occurrences), “GMSLR” (26 occurrences) and “SLE” (38 occurrences), which are all fairly well-known abbreviations for rather long phrases.

L11-12 *“This is because the dominant effect is reduction of area, not reduction of surface altitude, and the geographical variation of SMB must be taken into account.” This sentence could be more precise. In previous work geographical variation of SMB was also considered, even if in a simpler way. Nor does it seem that the dominant effect is the reduction in area. Rather, it seems that changes in atmospheric circulation act to mitigate the warming via increased cloudiness. The next sentence is already clearer, so I would suggest deleting this one.*

We have deleted it.

L15 *“owing to such effects.” This reference is not very clear, as increased cloudiness and precipitation would, in principle, help the ice sheet regrow. Consider rephrasing, or simply removing.*

Thanks for this point. We agree that it was confusing. We have rewritten the sentence, also following a similar comment by referee 1.

~**L118, Section 3.1** *Other boundary conditions aside from topography and SSTs should be*

explicitly mentioned here. Are greenhouse gas concentrations applied in FAMOUS-ice to be consistent with the applied SST fields, for example?

The BCs are introduced three paragraphs later in the same section (at line 143 of the submitted manuscript). Yes, the radiative forcing is consistent with the climate of the BCs. In the later paragraph we have now mentioned this, and made reference to Table 2 and Section 4.

Fig. 3 *This is an important figure, but feels a bit busy. Perhaps the color bars of each row could be placed vertically on the side? This would clean it up a bit and make more room for the panels themselves.*

We sympathise with this comment, and considered various designs, but couldn't find a better one for fitting this much information into one figure. If it were split into more than one figure, it would be harder to compare the corresponding cases. There is a different colour bar for each row, and a single row isn't high enough for a colour bar on its side. Without the colour bars between the rows, the individual panel could indeed be enlarged, but only in the north-south direction, which would make them look unlike Greenland as we know it.

Fig. 4 *This figure is hard to follow, as there is a lot of information. I would suggest revising colors and symbols to provide more clarity. For example, I think the colors for different forcing scenarios should be substantially different from the colors delineating model versions (low, medium, high albedo).*

As suggested, we have changed the colours for the scenario to be a different set from the colours for the albedo, and hope that this is less confusing. Also, we have adopted the suggestion of Referee 1 to put the key for line colours and symbols outside the panels, since it is common.

L269–270 *A stable but diminished ice sheet is consistent with previous work. Robinson et al. (2012) found that 10% of the ice sheet remained even above the tipping point for large scale melting (see state E3 shown in Fig. 4 of that reference for an approximate picture). It is clear that at some point retreat of the ice sheet to high elevation zones may lead to restabilization. I suggest reframing here.*

We have deleted the sentence here. We have made a remark along these lines in Section 3.5 (formerly Section 5).

L337–345 *Consider reframing title of this section and first paragraph along the lines of earlier comments.*

We have demoted Section 5 to a subsection (now 3.5), entitled “Discussion of reduced steady states”, and rewritten it following the comments by referees 1 and 2 under the heading **Threshold**.

L387–388, L395, etc. *I would remove the terms WOWS and NON, as they do not help comprehension beyond the already defined EWNS terms, and are used rather rarely in any case.*

We have removed “WOWS”, which was referred to only once after its definition. We have replaced “NON” with “no-north”, which may be clearer, and is useful because it stands for a group of three configurations and is referred to eight times.

L411–412 *I would point out that this seems to be an example of a tipping point being activated. This is, of course, not starting from present day, and so is not the same as what has been discussed until now. But it does show that the mechanisms for triggering self-sustained decline are present in the system.*

We have inserted, “Due to these feedbacks, there is more than one steady state for the given BCs.” We would rather not use the phrase “tipping point”, because that's popularly used for

unstable transitions in the opposite direction (as in Section 1.3), so its meaning here might not be clear.

L458 *Greenland or global warming → regional or global warming*

We have put “regional Greenland or global warming”.

Referee 3 (Xavier Fettweis)

(1) *In addition to Fig1, it should be interesting to show the differences/biases with the “reference”. In the legend, is it MAR forced by MIROC5? or ERA?*

We have included difference maps in Figure 1. MAR is forced by MIROC5, so the results of the two models are comparable.

(2), **Lines 183–187:** *The near-surface climate from RCMs forced by a GCM cannot be compared directly with the near-surface climate from the forcing GCM as RCMs simulate their own boundary layer climate and are even able to correct near-surface GCM biases. As FAMOUS–ice is forced by near-surface climate from GCM, it is normal that there are differences with the RCMs simulations. The RCM are only sensitive to the free atmosphere climate from GCMs. Therefore, this section should be a bit rephrased to explain this issue.*

Yes, we agree. We have changed this sentence to say, “A similarly large spread in SMB arises from the choice of Greenland model (FAMOUS–ice, MAR or RACMO), both because they simulate somewhat different regional climate in the free atmosphere and over land when given climate BCs from the same AOGCM, and because they have different SMB schemes.”

(3), **Lines 314–317:** *See **Low resolution**, above*

Large and irreversible future decline of the Greenland ice-sheet

Jonathan M. Gregory^{1,2,*}, Steven E. George^{1,*}, and Robin S. Smith^{1,*}

¹National Centre for Atmospheric Science, University of Reading, UK

²Met Office Hadley Centre, Exeter, UK

*All authors made essential contributions to this work.

Correspondence: Jonathan Gregory (j.m.gregory@reading.ac.uk)

Abstract.

We have studied the evolution of the Greenland ice-sheet under a range of constant climates typical of those projected for the end of the present century, using a dynamical ice-sheet model (Glimmer) coupled to an atmosphere general circulation model (~~FAMOUS-ice FAMOUS-ice~~ AGCM). The ice-sheet surface mass balance (SMB) is simulated ~~by the AGCM,~~
5 ~~including its within the AGCM by a multilayer snow scheme from snowfall and surface energy fluxes, including refreezing~~
~~and~~ dependence on altitude within AGCM gridboxes. Over millennia under ~~a any~~ warmer climate, the ice-sheet reaches a new steady state, whose mass is correlated with the ~~initial perturbation in SMB, and hence with the~~ magnitude of global climate change imposed. ~~If a climate that gives the recently observed SMB were maintained, GMSLR would reach 0.5–2.5 m. For any global warming exceeding 3 K, the contribution to GMSLR exceeds 5 m.~~ For the largest global warming considered (about
10 +5 K), the ~~contribution to global-mean sea-level rise (GMSLR) rate of GMSLR~~ is initially 2.7 mm yr⁻¹, and ~~the ice-sheet is eventually practically eliminated (giving eventually only a small ice-cap endures, resulting in~~ over 7 m of GMSLR). ~~For all RCP8.5 climates, final GMSLR exceeds 4 m. If recent climate were maintained, GMSLR would reach 1.5–2.5 m. Contrary to expectation from earlier work, we find no evidence for a.~~ ~~Our analysis gives a qualitatively different impression from previous work, in that we do not find a sharp~~ threshold warming that divides scenarios in which the ice-sheet suffers little reduction from
15 those in which it is mostly lost. ~~This is because the dominant effect is reduction of area, not reduction of surface altitude, and the geographical variation of SMB must be taken into account.~~ The final steady state is achieved by withdrawal from the coast in some places, and a tendency for increasing SMB due to enhancement of cloudiness and snowfall over the remaining ice-sheet
~~, through by~~ the effects of topographic change on atmospheric circulation, ~~outweighing the tendency for decreasing SMB from the reduction of surface altitude.~~ If late ~~twentieth-century 20th-century~~ climate is restored ~~, after~~ the ice-sheet ~~will not regrow~~
20 ~~to its present extent, owing to such effects, once its~~ mass has fallen below a threshold of about 4 m of sea-level equivalent, ~~it will not regrow to its present extent, because the snowfall in the northern part of the island is reduced once the ice-sheet retreats from there.~~ In that case, about 2 m of GMSLR would become irreversible. In order to avoid this outcome, anthropogenic climate change must be reversed before the ice-sheet has declined to the threshold mass, which would be reached in about 600
25 years at the highest rate of mass-loss within the likely range of the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.

1 Introduction

1.1 Mass-loss from the Greenland ice-sheet in recent decades

During 1961–1990 the Greenland ice-sheet had a roughly constant mass, in which ~~snow-accumulation~~ P ~~snowfall~~ was balanced
30 by the sum of surface ablation R (meaning all processes of mass-loss, predominantly liquid runoff due to melting) and solid discharge D of ice into the sea (forming icebergs). Over the last 30 years both ~~R and D~~ ~~ablation and discharge~~ have increased significantly while ~~P has not~~, giving a current rate of ~~snowfall~~ has not (Shepherd et al., 2012; van den Broeke et al., 2016; Bamber et al., 2018; M
35 ~~. In recent years, the~~ mass-loss ~~of about 250~~ from the Greenland ice-sheet of 239 ± 20 Gt yr⁻¹, ~~or (in 2012–2017, Shepherd et al., 2020)~~
~~, or about~~ 0.7 mm yr⁻¹ sea level equivalent (SLE) (Shepherd et al., 2012; van den Broeke et al., 2016; Bamber et al., 2018; Mouginit et al.,
~~This~~, accounts for about 20% of global-mean sea-level rise (GMSLR) ~~of recent years~~, most of which is due to thermal expansion of seawater (*i.e.* thermosteric) or mass-loss from glaciers.

The increase in ~~D~~ ~~discharge~~ is probably the ice-dynamical response of outlet glaciers to reduced buttressing by their ice-tongues, which have thinned due to basal melting by warmer sea-water (Holland et al., 2008). ~~Although ice discharge is projected to increase in coming decades with rising water temperature, it will decline on longer timescales as the ice-sheet~~
40 ~~thins at the coast and its outlet glacier termini retreat inland~~ (Nick et al., 2013; Fürst et al., 2015; Aschwanden et al., 2019).

The increase in ~~R~~ ~~ablation~~ causes 60% of the mass-loss (van den Broeke et al., 2016; Fettweis et al., 2017). It has been partly due to anthropogenic climatic warming, which is amplified at high northern latitudes, and partly to recent unusual atmospheric circulation (Tedesco et al., 2013; Fettweis et al., 2017; Pattyn et al., 2018; Trusel et al., 2018). ~~The~~ ~~In recent years, the~~ surface mass balance (~~SMB~~ $S = P - R$, ~~the net addition of mass~~) ~~in recent years where P is snowfall and R ablation~~, has fallen lower
45 than during the warm period in Greenland in the early ~~twentieth~~ ~~20th~~ century (Fettweis et al., 2017) and summer temperatures have risen higher (Hanna et al., 2012). Some recent summers have seen surface melting over practically the entire ice-sheet because of high air temperature, decreased cloudiness and reduction of albedo, the latter due to the increase of snow grain size and the exposure of bare ice, both caused by surface snow melting (Tedesco et al., 2013, 2016; Hofer et al., 2017; Trusel et al., 2018).

50 1.2 Projections of future mass-loss

The future of the Greenland ice-sheet is one of the large uncertainties in projections of GMSLR (Church et al., 2013; Clark et al., 2016). ~~Ice discharge is projected to increase in coming decades with rising water temperature, but it will decline on longer timescales as the ice-sheet thins at the coast and its outlet glacier termini retreat inland~~ (Nick et al., 2013; Fürst et al., 2015; Aschwanden et al.
55 ~~. On multicentennial timescales, SMB is dominant and the source of greater uncertainty~~ (Fürst et al., 2015).

Projections indicate that ~~R~~ ~~ablation~~ will increase non-linearly with temperature and more rapidly than ~~P~~ ~~snowfall~~, meaning that SMB will continue to decline and the rate of mass-loss will grow, especially under scenarios of high CO₂ emissions

(Gregory and Huybrechts, 2006; Fettweis et al., 2013; Vizcaíno et al., 2014; Pattyn et al., 2018; Rückamp et al., 2018; Golledge et al., 2019; Aschwanden et al., 2019). Recent projections of the contribution of the Greenland ice-sheet to GMSLR mostly lie within the likely ranges of the Fifth Assessment Report (AR5) of the Intergovernmental Panel on Climate Change (Church et al., 2013) viz. 0.04–0.12 m and 0.09–0.28 m by 2100 relative to 1986–2005 under scenarios RCP2.6 and RCP8.5 respectively. The range of uncertainty arises from the ~~model-spread-in-spread in global warming simulated by atmosphere–ocean general circulation models (AOGCMs) and in their~~ amplification of warming in Greenland relative to global warming, as well as the sensitivity of Greenland SMB to regional climate change (Gregory and Huybrechts, 2006; Fettweis et al., 2013).

Although substantial, ~~these contributions are~~ the contribution from the Greenland ice-sheet is only 10–30% of projected GMSLR.

~~The importance of the Greenland ice-sheet by 2100. Its importance~~ is greater on multicentury timescales, because its size (mass $M = 7.4$ m SLE) implies a large commitment to GMSLR. Thinning of the ice-sheet due to ~~positive ΔR (Δ denoting the difference from the initial state)~~ increasing ablation is affected by a positive feedback loop between SMB and elevation: as the surface elevation falls, the surface air temperature rises, and surface melting increases, magnifying ~~ΔR~~ the ablation increase. We refer to this as the local lapse-rate feedback. Another positive feedback on ~~ΔR~~ ablation is caused by the decrease in surface albedo due to melting, as in recent years (Tedesco et al., 2016). Despite ~~the~~ these feedbacks, a steady state could be regained with an ice-sheet of smaller mass but little loss of area if the reduction of SMB were compensated by the reduction of discharge resulting from thinning of outlet glaciers (Rückamp et al., 2018) (see Section 1.3). The reduction of mass would be mitigated if ~~$\Delta P > 0$, as projected by GCMs~~ snowfall increases, which is projected by AOGCMs.

On the other hand, previous work indicates there may be a threshold T_c of global-mean surface air temperature change ΔSAT (relative to pre-industrial) beyond which the ice-sheet will ~~vanish~~ be greatly reduced or vanish entirely (Huybrechts et al., 1991; Gregory et al., 2004; Gregory and Huybrechts, 2006; Robinson et al., 2012) (see Section 1.3). Levermann et al. (2013) estimated $T_c = 0.8$ – 2.2 K, using the model of Robinson et al. (2012) constrained by information from the last interglacial (Robinson et al., 2011). If this range is correct, limiting ΔSAT to 2.0 K in accordance with the Paris agreement or to its aspiration of 1.5 K could make a critical difference to whether T_c is exceeded (Pattyn et al., 2018). Loss of the Greenland ice-sheet would cause much greater GMSLR than from glacier mass-loss or thermosteric sea-level rise for similar degrees of warming (Church et al., 2013; Levermann et al., 2013) although, even in the most extreme scenarios, the complete removal of the ice-sheet would take at least a thousand years (*e.g.* Ridley et al., 2005; Aschwanden et al., 2019).

1.3 ~~Possibility-Discussion of irreversible mass-loss~~ the threshold warming

~~Because of the elevation and albedo feedbacks, SMB might remain negative in some deglaciated regions after CO_2~~ The rate of change of the mass of the ice-sheet is $dM/dt = S - D$, where D is discharge. In the unperturbed steady state $dM/dt = 0 \Rightarrow S = D$ *i.e.* fell and the climate cooled, meaning that the present SMB is balanced by discharge. In a warmer climate, ablation R and snowfall P both increase, but $\Delta R > \Delta P \Rightarrow \Delta S = \Delta P - \Delta R < 0$ (see references in Section 1.2), where Δ denotes the difference from the initial state. A new steady state can be achieved if $\Delta D = \Delta S$ *i.e.* if discharge reduces by as much as SMB, so that $D + \Delta D = S + \Delta S$.

Let us suppose that raising the global-mean SAT by T initially perturbs the SMB by an amount $\Delta S_T(T) < 0$. Further suppose that a new steady state can be achieved, with little change in ice-sheet ~~could not be regenerated in~~ area, in which discharge is reduced by marginal thinning, such that $\Delta D = \Delta S = \Delta S_T(T)$. The larger T , the greater the reduction in discharge needed to balance $\Delta S_T(T)$. The threshold $T = T_c$ is reached when there is just sufficient marginal thinning to cause all outlet glaciers to retreat from the coast, reducing discharge $D + \Delta D$ to zero. To attain a balance, SMB must also fall to zero, with $\Delta S = -S$. Hence $\Delta S_T(T_c) = -S$ defines T_c .

Any T exceeding T_c will give negative SMB, but discharge cannot be further reduced (*i.e.* below zero) to compensate. The unbalanced negative SMB will reduce the thickness of the ice-sheet and the altitude of the surface, making the SMB even more negative by the local lapse-rate feedback. If no other process is involved, the ice-sheet will be completely eliminated for any $T > T_c$ by this feedback loop, which is called the “small ice-cap instability”. The threshold has been estimated as $T_c = 1.9\text{--}4.5$ K relative to pre-industrial climate (Toniazzo et al., 2004), and may actually be a relict of a colder climate (Solgaard et al., 2013). This implies the existence of two steady states of (Gregory and Huybrechts, 2006; Meehl et al., 2007), by evaluating the warming required to reduce SMB to zero with the present-day surface topography. The same method gave $T_c = 2.1\text{--}4.1$ K in the AR5 (Church et al., 2013, Section 13.4.3.3).

Robinson et al. (2012) showed that this method of calculation overestimates the actual T_c for onset of the small ice-cap instability. One possible contribution to the difference is the reduction in SMB, neglected above, due to the local lapse-rate effect of the marginal thinning *before* the threshold is reached. Let us write this contribution to ΔS as ΔS_L . For steady state at the threshold, we now require $\Delta S_T(T_c) + \Delta S_L = -S \Rightarrow \Delta S_T(T_c) = -(S + \Delta S_L)$. Since $\Delta S_L < 0$, the right-hand side is less negative than before, so T_c is smaller.

An ice-sheet model is required to allow for ΔS_L in quantifying T_c , because the change in ice-sheet topography is determined simultaneously by SMB change and ice-dynamical change. With their model, Robinson et al. (2012) demonstrated that SMB may initially be positive but decline to zero as the topography changes, whereupon the instability is triggered, leading to the eventual loss of the ice-sheet. They determined their lower T_c by finding the final steady state for various T and versions of their coupled climate–ice-sheet model, and our approach is similar. The coupled system simulated by their and our models is considerably more complicated than this simplified conceptual treatment, which is intended to illustrate the idea. The actual outcome is affected by further feedbacks, both positive and negative, which we discuss later.

1.4 Possibility of irreversible mass-loss

If the ice-sheet ~~(either absent, or present as now)~~ in were removed then, even after CO_2 fell and global climate returned to pre-industrial climate, it might not be possible to regenerate it, because of greater ablation or reduced snowfall due to lower elevation and albedo in deglaciated regions (Toniazzo et al., 2004). If the ice-sheet did not regrow, it would imply that its pre-industrial steady state is a relict of a colder climate (Solgaard et al., 2013). Previous work shows there may be more than two steady states for pre-industrial climate (Charbit et al., 2008; Ridley et al., 2010; Solgaard and Langen, 2012; Robinson et al., 2012). Stable states of intermediate size (between zero and present-day) are possible because of the interaction of the ice-sheet with

its own climate through atmospheric dynamics, whereby its surface topography affects regional precipitation and temperature,
125 like mountains do. The existence of intermediate states means that partial loss of the ice-sheet could be irreversible.

It is the possibility of threshold behaviour (*i.e.* “tipping-points”) and irreversibility which makes the future of the Greenland ice-sheet of particular concern (Pattyn et al., 2018). Precautionary action to mitigate the threat of irreversible damage is a principle of the Framework Convention of Climate Change (Article 3.3), even when there is not full scientific certainty. The serious implications of the uncertainty are the motivation for the work presented in this paper, in which we reexamine the
130 future decline and possible recovery of the ice-sheet. Our conclusions differ in some critical ways from those of previous work, because of the greater complexity of the model, which we next describe.

2 Model

Previous work on the subject has used simplified climate models, or a small set of climate states, or been limited to a few centuries into the future. In the present work we use a dynamic ice-sheet model coupled to a atmosphere general circulation
135 model (AGCM) to study the transient and steady states of the ice-sheet over tens of millennia. ~~Our conclusions differ in some critical ways from those of previous work.~~

3 ~~Conceptual basis for the existence of a threshold-warming~~

~~The rate of change of the mass of the ice-sheet is $dM/dt = S - D$. In the unperturbed steady state $dM/dt = 0 \Rightarrow S = D$ *i.e.* SMB S is balanced by discharge D . In a warmer climate, ablation R and snowfall P both increase, but $\Delta R > \Delta P \Rightarrow \Delta S = \Delta P - \Delta R < 0$.
140 A new steady state can be achieved if $\Delta D = \Delta S$ *i.e.* if discharge reduces by as much as SMB.~~

~~Let us suppose that changing the global-mean SAT by T initially perturbs the ice-sheet area-mean specific SMB s by an amount $\Delta s(T)$. Specific SMB is the excess of snowfall over ablation at a location, and its area-integral is the quantity we usually call just “SMB” *i.e.* $S = As$, where A is ice-sheet area. The initial change in SMB is $\Delta S = A\Delta s(T)$. Further suppose that a new steady state can be achieved by marginal thinning of the ice-sheet, which will reduce D with little
145 change in A . However the elevation and albedo feedbacks may amplify the (negative) change in specific SMB, and require $\Delta D = \Delta S = A\Delta s(T)(1 + f(\Delta M))$ in the reduced steady state, where f is a function representing the feedbacks, with $f(0) = 0$ initially, and f increasing as ΔM becomes more negative.~~

~~This argument leads to the idea of a threshold T_c that gives a steady state with a loss ΔM_c of mass, such that $\Delta S = A\Delta s(T_c)(1 + f(\Delta M_c)) =$ in which SMB and D are both reduced to zero ($S + \Delta S = 0$, with $S = D$ and $\Delta S = \Delta D$, so $D + \Delta D = 0$), the latter achieved by
150 just sufficient marginal thinning to cause all outlet glaciers to retreat from the coast. Any T exceeding T_c will give negative SMB, but D cannot be further reduced (*i.e.* below zero) to compensate. Hence $T > T_c$ is expected to lead to the elimination of the ice-sheet by the so-called small ice-cap instability, wherein s must become increasingly negative as the surface gets lower. Therefore T_c is the greatest global warming that the ice-sheet can endure.~~

155 The threshold was estimated as 1.9–4.5 K relative to pre-industrial (Gregory and Huybrechts, 2006; Meehl et al., 2007), by assuming it equals the warming required to reduce SMB to zero with the present-day surface topography *i.e.* $\Delta S = A \Delta s(T_c) = -S$. This method overestimates T_c by neglecting the feedbacks represented by f . Although S may remain positive after the initial perturbation, it may become negative due to the feedbacks before a steady state is reached. To take $f(\Delta M)$ into account, ΔM must be predicted, for which Robinson et al. (2012) used a dynamical ice-sheet model, thus arriving at their smaller T_c .

3 Model

160 Typical AGCMs are not suitable for modelling ice-sheet SMB, because they do not have adequate treatments of albedo and hydrology, nor fine enough spatial resolution for the large gradients in topography and climate parameters across the margins of the ice-sheets, where much of the snowfall and snowmelt occurs (Vizcaíno, 2014). Specially developed regional climate models (RCMs) have proven very useful for high-resolution projections and process studies (*e.g.* MAR RCM, Fettweis et al., 2013; RACMO RCM, Noël et al., 2018) but they require lateral boundary conditions (BCs) from global GCMs/AGCMs, and cannot feed back on climate change outside their domain. Moreover, computational expense prevents the use of these RCMs in 165 studying ice-sheet evolution over millennia. The first such experiment, with MAR coupled to an ice-sheet model, was only 150 years long (Le clec’h et al., 2019). In multimillennial studies, empirical parametrisations for SMB as a function of surface air temperature (*e.g.* Reeh, 1989), precipitation etc. have often been applied. Being calibrated for observed climate, such schemes may be less reliable for simulations of very different climates of the future or past, and when used in coupling to an AGCM they imply surface energy and water fluxes which are unrelated to those within the AGCM, thus violating conservation.

2.1 FAMOUS-ice AGCM

For sufficient speed, we use the FAMOUS AGCM, which is the atmosphere component of the FAMOUS ~~atmosphere-ocean general circulation model (AOGCM)~~ AOGCM (Smith et al., 2008; Smith, 2012), itself a low-resolution version, at 7.5° longitude by 5° latitude, of the HadCM3 AOGCM (Gordon et al., 2000). ~~To simulate~~ For physical consistency, we calculate the SMB in the AGCM, but Greenland spans only seven gridboxes in longitude and five in latitude in the free atmosphere of the model, which is far from adequate for simulating the important effects of topographic gradients and snow hydrology for ice-sheets. ~~Therefore~~ Therefore in this work we use “FAMOUS-ice”, a new version of FAMOUS (~~version sgfjb, ?~~) (~~version xotzb, Smith et al., submitted~~), incorporating a multilayer surface snow scheme which calculates melting, refreezing of meltwater, runoff and SMB on “tiles” at a set of elevations within each AGCM gridbox (each tile covering a fraction of the gridbox area). This is similar to the ~~approach of Vizcaíno et al. (2013), method implemented for the Greenland ice-sheet in the Community Earth System Model (CESM) (Vizcaíno et al., 2013; Lipscomb et al., 2013; Muntjewerf et al., 2020) and we use the same ten elevations as them.~~ Smith et al. show the improvement in the cumulative distribution of area as a function of altitude (the hypsometry) that results from the subgridscale treatment. Below we summarise the FAMOUS-ice SMB and coupling schemes, of which further details are given by Smith et al.

185 For vertical interpolation of atmospheric variables from the AGCM gridbox elevation to the tile elevations we prescribe a lapse rate of 6 K km^{-1} for air temperature. This we obtained from the climate of 1980–1999 simulated by Fettweis et al. (2013) with the MAR RCM using sea-surface BCs (sea-surface temperature and sea-ice) from MIROC5, the AOGCM which Fettweis et al. found to give the most satisfactory SMB simulation. (~~The same uniform lapse rate is used e.g. by Aschwenden et al., 2019~~ →) Downwelling longwave radiation and specific humidity are vertically interpolated in FAMOUS–ice using gradients consistent with the prescribed lapse rate, but precipitation is not redistributed vertically, nor modified in phase. The same uniform air temperature lapse rate for Greenland is used e.g. by Aschwenden et al., 2019, and found by Sellevold et al. (2019) to give the most similar SMB gradient to RACMO in their CESM ice-sheet coupling, which, like our scheme, does not downscale precipitation.

We have paid particular attention to the treatment of the surface albedo of the Greenland ice-sheet, to which SMB is very sensitive. Bare ice has lower albedo than snow in FAMOUS–ice and snow albedo has different values for visible and near-infrared, both dependent on the snow-grain size, which is a prognostic that depends on the “ageing” of the surface snow by melting and refreezing following new snowfall. There is an uncertain parameter in the relationship between snow-grain size and albedo. In our experiments, we use three alternative parameter values that are consistent with observations of albedo. For convenience we refer to these as low, medium and high albedo, but the reader should keep in mind that the albedo is variable in each case. More details are given by ~~?~~Smith et al. (submitted).

Instead of simulating sea surface conditions by using the FAMOUS AOGCM, we use the AGCM alone, for both recent and future climate, with sea-surface BCs derived from AOGCM experiments of the Coupled Model Intercomparison Project Phase 5 (CMIP5) ~~for both recent and future climate, and atmospheric CO₂ concentration to give the corresponding radiative forcing (see Table 2 and the start of Section 3)~~. We use the AGCM for two reasons. First, the FAMOUS AOGCM has larger biases in its simulation of ~~the present-day-recent~~ climate than MIROC5 and the three other AOGCMs we use (CanESM2, HadGEM2-ES and NorESM1-M), which have all previously been selected as satisfactory for Greenland regional climate simulation (~~Fettweis et al., 2013; van Angelen et al., 2013~~)(see Fettweis et al., 2013, and van Angelen et al., 2013, for evaluation of their regional climate simulations). Second, this method allows us to investigate the uncertainty in Greenland ice-sheet projections that arises from the spread of climate projections given by AOGCMs for any given scenario. The AGCM sea-surface BCs are 20-year climatological monthly means, which lack interannual variability; we have checked that statistically indistinguishable results for the ice-sheet are obtained with the AGCM cycling through a 20-year series of monthly mean BCs for the same climate (Figure S1a).

By prescribing sea surface conditions, we exclude any climate interaction between the ice-sheet and the ocean, in particular, possible cooling of regional climate due to weakening of the AMOC caused by meltwater from the ice-sheet (*e.g.* Vizcaíno et al., 2010). There is wide uncertainty in this aspect of ocean climate change, whose implications for the ice-sheet could possibly be explored in further work by modifying the sea-surface temperatures in a range of ways to represent the effects of AMOC changes projected by AOGCMs (Stouffer et al., 2006; Gregory et al., 2016).

Figure 1. (a,b) ~~Comparison of~~ Greenland surface elevation above sea level in (a) FAMOUS-ice (medium albedo) and (b) observations (Bamber et al., 2001a, b). The white contour is the observed ice margin, the same in both maps. (c) Difference between (a) and (b), positive means FAMOUS-ice surface is higher. (d,e) ~~Comparison of specific~~ Specific surface mass balance (expressed as liquid water equivalent) for the climate of MIROC5 1980–1999 in (ed) FAMOUS-ice (medium albedo), (de) MAR. The black contour is the equilibrium line (where specific SMB is zero). (f) Difference between (d) and (e), positive means FAMOUS-ice SMB is more positive.

2.2 FiG coupling and spinup

We use the Glimmer-CISM community ice-sheet model (Rutt et al., 2009, <https://cism.github.io>) with the shallow-
220 ice approximation at 20 km grid-spacing ~~and no basal sliding.~~ Consequently the model does not simulate ice-streams or rapid ice-sheet dynamics, and it will inevitably underestimate the rate of ice-sheet mass-loss, especially in coming decades. This is acceptable because our aim is not to make realistic time-dependent projections, but to study the steady state obtained under constant climates.

Because the ice-sheet model lacks sufficient resolution and physical processes to simulate calving into fjords, we instantly
225 remove ice which flows beyond the present margin of the ice-sheet. This BC prevents a tendency for the ice-sheet to expand slightly, and ~~it~~ thus it makes the modelled ice edge coincide with the observed one. It becomes irrelevant in most of our experiments, when the ice-sheet contracts. For simplicity in the model we omit isostatic uplift, which in reality gives a negative feedback on ice-sheet mass-loss through the ~~elevation-SMB~~ local lapse-rate feedback, because it is not a large effect (*e.g.* 2% over 1000 years, Aschwanden et al., 2019) and does not seem necessary given that our scenarios are idealised in other ways as
230 well.

The AGCM and the ice-sheet model are coupled to make FAMOUS-ice-Glimmer (~~FiG, Gregory et al., 2012; Roberts et al., 2014; ?~~)
(FiG, Gregory et al., 2012; Roberts et al., 2014; Smith et al., submitted). After each AGCM year, the SMB simulated by the AGCM is interpolated horizontally and vertically (with the AGCM tile elevation as the vertical coordinate) to the ice-sheet surface topography, and the AGCM topography and land-surface properties are updated according to the ice-sheet model. When
235 the ice-sheet retreats, the newly exposed land is assigned the properties of bare soil, including a low snow-free albedo; its properties do not subsequently change because vegetation dynamics are not included in the model.

FiG runs at about 220 simulated AGCM years per wallclock day on six cores, with the AGCM consuming the great majority of the CPU time. Although this is fast for ~~a GCM~~ an AGCM, it is not fast enough for multimillennial experiments, ~~so we run 10 years of the ice-sheet model using.~~ Therefore, after each AGCM year's SMB, the ice-sheet model runs for ten years with
240 the resulting SMB field, depending on the assumption that the ~~elevation-SMB~~ local lapse-rate feedback will be negligible for changes in topography that occur within ~~a decade~~ that decade, before the AGCM runs again. We have verified that this 10:1 acceleration makes no significant difference to our results ([Figure S1a](#)). Hereafter by “year” in FiG experiments we mean an ice-sheet year except where otherwise stated.

Because our aim is to simulate ice-sheet response to climate change over millennia, we have to start from a coupled steady
245 state, with little long-term tendency in the ice-sheet topography. We initiate the ice-sheet model with observed topography

(Bamber et al., 2001a, b) and run FiG under the MIROC5 AOGCM climate of 1980–1999, during which period the ice-sheet was near to a steady state in reality (van den Broeke et al., 2016). In the first millennium the ice-sheet mass M increases by 0.1–0.2 m SLE. With medium and low albedo it subsequently decreases again more slowly, while with high albedo it continues to grow slowly and stabilises after 4 kyr at 0.3 m SLE above present-day (Figure ~~??~~S1b). The states obtained after
250 about 4 kyr of spin-up are used to initiate the experiments described in Section 3. In these states M is close to reality, and the topography similar to observed (Figure 1a,b,c), with the summit and southern dome altitudes being a few 100 m too low. (See also Appendix B concerning the constraint implied on albedo by the requirement of realistic M .)

2.3 Simulated surface mass balance for recent climate

Comparing the three choices of albedo in FAMOUS–ice with BCs for the MIROC5 1980–1999 climate, we find that lower
255 albedo produces lower SMB (the first group of cases in Table 1 differ significantly at the 10% level), because ablation is greater due to greater snowmelt, but snowfall is about the same (slightly larger with higher albedo because of greater ice-sheet area). For the same albedo (medium), the SMB is significantly lower with the CanESM2 and NorESM1-M historical climates than with MIROC5 because the ablation is larger, whereas the SMB is about the same with HadGEM2-ES as with MIROC5 (the second group in the table). This shows the influence of the different climate simulations of the AOGCMs.

260 ~~However, comparison of~~ A similarly large spread in SMB arises from the choice of Greenland model (FAMOUS–ice with MAR and RACMO for the same climates shows that the formulation of the regional climate model is at least as influential as the choice of climate BCs. MAR gives, MAR or RACMO), both because they simulate somewhat different regional climate in the free atmosphere and over land when given climate BCs from the same AOGCM, and because they have different SMB schemes. MAR has much larger SMB than FAMOUS–ice with the MIROC5 climate, because of smaller ablation, while RACMO ~~gives~~
265 has larger ablation than FAMOUS–ice with the HadGEM2-ES climate (the third group in the table). Comparison with MAR and RACMO for ERA-interim BCs (*i.e.* observationally derived, the fourth group in the table) suggests that FAMOUS–ice with high albedo is similar to both of them.

Regarding its geographical distribution, FAMOUS–ice SMB interpolated to the Glimmer grid compares favourably with the MAR simulation for the MIROC5 climate (Figure ~~1e,d,d,e,f~~). It shows positive and negative values of realistic magnitude, and
270 reproduces the important geographical features, including the confinement of negative SMB to the margins, especially on the west coast, the decrease in positive SMB towards the north-east, and the occurrence of greatest positive SMB in the strip of maximum snowfall along the south-east coast. We presume that the latter is not sufficiently intense in FAMOUS–ice because of the low resolution of the AGCM. The equilibrium line ~~altitude is similar in the two models~~ (black contour) is generally a little higher and further inland in FAMOUS–ice (see Smith et al., submitted, for details).

275 3 Mass-loss of the ice-sheet in warmer climates

We run a set of 47 FiG experiments to study the SMB change (Δ SMB), rate of mass-loss and eventual steady state of the Greenland ice-sheet, using the three different choices of FAMOUS–ice snow-albedo parameter, with 20-year climatological monthly

	Climate	Greenland model (albedo)	SMB	Snowfall	Ablation
1980-1999 climates					
<u>1</u>	MIROC5	FAMOUS-ice (low)	310 ± 10	693	383
	MIROC5	FAMOUS-ice (medium)	332 ± 11	697	364
	MIROC5	FAMOUS-ice (high)	414 ± 9	715	300
<u>2</u>	CanESM2	FAMOUS-ice (medium)	272 ± 21	681	409
	HadGEM2-ES	FAMOUS-ice (medium)	312 ± 20	705	393
	NorESM1-M	FAMOUS-ice (medium)	287 ± 16	721	434
<u>3</u>	MIROC5	MAR	437 ± 24	681	244
	CanESM2	MAR	410 ± 23	635	225
	HadGEM2-ES	RACMO	244 ± 25	660	416
	NorESM1-M	MAR	483 ± 16	691	208
<u>4</u>	ERA-interim	MAR	388 ± 23	637	249
	ERA-interim	RACMO	406 ± 22	683	277
MIROC5 2080-2099 climates					
<u>5</u>	RCP2.6	FAMOUS-ice (medium)	325 ± 14	704	379
	RCP4.5	FAMOUS-ice (medium)	150 ± 25	735	585
	RCP8.5	FAMOUS-ice (medium)	-207 ± 35	805	1013
Mean over AOGCM climates					
<u>6</u>	<u>1980-1999</u>	<u>FAMOUS-ice (medium)</u>	<u>307</u>	<u>703</u>	<u>395</u>
	<u>2080-2099 RCP2.6</u>	<u>FAMOUS-ice (medium)</u>	<u>212</u>	<u>746</u>	<u>533</u>
	<u>2080-2099 RCP4.5</u>	<u>FAMOUS-ice (medium)</u>	<u>60</u>	<u>777</u>	<u>716</u>
	<u>2080-2099 RCP8.5</u>	<u>FAMOUS-ice (medium)</u>	<u>-273</u>	<u>825</u>	<u>1098</u>

Table 1. Greenland area-integral surface mass balance (SMB), snowfall and ablation (all in Gtyr^{-1}) for FAMOUS-ice with MIROC5 AOGCM historical climate (with the three choices of FAMOUS-ice albedo), FAMOUS-ice with historical climates of other AOGCMs (FAMOUS-ice medium albedo only), the MAR and RACMO RCMs with the same AOGCM climates and with ERA-interim climate (from Table 2 of Fettweis et al., 2013), and FAMOUS-ice with MIROC5 AOGCM climate (medium albedo only) under RCP scenarios, and the FAMOUS-ice (medium albedo) mean over available AOGCMs for each climate (no HadGEM2-ES for RCP8.5, all four AOGCMs in other cases). The first column identifies the “groups” of results into which the table is divided; we refer to these group numbers in the text. Ablation is $\text{SMB} - \text{snowfall}$, mainly runoff from snowmelt, and including evaporation, sublimation, condensation and rainfall freezing in the snowpack (in the RCMs; all rainfall runs off in FAMOUS-ice). The \pm uncertainty shown for SMB is the standard error of the time-mean, estimated by assuming annual values to be independent. The SMB from FAMOUS-ice has smaller standard errors than from the RCMs for two reasons. First, the FAMOUS-ice [integrations simulations](#) exclude interannual variability due to SST and sea-ice by using climatological mean BCs. Second, the RCM time-means use 20 years of data, while we use [30-100](#) years [from for](#) the [second group of](#) FAMOUS-ice [MIROC5 1980-1999](#) simulations, [which supply our initial steady states](#), and [+00 from the first](#) 30 for other FAMOUS-ice simulations, [which are transient states](#).

CMIP5 scenario	years	CO ₂	ERF	notes
historical	1980-1999	402	2.1	
RCP2.6	2080-2099	402	2.1	
RCP4.5	2080-2099	650	4.7	
RCP8.5	2080-2099	1200	8.0	not with HadGEM2-ES
abrupt4xCO2	121-140	1200	8.0	CanESM2 and low albedo only
abrupt4xCO2	101-120	1200	8.0	HadGEM2-ES and low albedo only

Table 2. AOGCM climates used to supply sea-surface boundary conditions for the first set of FiG experiments. The BCs mostly determine the climate, ~~while with only a relatively small influence from~~ the CO₂ concentration (in ppm) ~~has a relatively small influence. For simplicity only three different~~ ~~This is “equivalent CO₂ concentrations were used”,~~ chosen ~~for RCP4.5 and RCP8.5~~ to give approximately the nominal effective radiative forcing ~~in RCPs at 2100~~ (ERF, W m⁻²) ~~of the RCPs, and with~~ all other forcing agents ~~were kept~~ as pre-industrial. ~~For simplicity, regarding 1980-1999 as “present day”, we decided to use the same concentration for historical and RCP2.6 simulations. We consider this acceptable because the AR5 median assessment of the net anthropogenic ERF in 2011 is 2.3 W m⁻², with a likely range of 1.1-3.3 W m⁻², and the difference between this and the nominal forcing of 2.6 W m⁻² under RCP2.6 at 2100 is small compared with the large systematic uncertainty. Similarly for simplicity we used the same CO₂ concentration for RCP8.5 and abrupt4xCO2.~~

mean sea-surface BCs taken from the four selected CMIP5 AOGCMs for five climate scenarios (Table 2). These five are the ~~recent past late 20th century~~ (1980-1999, called “historical”), the end of the 21st century under three RCP scenarios (representative concentration pathway, as in the AR5; van Vuuren et al., 2011), and quadrupled pre-industrial CO₂ (abrupt4xCO2, warmer than any RCP). The experiments have steady-state climates. This is unrealistic, but it simplifies the comparison, and is reasonable since no-one can tell how climate will change over millennia into the future. Our simulations should be regarded only as indicative, rather than as projections.

Each experiment begins from the FiG spun-up state for MIROC5 historical climate with the appropriate albedo parameter. Although in most cases there is a substantial instantaneous change in BCs when the experiment begins, the land and atmosphere require only a couple of years to adjust. ~~Being climatological means, the sea-surface BCs lack interannual variability, and we have checked that statistically indistinguishable results for the ice sheet are obtained by cycling through a 20-year series of monthly mean BCs.~~

3.1 Evolution of surface mass balance

Our set of BCs produces a wide range of global mean surface air temperature change Δ SAT of -1 to +5 K, relative to the MIROC5 historical climate. Some are negative because the historical climate is warmer in MIROC5 than in the other three AOGCMs. In warmer climates, snowfall and ablation are both increased (Table 1 ~~for MIROC5 with medium albedo, fifth group shows results with MIROC5 RCP climates, last group shows the mean over results for each of the available AOGCMs for each climate~~). In general, the greater the global warming, the more negative the Δ SMB -initially produced, relative to the time-mean MIROC5 historical state with the same albedo (Figure 2a). ~~For a given scenario, the AOGCMs give a range of Δ SAT, as is very~~


well-known (*e.g.* Collins et al., 2013). In our set of AOGCMs, NorESM1-M warms least, HadGEM2-ES most (Figure S2a). Δ SAT in FAMOUS-ice and the BCs is very highly correlated (Figure S2b). The spread of FAMOUS-ice results with BCs from different AOGCMs for a given warming is due to their different relationships between global Δ SAT and Greenland regional climate change (shown by the grey lines in Figure S2c).

300 Global warming under RCP2.6 is fairly small, leading to small Δ SMB, especially for MIROC5 (squares near 1.0 K in Figure 2a), although MIROC5 is in the middle of the range for RCP8.5 (squares near 3.5 K). For mean over AOGCM climates under RCP2.6, RCP4.5 and RCP8.5, FAMOUS-ice with medium albedo gives SMB change of of -95 , -247 and -580 Gt yr $^{-1}$ respectively with respect to the mean over AOGCM historical climates.

The greatest global warming is given by HadGEM2-ES abrupt4xCO $_2$. With low albedo, this climate produces the most
305 negative Δ SMB, of -1063 Gt yr $^{-1}$ (it changes from $+307$ to SMB of -756 Gt yr $^{-1}$) in the time-mean of the first 300 years, during which the topography hardly changes from its initial state change from the initial state is still quite small (Figure 3a1,b1). It is also the most negative Δ SMB of -1066 Gt yr $^{-1}$ relative to the MIROC5 historical climate with low albedo. Although this is a large perturbation, the Δ SMB, that of Aschwanden et al. (2019) for RCP8.5 is larger still, perhaps because they use a degree-day scheme and assume geographically uniform warming.

310 In our experiment, the specific SMB is strongly negative all around the margin and especially in the southern dome, where it has a local maximum in the historical climate (Figure 3a2,b2). The snowfall on the ice-sheet is $\sim 10\%$ larger in the abrupt4xCO $_2$ climate (Figures 3a4,b4). We note that the precipitation is $\sim 50\%$ larger, consistent with the warming in Greenland of 11 K and the increase of $\sim 5\%$ K $^{-1}$ found by previous studies *e.g.* Gregory and Huybrechts (2006), but the snow fraction declines from ~~88% to 67%~~ $\sim 90\%$ to $\sim 70\%$. The downwelling surface shortwave radiation in summer (June–August) is smaller because

Figure 2 (following page). Relationships between various quantities in the first set of experiments, with FiG under constant climates listed in Table 2, and run to a steady state, as shown in Figure 4b. All panels use the key of (a) for colours; (b–e) use the key of (a) for symbols. (a) Time-mean Δ SMB vs. Δ SAT, both for the first 300 years relative to the initial steady state under the historical MIROC5 climate with the same albedo parameter. The solid curve is the cubic relationship fitted by Fettweis et al. (2013) to MAR projections, and the dashed curves delimit the likely range of the AR5. (b) Trajectories of ice-sheet SMB (not Δ SMB) vs. mass M , shown as 200-year means for the first millennium, and 1000-year means thereafter. The trajectories begin at the symbols, with M close to the observed for the present day, and a wide range of SMB. They end with a wide range of M , but all have positive SMB. (c) Final steady-state M vs. time-mean Δ SAT in FAMOUS-ice for the first 300 years. (d) Final steady-state M of the ice-sheet vs. time-mean Δ SMB of the first 300 years. The vertical dashed lines mark the observational estimates of Δ SMB for the recent periods and studies shown in the key (van den Broeke et al., 2016; Mouginit et al., 2019; Shepherd et al., 2020); for van den Broeke et al. we used the steady-state SMB for 1961–1990 and the SMB trend for 1991–2015. The oblique solid and dashed lines are linear regressions of M vs. Δ SMB and vice-versa respectively for Δ SMB > -700 Gt yr $^{-1}$. The solid horizontal lines indicate the threshold of irreversibility for medium and low albedo, and the solid vertical lines translate them into Δ SMB thresholds, with uncertainty (± 2 standard deviations) shown by the grey band. (e) Trajectories of ice-sheet thickness (volume divided by area) vs. specific SMB for 1000-year means, beginning at the symbols. (f) Trajectories of M vs. ice-sheet area as grey lines, with the final configurations indicated by the symbols, and the fitted power-law relationship shown by the black line.







315 cloudiness is greater (Figures 3a3,b3). Both the increased snowfall and the reduced insolation tend to make ΔSMB positive, but ΔSMB is actually large and negative because of the overwhelming effect of increased downwelling surface longwave radiation, which is mainly due to the air above the ice-sheet being warmer, and partly to the increase in cloud cover.

We find that the relationship between ΔSAT and ΔSMB in the set of FiG experiments roughly follows the cubic formula (shown as the solid curve in Figure 2a) derived by Fettweis et al. (2013) for MAR projections and used in the AR5 for the
320 Greenland contribution to GMSLR. There is a small spread due to the choice of albedo parameter, and a larger spread due to ~~the AOGCM-dependent relationship between ΔSAT and Greenland climate change~~choice of AOGCM. The FiG ΔSMB mostly lies within the AR5 likely range (dashed curves). In the majority of cases SMB remains positive (Figure 2b), but because the ice-sheet was initially in balance, negative ΔSMB leads to loss of mass (Figure 4a). In the most extreme case, rapid retreat of the ice-sheet margin reduces the discharge by a third in the first century alone; this slightly offsets ΔSMB , giving ice-sheet
325 mass-loss of 2.5 mm yr^{-1} SLE.

Because of the effect of lowering topography, the SMB becomes more negative in most cases during the early centuries (Figure 2b). For the the 21st century, this effect is omitted in our experiments, since we instantaneously impose the climates from the end of the century on the initial state. This is an acceptable approximation because the effect is small on that timescale e.g. Edwards et al. (2014) give a best estimate of 4.3% for the consequent increment in the GMSLR contribution by 2100, but this increases with time, e.g. to 9.3% by 2150 (Le clec'h et al., 2019), and 9.6% by 2200 (Edwards et al., 2014). In the
330 28 cases with $\Delta\text{SMB} < -100 \text{ Gt yr}^{-1}$ in the time-mean of the first century of our experiments, ΔSMB becomes about 20% more negative on average during the second and third centuries due to the elevation-local lapse-rate feedback, about twice the size of the effect estimated by Edwards et al. (2014) for scenario A1B by 2200. Edwards et al. Thereafter the SMB becomes gradually more positive again (Figure 2b), because the area contracts, with the areas most prone to ablation being removed
335 most quickly, as happens with a retreating mountain glacier.

Figure 3 (following page). Illustrative states of the ice-sheet, all from coupled FAMOUS–ice–Glimmer experiments except for column (d), as follows: (a) initial ~~steady~~-state with HadGEM2-ES historical climate and low albedo; (b) initial state with HadGEM2-ES abrupt4xCO2 climate and low albedo; (c) transient state from the experiment of (b); (d) transient state of uncoupled Glimmer with the climate and 3D SMB of (b); (e) final state with CanESM2 abrupt4xCO2 climate and low albedo; (f) final state with MIROC5 RCP4.5 climate and medium albedo; (g,h) final states with MIROC5 historical climate and low albedo, regrown from transient states with $M = 3.83 \text{ m SLE}$ and $M = 4.03 \text{ m SLE}$ respectively in the experiment of (b). The quantity shown in each row in colours, and by contour lines in rows (3–4), is stated above its colour bar. Row (1) is an instantaneous state; rows (2–4) are time-means of 30 FAMOUS–ice years, equivalent to 300 FiG years. The ice-sheet edge is shown by a thick black line in rows (2–4). The numbers in the bottom right corner are in row (1) ice-sheet mass in m SLE, (2) ice-sheet area-integral SMB in Gt yr^{-1} , (4) ice-sheet area-integral snowfall in Gt yr^{-1} . The symbols in row (1) indicate steady-state configurations by the key of Figure 2f.



3.2 Final ice-sheet mass and global-mean sea-level change

The experiments continue until the ice-sheet reaches a steady state (defined as $|dM/dt| < 0.02 \text{ mm yr}^{-1}$ SLE over 2000 years). The longest experiments, which take 40 kyr (Figure 4b), are for large climate change (RCP8.5), which entails a large loss of mass, with high albedo, which causes a relatively slow rate of mass-loss. The shortest are the experiments in which a different historical climate from MIROC5 is applied, because the effect on SMB of differences among AOGCMs in their simulations of recent-late 20th-century climate is relatively minor.

There is a wide range of M in the final steady state (Figures 2b and 4b), between slightly greater than present-day (in some historical experiments), and almost zero (in abrupt4xCO2). With one exception, historical and RCP2.6 climates produce final M of 6 m SLE or more (implying GMSLR not exceeding 1.5 m), while RCP8.5 climates all produce final M of 3 m SLE or less (GMSLR exceeding 4 m). In all cases the SMB is finally positive (Figure 2b), and must be balanced by ice discharge, meaning that the ice-sheet does not retreat entirely inland.

There is a clear tendency for climates of greater ΔSAT to produce smaller ice-sheets, but the final M has quite a wide range for any given initial global-mean annual-mean ΔSAT within 1–4 K (Figure 2c). For given BCs, we have found in test experiments that the ice-sheet evolution follows somewhat different trajectories from slightly different initial states, but that they converge on very similar final states (Figure S1a). Thus, the scatter in Figure 2c is not random noise, but arises from the detailed interaction of the evolving ice-sheet topography with its regional climate, which depends on the choice of BCs. The final M depends on which AOGCM is used, because of their different patterns of SST and sea-ice change; this dependence is omitted if the warming is assumed to be uniform (e.g. Robinson et al., 2012; Aschwanden et al., 2019).

The Pearson product–moment correlation coefficient between final M and initial ΔSAT is -0.89 – -0.89 , and the Spearman rank correlation coefficient -0.83 – -0.83 . The correlation is similar if for ΔSAT we use Greenland area-mean summer-mean air temperature change, either at the surface or at 600 hPa (the latter as Fettweis et al., 2013) (Figure S2d,e,f). However, the relationship is better-defined using ΔSMB instead of ΔSAT (Figure 2d), with both product–moment and rank correlation coefficients of 0.92. If the initial ΔSMB is near zero, the ice-sheet changes little; the more negative the initial ΔSMB , the smaller the final M . Excluding the case with the most negative ΔSMB , a linear relationship is a fairly good fit. On the basis of the MAR simulations of Fettweis et al. (2013), the CMIP5-mean projection of ΔSMB for 2080–2099 climate is -242 Gt yr^{-1} under RCP4.5 and -710 Gt yr^{-1} under RCP8.5. According to the fit, the former implies eventual GMSLR of about 3 m, the latter about 7

With any choice of albedo, for any $T > 3 \text{ K}$, the final steady-state $M \lesssim 2 \text{ m SLE}$, meaning GMSLR exceeds 5 m.

It is important to note, however, that the spread of final M does not suggest any threshold in ΔSAT for the a sharply defined threshold in T beyond which a complete or nearly complete loss of the ice-sheet ensues (Figure 2c). According to Section 1.3, there should be no final For low and medium albedo, there is a fairly monotonic decline in size of the steady state from near present-day $M \simeq 7 \text{ m}$ at $T = 0 \text{ K}$ to $M < 1 \text{ m}$ for $T > 3 \text{ K}$. For high albedo, there might be a transition from $M \simeq 6 \text{ m}$ at $T = 2.0 \text{ K}$ to $M \simeq 3 \text{ m}$ at $T = 2.5 \text{ K}$ —to obtain a clearer description of the behaviour, more experiments are needed in this part of the diagram. In any case, the interval between temperatures giving a “large” and a “small” final ice-sheet for negative

370 ~~initial-SMB, implying ΔSMB below -360 Gt yr^{-1} for FiG (Table 1), but actually there are such states (Figure 2d). This is not the picture we expected, since it is contrary to the findings of previous work, and we discuss it later (Section 3.1). Instead, we find that~~ is wider in our results, or alternatively the mass interval between “large” and “small” is narrower, than in the results of Levermann et al. (2013) (their Figure 1C). All of the versions of their model have a sharp transition between $M > 6 \text{ m}$ and $M < 1 \text{ m}$ over a temperature interval which appears to be less than 0.1 K . Our model gives a qualitatively different impression of the transition.

If negative feedbacks were neglected, there would be no final ice-sheet for negative initial SMB, as described in Section 1.3. Actually all final states have positive SMB and non-zero M , although some have initially negative SMB (Figure 2b). In our model, if any climate warmer than historical is maintained indefinitely the ice-sheet will contract to a new non-zero steady state, whose size depends on the magnitude of the warming and the consequent SMB perturbation.

380 ~~Recent-Observational~~ analyses indicate that ~~the present-recent~~ ΔSMB (with respect to a steady state before the 1990s) is between -200 and -150 Gt yr^{-1} , with substantial interannual variation (*e.g.* van den Broeke et al., 2016; Mouginit et al., 2019; Shepherd et al., 2020; dotted lines in Figure 2d). If a climate giving such a ΔSMB were maintained it would eventually lead to GMSLR of ~~$1.50.5$~~ 2.5 m according to the linear fit (Figure 2d, allowing for the range of FiG initial M). On the basis of the MAR simulations of Fettweis et al. (2013), the CMIP5-mean projection of ΔSMB for 2080–2099 climate is -242 Gt yr^{-1} under RCP4.5 and -710 Gt yr^{-1} under RCP8.5. According to the fit, the former implies eventual GMSLR of about 3 m , the latter about 7 m .

3.3 Interaction of ice-sheet and climate during decline

To demonstrate the important influence of the climate–ice-sheet interaction, we repeat the HadGEM2-ES abrupt4xCO2 low-albedo experiment (the case of most negative ΔSMB) using Glimmer alone, uncoupled from the AGCM, forced by the AGCM SMB field (a function of geographical location and tile elevation) from the start of the FiG experiment. As the uncoupled experiment runs, the time-independent three-dimensional AGCM SMB field is continually interpolated onto the time-dependent ice topography using the same methods as in the FiG coupling. Thus the ~~elevation-local lapse-rate~~ feedback on SMB is included in the uncoupled experiment, but the ~~feedbacks of ice-sheet topography-regional climate feedbacks of topography and albedo~~ change on the atmospheric state and circulation are excluded.

395 The uncoupled Glimmer and FiG experiments begin from the same initial state and have the same initial rate of mass-loss, but soon diverge (the dotted red line and the lowest solid red line in Figure 4a). While the rate of mass-loss continuously decreases in the FiG experiment, it remains almost constant (2.1 – 2.6 mm yr^{-1} SLE) in the uncoupled experiment for about 2.5 kyr , and the ice-sheet is completely eliminated in 3.4 kyr (Figure 4b).

To understand the different behaviour, as an example we compare the state when $M = 2.38 \text{ m SLE}$, which is reached after 3600 years in the coupled experiment and 2020 years in the uncoupled. The coupled ice-sheet has a high central region (Figure 3c1), where specific SMB exceeds 0.25 m yr^{-1} LWE over about the same area as in the initial state (Figure 3b2,c2), surrounded closely by steep narrow margins with large negative specific SMB, giving negative area-integral SMB which is ~ 3 times smaller in magnitude than in the initial state under $4\times\text{CO}_2$. The regions where negative specific SMB appears were




Figure 4. Timeseries of Greenland ice-sheet mass with constant climates and FAMOUS-ice albedo indicated by the line colours according to the line key of (c). (a,b) First set of experiments, beginning from steady states for MIROC5 historical (1980–1999) climate, and continuing until a new steady state is reached under the scenarios indicated by the colours of the final symbols in (b) according to its the final symbol colour key in that panel. The solid and dashed lines are FiG experiments; the dotted line is the experiment with the uncoupled Glimmer ice-sheet model. The circles indicate transient and final states which provide the initial states for the second set of experiments. (c) Second set of FiG experiments, beginning from states of the same albedo, and continuing until a new steady state is reached under under the MIROC5 historical climate. The single high-albedo experiment begins from the low-albedo initial state of smallest mass. The experiments shown by dashed lines in (c) begin from the final states of the experiments shown by dashed lines in (b). In all panels, the FAMOUS-ice albedo is indicated by the line colours. In (b) and (c), the final symbols denote the configuration of the final steady states according to the symbol key of (e), the final states marked “e”–“h” are those shown in the columns indicated of Figure 3, and the two horizontal lines marked “Threshold” indicate the mass that divides transient states which regrow to nearly the initial steady-state mass (WOWS-EWNS or EWN configurations) from those which regrow only partially (NON-“no-north” configurations: EWS, EW and E).

near to equilibrium in the initial state, and the change is consistent with the elevation-local lapse-rate feedback due to the
405 lowered surface in the contracted margins. The area-mean ratio of changes in surface air temperature and surface elevation is
7.1 and 6.6 K km^{-1} within the initial and contracted ice-sheet extent respectively, close to the value of 6 K km^{-1} assumed in
the downscaling scheme. It is not uniform over the ice-sheet (Figure S3), but it is within in the range $4\text{--}8 \text{ K km}^{-1}$ over more
than half of the ice-sheet (considering either extent).

The uncoupled ice-sheet is similarly located in the north of Greenland, but has larger area and lower altitude (Figure 3d1).
410 Its specific SMB is negative ~~everywhere, and its~~. Its area-integral SMB (-991 Gt yr^{-1} , Figure 3d2) is more negative than in
the initial state (-756 Gt yr^{-1} , Figure 3b2), and ~ 4 times more than for the coupled ice-sheet (~~of the same M~~ (-262 Gt yr^{-1} ,
Figure 3e2c2). The much larger change exceeds the lapse-rate effect, and the area-mean specific SMB for any surface altitude
above 1000 m is more negative in the uncoupled case than the coupled. The main cause is greater downwelling shortwave radi-
ation at the surface in the uncoupled case (Figure 3c3,d3), due to lower cloud fraction. The region occupied by the contracted
415 ice-sheet coincides geographically with the high cold interior of the initial ice-sheet, where cloudiness is comparatively low,
but in the coupled case the cloudiness increases there as the ice-sheet becomes smaller and lower, giving a powerful negative
feedback on the mass-loss.

In the coupled experiment, the precipitation from the south-west advances inland, following the the margin of the contracting
ice-sheet (compare the grey contour line for 1 m yr^{-1} in Figure 3b4,c4). Consequently the precipitation on the ice-sheet is
420 about 15% greater in the coupled case. However, the snowfall is about 15% less in the coupled case (colours and numbers
in Figure 3c4,d4), because its surface is lower than in the initial climate, making the surface climate warmer and reducing
the snowfall fraction (to 64%). The uncoupled SMB has a larger snowfall fraction (84%) because the surface in the region it
occupies was initially much higher. The phase change of precipitation with elevation is omitted from the downscaling in the
coupling scheme (as mentioned earlier); including it in the uncoupled model would reduce the snowfall, and make its SMB
425 even more negative.

In summary, the uncoupled ice-sheet is eliminated rapidly through the small ice-cap instability (elevation-local lapse-rate
feedbacks from surface energy fluxes and temperature), whereas in the coupled case the decline is decelerated, and the ice-
sheet not completely ~~elimated~~eliminated, owing to negative feedbacks of topographic change on regional climate (changes in
cloudiness and precipitation). The comparison demonstrates the critical role of ice-sheet-climate interaction.

430 3.4 Final topography of the ice-sheet

According to the topographic features present, the final states can be put in five categories (indicated by symbols at the ends of
the trajectories in Figure 4b). In cases with small change in M , the final state is similar to the present day (configuration labelled
“EWNS” *e.g.* Figure 3a1). The northern portion (denoted “N”) is absent in some final states and the summit further south than
in the present day *e.g.* Figure 3f1 (EWS). Ice in the south (“S”) may become a separate ice-cap (as in Figure 3f1) or it may be
435 absent, resembling Figure 3h1 (EWN) and 3g1 (EW). In cases with the smallest final M , the north-western lobe (“W”) vanishes,
and ice remains only on the eastern mountains (E“E”). For example, in the experiment ending in Figure 3e1 (marked with “e”

in Figure 4b), the southern and north-western domes detach and vanish within 3 kyr. Subsequently contraction continues on all sides, but there is a slow small regrowth after the minimum mass is reached.

440 The transient and final states of all experiments lie close to a common power-law relationship between ice-sheet mass M and area A with $M \propto A^{1.31}$ (Figure 2f), similar to the exponent of 1.36–1.38 derived for glaciers from observations and theory (Bahr and Radić, 2012, and references therein). Final states with the same configuration have a characteristic deviation from the common relationship *e.g.* EWN states have greater M .

445 Because of the elevation-local lapse-rate feedback, the mass-loss sometimes accelerates by a few tenths of a mm yr^{-1} SLE while one of the outlying portions becomes separate or is eliminated, in a few cases after some millennia of relatively slow change. This is a similar phenomenon to the saddle collapse during the separation of the Laurentide and Cordilleran ice-sheets during the last deglaciation (Gregoire et al., 2012), but an order of magnitude smaller. For example, in the experiment ending in Figure 3f1 (the dotted green line, marked “f”, in Figure 4b), the rate of ice loss accelerates after 10 kyr, at the start of the retreat of the northern margin, which is completed by 15 kyr.

4 **Non-existence of a threshold warming**

450 3.1 Discussion of reduced steady states

In Section 1.3 we described why ~~ΔSAT that reduces Greenland SMB to zero is expected to~~ there might be a threshold **warming** ΔSAT beyond which the ice-sheet would be eliminated by the small ice-cap instability, whereas with smaller ΔSAT it would have mass and area little reduced from its present-day state. In Section 3 ~~we found no evidence for a threshold from FiG results, meaning that the conceptual basis for its existence is incorrect, instead of such a well-defined threshold, we found a range~~ of steady-state ice-sheet mass and area, generally smaller for larger ΔSAT . The ice-sheet endures, albeit in a much reduced state, even for ΔSAT giving large negative SMB, ~~because the expected runaway feedback (the small ice-cap instability) does not occur. The positive feedbacks on mass-loss due to reduction of thickness are weak, and overwhelmed by effects due to reduction of area. The conceptual model considers that the smallest possible steady state has almost the same area as now, but in FiG most steady states have smaller area. Because neither the initial specific SMB nor the change in it are geographically~~ uniform, some parts initial SMB.

460 For studying the evolution of the ice-sheet as its area A contracts, it is helpful to consider the specific SMB $s = S/A$, where obviously S and s have the same sign. We can write $\Delta s = \Delta s_T + \Delta s_L + \Delta s_C$, where Δs_T and Δs_L are the changes in specific SMB due to climate change and the local lapse-rate feedback, as in Section 1.3. When the warmer climate is initially imposed, $\Delta s_T < 0$, and the perturbation is amplified by $\Delta s_L < 0$ due to thinning of the ice-sheet.

465 The term Δs_C represents the effects of change in the climate experienced by the ice-sheet, arising both because the climate changes in all areas, and because the ice-sheet changes the areas it occupies. An important example of the latter is the retreat of the ice-sheet ~~can be removed more readily~~ margin (or a glacier tongue, in general) to higher altitude in a warmer climate, because this reduces the ablation while preserving the accumulation. In this and other cases, the climate effects can give $\Delta s_C > 0$. Thus they can counteract the local lapse-rate feedback $\Delta s_L < 0$, prevent a runaway feedback loop, and eventually

470 reverse the sign of Δs so that a steady state is reached with SMB and discharge in balance again, even if with greatly reduced area.

In cases where specific SMB is initially positive, it becomes more positive (Figure 2e), because the areas from which the ice-sheet retreats are predominantly those of relatively larger ablation or smaller snowfall. Consequently the area-integral SMB (the product of increasing specific SMB and decreasing area) changes relatively little (fairly vertical trajectories in Figure 2b).
475 For instance, under MIROC5 RCP4.5 climate with medium albedo, the initial S , P and R SMB, snowfall and ablation are 150, 735 and 585 Gt yr⁻¹ (Table 1). The final S SMB is the same as the initial because R and P ablation and snowfall both decrease by 115 Gt yr⁻¹ (Figure 2f2,f4), a larger fractional decline in R ablation (20%) than in P snowfall (15%). The steady state is achieved by the withdrawal of the margin from the coast in some sectors, reducing D discharge sufficiently (by 209 Gt yr⁻¹ or 60%) to balance the smaller S SMB.

480 In cases where specific and area-integral SMB are initially negative, they become positive (Figure 2b,e). This happens because P snowfall decreases less than R ablation. For instance, under HadGEM2 abrupt4xCO2 climate with low albedo, the initial S , P and R SMB, snowfall and ablation are -756, 797 and 1554 Gt yr⁻¹ (Figure 2b2,b4). The final state is a small eastern ice-cap (like Figure 2e1 but smaller) with S , P and R SMB, snowfall and ablation of 9, 31 and 21 Gt yr⁻¹; P snowfall is 3.9% and R ablation 1.4% of the initial value. The ice-cap receives greater precipitation and snowfall than the same region did
485 initially (compare Figure 2b4,e4), and has more cloud and less surface downwelling shortwave (Figure 2b3,e3), because of the effect of topography on atmospheric circulation and climate.

4 Threshold for irreversible mass-loss

Greenland ice-sheet mass-loss in the first set of experiments occurs on timescales which are comparable with or even longer than those of surface climate change and natural CO₂ removal. We therefore also consider whether the ice-sheet mass would
490 increase again if the climate cooled down. This will inform us about any irreversible commitment to GMSLR that might be incurred in coming decades despite subsequent CO₂ removal.

To study this question, we carry out a second set of FiG experiments, using MIROC5 1980-1999 BCs and recent CO₂ radiative forcing, starting from various transient and final steady states of the ice-sheet with reduced size from the first set of experiments. This is as if the climate instantaneously reverted to its late 20th-century-20th-century condition after many centuries in a high-
495 CO₂ warm steady state, during which the ice-sheet had been losing mass. The second set includes experiments with all three choices of albedo. All but one of the experiments with medium albedo (solid green lines in Figure 4c) begin from states of various mass along the trajectory of the CanESM2 RCP8.5 medium-albedo experiment (green line with circles in Figure 4b), whose final steady-state ice-sheet mass is 1.21 m SLE. All but one of those with low albedo (solid red lines in Figure 4c) begin from states of the HadGEM2-ES abrupt4xCO2 low-albedo experiment (red line with circles in Figure 4b), whose final mass
500 of 0.12 m SLE is the smallest of all in the first set. The single high-albedo experiment in the second set (solid blue line in Figure 4c) also begins from this minimal state. (The exceptions for medium and low albedo are the two experiments discussed

in Section 4.2 and shown with dashed lines in Figure 4c, which begin from the final states of the experiments shown with dashed lines in Figure 4b.)

4.1 Regrown steady states

505 In the initial state of all the experiments of the second set, the ice-sheet has a smaller mass than present, and it grows to reach a new steady state; there are none in which it continues to lose mass (Figure 4c). However, the mass of the regrown steady state depends on the initial state and the albedo.

With high albedo, the ice-sheet regrows, in about 50 kyr, from the minimal state to a steady state with the extent of the present day's (EWNS configuration, Figure 4c). Since this starting state is a limiting case, we assume that the ice-sheet would reach the same final state from any initial state, implying that this is only steady state for historical climate with high albedo. 510 Therefore the loss of the ice-sheet would be reversible, albeit on a long timescale, if the high albedo is realistic.

On the other hand, with the medium and low albedo, two distinct sets of steady states can be reached in the second set of experiments, one set with final mass of 7 m SLE or more, the other with final mass of 5–6 m SLE. Initial states are divided between these two sets of final states by a threshold of initial mass at 4.0 m SLE with the medium albedo, and 3.9 m SLE with 515 the low albedo.

Starting above the threshold, the ice-sheet regrows to the EWNS configuration with medium albedo as with high (Figure 4c), but with low albedo there are two steady states. The larger is the EWNS configuration (7.3 m SLE, Figure 3a1), while the smaller lacks the southern dome (7.0 m SLE, EWN configuration, Figure 3h1). ~~We will refer to these two configurations together as WOWS (“with or without S”).~~ The southern dome has positive SMB in the historical climate (Figure 3a2), but in 520 the warm climates it is readily lost due to increased ablation, and in the historical climate without the dome there is negative SMB inhibiting readvance at the new southern margin (Figure 3h2). The snowfall is however little changed in that region (Figures 3a4,h4). The southern dome is the last part of the ice-sheet to reappear with the high and medium albedoes (solid green and blue lines in Figure 4c after 30 kyr); ~~and perhaps it would do so with low albedo after sufficiently long as well.~~

Starting below the threshold, the ice-sheet attains steady states lacking the northern portion, which we will refer to collectively as ~~NON states (“no N”)~~ no-north states. The steady state with medium albedo has the EWS configuration (5.7 m SLE, like 525 Figure 3f1 in extent but thicker). With the low albedo, there are two steady states, having masses of 5.3 m SLE (EW configuration, Figure 3g1) and 5.0 m SLE (E configuration, like Figure 3e1 but much larger), which differ because the north-western dome is missing in the latter case. ~~Again, we suppose this dome might regrow in time, given that it~~ This dome is the last part to ~~do so~~ regrow with medium albedo.

530 Other authors have likewise found that the present state of the Greenland ice-sheet is not the only steady state under historical climate (Ridley et al., 2010; Solgaard and Langen, 2012; Robinson et al., 2012). A minimal state with ice solely or mostly in the east is a common feature of all these studies and ours. In other respects the steady-state configurations are dissimilar. The medium state of Robinson et al. (2012) most resembles our ~~NON~~ no-north states. Our results are more complex than others in showing five steady states. We suppose that this is because greater detail in the interaction of the ice-sheet topography with 535 atmospheric circulation and SMB can be simulated by FAMOUS-ice than by the simpler approaches of previous studies.

4.2 Vulnerability of the ice-sheet to irreversible loss

To summarise our second set of experiments: transient states which have passed below the threshold regrow to NON-no-north steady states, while those still above the threshold regrow to WOWS-EWNS or EWN steady states. Consistent with this, we note that all final states lying below the threshold in the first set of experiments are NON-no-north states (Figure 4b). States
540 taken from below the threshold on trajectories of rapid decline show no tendency for the northern portion to regrow, even after tens of millennia under historical climate. ~~The difference between initial states just above and just below the threshold is infinitesimal, yet sufficient to perturb the climate and SMB such as to make them diverge radically in outcome through,~~ Thus about 2 m of GMSLR will become irreversible once the Greenland ice-sheet mass drops below the threshold, if the medium or low albedo is realistic.

Under the same BCs, initial states which differ only slightly in topography (the minimum separation of our initial states in
545 M is actually 0.2 m SLE) can lead to final states which differ substantially (by more than 1 m SLE) because ice-sheet-climate feedbacks amplify the initially small difference in SMB. The probable reason is that ablation exceeds accumulation in the northern region without the ice-sheet (shown by negative SMB at the northern margin in Figure 3g2), partly because snowfall is reduced (Figure 3a4,g4). ~~Thus about 2 m of GMSLR will become irreversible once the Greenland ice-sheet mass drops below the threshold, if the medium or low albedo is realistic.~~

550 The low- and medium-albedo NON-steady-states-no-north steady states following regrowth are 1–2 m SLE above the threshold, and yet grow no further, ~~because they have lost the northern portion. This implies that, for states unlike states along trajectories of rapid decline having M in the same mass range *i.e.*~~ between the threshold mass (4 m SLE) and the NON-no-north mass (5–6 m SLE). The implication is that, for states in this mass range, the outcome depends on the history. To test this, we have conducted further experiments (dashed lines in Figure 4c) beginning from the two steady states in this range (large circles
555 at the end of dashed lines in Figure 4b), which were reached by slowly-slowly declining trajectories. These two are no-north (EWS) states. Initially the ice-sheet mass grows but, unlike when starting from rapidly declining transient states in this range, it soon becomes nearly constant at a slightly higher M than is reached from states below the threshold. ~~This difference. The difference in M~~ is due to a large southern dome ~~in these cases,~~ which was kept during the slow decline (along the dashed lines leading to the large circles in Figure 4b) but had been lost already in states of the same mass in the warmer climate producing
560 the fast decline (the solid lines with red and green circles in Figure 4b), and is not rebuilt in the historical climate. This result suggests that, for slow or quasi-static decline of the ice-sheet, the NON-no-north mass itself is the threshold of irreversibility.

Using the linear relationship between the initial rate of mass-loss and the final steady-state mass in the first set of experiments (solid line in Figure 2d), we can translate the threshold ~~ice-sheet mass (of irreversibility ($M=3.9-4.0$ m,~~ horizontal red and green lines), which applies during trajectories of rapid decline, into a threshold on ~~initial rate the rate~~ of loss (vertical red and green
565 lines). Under a warm climate which initially gives a more negative Δ SMB than the threshold rate, the ice-sheet will eventually decline to a state which is smaller than the threshold mass. Roughly estimating a range from the scatter in the relationship, the results suggest that the threshold Δ SMB lies between -500 and -150 Gt yr $^{-1}$ (Figure 2d). ~~Recent~~ Since recently observed Δ SMB (*e.g.* van den Broeke et al., 2016) is at the upper end of this range (~~van den Broeke et al., 2016~~). ~~Since the present-day rate of mass-loss is at the lower end of our scenarios,~~ *i.e.* a relatively small rate of decline, we recall from the previous paragraph

570 ~~that~~ the relevant threshold may instead be the ~~NON-no-north~~ steady-state mass of about 5.5 m SLE(~~as suggested in the previous~~
~~paragraph), in which~~. ~~In that~~ case the linear fit indicates that the ~~present-recently observed~~ Δ SMB is close to the threshold ~~for~~
~~partial-rate which will eventually lead to partially~~ irreversible loss of the ice-sheet.

If the ~~present-recently observed~~ rate of mass-loss of about 0.7 mm yr^{-1} SLE persisted, it would take ~~about 5000-4900~~ years
for the ice-sheet mass to reach the threshold of irreversibility, and ~~2500-about 2700~~ years to reach the ~~NON-no-north~~ steady-
575 state mass. At the highest rate of loss simulated in our experiments for the end of this century, of about 2 mm yr^{-1} SLE, it
would take 1700 years to reach the threshold. Allowing for systematic uncertainty, the AR5 predicted even larger rates of
mass-loss due to SMB perturbation, of up to about 6 mm yr^{-1} SLE by the end of the century, at which rate the threshold would
be reached in 600 years.

In order to avoid eventual irreversible ice-loss, the climate must be returned to near pre-industrial before the threshold mass
580 is reached. Reversing climate change requires extracting heat from the ocean, as well as removing the radiative forcing. If that
can be done at all, it could not be done instantaneously, and mitigating climate change in the short term will buy more time to
save the ice-sheet on the long term. Further simulations would be required to evaluate whether particular trajectories of future
climate would avoid irreversible ice-loss.

5 Conclusions

585 We have studied the multimillennial future evolution of the Greenland ice-sheet ~~in-response-to-for~~ various magnitudes of
anthropogenic climate change, in experiments with ~~constant climates using~~ an AGCM interactively coupled to a dynamic
ice-sheet model. For adequate resolution of gradients, especially at the margins of the ice-sheet, the surface mass balance is
simulated by the AGCM as a function of elevation within its gridboxes. Our aim is not to produce time-dependent projections
for coming centuries, but instead to investigate the long-term consequences for global-mean sea-level rise (GMSLR).

590 Under ~~constant~~ climates that are warmer than the late 20th century, the ice-sheet loses mass, its surface elevation decreases
and its surface climate becomes warmer. This gives a positive feedback on mass-loss, but it is outweighed by the negative
feedbacks due to declining ablation area and increasing cloudiness over the interior as the ice-sheet contracts. ~~Snowfall~~ ~~In~~
~~the ice-sheet area-integral, snowfall~~ decreases less than ablation because the precipitation on the margins is enhanced by the
topographic gradient, and moves inland as the ice-sheet retreats. Consequently after many millennia under a constant warm
595 climate the ice-sheet reaches a reduced steady state. ~~Final GMSLR is less than 1.5 m in most late-21st-century RCP2.6 climates,~~
~~and more than 4 m in all late-21st-century RCP8.5 climates. For warming exceeding 3 K, the ice-sheet would be mostly lost,~~
~~and its contribution to GMSLR would exceed 5 m.~~

Contrary to expectation based on work using simpler climate models (Huybrechts et al., 1991; Gregory et al., 2004; Gregory
and Huybrechts, 2006; Robinson et al., 2012; van den Broeke et al., 2016; Pattyn et al., 2018), we ~~find-no-evidence-for-a~~
600 ~~threshold-in-do not find a sharp threshold in regional~~ Greenland or global warming that divides scenarios in which the ice-
sheet suffers little reduction in its final steady state from those which it is mostly lost. ~~Our results give a qualitatively different~~
~~impression, because the transition occurs over a larger temperature interval, or involves a smaller mass-loss.~~ We think that

this difference arises from our using an AGCM, whose dynamics and physical detail are needed to simulate the response of snowfall and cloudiness to the evolving topography. Support for this hypothesis comes from comparison with an experiment using the uncoupled ice-sheet model, in which the surface mass balance evolves only through the ~~elevation-local lapse-rate~~ feedback, and ~~effects-of-atmospheric-circulation-change~~ regional climate feedbacks are omitted. In that case an almost constant rate of mass-loss is maintained for 3 kyr, during which the ice-sheet vanishes completely.

Under a warm climate, the final ice-sheet mass, and the entailed commitment to GMSLR, are well-correlated with the initial perturbation to surface mass balance, and hence with the magnitude of climate change imposed. The final mass is also affected by the geographical pattern of climate change. ~~If a climate like the present~~ According to a linear regression of our results, if a climate giving an SMB similar to that recently observed were maintained indefinitely, Greenland ice-sheet mass-loss would produce ~~1.50,5–2.5 m of GMSLR;~~ with the climate of the RCP8.5 scenario for the late 21st century, the ice-sheet would be almost completely eliminated, with over 7 m of GMSLR.

When transient and steady states of the ice-sheet obtained under warm climates are transplanted into the late ~~20th-century~~ 20th-century climate, as if subsequent anthropogenic climate change had been reversed, the ice-sheet regrows in all cases, over tens of millennia, but not necessarily to its present-day size (as also found by Charbit et al., 2008; Ridley et al., 2010; Robinson et al., 2012). The resulting steady states can be put in two groups, according to whether ice is present in the northern part of the island. If the ice-sheet retreats from this region, it may not regrow, because the snowfall is reduced there, meaning that about 2 m of GMSLR would become irreversible. This threshold size might eventually be reached with ~~the present-day late~~ 20th-century climate, and would be reached in about 600 years with the greatest rates of mass-loss projected for 2100 under RCP8.5 by Church et al. (2013). In order to avoid irreversible GMSLR, it would be necessary to restore the late 20th-century climate, in which the ice-sheet was near to mass balance, before the threshold is crossed.

The reliability of our conclusions depends on the realism of our model. There are systematic uncertainties arising from assumptions made in its formulation ~~(for example, the uniform-~~ The atmosphere GCM has low resolution and comparatively simple parametrisation schemes. The ice-sheet model does not simulate rapid ice-sheet dynamics; this certainly means that it underestimates the rate of ice-sheet mass-loss in coming decades, but we do not know what effect this has on the eventual steady states, which are our focus. The SMB scheme uses a uniform air temperature lapse rate and the omission of omits the phase change of precipitation in the downscaling) ~~from GCM to ice-sheet model.~~ The snow albedo is a particularly important uncertainty; with our ~~high-highest~~ choice of albedo, removal of the ice-sheet is reversible ~~under present-day climate.~~ Nonetheless,

Notwithstanding these limitations, our results demonstrate the importance of climate–ice-sheet interaction to projecting the future of the Greenland ice-sheet. It would obviously be useful if similar investigations were done ~~with-using~~ other models that couple an ice-sheet to an atmosphere GCM (perhaps as components of an AOGCM or Earth system model), especially with higher resolution in both the atmosphere and ice-sheet components. Even with ~~a-our~~ low-resolution GCMs ~~such as ours,~~ large ensembles of long experiments are computationally demanding, ~~meaning that and~~ our results give only ~~a sketchy-an~~ outline of possible behaviour, ~~and they-~~ They could be supplemented by using an emulator to explore a wider range of scenarios (Edwards et al., 2019).

Data availability. The model data used in this analysis will be made freely available for research at the Centre for Environmental Data Analysis (www.ceda.ac.uk) upon publication of this paper.

640 Appendix A: Relationship Technical sensitivity tests of albedo to steady-state historical M FAMOUS-ice-Glimmer

In order to test the sensitivity to certain technical changes, we ran three modified versions of the FAMOUS-ice-Glimmer experiment with CanESM2 RCP8.5 climate and medium albedo (from which the medium-albedo experiments of Section 4.1 begin, shown by a green line with circles in Figure 4b, and the solid black line in Figure S1a). The ice-sheet mass in each of the modified experiments differs by less than 0.2 m SLE from the standard experiment during the first 2000 years (Figure S1a).




Figure S1. (a) Timeseries of Greenland ice-sheet mass for the first 2000 ice-sheet years in experiments with CanESM2 RCP8.5 2081–2100 climate and FAMOUS-ice medium albedo as an example of sensitivity to technical modifications: different AGCM initial state, individual monthly means for the sea-surface BCs (“interannual variability”) rather than climatological monthly means, and synchronous coupling (one ice-sheet year per climate year, “no acceleration”) rather than 10:1 acceleration. The numbers in parentheses give the final steady-state mass. (b) Timeseries of Greenland ice-sheet mass with constant climate for 1980–1999 simulated by MIROC5 during FiG spinup integrations beginning from the observed topography (Bamber et al., 2001a, b). The crosses indicate the states from which the experiments of Section 3 were initiated.

645 The ice-mass $M(t)$ in the first modified experiment (dashed black line in Figure S1a) remains within ± 0.2 m of the standard experiment throughout its length, and is 0.1 m less than the standard in the final steady state. It is identical in forcing to the standard experiment but begins from a different atmosphere initial state of the same historical climate. Therefore its deviation from the standard experiment is due to chaotic unforced climate variability alone. The size of this unforced deviation is small compared with the differences of outcome due to climate and albedo among the experiments discussed in the paper, showing
650 that the forced differences are statistically significant.

For its sea-surface BCs, the second experiment cycles repeatedly through the 20-year series of individual monthly means that were used to make the 20-year climatological monthly means of the standard experiment. Thus it contains interannual variability in the climate. Its $M(t)$ (blue line in Figure S1a) is always within ± 0.4 m of and in the end 0.1 m less than the standard experiment's.

655 The third experiment has the same BCs as the second, and differs in addition from the standard experiment in that the ice-sheet model is run for only one year (not ten) after each FAMOUS-ice year. Because this version is almost ten times slower, we ran it for only 1700 years. During that period, it differed in $M(t)$ by less than 0.05 m from the standard experiment (red line in Figure S1a). The second and third experiments both have different SMB in every FAMOUS-ice year, but the acceleration (in the second experiment) makes these persist for a decade in the ice-sheet model. We think this explains the
660 greater excursions of the second experiment from the standard model.

Appendix B: Relationship of albedo to steady-state historical ice-sheet mass

Continuing the spinup experiments (which are among the experiments of Section 3), ice-sheet mass M remains at 7.7 m SLE with low albedo, with medium albedo it declines slightly to a steady state of 7.4 m SLE (very close to observed) over about 10 kyr, and with low albedo to 7.1 m SLE over 15 kyr (Figure ??S1b). These are small changes compared with those
665 simulated for 21st-century climate change (Section 3). Nonetheless, these small differences in M for low and high albedo from observations show that requiring a realistic steady state of the ice-sheet in a coupled model provides a strong constraint on the SMB simulation to which regional climate models such as MAR and RACMO are not subjected. A quadratic fit to the relationship between SMB and M in FiG steady states with MIROC5 historical climate gives $M = 7.9$ m SLE for the SMB of 437 Gt yr⁻¹ simulated by MAR for this climate.

670 An even higher choice of albedo in FiG gave SMB of 610 Gt yr⁻¹ and a steady-state M of 8.2 m SLE, and an even lower choice 195 Gt yr⁻¹ with M tending towards a steady state substantially below 7.0 m. These values of SMB approximately bound the range of SMB variations in the 20th century reconstructed with MAR (Fettweis et al., 2017, their Figure 8a), indicating that they could plausibly occur with historical climate and the present-day ice-sheet topography (as in MAR), but we excluded those choices of albedo because they would not be consistent with realistic M .




Figure S2. Timeseries of Relationships among air temperature change and Greenland ice-sheet mass with constant climate SMB change in CMIP5 AOGCM and FAMOUS-ice-Greenland experiments. The dotted line in (b) is 1:1. The grey lines in (c) are regression lines for 1980–1999 simulated by MIROC5 during FiG spinup integrations beginning the subset of data from each of the observed topography (Bamber et al., 2001a, b) four AOGCMs, indicated by the symbols along the top edge. The crosses indicate (d) is the states same as Figure 2c and repeated here for comparison. Changes are computed from which the experiments of Section 3 were initiated first 300 ice-sheet years and expressed relative to the MIROC5 historical climate with the same albedo parameter.

675 **Appendix C: ~~Alternatives to Δ SAT~~Alternative measures of air temperature change**

In Figure 2 we obtain global-mean annual-mean surface air temperature ~~change~~ (SAT) change, denoted Δ SAT, from FAMOUS-ice. It may also be obtained from the AOGCMs that supply the sea-surface BCs ~~-This~~ (Figure S2a). Separating the values by scenario reveals the AOGCM-dependence of global-mean SAT. NorESM1-M is cooler in general. HadGEM2-ES and CanESM2 have higher climate sensitivity, meaning that they warm more in response to forcing, while NorESM1-M has lower climate sensitivity and warms less.

680 Δ SAT in the CMIP5 AOGCMs is almost the same as in FAMOUS-ice (Figure ~~??a~~S2b). They differ because ~~surface-air temperature~~ SAT is not prescribed over land from the BCs in FAMOUS-ice.

We have investigated two other measures of air temperature change but neither is a better predictor than Δ SAT for the final ~~M-ice-sheet mass~~ (Figure ~~??e,d~~S2e,f).

685 ~~(a) Relationship between global-mean annual-mean surface air temperature change in FAMOUS-ice AGCM experiments and in the portions of the CMIP5 AOGCM experiments used to supply the FAMOUS-ice SST boundary conditions; (b,c,d) Relationships of Δ SMB to (b) FAMOUS-ice global-mean annual-mean SAT change (the same as Figure 2e and repeated here for comparison), (c) FAMOUS-ice Greenland-area-mean summer-mean (JJA) SAT change, (d) FAMOUS-ice area-mean summer-mean air temperature change at 600 hPa over Greenland. Changes are computed from the first 300 ice-sheet years of~~

690 ~~each FAMOUS-ice experiment and expressed relative to the MIROC5 historical climate with the same albedo parameter.~~

Appendix D: Lapse rate

In the downscaling of surface air temperature from FAMOUS gridboxes to FAMOUS-ice elevation tiles, we assume a uniform lapse rate of 6 K km^{-1} . Consequently this lapse rate is also used to predict the derivative of surface air temperature with respect to elevation change when Glimmer is run uncoupled from the AGCM. The derivative diagnosed from the coupled experiment

695 is shown in Figure S3.

Author contributions. SEG and RSS developed the model, SEG and JMG ran the experiments, JMG carried out the analysis and wrote the paper.

Competing interests. The authors have no competing interests.

Acknowledgements. We are grateful to John Church and Jason Lowe for often emphasising the importance of the long-term future of

700 Greenland and GMSLR and thus motivating the research. We thank the reviewers (Alex Robinson, Xavier Fettweis and one other) for their thoughtful and constructive comments, which helped us to improve the paper. This work was supported by NERC grants NE/P014976/1 and NE/I011099/1.




Figure S3. Change in surface air temperature divided by change in surface altitude (K km^{-1}) in the difference between the initial state of the experiment with HadGEM2-ES abrupt4xCO₂ climate and low albedo and the state after 3600 years (Figures 3b1,c1). The thick black line is the ice-sheet edge in the latter state.

References

- Aschwanden, A., Fahnestock, M. A., Truffer, M., Brinkerhoff, D. J., Hock, R., Khroulev, C., Mottram, R., and Khan, S. A.: Contribution of the Greenland ice sheet to sea level over the next millennium, *Sci. Adv.*, 5, eaav9396, <https://doi.org/10.1126/sciadv.aav9396>, 2019.
- Bahr, D. B. and Radić, V.: Significant contribution to total mass from very small glaciers, *The Cryosphere*, 6, 763–770, <https://doi.org/10.5194/tc-6-763-2012>, 2012.
- Bamber, J., Layberry, R. L., and Gogenini, S. P.: A new ice thickness and bed data set for the Greenland ice sheet 1: Measurement, data reduction, and errors, *J. Geophys. Res.*, 106, 33 773–33 780, <https://doi.org/10.1029/2001JD900054>, 2001a.
- 710 Bamber, J. L., Ekholm, S., and Krabill, W. B.: A new, high-resolution digital elevation model of Greenland fully validated with airborne laser altimeter data, *J. Geophys. Res.*, 106, 6733–6745, <https://doi.org/10.1029/2000JB900365>, 2001b.
- Bamber, J. L., Westaway, R. M., Marzeion, B., and Wouter, B.: The land ice contribution to sea level during the satellite era, *Environ. Res. Lett.*, 13, 063 008, <https://doi.org/10.1088/1748-9326/aac2f0>, 2018.

- Charbit, S., Paillard, D., and Ramstein, G.: Amount of CO₂ emissions irreversibly leading to the total melting of Greenland, *Geophys. Res. Lett.*, 35, <https://doi.org/10.1029/2008GL033472>, 2008.
- 715 Church, J. A., Clark, P. U., Cazenave, A., Gregory, J. M., Jevrejeva, S., Levermann, A., Merrifield, M. A., Milne, G. A., Nerem, R. S., Nunn, P. D., Payne, A. J., Pfeffer, W. T., Stammer, D., and Unnikrishnan, A. S.: Sea Level Change, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., Cambridge University Press, <https://doi.org/10.1017/CBO9781107415324.026>, 2013.
- 720 Clark, P. U., Shakun, J. D., Marcott, S. A., Mix, A. C., Eby, M., Kulp, S., Levermann, A., Milne, G. A., Pfister, P. L., Schrag, B. D. S. D. P., Solomon, S., Stocker, T. F., Strauss, B. H., Weaver, A. J., Winkelmann, R., Archer, D., Bard, E., Goldner, A., Lambeck, K., Pierrehumbert, R. T., and Plattner, G.-K.: Consequences of twenty-first-century policy for multi-millennial climate and sea-level change, *Nature Climate Change*, 6, 360–369, <https://doi.org/10.1038/NCLIMATE2923>, 2016.
- 725 Collins, M., Knutti, R., Arblaster, J. M., Dufresne, J., Fichefet, T., Friedlingstein, P., Gao, X., Gutowski, W. J., Johns, T., Krinner, G., Shongwe, M., Tebaldi, C., Weaver, A. J., and Wehner, M.: Long-term Climate Change: Projections, Commitments and Irreversibility, in: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Stocker, T. F., Qin, D., Plattner, G.-K., Tignor, M., Allen, S. K., Boschung, J., Nauels, A., Xia, Y., Bex, V., and Midgley, P. M., pp. 1029–1136, Cambridge University Press, <https://doi.org/10.1017/CBO9781107415324.024>, 2013.
- 730 Edwards, T. L., Fettweis, X., Gagliardini, O., Gillet-Chaulet, F., Goelzer, H., Gregory, J. M., Hoffman, M., Huybrechts, P., Payne, A. J., Perego, M., Price, S., Quiquet, A., and Ritz, C.: Effect of uncertainty in surface mass balance–elevation feedback on projections of the future sea level contribution of the Greenland ice sheet, *The Cryosphere*, 8, 195–208, <https://doi.org/10.5194/tc-8-195-2014>, 2014.
- Edwards, T. L., Brandon, M. A., Durand, G., Edwards, N. R., Golledge, N. R., Holden, P. B., Nias, I. J., Payne, A. J., 3, C. R., and Wernecke, A.: Revisiting Antarctic ice loss due to marine ice-cliff instability, *Nature*, 566, 58–64, <https://doi.org/10.1038/s41586-019-0901-4>, 2019.
- 735 Fettweis, X., Franco, B., Tedesco, M., van Angelen, J. H., Lenaerts, J. T. M., van den Broeke, M. R., and Gallée, H.: Estimating the Greenland ice sheet surface mass balance contribution to future sea level rise using the regional atmospheric climate model MAR, *The Cryosphere*, 7, 469–489, <https://doi.org/10.5194/tc-7-469-2013>, 2013.
- Fettweis, X., Box, J. E., Agosta, C., Amory, C., Kittel, C., Lang, C., van As, D., Machguth, H., and Gallée, H.: Reconstructions of the 1900–2015 Greenland ice sheet surface mass balance using the regional climate MAR model, *The Cryosphere*, 11, 1015–1033, <https://doi.org/10.5194/tc-11-1015-2017>, 2017.
- 740 Fürst, J. J., Goelzer, H., and Huybrechts, P.: Ice-dynamic projections of the Greenland ice sheet in response to atmospheric and oceanic warming, *The Cryosphere*, 9, 1039–1062, <https://doi.org/10.5194/tc-9-1039-2015>, 2015.
- Golledge, N. R., Keller, E. D., Gomez, N., Naughten, K. A., Bernales, J., Trusel, L. D., and Edwards, T. L.: Global environmental consequences of twenty-first-century ice-sheet melt, *Nature*, 566, 65–72, <https://doi.org/10.1038/s41586-019-0889-9>, 2019.
- 745 Gordon, C., Cooper, C., Senior, C. A., Banks, H., Gregory, J. M., Johns, T. C., Mitchell, J. F. B., and Wood, R. A.: The Simulation of SST, sea ice extents and ocean heat transports in a version of the Hadley Centre coupled model without flux adjustments, *Clim. Dyn.*, 16, 147–168, <https://doi.org/10.1007/s003820050010>, 2000.
- Gregoire, L. J., Payne, A. J., and Valdes, P. J.: Deglacial rapid sea level rises caused by ice-sheet saddle collapses, *Nature*, 487, 219–223, <https://doi.org/10.1038/nature11257>, 2012.
- 750 Gregory, J. M. and Huybrechts, P.: Ice-sheet contributions to future sea-level change, *Philos. Trans. R. Soc. London*, 364, 1709–1731, <https://doi.org/10.1098/rsta.2006.1796>, 2006.

- Gregory, J. M., Huybrechts, P., and Raper, S. C. B.: Threatened loss of the Greenland ice-sheet, *Nature*, 428, 616, 2004.
- Gregory, J. M., Browne, O. J. H., Payne, A. J., Ridley, J. K., and Rutt, I. C.: Modelling large-scale ice-sheet–climate interactions following glacial inception, *Clim. Past*, 8, 1565–1580, <https://doi.org/10.5194/cp-8-1565-2012>, 2012.
- 755 Gregory, J. M., Bouttes, N., Griffies, S. M., Haak, H., Hurlin, W. J., Jungclaus, J., Kelley, M., Lee, W. G., Marshall, J., Romanou, A., Saenko, O. A., Stammer, D., and Winton, M.: The Flux-Anomaly-Forced Model Intercomparison Project (FAFMIP) contribution to CMIP6: investigation of sea-level and ocean climate change in response to CO₂ forcing, *Geosci. Model Devel.*, 9, 3993–4017, <https://doi.org/10.5194/gmd-9-3993-2016>, 2016.
- Hanna, E., Mernild, S. H., Cappelen, J., and Steffen, K.: Recent warming in Greenland in a long-term instrumental (1881–2012) climatic
760 context: I. Evaluation of surface air temperature records, *Environ. Res. Lett.*, 7, 045 404, <https://doi.org/10.1088/1748-9326/7/4/045404>, 2012.
- Hofer, S., Tedstone, A. J., Fettweis, X., and Bamber, J. L.: Decreasing cloud cover drives the recent mass loss on the Greenland ice sheet, *Sci. Adv.*, 3, e1700 584, <https://doi.org/10.1126/sciadv.1700584>, 2017.
- Holland, D. M., Thomas, R. H., De Young, B., Ribergaard, M. H., and Lyberth, B.: Acceleration of Jakobshavn Isbrae triggered by warm
765 subsurface ocean waters, *Nature Geosci.*, 1, 659–664, <https://doi.org/10.1038/ngeo316>, 2008.
- Huybrechts, P., Letreguilly, A., and Reeh, N.: The Greenland ice sheet and greenhouse warming, *Palaeogeogr. Palaeocl.*, 89, 399–412, 1991.
- Le clec’h, S., Charbit, S., Quiquet, A., Fettweis, X., Dumas, C., Kageyama, M., Wyard, C., and Ritz, C.: Assessment of the Greenland ice sheet–atmosphere feedbacks for the next century with a regional atmospheric model coupled to an ice sheet model, *The Cryosphere*, 13, 373–395, <https://doi.org/10.5194/tc-13-373-2019>, 2019.
- 770 Levermann, A., Clark, P. U., Marzeion, B., Milne, G. A., Pollard, D., Radic, V., and Robinson, A.: The multimillennial sea-level commitment of global warming, *Proc. Natl. Acad. Sci. USA*, 110, 13 745–13 750, <https://doi.org/10.1073/pnas.1219414110>, 2013.
- Lipscomb, W. H., Fyke, J. G., Vizcaino, M., Sacks, W. J., Wolfe, J., Vertenstein, M., Craig, A., Kluzek, E., and Lawrence, D. M.: Implementation and Initial Evaluation of the Glimmer Community Ice Sheet Model in the Community Earth System Model, *J. Climate*, 26, 7352–7371, <https://doi.org/10.1175/JCLI-D-12-00557.1>, 2013.
- 775 Meehl, G. A., Stocker, T. F., Collins, W. D., Friedlingstein, P., Gaye, A. T., Gregory, J. M., Kitoh, A., Knutti, R., Murphy, J. M., Noda, A., Raper, S. C. B., Watterson, I. G., Weaver, A. J., and Zhao, Z.: Global climate projections, in: *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K. B., Tignor, M., and Miller, H. L., Cambridge University Press, 2007.
- 780 Mouginot, J., Rignot, E., Bjørk, A. A., van den Broeke, M., Millan, R., Morlighem, M., Noël, B., Scheuchl, B., and Wood, M.: Forty-six years of Greenland Ice Sheet mass balance from 1972 to 2018, *Proc. Natl. Acad. Sci. USA*, 19, 9239–9244, <https://doi.org/10.1073/pnas.1904242116>, 2019.
- Muntjewerf, L., Petrini, M., Vizcaino, M., Ernani da Silva, C., Sellevold, R., Scherrenberg, M. D. W., Thayer-Calder, K., Bradley, S. L., Lenaerts, J. T. M., and Lofverstrom, W. H. L. M.: Greenland Ice Sheet contribution to 21st century sea level rise as simulated by the
785 coupled CESM2.1-CISM2.1, *Geophys. Res. Lett.*, 47, e2019GL086 836, <https://doi.org/10.1029/2019GL086836>, 2020.
- Nick, F. M., Vieli, A., Andersen, M. L., Joughin, I., Payne, A., Edwards, T. L., Pattyn, F., and van de Wal, R. S. W.: Future sea-level rise from Greenland’s main outlet glaciers in a warming climate, *Nature*, 497, 235–238, <https://doi.org/10.1038/nature12068>, 2013.
- Noël, B., van de Berg, W. J., van Wessem, J. M., van Meijgaard, E., van As, D., Lenaerts, J. T. M., Lhermitte, S., Kuipers Munneke, P., Smeets, C. J. P. P., van Ulf, L. H., van de Wal, R. S. W., and van den Broeke, M. R.: Modelling the climate and surface mass balance of

- 790 polar ice sheets using RACMO2—Part 1: Greenland (1958–2016), *The Cryosphere*, 12, 811–831, <https://doi.org/10.5194/tc-12-811-2018>, 2018.
- Pattyn, F., Ritz, C., Hanna, E., Asay-Davis, X., DeConto, R., Favier, G. D. L., Fettweis, X., Goelzer, H., 0, N. R. G., Munneke, P. K., Lenaerts, J. T. M., Nowicki, S., Payne, A. J., Seroussi, A. R. H., Trusel, L. D., and van den Broeke, M.: The Greenland and Antarctic ice sheets under 1.5°C global warming, *Nature Climate Change*, 8, 1053–1061, <https://doi.org/10.1038/s41558-018-0305-8>, 2018.
- 795 Reeh, N.: Parameterization of melt rate and surface temperature on the Greenland ice sheet, *Polarforschung*, 59, 113–128, 1989.
- Ridley, J., Huybrechts, P., Gregory, J. M., and Lowe, J. A.: Elimination of the Greenland ice sheet in a high CO₂ climate, *J. Climate*, 18, 3409–3427, <https://doi.org/10.1175/JCLI3482.1>, 2005.
- Ridley, J., Gregory, J. M., Huybrechts, P., and Lowe, J.: Thresholds for irreversible decline of the Greenland ice sheet, *Clim. Dyn.*, 35, 1065–1073, <https://doi.org/10.1007/s00382-009-0646-0>, 2010.
- 800 Roberts, W. H. G., Valdes, P. J., and Payne, A.: Topography’s crucial role in Heinrich events, *Proc. Natl. Acad. Sci. USA*, 111, 16 688–16 693, <https://doi.org/10.1073/pnas.1414882111>, 2014.
- Robinson, A., Calov, R., and Ganopolski, A.: Greenland ice sheet model parameters constrained using simulations of the Eemian Interglacial, *Clim. Past*, 7, 381–396, <https://doi.org/10.5194/cp-7-381-2011>, 2011.
- Robinson, A., Calov, R., and Ganopolski, A.: Multistability and critical thresholds of the Greenland ice sheet, *Nature Climate Change*, 2, 429–432, <https://doi.org/10.1038/NCLIMATE1449>, 2012.
- 805 Rückamp, M., Falk, U., Frieler, K., Lange, S., and Humbert, A.: The effect of overshooting 1.5°C global warming on the mass loss of the Greenland ice sheet, *Earth Syst. Dynam.*, 9, 1169–1189, <https://doi.org/10.5194/esd-9-1169-2018>, 2018.
- Rutt, I. C., Hagdorn, M., Hulton, N. R. J., and Payne, A. J.: The Glimmer community ice sheet model, *J. Geophys. Res.*, 114, F02 004, <https://doi.org/10.1029/2008JF001015>, 2009.
- 810 Sellevold, R., van Kampenhout, L., Lenaerts, J. T. M., Noël, B., Lipscomb, W. H., and Vizcaino, M.: Surface mass balance downscaling through elevation classes in an Earth system model: application to the Greenland ice sheet, *The Cryosphere*, 13, 3193–3208, <https://doi.org/10.5194/tc-13-3193-2019>, 2019.
- Shepherd, A., Ivins, E. R., A. G., Barletta, V. R., Bentley, M. J., Bettadpur, S., Briggs, K. H., Bromwich, D. H., Forsberg, R., Galin, N., Horwath, M., Jacobs, S., Joughin, I., King, M. A., Lenaerts, J. T. M., Li, J., Ligtenberg, S. R. M., Luckman, A., Luthcke, S. B., McMillan, M., Meister, R., Milne, G., Mouginot, J., Muir, A., Nicolas, J. P., Paden, J., Payne, A. J., Pritchard, H., Rignot, E., Rott, H., Sørensen, L. S., Scambos, T. A., Scheuchl, B., Schrama, E. J. O., Smith, B., Sundal, A. V., van Angelen, J. H., van de Berg, W. J., van den Broeke, M. R., Vaughan, D. G., Velicogna, I., Wahr, J., Whitehouse, P. L., Wingham, D. J., Yi, D., Young, D., and Zwally, H. J.: A Reconciled Estimate of Ice-Sheet Mass Balance, *Science*, 338, 1183–1189, <https://doi.org/10.1126/science.1228102>, 2012.
- 815 Shepherd, A., Ivins, E., Rignot, E., Smith, B., van den Broeke, M., Velicogna, I., Whitehouse, P., Briggs, K., Joughin, I., Krinner, G., Nowicki, S., Payne, T., Scambos, T., Schlegel, N., Geruo, A., Agosta, C., Ahlstrøm, A., Babonis, G., Barletta, V. R., Bjørk, A. A., Blazquez, A., Bonin, J., Colgan, W., Csatho, B., Cullather, R., Engdahl, M. E., Felikson, D., Fettweis, X., Forsberg, R., Hogg, A. E., Gallee, H., Gardner, A., Gilbert, L., Gourmelen, N., Groh, A., Gunter, B., Hanna, E., Harig, C., Helm, V., Horwath, A., Horwath, M., Khan, S., Kjeldsen, K. K., Konrad, H., Langen, P. L., Lecavalier, B., Loomis, B., Luthcke, S., McMillan, M., Melini, D., Mernild, S., Mohajerani, Y., Moore, P., Mottram, R., Mouginot, J., Moyano, G., Muir, A., Nagler, T., Nield, G., Nilsson, J., Noël, B., Ootosaka, I., Pattle, M. E., Peltier, W. R., Pie, N., Rietbroek, R., Rott, H., Sørensen, L. S., Sasgen, I., Save, H., Scheuchl, B., Schrama, E., Schröder, L., Seo, K.-W., Simonsen, S. B., Slater, T., Spada, G., Sutterley, T., Talpe, M., Tarasov, L., Jan van de Berg, W., van der Wal, W., van Wessem, M., Vishwakarma, B. D.,
- 825

- Wiese, D., Wilton, D., Wagner, T., Wouters, B., and Wuite, J.: Mass balance of the Greenland Ice Sheet from 1992 to 2018, *Nature*, 579, 233–239, <https://doi.org/10.1038/s41586-019-1855-2>, 2020.
- Smith, R. S.: The FAMOUS climate model (versions XFXWB and XFHCC): description update to version XDBUA, *Geosci. Model Devel.*, 5, 269–276, <https://doi.org/10.5194/gmd-5-269-2012>, 2012.
- 830 Smith, R. S., Osprey, A., and Gregory, J. M.: A description of the FAMOUS (version XDBUA) climate model and control run, *Geosci. Model Devel.*, 1, 53–68, <https://doi.org/10.5194/gmd-1-53-2008>, 2008.
- Smith, R. S., George, S., and Gregory, J. M.: FAMOUS version xotzb (FAMOUS-ice): a GCM capable of energy- and water-conserving coupling to an ice sheet model, <https://doi.org/10.5194/gmd-2020-207>, submitted.
- 835 Solgaard, A. M. and Langen, P. L.: Multistability of the Greenland ice sheet and the effects of an adaptive mass balance formulation, *Clim. Dyn.*, 39, 1599–1612, <https://doi.org/10.1007/s00382-012-1305-4>, 2012.
- Solgaard, A. M., Bonow, J. M., Langen, P. L., Japsen, P., and Hvidberg, C. S.: Mountain building and the initiation of the Greenland Ice Sheet, *Palaeogeogr. Palaeoclimatol.*, 392, 161–176, <https://doi.org/10.1016/j.palaeo.2013.09.019>, 2013.
- Stouffer, R. J., Yin, J., Gregory, J. M., Dixon, K. W., Spelman, M. J., Hurlin, W., Weaver, A. J., Eby, M., Flato, G. M., Hasumi, H., Hu, A., 840 Jungclaus, J., Kamenkovich, I. V., Levermann, A., Montoya, M., Murakami, S., Nawrath, S., Oka, A., Peltier, W. R., Robitaille, D. Y., Sokolov, A., Vettoretti, G., and Weber, N.: Investigating the Causes of the Response of the Thermohaline Circulation to Past and Future Climate Changes, *J. Climate*, 19, 1365–1387, <https://doi.org/10.1175/JCLI3689.1>, 2006.
- Tedesco, M., Fettweis, X., Mote, T., Wahr, J., Alexander, P., Box, J. E., and Wouters, B.: Evidence and analysis of 2012 Greenland records from spaceborne observations a regional climate model and reanalysis data, *The Cryosphere*, 7, 615–630, [https://doi.org/10.5194/tc-7-](https://doi.org/10.5194/tc-7-615-2013) 845 615-2013, 2013.
- Tedesco, M., Doherty, S., Fettweis, X., Alexander, P., Jeyaratnam, J., and Stroeve, J.: The darkening of the Greenland ice sheet: trends, drivers, and projections (1981–2100), *The Cryosphere*, 10, 477–496, <https://doi.org/10.5194/tc-10-477-2016>, 2016.
- Toniazzo, T., Gregory, J. M., and Huybrechts, P.: Climatic impact of a Greenland deglaciation and its possible irreversibility, *J. Climate*, 17, 21–33, 2004.
- 850 Trusel, L. D., Das, S. B., Osman, M. B., Evans, M. J., Smith, B. E., Fettweis, X., McConnell, J. R., Noël, B. P. Y., and van den Broeke, M. R.: Nonlinear rise in Greenland runoff in response to post-industrial Arctic warming, *Nature*, 564, 104–108, <https://doi.org/10.1038/s41586-018-0752-4>, 2018.
- van Angelen, J. H., Lenaerts, J. T. M., van den Broeke, M. R., Fettweis, X., and van Meijgaard, E.: Rapid loss of firn pore space accelerates 21st century Greenland mass loss, *Geophys. Res. Lett.*, 40, 2109–2113, <https://doi.org/10.1002/grl.50490>, 2013.
- 855 van den Broeke, M. R., Enderlin, E. M., Howat, I. M., Kuipers Munneke, P., Noël, B. P. Y., van de Berg, W. J., van Meijgaard, E., and Wouters, B.: On the recent contribution of the Greenland ice sheet to sea level change, *The Cryosphere*, 10, 1933–1946, <https://doi.org/10.5194/tc-10-1933-2016>, 2016.
- van Vuuren, D. P., Edmonds, J., Kainuma, M., Riahi, K., Thomson, A., Hibbard, K., Hurtt, G. C., Kram, T., Krey, V., Lamarque, J.-F., Masui, T., Meinshausen, M., Nakicenovic, N., Smith, S. J., and Rose, S. K.: The representative concentration pathways: an overview, *Clim. Change*, 109, 5–37, 2011.
- 860 Vizcaíno, M.: Ice sheets as interactive components of Earth System Models: progress and challenges, *WIREs Climate Change*, <https://doi.org/10.1002/wcc.285>, 2014.
- Vizcaíno, M., Mikolajewicz, U., Jungclaus, J., and Schurgers, G.: Climate modification by future ice sheet changes and consequences for ice sheet mass balance, *Clim. Dyn.*, 34, 301–324, <https://doi.org/10.1007/s00382-009-0591-y>, 2010.

- 865 Vizcaíno, M., Lipscomb, W. H., Sacks, W. J., van Angelen, J. H., Wouters, B., and van den Broeke, M. R.: Greenland Surface Mass Balance as Simulated by the Community Earth System Model. Part I: Model Evaluation and 1850–2005 Results, *J. Climate*, 26, 7793–7812, <https://doi.org/10.1175/JCLI-D-12-00615.1>, 2013.
- Vizcaíno, M., Lipscomb, W. H., Sacks, W. J., and van den Broeke, M.: Greenland surface mass balance as simulated by the Community Earth System Model. Part II: Twenty-first-century changes, *J. Climate*, 27, 215–226, <https://doi.org/10.1175/JCLI-D-12-00588.1>, 2014.