

Dear Anonymous Reviewer 1, Dear Editor,

We would like to thank the reviewer for the insightful comments and good feedback provided on our manuscript. Below we provide answers to each comment. The original comments are numbered in **red**, and written in **bold**. Our answers are provided in black, and the manuscript text including the proposed changes is provided in *“quoted italics”*. For an easier comparison between this response letter and the original manuscript, we refer to the figures by their original numbering, despite new figures being presented here (and proposed for inclusion). New figures are presented and referenced in this letter as “new figure”, and their numbering will be adjusted in the revised manuscript.

### **Major issues:**

**1. This study uses an SSA model with a minimum ice thickness to simulate shallow ice flow around high-relief obstacles. In principle, I can see how this model is appropriate, but it is a fairly non-standard use of SSA. For one, it is unclear to what extent SSA applies for ice that changes from kilometer-scale thickness to meter scale thickness. Further, even if the exposure threshold you use is above the minimum imposed modeled thickness, the use of a minimum in of itself will potentially affect how ice flow occurs near the ice margin on the Nunatak (i.e. constant thickness at the minimum over space vs. gradually decreasing thickness over space). There are also missing details regarding the stress and flux boundary conditions at these sorts of margins. At a minimum, the model descriptions needs more details and justification for why we should believe SSA in such a situation (as opposed to, say, Full Stokes). It may also help to investigate how the results (particular in terms of simulated exposure timing) depend on the choices of minimum model ice thickness and the boundary conditions therein.**

We agree it is a non-standard use of SSA. Its use is justified by the lower computational cost of this approximation compared to a full-Stokes model, the target settings we aim to represent, and the model's dynamic adaptive meshing capability. We also agree that full-Stokes models would yield more accurate results, and had highlighted this in our original manuscript (L87). Based on the comments and suggestions above and from reviewers 2 and 3, we have expanded Sect. 2 to better justify the validity of the SSA approximation, clarify the boundary conditions used in our model, as well as its limitations compared to a full-Stokes model.

*“To better understand how ice flow interacts with the steep topography of nunataks and what impact this interaction has on ice surface elevation patterns, we perform a suite of numerical simulations using an idealised setup. We use Úa (Gudmundsson, 2020), an ice flow model that solves the shallow*

shelf, also referred to as shelfy stream approximation (SSA or SStA) of the Stokes equations (Cuffey and Patterson, 2010) on a horizontal, finite element mesh. Úa has been successfully applied to model the ice flow of idealised (e.g. Gudmundsson et al., 2012), modern (e.g. Miles et al., 2021), and palaeo-ice streams (e.g. Jones et al., 2021). Úa solves the ice surface and momentum equations simultaneously, and its finite element formulation allows for an adaptive mesh refinement in areas of particular interest, such as where the ice shallows around nunataks. For the modelled domain (which we describe below), an unstructured mesh was generated, which is refined during simulation time (including during spin up) based on a series of glaciological refinement criteria. Element size is refined according to ice thickness, from a maximum of 8 km down to 205 m around the interface between ice and nunatak, where ice thickness approaches the minimum (which we set as 1 m). The mesh is also refined (to 500 m) in a 4 km buffer zone centered at the grounding line. The mean element size after spin up was 740 m (414 m median).

In the vicinity of nunataks, where the ice is thinnest, the ice flow regime has a relevant vertical component, which is not captured by the SSA approximation. A more accurate representation of the ice dynamics for such regions can be achieved with full-Stokes models. However, such models are currently too computationally demanding when adopted for long simulation periods and multiple experiments (e.g. Schannwell et al., 2020). Over the thinnest-ice areas, the horizontal scales resolved in our model are of the order of a few hundred metres (commonly noted as  $O(10^2)$  m), while the vertical scale is of the order of a few metres ( $O(10^0)$  m). This yields an aspect ratio of  $O(10^{-2})$ , which falls within the range where shallow approximations are applicable ( $O(\leq 10^{-2})$ , e.g. Fowler and Larson, 1978). To our knowledge, no intercomparisons between simplified and full-Stokes models exist that focus on the representation of thickness gradients. At least within the context of MISMIP+ (Cornford et al., 2020), there are little variations between SSA (including the model used in our study), L1Lx, and higher-order models regarding simulated ice retreat. Full-Stokes models that participated in the MISMIP+ experiment also agreed with the simplified-physics models, indicating that other models should behave similarly. Instead, the MISMIP+ experiments highlighted the importance of the formulation of the sliding law (i.e., Weertman versus Coulomb-limited). The two sliding laws strongly differed in their rates of grounding line retreat, but it has been shown that such differences decrease with increasing spatial resolution (i.e., model grid/mesh refinement) at the grounding line (Gladstone et al., 2017). Here, we adopt a Weertman law for basal sliding and the required refinement at the grounding line to minimise sliding law issues.

At the free upper surface, the streamline upwind Petrov-Galerkin (SUPG) method is applied, which ensures model stability over regions of pronounced ice surface topography (Wirbel and Jarosch, 2020). In our experiments, the ice surface topography is steepest when the ice front retreats from the downstream end of the domain and in the vicinity of nunataks. When ice thins below the prescribed minimum (which we set to 1 m), the model uses the active-set method (Durand et al., 2009; Gudmundsson et al., 2012). In this method, violated ice thickness constraints are activated using the Lagrange multiplier approach (Ito and Kunisch, 2008), which is applied to the momentum equation,


*and ensures a better representation of the ice dynamics compared to simply resetting the thickness to the prescribed minimum value, as is commonly used in finite element models.*

*The model domain (Sect. 2.1) and spin up procedure (Sect. 2.2) are the same for all simulations. In a first set of simulations, we evaluate changes in ice surface elevation up and downstream of a single nunatak under three different thinning scenarios (Sect. 2.3). We then use the forcing that provides the highest ice-thinning rates to evaluate the impact of multiple nunataks and the width of glaciers between them on ice surface elevation patterns (Sect. 2.4). Finally, we repeat the last experiments for a series of regular meshes (without refinement) at horizontal resolutions commonly used in ice sheet models (Sect. 2.5). This final set of experiments assesses how well different grid resolutions resolve changes in ice surface elevation across steep marginal topography under thinning scenarios, and their implications for simulations of past ice surface elevations.”*


Finally, we agree that performing sensitivity tests for the minimum ice thickness is indeed helpful. We therefore carried out the same simulations as presented in Fig. 5, but adopted minimum thicknesses of 5 m and 0.5 m. We have added a new supplementary figure showing these results, where the sensitivity to a “crevassed” surrounding of the nunatak, as suggested by Reviewer 2 (comment 2), is also included. These new experiments are referenced in the main text when discussing our results regarding the difference in time of exposure:

*“The ice surface steepening and consequent mismatch between the up and downstream sides of the nunatak increases as the ice thins, until the downstream side becomes exposed. Exposure happens earlier downstream, as expected due to lower ice surface elevation, and an equidistant point upstream becomes exposed (or has its thinning stabilised) up to 14 kyr later than its downstream counterpart. The rates of thinning, and consequently the timing of bedrock exposure, are dependent on the choice of the basal sliding coefficient. An increase of ca. 50 % in ice thinning was observed between the higher sliding and control experiments, as well as between the control and lower sliding experiments (not shown). This pattern is expected given the influence of basal sliding on the initial ice sheet geometry (Fig. S5), and highlights that the exposure lags between up and downstream faces of a nunatak observed in the real world will be site dependent. The choice of minimum ice thickness allowed in our model (1 m) also influences the timing of exposure at a given point on the nunatak ([ref. to the new supplementary figure below]). A lower minimum thickness increased lag times, while a higher minimum thickness reduced them. The surroundings of nunataks are often crevassed, which results in a change in ice rheology. Hence, another test was carried out where the prescribed ice rheology becomes progressively softer towards the nunataks by a factor of ten ([ref. to the new supplementary figures below]). In this test, the most notable effect from the choice of ice rheology was a delay in the timing of exposure downstream of the nunatak relative to the control case, by 0.5-*

5 kyr, but the lag times were still of the same magnitude as the control case (i.e., between 2 and 10 kyr). While we use stabilisation of thinning upstream to determine..."



**New Supplementary Figure:** Spatial distribution of the rheology factor  $A$  for the 'crevassed' experiment ([ref. to new figure below]). The value for  $A$  away from the nunatak is that used in all other experiments ( $\log_{10}(A) \sim -8.5$ ).



**New Supplementary Figure:** As in Fig. 5 of the main text, but for minimum ice thickness experiments where the minimum thicknesses allowed in the model are 0.5 and 5 m, and a

“crevassed” experiment, where the ice is softer around the nunatak compared to the rest of the model domain (ref to Fig, above).

#### References not previously in the manuscript:

- Cornford, S.L., et al. (2020). Results of the third Marine Ice Sheet Model Intercomparison Project (MISMIP+). *The Cryosphere*, 14, 2283-2301.
- Durand, G., Gagliardini, O., De Fleurian, B., Zwinger, T., & Le Meur, E. (2009). Marine ice sheet dynamics: Hysteresis and neutral equilibrium. *Journal of Geophysical Research: Earth Surface*, 114, F03009.
- Fowler, A. C., & Larson, D. A. (1978). On the flow of polythermal glaciers-I. Model and preliminary analysis. *Proceedings of the Royal Society of London. A. Mathematical and Physical Sciences*, 363(1713), 217-242.
- Ito, K., & Kunisch, K. (2008). *Lagrange multiplier approach to variational problems and applications*. Society for Industrial and Applied Mathematics.
- Wirbel, A., & Jarosch, A. H. (2020). Inequality-constrained free-surface evolution in a full Stokes ice flow model (*evolve\_glacier v1.1*). *Geoscientific Model Development*, 13, 6425-6445.

**2. I come away from reading this study mostly convinced that there are potential issues with the way cosmogenic exposure ages are interpreted. However, I would also have liked to see a bit more concrete solutions/suggestions to solve this problem. Is the answer that every time someone measures a cosmogenic exposure ages, they will have to do detailed high-resolution modeling of ice flow in the region from that time period to place the specific measurements in context? This seems like quite a lot of work. Two less onerous possibilities (only one of which is currently mentioned in the manuscript) intrigue me: (1) can already-measured exposure ages be corrected using the results of this study? What would be necessary to do so? Does one need multiple ages from different parts of the same Nunatak to do such a correction? (2) It seems like you suggest in the discussion that sampling Nunatak at locations that are perpendicular to flow (i.e. on the "sides") would mitigate the age bias somewhat. However, most of your discussion of your experiments was focused on the difference between upstream and downstream. Overall, I think the focus of the discussion needs to be shifted towards providing potential ways forward on a solution, particularly for those who may not have the capability to do detailed simulations in support for exposure ages.**

We are reluctant to provide absolute values for corrections because regional settings such as mass balance, bedrock slope, and ice stream/glacier widths exert significant influence in the ice change history. Nevertheless, we have rewritten some parts of the discussion based on the proposed suggestions (detailed below), highlighting the recommendations that were already provided, and also the ways we believe the issues raised in our study can be best addressed:

- Sampling at nunatak flanks. Also addressing point 34, we have further included a comparison of cosmogenic-nuclide ages taken at the nunatak flanks (see new supplementary figure below), which complements Fig. S9, and refer to it in the same sentence:

*“We assess the degree to which our model results are reflected in empirical data by analysing all current cosmogenic nuclide exposure ages from nunataks in Antarctica (Balco, 2021, Heyman, 2021) that indicate ice recession since the Last Glacial Maximum (last 21 kyr). Choosing this time interval minimises the effect of inherited concentrations from prior exposure, and confirms that samples taken upstream and downstream at the same elevation interval show significant age differences. This pattern exists regardless whether results from  $^{10}\text{Be}$  (most ubiquitous) or  $^{14}\text{C}$  (least inheritance) are considered (Fig. S9). Age gradients are not as pronounced for upstream and downstream samples taken at nunatak flanks ([ref. to new supplementary figure below]).”*

- Sampling at areas where model predictions of lags can be tested would be useful for validating our results. For unpublished already-collected samples, we highlight that their position relative to flow should be reported (originally mentioned in L371 – 375). We have expanded on this sentence to draw the attention to the fact that the sample placement relative to ice flow should be considered when interpreting the data:


*“Such experiments would aid the validation and interpretation of our findings. Regardless, we recommend that in addition to reporting cosmogenic nuclide concentrations, apparent ages, and uncertainties, the sample location relative to ice flow near the nunatak should be specified. This additional information would be crucial in the interpretation of the local ice sheet history, and would contribute to a domain-wide comparison of field constraints and modelling results.”*

- Also, we see our paper as a call for closer collaboration between field scientists and ice sheet modellers in designing sampling strategies and interpreting results. We present this last point explicitly towards the close of our discussion section:

*“In summary, we advocate for a closer collaboration between ice sheet modellers and field scientists. This would allow for common goals to be defined and for both disciplines to help each other overcome their respective inherent limitations.”*

And in the end of our conclusions:

*“This should be achievable by a closer collaboration between ice sheet modellers and field scientists.”*



**New Supplementary Figure.** Sample location relative to flow and age distribution for  $^{10}\text{Be}$  (upper row) and  $^{14}\text{C}$  (lower row) as in Fig. S9, but considering only the samples taken at the nunatak flanks.

**3. I found the text of section 3 very dense with descriptions of results. Rather than re-describe a lot details that are in the figures, making more effort to synthesize results and describe their physical meaning would be welcome. Also, generally shortening section 3 would help as well.**

Wherever possible in section 3, we have made changes to be more concise in reporting our results, highlighting their physical meaning where they were not clear. However, we did not focus on them, as it would cause significant overlap with the discussion section, where we believe such focus is more fitting, and where we contextualise our results.

**Minor issues:**

**4. Throughout: new results should be described using the present tense, whereas prior studies should be described using past tense. There is some inconsistency throughout on this.**

We have revised the manuscript and changed accordingly.

**5. Line 15: while rapid ice flow through outlet glaciers...**

we used “efficient ice drainage” to highlight the fact that a constricted flow also contributes to the formation of a steeper ice surface elevation gradient, i.e., it is not necessarily the ice flow velocity that controls it.

**6. Line 15: what do you mean by "alleviated the differential response" - this is confusing**

we have rephrased the end of the sentence, which now reads:

*“A nunatak elongated transverse to ice flow, with an extensive subglacial reach, is able to increase ice retention and therefore impose steeper ice surface gradients, while efficient ice drainage through outlet glaciers produces gentler gradients.”*

**7. Line 23-43: These two paragraphs are quite general and can probably be shortened to one paragraphs**

We have merged both paragraphs, shortening the content by 25% without the loss of important information. We believe it is important to acknowledge the different time scales at which ice sheet modellers work, highlight the most recent improvements in ice sheet modelling, and show that a good representation of complex bedrock topography remains a knowledge gap. The first paragraph of the manuscript now reads:

*“Ongoing changes in climate are already causing significant mass loss and ice-margin retreat of both the Antarctic and Greenland ice sheets (Garbe et al., 2020; King et al., 2020). Near-future (2100 CE) projections of sea level rise point to ocean thermal expansion as the main cause (Oppenheimer et al., 2019), but over multi-centennial timescales, the sea level contribution from Antarctica is expected to become dominant (Pattyn and Morlighem, 2020). Numerical ice sheet modelling efforts are aimed at reducing uncertainty by better understanding the processes that lead to sea level rise, focusing on both shorter (Goelzer et al., 2020; Seroussi et al., 2020), and longer timescales (Pollard and DeConto, 2009; Albrecht et al., 2020). Recent efforts include improvements in key model components such as grounding line dynamics (e.g. Gladstone et al., 2017; Seroussi and Morlighem, 2018), coupling to solid Earth and sea level models (e.g. Gomez et al., 2020), and improved treatment of ice-ocean interaction processes (e.g. Reese et al., 2018; Kreuzer et al., 2020). The importance of bedrock topography (Morlighem et al., 2020) and grid resolution (Durand et al., 2011) have been acknowledged previously, and studied particularly for marginal regions of the ice sheet (e.g. Sun et al., 2014; Robel et al., 2016; Favier et al., 2016). Spatial variations in bedrock topography (Robel et al., In Review) and the resulting basal and lateral drag exerted at the ice-bedrock interface for different spatial scales (Jamieson et al., 2012, 2014; Åkesson et al., 2018; Jones et al., 2021), can slow down or even stabilise grounding line retreat. Near-marginal regions with large subglacial topographic relief, such as the overridden mountain ranges that fringe the glaciated cratons of Greenland and East Antarctica (Howat et al.,*

2014; Burton-Johnson et al., 2016), therefore require suitable consideration when evaluating ice loss beyond this century.”

**8. Line 27: ice loss from marine-terminating outlet glaciers**

We have changed it accordingly.

**9. Line 40: drag is exerted on fjords?**

*This statement has been removed after the suggestions from comment 7*

**10. Line 56: setting where rock samples are acquired**

We have added “rock samples” accordingly.


**11. Line 69: that there is no systematic approach to selecting the sampling**

We have changed accordingly.

**12.** The following comments are grouped together for simplicity:

- **Figure 2: is the coloring of the bars necessary? The information seems redundant with the radial axis.**
- **Figure 2: later in the discussion you seemingly suggest that you might also have information on ages for each of these samples. If so, it would seem to make sense (or at least be interesting) to incorporate this extra data in this figure. Even if there is sampling bias, it would be interesting to show if there was a systematic difference in exposure ages in upstream and downstream samples. However, if this data is not available, that's not an issue.**

We have removed the colouring, and added a dashed line to help identify the threshold for considering a sample to be upstream or downstream. This was also done to the supplementary figures, for consistency. We have also added two panels to Fig. 2, showing the kernel density plots for  $^{10}\text{Be}$  and  $^{14}\text{C}$  ages considering all samples, as suggested:



**Figure 2.** (a) Polar histogram showing the location of cosmogenic  $^{10}\text{Be}$  and  $^{14}\text{C}$  samples from boulders in Antarctica (Heyman, 2021; Balco, 2021) with ages younger than the Last Glacial Maximum, relative to the nearest nunatak summit and its adjacent ice flow direction ( $n = 191$ ; sample duplicates were excluded). The difference in direction was computed between a sample position relative to the nunatak summit (identified in BedMachine-Antarctica; Morlighem et al., 2020) and ice flow (Mouginot et al., 2012) near the nunatak summit. Summits were identified through a morphological feature map (Wood, 1996; see supplementary material). The area of each slice is proportional to the number of samples within that category, and each category spans a  $15^\circ$  arc. In this figure,  $0^\circ$  ( $180^\circ$ ) implies that the sample was taken downstream (upstream), directly aligned with the ice flow. (b,c) kernel density function of  $^{10}\text{Be}$  (b) and  $^{14}\text{C}$  (c) apparent exposure ages from samples shown in panel (a). Dashed lines show the median age, and shading shows the uncertainty interval based on the median uncertainty of their respective ages.

**13. Line 78: To perform these tests**


We have changed accordingly.

**14. Line 104: how is the ice front simulated?**


The ice front is treated as an equilibrium between the pressure exerted by the ice and the ocean (set at 0 m, as shown by the dashed line in Fig. 3b). The ice front is considered to retreat when the thickness reaches the prescribed minimum thickness, for which the active-set method is used. This is now included in the manuscript, see answer to comment 1.

**15. Line 118: I felt at this point like it would be helpful to have a figure just showing the bed geometries used in the simulations because it was hard to visualize from the text alone**

We have added a new figure showing bed elevation for both nunatak shapes. We added the profile lines and points used in Fig. 4, and removed its inset plots. We believe this change has made Fig. 4 cleaner, and addresses the issue raised in comment 25 regarding the meaning of the different lines.



**New Figure.** Bedrock elevation (in metres above sea level, m a.s.l.) used in the experiments where the nunatak is placed (a) transverse and (b) parallel to ice flow. Ice flow in this figure is from left to right. Transect lines show the position of the transects presented in Fig. 4a,b, and coloured circles show the locations for the ice surface evolution analysis presented in Fig. 4c,d following the same colours as the lines therein.



**Figure 4.** Surface elevations after 0, 10, and 20 kyr for the ‘thw’ experiment with a nunatak elongated (a) transverse and (b) parallel to ice flow at 0, 10, and 15 km from the centreline; dark grey lines show the respective ice surface elevation for experiments without a nunatak; (c,d) evolution of the ice surface elevation difference between six pairs of equidistant points up and downstream from the nunatak, along the centre line for the experiments showcased in (a,b) respectively. In short, this figure shows to which extent a single nunatak is able to influence ice surface elevation in space (along and perpendicular to flow) and through time, how ice surface elevation is linked to a nunatak’s subglacial topography, and how the ice surface elevation mismatch evolves differently depending on the distance from the nunatak.

## 16. Line 121: idealized nunatak dimensions

We have changed accordingly.

**17. Line 122: are constrained by 33 real nunatak topographic profiles from Antarctica**

We have changed accordingly.

**18. Line 128: is there any expectation that your results are dependent on the sliding law, or coefficient chosen? Would more rapid sliding give a drastically different result?**

The fact that the sliding coefficient used was chosen to match that of an inversion, which also used Weertman's sliding law, prevents us from reliably choosing an equivalent coefficient for a different sliding law. However, the greatest differences between Coulomb-limited and Weertman-type laws occur in the vicinity of the grounding line. Given our mesh refinement over such regions, these differences are expected to be small (see Gladstone et al., 2017). This is now mentioned in our updated Sect. 2 (see response to comment 1).

We have also provided a comparison of total ice thinning, as opposed to time of exposure, for the different-sliding 'thw' scenario experiments, for which the initial states were originally presented in the supplement. The choice for ice thinning is motivated by the fact that their initial ice surface elevation is significantly different from one another. We mention it in the discussion (Sect. 4.1, second paragraph; see response to comment 1).

**19. Equation 2: is  $L_x = L \cdot x$  or  $L_x$ ?**

It is  $L_x$ . Thank you for spotting this typo.

**20. Line 145-146: confusing sentence**

We have slightly rephrased it as:

*"The average ice surface slope between the ice divide and the grounding line (which after spin up was located at  $x = 136$  km; Fig. 3b), is  $\sim 1.3$  %."*

**21. Section 2.5: here and elsewhere you say that typical paleo-ice sheet mode configurations are  $>5$  km. For ice sheet wide simulations, sure, this is true, but I think for individual catchments, there are plenty of people doing  $\sim 1$  km and higher resolution simulations, even for paleo simulations. Its pretty well within the realm of current SSA models to simulate such resolution on a spatially limited basis (even without adaptive meshing or other sophisticated techniques) for 10's of kyr.**

That is correct, thanks for highlighting this point. We meant indeed ice sheet-wide simulations, and have clarified it in the sentence:

*"In a final series of sensitivity experiments, we assess how well regular-spaced grids of coarser resolutions typically used in continental-scale ice sheet models (5, 10, and 20 km) resolve the ice surface elevation pattern around nunataks compared to the solution using an unstructured, locally refined mesh."*

**22. Line 190-193: I'm a bit confused because the ice intersects "above" the lowest ice surface elevation both upstream and downstream. Either this is a mistake, or more explanation is needed because it is counterintuitive that ice thickness would be higher downstream.**

Based on this comment and comment 3, we have rewritten the sentence to be more concise:

*"At the start of the thinning experiments where the nunatak is elongated transverse to flow, the effect of the nunatak on the ice surface elevation is seen up to 30 km away from its summit along flow, and 15 km perpendicular to flow. The ice surface directly upstream of the nunatak is 360 m above the general ice sheet surface (considered to be the elevation 15 km away from the centreline), while downstream the lowest ice surface elevation (located 3 km downstream of the summit) is 100 m lower than the general ice surface elevation."*

We have also provided a brief discussion regarding possible explanations for this phenomenon in the end of the first paragraph of Sect. 4.1:

*"A relative increase in the ice surface elevation was observed immediately downstream of the nunatak, while the lowest elevation attained was located ~3 km downstream of the nunatak summit at the start of the simulations. Although a similar effect can be observed on the lee side of isolated nunataks in Dronning Maud Land, MacRobertson Land, and the Transantarctic Mountains (Howat et al., 2019), we cannot rule out that this is a numerical effect, caused by artificial advection of ice downstream from the regions where the minimum thickness constraint is violated and thus numerically modified (see Jarosch et al., 2013)."*

**23. Line 199-201: confusing sentence**

We have rephrased the sentence, which now reads:

*"In both cases, changes in ice surface elevation perpendicular to the centreline, caused by the nunatak presence, stand out from the general ice surface elevation for the entire extent of the subglacial nunatak obstruction (cf. Figs. 4a,b, and [ref. to new figure showing the bed elevation]a,b))."*

**24. Line 213: this connection between slope and exposure is not very clear. Could be described more**

We have rewritten the mentioned sentences, also considering comment 3. This part now reads:

*“Both ice thinning experiments detailed in Fig. 4a,b reveal an overall steepening of the ice surface gradient during the 20 kyr thinning period (profiles from red to green). The evolution of this steepening is illustrated in Fig. 4c,d, where points equidistant from the nunatak summit (upstream and downstream) show surface elevation differences that increase through time. For equidistant locations closer to the nunatak summit (< 2.5 km and < 7.5 km for Figs. 4c and d, respectively) the increase in steepening (i.e., in elevation difference) is disrupted when the downstream location becomes exposed, since its elevation no longer changes while the equivalent point upstream is still ice covered and thinning.”*

**25. Figure 4: it is not clear what all the different lines meaning**

We have removed the insets, and put the lines in the bed elevation figure, requested in comment 15.

**26. Figure 4: why does it appear like the "idealized" nunatak (brown in panels a and b) is bumpy? I'm guessing a plotting artifact, but its a bit distracting**

It was indeed a plotting artifact, and we thank the reviewer for spotting that. We believe we have improved it in the updated plot, as shown in our answer to comment 15, where we presented the changes to Figure 4.

**27. Figure 6: difference with respect to what?**

Figure 6 shows the difference between the three-nunatak experiments with respect to the one-nunatak control case, shown in panel (e). We have rephrased the caption to clarify this:

*“(a-d) Difference in surface elevation (m) between each three-nunatak ‘thw’ experiment of varying glacier widths (0, 5, 10, 15 km, respectively) with respect to the one-nunatak ‘thw’ control run (shown in panel e for reference), after 20 kyr of simulation”*

**28. Line 287: with-width**

Typo fixed accordingly.

**29. Line 294: response to what?**

To the nunataks. We have rephrased the sentence for clarity:

*“Our modelling experiments demonstrate that the magnitude of the ice surface elevation response due to the presence of nunataks is proportional to their ability to obstruct or constrict ice flow.”*

**30. Line 299: In their fjord experiment, Frank**

We have changed accordingly.

**31. Line 309: between ice thickness on the up and downstream sides o the nunatak increases...**


We have changed accordingly.

**32. Line 327-328: would this add or subtract to the effect from flow around the nunatak? Could the whole exposure age discrepancy be compensated by this, or made even more severe? Seems like its worthy of a test in your model.**

We have performed two additional simulations, where the SMB profile is smoother at the upstream side of the domain, and decays rapidly downstream, through the following equation (cf. Eq. 2 of the main text):

$$SMB_0 - (SMB_0 - SMB_e) * \tan(|x|/L_x) * \sin^2(|x|/L_x) + b(t)$$

We prescribe two different values for  $SMB_0$ ,  $-0.92 \text{ ma}^{-1}$  and  $-0.78 \text{ ma}^{-1}$ , and call the respective experiments 'margin1' and 'margin2'. While 'margin1' ensures that the integrated SMB at  $t=0$  kyr is the same as in the 'thw' scenario (referred here as 'control'), 'margin2' is designed so that the SMB upstream is close to zero. The new figure below shows how ice thinning is impacted by this different SMB curve, as in Fig. 4a of the manuscript. Overall, the smoother profile at the nunatak vicinity caused the surface elevation differences up and downstream to be smaller than in the 'control' run, although the general surface elevation is lower. Thus, the differences in ice surface elevation seem to be more sensitive to the gradient in SMB rather than the absolute values themselves.



**New Supplementary Figure.** (a) different surface mass balance (SMB) profiles analysed. Solid lines show the profile at the start of the simulations ( $t = 0$  kyr), while dashed lines show the profile at the end ( $t = 20$  kyr). The ‘thw’ profile (see Fig. 3a of the main text) is shown here as ‘control’. (b – d) surface elevation profiles after 0, 10, and 20 kyr for each experiment referred in panel (a), as in fig. 4a of the main text. Grey lines show the same “no nunatak” experiment as in Fig. 4a for an easier comparison between all profiles.

We have added the above information to the supplementary material, and refer to it in the main text, adding to the SMB discussion highlighted by the reviewer (Sect. 4.1, last paragraph), as follows:

*“A final set of sensitivity experiments ([ref. to new supplementary figure shown above]) shows that the surface elevation difference between locations equidistant up and downstream of a nunatak is more sensitive to the spatial gradient of prescribed SMB than to the absolute values themselves.”*

**33. Line 338: where does the 400 m figure come from?**

It comes from the ice surface elevation difference at the ice-bedrock interface, and the general ice surface unaffected by the nunatak. We have referred to Fig. 4 to make it clearer.

**34. Line 361-364: I referenced this above, but this point seems very important and could do with more references to results, since you should be able to show this in your simulations**

We have added a reference to Fig. 4, where it is possible to compare the patterns of ice surface elevation to the sentence in question:

*“The areas of nunataks located perpendicular to ice flow (i.e., their flanks), where the subglacial signature of a nunatak is not as pronounced, are likely to provide more accurate estimates of regional ice sheet thickness change, as these areas are less impacted by the differences in ice surface steepening (cf. dashed coloured lines and dotted black line in Fig. 4a,b, and their respective locations in Fig. 2)”*

We believe that this, combined with the other changes listed under our response to comment [2](#), have addressed this remark.

We thank the reviewer for the thorough and insightful review of our manuscript, and look forward to the opportunity to present a significantly updated manuscript based on the revisions outlined above.