

# Evaporation over a glacial lake ~~glacial lakes~~ in Antarctica

Elena Shevnina<sup>1</sup>, Miguel Potes<sup>2</sup>, Timo Vihma<sup>1</sup>, Tuomas Naakka<sup>1</sup>, Pankaj Ramji Dhote<sup>3</sup>, Praveen Kumar Thakur<sup>3</sup>

<sup>1</sup> Finnish Meteorological Institute, Helsinki, Finland-

<sup>2</sup> Institute Earth Science, Evora, Portugal

5 <sup>3</sup> Indian Institute of Remote Sensing, Dehradun, India

Correspondence: [elena.shevnina@fmi.fi](mailto:elena.shevnina@fmi.fi)

**Abstract.** The study provides estimates of summertime evaporation over a glacial lake located in the Schirmacher oasis, Dronning Maud Land, East Antarctica. Lake Zub/Priyadarshini is the second largest lake in the oasis, and its maximum depth is 6 m. The lake is among the warmest glacial lakes, and it is free of ice during almost two summer months. The summertime evaporation over the ice-free lake was measured using the eddy covariance (EC) method, and estimated on the basis of the bulk-aerodynamic method and four combination equations. We used meteorological and hydrological measurements collected during a field experiment carried out in 2018. The EC method was considered the most accurate, and the evaporation was estimated to be 114 mm for the period from 1 January to 7 February 2018 (38 days) on the basis of this method. The average daily evaporation was estimated to be 3.0 mm day<sup>-1</sup> in January 2018. The largest changes in daily evaporation were driven by synoptic-scale atmospheric processes rather than local katabatic winds. The bulk-aerodynamic method suggests the average daily evaporation to be 2.0 mm day<sup>-1</sup>, which is over 32 % less than the results based on the EC method. This method is much better in producing the day-to-day variations in evaporation compared to the combination equations, which underestimated the evaporation over the lake open water table by over 40–72 %. We also suggested a new combination equation to evaluate the summertime evaporation of Lake Zub/Priyadarshini from meteorological observations from the nearest site. The performance of the new equation is better than the performance of the indirect methods considered. After this equation, the evaporation over the period of the experiment was 124 mm, which is only 9 % larger than the result according to the EC method.

~~**Abstract.** The study provides estimates of summertime evaporation over ice free surface of Lake Zub/Priyadarshini located in the Schirmacher oasis, Dronning Maud Land, East Antarctica. Lake Zub/Priyadarshini is the second largest lake in the oasis, its maximum depth is 6 m. The lake is among the warmest glacial lakes, and it is free of ice during almost two summer months. The summertime evaporation over the open water table of the lake was estimated after the eddy covariance (EC) method, the bulk aerodynamic method and Dalton type empirical equations. We used special meteorological and hydrological measurements collected during the field experiment carried out in 2018 in addition to the standard observations at the nearest meteorological site. The EC method was considered as the most accurate given a reference for other estimates of evaporation over the lake water surface. We estimated the evaporation over the ice free lake surface as 114 mm in the period from 1 January to 7 February 2018 (38 days) after the direct EC method. The average daily evaporation is estimated to be 3.0 mm day<sup>-1</sup> in January 2018. The largest changes in the daily evaporation were driven by the synoptic-scale atmospheric processes rather than local katabatic winds. The bulk aerodynamic method suggests the average daily evaporation to be 2.0 mm day<sup>-1</sup>, and it is over 30 % less than the EC method. This method is much better in producing the~~

day-to-day variations in evaporation compared to the Dalton type semi-empirical equations, which underestimated the evaporation over the lake open water table for over 40–72 %. We also suggested a linear empirical relationship to evaluate the summertime evaporation of Lake Zub/Priyadarshini from the observations at the nearest meteorological site and surface water temperature. After this method, the evaporation over the period of the experiment is 120 mm, and it is only 5 % larger than the result according to the EC method. We also estimated the daily evaporation from the ERA5 reanalysis, which suggested the average daily evaporation during austral summer (December–February) 2017–2018 to be 0.6 mm day<sup>-1</sup>. It is only one fifth of the evaporation estimated with the direct EC method.

## 1 Introduction

Liquid water is increasingly more present over margins of glaciers and ice sheets, and over the surface of the Arctic Sea ice and Antarctic ice shelf due to rise of near-surface air temperatures enhancing snow and ice melt. A large part of the melt water accumulates in a population of glacial lakes and streams, which are typical for the lowermost (melting) zone of glaciers and ice sheets, where the amount of liquid water is sufficient for both the surface and subsurface water runoff (Golubev, 1976). The area of the melting zone is evaluated from in-situ data gathered during glaciological surveys or from remote sensing data. The total area of the melting zone over the Antarctic ice sheet was estimated to be over  $92.5 \pm 13$  thousands square kilometers based on the in-situ data collected during the period of 1969–1978 (Klokov, 1979). Estimations of the area of the melting zone in Antarctica are also available from microwave remote sensors for the summers in the period of 1979/80–2005/06, and already during this period the melting zone has covered over 25 % of the entire continent in at least five summers (Picard et al., 2007).

Recently, remote sensors and geophysical surveys have yielded evidence on a large number of glacial lakes in Greenland and Antarctica (Leeson et al., 2015; Arthur et al., 2020). In 2017, remote sensing data allowed the detection of more than 65000 glacial (supraglacial) lakes located over the East Antarctic coast during the peak melting season (Stokes et al., 2019). The total area of these supraglacial lakes was over 1300 km<sup>2</sup>, and most of them were located at low elevations. Glacial lakes are connected by ephemeral streams into a hydrological network that may develop rapidly in the melting season (Lehnherr et al., 2018; Hodgson, 2012). During 2007 – 2016, the mass loss from the Antarctic ice sheet tripled relative to 1997 – 2006 (Meredith et al., 2019), and it explains the observed changes in physiographic parameters (volume, depth and surface area) of many of the glacial lakes located in the East Antarctic oases (Levy et al., 2018; Boronina et al., 2020). Glacial lakes are a well-known indicator for climate change (Verleyen et al., 2003; Williamson et al., 2009; Verleyen et al., 2012). The possible effects of the glacial lakes on global sea level rise are not clear because the processes and mechanisms driving meltwater production, accumulation and transport in the glacial hydrological network are not fully understood (Bell et al., 2017; Bell et al., 2019).

With rising near-surface air temperatures and enhanced snow and ice melt, liquid water is increasingly more present over the margins of glaciers, ice sheets, and over the surface of the Arctic sea ice and Antarctic ice shelf. A large part of melt water accumulates in a population of glacial lakes (Golubev, 1976; Hodgson, 2012). The glacial lakes are typical for the lowermost (melting) zone of glaciers and ice sheets, where the amount of liquid water is sufficient for surface/subsurface runoff

70 (Golubev, 1976). The area of the melting zone is evaluated from in-situ data gathering during glaciological surveys or from remote sensing data. The total area of the melting zone over the Antarctic ice sheet was estimated over  $92.5 \pm 13.0 \times 10^3$  km<sup>2</sup> based on the in-situ data collected during the period of 1969–1978 (Klokov, 1979). Estimations of the area for the melting zone in Antarctica are also available from the microwave remote sensors for the summers in the period of 1979/80–2005/06, and the melting zone expands over 25 % of the continent at least five times (Picard et al., 2007).

75 Recently, remote sensors and geophysical surveys have yielded evidence on a large number of glacial lakes in Greenland and Antarctica (Leeson et al., 2015; Arthur et al., 2020). Stokes et al. (2019) used remote sensing data to detect water bodies over the East Antarctic coast, and more than 65000 glacial (supraglacial type) lakes were found in the peak melting season 2017. The total area of these supraglacial lakes is over 1300 km<sup>2</sup>, and most of them are located at low elevations. Possible effects of glacial lakes on the global sea level rise are not clear because the processes and mechanisms driving the meltwater  
80 production, accumulation and transport in the glacial hydrological network are not fully understood (Bell et al., 2017; Bell et al., 2019). The glacial lakes are a well known indicator for climate change (Verleyen et al., 2003; Williamson et al., 2009; Verleyen et al., 2012). During 2007–2016 mass loss from the Antarctic ice sheet tripled relative to 1997–2006 (Meredith et al., 2019), which may partly explain the observed changes in lakes physiographic parameters (volume, depth and surface area) for many glacial lakes located in the East Antarctic oases (Levy et al., 2018; Boronina et al., 2020). The glacial lakes  
85 are connected by ephemeral streams into a hydrological network that may rapidly develop in the melting season (Lehnherr et al., 2018). After retreating of the glaciers the glacial lakes have become landlocked. The lakes of the landlocked type occupy local relief depressions over deglaciaded areas also named as oases in Antarctica (Simonov, 1971; Hodgson, 2012).

Among others, a modelling approach can help has been applied to understand how climate warming changes the amount of liquid water seasonally formed in the glacial hydrological network including the lakes and streams. The mass (or water) in particular, the water balance equation of a lake is among the models applied to evaluate the allows estimating the volume of  
90 at the lake from known inflow and outflow terms (precipitation, evaporation, surface/subsurface inflow/outflow runoff, water withdrawal) measured or modelled (Chebotarev, 1975; Mustonen, 1986). In Antarctica, various different processes drive the water exchange in the local lakes, and their mass (water) budget is closely linked to the heat budget (glacial and landlocked lakes (Simonov, 1971; Krass, 1986; Shevnina and Kourzeneva, 2017), and different numbers), and it need to different  
95 number of the terms which are important while estimating their volume depending on whether the lake is of the glacial type or the land-locked type. However, for all local lakes, the estimates of the water budget in the water budget of various lakes (Shevnina and Kourzeneva, 2017). The estimates of the water transport scale for the lakes (and their water budget) are sensitive to uncertainties inherent into the methods applied to evaluate evaporation for both glacial and landlocked lakes (Shevnina et al., 2021). This study suggested the estimations for the uncertainties inherent in the indirect methods applied to  
100 simulate the evaporation over the ice free water table of the glacial lake located in Antarctica.

Performing direct measurements of evaporation is difficult in practice, and therefore various indirect methods are used to evaluate the evaporation over the lakes. Finch and Calver (2008) categorize such methods into seven major models (approaches) needing various meteorological and hydrological measurements, and each approach has inherent strengths and

weaknesses. The pan evaporation approach has good accuracy, however the maintenance of instruments is difficult to perform in remote locations, such as Antarctica. The mass (water) balance approach needs observations on the terms of the lake water budget (precipitation, surface/subsurface inflow/outflow runoff, water extraction, etc.) and knowledge of the lake's physiography (volume and surface area) to estimate the evaporation together with the discrepancy term. The discrepancy term depends on the uncertainties inherent in the hydrological and meteorological measurements and in the methods applied to estimate the terms of the lake's water budget (Finch and Calver, 2008). The application of the mass balance method for lakes located in Antarctica is not possible due to the lack of the hydrological observations. In the energy budget approach, evaporation from a lake is estimated as the term required to close the energy budget when all other terms of the budget are known (similarly to the mass balance approach). It needs a large number of observations with a high frequency of the measurements for temperature, wind speed, humidity and radiation fluxes (Finch and Calver, 2008). In the bulk aerodynamic approach, the evaporation is calculated on the basis of data from the Earth surface properties (surface temperature and specific humidity as well as roughness lengths for momentum and moisture/heat) and atmospheric variables (wind speed, specific humidity and air temperature) in the lowermost part of the atmospheric boundary layer. In addition to observational studies on evaporation and associated latent heat flux, the bulk method is the cornerstone for parameterization of the turbulent fluxes of momentum and sensible heat in numerical weather prediction and climate models (Brunke et al., 2003). For Antarctic applications of the bulk method for evaporation and latent heat flux on the basis of in-situ and remote-sensing observations, see Broun et al. (2001), Vihma et al. (2002), Favier et al. (2011), and Boisvert et al. (2020). The combination equation approach includes the elements of both energy balance and mass-transfer approaches in the estimation of evaporation. The Penman equation (Penman, 1948) is among the most famous presenting this approach, where evaporation is calculated from the simultaneous solution of diffusion equations for heat and water vapor, and the energy balance equation (Finch and Calver, 2008). A more general form of the combination equation is given by the Penman-Monteith equation (Monteith, 1965), which was developed to describe evaporation from plants (evapotranspiration). There are also a number of empirical formulas that need additional information on lake surface area, radiation, daily minimum and maximum air temperatures, etc. (Hojjati et al., 2020; Zhao et al., 2013) or require only the air temperature and relative humidity to be known (Konstantinov, 1968). The disadvantage of the empirical and combination equation approaches is that their application is limited by the features of the location where the empirical coefficients were estimated, and there are no regional values suggested for the Antarctic continent (Finch and Hall, 2001). The combination equations are also named as the Dalton-type equations in Odrova (1979). In this study, we estimated the uncertainties inherent in four equations while estimating the summertime evaporation over lakes located in Antarctica. The equations that were used were by Borghini et al. (2013) and Shevnina and Kourzeneva (2017), however, the uncertainties inherent in the estimations are not yet known. Edinger et al. (1968) formulated a basis for the equilibrium temperature approach, where fluctuations in water temperature are driven by heat exchange. The empirical factors approach allows converting the estimations of evaporation from one type of the land surface to evaporation from another one with empirical coefficients (or factors), which are regionally specific

(Finch and Calver, 2008). The estimates of the evaporation are also available from atmospheric reanalyses which share results of simulations carried out applying numerical weather prediction models. Also in the most recent global atmospheric reanalysis, the ERA5 of the European Centre for Medium-Range Weather Forecasts (Hersbach et al., 2020), the evaporation is estimated based on short-term weather forecasts applying the bulk aerodynamic method.

The eddy covariance (EC) method is recognized as the most accurate method in estimation of evaporation. This method has been introduced more than 30 years ago (Stannard and Rosenberry, 1991; Blanken et al., 2000; Aubinet et al., 2012), but it is rarely used in remote regions. The turbulence measurements require special instruments and sensors which are difficult to maintain and operate in places such as Antarctica.

Performing the direct measurements of evaporation is practically difficult, and various methods are applied to estimate over various types of the land surface including the lakes. The methods are generally indirect because they are like narrow or “pointed” measurements made by an instrument, and/or the evaporation is calculated from measured meteorological variables (Guidelines, 2008). Many of these methods require special instruments and sensors for humidity, wind speed and temperature (Brutsaert, 1982; Finch and Hall, 2001); the turbulence measurements (ie. the EC method); profile measurements (ie. the aerodynamic methods) and the measurements at various heights (ie. Bowen-ratio based energy-balance methods). Among others, the eddy covariance (EC) method is recognized as most accurate in estimations of the evaporation. This method has been introduced for more than 30 years (Stannard and Rosenberry, 1991; Blanken et al., 2000; Aubinet et al., 2012). The turbulence (EC) measurements are direct measurements of the vertical flux of water vapour occurring over the lake surface. Assuming that the flux at the measurement height is the same as at the surface (or low as in our field experiment), the EC measurements are direct measurements of local evaporation over the lake. In this study, we assumed that the point measurement of the EC measuring system was a direct measurement of the lake evaporation.


The Dalton type semi-empirical equations allow us to calculate the evaporation from the meteorological observations collected at monitoring sites (Braslavskiy, 1966; Keijman, 1974; Sene et al., 1991; Shuttleworth, 1993; Majidi et al., 2015). The empirical coefficients in these equations should only be used upon conditions that they were determined (Finch and Hall, 2001). Estimates of the evaporation (or sublimation) over the Antarctic areas demonstrate a huge variation range (Thiery et al., 2012). The evaporation over the lakes located in Antarctica are evaluated with the semi-empirical equation with the coefficients estimated for different climate zones (Borghini et al., 2013; Shevnina and Kourzeneva, 2017). In this study, we estimated the uncertainties in the Dalton-type equations applied to calculate evaporation over the lakes located in the Antarctica.

This study addresses summertime evaporation over the ice-free water surface of a glacial lake evaluated by applying various methods, namely, namely the eddy covariance, the bulk-aerodynamic and combination and Dalton type semi-empirical equations. The EC measurements are used as a reference to evaluate the uncertainties in the estimates based on with the bulk aerodynamic method and the combination semi-empirical equations. This information is beneficial, as EC measurements over glacial lakes are rarely available, and other estimates have to be used. The field experiment was carried out on the shore of the large Lake EC measurements of evaporation were collected during a field experiment on the glacial Lake

Zub/Priyadarshini located in the Schirmacher oasis, East Antarctica, ~~from 1 of January to 8 of February 2018. We also suggested the empirical relationship from the daily series of evaporation estimated after the direct EC method and measurements at the nearest meteorological site.~~

## 175 2 The study area, weather and lakes

The Schirmacher oasis ( $70^{\circ} 45' 30''$  S,  $11^{\circ} 38' 40''$  E) is located approximately 80 km from the coast of the Lazarev Sea, Queen Maud Land, East Antarctica (Fig. 1a). The oasis is the ice-free area elongated in a narrow strip around 17 km long and 3 km wide ~~from west-north-west to east-north-east~~ West-Northwest to East-North-East, and its total area is  $21 \text{ km}^2$  (Konovalov, 1962). The relief is hillocks with absolute heights up to 228 m above sea level. The oasis separates the continental ice sheet from the ice shelf, and the region allows studies on deglaciation processes and ~~the~~ continental ice sheet mass balance components including melting and liquid water runoff (Klokov, 1979; Srivastava et al., 2012).-





185

**Figure 1. The lakes in the study region: (a) Location of the Schirmacher oasis (SA) in Antarctica; (b) the lakes in SA (SCAR ADD) with Landsat Image Mosaic of Antarctica, LIMA (<https://lima.usgs.gov/>) given as the background; (c) the observational network in the catchment of Lake Zub/Priyadarshini with a Google Earth image given as the background.**

190

The location and boundary ~~climate~~ of the lakes are given in the Scientific Committee on Antarctic Research (SCAR) Antarctic Digital Database (ADD): <https://www.add.scar.org/>, last access February 8, 2022. In this dataset the location of the lakes in the Schirmacher oasis were systematically shifted to the LIMA composite (Fig. 1 b and the red lines in Fig. 1c). The climate of the oasis is characterized by low air humidity and temperature, and persistent (katabatic) wind blowing most of the year. This easterly-~~south-easterly~~~~southeasterly~~ wind blows from the continental ice sheet, and advects cold continental air masses to the oasis (Bormann and Fritzsche, 1995). There are two meteorological sites operating in the Schirmacher oasis: ~~the~~. The observations were started in 1961 at ~~the~~ Novolazarevskaya (Novo) meteorological site (70°46'36"S, 11°49'21" E, 119 m asl, ~~WorldWord~~ Meteorological Organization (WMO) number 89512). ~~The~~ Maitri meteorological site (70°46'00"S, 11°43'53" E, 137 m asl, WMO number 89514) opened in 1989, and is located ~~over~~ 5.5 km from ~~the~~ Novo site.

195 Both meteorological sites are included in a long-term monitoring network, and their measurements are done according to ~~the~~ WMO's standards ~~standards by WMO~~ (Turner and Pendlebury 2004). The meteorological data gathered at these two stations are available from the British Antarctic Survey Dataset (<https://www.bas.ac.uk>, last accessed 14.12.2018). Table 1 shows weather conditions during the austral summer 2017–2018 and averaged over the period of 1961–2010 according to the observations at ~~the~~ Novo site (the data given by the Arctic and Antarctic Research Institute at

200 [http://www.aari.aq/default\\_ru.html](http://www.aari.aq/default_ru.html), last access ~~December 7, 07.12.2021~~).

**Table 1. The monthly minimum, mean and maximum values for the meteorological parameters calculated for the period of 1961 – 2010 (the values are separated by a slash), and their monthly average calculated for the austral summer 2017 – 2018. The values are evaluated from the observations at the Novo site.**

**Table 1. Basic statistical characteristics for the meteorological parameters observed in the summer months for the period of 1961 – 2010 and 2017 – 2018 (based on the observations at Novo site):**

Parameter	Period	December	January	February
Air temperature, °C	1961–2010	-3.9 / -1.0 / 1.5*	-2.5 / -0.4 / 1.4	-4.7 / -3.3 / -1.0
	2017	-0.1	-	-

	2018	—	-1.3	-3.0
Relative Humidity,%	1961—2010	47/56/69	49/56/66	41/49/59
	2017	50	—	—
	2018	—	57	49
Atmospheric pressure, Pa	1961—2010	965.1/974.7/990.6	963.8/975.6/985.8	964.1/973.3/987.4
	2017	970.2	—	—
	2018	—	969.9	966.6
Wind speed, ms <sup>-1</sup>	1961—2010	4.3/7.4/10.3	3.1/7.0/10.4	5.8/9.4/13.1
	2017	7.0	—	—
	2018	—	6.2	9.4
Soil surface temperature, °C	1961—2010	3.0/6.7/10.0	3.0/6.7/11.0	-2.0/0.2/4.0
	2017	5.0	—	—
	2018	—	3.0	0.0
Precipitation, mm	1961—2010	0.0/5.3/54.8	0.0/2.6/38.0	0.0/2.9/25.9
	2017	1.9	—	—
	2018	—	10.9	4.6

\*Min / Mean / Max

The field experiment ~~lasted 38 days~~ lasts 38 day in January–February, 2018. ~~Generally, the~~ The weather during the experiment was colder and less windy, while the ~~relative humidity and amount of the precipitation~~ precipitation and relative humidity were close to the ~~monthly means estimated for the period 1961–2010~~ long-term mean values (Table 1). According to ~~data from the~~ Novo meteorological site, ~~during the period of the campaign the~~ the daily air temperatures ranged from -8.3 to 2.8 °C, ~~and~~ the wind speed from 1.5 to 14.3 ms<sup>-1</sup>, with an average of 6.2 ms<sup>-1</sup>. The observations at ~~the~~ Maitri site were very similar to those at ~~the~~ Novo site, with the correlation coefficient between the daily series of ~~the~~ air temperature, relative humidity and wind speed ~~varying~~ vary from 0.95 to 0.98. According to the Maitri meteo site, the wind speed varied from 1.6 to 14.4 ms<sup>-1</sup>, with an average of 6.7 ms<sup>-1</sup>. The air temperature ranged from -8.3 to 2.1 °C, with an average of 1.5 °C. The average relative humidity during the summer was 54 %.

~~To plan the field experiment we used 6-hour synoptic observations at the Novo site available from the British Antarctic Survey Dataset (<https://www.bas.ac.uk>, last access 14.12.2018) covering the period 1998–2016 to calculate the wind direction and frequency of wind speed anomalies over the multi-year means for eight ranges (Fig. 1). The positive anomalies in the wind speed suggest that the observed wind speed is higher than the mean value. Generally, the prevailing wind direction is ranged from 120 to 140° (Fig. 1), and the positive wind speed anomalies are typical for this range, ie. one can expect the majority of strong winds from these directions. We accounted for these circumstances in choosing the location to deploy the EC-measuring systems, to aim the water vapor sensor, and to design its maintenance system to sustain the local~~



winds during the field experiment.




Figure 1: Wind direction (x-axis) and frequency of wind speed anomalies (y-axis) according to the observations at Novo meteorological site: (a) December; (b) January.


More than 300 lakes are mapped in the Schirmacher oasis (Fig. 1 b), and many of the lakes stay free of ice in the summertime for almost two months (Simonov, 1971; Richter and Borman, 1995; Kaup and Haendel, 1995; Kaup, 2005; Phartiyal et al., 2011). The hydrological cycle and changes of the lakes' volume are modulated by the seasonal weather cycle (Sokratova, 2011; Asthana et al., 2019). The physiography of the lakes is available from bathymetric surveys for only the largest lakes (Simonov and Fedotov, 1964; Loopman et al., 1988; Khare et al., 2008; Dhote et al., 2021). This study focuses on Lake Zub (also known as Lake Priyadarshini), and hereafter we will use both names of the lake, which is among the largest and warmest water bodies of the Schirmacher oasis. The lake's surface area is  $35 \times 10^3 \text{ m}^2$ , its volume is over  $10 \times 10^3 \text{ m}^3$ , and the maximum depth is 6 m (Khare et al., 2008; Dhote et al., 2021). Lake Zub/Priyadarshini occupies a local depression and is fed by two inflow streams present in warm seasons. The outflow from the lake occurs via a single stream. The lake stays free of ice for almost two summer months from mid-December to mid-February (Sinha and Chatterjee, 2000). The water level (and volume) of Lake Zub/Priyadarshini has been reducing continuously, and in 2018 the lake water level lowered by approximately 0.4 m (Dhote et al., 2021). The lake is used as the water supply for the year-round Indian scientific base Maitri.

More than 300 lakes are mapped in the oasis (Fig. 2 b), and many of the lakes are ice free in summertime for almost two months (Map of SA, 1972; Kaup et al., 1988; Richter and Borman, 1995; Kaup et al., 1995; Kaup, 2005; Khare et al., 2008;

Phartiyal et al., 2011). The physiography of the lakes is available from bathymetric surveys for only the largest lakes (Simonov and Fedotov, 1964; Loopman et al., 1988; Sokratova, 2011). The hydrological cycle and changes of the volume in these lakes are modulated by the seasonal weather cycle (Asthana et al., 2019).

240 Gopinath et al. (2020) used water samples collected from 12 lakes (including Lake Zub/Priyadarshini) located in the Schirmacher Oasis to recognize major sources of water in the lakes. The samples were analysed with the isotope method (Ellehoj et al., 2013), and the isotopic concentrations show that Lake Zub/Priyadarshini is mostly sourced by the melting of the adjacent glaciers. For landlocked lakes, the major source of water is melting seasonal snow cover (Shevnina and Kourzeneva, 2017; Hodgson, 2012). It allows us to suppose that Lake Zub/Priyadarshini is the glacial type, as it is not the landlocked type as given in Phartiyal et al. (2011). Lake Zub/Priyadarshini is the lowest in the chain of the glacial lakes  
245 sourced by the ice/snow melting in the lowermost zone of the glaciers, and we estimated that more than 60 % of its catchment area is covered by rocks. This allows for the specific thermal regime and water balance of this glacial lake, which is among the warmest in the oasis: its water temperature rises up to 8 – 10 °C in January (Ingole and Parulekar, 1990). Such water temperatures are typical for the landlocked lakes (Simonov, 1971).

Lake Zub/Priyadarshini presents ideal conditions to study evaporation over a glacial lake, and to plan the field experiment,  
250 we accounted for the location to set up the EC measuring systems. Selection of the exact site for EC measurements requires, among others, data on the prevailing winds and their fetch over the lake, and naturally also the accessibility for regular maintenance. To evaluate the prevailing wind direction, we used 6-hourly synoptic observations at the Novo site available from the British Antarctic Survey Dataset (<https://www.bas.ac.uk>, last accessed December 14, 2018) covering the period 1998–2016. We calculated the number of cases when wind was blowing from 36 sectors each 10 degrees wide, and then  
255 defined the prevailing wind directions (marked with the black arrows in Fig. 2). The prevailing wind directions range from 110 to 140°.



**Figure 2: Wind direction and wind speed anomalies for two austral summer months (December and January).**

We also calculated the wind speed anomalies of each 10-degree sectors given in colour codes in Fig. 2. The positive wind speed anomalies are often observed within the range of the prevailing wind directions (marked with orange, yellow, red, brown and black in the legend of Fig. 2). Therefore, one can expect the majority of strong winds from these directions. The region of the study is featured by persistent katabatic winds blowing from the continental interior. Fig. 2 shows that almost all winds come from a direction that would be the direction of katabatic winds. However, it is not guaranteed that all these winds are entirely of katabatic origin, and some winds may be driven by a combined effect of katabatic and synoptic forcing.

**Figure 2. The lakes in the study region: (a) Location of the Schirmacher oasis (SA) in Antarctica; (b) the lakes in SA (after Map of SA, 1972) with Landsat Image Mosaic of Antarctica, LIMA given as the background; (c) the temporal observational network in the catchment of Lake Zub/Priyadarshini.**

This study focuses on Lake Zub/Priyadarshini, which is among the largest and warmest water bodies of the Schirmacher oasis. Water in the lake is mainly sourced by the continuous the glacial melting water (Gopinath et al., 2020). Lake is not fully lost the connection to the glacier, and its melting is still a major inflow term of the water budget of the lake. It allows us to suppose that Zub/Priyadarshini is the glacial type (not the landlocked type as given in Phartiyal et al., 2011). This lake is the lowest in the chain of the glacial lakes over the continental ice-sheet. The lake catchment includes a low portion of glaciated area, and results in a specific thermal regime and water budget of the lake. The water temperature in the lake rises up to 8–10 °C in January (Ingle and Parulekar, 1990), and it's typical for the landlocked lakes (Simonov, 1971).

275 The lake's surface area is  $33.9 \times 10^3 \text{ m}^2$ , its volume is over  $10.0 \times 10^3 \text{ m}^3$ , and maximal depth of the lake is 6 meter (Khare et al., 2008). Lake Zub/Priyadarshini occupies a local depression, and is fed by two inflow streams present in warm seasons. The outflow from the lake occurs via a single stream. The lake stays free of ice for almost two summer months from mid-December to mid-February, and it has no significant thermal stratification withing this period (Sinha and Chatterjee, 2000). The lake is used as the water supply of the year-round scientific base Maitri (India).

## 280 | 3: Data and Methods

### 3.1 Data

In the field experiment in 2017–2018, we collected the hydrological and meteorological observations needed to evaluate the water balance terms of Lake Zub/Priyadarshini. The hydrological network included water level/temperature gauges, water discharge/level gauges and an evaporation gauge (Fig. 2 c). In this study we used only those data required in the evaluation  
285 of only one term of the water budget of a lake, namely evaporation. The evaporation gauge was a flux tower equipped with an Irgason device by Campbell Scientific. The Irgason consists of a 3D sonic anemometer and two gas analysers measuring  $\text{CO}_2/\text{H}_2\text{O}$  concentrations, and it is one of the EC devices taking this kind of measurements (in this study, we named our EC station as Irgason). The Irgason was deployed on the shore of the lake to collect high-frequency data on wind speed/direction and water vapor concentration needed to evaluate evaporation with the EC method (Fig. 3 a). The flux tower was placed 5–6  
290 m inland of the shoreline of Lake Zub/Priyadarshini for the period of 1 January to 7 February 2018 (Shevnina, 2019). The meteorological parameters (air temperature, wind speed and relative humidity) were measured simultaneously at the Maitri meteorological site and at the evaporation gauge located on the lake shore (Irgason in Fig. 1 c). The data gathered by the sensors cover various observational periods (Table 2). The shortest 14-day period with the measurements is available for the iButton sensor, and this period lasted from January 27 to February 9, 2008.

295 In the field experiment 2017 – 2018, we collected the hydrological and special meteorological observations needed to evaluate the water balance terms of Lake Zub/Priyadarshini. The temporal hydrological network included water level/temperature gauges, water discharge/level gauges and an evaporation gauge (Fig. 2 c). In this study we used only those data required in evaluation of only one term of the water budget of a lake namely the evaporation. The evaporation gauge was a flux tower equipped with the Irgason EC measurement device by Campbell Scientific (user manual available at  
300 [https://s.campbellsci.com/documents/ca/manuals/irgason\\_man.pdf](https://s.campbellsci.com/documents/ca/manuals/irgason_man.pdf), last access 09.07.2021). The Irgason consists of a 3D sonic anemometer and two gas analysers measuring  $\text{CO}_2/\text{H}_2\text{O}$  concentrations, It was deployed on the shore of the lake to collect high-frequency data on the wind speed/direction and water vapor concentration needed to evaluate the evaporation with the EC method. The flux tower was placed 5–6 m to the shoreline of Lake Zub/Priyadarshini for the period of 1 January to 7 February, 2018 (Shevnina, 2019). Irgason was deployed on the boom at the height of 2 meter, and it was fixed with 6  
305 metal guidelines angled  $120^\circ$  to each other (Fig. 3 a) The field experiment lasted for 38 days, and the meteorological parameters (air temperature, wind speed and relative humidity) were measured simultaneously at the Maitri meteo site and at

the evaporation gauge located on the lake shore (Irgason in Fig. 2 c). The data gathered by the various sensors cover observational periods lasting from 14 to 45 days (Table 2).

**Table 2. The hydrological and meteorological data collected during the field experiment in the summer 2017–2018: “–” no information available.—2018–**


<u>Site / Sensor</u> (Fig. 1 c)	<u>Elevation,</u> <u>m</u>	<u>Measured variables</u>	<u>Accuracy /</u> <u>(Resolution)</u>	<u>Time series</u> <u>used in the</u> <u>analysis</u>	<u>Period</u>
<u>Irgason site</u>	<u>124</u>	<u>air temperature, °C;</u> <u>H<sub>2</sub>O concentration, g/m<sup>3</sup>;</u> <u>wind speed, ms<sup>-1</sup></u>	<u>±0.15 / (0.025)</u> <u>±0.037 / (0.00350)</u> <u>=</u>	<u>30 minute</u>	<u>01.01.2018 –</u> <u>07.02.2018</u>
<u>Hobo</u>	<u>122</u>	<u>water temperature, °C</u> <u>barometric pressure, Pa</u>	<u>±0.44 / (0.10)</u> <u>=</u>	<u>daily average</u> <u>(not used in this</u> <u>study)</u>	<u>30.12.2017 –</u> <u>09.02.2018</u>
<u>iButton</u>	<u>122</u>	<u>water temperature, °C</u>	<u>±0.5 / (0.5)</u>	<u>daily average</u>	<u>27.01.2018 –</u> <u>09.02.2018</u>
<u>Maitri site</u>	<u>137.5</u>	<u>air temperature, °C;</u> <u>relative humidity, %;</u> <u>wind speed, ms<sup>-1</sup></u>	<u>±0.2 / (-)</u> <u>±1 / (-)</u> <u>±0.5 / (-)</u>	<u>daily average</u>	<u>01.12.2017 –</u> <u>28.02.2018</u>

Table 2 shows the information on the accuracy and resolution of the sensors according to the technical specifications given by the manufacturers. Ramesh and Soni (2018) give the information for the sensors installed at the Maitri site. The elevation of the lake water level was measured by the geodetic instrument Leica CS10 during the installation of the Solinst water level logger on 30 December 2017 (Fig. 1 c). The same instrument was used to measure the elevation of the Maitri site in January 2018 (Dhote et al., 2021). The elevation of the lake water level was 122.3 m (WGS84 ellipsoid vertical datum), we further used this elevation while calculating the elevation for the Hobo, iButton and Irgason temperature sensors.

310



a)



315


Site / Sensor (Fig. 2-c)	Elevation, m	Measured variables	Period	Time series used in the analysis
Irgason-site	125.5	Air temperature, °C; H <sub>2</sub> O concentration, g/m <sup>3</sup> ; 3D wind speed, ms <sup>-1</sup>	01.01.2018—07.02.2018	30-minute
Hobo	123	Water temperature, °C	30.12.2017—09.02.2018	Daily average
Solinst	119.5	—	01.01.2018—15.12.2018	

iButton	123	—	27.01.2018—09.02.2018	
Maitri site	137.5*	Air temperature, °C; relative humidity, %; wind speed, ms <sup>-1</sup>	01.12.2017—28.02.2018	Daily average

\* measured during the summer season 2017—2018 by the geoesic instrument Leica CS10, the elevation is given in WGS84 vertical datum.





(a)




(b)

Wind Speed [m/s]



(c)

X90 [m]





**Figure 3: -The experiment on the coast of Lake Zub/Priyadarshini: (a) Irgason deployed on the lake shore (06.01.2018); (b) wind speed and direction measured at the Irgason site, dashed line indicates the footprint wind sector; (c) the footprint length estimate (X90).**

The footprint is an important concept for evaluating fluxes correctly with the EC method. The footprint is defined as the area upwind of the EC station where the fluxes observed at the station originate from. Hence, the footprint area depends on the location of the EC station, the height of its sensors, the roughness of the upwind surface, and the stratification of the upwind atmospheric surface layer (Kljun et al., 2004; Burba, 2013). The Irgason was settled at the height of 2 metres above the ground, which yields footprint lengths of less than 200 metres, which in this study was defined as X90 and represented 90 % of the cumulative contribution to the fluxes (Fig. 3 c). This distance is less than twice of that between the Irgason and the shore of Lake Zub/Priyadarshini in an east-southeast direction (Fig. 1 c), and it ensures that the measured data is representative only for the lake and free of contamination from the upwind shore. The tower height of 2 m generates a blind zone near the tower, so that the stones on the downwind shore do not affect the fluxes. The location of the EC tower accounted for the prevailing wind directions (Fig. 2) meaning that the footprint area is mainly represented by the lake surface. We filtered out data outside the footprint (Fig. 3 b). Gaps in the wind direction were replaced with the average values of the neighboring 30-minute blocks. The Irgason's raw data consisted of values measured at a frequency of 10 Hz. We used these raw data to calculate a 30-minute time series of evaporation, turbulent fluxes of momentum, sensible heat and latent heat, as well as air temperature, wind speed, and wind direction. The daily evaporations were calculated as a sum of the 30-minute time series. The low observation height of 2 m guarantees that the vertical divergence of the water vapour flux is negligible, and therefore the water vapour flux observed at the height of 2 m represents the surface evaporation.

To allow the estimation of evaporation by the combination equations, measurements of the water temperature are needed. We measured the water temperature of the lake's surface with two sensors: the iButton temperature sensor was installed in Lake Zub/Priyadarshini in the depth of 0.2 metres and was placed ahead of the EC station (Irgason) toward the prevailing wind directions. The Hobo temperature sensor was deployed in the depth of 0.2 metres in the end of the stream inletting the neighbouring lake (Fig. 1 c). This stream is an outlet of Lake Zub/Priyadarshini, and we assumed that the observations collected by the Hobo were representative for the stream more than for the neighbouring lake itself. The accuracy of both temperature sensors is similar, and the resolution of the Hobo temperature sensor is better than the iButton's resolution (Table 2). The lake water temperature was measured every 10 minutes, and we further calculated the daily average time series of the water temperature in the lake. The Hobo consists of two sensors measuring temperature and barometric pressure allowing us to evaluate the water level/stage, and we used only the temperature measurements in this study. Sinha and Chatterjee (2000) reported that Lake Zub/Priyadarshini was thermally homogeneous down to the bottom almost from mid-January 1996 to mid-February 1997, and we assumed that the lake had no thermal stratification during the whole field experiment in 2018.

For the EC method, the footprint is an important concept to evaluate the fluxes correctly. The footprint is defined by a sector of wind direction covering the source area, and the footprint length depends on the height of the sensors (Kljun et al., 2004;

Burba et al., 2016). The location of the Igrason done accounting to the prevailing wind directions (Fig. 2), so that for most of the time the wind is blowing from the source area covered by the lake surface, and we filtered out data outside the footprint (Fig. 3 b). The gaps in the wind direction were replaced with average values of the neighbour 30-minute blocks. The Irgason was settled at the height of 2 meter above the ground, and it allows for footprint lengths less than 200 meter (Fig. 3 c), with only one exception during the whole experiment. This distance is twice less than those between the Irgason and the shore of Lake Zub/Priyadarshini, in east-southeast direction (Fig. 2 c). This condition ensures that the retained data is representative only from the lake and free of contamination from the shores. The height of the Irgason allows for a blind zone near the tower, therefore the stones on the lake shore do not affect the fluxes. The raw data by Irgason consisted of the values measured at a frequency of 10 Hz. We used these raw data to calculate a 30-minute series of the evaporation, turbulent fluxes of momentum, sensible heat and latent heat, as well as air temperature, wind speed, and wind direction. The daily evaporation were calculated as a sum of the 30-minute series, and the combination of the EC footprint and the small dimension of the lake allow us to consider these measurements as direct measurements of the evaporation over the lake surface.

To allow the estimation of the evaporation after the Dalton type empirical equations, the measurements of the water temperature are needed. Therefore, we measured the water temperature of the lake's surface with three sensors installed in Lake Zub/Priyadarshini (Fig. 2 c) at the depths of 0.2 meter (Hobo, iButton) and 3.9 meter (Solinst). The sensors Solinst, Hobo and iButtons were installed at different depths, and it is the result of the misunderstanding between hydrologists working in the field. We expected that the water temperature in the lake is measured by a chain of sensors instead of the Solinst. The Hobo sensor was deployed on the stream inletting the neighbouring lake, and its measurements were considered to be representative for the water level and water temperature in the stream more than for the neighbouring lake itself. We assumed that Lake Priyadarshini is thermally homogeneous down to the bottom (Sinha and Chatterjee, 2000). In summers, the thermal homogeneity is typical for the lakes of the similar morphology located in the Larsemann Hills oasis, East Antarctica (Shevnina and Kourzeneva, 2017). The lake surface temperature was measured every 10 minute, and we calculated further the daily average series of the water temperature in the lake. Two temperature sensors (Solinst and Hobo) also measured the barometric pressure allowing us to evaluate the water level/stage in Lake Zub/Priyadarshini, however, we did not utilize these data in this study focused on the evaporation only.

Figure 4 a shows the daily time series of the lake water temperature, air temperature and wind speed calculated from the measurements done by the sensors during the period of the experiment. The best agreement was found for the water temperature measured by the sensors Hobo and iButton, the correlation coefficient for these series equals to 0.89. The water temperature measured by Solinst sensor is systematically lower than those measured by Hobo and iButtons (Fig. 4 a). This circumstance is likely connected to the effect of the cold water incoming with the inflow stream, which is incoming close to the deploying place of Solinst temperature sensor. This inflow stream results from the small glacial lake located upstream of Lake Zub/Priyadarshini, and the water in the upstream lake is colder than in Lake Zub/Priyadarshini itself.




Figure 4: Daily time series of (a) the lake surface water temperature measured by the Hobo (blue), by the iButton (red) and Solinst (grey), and the air temperature measured at the Maitri site (black).

(b) the 2-meter wind speed estimated with the logarithmic profile after the measurements at Maitri site. The red dotted lines show the daily minimum and maximum water temperatures measured by iButton temperature sensor.

Figure 4 shows the daily time series of the lake water temperature, air temperature and wind speed calculated from the measurements done by the sensors during the period of the experiment. The correlation coefficient equals to 0.89 for the series of the water temperature measured by two temperature sensors (Hobo and iButton). We further used the measurements collected by the Hobo temperature sensor to estimate the evaporation over Lake Zub/Priyadarshini in January 2018. In our calculations based on the combination equations we applied the data collected by the meteorological sensors installed both at Maitri and Irgason sites, and the meteorological sensors are deployed at the different elevations (Table 2). The elevation of the temperature sensor and gas analyser of the Irgason is lower than the sensors at Maitri site, and therefore we used the logarithmic approximation of the wind profile to correct the wind speed data measured at the Maitri site, for which we estimated a constant aerodynamic roughness length of 0.002 m (Stull, 2017). We did not use any height correction for the

data on the relative humidity and air temperature since their changes with elevation are negligible in our case (Tomasi et al., 2004).

In our calculations after the Dalton-type equations we applied the data collected by the meteorological sensors installed both at Maitri and Irgason sites. The meteorological sensors are installed at different heights: the Irgason's sensors are deployed at the height of 2 meters over the ground (the lake water table), and the sensors at Maitri site are mounted on the mast at the height of 6 meters over the ground. It requires applying the transformation to the wind speed measured at the Maitri site before using these measurements in the Dalton-type equations. Further, we used the logarithmic approximation of the wind profile to correct the wind speed data measured at the Maitri site, where the roughness length constant equaling to 0.0024 meter (as suggested: <https://wind-data.ch/tools/profile.php?lng=en>, last access 15.10.2021). We did not use any transformation for the data on the relative humidity and air temperature since their changes with elevation are negligible in our case (Tomasi et al., 2004).

### 3.2 Methods

To evaluate the evaporation with the direct EC method, we used the data collected by the Irgason installed on the shore of Lake Zub/Priyadarshini. The Irgason raw data were measured with a frequency of 10 Hz, which were further analysed in the following steps. First, the bad data with less than 50 % of total 10 Hz measurements were excluded. Second, we excluded all data automatically flagged for low quality, and the data with a gas signal strength less than 0.7 (or 70 % of the strength of a perfect signal). The gas signal strength is usually lower than 0.7 during rainfalls, which were not observed in January, 2018 in the Schirmacher oasis. Generally, rainfalls are rare along the East Antarctic coast where rainfall occurs 22 days per year at most (Vignon et al., 2021). In the third step, the spikes were removed applying the method by Vickers and Mahrt (1997), fixing the threshold window of 3.5 standard deviation for horizontal wind speed, CO<sub>2</sub> and H<sub>2</sub>O and 5.0 for vertical wind speed. This procedure was repeated up to 20 times or until no more spikes were found. Finally, we obtained, among others, the 30-minute fluxes of momentum, sensible heat and latent heat (evaporation), as well as the water vapor and carbon dioxide concentrations (see the Supplement). The evaporation over the lake was calculated only by those values collected within the footprint of the ice-free surface of the lake. Therefore, we filtered the data outside the footprint which covered the wind directions within the range of 105 – 240° (Fig. 3 b). We excluded 18 % of the total data from further consideration after the three step filtering. To fill these gaps we replaced the excluded values by the mean value, which was estimated from the time series of 30-minute values. We also evaluated the relative humidity from the water vapor concentration as given by Hoeltgebaum et al. (2020).

The daily evaporation over ice free surface of Lake Zub/Priyadarshini was evaluated with the direct and indirect methods. The indirect methods are the bulk aerodynamic method and Dalton-type semi-empirical equations. To evaluate the evaporation with the direct EC method, we used the data collected by the Irgason instrument installed on the shore of Lake Zub/Priyadarshini. The Irgason raw data were measured with a frequency of 10 Hz, and they required post processing. We

435 followed Potes et al. (2017) in the post processing procedure of the raw data, which were further filtered in three steps: in the first step, the bad data with less than 50 % of total 10-Hz measurements were excluded; on the second step, we excluded all data automatically flagged for the low quality, and the data with the gas signal strength less than 0.7 (or 70 % of the strength of a perfect signal). The gas signal strength is usually lower than 0.7 during rain, which is not observed in Antarctica. On the third step, the raw data were processed to remove spikes after the method by Vickers and Mahrt (1997). This procedure was repeated up to 20 times, or until no more spikes were found. Finally, the 30-minute values of the atmospheric fluxes (the momentum flux, the sensible heat flux, the latent heat flux), the water vapor concentration, the specific humidity, various turbulence parameters and evaporation were obtained (see the Supplement).

440 To account for only those values collected within the lake surface area, we also filtered the data outside the footprint, which covered the winds with the direction ranging from 105 to 240° (Fig. 3 b). We excluded 18 % of the total measurements from further consideration after three steps filtering, and these gaps were replaced with the mean and median values. Finally, the daily evaporation over the lake surface was estimated as the sum of 30-minute values in each day of the experimental period. We also evaluated the relative humidity from the water vapor concentration as given by Hoeltgebaum et al. (2020) to compare with the relative humidity measured at the Maitri site.

445 Uncertainties in the estimation of evaporation by after any method include the instrumental errors associated with the specific instrument. Aubinet et al., (2012) suggest three methods allowing the quantification of to quantify the uncertainty of the EC method. In this study, we applied the paired tower method to evaluate the instrumental uncertainties of the EC method taking advantage of an intercomparison campaign in Alqueva reservoir, Portugal, in October 2018. The instrumental error does not depend on the region where the instrument will be used, and therefore the intercomparison may be done elsewhere. The relative instrumental error estimated in this intercomparison campaign was 7 % (see the Annex). The uncertainties of the EC method also include the errors due to the filtering of measurements within within the footprint area. In our study, 18 % of gaps were filtered, and we filled these data with the mean and median values. The large number of filters and corrections that we applied to the EC data allowed us to reduce the errors and uncertainties. Even the EC method itself has some errors and uncertainties but it is the most versatile and accurate method to measure the evaporation.

455 In the bulk aerodynamic approach, evaporation aerodynamic method, the evaporation ( $\text{kg m}^{-2} \text{s}^{-1}$ ) is defined as the vertical surface flux of water vapor due to atmospheric turbulent transport. It is calculated from the difference in specific humidity of between the surface (i.e., ice or water for which the specific humidity equals the saturation specific humidity that depends on the surface temperature), and the air, as well as the factors that affect the intensity of the turbulent mixing exchange: wind speed, surface roughness, and thermal stratification (Boisvert et al., 2020; Brutsaert, 1985). In our study, the evaporation after the bulk-aerodynamic method was calculated as follows :

The evaporation based on the bulk-aerodynamic method is calculated as follows:

$$E = \rho C_{Ez} w_z (q_s - q_{az}) \quad (1)$$

$$E = \rho C_{Ez} (q_s - q_{az}) w_z \quad (1)$$

465 where E is the evaporation (in  $\text{kg m}^{-2} \text{s}^{-1}$ , which we in the following convert to  $\text{mm day}^{-1}$ ,  $\text{kg s}^{-1}$ );  $\rho$  is the air density, ( $\text{kg m}^{-3}$ );  $C_{Ez}$  is the turbulent transfer coefficient for moisture unitless),  $q_s - q_s$  is the saturation specific humidity at water surface of the lake ( $\text{kg/kg}$ ),  $q_s$  corresponding to the lake surface temperature,  $q_{az}$  is the air saturation specific humidity ( $\text{kg/kg}$ ) specific humidity, and  $w_z$  is the wind speed (m). The subscript z refers to the observation height (here 2 m). The

470 meters). For the turbulent transfer coefficient for moisture depends on the atmospheric stratification: for  $C_{Ez}$  under neutral stratification ( $C_{EzN}$ ); we applied the value of 0.00107 based on previous measurements over a boreal lake (Heikinheimo et al., 1999; Venäläinen et al., 1998). It allows us to better take into account the different regime of turbulent mixing over a small lake compared to the sea (Sahlée-Sahlee et al., 2014). As the stratification is not always neutral, we took into account the effects of stratification on the turbulent transfer coefficient  $C_{Ez}$  as follows:

Since the stratification of the atmosphere is not always neutral, we took into account its effects on the turbulent transfer coefficient as follows:

$$475 \quad C_{Ez} = \frac{C_{DzN}^{1/2} C_{EzN}^{1/2}}{\left[ 1 - \left( \frac{C_{DzN}^{1/2}}{k} \right) \psi_m \left( \frac{z}{L} \right) \right] \left[ 1 - \left( \frac{C_{EzN}^{1/2}}{k} \right) \psi_q \left( \frac{z}{L} \right) \right]} \quad (2)$$

where,  $C_{DzN}$  is the neutral drag coefficient for the lake surface,  $k$  is the von Karman constant (0.4),  $\psi_m$  and  $\psi_q$  are empirical stability functions ; and  $L$  is Obukhov length ( in meters):

480 where  $C_{DzN}$  is the neutral drag coefficient for the lake surface (0.00181; Heikinheimo et al. (1999)),  $k$  is the von Karman constant (0.4), and the effects of thermal stratification are presented by the empirical functions ( $\psi_m$  and  $\psi_q$ ) depending on the Obukhov length ( $L$ ). For  $\psi_m$  and  $\psi_q$ , we used the classic form by Businger et al. (1971) for unstable stratification and that of Holtslag and de Bruin (1988) for stable stratification. The values by Heikinheimo et al. (1999) were given for  $z = 3$  meters, and converted to our observation height of 2 meters using Launiainen and Vihma (1990), and the same algorithm was applied to iteratively solve the interdependency of the turbulent fluxes and  $L$ . The latent heat flux is obtained by multiplying the evaporation rate by the latent heat of vaporizations.

485 The Dalton type semi-empirical equations allow calculation of the evaporation from a wind function and a gradient of the temperature of water surface and ambient air measured at the height 2 meters:

$$L = - \frac{\rho c_p u_*^3 \theta_z}{k g H} \quad (3)$$

490 where  $\rho$  is the air density,  $c_p$  is the specific heat,  $u_*$  is the friction velocity,  $\theta_z$  is the air potential temperature,  $g$  is the acceleration due to gravity, and  $H$  is the surface sensible heat flux. The Obukhov length (Obukhov, 1946) is the key element of the Monin-Obukhov similarity theory (Monin and Obukhov, 1954; Foken, 2006), and needed to adjust the bulk transfer coefficients to the actual stratification in the atmospheric surface layer. In our calculations, the neutral drag coefficient equals to 0.00181 as suggested by Heikinheimo et al. (1999). For  $\psi_m$  and  $\psi_q$ , we used the classic form by Businger et al. (1971) for unstable stratification and that of Holtslag and de Bruin (1988) for stable stratification. The values by Heikinheimo et al. (1999) were given for  $z = 3$  meters, and converted to our observation height of 2 meters using Launiainen and Vihma (1990), and the same algorithm was applied to iteratively solve the interdependency of the turbulent fluxes and  $L$ .  
495 The latent heat flux is obtained by multiplying the evaporation rate by the latent heat of vaporization.

Most of the empirical equations are based on a simple mass transfer relation between the evaporation rate and the water deficit and wind conditions. The general form of the relation reads as  $E = K w_2 (e_s - e_2)$ , where  $K$  is empirical function approximated with a small number of coefficients. Among others, Shuttleworth (1993) suggests two mass transfer equations for the estimation of evaporation from the surface of lakes and ponds depending on their surface area. In this study, we used his formula for water bodies in the range of  $50 \text{ m} < A^{0.5} < 100 \text{ km}$  located in regions with a relatively arid climate. The equation reads as  $E = 2.909 A^{0.05} w_2 (e_s - e_2)$ , where  $E$  is the evaporation in  $\text{mm day}^{-1}$ ;  $A$  is the surface area in  $\text{m}^2$ ;  $w_2$  is 2-metre wind speed in  $\text{ms}^{-1}$ ; and  $e_s$  and  $e_2$  are the surface water and air vapor saturation pressure in kPa. In this study, we used this formula to estimate the daily evaporation from Lake Zub/Priyadarshini, whose surface area is estimated as  $350\,000 \text{ m}^2$  in 2016 (Dhote et al., 2021). The method by Shuttleworth (1993) has been used to evaluate evaporation over small lakes located in Antarctica (Boghini et al., 2013), however the scope of uncertainties inherent in the method is not known.

Penman (1948) first suggested taking the elements of the mass transfer and energy budget approaches into the estimation of evaporation from open water, and his formula is one of the combination equations (Shuttleworth, 1993; Finch and Calver, 2008). In this study, we applied three combination equations to calculate daily evaporation:  $E = 0.26 (1 + 0.54 w_2)(e_s - e_2)$  and  $E = 0.26 (1 + 0.86 w_2)(e_s - e_2)$  adopted from Tanny et al. (2008), where these formulas are referred to Penman (1948) and Doorenbos and Pruitt (1975) respectively. These equations are among those most often used in hydrological practice (Finch and Calver, 2008), and therefore we have chosen them in this study. We also used the formula  $E = 0.14 (1 + 0.72 w_2)(e_s - e_2)$ , which has been applied to evaluate evaporation from lakes located in northern Russia (Odrova 1979). In these equations,  $e_s$  and  $e_2$  are the surface water and air vapor saturation pressure (millibars), and we calculated them according to Tetens's formula given in Stull (2017). The method by Odrova (1979) has been used in estimations of evaporation over glacial lakes located in Antarctica (Shevnina and Kourzeneva, 2017), but the method's uncertainties have not been estimated. We calculated daily evaporation separately using the meteorological observations collected at the Maitri site and at the lake shore (Irgason site).

where,  $E$  is daily evaporation,  $\text{mm day}^{-1}$ ;  $e_s$  is the water vapor saturation pressure;  $e_2$  is the water vapor pressure at 2 meters height;  $C$  is a coefficient (or a function) depend on meteorological conditions (or a linear wind function with two parameters computes boundary layer transfer coefficients (Tanny et al., 2008)). The  $C$  is evaluated from observations with empirical approximations (Finch and Hall, 2001).

In this study, we applied three semi-empirical equations to calculate the daily evaporation rate suggested by Penman (1948), by Doorenbos and Pruitt (1975) and by Odrova (1979), the Eqs. 4–6 respectively:

$$E = 0.26(1+0.54 w_2)(e_s - e_2) \quad (4)$$

$$E = 0.26(1+0.86 w_2)(e_s - e_2) \quad (5)$$

$$E = 0.14(1+0.72 w_2)(e_s - e_2) \quad (6)$$

where  $E$  is the evaporation expressed in  $\text{mm day}^{-1}$ ;  $w$  is the wind speed measured at the 2 meters height;  $e_s$  and  $e_2$  are water and air vapor saturation pressure, given in millibars (calculated according to the Tetens's formula in Stull, 2017). The approximations by Penman (1948) and Doorenbos and Pruitt (1975) are among the methods which are most often methods used in hydrological practice (Finch and Hall, 2001), therefore we have chosen them in this study. The method by Odrova (1979) is used to evaluate the daily evaporation over the lakes in Antarctica, however the scope of uncertainties of this method is not estimated previously (Shevnina and Kourzeneva, 2017). After the semi-empirical equations, we calculated



the daily evaporation separately using the meteorological observations collected at the Maitri site and at the lake shore (Igrason site). After the bulk aerodynamic method, the daily series of the evaporation were evaluated from the 30-minute series of the meteorological data collected at both sites:

The empirical coefficients in the combination equations usually limit their applicability to the region where such coefficients are obtained (Finch and Hall, 2005). The empirical coefficients in four selected equations are evaluated from data gathered in regions with different climates, and therefore they probably will not be applicable for lakes located in Antarctica. In this study, we suggested two regional empirical relationships based on the daily series of evaporation estimated by the direct EC method and the meteorological observations at the Maitri site, which is the nearest meteorological site to the lake. In the first relationship, evaporation ( $E$ ,  $\text{mm day}^{-1}$ ) was evaluated with the linear model  $(a + b w_2)(e_s - e_2)$ , where  $a$  and  $b$  are fitted with empirical coefficients, and  $(e_s - e_2)$  is expressed in mbar. The second relationship reads as  $E = a w_2^b (e_s - e_2)$ . The efficiency of fitting the coefficients were performed on the same data for the experiment (lasting 38 days).

The empirical coefficients in the Dalton type empirical equations usually limit their applicability to the region where such coefficients are obtained (Finch and Hall, 2005). The empirical coefficients in the equations (4-6) are evaluated from the data gathered in regions with different climates, and therefore they probably will not be applicable for lakes located in Antarctica. In this study, we suggested the regional empirical relationship using the daily series of the evaporation estimated after the direct EC method and observations at the meteorological site nearest to the lake. The evaporation ( $\text{mm day}^{-1}$ ) was evaluated with the linear regression model  $a + b_1 w_2 + b_2 (e_s - e_2)$ , where  $(e_s - e_2)$  is expressed in mbar. The efficiency of the relationship was estimated with the cross validation procedure, where the whole period with observations (38 day) was divided into two sub periods 19 day each. The daily evaporations within the first period were used to estimate the empirical coefficients, and then the daily evaporation in the second period were used as the independent data while estimating the efficiency of the empirical relationship. Then, the procedure was applied *vice-versa*: the values for the empirical coefficients were evaluated from the evaporation over the second sub period, and the efficiency of the relationship was estimated using the evaporation in the first sub period.

Evaporation by The evaporation after the indirect methods were compared to the direct (EC) method in order to find the method with the lowest scope of range in the uncertainties, and, therefore, the method of the highest efficiency. We applied

the Pearson correlation coefficient ( $PR$ ), the root square standard error ( $RMSE = \sqrt{\sum_1^n (E_{EC} - E_{mod})^2}$ ) and the  $s/\sigma$

criteria ( $SSC$ ) to evaluate the scope. To estimate the efficiency of the uncertainties inherent in the indirect methods. The  $SSC$

reads as follows (Popov, 1979):  $s = \sqrt{\sum_{i=1}^n (E_{EC}^i - E_m^i)^2 / (n-m)}$ , and  $\sigma = \sqrt{\sum_{i=1}^n (E_{EC}^i - \bar{E}_{EC})^2 / n}$ . In these formulas,  $\bar{E}$

is the mean evaporation, (mm);  $n$  is the length of the series (38), and  $m$  is the number of empirical coefficients in the relationships (equal to 2). Overall, a new method is acceptable for further use in hydrological practice if the  $SSC$  value is less than 0.8 (Popov, 1979). bulk aerodynamic method and semi-empirical equations, we used the Pearson correlation coefficient

and the Nash-Sutcliffe efficiency index (Nash and Sutcliffe, 1970) as given by Tanny et al. (2008):-

$$NSS = 1 - \frac{\sum_{i=1}^n (E_{EC}^i - E_m^i)^2}{\sum_{i=1}^n (E_{EC}^i - E_{EC})^2}, \quad (7)$$

where  $E_{EC}$  and  $E_m$  are the evaporation estimates after the direct method and after the indirect method correspondingly;  $\bar{E}_{EC}$  is an average daily evaporation over the observational period (ie. 38 days). The values of the  $NSS$  can range from  $-\infty$  to 1, and  $NSS = 1$  indicates to a perfect match of the data modeled after the indirect methods to the data after the EC method; and  $NSS = 0$  indicates that the indirect methods are as accurate as the average of the EC data.

We also applied the  $s/\sigma$  criteria after Popov (1979):

$$s = \sqrt{\frac{\sum_{i=1}^n (E_{EC}^i - E_m^i)^2}{(n-m)}}, \quad (8)$$

$$\sigma = \sqrt{\frac{\sum_{i=1}^n (E_{EC}^i - \bar{E}_{EC})^2}{n}}, \quad (9)$$

where  $E_{EC}$  and  $E_m$  are the evaporation estimates after the direct method and after the indirect method correspondingly,  $n$  is the length of series, and  $m=2$  (a number of the empirical coefficients in the empirical relationship). The criterion value less than 0.8 shows that the indirect method is acceptable for estimations of the evaporation against the EC method.

The region of the study is featured with the persistent katabatic winds blowing from the continental interior. Fig. 3 b shows that almost all winds come from a direction that would be the direction of katabatic winds. However, it is not guaranteed that all these winds are entirely of katabatic origin. Some may be driven by a combined effect of katabatic and synoptic forcing.

## 4 Results

### 4.1 Evaporation

We considered the direct EC method as the most accurate, providing the reference estimates for the daily evaporation over the lake surface (Finch and Hall, 2005; Tanny et al., 2008; Rodrigues et al., 2020). According to the EC method, the daily evaporation varied from 1.5 to 5.0 mm day<sup>-1</sup> with the average being equal to 3.0 mm day<sup>-1</sup>, and the standard deviation was  $\pm 1.1$  mm day<sup>-1</sup>. The average was calculated by dividing 114 mm of evaporated water (which is the sum of the 30-minute series of evaporation) by the number of days in the observational period (which is 38). In this study we used two methods to fill those 18% of the gaps in the 30-minute series: by the median and mean values, however the results differ for only 2 mm. Therefore we decided to use the only median value to fill whole gaps in the 30-minute series of the evaporation. The sum of the evaporation over the period of the field experiment is 94 mm, if we simply excluded the whole gaps in the 30-30-minute series.

We estimated the uncertainties inherent in the indirect methods by comparing their results with those based on the EC method. The bulk aerodynamic method suggests the average daily evaporation was to be 2.0 mm day<sup>-1</sup> calculated by the bulk aerodynamic method with the mass transfer coefficients after Heikinheimo et al. (1999), and this value is approximately 30, which is over 32 % less than those estimated by the result based on the EC method, and it is the best estimate among the. This the best estimates for the average daily evaporation among other indirect methods (bold notation in Table 34). All combination Dalton type semi-empirical equations underestimated the evaporation over the lake surface by over 30 – 75 for over 40 – 72 %, and the method by after Odrova (1979) yielded the greatest maximal underestimation of the


mean daily evaporation over the lake surface. The uncertainties in the estimates by indirect methods are approximately the same for both cases of the input data (Maitri and Irgason).

**Table 34. The daily evaporation (mm day<sup>-1</sup>) over the surface of calculated after indirect methods for the experiment on Lake Zub/Priyadarshini for the period of 38 days of the field experiment (01.01.2018 – 07.02.2018): SD is the standard deviation; r is ratio the sum  $E_{EC}$  divided by the sum  $E_m$ .**


Method	Input data: Irgason site				Input data: Maitri site			
	Min/Max	Mean $\pm$ SD	Sum	r	Min/Max	Mean $\pm$ SD	Sum	r
Bulk aerodynamic (Heikinheimo et al., 1999)	0.6 / 3.5	2.0 $\pm$ 0.8	78	1.5	0.7 / 2.9	1.9 $\pm$ 0.6	72	1.6
Shuttleworth, 1993	0.2 / 1.8	1.0 $\pm$ 0.4	38	3.0	0.1 / 1.9	0.9 $\pm$ 0.4	36	3.2
Penman, 1948	0.0 / 2.0	1.3 $\pm$ 0.5	2.4	1.9	0.1 / 2.2	1.2 $\pm$ 0.5	46	2.5
Doorenbos and Pruitt, 1975	0.0 / 2.9	1.8 $\pm$ 0.8	68	1.7	0.2 / 3.2	1.7 $\pm$ 0.7	66	1.4
Odrova, 1979	0.1 / 1.3	0.8 $\pm$ 0.3	32	3.6	0.1 / 1.6	0.8 $\pm$ 0.3	32	3.6
Methods	Input data: Irgason-site				Input data: Maitri-site			
	Min/Max	Mean $\pm$ SD*	Sum	k**	Min/Max	Mean $\pm$ SD	Sum	k
Bulk-aerodynamic method	0.6 / 3.5	2.0 $\pm$ 0.8	78	1.5	0.7 / 2.9	1.9 $\pm$ 0.6	72	1.6
Penman, 1948	0.0 / 2.0	1.3 $\pm$ 0.5	48	1.9	0.1 / 2.2	1.2 $\pm$ 0.5	46	2.0
Doorenbos and Pruitt, 1975	0.0 / 2.9	1.8 $\pm$ 0.8	68	1.4	0.2 / 3.2	1.7 $\pm$ 0.7	66	1.4
Odrova 1979	0.1 / 1.3	0.8 $\pm$ 0.3	32	2.9	0.1 / 1.6	0.8 $\pm$ 0.3	32	2.9

\*SD is the standard deviation; \*\* k is ratio  $E_{EC}/E_m$ , where  $E_{EC}$  and  $E_m$  are the evaporation estimates after the direct method and after the indirect method correspondingly.


Figure 5 shows the daily evaporation estimated by the direct EC against those estimated by the four indirect methods calculated based on applying the meteorological observations collected at two measurement sites: Maitri and Irgason. There was not a large difference in the results, and therefore we can recommend using the meteorological observations gathered by the nearest site in further estimation of evaporation. Table 5 gives a summary of the scope the evaporation. The efficiency of the indirect methods to model the day-by-day series of the uncertainties and efficiency of the indirect methods to model the day-by-day series of the evaporation with the selected criteria. evaporation was quantified by applying the Pearson correlation coefficient (R), the Nash-Sutcliffe index (NSI) and the  $s/\sigma$  criteria (SSC), and Table 5 shows the values of these criteria.




a)




b)




c)




d)




(a)



(b)



(c)



(d)

**Figure 5: Scatter plots of the daily evaporation estimated with the indirect methods (Y-axis) against the direct EC method (X-axis): (a) the bulk aerodynamic; (b) Penman; (c) Doorenbos and Pruitt; (d) Odrova.  $R^2$  refers to the determination coefficient. The red dots indicate the estimates of the evaporation with the meteorological parameters measured at the WMO synoptic site Maitri, which is the nearest site to Lake Zub/Priyadarshini. The black dots indicate the estimates of the evaporation done with the meteorological parameters measured at the lake shore (Irgason site).**

615

The bulk aerodynamic method gave the best fit to the EC method according to all criteria (bold notation in Table 45). As one can expect, the efficiency of the Dalton-type semi-empirical equations is poor: the correlation coefficient varied from 0.33±2

620 to 0.5534, and both the *RMSE\_NSS* and SSC criteria indicate the indicated a low ability of the methods to estimate in estimations of the daily evaporation.-


**Table 45. The efficiency of the indirect methods with the Pearson correlation coefficient (*PRR*), the root square standard error (*RMSE\_Nash-Sutcliffe index (NSI)*) and the  $s/\sigma$  s-sigma criteria (*SSC*).**

Method	Input data: Irgason site			Input data: Maitri site		
	<i>PR</i>	<i>RMSE</i>	<i>SSC</i>	<i>PR</i>	<i>RMSE</i>	<i>SSC</i>
Bulk aerodynamic (Heikinheimo et al., 1999)	<b>0.87</b>	<b>1.0</b>	<b>1.1</b>	<b>0.88</b>	<b>1.1</b>	<b>1.2</b>
Shuttleworth, 1993	0.55	2.1	2.3	0.39	2.2	2.3
Penman, 1948	0.35	1.8	2.0	0.41	2.1	2.0
Doorenbos and Pruitt, 1975	0.43	1.3	1.6	0.46	1.6	1.6
Odrova 1979	0.35	2.2	2.4	0.45	2.4	2.4

625 The bulk-aerodynamic method also yields the best estimates for day-to-day time series of evaporation (Table 4). However, even this method cannot be suggested to match the daily evaluation of evaporation using the meteorological observations at the Maitri site (Fig. 6). The mean difference between the daily evaporation estimated by the EC and the bulk-aerodynamic method is 0.6 mm day<sup>-1</sup>, and it is the greatest on those days when the wind speeds are 6 – 7 m s<sup>-1</sup>. Therefore, the relationship between evaporation and 2-meter wind speed and saturation deficit was approximated by the formula reading as  $E = a + bw_2(e_s - e_2)$ , and it's similar to the combination equations (given in form  $E = a(1 + abw_2)(e_s - e_2)$  in Table 3), where the saturation deficit ( $e_s - e_2$ ) is expressed in (kPa), and two empirical coefficients (*a* and *b*) were evaluated from the series of the evaporation (after the EC method) and the wind speed and air temperature observations done at Maitri site, which is nearest to Lake Zub/Priyadarshini. The daily series for the period lasting from 01.01.2018 to 07.02.2018 was used in the fitting procedure. Figure 6 shows the daily evaporation estimated by the EC method, by the bulk aerodynamic method and new combination equation with two empirical coefficients fitted from the observations.

Methods	Input data: Irgason site			Input data: Maitri site		
	<i>R</i>	<i>NSI</i>	<i>SSC</i>	<i>R</i>	<i>NSI</i>	<i>SSC</i>
Bulk aerodynamic-	<b>0.87</b>	<b>-0.1</b>	<b>1.1</b>	<b>0.88</b>	<b>-0.5</b>	<b>1.2</b>
Penman, 1948	0.33	-2.7	2.0	0.41	-2.8	2.0
Doorenbos and Pruitt, 1975-	0.40	-1.3	3.3	0.46	-1.3	3.3
Odrova 1979	0.37	-4.2	2.3	0.45	-4.1	2.3

635 The bulk aerodynamic method also allow the best estimates for the day-by-day series of the evaporation (Table 5), however even this method cannot be suggested to evaluate the daily evaporation using the meteorological observations at Maitri site (Fig. 6). The mean difference between the daily evaporation estimated after the EC and bulk-aerodynamic method is  $0.6 \text{ mm day}^{-1}$ , and it takes maximum in days with the winds with speed of  $6-7 \text{ m s}^{-1}$  (Fig. 6 and Fig. 4 b).



640 **Figure 6. The daily time series of evaporation ( $\text{mm day}^{-1}$ ) calculated by the after the direct EC method (black), by the indirect bulk-aerodynamic method (blue) and by new combination equation (red) applying the meteorological measurements at the Maitri site (red).**

645 The daily evaporation was estimated to be  $3.3 \pm 1.6 \text{ mm day}^{-1}$  (where the numbers represent the mean and standard deviation, respectively) by the equation  $E = -0.33 + 0.60w_2(e_s - e_2)$ ; and sum of the evaporation for the period 38 days by this method differs for less than 10 % from those estimated by the EC method. It is the lowest difference for the indirect methods considered; the Pearson correlation coefficient and the mean root square standard error are estimated to be 0.59 and 1.0, respectively. These scopes allow us to consider this equation the second best among the indirect methods (Table 3), the only bulk aerodynamic method showing the better scope. However, these estimates are done on the similar data as the empirical coefficients were fit, and the independent data are needed.




The efficiency of the empirical formula  $E = -0.33 + 0.60w_2(e_s - e_2)$  with the independent data was estimated from the wind speed and air temperature measured at Irgason site (Fig. 1 c). We also used the lake water surface temperature measured at iBuntton site for the period of 27.01.2018 – 07.02.2018 (or 12 days); the daily series of the evaporation were calculated with this formula and then they were compared with those estimated after the EC method. The Pearson correlation coefficient and the mean root square standard error are estimated to be 0.68 and 1.3, respectively. The sum of the evaporation for the period 12 days by this method is over 30 % higher than those estimated by the EC method. It shows that new combination formula may tend to overestimate the evaporation.

## 4.2 Impact of katabatic winds on evaporation

### 4.2 Empirical models

The empirical coefficients limite applications of the Dalton type equations to regions where these coefficients are obtained, and there are no any suggestions given for Antarctica (Finch and Hall, 2001). We further suggested the regional empirical relationship to applying the data on the daily evaporation estimated after the direct EC method and observations collected at the meteorological site nearest to the lake. The evaporation ( $\text{mm day}^{-1}$ ) was evaluated with the linear regression model  $a + b_1w_2 + b_2(e_s - e_2)$ , where  $(e_s - e_2)$  is expressed in mbar. We estimated the empirical coefficients in this relationship based on the whole observations as well as on two subsets collected in two periods. To evaluate how effective the empirical relationship in simulating observations is, the verification with the independent observations is needed. In case of using whole observations in fitting of the empirical coefficients, it was not possible to fully estimate the quality of the model since no independent measurements of evaporation were in rest. Therefore, the verification of the fitted regression models was performed applying the cross validation procedure: the empirical coefficients were estimated with the data collected during the period of 01.01.2018 – 19.01.2018 (19 days), and this linear regression was applied to simulate the daily evaporation for the period of 20.01.2018 – 07.02.2018 (Fig. 7). Then, the procedure was repeated backward: the linear regression was evaluated from the data collected over the period of 20.01.2018 – 07.02.2018, and it was used to model the daily evaporation for the period of 01.01.2018 – 19.01.2018.



**Figure 7. The daily series of the evaporation evaluated after the direct EC method (black) and applying the linear regression with the empirical coefficients estimated from data collected during various periods: 01.01.2018–07.02.2018 (grey), 01.01.2018–19.01.2018 (blue) and 20.01.2018–07.02.2018 (red).**

675

Table 6 shows the estimates of the empirical coefficients in the linear relationship  $a + b_1 w_2 + b_2 (e_s - e_2)$ , which were calculated applying three different subsets of data (Fig. 7). The parameter  $b_1$  is very similar, being estimated from three different subsets. The estimates of two parameters ( $a$  and  $b_2$ ) are also similar for the subsets 1 and 2. The estimates of the parameters varied substantially between the subset 3 and other two subsets. The value of the Pearson correlation coefficient is highest for the subset 2, when the value of the residual standard error is minimal.

680

**Table 6. Estimates of the efficiency indexes ( $R^2$ ,  $R$ ) and empirical coefficients ( $a$ ,  $b_1$ ,  $b_2$ ) in the linear regression model to evaluate the daily series of evaporation based on the observations at Maitri site.**

Subset of observations	$a$	$b_1$	$b_2$	$R^2$	$R$	$RSE$	$RMSE$	$N$
Subset 1: 38 days	-0.37	0.40	0.88	0.40	0.37	0.84	—	35
Subset 2: 19 days	-0.46	0.45	0.86	0.56	0.50	0.82	0.87	16

Subset 3: 19 days	-1.20	0.44	1.21	0.28	0.18	0.89	0.91	16
-------------------	-------	------	------	------	------	------	------	----

Notation: *RSE* or a residual standard error is the average variation of points around the fitted regression line (the lower the

685 *RSE*, the better model is); *RMSE* or a root square standard error is estimated as follows:  $\sqrt{\sum_1^n (E_{EC} - E_{mod})^2 / n}$ ; *N* is the

degree of freedom calculated as the length of the subset minus the number of empirical coefficients in the linear regression.

Further the day-by-day series of the evaporation were estimated with the empirical coefficients evaluated for the subset 2  
 (boulded values in the Table 6) for the whole of the field experiment. The sum of evaporation over the period of 38 days is  
 120 mm, and it is over 5 % larger than the sum estimated after the direct EC method. The daily evaporation varies from 1.7  
 690 to 5.1 mm day<sup>-1</sup>, with the average taking 3.2 mm day<sup>-1</sup> and the standard deviation 0.8 mm day<sup>-1</sup>, and it is only a bit larger than  
 for the EC method.

### 4.3 Impact of katabatic winds on evaporation

The study region is dominated by winds from the south-easterly sector (Fig. 3 b). This corresponds to the  
 695 katabatic winds, which the Coriolis force has turned left from the direct down-slope direction. To better understand the  
 impact of katabatic winds, we carried out further analyses on the wind conditions in the study region. We calculated the  
 geostrophic wind fields for each day of the study period from the mean sea level pressure fields estimated from the ERA5  
 reanalysis. The results demonstrated that the geostrophic (synoptic) wind was mostly from the east, i.e., some 45 degrees  
 right from the mean direction of the observed near-surface wind. This deviation angle may partly result from the Ekman  
 700 turning in the atmospheric boundary layer, which over an ice sheet with a rather small aerodynamic roughness may  
 contribute some 20 degrees, and from the katabatic forcing. In any case, in most cases the observed near-surface winds  
 resulted from the combined effects of synoptic and katabatic forcing, which supported each other. Hence, it is very difficult  
 to robustly distinguish the impact of katabatic forcing on the near-surface winds over the lake.

However, the geostrophic wind direction was distinctly different, 240 – 350<sup>o350o</sup>, in the following days: 6, 8 – 10, 19 and 25 –  
 705 27 January. These days were related to transient cyclones centred north-west of the lake or high-pressure  
centres north-east of the region under the study. During the days, the wind speed over the lake was strongly  
 reduced (Table 57), as the katabatic and synoptic forcing factors opposed each other. The lake surface temperature was  
 higher than usual, but the air temperature was lower. The latter is partly because, during events when the geostrophic and  
 katabatic forcing factors support each other (sector 60 – 130°), the strong wind effectively mixes the atmospheric boundary  
 710 layer. In stably stratified conditions, which prevail over the ice sheet, vertical mixing results in higher near-surface air  
 temperatures (Vihma et al., 2011). In addition, the adiabatic warming during the downslope flow is a major factor  
 contributing to higher air temperatures (Xu et al., 2021). The impact of adiabatic warming is also seen as lower relative  
 humidity in cases when the geostrophic wind is from the sector 60 – 130<sup>o130o</sup>. Related to the compensating effects of air  
 temperature and relative humidity, the specific humidity was not sensitive to the geostrophic wind direction. The effect of

715 the wind speed dominated the effect of the lake surface temperature (which controls  $q_{ses}$  in Eq. (q3)), and evaporation was strongly reduced when the geostrophic wind was from the sector 60 – 130°<sup>±30°</sup> (Table 57).

**Table 57. The mean values of the evaporation ( $E_{EC}$ ), the wind speed ( $w_2$ ), air specific humidity ( $Q_2$ ), and lake surface temperature ( $w_s$ ), and air temperature ( $t_2$ ) calculated over the days when the geostrophic wind direction was 60 – 130° and when it was 240 – 350°.**

Geostrophic wind dir.	Evaporation (mm day <sup>-1</sup> )	$w_2$ (m s <sup>-1</sup> )	$Q_2$ (g kg <sup>-1</sup> )	$t_w$ (°C)	$t_2$ (°C)
60 – 130°	3.1	6.9	2.0	3.6	-0.2
240 – 350°	1.3	2.3	2.0	4.8	-2.8

Geostrophic wind dir.	Evaporation (mm day <sup>-1</sup> )	$w_2$ (m s <sup>-1</sup> )	$Q_2$ (g kg <sup>-1</sup> )	$w_s$ (°C)	$t_2$ (°C)
60 – 130°	3.1	6.9	2.0	3.6	-0.2
240 – 350°	1.3	2.3	2.0	4.8	-2.8

720 The katabatic wind was a quasi-persistent feature during the study period, and the major changes in the evaporation were driven by changes in the synoptic scale wind direction, which affected the local wind speed.

## 5 Discussion

725 Our study yielded estimates of evaporation over a glacial lake in the summer based on direct EC measurements during a field experiment lasting 38 days. These direct estimates of evaporation were considered as the reference when estimating the uncertainties inherent in the indirect methods including the bulk-aerodynamic method and four combination equations. The results based on the bulk-aerodynamic method reached the best skill scores based on the efficiency indexes, however, this method underestimated the daily evaporation by over 30 %. The efficiency of all selected combination equations was low: they underestimated the mean daily evaporation by up to 72 %. The empirical coefficients for the combination equation were fitted from the series of the evaporation (by EC method) and the meteorological observations at the station nearest to the lake site. This combination equation can be potentially used in estimations of the evaporation over the ice-free glacial lakes located in Schirmacher oasis. However, in this study the estimations of the daily evaporation and efficiency indexes were performed on the same data for the experiment (lasting 38 days). Also, we estimated the efficiency using the independent data on the air temperature, wind speed and lake surface temperature. The estimations of efficiency indexes were also done with the full independent data including the evaporation estimated by the EC method, therefore we would not suggest applying these coefficients as the regional references without further analysis. In this study, we did not estimate the evaporation using the energy balance method, but plan to further evaluate the uncertainties inherent also in this method while estimating the evaporation over the glacial lakes located in Antarctica.

740 At monitoring sites, evaporation over lakes is in practice measured with evaporation pans, which are not fully applicable in polar regions. The EC measurements require specific equipment not always possible to deploy and operate in the remote Antarctic continent. Hence, evaporation (or sublimation) over lakes is usually estimated only indirectly on the basis of regular or campaign observations or numerical model experiments. There are only a few studies of evaporation over lakes located in Antarctica. Borghini et al. (2013) propounded estimates of evaporation over a small endorheic lake located on the

745 shore of Wood Bay, Victoria Land, East Antarctica (70° S). This lake is of 0.8 m depth, and by early 2000s its surface area has decreased to half of the value in late 1980s. The lake is of the landlocked type, and Borghini et al. (2013) used the method by Shuttleworth (1993) to estimate the evaporation from the lake surface during a couple of weeks in December 2006. They estimated the mean daily evaporation as  $4.7 \pm 0.8 \text{ mm day}^{-1}$ ; and such an evaporation rate results in loss of over  $40 \pm 5 \%$  of the total volume of the lake during the observation period. The lake studied by Borghini et al. (2013) differs from Lake Zub/Priyadarshini, but the daily evaporation rates are of the same order of magnitude, and although one could expect a much larger evaporation from landlocked lakes than glacial lakes. Our results show that method by Shuttleworth (1993) underestimates the evaporation of lakes located in the Schirmacher oasis by over 60 %.

750 The estimations of the lake volume of the glacial lakes and the time scale of their water exchange are sensitive to the uncertainties inherent in various methods applied to evaluate evaporation (Shevnina et al., 2021). Our study yielded estimates of the evaporation over glacial lakes in summer after the direct EC method, and the results are based on the data collected during the field experiment lasting 38 days. These estimates of the evaporation were considered as most accurate (or reference) while estimating the uncertainties inherent in the indirect methods including the bulk aerodynamic method and three Dalton type empirical equations. The results after the bulk aerodynamic method reached the best skill scores based on the efficiency indexes, however it underestimated the daily evaporation of over 30 %. The efficiency of the selected Dalton type semi-empirical equations was low, and they underestimated the mean daily evaporation up to 72 %. We suggested the regional empirical relationship to simulate the evaporation from the observations at the nearest meteorological site and water temperatures measured in the lake. We suggested applying this regional empirical relationship in simulations of day-by-day series of evaporation over the ice-free surface of the lakes in Antarctica. We did not apply the energy balance method in this study, and we plan to further use also this method in estimations of the evaporation over the glacial lakes. It allows evaluation of the uncertainties inherent also to this method.

765 At monitoring sites, the evaporation over the lakes is practically measured with evaporation pans, which are not fully applicable in polar regions. The EC method requires specific equipment not always possible to deploy and operate in the remote Antarctic continent. Hence, the evaporation (or sublimation) over the lakes is usually estimated only indirectly on the basis of the regular or campaign observations or numerical model experiments. There are only a few studies of the evaporation over the lakes located in Antarctica. Borghini et al. (2013) propounded estimates of evaporation over a small endorheic lake located on the shore of the Wood Bay, Victoria Land, East Antarctica (70° S). This lake is of 0.8 m depth and its surface has decreased more than twice since late 1980s to early 2000s (from  $4.0 \times 10^3$  to  $2.0 \times 10^3 \text{ m}^2$ ). The lake is the landlocked type, and therefore the evaporation is an important outflow term of the lake's water budget. The authors use the semi-empirical equation given in Shuttleworth (1993) to calculate the evaporation from the ice-free water surface with data on the water temperature, air temperature, and wind speed collected during a couple of weeks in December, 2006. The mean daily evaporation was estimated to be  $4.7 \pm 0.8 \text{ mm day}^{-1}$ . These estimates result in the loss of the liquid water at  $40 \pm 5 \%$  of the total volume of the lake during the observational period. Although the lake studied by Borghini et al. (2013) differs from Lake Zub/Priyadarshini, the daily evaporation rates are of the same order of magnitude even one can expect a much larger

~~evaporation over the surface of the landlocked lakes rather than over the glacial lakes.~~

780 Shevnina and Kourzeneva (2017) used two indirect methods to evaluate ~~the~~ daily evaporation for two lakes located in the Larsemann Hills oasis, East Antarctica (69° S). Lake Progress and Lake Nella/Scandrett are of the glacial type, however, they are much deeper and larger in ~~the~~ volume than Lake Zub/Priyadarshini, and over 30—70 % ~~of~~ their catchments are covered by the glacier. The thermal regime of these glacial lakes is also different: Lake Nella/Scandrett and Lake Progress have partially lost their ice cover in austral summers when their surface water temperature is only 2–3 °C, which is lower than the water temperature over the surface of Lake Zub/Priyadarshini. The daily evaporation was estimated to be 1.8 mm day<sup>-1</sup> and 1.4 mm day<sup>-1</sup> on the basis of the energy budget method (Mironov et al., 2005) and ~~by the equation of the semi-~~  
785 ~~empirical equation after~~ Odrova (1979), respectively. ~~Shevnina and Kourzeneva (2017) It is~~ concluded that ~~the~~ daily evaporation over ~~the~~ glacial lakes is underestimated by both ~~of these~~ indirect methods. Our results prove that the uncertainties ~~inherent in the method by of the semi-empirical equation after~~ Odrova (1979) are the largest among other considered methods.-


790 ~~Faucher et al. (2019) evaluated the annual cycle of the terms of the water balance equation written for Lake Untersee, Dronning Maud Land, East Antarctica (71° S). Lake Untersee is perennially frozen year-round; it is the glacial type lake directly attached to the continental ice sheet, not being the landlocked.. The sublimation (evaporation) over the lake surface was estimated in terms of its water budget. These estimations were based on two years of in situ measurements using snow sticks. Faucher et al. (2019) estimated the water losses from the ice-covered surface of the lake due to sublimation to be from 400 to 750 mm year<sup>-1</sup>. The daily evaporation from the lake surface was approximately 1.1–2.1 mm day<sup>-1</sup>. Dhote et al. (2021)~~  
795 ~~estimated the summertime evaporation over Lake Zub/Priyadarshini using the meteorological observations collected at the Maitri site. The sum of the evaporation over the lake surface was estimated to be 167 mm for January and February 2018 (or 2.8 mm day<sup>-1</sup>), and this estimate is very close to those based on the EC method given in this study.~~

800 ~~Faucher et al. (2019) evaluate the annual water budget for Lake Untersee, the Dronning Maud Land, East Antarctica (71° S). Lake Untersee is perennially frozen year-round, it is the glacial type lake directly attached to the continental ice sheet. Lake Untersee is not the landlocked lake as mentioned in Faucher et al. (2019). The sublimation (evaporation) over the lake surface is estimated among other terms of its water budget. These estimations are based on two years of the in-situ measurements using the snow sticks. The authors estimate the water losses from the ice covered surface of the lake due to sublimation from 400 to 750 mm year<sup>-1</sup>. The daily evaporation from the lake surface is approximately 1.1–2.1 mm day<sup>-1</sup>.~~

805 ~~This study focused~~~~focuses~~ on the evaporation over a glacial lake ~~located in the Schirmacher oasis, East Antarctica. Over 65 thousand glacial lakes have been detected in the coastal region via satellite remote sensing in austral summer 2017, and most of them spread over the ice shelf and the margins of the continental ice sheet (Stokes et al., 2019). The total area of which is ice free during almost 2 months in summer. Seasonal presence of the liquid water (ie. in glacial lakes in vicinity of the Schirmacher oasis was over 72 km<sup>2</sup> in January 2017 (Fig. 7), and the two largest glacial lakes being of a similar size as the Schirmacher oasis itself. During warm periods, a high number of glacial lakes (or melt ponds) are recognized and iceed~~  
810 ~~“swamps”)~~ over the ~~margins of the Greenland ice sheet (How et al., 2021), and melt ponds are very common also on the~~

815 surface of Arctic sea ice (Lu et al., 2018). The glacial lakes may exist over the snow/ice covered surface for 1 – 3 months, and their presence has changed land cover properties and affected the surface heat budget/ice/snow covered land surface affects the surface-atmosphere moisture exchange. A proper description of the land cover is a crucial element of numerical weather prediction (NWP) predictions (NWP) and climate models, where the overall characteristics of the land cover are represented by the surfaces covered by ground, whether vegetation, urban infrastructure, water (including lakes), bare soil or other. Various parameterization schemes (models) are applied to describe the surface-atmosphere moisture exchange and surface radiative budget (Viterbo, 2002). Lakes have been recently included in the surface parameterization schemes of many NWP models NWP (Salgado and Le Moing, 2010; Dutra et al., 2010; Balsamo et al., 2012) with known external parameters (location, mean depth) available from the Global Lake Database, GLDB (Kourzeneva, 2010; Kourzeneva et al., 820 2012). The newest version of the GLDB includes glacial lakes information on only a few glacial lakes is included in the newest GLDBv3 version, and not any lakes found in Antarctica (Toptunova et al., 2019). In future studies, it is important to understand how Over 65 thousand glacial lakes have been detected over the East Antarctic coast via satellite remote sensing in austral summer 2017, and most of them spread over the ice shelf and margins of the continental ice sheet (Stokes et al., 2019). For example, the total area of the glacial lakes affect the regional air moisture transport in vicinity of the Schirmacher 825 oasis was over 72 km<sup>2</sup> in January 2017 (Fig. 8), two largest glacial lakes are of the similar size as the Schirmacher oasis itself. Such amount of the liquid water over the polar regions and local weather/ice/snow covered region may contribute the additional source of the uncertainties inherent in the NWP.





**Figure 78.** The glacial lakes over the surface of an ice shelf in the vicinity of the Scirmacher oasis, East Antarctica.

830 Estimates of evaporation are available from atmospheric reanalyses which share results of simulations done by NWP models. As for other reanalyses, ERA5 does not assimilate any evaporation observations, and the evaporation is based on 12-hour forecasts of an NWP model by applying the bulk-aerodynamic method. The results naturally depend on the presentation of the Earth's surface in ERA5, and in Dronning Maud Land, the surface type is ice and snow with no lakes. Therefore, the estimate of evaporation does not include evaporation from liquid water surfaces. We estimated the daily evaporation also

835 from ERA5, and the results suggest that the evaporation during summer (December – February) 2017 – 2018 was 0.6 mm day<sup>-1</sup>. This is only one fifth of the evaporation estimated with the direct EC method.

The estimates of the evaporation are also available from atmospheric reanalyses which share results of simulations done by the NWP. The most recent global atmospheric reanalysis is ERA5 of the European Centre for Medium-Range Weather Forecasts (Copernicus Climate Change Service, <https://climate.copernicus.eu/>, last access 09.07.2021; Hersbach et al.,

840 2020). As other reanalyses, ERA5 does not assimilate any evaporation observations, but evaporation is based on 12-h accumulated NWP forecasts applying the bulk aerodynamic method. The results naturally depend on the presentation of the Earth surface in ERA5, and in the Dronning Maud Land, the surface type is ice and snow with no lake. Therefore the

estimate of the evaporation does not include evaporation from liquid water surface. We estimated the daily evaporation also from the ERA5, and the results suggest that the evaporation during summer (DJF) 2017–2018 was 0.6 mm day<sup>-1</sup>. It is only one fifth of the evaporation estimated with the direct EC method.

Naakka et al. (2021) estimated the evaporation over the Antarctic region from the ERA5 reanalysis for five domains, including the East Antarctic slope where the Schirmacher oasis is located. There the average daily evaporation in summer is 0.3 mm day<sup>-1</sup>, and this is reasonable for the ice/snow covered surface. In summertime, the presence of the liquid water over ice/snow covered surface changes the fraction of the lakes over the East Antarctic slope, and it is 6–8 % of the region in the vicinity of the Schirmacher oasis (Fig. 78). The increasing numbers of the glacial lakes over the surface of the East Antarctic slope affects the surface-atmosphere moisture interactions, and it also changes the regional evaporation not accounted for by the numerical weather prediction systems and climate models. We assumed that the 0.3 mm of ERA5 is a fair value for the ice sheet in the East Antarctic slope and that 3 mm is a representative value for the glacial lakes, and it may add up to 0.16–0.22 mm to the regional summertime evaporation over the margins of the East Antarctic slope. These numbers seem to be insignificant for the mass balance of the Antarctic ice sheet and ice shelves. However, we suggested more comprehensive research to better understand and quantify the impact of glacial lakes on the surface heat budget and atmospheric moisture transport in the summer-atmosphere moisture exchange and surface radiative budget in summer periods.

## 6 Conclusions

This study suggested the estimates of summertime evaporation over an ice-free surface of Lake Zub/Priyadarshini the glacial lake applying the direct EC method. Evaporation was also evaluated using six indirect methods only needing as input a few hydrometeorological parameters monitored at selected sites (e.g., WMO stations). Our study focused on the glacial Lake Zub/Priyadarshini has located in the Schirmacher oasis, Dronning Maud Land, East Antarctica. The catchment of the lake includes less than 30 % of the area covered by glaciers, which with the glacier, and results in a specific thermal regime and water balance for the lake. We estimated the evaporation over ice free lake surface as 114 mm in the period from 1 January to 7 February 2018 on the basis of the direct EC method. The daily evaporation is estimated to be 3.0 mm per day in January 2018. The largest changes in the daily evaporation were driven by the synoptic-scale atmospheric processes rather than local katabatic winds. This study gave the estimations of the uncertainties inherent in the indirect methods applied to evaluate summertime evaporation over a lake surface. The bulk aerodynamic method suggests the average daily evaporation to be 2.0 mm day<sup>-1</sup>, which is over 32 % less than the result based on the EC method. Four selected combination of the selected Dalton-type semi-empirical equations underestimated the evaporation over the lake surface by over 40–72 %. We suggested a new combination equation to evaluate applying the new regional empirical relationship while estimating the summertime evaporation of Lake Zub/Priyadarshini from meteorological observations from the nearest site. The performance of the new

875 ~~equation is better than the performance of the indirect methods considered over the ice-free glacial lakes located in~~  
~~Antarctica.~~ We ~~also~~ stress the need for accurate measurements of the ~~lake water surface temperature to allow better~~  
~~estimates~~ surface water temperature in local lakes to support studies of lake water budget and evaporation (sublimation).

The evaporation results were not sensitive to differences in the data collected at the meteorological site nearest to the lake  
and the site located on the lake shore. Hence, we suggest using the synoptic records at the meteorological site Maitri to  
880 evaluate the evaporation over the surface of Lake Zub/Priyadarshini. Field experiments are needed to make analogous  
comparisons of meteorological conditions between other glacial lakes and the permanent observation stations nearest to  
them. The water balance terms of ~~the~~ glacial lakes (including evaporation) are closely connected to their thermal regime; and  
coupled thermophysical and hydrological models are needed to predict the amount of water in these lakes. Our results also  
demonstrated the need to present glacial lakes in atmospheric reanalyses as well as ~~NWPs~~ NWP and climate models. Ignoring  
885 them in a lake-rich region, such as the Schirmacher oasis, results in a large underestimation of regional evaporation in ~~the~~  
~~summer.~~

## Annex.

890 **To evaluate the uncertainties of the EC method with the ~~paired tower method~~ method-of-paired-tower: the  
intercalibration experiment at Alqueva reservoir, Portugal.**

The eddy covariance method has some errors and uncertainties associated with the nature of the measurement and the  
instrument system. ~~Therefore,~~ therefore, the results need to be treated with special attention. Nevertheless, the complexity of  
the method, namely the filters and corrections that this method requires (see Section 3.3), make it possible to reduce the  
errors and uncertainties. According to Aubinet et al. (2012), there are three methods to quantify the total random uncertainty  
895 for the eddy covariance method: the paired tower, 24 h differencing, and the model residual. In our study we apply the paired  
tower method to evaluate the errors of the Irgason installed on the shore of Lake Zub/Priyadarshini. The intercalibration  
experiment lasted from 12 October to 25 October, 2018, and during this period two Irgason instruments were deployed on a  
floating platform in Alqueva artificial lake located southeast of Portugal.-

The floating platform (38.2° N; 7.4° W) has been operating continuously since April 2017, and in this experiment, two eddy  
900 covariance stations (Irgason) were installed on the height of 2.0 m next to each other facing the same footprint (Fig. A1). In  
this experiment, we compare the measurements of the Irgason of the Finnish Meteorological Institute (FMI) to those  
collected by the Irgason of the Institute of Earth Sciences (ICT), University of Évora. Taking ~~advantage of the fact~~  
~~advantage~~ that both instruments are identical, the settings were set exactly the same. The standard gas zero and span  
calibration was performed before the experiment. The raw measurements from both instruments were post-processed  
905 applying the algorithm given in Potes et al. (2017). It allows precise estimates of random instrument uncertainty, rather than  
total random uncertainty which demands that both instruments are in the same area but with different footprints (Dragoni et  
al., 2006).-



Figure A1: The instruments installed in Alqueva reservoir (Portugal) for the intercalibration. The left instrument belongs to [the](#) Institute of Earth Sciences, and the instrument on the right belongs to [the](#) Finnish Meteorological Institute.-

910

Figure A2 shows a scatter plot between 30-minute evaporation evaluated [from](#)[after](#) the measurements of two instruments during the intercomparison campaign that took place in Alqueva reservoir. The correlation coefficient between the evaporation calculated [by two Irgason](#)[after two Irgason](#) is over 0.98, and it suggests strong agreement between the measurements. Figure A3 presents the frequency distribution of the 30-minute evaporation random instrument uncertainty ( $\epsilon_F$ ) during the intercomparison campaign (see the Eq. 9 from Dragoni et al., 2007). The random instrument error in 30-minute evaporation, estimated as the standard deviation of the evaporation random instrument uncertainty ( $\epsilon_F$ ), is 0.004324 mm. Thus, in relative terms, the intercomparison campaign allows [obtaining to obtain](#) an estimate of [at](#)[the](#) random instrument error of 7.0 %. This value is below [other](#)[another](#) studies presented by several authors, namely: Eugster et al. (1997), that used the same approach of the paired towers in [Alaskan](#)[Alaska](#) tundra, and obtained 9 % for latent heat flux; Finkelstein and Sims (2001), that present a value between 14 and 35 % for latent heat flux in forest and agricultural sites; [and](#) Salesky et al. (2012), that found typical errors of 10 % for [the](#) heat flux.

915

920





Figure A2: Scatter plot between 30-minute evaporation from both instruments: **the Y-axis shows** the values estimated after the measurements by the Irgason of **the FMI**, and **the X-axis shows** the values after the ~~measurements~~ of the Irgason of **the ICT**.

925





**Figure A3: Frequency distribution of the 30-minute evaporation random instrument uncertainty ( $\epsilon_F$ ).**

**Acknowledgement.** The study was funded by the Academy of Finland (contract number 304345), [the H2020 project PolarRES \(contract 101003590\)](#) and the COST Snow Action ES1404. The measurement campaigns were supported by the Finnish Antarctic Research Program, the Russian Antarctic Expedition, and the Indian Antarctic expedition. We thank Daniela Franz, Ekaterina Kourzeneva and Rui Salgado for the discussions during the 5<sup>th</sup> and 6<sup>th</sup> Workshops on Parameterization of Lakes in Numerical Weather Prediction and Climate Modelling (October 2017, Berlin, Germany, and

October 2019, Toulouse, France). We thank the participants of two scientific conferences held in St. Petersburg in 2020 and  
935 Lisbon (on-line) in 2021 for their questions and comments. Our special thanks to Alexander Piskun, who shared with us his  
private collection of Bulletins of the Soviet Antarctic Expedition (1960–1986). We are grateful to Indian Space Research  
Organisation and National Centre for Polar and Ocean Research, Goa, for constant support and encouragement during this  
work. We thank the editor and two anonymous referees for their fruitful comments and suggestions allowing the  
improvement of the presentation of our results.

940

**Supplement.** ES attached the calculation of the evaporation with the combination equations in Table 3 (combinationsemi-  
empirical equations (Dalton equations\_results.csv) and the code (results\_code.r).code used to evaluate the estimates of the  
uncertainties inherent various methods. MP has attached the post processed byafter the EC method  
(20180101\_20180207\_EC\_FLUX.txt). TV attached the calculations byafter the bulk aerodynamic method  
945 (Bulk\_method\_results\_Irgason\_input.txt and Bulk\_method\_results\_Maitri\_input.txt). PD attached the have attached  
meteorological data measured at the MaitriMaitry site (Meteorological\_Parameters\_Summer\_2017-18.xlsx). TN provided;  
and the water surface temperature measured by Solinst instrument (Lake\_Temperature\_Solinst\_Jan-Feb2018.xlsx). TN  
provides the series of the daily evaporation from the ERA5 reanalysis at the grid note nearest to the Novo meteorological site  
(Evaporation\_Schirmacher\_Oasis\_from\_ERA5.csv). The supplement related to this article is available online at:-

950

**Data and code availability.** The data and code used in this study are available in the Supplement. We also used two datasets  
stored at zenodo: <http://doi.org/10.5281/zenodo.3469570> and <http://doi.org/10.5281/zenodo.3467126>.-

**Author's contribution.** ES initiated the manuscript, and collected the data in the field experiment 2017–2018, and. She also  
955 calculated the evaporation applying the combinational equations, their uncertainties and efficiency indexessemi-empirical  
equations. MP supervised the EC measurements in the field, then he calculated the evaporationhe carried out the calculations  
applying the EC method, and he analysed the data collected during the intercalibration campaign. (including the  
intercalibration campaign). TV contributed towith the estimations of evaporation applying the bulk aerodynamic method. TN  
contributed with analyses of evaporation based on ERA5. TV and TN made the analysis of the impact of the katabatic winds.  
960 PD and PKT contributed with the analysis of the meteorological observations at the Maitri site. All authors contributed to  
writing of the manuscript.-

**Competing interests.** The authors declare that they have no conflictseonffiet of interest.

## References

965 Asthana, R., Shrivastava, P.K., Srivastava, H.B., Swain, A.K., Beg, M.J., and Dharwadkar, A.: Role of lithology, weathering



- and precipitation on water chemistry of lakes from Larsemann Hills and Schirmacher Oasis of East Antarctica, *Adv Polar Sci*, 2019, 30(1): 35-51, doi: 10.13679/j.advps.2019.1.00035, 2019.
- Arthur, J. F., Stokes, C. R., Jamieson, S. S. R., Carr, J. R., and Leeson, A. A.: Distribution and seasonal evolution of supraglacial lakes on Shackleton Ice Shelf, East Antarctica, *The Cryosphere*, 14, 4103–4120, <https://doi.org/10.5194/tc-14-4103-2020>, 2020.
- 970 Aubinet M., Vesala, T., Papale, D. (Eds): *Eddy Covariance: A Practical Guide to Measurement and Data Analysis*, ISBN 978-94-007-2350-4e-ISBN, doi: 10.1007/978-94-007-2351-1, 2012.
- Balsamo, G., Salgado, R., Dutra, E., Boussetta, S., Stockdale T., Potes M.: On the contribution of lakes in predicting near-surface temperature in a global weather forecasting model, *Tellus A: Dynamic Meteorology and Oceanography*, 64:1, doi: 10.3402/tellusa.v64i0.15829, 2012.
- 975 Bell, R., Chu, W., Kingslake, J., Das, I., Tedesco, M., Tinto, K. J., Zappa, C.J., Frezzotti, M., Boghosian, A., Lee, W.S.: Antarctic ice shelf potentially stabilized by export of meltwater in a surface river, *Nature*, 544, 344–348, doi: 10.1038/nature22048, 2017.
- Bell, R., Banwell, A., Trusel, L., Kingslake, J.: Antarctic surface hydrology and impacts on the ice-sheet mass balance. *Nature climate change*, doi: 10.1038/s41558-018-0326-3, 2019.
- 980 Boisvert, L., Vihma, T., Shie, C.L.: Evaporation from the Southern Ocean estimated on the basis of AIRS satellite data. *Journal of Geophysical Research: Atmospheres*, 125, e2019JD030845, <https://doi.org/10.1029/2019JD030845>, 2020.
- Boronina, A.S., Popov, S.V., Pryakhina, G.V.: Hydrological characteristics of lakes in the eastern part of the Broknes Peninsula, Larsemann Hills, East Antarctica, *Led i Sneg*, 59 (1), 39–48, doi: 10.15356/2076-6734-2019-1-39-48, 2019. (In Russian).
- 985 Bormann, P., Fritzsche, D.: *The Schirmacher Oasis, Queen Maud Land, East Antarctica, and Its Surroundings*. Justus Perthes Verlag Gotha, Darmstadt, 448 pp., 1995.
- Borghini, F., Colacevich, A., Loisel, S.A., Bargagi, R.: Short-term dynamics of physico-chemical and biological features in a shallow, evaporative antarctic lake. *Polar Biol*, 36, 1147–1160, <https://doi.org/10.1007/s00300-013-1336-2>, 2013.
- 990 Blanken, P. D., Rouse, W. R., Culf, A. D., Spence, C., Boudreau, L. D., Jasper, J. N., Kochtubajda, B., Schertzer, W.M., Marsh, P., and Verseghy, D.: Eddy covariance measurements of evaporation from great slave Lake, northwest territories, Canada. *Water Resources Research*, 36(4), 1069–1077, <https://doi.org/10.1029/1999WR900338>, 2000.
- [Braun, M., H. Saurer, S. Vogt, et al., : The influence of large-scale atmospheric circulation on the surface energy balance of the King George Island ice cap. \*Int. J. Climatol.\*, 21, 21–36, doi: 10.1002/joc.563, 2001.](#)
- 995 [Brunke, M. A., Fairall, C. W., Zeng, X., Eymard, L., & Curry, J. A. \(2003\). Which bulk aerodynamic algorithms are least problematic in computing ocean surface turbulent fluxes? \*Journal of Climate\*, 16\(4\), 619– 635. <https://doi.org/10.1175/1520-0442>.](#)
- [1. Braslavskiy, A. Calculation on evaporation rate from freshwater reservoirs located on flat areas. In book-](#)

- 1000 [Materials for a committee on study of methods to calculate an evaporation from water/land surfaces. GGI, Valday, 1966. \(in Russian\)](#)
- Brutsaert, W.: Evaporation into the atmosphere - theory, history and applications, Dordrecht, Holland: D Reidel Publishing Company, 299 pp., 1982.
- Burba, G.: Eddy Covariance Method for Scientific, Industrial, Agricultural, and Regulatory Applications: A Field Book on Measuring Ecosystem Gas Exchange and Areal Emission Rates. LI-COR Biosciences, Lincoln, NE, USA, 331 p., 2013.
- 1005 Businger, J. A., Wyngaard, J. C., Izumi, Y., and Bradley, E. F.: Flux-Profile Relationships in the Atmospheric Surface Layer, *Journal of Atmospheric Sciences*, 28(2), 181–189, [https://doi.org/10.1175/1520-0469\(1971\)028<0181:FPRITA>2.0.CO;2](https://doi.org/10.1175/1520-0469(1971)028<0181:FPRITA>2.0.CO;2), 1971.
- [Chebotarev A.I. Hydrology. Leningrad: Hydrometizdat; 1975. 544 p. \(in Russian\)](#)
- [DorenbosDørenbos, J., Pruitt, W.O.: Crop water requirements, FAO Irrigation and Drainage Paper No. 24 FAO Rome, 179](#)
- 1010 [pp., 1975.](#)
- Dragoni, D., Schmid, H.P., Grimmond, C.S.B., Loescher, H.W.: Uncertainty of annual net ecosystem productivity estimated using eddy covariance flux measurements. *J Geophys Res Atmos*, 112, D17102, 2007.
- Dutra, E., Stepanenko, V. M., Balsamo, G., Viterbo, P., Miranda, P. M. A., Mironov, D., and Schär, C.: An offline study of the impact of lakes on the performance of the ECMWF surface scheme. *Boreal Environment Research*, 15, 100-112, 2010.
- 1015 [Edinger, J. E., Duttweiler, D. W., Geyer, J. C.: The response of water temperature to meteorological conditions. \*Water Resources Research\*, 4, 1137-1143, 1968.](#)
- [Ellehoj, M.D., Steen-Larsen, H.C., Johnsen, S.J., Madsen, M.B.: Ice-vapor equilibrium fractionation factor of hydrogen and oxygen isotopes: Experimental investigations and implications for stable water isotope studies. \*Rapid Commun. Mass Spectrom.\*, 27\(19\): 2149—2158, 2013.](#)
- 1020 Eugster, W., McFadden, J.P., Chapin, E.S.: A comparative approach to regional variation in surface fluxes using mobile eddy correlation towers, *Bound Layer Meteorol*, 85, 293–307, 1997.
- Hersbach, H, Bell, B, Berrisford, P, et al.: The ERA5 global reanalysis. *Q J R Meteorol Soc.*, 146, 1999–2049. <https://doi.org/10.1002/qj.3803>, 2020.
- [Holtslag, A. A. M., De Bruin, H. A. R.: Applied Modeling of the Nighttime Surface Energy Balance over Land, \*Journal of Applied Meteorology and Climatology\*, 27\(6\), 689-704, 1988.](#)
- 1025 [Faucher, B., Lacelle, D., Fisher, D., Andersen, D., McKay, C.: Energy and water mass balance of Lake Untersee and its perennial ice cover, East Antarctica, \*Antarctic Science\*, 31\(5\), 271-285. doi:10.1017/S0954102019000270, 2019.](#)
- [Favier, V., C. Agosta, C. Genthon, L. Arnaud, A. Trouvillez, and H. Gallée: Modeling the mass and surface heat budgets in a coastal blue ice area of Adelie Land, \*Antarctica J. Geophys. Res.\*, 116, F03017, doi:10.1029/2010JF001939, 2011.](#)
- 1030 Finch, J. W., Hall, R. L.: Estimation of Open Water Evaporation: A Review of Methods, R&D Technical Report W6-043/TR. Environment Agency, Bristol, 155 pp., 2001.

- Finch, J. W., Hall, R. L.: Evaporation from Lakes. Encyclopedia of Hydrological Sciences, Part 4 Hydrometeorology Centre for Ecology and Hydrology, Wallingford. pp. 635–646, 2005.
- 1035 [Finch, J.W., Calver, A.: Methods for the quantification of evaporation from lakes. Prepared for the World Meteorological Organization's Commission for Hydrology, Oxfordshire, UK, 41 p., 2008.](#)
- Finkelstein, P. L., Sims, P. F.: Sampling error in eddy correlation flux measurements, *J. Geophys. Res.*, 106, 3503–3509, 2001.
- [Foken, T.: 50 Years of the Monin–Obukhov Similarity Theory. \*Boundary-Layer Meteorol\* 119, 431–447, <https://doi.org/10.1007/s10546-006-9048-6>, 2006.](#)
- 1040 Gopinath, G., Resmi, T. R., Praveenbabu, M., Pragatha, M., Sunil, P. S., Rawat, R.: Isotope hydrochemistry of the lakes in Schirmacher Oasis, East Antarctica, *Indian Journal of Geo Marine Sciences*, 49 (6), 947–953 p., <http://nopr.niscair.res.in/handle/123456789/54944>, 2020.
- Ingole, B.S., and Parulekar, A.H.: Limnology of Priyadarshini Lake, Schirmacher oasis, Antarctica. *Polar record*, 26 (156), 13–17, 1990.
- 1045 Golubev G.N. Hydrology of glaciers. *Gidrometeoizdat, Leningrad*, 128 pp., 1976, (in Russian).
2. ~~[Guide to meteorological instruments and methods of observations, WMO No 8: Geneva, Switzerland, ISBN 978-92-63-10008-5, 681 p., 2008.](#)~~
- Heikinheimo, M., Kangas, M., Tourula, T., Venäläinen, A., Tattari, S.: Momentum and heat fluxes over lakes Tämnaaren and Råksjö determined by the bulk-aerodynamic and eddy-correlation methods, *Agricultural and Forest Meteorology*, 98–99, ~~521–521~~–534, [https://doi.org/10.1016/S0168-1923\(99\)00121-5](https://doi.org/10.1016/S0168-1923(99)00121-5), 1999.
- 1050 Hodgson, D. A.: Antarctic lakes, in: *Encyclopedia of Lakes and Reservoirs. Encyclopedia of Earth Sciences Series*, Springer, Dordrecht, doi:10.1007/978-1-4020-4410-6, 2012.
- Hoeltgebaum L.E. B., Diniz A. L., Dias, N. L. C., Intercomparação de sensores de temperatura e umidade relativa para uso em campanha micrometeorológica, *Ci. e Nat.*, Santa Maria v.42, Special Edition: Micrometeorologia, e18, 1055 doi:10.5902/2179460X46565, 2020. Available online <https://periodicos.ufsm.br/cienciaenatura/article/view/46565/html>, last access 16.06.2021. (in Portuguese).
- [Hojjati, E., Mahtabi, G., Taran, F., Kisi, O.: Estimating evaporation from reservoirs using energy budget and empirical methods: Alavian dam reservoir, NW Iran. \*Italian Journal of Agrometeorology\* \(2\): 19–34. doi: 10.13128/ijam-1033, 2020.](#)
- 1060 [How, P., Messerli, A., Mätzler, E., Santoro, M., Wiesmann A., Caduff R., Lagley K., Bojesen M/, Paul F., Käb, A., Carrivick J.: Greenland-wide inventory of ice marginal lakes using a multi-method approach. \*Sci Rep\* 11, 4481 <https://doi.org/10.1038/s41598-021-83509-1>, 2021.](#)
- Kaup, E.: Development of anthropogenic eutrophication in Antarctic lakes of the Schirmacher Oasis, *Verhandlungen Internationale Vereinigung der Limnologie*, 29 (2), 678–682, 2005.
- Kaup, E., Haendel, D.: Snow and ice cover of water bodies. In: P. Bormann and D. Fritsche (Eds): *The Schirmacher Oasis*,

- 1065 Queen Maud Land, East Antarctica, Gotha, Justus Perthes Verlag, 279–285, 1995.
- Khare, N., Chaturvedi, S. K., Saraswat, R., Srivastava, R., Raina, R., Wanganeo, A.: Some morphometric characteristics of Priyadarshini water body at Schirmacher Oasis, Central Dronning Maud Land, Antarctica with special reference to its bathymetry. *Indian Journal of Marine Sciences*, 37(4), 435–438, 2008.
- 1070 3. [Kingslake, J., Ely, J., Das, I., Bell R.E.: Widespread movement of meltwater onto and across Antarctic ice shelves. Nature 544, 349–352, <https://doi.org/10.1038/nature22049>, 2017.](#)
- Klokov, V.D.: Surface melting and liquid water runoff from the Antarctic ice-sheet. Leningrad, Gidrometeoizdat, 126 pp., 1979. [\(in Russian\)\[in Russian\]](#)
- [Konstantinov, A. R.: Evaporation Under Natural Conditions, Israel Program for Scientific Translation, Jerusalem, 523 p., 1968.](#)
- 1075 Kourzeneva, E.: External data for lake parameterization in numerical Weather Prediction and climate modeling. *Boreal Env. Res.* 15, 165–177, 2010.
4. [Kourzeneva, E., Asensio, H., Martin, E., Faroux, S.: Global gridded dataset of lake coverage and lake depth for use in numerical weather prediction and climate modelling, Tellus A., 64, 15640, <https://doi.org/10.3402/tellusa.v64i0.15640>, 2012.](#)
- 1080 5. [Keijman, J. Q.: The estimation of the energy balance of a lake from simple weather data. Boundary-Layer Meteorol., 7, 399–407, doi: 10.1007/BF00240841, 1974.](#)
- Kljun, N., Calanca, P., Rotach, M.W., et al.: A Simple Parameterisation for Flux Footprint Predictions. *Boundary-Layer Meteorology*, 112, 503–523, <https://doi.org/10.1023/B:BOUN.0000030653.71031.96>, 2004.
- Konovalov, G.V.: Geomorphological description of the Schirmacher oasis and surrounds. *Bulletin of the Soviet Antarctic Expedition*, 37, 8 – 13, 1962. (In Russian).
- 1085 Krass M.S. : Thermophysics of lakes in Antarctic oases. Reports of the Russian Academy of Science to Antarctic committee, Vol.25, 99–125, 1986. (In Russian).
- Launiainen, J., Vihma, T.: Derivation of turbulent surface fluxes — An iterative flux-profile method allowing arbitrary observing heights, *Environmental Software*, 5(3), 113–114, [https://doi.org/10.1016/0266-9838\(90\)90021-W](https://doi.org/10.1016/0266-9838(90)90021-W), 1990.
- 1090 Leeson, A.A., Shepherd, A., Briggs, K., Howat, I., Fettweis, X., Morlighem, M., Rignot, E.: Supraglacial lakes on the Greenland ice sheet advance inland under warming climate. *Nature Climate Change*, 5(1), 51–55, <https://escholarship.org/uc/item/94m0x351>, 2015.
- Lehnerr, I., St. Louis, V.L., Sharp, M. et al.: The world’s largest High Arctic lake responds rapidly to climate warming. *Natural Communication*, 9, 1290. <https://doi.org/10.1038/s41467-018-03685-z>, 2018.
- 1095 Levy, J.S., Fountain, A.G., Obryk, M.K., et al.: Decadal topographic change in the McMurdo Dry Valleys of Antarctica: Thermokarst subsidence, glacier thinning, and transfer of water storage from the cryosphere to the hydrosphere. *Geomorphology*, 323: 80–97, <https://doi.org/10.1016/j.geomorph.2018.09.012>, 2018.
- Loopman, A., Kaup, E., Klokov, V., Simonov, I., and Haendel D.: The bathymetry of some lakes of the Antarctic oases Schirmacher and Untersee, in *Limnological Studies in Queen Maud Land (East Antarctic)*, Ed. by J. Martin (Valgus,

- 1100 Tallinn), 6–14, 1988.
- [Lu, P., Leppäranta, M., Cheng, B., Li, Z., Istomina, L., and Heygster, G.: The color of melt ponds on Arctic sea ice. \*The Cryosphere\*, 12, 1331–1345, <https://doi.org/10.5194/tc-12-1331-2018>, 2018.](#)
- 1105 [6. Majidi, M.; Alizadeh, A.; Farid, A. and Vazifedoust, M.: Estimating Evaporation from Lakes and Reservoirs under Limited Data Condition in a Semi-Arid Region \*Water Resour. Manage.\*, 29, 3711–3733, doi: 10.1007/s11269-015-1025-8, 2015.](#)
- [7. Map of the Schirmacher oasis: scale 1:25000. Ministry of Merchant Fleet of the U.S.S.R. \(Russia\), 3 sheets, 1972.](#)
- Meredith, M., M. Sommerkorn, S. Cassotta, C. Derksen, A. Ekaykin, A. Hollowed, G. Kofinas, A. Mackintosh, J. Melbourne-Thomas, M.M.C. Muelbert, G. Ottersen, H. Pritchard, and E.A.G. Schuur, 2019: Polar Regions. In: IPCC Special Report on the Ocean and Cryosphere in a Changing Climate [H.-O. Pörtner, D.C. Roberts, V. Masson-Delmotte, P. Zhai, M.
- 1110 Tignor, E. Poloczanska, K. Mintenbeck, A. Alegría, M. Nicolai, A. Okem, J. Petzold, B. Rama, N.M. Weyer (eds.)]. In press.
- Mironov, D., Golosov, S., Heise, E., Kourzeneva, E., Ritter, B., Scheider, N. and Terzhevik, A. 2005. FLake – a lake model for environmental applications. In: Proceedings of the 9th Workshop on Physical Processes in Natural Waters (eds. A. Folkard and I. Jones). Lancaster University, Lancaster, 73 pp.
- 1115 [Monin, A.S. and Obukhov, A.M.: Basic laws of turbulent mixing in the surface layer of the atmosphere. \*Contrib. Geophys. Inst. Acad. Sci. USSR\*, 151\(163\), p.e187, 1954.](#)
- [Monteith, J.: Evaporation and the environment. In: 19th Symposium of the Society of Experimental Biology. Cambridge, UK Cambridge University Press, 205-234, 1965.](#)
- [Mustonen S. \(Ed.\), \*Sovellettu hydrologia. Vesiyhdistys, Helsinki, pp. 291-323, 1986. \(in Finnish\).\*](#)
- 1120 Naakka, T., Nygård, T., Vihma T.: Air moisture climatology and related physical processes in the Antarctic on the basis of ERA-5 reanalysis, 34(11), 4463-4480, <https://doi.org/10.1175/JCLI-D-20-0798.1>, 2021.
- [Obukhov, A.M.: Turbulence in an Atmosphere with a Non-Uniform Temperature. \*Boundary-Layer Meteorology\*, 2, 7-29, <https://doi.org/10.1007/BF00718085>, 1946.](#)
- 1125 [8. Nash, J.E., Sutcliffe, J.V.: River flow forecasting through conceptual models: Part I—A discussion of principles. \*J. Hydrol.\* 10, 282–290. doi: 10.1016/0022-1694\(70\)90255-6, 1970.](#)
- Odrova, T.: Hydrophysic of water reservoirs. Leningrad, Gidrometizdat, 312 p., 1979. (In Russian)
- Penman, H.L.: Natural evaporation from open water, bare soil and grass. *Proc. Roy. Soc. London A*, 194, 120–145, <http://doi.org/10.1098/rspa.1948.0037>, 1948.
- Phartiyal, B., Sharma, A., Bera, S.K.: Glacial lakes and geomorphological evolution of Schirmacher Oasis, East Antarctica during Quaternary. *Quaternary International* 23, 128–136, doi:10.1016/j.quaint.2010.11.025, 2011.
- 1130 Popov, E. G.: Hydrological forecasts, Leningrad, Gidrometeoizdat, 257 p., 1979. (in Russian).
- Potes, M., Salgado, R., Costa, M.J., Morais, M., Bortoli, D., Kostadinov, I., Mammarella, I.: Lake–atmosphere interactions at Alqueva reservoir: a case study in the summer of 2014, *Tellus A: Dynamic Meteorology and Oceanography*, 69:1, doi: 10.1080/16000870.2016.1272787, 2017.

- 1135 Picard, G., Fily, M., Gallee, H.: Surface melting derived from microwave radiometers: A climatic indicator in Antarctica. *Annals of Glaciology*, 46, 29–34. doi:10.3189/172756407782871684, 2007.
- [Ramesh, K.J., Soni, V.K.: Perspectives of Antarctic weather monitoring and research efforts. Pol. Sci. 18, 183–188, https://doi.org/10.1016/j.polar.2018.04.005, 2018.](https://doi.org/10.1016/j.polar.2018.04.005)
- 1140 Richter, W., Borman, P.: Geomorphology of the Schirmacher oasis, In: P. Bormann and D. Fritsche (Eds). *The Schirmacher Oasis, Queen Maud Land, East Antarctica*, Gotha: Justus Perthes Verlag, 171–206, 1995.
- Rodrigues, C. M., Moreira, M., Guimarães, R. C., Potes, M.: Reservoir evaporation in a Mediterranean climate: comparing direct methods in Alqueva Reservoir, Portugal, *Hydrol. Earth Syst. Sci.*, 24, 5973–5984, <https://doi.org/10.5194/hess-24-5973-2020>, 2020.
- 1145 [Salgado, R., Le Moigne, P.: Coupling of the FLake model to the Surfex externalized surface model. Boreal Environment Research, 15:2, 231-244, 2010.](https://doi.org/10.1007/s10077-010-0231-2)
- [Sahlée, E., Rutgersson, A., Podgrajsek, E., Bergström, H. : Influence from surrounding land on the turbulence measurements above a Lake. Boundary-Layer Meteorol., 150:235–258, DOI 10.1007/s10077-014-0235-2, 2014.](https://doi.org/10.1007/s10077-014-0235-2)
- Salesky, S., Chamecki, M., and Dias, N.: Estimating the random error in eddy covariance based fluxes and other turbulence statistics: the filtering method, *Bound. Lay. Meteorol.*, 144, 113–135, 2012.
- 1150 SCAR Antarctic Digital Database (ADD): available at <https://data.bas.ac.uk/collections/e74543c0-4c4e-4b41-aa33-5bb2f67df389/>, last access 18.10.2021.
9. [Sene, K. J., Gash, J. H., McNeil, D. D.: Evaporation from a tropical lake: comparison of theory with direct measurements. Journal of Hydrology, 127, 193–217, 1991.](https://doi.org/10.1007/BF02671991)
- Simonov I.M.: Oases of East Antarctica. *Gidrometeoizdat*. 1971. 176 pp., 1971. (in Russian).
- 1155 Simonov, I.M., Fedotov, V.I.: Oзера oasisa Schimachera. [Lakes of the Schirmacher oasis]. *Informazioni bulletin Sovetskoy Antarticheskoy Expedicii*, 47, 19–23, 1964. (In Russian).
- Sinha R., Chatterjee A.: Thermal structure, sedimentology, and hydro-geochemistry of Lake Priyadarshini, Schirmacher oasis, Antarctica. Sixteenth Indian Expedition to Antarctica, Scientific Report, Department of Ocean Development, Technical Publication No. 14, 36 pp., 2000.
- 1160 Sokratova, I. N.: Hydrological investigations in the Antarctic oases, *Russ. Meteorol. Hydrol.*, 36, 207, doi:10.3103/S1068373911030083, 2011. (in Russian).
- Shevnina, E., Kourzeneva, E.: Thermal regime and components of water balance of lakes in Antarctica at the Fildes peninsula and the Larsemann Hills. *Tellus A: Dynamic Meteorology and Oceanography*, 69(1), 1317202, doi:10.1080/16000870.2017.1317202, 2017.
- 1165 Shevnina, E.: 3D wind speed and CO<sub>2</sub>/H<sub>2</sub>O concentration measurements collected during austral summer 2017/2018 over an ice free surface of a shallow lake located in the Schirmacher oasis, East Antarctica. [Data set]. Zenodo. <http://doi.org/10.5281/zenodo.3469570>, 2019.

- Shevnina, E., Kourzeneva, E., Dvornikov, Y., and Fedorova, I.: Retention time of lakes in the Larsemann Hills oasis, East Antarctica, *The Cryosphere*, 15, 2667–2682, <https://doi.org/10.5194/tc-15-2667-2021>, 2021.
- 1170 Shuttleworth, W. J.: Evaporation. In: Maidment, D.R., (Eds): *Handbook of Hydrology*, New York: McGraw-Hill, 4.1-4.53, 1993.
- Srivastava, A. K., Ingle, P. S., Lunge, H. S., Khare, N.: Grain-size characteristics of deposits derived from different glacial environments of the Schirmacher Oasis, East Antarctica. *Geologos* 18, 4, 251–266, doi: 10.2478/v10118-012-0014-0, 2012.
- 1175 Stannard, D. I., and Rosenberry, D. O.: A comparison of short-term measurements of lake evaporation using eddy correlation and energy budget methods. *Journal of Hydrology*, 122 (1-4), 15–22. [https://doi.org/10.1016/0022-1694\(91\)90168-H](https://doi.org/10.1016/0022-1694(91)90168-H), 1991.
- Stokes, C. R., Sanderson, J. E., Miles, B. W. J. et al.: Widespread distribution of supraglacial lakes around the margin of the East Antarctic Ice Sheet, *Scientific Reports*, 9, 13823, doi: 10.1038/s41598-019-50343-5, 2019.
- Stull, R.: *Practical Meteorology: An Algebra-based Survey of Atmospheric Science -version 1.02b*. Univ. of British  
 1180 Columbia. 940 p. 2017.
- Tanny, J., Cohen, S., Assouline, S., Lange, F., Grava, A., Berger, D., Teltch, B., Parlange, M.B.: Evaporation from a small water reservoir: Direct measurements and estimates. *Journal of Hydrology*, 351, 218–229, 2008.
- 1185 10. Thiery, W., Gorodetskaya, I. V., Bintanja, R., Van den Broeke, M. R., Reijmer, C., Kuipers, P., Munneke, N., Van Lipzig, P. M.: Surface and snowdrift sublimation at Princess Elisabeth station, East Antarctica. *Cryosphere*, 6 (2), 841–857, 2012.
- Tomasi, C., Cacciari, A., Vitale, V., Lupi, A., Lanconelli, C., Pellegrini, A., Grigioni, P.: Mean vertical profiles of temperature and absolute humidity from a 12-year radiosounding data set at Terra Nova Bay (Antarctica). *Atmospheric Research*, 71, 3, p. 139-169, doi: 10.1016/j.atmosres.2004.03.009, 2004.
- Toptunova, O., Choulga, M., Kurzeneva, E.: Status and progress in global lake database developments, *Adv. Sci. Res.*, 16,  
 1190 57–61, <https://doi.org/10.5194/asr-16-57-2019>, 2019.
- Turner, J., Pendlebury, S. F.: *The International Antarctic Weather Forecasting Handbook*. British Antarctic Survey, 663 p., 2004.
- 1195 11. Vihma, T., E. Tuovinen, and H. Savijärvi.: Interaction of katabatic winds and near-surface temperatures in the Antarctic, *J. Geophys. Res.*, 116, D21119, doi:10.1029/2010JD014917, 2011.
12. Viterbo, P.: A review of parametrization schemes for land surface processes. ECMWF, Reading, England, 2002. Available online: <https://www.ecmwf.int/sites/default/files/elibrary/2002/16960-review-parametrization-schemes-land-surface-processes.pdf>, last access: 25.05.2021.
- Verleyen, E., Hodgson, D.A., Vyverman, W., Roberts, D., McMinn, A., Vanhoutte K., Saabe K.: Modelling diatom responses to climate induced fluctuations in the moisture balance in continental Antarctic lakes. *Journal of Paleolimnology*,  
 1200 30, 195–215, <https://doi.org/10.1023/A:1025570904093>, 2003.
- Verleyen, E., Hodgson, D., Gibson, J., Imura, S., Kaup, E., Kudoh, S., Wever, D. A., Hoshino, T., McMinn, A., Obbels, D., Roberts, D., Roberts, S., Saabe, K., Souffreau C., Tavernier I., van Niewenhuyze, W., van Ranst E., Vindevogel, N.,



- Vyverman, W.: Chemical limnology in coastal East Antarctic lakes: Monitoring future climate change in centers of endemism and biodiversity. *Antarctic Science*, 24(1), 23–33. doi:10.1017/S0954102011000642, 2012.
- 1205 Venäläinen, A., Heikinheimo, M. Tourula, T.: Latent heat flux from small sheltered lakes. *Bound-Lay Meteorol.* 86, 355–377, doi: 10.1023/A:1000664615657, 1998.
- Vickers, D., Mahrt, L.: Quality control and flux sampling problems for tower and aircraft data. *Journal of Atmospheric and Oceanic Technology*, 14(3), 512–526, [https://doi.org/10.1175/1520-0426\(1997\)014%3C0512:QCAFSP%3E2.0.CO;2](https://doi.org/10.1175/1520-0426(1997)014%3C0512:QCAFSP%3E2.0.CO;2), 1997.
- 1210 [Vignon, É., Roussel, M.-L., Gorodetskaya, I. V., Genthon, C., Berne, A.: Present and future of rainfall in Antarctica. \*Geophysical Research Letters\*, 48, e2020GL092281, <https://doi.org/10.1029/2020GL092281>, 2021.](#)
- [Vickers, D., Mahrt, L.: Quality control and flux sampling problems for tower and aircraft data. \*Journal of Atmospheric and Oceanic Technology\*, 14\(3\), 512–526, \[https://doi.org/10.1175/1520-0426\\(1997\\)014%3C0512:QCAFSP%3E2.0.CO;2\]\(https://doi.org/10.1175/1520-0426\(1997\)014%3C0512:QCAFSP%3E2.0.CO;2\), 1997.](#)
- [Vihma, T., J. Uotila, B. Cheng, and J. Launiainen: Surface heat budget over the Weddell Sea: buoy results and comparisons with large-scale models. \*J. Geophys. Res.\*, 107 \(C2\), 3013, doi: 10.1029/2000JC00037, 2002.](#)
- 1215 [Vihma, T., E. Tuovinen, and H. Savijärvi.: Interaction of katabatic winds and near-surface temperatures in the Antarctic. \*J. Geophys. Res.\*, 116, D21119, doi:10.1029/2010JD014917, 2011.](#)
- [Viterbo, P.: A review of parametrization schemes for land surface processes. ECMWF, Reading, England, 2002. Available online: <https://www.ecmwf.int/sites/default/files/elibrary/2002/16960-review-parametrization-schemes-land-surface-processes.pdf>, last access: 25.05.2021.](#)
- 1220 [Webb, E. K.: On Estimating Evaporation with Fluctuating Bowen Ratio, \*J. Geophys. Res.\* 65, 3415–3417, 1960.](#)
- Williamson, C.E., Saros, J.E., Vincent, W.F. Smol , J.P.: Lakes and reservoirs as sentinels, integrators, and regulators of climate change. *Limnology and Oceanography*, 54, 2273–2282, 2009.
- Xu, M., L. Yu, K. Liang, T. Vihma, D. Bozkurt, X. Hu, Q. Yang: Dominant role of vertical air flows in the unprecedented warming on the Antarctic Peninsula in February 2020. *Communications Earth and Environment*, 2, 133, <https://doi.org/10.1038/s43247-021-00203-w>, 2021.
- 1225 [Zhao, L., Xia, J., Xu, C. et al.: Evapotranspiration estimation methods in hydrological models. \*J. Geogr. Sci.\* 23, 359–369, <https://doi.org/10.1007/s11442-013-1015-9>, 2013.](#)