

Resolving GIA in response to modern and future ice loss at marine grounding lines in West Antarctica

5 Jeannette Xiu Wen Wan¹, Natalya Gomez¹, Konstantin Latychev², Holly Han¹

¹McGill University, Department of Earth and Planetary Sciences, Montreal, Canada

²Harvard University, Department of Earth and Planetary Sciences, Cambridge, Massachusetts

Correspondence to: Natalya Gomez (natalya.gomez@mcgill.ca)

10 **Abstract.** Accurate glacial isostatic adjustment (GIA) modeling in the cryosphere is required for interpreting satellite, geophysical and geological records and to assess the feedbacks of Earth deformation and sea level change on marine ice-sheet grounding lines. Assessing GIA in areas of active ice loss in West Antarctica is particularly challenging because the ice is underlain by laterally varying mantle viscosities that are up to several orders of magnitude lower than the global average, leading to a faster and more localized response of the solid Earth to ongoing and future ice sheet retreat and necessitating GIA
15 models that incorporate 3-D viscoelastic Earth structure. Improvements to GIA models allow for computation of the viscoelastic response of the Earth to surface ice loading at sub-kilometre resolution and ice-sheet models and observational products now provide the inputs to GIA models at comparably unprecedented detail. However, the resolution required to capture GIA in models remains poorly understood, and high-resolution calculations come at heavy computational expense. We adopt a 3-D GIA model with a range of Earth structure models based on recent seismic tomography and geodetic data to
20 perform a comprehensive analysis of the influence of grid resolution on predictions of GIA in the Amundsen Sea Embayment (ASE) in West Antarctica. Through idealized sensitivity testing down to sub-kilometer resolution with spatially isolated ice loading changes, we find that a grid resolution of ~ 3 times the radius of the load is required to accurately capture the elastic response of the Earth. However, when we consider more realistic, spatially coherent ice loss scenarios based on modern observational records and future ice sheet model projections and adopt a viscoelastic Earth, we find that errors of less than 5%
25 along the grounding line can be achieved with a 7.5 km grid, and less than 2% with a 3.75 km grid, even when the input ice model is on a 1 km grid. Furthermore, we show that low mantle viscosities beneath the ASE lead to viscous deformation that contributes to the instrumental record on decadal timescales and equals or dominates over elastic effects by the end of the 21st century. Our findings suggest that for the range of resolutions of 1.9-15 km that we considered, the error due to adopting a coarser grid in this region is negligible compared to the effect of neglecting viscous effects and the uncertainty in the adopted
30 mantle viscosity structure.

1 Introduction

Changes in sea level in response to ice-mass loss are spatially variable because of glacial isostatic adjustment (GIA), which is the deformational, gravitational, and rotational response of the viscoelastic solid Earth to changes in surface ice and water distribution. The response of the bedrock begins with an instantaneous elastic response of the solid planet's lithosphere and mantle, and transitions to a longer timescale viscous relaxation of the mantle towards isostatic equilibrium. GIA models produce predictions of changes in the height of the sea surface equipotential and solid Earth surface (i.e. sea-level changes) in response to surface ice loading changes, which are in turn used to interpret satellite, geophysical and geological records and serve as input to models of ice-sheet dynamics.

40

Accurate GIA modelling is required to constrain the sea level and solid earth feedbacks on ice dynamics in the coming centuries, especially along unstable marine-grounded ice fronts in Antarctica where bedrock uplift and gravitational drawdown of the sea surface due to ice loss act as a negative feedback to stabilise the retreat of marine-grounded ice-sheet grounding lines (e.g. Gomez et al., 2010; 2013; 2015; De Boer et al., 2014; Konrad et al., 2015; Larour et al., 2019). Furthermore, the GIA response to past and modern ice cover changes makes a significant contribution to satellite records of modern mass changes in marine sectors of the West Antarctica that are actively experiencing ice loss (e.g. King et al., 2012; the IMBIE team, 2018).

GIA modelling in Antarctica is complicated by the fact that the continent is characterised by strong lateral variability in lithospheric thickness and upper mantle viscosity, with low viscosities in the west and high viscosities in the east (e.g. An et al., 2015a; Heeszel et al., 2016; Lloyd et al., 2020). In particular, the low-viscosity mantle and thin lithosphere observed under the West Antarctic Ice Sheet (WAIS) identified from increasingly resolved seismic tomography and geodetic and geologic constraints (Ritzwoller et al., 2001; Morelli and Danesi, 2004; Kaufmann et al., 2005; Nield et al., 2014; Heeszel et al., 2016; Barletta et al., 2018; Shen et al., 2018; Lloyd et al., 2020) leads to a more spatially localised (short wavelength) and faster viscoelastic response to surface loading than cratonic regions covered by Late Pleistocene ice sheets (e.g. Hay et al., 2017; Powell et al., 2020). Over West Antarctica, upper mantle viscosities are thought to vary by several orders of magnitude over short spatial scales (~100s of km) reaching as low as 10^{18} Pa s in the Amundsen Sea Embayment (ASE) beneath areas of active marine ice loss (e.g. Nield et al., 2014; Barletta et al., 2018). This implies that viscous effects due to 20th century and more recent ice loss will become significant on annual to decadal timescales and accelerate during the timeframe of instrumental records (Barletta et al., 2018; Powell et al., 2020). Viscous deformation due to ongoing ice loss also has the potential to influence ice sheet grounding lines in the coming centuries (Gomez et al., 2015; Kachuck et al., 2020; Coulon et al., 2021) but has not been included in recent high resolution coupled ice sheet – sea level model projections (Larour et al., 2019).

65 To accurately capture the timing and wavelength of GIA effects across Antarctica, models must be capable of both accounting
for 3-D earth structure (i.e., 3-D GIA models such as Latychev et al., 2005 or van der Wal et al., 2015), and be of sufficient
spatiotemporal resolution to capture the geometry of grounded ice cover. Commonly, GIA, ice-sheet and coupled ice-sheet –
GIA modelling (e.g. Gomez et al., 2015; Konrad et al. 2015) studies of the sea level change in response to modern and future
70 ice loss have been performed with only 1-D (radially varying) Earth structure models (e.g., Kendall et al., 2005; Spada and
Stocchi, 2007; Adhikhari et al., 2016), or with coarse spatial resolutions of > 20 km due to the computational expense. GIA
models capable of km- to sub-km-resolution have also been developed (e.g. the 1-D GIA model by Adhikhari et al., 2016; the
3-D GIA model by Latychev, et al. 2005 with updates described in supplementary materials of Gomez et al., 2018). For
computational efficiency, some of these models implement regional grid refinement techniques which allow for a higher
resolution along ice retreat margins. Alongside this, improvements in ice-sheet models (e.g. Nowicki et al., 2016; Seroussi et
75 al., 2020; DeConto et al., 2021) and observational products (e.g. Studinger, 2014; Bamber et al., 2020; Smith et al., 2020;
Morlinghem et al., 2020) allow similarly high-resolution km- to sub-km- ice thickness and bedrock topography datasets that
serve as input to GIA models. These advancements allow for GIA models to capture short-wavelength bedrock deformation
and input ice loading changes at unprecedented detail, but at heavy computational expense, particularly for global 3-D GIA
models.

80

It is well-established that dynamic ice-sheet models are sensitive to the chosen grid resolution (e.g. Durand et al., 2009; Van
den Berg & Van de Wal, 2006), requiring km to sub-km resolution to accurately represent ice dynamics and grounding line
migration in some applications (e.g. Gladstone et al., 2012; Pattyn et al., 2013; Cornford et al., 2016). It has also been suggested
that as fine as 1-km resolution bedrock topography may be needed to capture the influence of fine-scale topographic features
85 on the ice-sheet evolution (Durand et al., 2009) and high resolution may also be needed to represent some embayment walls
and pinning points that act to slow down retreat (e.g. Favier et al., 2012; Joughin et al., 2014; Berger et al., 2016).

While topographic features themselves can be very fine scale, the changes in bedrock elevation and sea level in response to
ice cover changes tend to be longer wavelength, and the corresponding spatial resolution required to accurately resolve these
90 changes in GIA models and their influence on ice dynamics remains poorly understood. Larour et al. (2019) suggested that
kilometer-scale resolution may be required to capture the elastic component of deformation in response to ice loss. However,
the idealized tests they performed considered an isolated, and increasingly localized load, and their conclusion may not hold
for more realistic, spatially coherent ice loss scenarios. Furthermore, their model did not include viscous deformation in
response to ongoing ice loss during the simulation, or account for lateral variations in Earth structure. There have been limited
95 studies assessing the length scale of realistic viscoelastic bedrock response beneath the structurally complex WAIS, though a
recent high-resolution bedrock deformation modelling study by Zwinger et al. (2020) suggests a convergence in modelled
deformation at resolutions finer than 5 km. The broad spatial nature of bedrock deformation and the spatially coherent nature

of ice-sheet retreat, which becomes less localised over longer timescales, suggest that sub-km to km grid resolution, which comes at great computational cost, may not be necessary for accurate GIA model calculations.

100

The aim of this study is to assess the sensitivity of predictions of GIA in response to modern and future ice loss to spatial resolution in the rheologically complex, marine sectors of the WAIS. We build a 3-D viscosity model based on the most recent Antarctic-wide seismic tomography model (Lloyd et al., 2020) to serve as input to a 3-D finite volume, global sea level model (Latychev et al., 2005) to assess the performance of 3-D GIA model predictions across surface grid resolutions of 1.9-15 km.

105

We repeat calculations with a range of Earth models, considering the contribution from elastic and viscous deformation separately. We focus on the response to observed modern ice loss over the last two decades (Shepherd et al., 2019) and projected future ice sheet retreat in the coming century (Golledge et al., 2019; DeConto et al., 2021) in the Amundsen Sea Embayment of West Antarctica. Our study is motivated by the following questions: What 3-D grid resolution is necessary to adequately capture the elastic and viscous uplift and associated gravitational and rotational effects in response ice loading changes? How significant is the effect of grid resolution compared to sources of uncertainty and simplifications made in some previous modelling, in particular the neglect of viscous deformation.

110

2 Methods

To investigate the influence of GIA model grid resolution, we first conduct idealized load sensitivity tests over a range of surface grid resolutions from 7.5 to 0.5 km, for the instantaneous removal of cylindrical loads from 0.5 to 16 km in radius (Section 3). We then widen our “aperture” to assess the model grid resolution required to accurately capture GIA due to modern ice-sheet cover changes from satellite observations (Shepherd et al., 2019) and future ice loss from ice-sheet model projections (Golledge et al., 2019; DeConto et al., 2021) in the rapidly evolving Amundsen Sea Embayment of West Antarctica (Section 4). We choose to locate our study region (light blue square in Fig. 1a) on the ASE both because of the ongoing ice loss and vulnerability to large-scale future marine ice-sheet retreat in the region, and because the region is characterized by low upper mantle viscosities and a thinned lithosphere (e.g. Barletta et al., 2018) making the ice there sensitive to solid-earth ice-sheet feedbacks. In the ASE, the Pine Island and Thwaites Glaciers together contributed 95 Gt/year of the 159 ± 8 Gt/year total WAIS mass flux in 2017 (Rignot et al., 2019), with studies estimating that collapse of Thwaites Glacier is already underway (Joughin et al., 2014). Accurate GIA predictions are critical to assess rates of future ice-sheet retreat and associated sea level changes making it an ideal location to study the effects of grid resolution on modelled GIA. To represent the radially and laterally variable Earth rheology in this region, we use a viscoelastic Earth rheology and a range of 3-D viscosity structure models in Antarctica derived from seismic tomography (An et al., 2015a; Heeszel et al., 2016; Lloyd et al., 2020). We adopt a range of 3-D Earth model grids with surface resolutions from 1.9-15 km and compare results to first assess the resolution required to capture the elastic deformation associated with ice loss. Lastly, we repeat these experiments with viscoelastic

120

125

130 effects and compare results to the elastic calculations to assess the contribution of viscous effects to modern and future sea level, and the model resolution required to capture these effects. In the sections that follow, we describe the adopted 3-D GIA model, computational grids, Earth rheological models, and adopted modern and future ice loss scenarios.

2.1 3-D GIA Model

We compute GIA predictions with a global 3-D finite volume sea level and Earth deformation model (Latychev et al., 2005) 135 capable of regional grid refinement (Gomez et al., 2018). The model solves the sea level equation (Kendall et al., 2005) over a 3-D global spherical tetrahedral grid defined from the surface to the core-mantle boundary (CMB) that allows us to resolve the laterally and radially varying Earth structure, which are a strong feature in Antarctica. We also adopt this 3-D GIA model because it is capable of regional grid refinement to achieve regional resolution at sub-km scale in regions of interest within a lower resolution globe (Section 2.2). The model computes gravitationally self-consistent solutions for the sea level equation, 140 incorporating effects of time-varying shorelines, Earth rotation changes and viscoelastic deformation of the Earth assuming an elastically compressible Maxwell viscoelastic rheology. The GIA model requires two main inputs, a 3-D Earth model of viscoelastic rheological properties and a time series of ice thickness changes. These components are described in the following sections. The model also requires global topography as an initial boundary condition, including the elevation of the bedrock beneath the ice. Topography globally outside of Antarctica is set to etopo2 (NOAA National Geophysical Data Centre, 2006) 145 in all experiments with a realistic loading scenario, and the Antarctic bedrock elevation for each of the experiments is described below. Note that we adopt a standalone GIA model throughout this study with the purpose to inform the setup of coupled ice sheet - sea level modeling studies, but we do not model the feedback of GIA on ice sheet dynamics explicitly. However, other work (e.g. Kachuck et al., 2020; Gomez et al., 2015) suggests that the scale of differences in GIA simulated at different resolutions and adopting different Earth structure models here will be large enough to alter the timing and magnitude of 150 grounding line migration in a coupled modelling context.

2.2 Computational model grid: regional grid refinement

To compute GIA model predictions, we construct model grids of various surface resolutions (Fig. 1) with the regional grid refinement process detailed in the supplementary material of Gomez et al. (2018). Grid refinement is achieved by incrementally bisecting grid edges in the selected region to achieve a desired resolution, and a final smoothing operation along the region 155 boundary to ensure a well-behaved transition. We perform calculations on a base grid with a global surface resolution of 15-km, which consists of 20 million nodes and 70 radial layers between the core-mantle boundary and the Earth's surface. The radial layers of the grid are defined to respect the unconformities in material properties of the radially varying (1-D) seismic reference model STW105 (Kustowski et al., 2008), with the shallowest layers at 12, 25 and 43 km depth. Eight regionally refined grids are constructed from this base grid: five for the idealized load sensitivity tests at 7.5, 3.75, 1.75, 1 to 0.5 km 160 surface grid resolution over a minimum 0.3-degree radius region around the test load, and three for the more realistic calculations from observed modern and future model projected ice loss, at 7.5, 3.75 and 1.9 km surface resolution in

incrementally smaller regions converging over the ASE (Fig. 1). The highest-resolution 1.9 km grid over the ASE has ~ 29 million nodes, which takes ~ 65 CPU hours to run per time step on a high performance computing cluster. As our study focusses on surface grid resolution, the grid refinement is limited to the surface and a few layers down to ~ 10 km. Testing with deeper
165 grid refinement during experiment design process indicated that this was sufficient, and our results indicate that km-scale model resolution is only needed at the surface to accurately capture the geometry of surface loading.

2.3 Earth rheological model

The spatial pattern and amplitude of surface deformation in response to ice loading is dependent on the underlying Earth
170 structure. For the idealized sensitivity tests in section 3, we adopt a purely elastic Earth model with a 1-D elastic and density structure. In section 4, we move to a set of realistic simulations using observed or modelled AIS ice loading, adopting 3-D viscoelastic Earth models with a range of viscosity structures constrained by seismic tomography (An et al., 2015a,b; Lloyd et al., 2020) and informed by GNSS-inferences of local mantle structure. The elastic and density structure for these models are based on seismic reference model STW105 (Kustowski et al., 2008). Laterally varying lithospheric thickness (Fig. S1(d)) in
175 all simulations is a composite of a regional lithospheric thickness model by An et al. (2015a) over Antarctica, and a global lithospheric thickness model by Conrad and Lithgow-Bertelloni (2006) everywhere else. Over Antarctica, lithospheric thickness is scaled to have an average of 96 km, resulting in a minimum of 40 km, as was done in Hay et al. (2017).

3-D mantle viscosity variability is estimated from seismic velocity models by scaling isotropic seismic shear wave velocity
180 anomalies to a viscosity variability term using the method developed by Ivins and Sammis (1995). This term defines the variability of mantle viscosity with reference to a chosen 1-D viscosity profile. We follow the same procedure as described in Latychev et al. (2005), Austermann et al. (2013) and Gomez et al. (2018), whereby lateral variations in mantle viscosity are established by sequentially converting the field of relative variations in isotropic shear wave velocity to density, temperature and eventually viscosity anomalies. The final conversion adopts a factor that scales the dependence of viscosity to temperature
185 variations, determining the peak-to-peak lateral variation in viscosity. A different scaling factor is applied to regional and global seismic velocity models to account for the amplitude differences between the different seismic velocity models.

To address the substantial uncertainty in Earth structure, we repeat our simulations with three different 3-D viscosity models: EM1_L (Figs. 1c, d, S1), EM1_M, and EM2. EM1_M and EM1_L adopt the latest high-resolution 25 km Antarctic seismic
190 tomography model by Lloyd et al. (2020) (ANT-20) in the region south of 45°S and extending from the surface down to the transition zone, and S362ANI (Kustowski et al., 2008) for the rest of the globe. ANT-20 is provided on a 25 km grid at the surface, with distance between the points decreasing with depth, and provided at depth slices in 5 km intervals from 0 to 800 km depth. S362ANI is a 3-D global anisotropic shear wave velocity model for the whole mantle, extending from 25 km depth to the core-mantle-boundary defined at 2891 km depth, provided at a 2° lateral grid resolution at the surface. We note, however,

195 that the spacing of the seismic model grid is distinct from the scale of the Earth structure variations captured by the model. While quantitatively assessing the latter remains an outstanding challenge in seismic tomography, we expect the resolution of the ANT-20 model to be $\mathcal{O}(>100 \text{ km})$ in the upper mantle and coarser at greater depths, and thus well represented by the GIA model grid. The two variations EM1_M and EM1_L were scaled to represent a moderate range of viscosities across Antarctica that match regional averages, and a lower-viscosity endmember adjusted to match GPS-derived inferences of minimum
200 viscosity beneath WAIS (Nield et al., 2014; Zhao et al., 2017; Barletta et al., 2018).

The viscosity variations in EM1_M are more moderate with scaling factors of 0.0263 for ANT-20 and 0.035 for S362ANI, both close to the preferred value in Kaufmann et al. (2005). These viscosity variations are superimposed on a 1-D viscosity profile of $5 \times 10^{20} \text{ Pa s}$ in the upper mantle and $5 \times 10^{21} \text{ Pa s}$ in the lower mantle typical in most GIA-based inferences of mantle
205 viscosity (e.g., Mitrovia and Forte, 2004). As noted, the viscosity variations in EM1_L (Figs. 1c, d, S1) are of higher amplitude with lower viscosities under WAIS and the ASE. EM1_L adopts larger scaling factors of 0.033 for ANT-20 and 0.04 for S362ANI and an accompanying 1-D viscosity profile of $1 \times 10^{20} \text{ Pa s}$ in the upper mantle, which is aligned with Lambeck et al., (2014) estimates of 1-D upper mantle viscosity using far-field sea-level proxy records, and $5 \times 10^{21} \text{ Pa s}$ in the lower mantle. The larger scaling factors applied in EM1_L were calibrated to best reflect the absolute upper mantle viscosity estimates from
210 dynamically derived GPS bedrock uplift rates in the WAIS at three locations: $\sim 6 \times 10^{17}$ to $2 \times 10^{18} \text{ Pa s}$ at the northern Antarctic Peninsula (Nield et al., 2014), $\sim 2 \times 10^{19}$ to $2 \times 10^{20} \text{ Pa s}$ at the Fleming Glacier in central Antarctic Peninsula (Zhao et al., 2017), and $\sim 2.5 \times 10^{18}$ to $4 \times 10^{19} \text{ Pa s}$ at the Amundsen Sea Embayment (Barletta et al., 2018). Figures 1c, d and S1 shows the resulting low viscosity earth model structure (EM1_L), which has the lowest viscosity at Marie Byrd land of $\sim 9 \times 10^{17}$ to $7 \times 10^{18} \text{ Pa s}$ in the upper mantle.

215

Lastly, the EM2 model, also adopted in Hay et al. (2017), Gomez et al. (2018) and Powell et al. (2020), is a combination of three seismic models: S40RTS (Ritsema et al., 2011) globally, a model by An et al. (2015a) in the East Antarctica and Antarctic Peninsula, and the model by Heeszel et al (2016) for West and Central Antarctica. The full construction of EM2 is detailed in Hay et al. (2017).

220 **2.4 Ice model and topography**

We consider three ice melt scenarios with resolution ranging from 1- to 5 km in the ASE. The total ice thickness change from start to end of each scenario is shown in Figs. 2a-c.

For observations of modern ice loss, we adopted a reconstruction we term ICE-SH from Shepherd et al. (2019) of surface
225 elevation change (Δh) from 25 years (1992-2017) of multi-mission satellite altimetry data resolved over a 5-km grid at 5-year intervals. We treat Δh as a proxy for ice thickness change (Carrivick et al., 2019), apply the Bedmap2 (Fretwell et al., 2013) grounded ice mask and saturate ice thickness change $> 20 \text{ m/yr}$ to control against spurious data points. Initial ice thickness is

given by Bedmap2. Observations of ice surface elevation changes in Antarctica are continuously improving in resolution, and currently range from metre-scale resolution over short observational tracks (e.g. Studinger et al., 2014), to sub-km to km-scale resolution at the regional scale (e.g. Bamber et al., 2020), to ~ 5 to 35 km from radar and laser satellite altimetry derived records over the whole Antarctic (e.g. Martin Español et al., 2016; Schröder et al., 2019; Shepherd et al., 2019; Smith et al., 2020). ICE-SH was selected from the available observational datasets due to the 5-km resolution being the highest of its class of available satellite altimetry derived-records providing decadal time span surface elevation change records covering the whole Antarctic.

235

For Antarctic evolution over the next century, we apply modelled ice thickness changes from two Antarctic-wide ice-sheet model projections: (1) ICE-GOL which predicts AIS evolution under RCP 8.5 and including meltwater feedbacks from 2000 to 2100 at 5-km resolution over 5-year intervals (Golledge et al., 2019), and (2) ICE-RD which predicts AIS evolution from 1950 to 2100 at 10-km resolution over the whole AIS with a nested 1-km resolution simulation over ASE at annual resolution (Extended Data Fig. 5 from DeConto et al., 2021). For ICE-RD, we ran simulations at yearly intervals from 1950-2100 but the interval from 1950-2000 is a period of ice model initialization and we therefore focus on the period between 2000 and 2100 in our results. We also take initial Antarctic bedrock topography from the ice models. Further information of each model can be found in the corresponding references. In selecting these scenarios, the goal is to provide a representative sample of spatially coherent ice-sheet retreat scenarios at high resolution from the literature, rather than to capture all possible projected future ice loss scenarios.

245

3 Idealized experiments: sensitivity of elastic uplift predictions to grid resolution

Our main goal in this analysis is to assess the relationship between grid resolution and elastic GIA predictions, and to identify, for a given load dimension, the grid resolution required to accurately model the associated GIA response. Realistic ice retreat has complex geometry, making it difficult to pinpoint the cause of GIA inaccuracies due to resolution, which may be due to poor representation of the ice load, or numerical errors in representation of the response to Earth loading. To isolate the effect of changing grid resolution on GIA predictions, we first perform a suite of idealized sensitivity tests modelling the instantaneous elastic deformation from unloading of an isolated cylindrical ice load with differing surface grid resolutions that are iteratively bisected from 7.5 km down to 0.5 km. We chose to perform the test with short wavelength, spatially isolated ice loading changes because these would be most poorly represented by a coarse grid compared to coherent ice loss over a broader area. Furthermore, these tests with idealized loads are less computationally costly and enable a systematic assessment reaching higher spatial resolution. In total, 85 GIA simulations were run using 17 ice cylinders of height 100 m and radii ranging from 0.5 to 16 km (0.5, 1, 2, ..., 15, 16), across five different grid surface resolutions: 0.5, 1, 1.75, 3.75 and 7.5 km.

255

The simulations are performed with the purely elastic 1-D earth model (Sect. 2.3), and an idealized topography of 3800 m south of 24.5 °S and - 835 m everywhere else to reflect the 30:70 land to sea ratio on earth (Fig. S2). A radially symmetric ice sheet with steady-state Antarctic ice dome profile (Paterson and Colbeck, 1980) sits on top of this topography extending from the south pole to 69°S, with a maximum height of 3500 m. We also consistently place the centre of the cylindrical load on an arbitrary model grid node in the ASE (76°S 150°W).

3.1 Idealized experiment results

Figure 3 summarises the error in predicted elastic GIA response with varying load radius and grid resolution. Figure 3a shows the bedrock deformation along a transect from the centre of the load for ice cylinders of 2, 5 and 10 km radius, with maximum bedrock uplift predicted on the finest 0.5 km grid of 48, 120 and 185 mm, respectively. Figure 3b indicates whether the grid over- (blue) or under- (red) represents the mass of the load within the model (i.e. where the “Mass Factor” is less or greater than 1). Errors in the GIA prediction compared to the result for the finest resolution 0.5 km grid (yellow lines) are typically the largest at the load centre where they underestimate the magnitude of peak displacement. Although a coarser grid may either under- or over-estimate the mass of loading represented in our model (Fig. 3b), it will always dampen the magnitude (effective height) of the load by spreading the load area over a larger grid region. For example, a 5 km radius load will be represented by 3 grid nodes on a 7.5 km grid, resulting in an overestimated 11.25 km radius loading footprint. For certain radius and grid combinations, the wider load footprint on a coarser grid leads to another zone of peak error occurring outside the load edge (e.g., compare dashed black line to yellow line in Fig. 3a). Even further from the load, the magnitude of deformation decreases and the results from various grid resolutions begin to converge.

These sensitivity test results highlight that the accuracy of predictions depends on the placement of the edge of the load relative to grid nodes and find that the load will be best captured if its edge lies sufficiently close to a grid node (e.g., in Fig. 3b the 1.75 km grid more closely captures the mass of a 2 km radius load than the 3 km radius load). Finally, the grid is unable to resolve the load when the grid resolution is more than approximately three times the radius of the load. This is illustrated, e.g., in the black solid line in Fig. 3a, where unloading a 2 km radius load on a 7.5 km grid resulted in no deformation, whereas a 3 km radius load is captured with a 7.5 km grid.

To quantify grid-related error, we calculate the difference between a given simulation and the corresponding simulation with the finest (0.5 km) resolution. We plot the root mean square error (RMSE; Fig. 3c) as an absolute measure of error and the average and standard deviation of the absolute percentage error (Figs. 3d, e) as a relative measure of error, beneath and within 2-km of the loaded region. Fig. 3c shows that the magnitude of RMSE remains relatively constant. This RMSE remained between ~10-20 mm for a 3.75 km grid, and ~ 20-40 mm for a 7.5 km grid, for example. As the load radius increases, the magnitude of deformation increases as well. However, the error due to grid resolution becomes less significant relative to the total deformation (i.e. the percentage error in Figs. 3d, e decreases).

While the dependence of grid performance on load position relative to grid nodes complicates matters, in order to arrive at an approximate relationship between grid resolution and load size, we assume a linear relationship between the two, which allows us to estimate, for this GIA model, a threshold beyond which grid-related error becomes sufficiently low to no longer merit a further refinement in grid resolution (Figs. 3d, e). For example, in the cross-sectional view of deformation in Fig. 3a, the 10 km radius load deformation is equally well represented by grid resolutions between 0.5 to 1.75 km. Considering the average absolute percentage error (Figs. 3d, e), we found that a 3:1 ratio (represented visually on Figs. 3b-e in the form of a black dashed line) between load radius and grid resolution (6:1 ratio between load diameter and grid resolution) brings the error to $< 7 \pm 3 \%$ (where 3% represents one standard deviation of the absolute percentage error calculated within 2 km of the load region). Furthermore, the mass of the load is accurately captured with no more than $\sim 7.5\%$ error with this 3:1 ratio (Fig. 3b).

The results from this analysis of spatially isolated cylindrical loads provide a rough estimate of the magnitude of error one can expect from a given model resolution and loading scenario, and can serve as a guide for selecting the appropriate grid resolution for a given problem. For example, for an input load with significant isolated locations of ice loss ~ 3 km in radius (or a ~ 6 km in diameter), a grid resolution of 1 km should be adopted. However, these idealized cylindrical load experiments are unlikely to capture the sensitivity of GIA predictions to grid resolution when realistic ice loss geometries are adopted. Such geometries are rarely characterized by spatially localized loads. Furthermore, these experiments capture only elastic deformation and neglect viscous effects, which can be significant on short timescales in low viscosity zones of the West Antarctic. In the following sections we explore how the dependence on grid resolution of GIA model predictions identified here changes when more realistic ice loss geometries and 3-D viscoelastic Earth structure are adopted.

4 Results with realistic modern and future ice loss in the Amundsen Sea

In this section, we consider the importance of grid resolution error for more realistic, spatially coherent modern and future ice loss scenarios. We begin with a consideration of the influence of grid resolution on sea level change in simulations adopting a purely elastic Earth model in section 4.1. Following this, we adopt 3-D viscoelastic Earth models to consider the contribution to sea level change from viscous deformation.

4.1 Influence of grid resolution on predictions of elastic deformation

Figures 2d-f show predicted sea level change in the Amundsen Sea Embayment of West Antarctica adopting an elastic Earth model for three different ice retreat scenarios, performed on a grid resolution of 1.9 km: one observationally constrained from 20 years of satellite altimetry data from 1997 to 2017 (ICE-SH; Fig. 2a), and two projected from dynamic ice sheet models for the coming century (ICE-GOL and ICE-RD; Figs. 2b-c). Sea level fall is predicted in the entire study region in all scenarios associated with the combination of sea surface subsidence and elastic bedrock uplift due to ice loss – the latter being the

dominant signal. Earth rotational effects are included in the predictions but are negligible compared to the other effects in the vicinity of the ice loss. For the modern, the maximum sea level fall from 1992 – 2017 reaches 0.68 m (Fig. 2d), while for
325 future ice loss projections, the sea level fall reaches up to 9.06 m and up to 12.8 m from 2000 – 2100 for ICE-GOL and ICE-RD respectively. Note that in addition to signal coming from local ice loss in the ASE, ice outside this region of interest also contributes a broad signal of smaller magnitude (see Supplementary Section S1).

To explore the resolution dependence of sea level predictions that adopt an elastic Earth model, we repeat the calculations in
330 Figs. 2d-f with a surface grid resolution ranging between 1.9 and 15 km. Fig. 4a-i, shows the difference between results for simulations performed at 1.9 km grid resolution relative to coarser resolutions. The coarser grid simulations tend to underestimate the magnitude of ice unloading and associated sea level fall in most of the domain (red regions in Fig. 4).

The highest grid resolution error occurs at the periphery of the load within a few kilometres of the final grounding line position
335 rather than at the location of maximum deformation (compare Figs. 4 to Fig.2). This suggests that high resolution is necessary for better representation of the load at the grounding line, rather than for representation of the smoother response of the solid earth – a finding consistent with the idealized experiments discussed in section 3 (see Fig. 3b). For example, for the ICE-GOL ice loss scenario, the greatest difference between 1.9 and 15 km grid simulations ranges occurs along the entire final grounding line (Fig. 4b), but the maximum sea level fall of over 9 meters occurs only on a concentrated region ~ 2 km away from the
340 grounding line (Fig. 2e).

The error decreases with increasing resolution, with minimal differences between the 1.9 km and 3.75 km grid resolutions. The maximum absolute error in the case of a 15 km grid (i.e. the maximum difference between the 15 km and 1.9 km resolution cases) is 44 cm at 2100 in ICE-RD (Fig. 4c), 47 cm at 2100 in ICE-GOL (Fig. 4b) and 9.1 cm after 25 years of modern melt
345 in ICE-SH (Fig. 4a). That is 3.4%, 5.2% and 13% of the peak elastic sea level fall predicted at that time for each respective scenario. The errors are approximately an order of magnitude smaller when a grid resolution of 3.75 km is adopted: 9 cm for ICE-RD, 5 cm for ICE-GOL and 0.7 cm for ICE-SH, or 0.7%, 0.4% and 1.0% of the peak elastic sea level fall respectively.

Since maximum grid resolution error is concentrated along the grounding line for elastic runs, in Fig. 5 we explore how the
350 error evolves during the ICE-RD simulation along a 10-km region bounding the grounding line. The error increases in absolute magnitude with increasing ice loss (“Error” in Fig. 5) but the relative error decreases across the same runs (“Percentage Error” in Fig. 5). In the case of 15 / 7.5 / 3.75 km grid resolutions, the peak error is ~10 / 5 / 1 cm at the 25 year mark of the simulations, and ~50 / 15 / 5 cm at the 150 year mark (whiskers in Fig. 5a top row). In contrast, the percentage, peaks at 20 / 6 / 1.5% of the signal at 25 years and drops to < 5 / < 2 / < 0.3% after 150 years (Fig. 5). This decrease in percent error with time reflects
355 that the ice geometry changes become broader in wavelength and can therefore be resolved by a coarser grid compared to the more spatially isolated changes occurring earlier in the simulation. Given the uncertainty in modelled and observed ice loss

and bedrock elevation in Antarctica (e.g. Morlinghem et al., 2020), we suggest that for most applications, errors of less than 5% can be achieved with a 7.5 km grid, and errors of less than 2% with a 3.75 km grid.

4.2 Contribution of viscous earth deformation to sea level predictions

360 So far we have focussed on the resolution dependence of the contribution to sea level change from elastic Earth deformation, as this has been a focus of recent literature (Larour et al., 2019). However, the Antarctic Ice Sheet is underlain by strongly laterally varying viscosity structure, and the Amundsen Sea region in particular is underlain by a low viscosity zone and thinned lithosphere (e.g. Barletta et al., 2018; Lloyd et al., 2020). Viscous deformation associated with ongoing ice loss is neglected in Larour et al. (2019) but is expected to be significant on decadal to centennial timescales in this region. In Figs. 2g-i the calculations of sea level change associated with the three ice loss scenarios shown in Figs. 2d-f are repeated with the 3-D viscoelastic Earth EM1_L described earlier. As with the elastic case, sea level falls beneath regions that experience ice loss in all three cases, but the magnitude of the sea level fall is significantly larger than predictions based on an elastic Earth model (compare Figs. 2g-i to Figs. 2d-f). In particular, peak sea level fall in this case reaches -0.79 m over 25 years in the ICE-SH ice loss scenario, and -14.9 m and -29.1 m from 2000 to 2100 for ice loss scenarios ICE-GOL and ICE-RD, respectively. The latter (Fig. 2i) is more than double the sea level calculated with the elastic Earth model (Fig. 2f).

Figure 6a-c shows the contribution of viscous Earth deformation to the sea level predictions, calculated by taking the difference between the full viscoelastic calculation shown in Figs. 3g-i and the calculation with an elastic Earth model shown in Figs. 3d-f. Over the 25 year modern ice loss scenario (Fig. 6a), viscous effects contribute up to 12 cm, or 15% of the peak viscoelastic prediction. In the future projections, the viscous contribution reaches up to 6 m of sea level fall, 41% of the peak prediction, within 100 years in predictions based on ICE-GOL and up to -17.7 m, or 61% of the peak viscoelastic signal for ICE-RD, making viscous effects the dominant contributor over elastic effects in this latter case. For the future projections, compared with the elastic signal (Figs. 3e-f), the zones of maximum viscous uplift and sea level fall (i.e. zones of intense red in Figs. 6b-c) are centered farther out beneath regions that experienced ice mass loss sooner in the simulation and have had more time for viscous deformation to occur (Figs. 6a-c), but as we highlight below, substantial viscous deformation still occurs along the current grounding line in the simulation. This is less evident in the modern because migration of the location of maximum ice mass loss is minimal. Faint blue areas further from the region of ice retreat in Fig. 6b indicate a sea level rise due to peripheral bulge subsidence, a viscous process that results from the return flow of mantle material assuming viscous incompressibility.

4.3 Influence of resolution on viscoelastic sea level predictions

385 In Fig. 7, we repeat the assessment of grid resolution error in Fig. 4, but with a viscoelastic rheology based on the 3-D earth model EM1_L. With the inclusion of viscous behaviour, the magnitude of the grid resolution error is similar to the elastic case (compare the range of errors on Figs. 7 and 4) but the spatial pattern of the error becomes more complex. The maximum error is no longer solely concentrated along the current grounding line since the solid earth continues to respond viscously to the

390 poorly resolved loading changes along previous locations of the grounding line. This is particularly evident in the ICE-RD simulations where the grounding line retreats across a large area. In this case, the grid resolution error over the region of past ice loss and grounding line migration is equal to or larger than the error along the active grounding line (Figs. 7b,c). The error increases during the simulation as viscous deformation builds, and suggests it also has a dependence on the 3D viscosity structure both due to lateral variation captured and the impact of a lower than average viscosity upper mantle in the WAIS, which we explore briefly in Section 4.3.

395

Note that in the blue region of Fig. 7c, the sign of the error is not the same as in Fig. 7f because ice retreat is not consistent within this particular region in ICE-RD. Specifically, between the years ~2020 to 2050 in ICE-RD, the grounding line in this blue region stays relatively fixed, experiencing multiple episodes of localized ice retreat and re-advance (unloading and loading). Situations of re-advance tend to occur at lateral scales < 15 km, such that these sub-grid scale movements were not 400 adequately captured with a coarser, 15 km grid. Along the final grounding line for ICE-RD, the coarser grids consistently result in a lower magnitude sea level fall as in the elastic case.

4.4 Earth model uncertainty

To investigate the influence of uncertainty in prescribed mantle viscosity structure, we compare simulations adopting five different earth model configurations: a globally averaged 1-D viscoelastic earth model, as used in EM1_M described in section 405 2.3, and three 3-D mantle viscosity configurations derived from seismic tomography models (EM1_L; EM1_M; EM2; see Sect. 2.3). Figs. 6d-f shows the difference in sea level predicted using the EM1_L and EM1_M models, which are based on the same underlying 3-D seismic velocity models but with a varying viscosity scaling factor. EM1_L (shown in Figs. 6a-c and 2g-i) has the lowest viscosity upper mantle beneath the ASE. Red regions in Fig. 6d-f indicate locations of higher predicted sea-level fall due to lower mantle viscosity in EM1_L, which results in shorter time scale viscous response. Differences reach 410 up to 5 cm after 25-years with ICE-SH, and up to 2.3 m and 5.8 m between 2000 and 2100 relative for ice loss scenarios ICE-GOL and ICE-RD, respectively. The simulation with EM2, a 3-D mantle viscosity model built from a different seismic tomography dataset, produced deformation with magnitudes intermediate to the simulations with EM1_L and EM1_M. We note that the rheological model for this area is uncertain, and our experiments do not comprehensively capture the full range of this uncertainty (see Whitehouse et al., 2019 for a more detailed discussion).

415 4.5 Time evolution of influence of grid resolution and viscous effects

To compare the relative contributions of grid resolution and Earth model differences over time, we extract predicted sea level time series from all simulations with the ICE-RD ice loading scenario and elastic and viscoelastic Earth models at two locations in Fig. 8: (X) the region experiencing the largest viscous uplift by 2100 (blue star), and (Y) the location of maximum ice loss at the 2100 grounding line (red star). Note that the shaded grey region from 1950-2000 in Figs. 8b and c represents a time of 420 hindcast spin up to the year 2000 in the ice sheet model simulation (see DeConto et al., 2021) rather than a realistic

representation of the ice cover changes in this region over this time period. In the case of the average 1-D viscoelastic earth model, the sea level response is similar to the elastic case (Fig. 8, compare red and black lines) because the upper mantle viscosity is set to 5×10^{20} Pa s and thus a significantly longer timescale is required for viscous effects to become significant. Differences between these two simulations and any of the simulations adopting 3-D Earth structure is larger. For example, the
425 differences between simulations using EM1_L and the global average 1-D viscoelastic earth model, reach up to 16.5 m and 12.7 m at the sites shown in Figs. 8b and c, respectively, by 2100.

To briefly explore the distinction between the impact on predictions of adding lateral variations in Earth structure and the impact of adopting a lower-than-average 1-D viscosity model representative of the structure underlying the WAIS, we ran an
430 additional simulation adopting a 1-D viscoelastic earth model for West Antarctica derived in Powell (2021). This model has a viscosity of 3.2×10^{19} Pa s in the shallow upper mantle, 1.3×10^{20} Pa s in the deep upper mantle, 2.0×10^{20} Pa s in the transition zone, and 5×10^{21} Pa s in the lower mantle. The predicted sea level changes with the lower viscosity 1-D Earth model shown by the purple lines in Fig. 8 lie closer to the 3-D earth model results than the global average 1-D earth model at sites X and Y. The degree to which the response of the laterally varying Earth structure in this region can be captured with a radially varying
435 approximation is explored in more detail in Powell (2021) and Blank et al. (2021).

Starting early in the century, the influence of grid resolution becomes smaller than the effect of adopting different Earth models (compare the differences between the dashed and solid lines to differences between different colored lines in Fig. 8). Using the 1-D average earth model (Fig. 8 red lines), viscous effects start to emerge after 50 years and reach only 4% of the peak
440 signal by the end of the run at year 2100. Nevertheless this signal is more significant than the error incurred by using 15 km grid resolution versus of 1.9 km grid resolution by 2040. With a 3-D earth rheology and low viscosities beneath the ASE, viscous effects are pronounced within decades in the simulation (blue and green lines in Fig. 8) and become larger than the difference in predictions based on 15 km and 1.9 km grid resolutions within 25-30 years and before substantial ice loss has occurred in the simulation.

445 Fig. 9 provides a more detailed picture along the grounding line of the contributions to differences in predicted sea level described in the preceding sections. We consider the impact of each factor on the predicted sea level signal at the end of the simulations, plotting the distribution of differences between simulations across all grid points within 10-km of the final grounding line. In interpreting Fig. 9, note once again that in viscoelastic runs, the region of maximum grid-resolution error does not necessarily occur near the grounding line (e.g. Figs. 7c, f). To visualise the distribution, we plot a classic box-whisker
450 diagram where the edges of the boxes represent (from left to right) the 25th percentile, median and 75th percentile of the distribution, whilst the whiskers represent the “minimum” and “maximum” (25th and 75th percentiles minus 1.5 times the interquartile range), overlain by a density curve. A box-whisker plot was chosen to mitigate against the effect of outlier points, which we plot as hollow diamonds.

Note that the error due to grid resolution consistently has a unimodal distribution that peaks at $\sim 0-1\%$ near the grounding line across a range of durations from 25-years of ICE-SH, to 100 and 150 years (starting from 1950) of ICE-RD. The hollow diamonds show that a significant number of points are statistical outliers, which is likely due to the fact that predicted sea-level change is low in magnitude at some regions along the grounding line that experience less ice-loss, causing even a small magnitude of error to contribute a large percentage error. Inclusive of these outliers, the range of percent error due to a 15-km grid peaks at $\sim 20\%$ after 25-years of the ICE-SH simulation, but decreases to less than $\pm 8\%$ by 100-years of the ICE-RD simulation as the magnitude of predicted sea level fall becomes larger across the entire region and the spread of outlier points diminishes. We conclude that the range denoted by the box-whisker plot likely provides a more accurate assessment of the error contributed by each factor in Fig. 9 in zones of active grounding line migration. However, in the following paragraphs we continue to describe the range of errors inclusive of outlier points so as to not under-estimate the possible spread.

Our results indicate that the differences due to the choice of adopted Earth model equals and, in most cases, exceeds the size of the error due to grid resolution near the grounding line by the end of all our simulations. Over the 25-year modern observed ice loss scenario, the difference in predictions associated with earth model configuration lies between $\sim 2-10\%$ within 10-km of the grounding line, which is within the range of error due to insufficient grid resolution in a viscoelastic run, which ranges from $\pm \sim 20\%$ with a 15 km grid, $\pm 6\%$ with a 7.5 km grid, and $< 3\%$ with a 3.75 km grid at all grid points (range of the diamonds outside the shaded distributions in Fig. 9a, though noting the discussion above, the percent error is substantially smaller than these end member values at most points). However, with more ice loss over longer timescales, the difference due to adopted Earth model far exceeds the grid resolution error (Figs 9b, c).

If we look beyond the grounding line and consider the difference in predicted sea level between different adopted 3-D viscosity models in the entire study region (difference in results with EM1_L and EM1_M, plotted in Fig. 6 d-f at 2100), the lower viscosity model results in additional viscous deformation that is up to 10.2 %, 21.4% and 20.9 % of the total signal after 25-years of ICE-SH, 100-years of ICE-RD and 150-years of ICE-RD respectively. In all cases, the error due to neglecting viscous effects far surpasses the error due to grid resolution with a 15-km grid (compare the bottom rows to top 3 rows of Figs 9b and c)., with the lower viscosity EM1_L model resulting in an additional viscous deformation that is up to 23.8 %, 58.9 %, 62.4 % of the elastic signal near the grounding line after 25-years of ICE-SH, and 100-years of ICE-RD and 150-years of ICE-RD respectively.

5 Discussion

Our study provides an assessment of the model grid resolution needed to capture decadal to centennial-scale GIA in the vicinity of active ice loss. We targeted the ASE in West Antarctica as our study location as it is a region with ongoing and projected

marine ice sheet retreat and where low mantle viscosity and thin lithosphere result in a rapid and localised GIA response to ice loading. Whilst we focus on the ASE where there is both ongoing and expected future ice loading and complex Earth structure, since the resolution error is primarily associated with the representation of the ice load, the general conclusions on the resolution required and results of the sensitivity experiments can be applied to any area of active, localised ice loading, for example in other parts of Antarctica, in Greenland or in the vicinity of smaller glaciers. We adopted a 3-D GIA model to accurately capture the viscoelastic response at high resolution, including the complexity introduced by laterally varying earth rheology in the region. Accurate assessments of solid earth deformation from past and present ice evolution are important for constraining the negative sea level - solid earth feedback on ice sheet retreat, and more accurate interpretation of geophysical observables. For the former, our study focusses on the sensitivity of GIA predictions along the ice sheet grounding line where this feedback occurs in Fig. 9.

5.1 Influence of grid resolution

For our suite of simulations with elastic and viscoelastic Earth models, modern and 21st century ice loss scenarios, and surface grid resolution ranging between 15 and 1.9 km, we found that improvements in the accuracy of model predictions with increasing grid resolution was limited, remaining within centimetres to decimetres at most at the grounding line. Furthermore, our results converged at higher resolutions, with errors from a 3.75 km grid resolution reaching at most 6 cm within 10 km of the grounding line in all simulations, even when the input ice sheet model results were available at 1 km resolution in the case of ICE-RD. The error introduced in assuming an elastic Earth model and neglecting viscous deformation in the ASE builds to an order of magnitude or more larger than the grid resolution error within three to four decades, and up to tens of meters by the end of the century. In addition, predictions adopting different 3-D Earth models that reflect the uncertainty in viscoelastic Earth structure in the region diverge by up to 1 meter within 50 years and upwards of 2-3 meters after 100 years in the simulation.

For coupled ice-sheet - GIA model applications, our results suggest that adopting high resolution in the ice sheet model does not require a similarly high-resolution GIA model. In our simulations, a 3.75 km grid was sufficient to bring errors to < 2% along the grounding line for all scenarios (Fig. 9). Furthermore, this percentage decreased over time in our simulations, and would continue to decrease multi-century and millennial simulations as the magnitude of viscous deformation and the scale of the ice loss continue to grow. While bedrock topography has smaller scale features (Morlinghem et al., 2020), our results suggest that the GIA signal is less localized and may be computed at lower resolution relative to the ice sheet dynamics and then interpolated and added to the initial topography on the higher resolution ice sheet model grid, as is done in, e.g., Gomez et al. (2015) and DeConto et al. (2021).

Our results showing that the location of maximum error consistently lies along the load edge for elastic model runs (Fig. 4a-c) suggest that the error due to coarse model resolution is predominantly a result of poor representation of surface ice cover

520 changes rather than representation of the smoother response of the solid Earth. For the latter, we would expect the error to occur instead at the location of maximum GIA response (compare differing spatial patterns in Fig. 2e, f to 4b, c). When the viscous response is incorporated, the time-evolving nature of viscous deformation leads to an additional peak in grid resolution error at locations predominantly downstream of the grounding line due to inaccurate representation of past loading (Fig. 7). This additional zone of error will not affect active ice sheet grounding lines (though it may be important for interpretation of modern records), and while the spatial pattern of the error differs, the magnitude of the error due to grid resolution was similar across both elastic and viscoelastic simulations. However, it is important to note that deformation downstream of the grounding line can be important for ice sheet stabilization if it causes re-grounding of an ice shelf.

Our findings on the size and source of resolution error are in contrast to recent work by Larour et al., (2019) who suggested that kilometre resolution was required to capture elastic deformation. This discrepancy may be due in part to Larour et al. (2019) considering only point loads in their idealized resolution experiment, while our conclusions are based on more realistic, spatially coherent ice loss scenarios. Differences may also arise due to the nature of the computational grid and processing of inputs (see section 5.3) which should be explored in more detail in future GIA model inter comparison efforts. Nevertheless, our predictions based on an elastic Earth model converge to theirs for more spatially broad loads.

535 One possible limitation in this study is we do not reach sub-km grid resolution in our GIA model, and our highest resolution ice model is 1 km. In sensitivity tests with idealized loading scenarios in Section 3 we adopted a grid resolution as low as 0.5 km grid and found that a minimum 1:3 ratio between grid resolution and load radius was required for results to incur an error of $< 5 \pm 3$ (σ) % along the load edge, suggesting that a 3.75km grid would be unable to capture a spatially isolated, < 1 km radius ice unloading event. That we did not see significant error at this resolution in the realistic simulations indicates that the ice cover changes are spatially coherent and there are no significant spatially isolated ice unloading events (i.e. no ice thickness changes occurring over only a few grid points) predicted in the 1-km resolution ICE-RD ice model simulation (Fig. 5).

To further investigate if short-wavelength, spatially isolated ice loss scenarios exist over Antarctica, we assessed the surface elevation change observables from 40 and 25 years of multi-mission satellite altimetry data by Schröder et al., (2019) and Shepherd et al. (2019) respectively, and 15 years of airborne laser altimetry from Operation IceBridge (OIB ATM L4; Studinger, 2014 (Updated 2018)). While a more detailed investigation is merited, in our initial analysis of these datasets we observed that spatially isolated ice loss events have a lower magnitude, only persist over short timescales, and found no evidence of high magnitude, short-wavelength ice loss occurring with spatial scales < 5 km. Thus, we expect that spatially isolated ice unloading occurs rarely and will not have a significant impact on the overall accuracy of GIA model results in a given region. Nonetheless, with improving observational products and ice sheet model resolutions, we expect to obtain regional-scale ice loading grids of sub-km resolution that may warrant further study with a sub-km GIA model grid (e.g. Durkin et al., 2020).

5.2 Influence of viscous deformation and earth model uncertainty

555 Within decades in the ASE, viscous deformation is a significant contributor to the GIA signal (Hay et al., 2017; Barletta et al.,
2018; Powell et al., 2020; Kachuck et al., 2020). The GIA response can be decomposed into the following: perturbation of the
sea surface equipotential, elastic deformation and viscous deformation of the solid earth surface. Previous studies have isolated
the GIA signals from each of these components to assess the relative importance of each factor on ice sheet dynamics. Over
560 decadal to centennial timescales, Larour et al. (2019) show that purely elastic deformation was more significant than the sea
surface perturbation on continental-scales, while Kachuck et al. (2020) found that viscous deformation is more significant than
either elastic deformation or the perturbation to the sea surface. In this study, we have confirmed that viscous deformation
effects are significant within decades, particularly in the low viscosity region of the ASE, where the viscous component of
deformation can reach multi-metre scales by the end of the century. This body of work highlights the importance of
incorporating viscous behaviour in GIA modelling applications in regions of low mantle viscosity.

565

Complicating efforts to accurately characterise the viscous deformation behaviour is the uncertainty in Earth's viscosity
structure. The timescale of viscous solid-earth deformation on ice sheet dynamics is strongly dependent on the assumed earth
rheology. The global average mantle viscosity of $\sim 10^{21}$ Pa s (Forte and Mitrovica, 1996) corresponds to response times from
centuries to millennia, whilst recent seismic (Lloyd et al., 2020) and GPS observations suggest an upper mantle viscosity under
570 the ASE as low as $\sim 10^{18}$ Pa s (Barletta et al., 2018). Rapid viscous uplift response was similarly identified in Kachuck et al.
(2020) which used a 2-D GIA model of mantle viscoelastic deformation and found that sea level fall associated with
viscoelastic mantle deformation led to a 30% reduction in modelled ice sheet volume loss by 2150. Our study compares results
generated with three 3-D earth rheology models; EM1_M and EM2 have a comparable viscosity range, while EM1_L has the
lowest viscosity values under the ASE. These results demonstrate that uncertainties regarding the effective mantle viscosity
575 are significant and can contribute up to multiple metres of uncertainty by the end of a 100-year simulation (Fig. 6). Furthermore,
additional uncertainty arises from the model of viscoelastic behaviour. We adopt a viscous Maxwell rheology, but studies
suggest that incorporating a short-term transient component of deformation may result in even faster viscous deformation (e.g.
Pollitz, 2019).

580 Finally, we note that the required resolution of the GIA model grid will depend on the resolution of the seismic model used to
construct the 3D Earth structure model. Earth structure is currently resolved in seismic tomography models in Antarctica at
length scales of $\mathcal{O}(100 \text{ km})$ or greater (Lloyd et al., 2020; Lucas et al., 2020), but as further improvements in the resolution of
seismic tomography emerge, variations in earth properties at even shorter spatial scales may be revealed and need to be
represented in GIA models. However, given the smooth nature of the GIA response, we expect that the wavelength of ice
585 loading variations will remain the determining factor of surface GIA model grid resolution requirements.

5.3 On GIA model setup

In choosing a method for representing a finer resolution load grid onto a coarser model grid, we found that it is important to consider how the model itself discretizes the load, and the input load interpolation schemes. Here, it is worth noting that our GIA model grid is a tetrahedral grid (triangular grid on surface), and these findings may not translate perfectly to other model grid compositions. Our GIA model grid consists of a uniform global tetrahedral grid that allows for regional patches of refined resolution (also uniform) but does not permit matching of model grid nodes to the input grid. For our experiments, by comparing the volumes of the input ice calculated on the input and GIA model grids, we found that the in-built Poisson interpolation scheme (Latychev et al., 2005) performed better in interpolating the finest resolution load grid onto the model grid compared to other tested schemes, suggesting that an understanding of the method in which the load is mapped onto the model grid nodes is important. Additionally, we note that considerations such as the resolution of the input ice sheet model and treatment of the ice cover outside the region of interest also have an influence on the final GIA model predictions (see Supplementary Section S1) and should be explored further in future studies.

6 Conclusion

In this study, we present a comprehensive analysis of the influence of grid resolution on modelled GIA effects in response to ice cover changes in the ASE over the modern satellite era and through the 21st century. We adopt a range of Earth models including models that capture lateral variations in Earth structure based on seismic tomography and GNSS analyses. These experiments showed that: (1) the grid resolution error introduced through adopting a 15 km grid relative to a 1.9 km model grid remains within centimetres to decimetres (which can go up to 15% of the total signal along the grounding line) throughout our simulations; (2) the grid resolution error is the highest along load edges for purely elastic deformation cases, and along past and current grounding line positions for viscoelastic Earth models, and is primarily associated with the representation of the surface load; (3) results with grid refinement beyond 3.75 km converged in our simulations, even when adopting a 1 km resolution input load, and this likely represents a conservative lower bound since the next coarser grid we considered was 7.5 km. The errors associated with the choice of grid resolution will decrease with time for longer simulations as the extent and magnitude of ice loss and associated GIA response increase. Furthermore, comparison of simulations adopting elastic and 3-D viscoelastic Earth models demonstrate that the contribution of viscous deformation can be up to tens of metres over the 21st century, or > 50% of the total deformation signal. Additionally, uncertainties in earth properties can contribute up to several metres of error. This indicates the importance of considering viscous uplift when modelling GIA over decadal to centennial timescales in the ASE. In comparison, the error due to grid resolution is negligible for grids of spacing of 3.75 km and less.

To supplement these findings with realistic loading, we conducted a sensitivity test with cylindrical loads with radii from 16 km to 0.5 km and grid resolutions from 7.5 to 0.5 km. These indicate a minimum 1:3 ratio between the required grid resolution and the load radius (or 1:6 with load diameter) to minimise grid resolution error. However, no significant spatially isolated

loads occur in our adopted observation- and model-based ice loss scenarios, and a preliminary examination of other ice observation and modelled products suggest that significant ice loss with < 5 km wavelength is rare in the ASE. These results, taken together, support the conclusion that km-scale resolution in GIA modelling is generally not necessary. However, as higher resolution sub-km ice observational and dynamic ice model grid products are released, this guidance may have to be revisited.

Code/Data Availability

We will make all model output from the sensitivity tests and more realistic simulations available on a public repository at https://osf.io/2vmqh/?view_only=4d08e562720941688e0644b80781eeaa. The 3-D GIA model adopted here has been used in numerous previous studies, questions regarding the model or requests for additional output can be discussed with the corresponding author and K.L., the developer of the code. Additional data related to this paper may be requested from the authors.

Author Contribution

J.X.W.W. and N.G. developed the ideas and experiments in the study with input from K.L. and H.K.H.. K.L. contributed to designing the experimental set up. J.X.W.W. performed the simulations and analysis. J.X.W.W. and N.G. wrote the original text and all authors contributed to revisions.

Competing interests

The authors declare they have no competing interests.

References

- Adhikari, S., Ivins, E. R., and Larour, E.: ISSM-SESAW v1. 0: Mesh-based computation of gravitationally consistent sea-level and geodetic signatures caused by cryosphere and climate driven mass change, *Geoscientific Model Development*, 9, 1087-1109, <https://doi.org/10.5194/gmd-9-1087-2016>, 2016.
- An, M., Wiens, D. A., Zhao, Y., Feng, M., Nyblade, A. A., Kanao, M., Li, Y., Maggi, A., and L  v  que, J. J.: S-velocity model and inferred Moho topography beneath the Antarctic Plate from Rayleigh waves, *Journal of Geophysical Research: Solid Earth*, 120, 359-383, <https://doi.org/10.1002/2014JB011332>, 2015.
- An, M., Wiens, D. A., Zhao, Y., Feng, M., Nyblade, A., Kanao, M., Li, Y., Maggi, A., and L  v  que, J. J.: Temperature, lithosphere-asthenosphere boundary, and heat flux beneath the Antarctic Plate inferred from seismic velocities, *Journal of Geophysical Research: Solid Earth*, 120, 8720-8742, <https://doi.org/10.1002/2015JB011917>, 2015.

- Austermann, J., Mitrovica, J. X., Latychev, K., and Milne, G. A.: Barbados-based estimate of ice volume at Last Glacial Maximum affected by subducted plate, *Nature Geoscience*, 6, 553-557, <https://doi.org/10.1038/ngeo1859>, 2013.
- Bamber, J. L. and Dawson, G. J.: Complex evolving patterns of mass loss from Antarctica's largest glacier, *Nature Geoscience*, 13, 127-131, <https://doi.org/10.1038/s41561-019-0527-z>, 2020.
- 650 Barletta, V. R., Bevis, M., Smith, B. E., Wilson, T., Brown, A., Bordoni, A., Willis, M., Khan, S. A., Rovira-Navarro, M., Dalziel, I., Smalley, R., Jr., Kendrick, E., Konfal, S., Caccamise, D. J., 2nd, Aster, R. C., Nyblade, A., and Wiens, D. A.: Observed rapid bedrock uplift in Amundsen Sea Embayment promotes ice-sheet stability, *Science*, 360, 1335-1339, <https://doi.org/10.1126/science.aao1447>, 2018.
- 655 Berger, S., Favier, L., Drews, R., Derwael, J.-J., and Pattyn, F.: The control of an uncharted pinning point on the flow of an Antarctic ice shelf, *Journal of Glaciology*, 62, 37-45, <https://doi.org/10.1017/jog.2016.7>, 2016.
- Blank, B., Barletta, V., Hu, H., Pappa, F., & van der Wal, W.: Effect of lateral and stress-dependent viscosity variations on GIA induced uplift rates in the Amundsen Sea Embayment. *Geochemistry, Geophysics, Geosystems*, 22.9, <https://doi.org/10.1029/2021GC009807>, 2021
- 660 Carrivick, J. L., Davies, B. J., James, W. H., McMillan, M., and Glasser, N. F.: A comparison of modelled ice thickness and volume across the entire Antarctic Peninsula region, *Geografiska Annaler: Series A, Physical Geography*, 101, 45-67, <https://doi.org/10.1080/04353676.2018.1539830>, 2019.
- Conrad, C. P. and Lithgow-Bertelloni, C.: Influence of continental roots and asthenosphere on plate-mantle coupling, *Geophysical Research Letters*, 33, <https://doi.org/10.1029/2005GL025621>, 2006.
- 665 Cornford, S., Martin, D., Lee, V., Payne, A., and Ng, E.: Adaptive mesh refinement versus subgrid friction interpolation in simulations of Antarctic ice dynamics, *Annals of Glaciology*, 57, 1-9, <https://doi.org/10.1017/aog.2016.13>, 2016.
- De Boer, B., Stocchi, P., and Van De Wal, R.: A fully coupled 3-D ice-sheet–sea-level model: algorithm and applications, *Geoscientific Model Development*, 7, 2141-2156, <https://doi.org/10.5194/gmd-7-2141-2014>, 2014.
- 670 DeConto, R. M., Pollard, D., Alley, R. B., Velicogna, I., Gasson, E., Gomez, N., Sadai, S., Condron, A., Gilford, D. M., Ashe, E. L., Kopp, R. E., Li, D., and Dutton, A.: The Paris Climate Agreement and future sea-level rise from Antarctica, *Nature*, 593, 83-89, <https://doi.org/10.1038/s41586-021-03427-0>, 2021.
- Durand, G., Gagliardini, O., Zwinger, T., Le Meur, E., and Hindmarsh, R. C.: Full Stokes modeling of marine ice sheets: influence of the grid size, *Annals of Glaciology*, 50, 109-114, <https://doi.org/10.3189/172756409789624283>, 2009.
- Durkin IV, W., Wilson, T.J., Hansen, J.S., Willis, M. and Bevis, M.G., 2020, December. Reevaluating the Elastic Response to Ice Mass Change in Antarctica. In AGU Fall Meeting Abstracts (Vol. 2020, pp. G012-0023).
- 675 Favier, L., Gagliardini, O., Durand, G., and Zwinger, T.: A three-dimensional full Stokes model of the grounding line dynamics: effect of a pinning point beneath the ice shelf, *The Cryosphere*, 6, 101-112, <https://doi.org/10.5194/tc-6-101-2012>, 2012.
- 680 Fretwell, P., Pritchard, H.D., Vaughan, D.G., Bamber, J.L., Barrand, N.E., Bell, R., Bianchi, C., Bingham, R.G., Blankenship, D.D., Casassa, G. and Catania, G.: Bedmap2: improved ice bed, surface and thickness datasets for Antarctica. *The Cryosphere*, 7, 375-393, <https://doi.org/10.5194/tc-7-375-2013>, 2013.
- Forte, A.M. and Mitrovica, J.X.: New inferences of mantle viscosity from joint inversion of long-wavelength mantle convection and post-glacial rebound data, *Geophysical Research Letters*, 23, 1147-1150, <https://doi.org/10.1029/96GL00964>, 1996

- 685 Gladstone, R. M., Payne, A. J., and Cornford, S. L.: Resolution requirements for grounding-line modelling: sensitivity to basal drag and ice-shelf buttressing, *Annals of glaciology*, 53, 97-105, <https://doi.org/10.3189/2012AoG60A148>, 2012.
- Golledge, N. R., Keller, E. D., Gomez, N., Naughten, K. A., Bernales, J., Trusel, L. D., and Edwards, T. L.: Global environmental consequences of twenty-first-century ice-sheet melt, *Nature*, 566, 65-72, <https://doi.org/10.1038/s41586-019-0889-9>, 2019.
- 690 Gomez, N., Latychev, K., and Pollard, D.: A coupled ice sheet–sea level model incorporating 3D earth structure: variations in Antarctica during the last deglacial retreat, *Journal of Climate*, 31, 4041-4054, <https://doi.org/10.1175/JCLI-D-17-0352.1>, 2018.
- Gomez, N., Mitrovica, J. X., Huybers, P., and Clark, P. U.: Sea level as a stabilizing factor for marine-ice-sheet grounding lines, *Nature Geoscience*, 3, 850-853, <https://doi.org/10.1038/ngeo1012>, 2010.
- 695 Gomez, N., Pollard, D., and Holland, D.: Sea-level feedback lowers projections of future Antarctic Ice-Sheet mass loss, *Nat Commun*, 6, 8798, <https://doi.org/10.1038/ncomms9798>, 2015.
- Gomez, N., Pollard, D., and Mitrovica, J. X.: A 3-D coupled ice sheet–sea level model applied to Antarctica through the last 40 ky, *Earth and Planetary Science Letters*, 384, 88-99, <https://doi.org/10.1016/j.epsl.2013.09.042>, 2013.
- 700 Hay, C. C., Lau, H. C., Gomez, N., Austermann, J., Powell, E., Mitrovica, J. X., Latychev, K., and Wiens, D. A.: Sea level fingerprints in a region of complex Earth structure: The case of WAIS, *Journal of Climate*, 30, 1881-1892, <https://doi.org/10.1175/JCLI-D-16-0388.1>, 2017.
- Heeszel, D. S., Wiens, D. A., Anandakrishnan, S., Aster, R. C., Dalziel, I. W., Huerta, A. D., Nyblade, A. A., Wilson, T. J., and Winberry, J. P.: Upper mantle structure of central and West Antarctica from array analysis of Rayleigh wave phase velocities, *Journal of Geophysical Research: Solid Earth*, 121, 1758-1775, <https://doi.org/10.1002/2015JB012616>, 2016.
- 705 Joughin, I., Smith, B. E., and Medley, B.: Marine ice sheet collapse potentially under way for the Thwaites Glacier Basin, West Antarctica, *Science*, 344, 735-738, <https://doi.org/10.1126/science.1249055>, 2014.
- Kachuck, S. B., Martin, D. F., Bassis, J. N., and Price, S. F.: Rapid viscoelastic deformation slows marine ice sheet instability at Pine Island Glacier, *Geophysical Research Letters*, 47, e2019GL086446, <https://doi.org/10.1029/2019GL086446>, 2020.
- Kaufmann, G., Wu, P., and Ivins, E. R.: Lateral viscosity variations beneath Antarctica and their implications on regional rebound motions and seismotectonics, *Journal of Geodynamics*, 39, 165-181, <https://doi.org/10.1016/j.jog.2004.08.009>, 2005.
- 710 Kendall, R. A., Mitrovica, J. X., and Milne, G. A.: On post-glacial sea level–II. Numerical formulation and comparative results on spherically symmetric models, *Geophysical Journal International*, 161, 679-706, <https://doi.org/10.1111/j.1365-246X.2005.02553.x>, 2005.
- King, M. A., Bingham, R. J., Moore, P., Whitehouse, P. L., Bentley, M. J., and Milne, G. A.: Lower satellite-gravimetry estimates of Antarctic sea-level contribution, *Nature*, 491, 586-589, <https://doi.org/10.1038/nature11621>, 2012.
- 715 Konrad, H., Sasgen, I., Pollard, D., and Klemann, V.: Potential of the solid-Earth response for limiting long-term West Antarctic Ice Sheet retreat in a warming climate, *Earth and Planetary Science Letters*, 432, 254-264, <https://doi.org/10.1016/j.epsl.2015.10.008>, 2015.
- Kustowski, B., Ekström, G., and Dziewoński, A.: Anisotropic shear-wave velocity structure of the Earth's mantle: A global model, *Journal of Geophysical Research: Solid Earth*, 113, <https://doi.org/10.1029/2007JB005169>, 2008.

- 720 Ivins, E.R. and Sammis, C.G.: On lateral viscosity contrast in the mantle and the rheology of low-frequency geodynamics. *Geophysical Journal International*, 123, 305-322, <https://doi.org/10.1111/j.1365-246X.1995.tb06856.x>, 1995.
- Lambeck, K., Rouby, H., Purcell, A., Sun, Y., and Sambridge, M.: Sea level and global ice volumes from the Last Glacial Maximum to the Holocene, *Proc Natl Acad Sci U S A*, 111, 15296-15303, <https://doi.org/10.1073/pnas.1411762111>, 2014.
- 725 Larour, E., Seroussi, H., Adhikari, S., Ivins, E., Caron, L., Morlighem, M., and Schlegel, N.: Slowdown in Antarctic mass loss from solid Earth and sea-level feedbacks, *Science*, 364, <https://doi.org/10.1126/science.aav7908>, 2019.
- Latychev, K., Mitrovica, J. X., Tromp, J., Tamisiea, M. E., Komatitsch, D., and Christara, C. C.: Glacial isostatic adjustment on 3-D Earth models: a finite-volume formulation, *Geophysical Journal International*, 161, 421-444, <https://doi.org/10.1111/j.1365-246X.2005.02536.x>, 2005.
- 730 Lloyd, A., Wiens, D., Zhu, H., Tromp, J., Nyblade, A., Aster, R., Hansen, S., Dalziel, I., Wilson, T., and Ivins, E.: Seismic structure of the Antarctic upper mantle imaged with adjoint tomography, *Journal of Geophysical Research: Solid Earth*, 125, <https://doi.org/10.1029/2019JB017823>, 2020.
- 735 Martín-Español, A., Zammit-Mangion, A., Clarke, P.J., Flament, T., Helm, V., King, M.A., Luthcke, S.B., Petrie, E., Rémy, F., Schön, N. and Wouters, B.: Spatial and temporal Antarctic Ice Sheet mass trends, glacio-isostatic adjustment, and surface processes from a joint inversion of satellite altimeter, gravity, and GPS data, *Journal of Geophysical Research: Earth Surface*, 121, 182-200, <https://doi.org/10.1002/2015JF003550>, 2016.
- Mitrovica, J. and Forte, A.: A new inference of mantle viscosity based upon joint inversion of convection and glacial isostatic adjustment data, *Earth and Planetary Science Letters*, 225, 177-189, <https://doi.org/10.1016/j.epsl.2004.06.005>, 2004.
- Morelli, A. and Danesi, S.: Seismological imaging of the Antarctic continental lithosphere: a review, *Global and Planetary Change*, 42, 155-165, <https://doi.org/10.1016/j.gloplacha.2003.12.005>, 2004.
- 740 Morlighem, M., Rignot, E., Binder, T., Blankenship, D., Drews, R., Eagles, G., Eisen, O., Ferraccioli, F., Forsberg, R., and Fretwell, P.: Deep glacial troughs and stabilizing ridges unveiled beneath the margins of the Antarctic ice sheet, *Nature Geoscience*, 13, 132-137, <https://doi.org/10.1038/s41561-019-0510-8>, 2020.
- NOAA National Geophysical Data Centre: 2-minute gridded global relief data (ETOPO2) v2, 2006.
- 745 Nield, G. A., Barletta, V. R., Bordoni, A., King, M. A., Whitehouse, P. L., Clarke, P. J., Domack, E., Scambos, T. A., and Berthier, E.: Rapid bedrock uplift in the Antarctic Peninsula explained by viscoelastic response to recent ice unloading, *Earth and Planetary Science Letters*, 397, 32-41, <https://doi.org/10.1016/j.epsl.2014.04.019>, 2014.
- Nowicki, S. M. J., Payne, T., Larour, E., Seroussi, H., Goelzer, H., Lipscomb, W., Gregory, J., Abe-Ouchi, A., and Shepherd, A.: Ice Sheet Model Intercomparison Project (ISMIP6) contribution to CMIP6, *Geosci Model Dev*, 9, 4521-4545, <https://doi.org/10.5194/gmd-9-4521-2016>, 2016.
- 750 Paterson, W. and Colbeck, S.: Ice sheets and ice shelves, in: *Dynamics of snow and ice masses*, Academic Press, 1-78, 1980.
- Pattyn, F., Perichon, L., Durand, G., Favier, L., Gagliardini, O., Hindmarsh, R. C., Zwinger, T., Albrecht, T., Cornford, S., and Docquier, D.: Grounding-line migration in plan-view marine ice-sheet models: results of the ice2sea MISMIP3d intercomparison, *Journal of Glaciology*, 59, 410-422, <https://doi.org/10.3189/2013JoG12J129>, 2013.
- 755 Pollitz, F. F.: Lithosphere and shallow asthenosphere rheology from observations of post-earthquake relaxation, *Physics of the Earth and Planetary Interiors*, 293, 106271, <https://doi.org/10.1016/j.pepi.2019.106271>, 2019.


- Powell, E., Gomez, N., Hay, C., Latychev, K., and Mitrovica, J.: Viscous effects in the solid Earth response to modern Antarctic ice mass flux: Implications for geodetic studies of WAIS stability in a warming world, *Journal of Climate*, 33, 443-459, <https://doi.org/10.1175/JCLI-D-19-0479.1>, 2020.
- 760 Powell, E.: Glacial Isostatic Adjustment in a region of complex Earth structure: The case of WAIS, Doctoral dissertation, Harvard University Graduate School of Arts and Sciences, Chapter 4, 2021.
- Rignot, E., Mouginot, J., Scheuchl, B., van den Broeke, M., van Wessem, M. J., and Morlighem, M.: Four decades of Antarctic Ice Sheet mass balance from 1979-2017, *Proc Natl Acad Sci U S A*, 116, 1095-1103, <https://doi.org/10.1073/pnas.1812883116>, 2019.
- 765 Ritsema, J., Deuss, a. A., Van Heijst, H., and Woodhouse, J.: S40RTS: a degree-40 shear-velocity model for the mantle from new Rayleigh wave dispersion, teleseismic traveltime and normal-mode splitting function measurements, *Geophysical Journal International*, 184, 1223-1236, <https://doi.org/10.1111/j.1365-246X.2010.04884.x>, 2011.
- Ritzwoller, M. H., Shapiro, N. M., Levshin, A. L., and Leahy, G. M.: Crustal and upper mantle structure beneath Antarctica and surrounding oceans, *Journal of Geophysical Research: Solid Earth*, 106, 30645-30670, <https://doi.org/10.1029/2001JB000179>, 2001.
- 770 Schröder, L., Horwath, M., Dietrich, R., Helm, V., Broeke, M. R., and Ligtenberg, S. R.: Four decades of Antarctic surface elevation changes from multi-mission satellite altimetry, *The Cryosphere*, 13, 427-449, <https://doi.org/10.5194/tc-13-427-2019>, 2019.
- Seroussi, H., Nowicki, S., Payne, A. J., Goelzer, H., Lipscomb, W. H., Abe-Ouchi, A., Agosta, C., Albrecht, T., Asay-Davis, X., and Barthel, A.: ISMIP6 Antarctica: a multi-model ensemble of the Antarctic ice sheet evolution over the 21st century, *The Cryosphere*, 14, 3033-3070, <https://doi.org/10.5194/tc-14-3033-2020>, 2020.
- 775 Shen, W., Wiens, D. A., Anandakrishnan, S., Aster, R. C., Gerstoft, P., Bromirski, P. D., Hansen, S. E., Dalziel, I. W., Heeszel, D. S., and Huerta, A. D.: The crust and upper mantle structure of central and West Antarctica from Bayesian inversion of Rayleigh wave and receiver functions, *Journal of Geophysical Research: Solid Earth*, 123, 7824-7849, <https://doi.org/10.1029/2017JB015346>, 2018.
- 780 Shepherd, A., Gilbert, L., Muir, A. S., Konrad, H., McMillan, M., Slater, T., Briggs, K. H., Sundal, A. V., Hogg, A. E., and Engdahl, M. E.: Trends in Antarctic Ice Sheet elevation and mass, *Geophysical Research Letters*, 46, 8174-8183, <https://doi.org/10.1029/2019GL082182>, 2019.
- Smith, B., Fricker, H. A., Gardner, A. S., Medley, B., Nilsson, J., Paolo, F. S., Holschuh, N., Adusumilli, S., Brunt, K., Csatho, B., Harbeck, K., Markus, T., Neumann, T., Siegfried, M. R., and Zwally, H. J.: Pervasive ice sheet mass loss reflects competing ocean and atmosphere processes, *Science*, 368, 1239-1242, <https://doi.org/10.1126/science.aaz5845>, 2020.
- 785 Spada, G. and Stocchi, P.: SELEN: A Fortran 90 program for solving the “sea-level equation”, *Computers & Geosciences*, 33, 538-562, <https://doi.org/10.1016/j.cageo.2006.08.006>, 2007.
- Studinger, M.: IceBridge ATM L4 surface elevation rate of change, NASA National Snow and Ice Data Center Distributed Active Archive Center, Boulder, Colorado USA, 10, 2014.
- 790 The IMBIE team: Mass balance of the Antarctic Ice Sheet from 1992 to 2017. *Nature* 558, 219–222, <https://doi.org/10.1038/s41586-018-0179-y>, 2018.
- Van den Berg, J., Van de Wal, R., and Oerlemans, J.: Effects of spatial discretization in ice-sheet modelling using the shallow-ice approximation, *Journal of glaciology*, 52, 89-98, <https://doi.org/10.3189/172756506781828935>, 2006.

795 van der Wal, W., Whitehouse, P. L., & Schrama, E. J.: Effect of GIA models with 3D composite mantle viscosity on GRACE mass balance estimates for Antarctica, Earth and Planetary Science Letters, 414, 134-143, <https://doi.org/10.1016/j.epsl.2015.01.001>, 2015.


Whitehouse, P. L., Gomez, N., King, M. A., & Wiens, D. A.: Solid Earth change and the evolution of the Antarctic Ice Sheet, Nature Communications, 10, 503, <https://doi.org/10.1038/s41467-018-08068-y>, 2019.

800 Zhao, C., King, M. A., Watson, C. S., Barletta, V. R., Bordoni, A., Dell, M., and Whitehouse, P. L.: Rapid ice unloading in the Fleming Glacier region, southern Antarctic Peninsula, and its effect on bedrock uplift rates, Earth and Planetary Science Letters, 473, 164-176, <https://doi.org/10.1016/j.epsl.2017.06.002>, 2017.

805 Zwinger, T., Nield, G. A., Ruokolainen, J., and King, M. A.: A new open-source viscoelastic solid earth deformation module implemented in Elmer (v8. 4), Geoscientific Model Development, 13, 1155-1164, <https://doi.org/10.5194/gmd-13-1155-2020>, 2020.




810 **Figure 1. Grid and Earth model configuration. (a-b) Configuration of the tetrahedral grid in the finite-volume 3-D GIA model with regional refinement, used for observational and modelled ice loading scenarios. (a) shows a cross-sectional view of the regional refinement along ASE. (b) indicates areas of grid refinement across Antarctica with a surface grid resolution of 7.5 km over all Antarctica in black, 3.75 km over a section of the West Antarctica in magenta and 1.9 km in the ASE (light blue square). (c-d)**
 815 **Logarithmic viscosity perturbation map at depth 200 km for low upper mantle viscosity model EM1_L over (c) Antarctica and (d) our study region in the Amundsen Sea Embayment. Values are relative to reference 1-D model with upper mantle viscosity of 1×10^{20} Pa s, and lower mantle viscosity of 5×10^{21} Pa s. The black line delimits the Antarctic grounding line, and the grayline shows the bedrock topography contour at 0 m from Bedmap2 (Fretwell et al., 2013) highlighting marine-based sectors of ice. Transparent patch in (c) contains no data on mantle viscosity as it the region contains lithosphere at 200 km depth.**




820

Figure 2. Ice loading scenarios and corresponding elastic sea level predictions in the ASE. (a – c) Total ice thickness change in meters predicted from (a) 1997 to 2017 in the observation-based ICE-SH ice model (Shepherd et al., 2019), and from 2000 to 2100 in the (b) ICE-GOL (Golledge et al., 2019) and ICE-RD (DeConto et al., 2021) ice model projections. (d – f) show the predicted sea level change in meters with an elastic earth model associated with the ice cover changes shown in (a-c). (g– i) as in (d-f) but adopting 3-D

825 viscoelastic earth model EMI_L. All sea level predictions were performed on a 1.9 km resolution grid. The black and blue line indicates final and initial grounding lines, respectively, for each simulation. Each frame is annotated with the maximum and minimum value within the frame. Note that the colour bars change across each frame.



830 **Figure 3. Idealized sensitivity experiment of the effect of surface grid resolution on GIA model calculations with elastic bedrock**
 835 **deformation due to instantaneous removal of cylindrical ice loads. Cylinders are all of unit height 100 m, and radius from 0.5 to 16**
km. Five grid resolutions applied within an area of minimum 40km width were tested: 0.5, 1, 1.75 km, 3.75 and 7.5 km (Figure S1).
(a) Transect of bedrock deformation for removal of ice cylinders with unit height and radii of 2 km (solid lines), 5 km (dotted) and
10 km (dashed lines). (b) – (e) Results of a suite of simulations adopting ranges of ice cylinder radii and grid resolutions. (b) Colors
indicate 1 minus the Mass Factor, [1 – Mass Factor], where the Mass Factor is the ratio of the theoretical mass of the load and the
mass of the load represented on the given model grid. 0 represents a scenario where the model grid perfectly represents the mass of
 840 **the idealized load, whilst positive (blue) and negative (red) values indicate the load mass is over- and under-represented by model**
grid resolution, respectively. (c) Root mean square error across the suite of simulations (mm). (d) Average absolute percentage error
(%). (e) Standard deviation of the absolute percentage error (%) of the given test from the finest 0.5 km resolution model result,
calculated within 2 km of the loaded region. Dashed black lines represent the 1:3 ratio between idealized load cylinder radius and
surface grid resolution whereby average absolute percentage error becomes $< 7 \pm 3$ (σ) % for all scenarios.




845

Figure 4. Influence of grid resolution on elastic sea level predictions in ASE. Difference in predicted sea level change in meters between (a-c) 1.9 and 15 km, (d-f) 1.9 and 7.5 km; and (g-i) 1.9 and 3.75 km resolution GIA model simulations with a purely elastic Earth model across the times indicated at the top of the column for ice loading scenarios (from left – right) ICE-SH ICE-GOL and ICE-RD. For each ice retreat scenario there is a different colour bar since the magnitude of error due to grid resolution differs. In


some panels, the colour bar is saturated. The black and blue line indicates final and initial grounding lines, respectively, for each simulation, and each frame is annotated with the maximum and minimum values within the frame.



850 **Figure 5. Evolution of error in elastic sea level predictions due to grid resolution from 1950 to 2100 with the 1km input resolution**
ICE-RD ice model. (a) Box-whisker plots of the error and percent error (see methods) calculated from the difference in predicted
sea level changes from the start of the simulation to the indicated time within 10 km of the grounding line at that time between a
simulation with 1.9 km resolution and simulations adopting 15 km (light pink), 7.5 km (medium pink) and 3.75 km (dark pink) grid
855 **resolutions. The box represents (from left to right) the 25th percentile, median and 75th percentile of the distribution, whilst the**
whiskers represent the "minimum" (25th percentile - 1.5 x the interquartile range) and "maximum (75th percentile - 1.5 x the
interquartile range). Error (m) is the difference between sea-level predictions from the higher - lower resolution run. Percentage
Error (%) is calculated as $100 * (SL_{1.9km} - SL_{lowres}) / SL_{1.9km}$ for each grid point.




860 **Figure 6. Influence of incorporating viscous behaviour and uncertainty in viscoelastic Earth structure on sea level predictions. Frames (a-c) shows the difference in sea level change predicted from simulations adopting 3-D viscoelastic earth model EM1_L and an elastic Earth model. Frames (d-f) shows the difference in sea level change predicted from simulations adopting two different 3-D viscoelastic earth models EM1_L and EM1_M. Note the difference in scale. Time frames and ice models are as indicated at the top of the columns. In each frame, the black and blue line indicates final and initial grounding lines respectively for each simulation and is annotated with the maximum and minimum data value within the frame. All simulations adopt a 1.9 km grid resolution.**




865

Figure 7. Influence of grid resolution on viscoelastic sea level predictions in ASE. As in Figure 4 but adopting 3-D viscoelastic Earth model EMI_L.



870 **Figure 8. Evolution of site-specific sea level in simulations adopting a range of model resolutions and Earth models. (a) ice thickness**
change from 2000 to 2100 predicted in the ICE-RD simulation. This frame is identical to figure 2c. (b) colored lines show predicted
sea level change, in meters, at Site X that experiences the maximum viscous uplift in the 1.9 km resolution simulation, shown by the
blue star in frame (a) in the simulations with a purely elastic solid earth response (black lines), viscoelastic solid earth response based
on a global average 1-D earth model (red lines), a 1-D earth model best-fit for the WAIS from Powell et al. (in review) (purple lines),
875 **a low viscosity 3-D earth model EM1_L (dark green lines), a moderate viscosity 3-D earth model EM1_M (light green lines) and 3-**
D viscosity earth model EM2 (blue lines). Solid lines are for simulations performed at 1.9 km resolution, and dashed lines adopt a
15km resolution. (c) is as in frame (b) but for Site Y along the final grounding line position at 2100 that experiences the greatest ice
thickness change, labeled by the red star in frame (a).



885 **Figure 9. Comparison of factors contributing to differences in sea-level predictions in this study. Each distribution represents the influence of the specific factor across points within 10-km of the grounding line at the specified year of the model run, as a percentage of the total sea-level change at that time for a) ICE-SH ice model from 1992 to 2017; b) ICE-RD ice model from 1950 to 2050; c) ICE-RD ice model from 1950 to 2100. Six factors are compared in this figure, as labelled on the right. To visualise the distribution, we plot a classic box-whisker diagram overlain with a density curve. The edges of the box represents (from left to right) the 25th percentile, median and 75th percentile of the distribution, whilst the whiskers represent the “minimum” (25th percentile – 1.5 x the interquartile range) and “maximum” (75th percentile + 1.5 x the interquartile range). The diamonds outside the whiskers represent outliers.**