

Towards accurate quantification of ice content in permafrost of the Central Andes - Part II: an upscaling strategy of geophysical measurements to the catchment scale at two study sites

Tamara Mathys¹, Christin Hilbich¹, Lukas U. Arenson², Pablo A. Wainstein², and Christian Hauck¹

¹University of Fribourg, Department of Geosciences, Switzerland

²BGC Engineering Inc., Canada

Correspondence: Tamara Mathys (tamara.mathys@unifr.ch)

Abstract. With ongoing climate change, there is a pressing need to better understand how much water is stored as ground ice in areas with extensive permafrost occurrence, and how the regional water balance may alter in response to the potential generation of melt water from permafrost degradation. However, field-based data on permafrost in remote and mountainous areas such as the South-American Andes is scarce and most. Most current ground ice estimates are based on broadly generalised assumptions such as volume-area scaling and mean ground ice content estimates of rock glaciers. In addition, ground ice contents in permafrost areas outside of rock glaciers are usually not considered, resulting in a significant uncertainty regarding the volume of ground ice in the Andes, and its hydrological role. In part I of this contribution, Hilbich et al. (submitted) present an extensive geophysical data set based on Electrical Resistivity Tomography (ERT) and Refraction Seismic Tomography (RST) surveys to detect and quantify ground ice of different landforms and surface types in several study regions in the semi-arid Andes of Chile and Argentina with the aim to contribute to the reduction of this data scarcity. In part II we focus on the development of a methodology strategy for the upscaling of geophysical-based geophysics-based ground ice quantification to an entire catchment to estimate the total ground ice volume (and its estimated approximate water equivalent) in the study areas. In addition to the geophysical data, the upscaling approach is based on a permafrost distribution model and classifications of surface and landform types. Where available, ERT and RST measurements were quantitatively combined to estimate the volumetric ground ice content using petrophysical relationships within the Four Phase Model (Hauck et al., 2011). In addition to introducing our upscaling methodology, In this paper, we introduce our upscaling strategy, and we demonstrate that the estimation of large-scale ground ice volumes can be improved by including (i) non-rock glacier permafrost occurrences, and (ii) field evidence through a large number of geophysical surveys and ground truthing information. The results of our study indicate, that (i) conventional ground ice estimates for rock-glacier dominated catchments without in-situ data may significantly overestimate ground ice contents, and (ii) substantial volumes of ground ice may also be present in catchments where rock glaciers are lacking.

1 Introduction

In many arid and semiarid areas around the world, mountain regions play a significant role for controlling downstream water supply. At high altitudes, runoff is delayed by glaciers, which consequently act as a major source of water for agriculture, power generation, mining and drinking water during the summer (Urrutia and Vuille, 2009; Salzmann et al., 2013; Rangescroft et al., 2015; García et al., 2017; Hoelzle et al., 2019). With continued climate change further enhancing the recession of glaciers globally, their contribution to the summer runoff will eventually decline, whereby the timing of this decline differs globally based on the region (IPCC, 2019). Therefore, water availability, and specifically its timing, will drastically change. In this context, the permafrost distribution, the corresponding ground ice content and its degradation is of increasing importance. ~~Particularly~~ as it is currently debated whether permafrost ground ice can be considered as a significant water reservoir and as an alternative resource of fresh water that could potentially moderate water scarcity during dry seasons in the future (Brenning, 2005; Azócar and Brenning, 2010; Duguay et al., 2015; Hoelzle et al., 2017; Jones et al., 2019; Liaudat et al., 2020). Thus, there is a pressing need to better understand i) how much water is stored as ground ice in areas with extensive permafrost occurrence and ii) how the regional water balance may alter in response to the potential generation of melt water from permafrost degradation.

In the arid and semi-arid Andes, permafrost and specific permafrost landforms such as rock glaciers are known to be well developed and abundant (Azócar and Brenning, 2010; Jones et al., 2018a, 2019; Schaffer et al., 2019; Masiokas et al., 2020). Several recent studies attempt to estimate and model water volumes and the potential hydrological significance of rock glaciers by quantifying their ground ice contents (Azócar and Brenning, 2010; Rangescroft et al., 2015). These studies hypothesise that a significant amount of water may be stored in the ice-rich layers of rock glaciers, ~~however,~~ ~~However,~~ those assertions are typically not supported by in-situ measurements. If present (and future) periglacial protection laws in the South American Andes (e.g. Chile and Argentina) are to be practically adopted, a more precise understanding of the distribution, ground ice content and thermal state of permafrost is necessary. Most of the large-scale studies addressing ground ice volumes in rock glaciers are based on remote sensing data and use empirical rules of thumb to estimate the ground ice content without any ground truthing as validation. Due to the lack of in-situ investigations and subsurface information, commonly a mean volumetric ice content of 40 % to 60 % is assumed for the entire rock glacier area (Brenning, 2005; Rangescroft et al., 2015; Bodin et al., 2010; Azócar and Brenning, 2010; Rangescroft et al., 2015; Jones et al., 2019). This can be problematic as the ground ice content of rock glaciers has been shown to vary considerably from case to case (~~Arenson and Jakob, 2010; Hauck et al., 2011; Mollaret et al., 2020; ?, Hilbich et al., submitted~~) (~~Arenson and Jakob, 2010; Hauck et al.,~~ and also along longitudinal profiles of a single rock glacier (~~Jones et al., 2019; ?~~)(~~Jones et al., 2019; Halla et al., 2021~~). Furthermore, while rock glaciers are the focus of several studies assessing the hydrological importance of permafrost in semi-arid regions, knowledge about the permafrost distribution outside of rock glaciers is still extremely limited (Arenson and Jakob, 2010; Duguay et al., 2015), even though ice-rich ~~ground-ice permafrost~~ has been shown to be present in areas devoid of rock glaciers (García et al., 2017; Schaffer et al., 2019). The water equivalent stored in ice-rich permafrost zones can thus be expected to be significantly higher than calculated from rock glaciers alone (Arenson and Jakob, 2010; García et al., 2017; Baldis and Liaudat, 2020). Clearly, field based studies are needed to fully assess the partitioning of the different permafrost landforms to the

total ground ice volume stored in permafrost (e.g. Schaffer et al., 2019; Croce and Milana, 2002; Azócar and Brenning, 2010; Arenson and
(e.g. Schaffer et al., 2019; Croce and Milana, 2002; Azócar and Brenning, 2010; Arenson and Jakob, 2010; Monnier and Kinnard, 2013; I
. Moreover, the few existing (statistical) permafrost distribution models of the Central Andes (and elsewhere) are commonly
based on the presence and distribution of active rock glaciers and can therefore not easily be extended to other types of surface
60 covers (Arenson and Jakob, 2010; Bodin et al., 2010; Azócar et al., 2017; Esper Angillieri, 2017). This bias is caused by the
inability of available remote sensing data to detect permafrost from space outside of clear geomorphic indicators, such as rock
glaciers.

In order to remove this bias and estimate the total ground ice volume more accurately, Hilbich et al. (submitted) presented
an extensive geophysical data set from several permafrost regions in the Central Andes of Chile and Argentina consisting of
65 Electrical Resistivity Tomography (ERT) and Refraction Seismic Tomography (RST) measurements, which can be used for the
detection of permafrost and the quantification of the ground ice content. In the absence of boreholes, geophysical investigations
are a feasible and cost-effective technique to detect and quantify ground ice occurrences within a variety of landforms and sub-
strates (e.g. Hauck and Kneisel, 2008a; Hilbich et al., 2009, 2011; Monnier and Kinnard, 2013; Mewes et al., 2017; Pellet and Hauck, 201
(e.g. Hauck and Kneisel, 2008a; Hilbich et al., 2009, 2011; Monnier and Kinnard, 2013; Mewes et al., 2017; Pellet et al., 2016; Mollaret e
70 . Several combinations of geophysical measurements have been used in this context. Monnier and Kinnard (2013) used ground-
penetrating radar to estimate the ground ice content in a rock glacier in the Central Andes of Chile. Halla et al. (2020, 2021)
used ERT and RST data in combination with the so-called 4-phase model (Hauck et al. 2011) to calculate the ground ice con-
tent within a rock glacier complex in Argentina. In further studies on rock glaciers in other mountain ranges, electromagnetic
(Bucki et al., 2004), gravimetric (Hausmann et al., 2007) and multitudes of geophysical techniques (Maurer and Hauck, 2007;
75 Buchli et al., 2018) were applied, often in combination with ground truth data from boreholes.

Finally, and in addition to the geophysical surveys and local ground ice calculations themselves, upscaling techniques are
needed to estimate the ground ice content over larger scales than the actual profile lengths of the geophysical surveys, such
as a whole watershed (e.g. Minsley et al., 2012; Hubbard et al., 2013; Dafflon et al., 2017). Upscaling is needed when water
balances and potential future changes on larger spatial scales have to be assessed. In this study we aim to bridge the scale
80 gap between the individual geophysical profiles and the catchment scale by establishing an upscaling methodology strategy,
which is then applied to two study sites in the Central Andes of Chile and Argentina. The geophysical data and analyses are
hereby based on the companion paper in part I (Hilbich et al., submitted). We present first estimates of total ground ice content
contents (and water equivalent equivalents) of a) a rock glacier dominated (Site A) and b) a rock glacier free site (Site D). We
then compare the results of our proposed upscaling methodology strategy to conventional estimates based on remotely sensed
85 data and the empirical rule of thumb established by Brenning (2005).

2 Study Sites

The field data of the two sites were acquired in summers 2018 and 2019 (Southern Hemisphere) in the framework of envi-
ronmental impact assessment studies (EIA) in mining environments. Profile locations were chosen according to the probable

presence of frozen ground, but also according to easy access and safety within the mines. Locations are therefore not always
90 optimally situated with respect to representativity in an upscaling context. ~~The~~ However, the location of geophysical surveys and other field activities were chosen in such a way that they were not located in regions of active mining activities, which could have impacted the study results. Site D is still in an explorative phase, and thus not an active mining site, where impacts are limited to the construction of roads. Site A surrounds an active mining pit. However, we only consider geophysical profile results from areas that are far away from any disturbances, or are only affected by small scale surface disturbances, such as
95 access roads or drilling platforms. Therefore, the local mining context has no further relevance for the scientific content of this paper and is ~~therefore~~ omitted in the following. The mining infrastructure, however, enabled access to high-altitude permafrost environments and made the collection of validation data possible.

2.1 Rock glacier free site - Site D

The rock glacier free site is located between 3800 to 5400 m a.s.l., about 140 km southeast of the city of Copiapò (cf. Site D
100 in Figure 1). The climate of Site D is classified as semi-arid, typical ~~of for the~~ high altitude Central Andes. Precipitation and relative humidity are low, while solar radiation is high. Mean annual air temperature (MAAT) calculated from a meteorological weather station that was installed on site at an altitude of 5012 m a.s.l. in 2015 is -5.4 °C (for the measurement period of 2015 - 2017). Using long-term representative regional climate stations in the vicinity of the site, a MAAT of -7.3 °C was estimated for the period of 1978 to ~~2015 (Devine et al., 2019).~~ 2015. Mean annual precipitation estimated from the same long-term
105 series is 131 mm, with significant variability (annual values ranging from 0 mm to 738 mm) (Devine et al., 2019). Site D is characterized by a uniform landscape consisting of a widespread fine-grained sediment cover. Rock glaciers are absent at this study site ~~and geomorphic~~. Geomorphic surface indicators for permafrost mostly consist of widespread gelifluction lobes and in some cases weakly developed patterned ground. ~~The~~

The main host rock at Site D consists of Late Paleozoic (Permian - Triassic) rhyolites and andesites, which are overlain by
110 Jurassic and Cretaceous sediments and conglomeratic clastic rocks that are strongly silicified and altered. Furthermore, Site D is located on a large hydrothermal alteration zone characterized by high sulphidation epithermal alterations and porphyry alterations. The bedrock is in general highly fractured as a result of the numerous fault systems that cross the area (Devine et al., 2019). The results of the geophysical surveys presented in part I (cf. Hilbich et al., submitted) point to largely homogeneous sub-surface conditions with significant ground ice occurrences (mostly in terms of a thin, ice-rich layer varying in thicknesses of
115 approximately 2-5 m). Furthermore, ground ice is expected to be present as well in the highly fractured and hydrothermally altered bedrock at greater depths ~~(?)~~ (Hilbich and Hauck, 2018).

~~Overview maps of study sites A (rock-glacier dominated site, including various sub-study areas) and D (rock-glacier free site) in the Central Andes of Chile/Argentina. The permafrost zonation index after Gruber (2012) shows the modelled permafrost distribution within the Central Andes. Black lines indicate the position of the geophysical profiles. Map data: Site A: Google, ©2021, Maxar Technologies; Site D: DigitalGlobe, accessed in GlobalMapper V22 (Blue Marble Geographics), date unknown~~
120




Figure 1. Overview maps of study sites A (rock-glacier dominated site, including various sub-study areas A1-A4) and D (rock-glacier free site) in the Central Andes of Chile/Argentina. The permafrost zonation index after Gruber (2012) shows the modelled permafrost distribution within the Central Andes. Black lines indicate the position of the geophysical profiles. Map data: Site A: Google, ©2022, Maxar Technologies; Site D: DigitalGlobe, accessed in GlobalMapper V22 (Blue Marble Geographics), date unknown

2.2 Rock glacier dominated site - Site A

Contrasting Site D, we choose a second field site with an abundance of specific permafrost landforms including rock glaciers and talus slopes. Site A is located in the upper part of the Choapa Valley, about 200 km north of Santiago de Chile (Figure 1) and was previously studied by Monnier and Kinnard (2013). The geophysical measurements are located at an altitudinal range of 3560 to 3850 m a.s.l., whereas the maximum altitude of Site A is around 4300 m a.s.l. The climate of site A is arid-semi-arid with a short rainy season between May and August. The annual precipitation ranges between 200 mm and 800 mm with an average of 334 mm for the period of 1961-90 (Monnier and Kinnard, 2013). MAAT is estimated to be around +0.5 °C in 2010

at an altitude of 3700 m a.s.l., which is a clear indicator that permafrost would not form under current climatic conditions and it is naturally in a degrading state. ~~Site A contains various distinctive landforms.~~ (Monnier and Kinnard, 2013). The most important periglacial landforms present at Site A are rock glaciers with about 15 rock glaciers that have been identified in the area of interest. ~~In general, variable~~ Variable ground ice contents were found in the different rock glaciers and other landforms of Site A (Monnier and Kinnard, 2013, Hilbich et al., submitted.). Areas outside of rock glaciers are most likely unfrozen and free of permafrost, as seen by the relatively low electrical resistivities found in such areas (Hauck et al. 2017, Hilbich et al., submitted), and as confirmed by several test pits.

135 3 Methodology

~~The~~ The geology of Site A is characterized by volcanic (both intrusive and extrusive) as well as sedimentary rocks aging from the Middle Triassic until recent. A large part of the study area has been mapped as quartz-diorites and andesites. Quaternary deposits consist of fluvioglacial sediments and morainic deposits. Similar to Site D, several faults associated with alteration and mineralization are located at Site A (Tapia et al., 2016).

140 3 Methods

The data sets used for this study consist of (i) geophysical measurements (Electrical Resistivity Tomography, ERT, and Refraction Seismic Tomography, RST), which are part of the large data set presented in the companion paper of Hilbich et al. (submitted), (ii) ground truth data in the form of test pits, boreholes and natural outcrops (unpublished data provided by BGC Engineering Inc. (BGC)), and (iii) geomorphological maps produced during fieldwork in the framework of this study. All data are then combined within a workflow for upscaling from the individual geophysical profile to the catchment scale (Figure 2). The upscaling approach is based on the assumption, that the geophysical profiles can be considered representative for larger areas with comparable near-surface substrate or for similar landforms in similar topo-climatic locations, by this representing comparable permafrost occurrences with similar ground ice contents. The workflow consists of five steps (Figure 2): (1) creating a background permafrost distribution model, (2) modelling quantitative ground ice contents for each profile, (3) ~~partitioning of the study site into different surface/subsurface classes,~~ quantitative ground ice content estimations for each profile, upscaling classes, and (4) establishing a conceptual soil stratigraphy model for each upscaling class. In a final step (5), the ground ice content ~~estimations are integrated over estimates are upscaled to~~ estimations are upscaled to the whole study area ~~according to the soil stratigraphy model and the respective abundance of each upscaling class~~ using the upscaling classes as a basis. While steps (3) and (4) are closely related to the geophysical data and corresponding ground ice content calculations (see section 3.1), steps (1) and (2) are more flexible and can be based on different approaches. Any permafrost distribution model may be used to distinguish between permafrost classes and non-permafrost classes (1) and any field information and remote sensing data can be used to classify the surface into upscaling classes (2). In the following, the different steps of the strategy are further

explained, before we show the results for the study site classification and the ice content quantification in the results section (section 4).




Figure 2. Workflow of the upscaling methodology strategy used in this study. Step 1, we create a simple permafrost distribution model to exclude areas with low permafrost probability from further processing. Then, we use geophysical data, and where applicable, model the ground ice content using the 4PM (Step 2). Then, in Step 3, we define upscaling classes using information from different sources such as aspect and slope maps derived from DEMs, field notes and observations. In Step 4, a conceptual soil stratigraphy model is established using the geophysical results as input for subsurface layers and corresponding ice contents for each upscaling class, which serves as the basis for the upscaling of the ice content to the catchment scale (Step 5).

160 3.1 Geophysical datasets and validation

In part I of this study, Hilbich et al. (submitted), provide the details about [geophysical](#) data acquisition and processing as well as data interpretation [and ice content modelling](#) in the context of a comparison of several permafrost field sites in the Central Andes. It has to be noted that the geophysical profiles were not planned specifically for the purpose of upscaling to larger regions. However, the extensive data set allowed us to develop our [methodology-strategy](#) and apply it to the two field sites
165 described above. Several sources of ground truthing were available for the validation of the ~~interpretation of the geophysical surveys~~[geophysical results](#). At Site A, a large set of borehole data exist (Koenig et al., 2019), and the results from the geophysical surveys are in good agreement with the available borehole data (Hilbich et al., submitted; ~~Hauck et al. (2017); ?~~[Hauck et al. \(2017\); Hilbich and Hauck \(2018\)](#)). At Site D, no geotechnical boreholes exist for calibrating the geophysical lines, ~~however,~~ [However](#), several test pits were excavated, which largely confirmed the findings from the geophysical sur-
170 veys. Additionally, several natural outcrops were present in the vicinity of the profiles and further served as validation data (~~Hauck et al., 2017; ?~~). ~~We refer the reader to the companion paper (Hilbich et al., submitted) regarding geophysical data acquisition and processing, as well as the ground ice quantification from co-located ERT and RST tomograms by applying the so-called 4-phase model (4PM, Hauck et al., 2014) (Hauck et al., 2017; Hilbich and Hauck, 2018).~~ [Ground truthing data locations are shown in Figure 3.](#)

175 3.2 Permafrost distribution models

For the sake of simplicity and to use our available ground truthing data effectively, we chose a multiple regression approach that was first introduced by Hoelzle (1994) to generate a permafrost distribution model for our field sites. In contrast to e.g. Hoelzle (1994), who used data from the bottom temperature of the snow cover (~~BTS~~), we use the ~~results of the geophysical measurements~~[geophysical results](#), boreholes and test pits as point input for the response variable Y, indicating likely permafrost
180 occurrence [or absence](#). Elevation and potential incoming solar radiation (PISR) were chosen as the predictor variables (equation 1). [PISR was calculated using ArcGIS Pro's "Area Solar Radiation \(Spatial Analyst\)" tool, which uses a DEM as input and includes the latitude of the study sites for calculations of solar declination and solar position to derive PISR. In total, 50 point indicators for permafrost occurrence were used for the regression models. For Site D, these points correspond to either a location on a geophysical profile or a test pit located at elevations ranging from 4400 m a.s.l. to 5151 m a.s.l. and comprise both permafrost and non-permafrost observations. A standard error of 0.58 and an R² of 0.6 was calculated. For Site A, point indicators correspond to points on the geophysical profiles or data from boreholes ranging from 3200 m a.s.l. to 4000 m a.s.l. A standard error of 0.25 and R² of 0.72 was calculated. These regression statistics resemble the values used for models in other studies \(e.g. Hoelzle \(1994\)\), and are therefore considered to be acceptable.](#)

$$190 \quad y(x) = b_0 + b_1 * elevation(x) + b_2 * PISR(x) \quad (1)$$

[where \$b_{0,1,2}\$ = regression coefficients](#)




Figure 3. Comparison of the permafrost distribution model developed in this study (b and d, DEM 5 m resolution) with the PZI model by Gruber (2012) (a and c, based on SRTM30 data set, <1 km resolution). Locations of boreholes (Site A) and test pits (Site D) are marked by red points on all maps. Map data: DigitalGlobe, accessed in GlobalMapper V22 (Blue Marble Geographics), date unknown

The resulting permafrost distribution model delineates areas where permafrost occurrence is probable, possible or unlikely, analogous to similar approaches in earlier studies (Hoelzle, 1994; Gruber and Hoelzle, 2001; Arenson and Jakob, 2010; Esper Angillieri, 2017; Kenner et al., 2019). The main purpose of the model in the context of this study is to provide a coarse overview of the expected permafrost distribution of the study region to be able to directly exclude those areas from further processing, where permafrost is highly unlikely (cf. yellow-white areas in Figure 1a, 2a and 3-3 b and d). More complex, statistical permafrost distribution models for the Central Andes have been presented (e.g. Bodin et al., 2010; Azócar et al.,

2017), ~~however,~~ However, they cover only parts of our study sites. A comparison of our model approach with the model by Azócar et al. (2017) for Site A and with an unpublished model (L. U. Arenson, personal communication), following the approach presented in Arenson and Jakob (2010) for Site D, showed a good agreement regarding the areas where permafrost is estimated to be absent. A comparison between the permafrost distribution model developed in this study and the global permafrost zonation index (PZI) developed by Gruber (2012) is shown in Figure 3. For Site D, both models predict (high) probability for permafrost occurrence. ~~However, the PZI distinguishes between areas of very high probability to the West of the site and areas with medium probability for permafrost to the North of the site, whereas our model predicts a high probability for permafrost everywhere~~ at the location of the geophysical profiles (black lines in Figure 3). For Site A, the PZI does not cover the entire study site area, probably as a result of the lower altitude which is not taken into account in the PZI. At this site, the PZI predicts a low probability for permafrost occurrence at all subsites. The model developed in this study also delimits large portions of the study site as unlikely for permafrost. Nevertheless, ~~the model it~~ also suggests the possibility of permafrost (light blue areas) for most of the areas, where the geophysical profile lines are located. A higher probability for permafrost is here only modelled for steep and high ~~bedrock slopes~~ altitude (> 3800 m a.s.l) bedrock slopes here. As a consequence of these clear differences between the two sites, the entire area of Site D was considered for the following steps of the strategy, whereas only rock glaciers and talus slopes were considered at Site A.

~~Comparison of the permafrost distribution model developed in this study (a and c, DEM 5 m resolution) with the PZI model by Gruber (2012) (b and d, based on SRTM30 data set, <1 km resolution). Map data: DigitalGlobe, accessed in GlobalMapper V22 (Blue Marble Geographics), date unknown~~

3.3 Study site classification

After excluding areas where permafrost is highly unlikely based on the distribution model, the study sites are subdivided and categorized into different classes, which are hereafter called *upscaling classes*, ~~and which~~ These classes are later used to upscale the estimated ground ice content at from the geophysical profiles to a larger area. Surface characteristics are feasible suitable data sets for upscaling approaches of geophysical data because the classification can be performed over larger areas ~~by~~ using remote sensing data (e.g. Dafflon et al., 2017). In general, we distinguished different upscaling classes based on factors that are known to strongly influence the thermal regime of high mountain permafrost. The main classification criterion is the near-surface substrate type (e.g., coarse blocky, ~~bedrock~~ fine-grained sediment, etc.), as it has been confirmed to be an important factor for the thermal regime and thus permafrost occurrence besides topo-climatic factors. For example, higher ground ice contents have been shown to be present in areas with blocky surface and sub-surface material (e.g. Schneider et al., 2012; Staub, 2015; Gubler et al., 2013). Further, the presence of specific landforms such as active and relict rock glaciers, talus slopes or gelifluction lobes is used as an important geomorphic indicator for permafrost and ice-rich zones, as they have been shown to possess similar ground ice characteristics for a given region (cf. Hilbich et al., submitted). ~~Finally, in order to delimit bedrock areas from areas considered to contain a sediment cover, we use surface slope angles > 30° as a rough indicator for pure bedrock.~~

In addition, a good general understanding of the study site through field observations and any additional field data will strongly support the classification process, especially in heterogeneous mountain terrain. For Site D, a few general observations made on site and through the analysis of the geophysical data were used to define the upscaling classes as well as the choice of the conceptual models and the ground ice content estimations described in the following section. ~~For example At~~
235 ~~Site D,~~ bedrock, as observed in outcrops during the field campaign, is highly fractured and ice-rich ~~and~~; frozen conditions can be assumed for larger depths and were also observed close to the surface (e.g. shallow active layer) in the various outcrops throughout the study site (Hilbich et al., submitted). Likewise, field observations and the ~~results from the geophysics~~ geophysical results have shown that the surface as well as the subsurface conditions are typically homogeneous, ~~and that frozen conditions can~~. Frozen conditions can therefore be assumed for the entire study site.

240 The resulting classes are ~~then~~ further divided into smaller areas with presumably similar ground ice contents, based on the results of the geophysical surveys and considering the exposition (aspect) of the area. ~~The result of this classification is shown for Site D in Figure 6. Table 1 lists the factors taken into consideration for the resulting upscaling classes and subdivisions for Site D as well as the available geophysical ground truthing data. For detailed explanations on the ERT and RST tomograms we refer the reader to ?.~~ A complete explanation of the different steps taken during the classification process is given in section
245 4.1.

~~Definition of the upscaling classes used for Site D:~~

3.4 Quantification of ground ice contents per class

For each upscaling class, ~~as listed in Table 1,~~ a conceptual soil stratigraphy model was developed to further simplify the complex subsurface conditions into layered models with varying ground ice contents. These conceptual stratigraphies represent
250 a simplified version of representative ERT tomograms, ~~and~~. They form the basis for the upscaling of the ground ice content ~~calculation of to~~ the entire study site. Where available, the 4PM results are used for the estimation of the ice content of each layer. ~~As the~~ Where RST data and hence 4PM calculations are ~~based on coincident ERT and RST measurements (cf. Hauck et al. 2011) and RST surveys were not conducted at all ERT profile lines in the study area (cf. Hilbich et al., submitted), not available,~~ estimations are based on an interpretation of the resistivity distribution of the subsurface alone, in combination with
255 ground truthing data, ~~wherever RST data are absent~~. Figure 4 and Figure 5 show examples of such soil stratigraphy models for both study sites and both cases (with and without coincident RST data). The total ground ice content (in %) was then calculated using equation (2), which calculates the sum of the ground ice contents x_i within each of the (total number of n) identified subsurface layers (i):

$$\sum_{i=1}^n X_i = (\text{thickness} * \text{area} * d \times A \times \text{ice content}) \quad (2)$$

260

where d = depth (m), A = area (m²), ice = ice content (%)




Figure 4. Examples of simplified soil stratigraphy models from both study sites (a = Site D, b = Site A). In both cases 4PM results are not available and ground ice contents for the different layers are estimated from the ERT tomogram and based on ground truthing from test pits and outcrops.

A minimum, mean and maximum ground ice content was considered for each layer in order to account for the uncertainties resulting from the approximation of a homogeneous thickness, area and ice content of each layer. For the (ice-rich) layers we use the same minimum, mean and maximum ice content estimates as applied in the so-called zone-of-interest (ZOI) in part I of this study (Hilbich et al., submitted). Ice content estimations for layers that are not considered in detail in part I are mean values




Figure 5. Examples of soil stratigraphy models for both study sites, where the 4PM was applied. a and b): soil stratigraphy models, (c and d): corresponding 4PM results (ERT, RST and ice content) (Hilbich et al., 2018). Here, ground ice content estimations stem from the 4PM results: The ice content values for the ice-rich layers correspond to the results from part I of this study (Hilbich et al., submitted). Ice content estimations for layers that are not considered in detail in part I are mean values from the 4PM results (cf. Hilbich and Hauck (2018); Hilbich et al. (2018))

from the 4PM results (cf. Hilbich and Hauck (2018); Hilbich et al. (2018)). Finally, the water equivalent (w.e.) was calculated for both sites assuming an ice density conversion factor of 0.9 g cm^{-3} (e.g. Paterson, 1994). ~~In order to compare the sites and different landforms directly, the resulting ice contents and water equivalents are in a final step expressed as unit per 1 km^2 .~~

270 ~~Examples of soil stratigraphy models for both study sites (a = Site D, b = Site A), where 4PM results are not available. Here, ground ice contents are estimated for the different layers based on expert knowledge and ground truthing from test pits and outcrops.~~

275 ~~Examples of soil stratigraphy models for both study sites, where the 4PM was applied. a and b): soil stratigraphy models, (c and d): corresponding 4PM results (ERT, RST and ice content) (Hilbich et al., 2018). Here, ground ice content estimations stem from the 4PM results: The ice content values for the ice-rich layers correspond to the results from part I of this study (Hilbich et al., submitted). Ice content estimations for layers that are not considered in detail in part I are mean values from the 4PM results (cf. Hilbich et al. (2018)).~~

4 Results

280 The methodology approach was applied to both study sites, in order to test it for two cases that differ significantly in their surface and subsurface characteristics and corresponding permafrost distribution and conditions. As can be seen from the permafrost distribution models (Figure 3), Site D is located in a zone with high permafrost probability and significant potential for the occurrence of ground ice (Figure 3, ~~1a;~~ a, b), despite the absence of ice-rich landforms such as rock glaciers(?). On the contrary, permafrost ground ice is limited to rock glaciers and possibly talus slopes at Site A (Hilbich et al., 2018) (cf. Figure 3, ~~2a,b;~~ c,d).

285 For both sites, we calculated total volumetric ground ice estimates (section 4.3.4.2) according to two different scenarios (S) with different assumptions regarding the spatial extent of ice-bearing layers for each site. For Site D, the two scenarios differ with respect to their investigation depths:

(S1)

290 (SD1) Calculation of ground ice content contained in the uppermost 10 meters. This scenario gives a relatively good estimate for the ground ice content in the near-surface layer of the study site, where the reliability of the geophysical surveys is ~~still high.~~

(S2) high.

295 (SD2) Calculation of the ground ice content down to a maximum depth of 30 m. Obviously, in areas with potentially deep permafrost occurrences, the total estimated ground ice content volume depends on the depth considered. This second scenario considers a larger investigation depth, which corresponds to three times the extent of scenario ~~S1~~ SD1 and may be more relevant for modelling purposes in the context of permafrost degradation and its impact on the water balance. Frozen ground conditions to even larger depths can be assumed for Site D, however, uncertainties regarding subsurface conditions (porosity, ice content, etc.) at larger depths are high.

For the rock glacier Site A, where permafrost is limited to rock glaciers and possibly talus slopes, the following scenarios were considered:

300 ~~(S3)~~

(SA1) Calculating the ice content of each sub-catchment by only considering the identified rock glaciers.

~~(S4) As S3~~

(SA2) As SA1, but including talus slopes as additional potentially ice-rich landforms, due to their coarse-blocky surface layer.

In the following sections, we will illustrate the individual steps of the upscaling ~~methodology in detail only~~ strategy in
305 detail mainly for Site D. However, the fundamental differences in ground ice distribution of the sites result in slightly different upscaling approaches for Site A and D. Therefore, we will also shortly explain the most important assumptions made for the ground ice content quantification of Site A.

4.1 ~~Permafrost distribution model~~

~~The permafrost distribution model developed in the framework of this study mainly served to identify likely unfrozen areas
310 that can be excluded from the overall area for the upscaling (Figure 3). In total, 50 point indicators for permafrost occurrence were used for the regression models. These points correspond to either a location on a geophysical profile or a test pit located at elevations ranging from 4400 m a.s.l. to 5151 m a.s.l. and comprise both permafrost and non-permafrost observations. A standard error of 0.58 and an R-squared of 0.6 was calculated. These regression statistics resemble the values used for models in other studies (e.g. Hoelzle (1994)), and are therefore considered to be acceptable. The model indicates a high probability
315 of permafrost for the entire area of Site D, but less probable permafrost conditions at Site A. This result is very similar to the results for the Permafrost Zonation Index (Gruber 2012) (shown in Figure 3 b and d) and an unpublished model of the region (L. U. Arenson, personal communication), which estimates that more than 90 % of Site D is underlain by permafrost. As a consequence of these clear differences between the two sites, the entire area of Site D was considered for the following steps of the methodology, whereas only rock glaciers and talus slopes were considered at Site A.~~

320 4.1 Classification into upscaling classes

The upscaling classes determined for Site D are listed in Table 1, together with their specific surface characteristics, permafrost probability, aspect and the tomogram ~~that is representative of~~ representative for each upscaling class. These ERT tomograms can be found in the Appendix (Figure A1). ~~Further,~~ For a more detailed discussion of the individual ERT and RST tomograms see Hilbich et al., submitted, and Hilbich and Hauck (2018).

325 For the classification, Site D was split into sub-catchment ~~1-D1~~ 1-D1 to the west of the main ridge, and sub-catchment ~~2-located on D2 located to~~ 2-located on D2 located to the east (Figure ~~1a and Figure 36~~). For the classification, the surface types and landforms were mapped manually based on satellite images and geomorphological mapping during the field ~~campaign~~ campaigns and using a DEM to delimit

Upscaling class		landform, substrate/ Geomorphology	Permafrost probability	Aspect	Other Characteristics	Tomogram
1	a	Gelifluction lobes	High	N / NW	Fine-grained sediments	D01
	b	Fine sediments, less distinctive gelifluction lobes	High	S / SW	Fine-grained sediments	D02
2	a	Fine Sediments, no special geomorphology	High	SE	High altitude	D03 & D04
	b	Fine Sediments, no special geomorphology	High	SE	All tomograms in this region contain a conductive layer	D05, D06, D07 & D08
	c	Fine sediments, gelifluction lobes	High	N	Ice-poor bedrock layer beneath sediments	Part of D07
3	a	Bedrock (highly-fractured) with shallow sediment cover	High	W / NW	Slope >20° and <30° Ice-rich +/- 2m sediment cover	D09
	b	Bedrock (highly fractured) with shallow sediment cover	High	N or E	Slope >20° and <30° Ice poor	Based on parts of D07
4	a	Bedrock	High	W / NW	Ice-rich	No tomogram
	b	Bedrock	High	N	Ice-poor	No tomogram

Table 1. Definition of the upscaling classes used for Site D.

bedrock areas based on a slope threshold. At Site D ~~they~~ surface types and landforms include fine-grained (colluvial) sediments, gelifluction lobes, bedrock with a sediment veneer (defined as approximately <2 m sediment covering the bedrock), and pure bedrock. Sub-catchment 1-D1 is characterized by a widespread fine-grained sediment cover, ~~which~~. The sub-catchment is further divided into a) the northwestern slope ~~of the sub-catchment~~ with fine-grained sediment with and gelifluction lobes (upscaling class 1a, with D01 serving as reference tomogram), and b) the southwestern slope dominated by fine-grained sediment without gelifluction lobes (upscaling class 1b, with D02 serving as reference tomogram). Profile D09 serves as a reference in sub-catchment 1-reference for slopes steeper than 20° which were classified as bedrock with a thin sediment cover (sediment veneer) ~~;-based on field observations and~~ defined as upscaling class 3a. Hereby, a thin sediment layer of 1.5 - 2 m above a highly fractured, likely ice-rich bedrock was observed at an outcrop close to profile D09. ~~A small area on the south-east of the site~~ The area of sub-catchment D1 with slope angles >30°, is mapped as upscaling-class 4a corresponding to pure bedrock (slope >upscaling class 4a). This threshold is based on field observations and mapping, where an absence of significant sediment covers were observed in areas steeper than 30°). It is assumed to be ice-rich in the uppermost layer and ice-poor at larger depths, further considering the results of profile D09.

A large central area of sub-catchment 2-D2 consists of colluvial, fine-grained sediment that does not contain any special specific landform. Profiles D03, D04, D05, D06, D07 and D08 are all located in this area (Figure 6). The results from the geophysical surveys point to differences between the profiles located at lower elevations (in the center of ~~the~~ sub-catchment D2) and profiles located higher up, in steeper slopes. Therefore, this colluvial, fine-grained sediment area was further sub-

345 divided into upscaling classes 2a and 2b. Zone 2a contains the profiles D05, D06, D07 and D08, which are all characterized by a distinctive conductive layer whose thickness is probably overestimated by the inversion process (cf. [Part I of this contribution, Hilbich et al., submitted](#)). Zone 2b hosts the profiles D03 and D04, which do not contain said conductive layer. At the foot of the local mountain peak (~~Zones 4a and 4b) gelifluction lobes and sediments (darker color) representing gelifluction morphologies~~ [gelifluction lobes](#) were observed during the field work ([Zone 2c](#)). Profile D07, which reaches the lower boundary of these

350 gelifluction lobes, may point to less ice-rich bedrock conditions compared to what has been found in D09 in sub-catchment ~~4-D1~~ [4-D1](#). Thus, the right-hand part of D07 is taken as a reference for the ground ice calculations for this area, defined as Zone 2c.

Similar to sub-catchment ~~4-D1~~ [4-D1](#), slopes steeper than 20° were mapped as bedrock with a shallow sediment cover. This concerns [a small section \(Zone 3b₂\) next to Zone 2b, but](#) mostly the northern slope ~~to the east of the local mountain peak~~ [to the east of the local mountain peak](#). Because of its northern aspect, and resulting higher exposure to solar radiation, this area ([Zone 3b₁](#)) is assumed to be less ice-rich than

355 its ~~southmost western / south-western-northwestern facing~~ [southmost western / south-western-northwestern facing](#) counterpart in sub-catchment ~~4-D1~~ [4-D1](#). The steepest slopes (> 30°) located at the top of ~~peak on the peak in~~ [peak on the peak in](#) the south-east were mapped as pure bedrock (Zone 4b). Following the argumentation for Zone 3b, this area is also considered to be less ice-rich than the pure bedrock of sub-catchment ~~4-D1~~ [4-D1](#).

The geomorphological mapping and classification for Site A ([not shown in the scope of this paper](#)) yielded areas of fine-grained sediments with slopes < 20° (Class 1), bedrock with a shallow sediment cover (Class 2) (slope \leq 20° and < 30°),

360 and pure bedrock (Class 3) (slopes > 30°), in addition to the mapped rock ~~glacier and talus slope areas glaciers and talus slopes~~ [glacier and talus slope areas glaciers and talus slopes](#) for each sub-catchment (cf. [Figure 1](#)). However, as the permafrost distribution modelling showed that permafrost can be assumed to be absent outside of rock glaciers and talus slopes at Site A, the resulting upscaling classes containing ice only include rock glaciers (scenario ~~S3SA1~~ [S3SA1](#)), or rock glaciers and talus slopes (~~S4~~ [S4](#)) (~~not shown~~ [scenario SA2](#)). Each rock glacier for which the ice content was modelled using the 4PM (cf. [Hilbich et al., submitted](#)) is considered an own upscaling class

365 (Class 4x) with a specific subsurface soil stratigraphy and ice content, accounting for their heterogeneity [which was observed through the geophysical surveys](#). Rock glacier areas ~~, where no geophysical measurements are available, without geophysical measurements~~ were first categorized into intact (Class 5, potentially ice-rich) or relict (Class 6, unfrozen), using the rock glacier classification made by Azócar et al. (2017). Talus slope areas were divided into two upscaling classes, differing between north- (Class 7) or south-facing slopes (Class 8), to account for possibly lower ice contents in north-facing slopes.

370 [The classification process can be summarized by the following steps:](#)

1. [Identifying and isolating areas \(polygons\) with slopes > 30 ° using a DEM \(5 m resolution\) to map bedrock areas, as well as slopes with > 20 ° and < 30 ° to map bedrock with a sediment veneer.](#)
 2. [Mapping of any other surface types and landforms observed in the field and on satellite pictures.](#)
 3. [Checking the geophysical results for the different surface types and landforms, if available. If more than one geophysical profile is present for one surface type class in a sub-catchment \(e.g. the fine-grained sediment area in the center of sub-catchment D2\), the profiles are checked for whether they suggest homogeneous subsurface conditions for the surface type class, or not.](#)
- 375




Figure 6. Map showing the different upscaling class areas for Site D (c). The photographs (d.) illustrate the general surface types /conditions for some of the most important upscaling classes - Classes 1a (highly fractured bedrock (3a)), 1b (widespread uniform fine-grained sediment (2a, 2b)) and 3a (gelifluction lobes (1a)). Also shown are situated in sub-catchment 1 - the slope (a.) and aspect (b.) maps used for the study site classification. Map data: DigitalGlobe, accessed in GlobalMapper V22 (Blue Marble Geographics), date unknown.

380

4. If the profiles do not suggest homogeneous subsurface conditions, the area is further subdivided into smaller upscaling classes. For this, any additional available information that can be used to explain the differences seen in the geophysical results can be used.
5. If no geophysical profile is present in a delimited area (e.g. upscaling class 3b_{1,2} in sub-catchment D2), the information for the subsurface conditions is taken from a profile with similar surface conditions (surface type, altitude,...). Here, we

again rely strongly on field observations and expert knowledge in order to choose the most representative profile for the definition of the subsurface stratigraphy and ice contents.

385 6. Finally, a soil stratigraphy model is established for each upscaling class based on the representative geophysical profile chosen in points 2-4.

4.2 Ground ice content quantification

The ground ice content was quantified for each upscaling class based on the conceptual soil stratigraphy models (cf. Figure 4, Figure 5). ~~In cases with insufficient~~ To do this, Site A was partitioned into smaller sub-catchments as a result of its larger
390 size in comparison to Site D. The partitioning into A1, A2 and A3/4 is a function of the spatial "clustering" of geophysical profiles in Site A. Site D was not further subdivided for the calculation of the total ground ice contents and water equivalents as the total area is comparatively small and because the geophysical profiles are located in closer proximity. In cases where (i) geophysical profiles are not available, (ii) the investigation depth of the geophysical surveys is insufficient, or where certain layer boundaries could not be resolved, additional assumptions had to be made for individual layers of the upscaling classes.
395 Table 2 summarizes the generalized layer structure (active layer, ice-rich sediment layer and ~~the~~ bedrock layer) including maximum thicknesses and mean ice contents for such cases at Site D.

Layer	Max. depth (m)	Ice content (%)
Active layer	1	0 % (per definition)
Sediment	Depending on ERT results	Ice-rich (= max. 85%), or ice-poor (= max. 20%), depending on interpretation of resistivity measurements (ERT)
Highly fractured bedrock	20	Ice-rich (= max. 12%) or ice-poor (= max. 6 %), depending on interpretation of resistivity measurements (ERT)
(more) Consolidated bedrock	30	Ice-poor (= max. 6 %)

Table 2. General assumptions of layers used in the conceptual soil stratigraphy models for the ground ice content calculations for Site D, where (i) no 4PM results exist, (ii) ground truthing data is absent and/or (iii) the investigation depth from the geophysical methods is not sufficient.

For Site A, ground ice contents for the ice-rich core of rock glaciers (Class 4x) were taken from the 4PM results (~~cf. where available, as published in~~ Hilbich et al., submitted), ~~where available~~. The bedrock below ~~this the~~ frozen layer is assumed to be unfrozen in the entire study area. Where no geophysical measurements are available for ice content estimations, the ice content
400 of the ice-rich core of intact rock glaciers (Class 5) ~~are is~~ derived using average values ~~from Site A~~ for rock glaciers that are situated in similar settings (aspect, altitude, permafrost probability) at Site A. Relict rock glaciers (Class 6) are considered to not contain ground ice anymore. For scenario ~~S4SA2~~, talus slopes were furthermore considered to contain possible ice-bearing layers (~~(Gude et al., 2003a; Hauck and Kneisel, 2008b; Mollaret et al., 2020)~~)(Gude et al., 2003b; Hauck and Kneisel, 2008b; Mollaret et al.,
. Here, we assume lower ice contents for north-facing talus slopes (Class 7) ranging from 0 % to 10 % than for south-facing
405 slopes (Class 8), where we applied an ice content range of 5 % to 20 %. These assumptions stem from the hypothesis that talus




Figure 7. Assumed min, mean and max ice content stratigraphies for all upscaling classes (x-axis = ice content (%), y-axis = depth (m)), based on the geophysical results, and calculated ground ice content for each upscaling class. The red shaded parts of the plots indicate where assumptions were made based on Table 2. The resulting min, mean and max water contents for each class are shown in m.w.e. in the boxplots. The map shows the m.w.e. distribution for site D.

slopes **could potentially store higher** can potentially store significant ice volumes as a result of their blocky (insulating) surface layer that allows for more effective cooling (e.g. Wicky and Hauck, 2017). However, the geophysical and 4PM results on talus slopes presented in Part I of this study (Hilbich et al., submitted) are not conclusive in terms of ice content. The numbers

presented in this part of the study should, therefore, be viewed with care. More research is clearly needed to better estimate the
 410 ice content in talus slopes, as the current data coverage is very scarce.




Figure 8. Calculated minimum/maximum water equivalents (m w.e.) for the different scenarios (SD1 - SA2), as well as for rock glaciers and talus slopes at Site A only. For SD1 and SD2, the black bars indicate the uncertainty resulting from using different delineation scenarios for the upscaling classes (see Discussion)

Figure 7 shows the ice content percentage profiles of the soil stratigraphies resulting from Step 4 of our workflow (see Figure 2) until a maximum depth of 30 m together together with the corresponding calculated water equivalents (min, mean and max values) for each upscaling class for Site D. It also shows a map that indicates the distribution of m w.e. in the study site. The calculated total ground ice contents per upscaling class were then summed up for the entire considered area and converted into water equivalent per 1 km² (equivalents (m w.e.) (Figure 8). The total ground ice volumes on the catchment scale calculated for Site D (total area of 4.02 km²) range between 0.0037-0.0095-0.0033-0.0085 km³ (=4.11-10.57 m w.e.) for an investigation depth of 10 m (scenario S1) and 0.0065-0.016-SD1) and 0.0059-0.015 km³ (=7.3-18.4 m w.e.) for an investigation depth of 30 m (scenario S2SD2), respectively. These results equal to a water equivalent per km² ranging from 1.02-2.61 ranging from 0.83-2.13 m w.e. for a depth of 10 m, and 1.81-4.57-1.47-3.71 m w.e. for the 30 m investigation depth. For Site A, the total

420 ground ice volumes ~~on the catchment scale (with (for~~ a total area of 15.32 km²) range between 0.0010-0.0047 km³ ~~(=1.11-5.2~~
~~m w.e) for scenario 3 for scenario SA1~~ and 0.0013-0.006 km³ ~~(=1.46-6.62 for scenario SA2. This equals a water equivalent~~
~~of 0.06-0.27 m w.e) for scenario 4. Here, the water equivalent per km² amounts to 0.10-0.43. for SA1, and 0.084-0.39 m w.e;~~
~~considering the entire study area. for SA2~~ (Figure 8). Values for the Site A sub-catchments are in a similar range. Therefore,
425 non-rock glacier catchments may also contain substantial volumes of ground ice. It is, however, important to note that this
ground ice does not necessarily influence the water balance and probably only marginally influences the annual watershed
hydrology (see ~~discussion in section 5~~conclusion in section 6).

~~Calculated minimum/maximum water equivalents (m w.e) per 1 km² for the different scenarios (S1-S4), as well as for~~
~~rock glaciers and talus slopes at Site A only. Site A was partitioned into smaller sub-catchments as a result of its larger size~~
430 ~~in comparison to Site D. The partitioning into Sub 1, Sub 2 and Sub3/4 is a function of the spatial "clustering" of geophysical~~
~~profiles in Site A. Site D was not further subdivided for the calculation of the total ground ice contents/water equivalents as the~~
~~total area is comparatively small and because the geophysical profiles are located in closer proximity. For S1 and S2, the black~~
~~bars indicate the uncertainty resulting from using different delineation scenarios for the upscaling classes (see Discussion)~~

~~To our knowledge, permafrost ground ice outside of rock glaciers has not yet been quantified or documented in discussions~~
435 ~~related to permafrost in the Andes before. With the upscaling methodology established in the framework of this study, ground~~
~~ice content estimations are now possible also outside of rock glacier dominated catchments, and the relative importance of~~
~~different permafrost occurrences for total ground ice volumes can be evaluated. For Site D with a total area of 4.02 km², the~~
~~ground ice volume estimates are 0.0037-0.0095 km³ for an investigation depth of 10 m (scenario S1) and 0.0065-0.016 km³~~
~~for an investigation depth of 30 m (scenario S2), which convert to a water equivalent of 4.1-10.5 m w.e. and 7.3-18.4 m w.e.~~
440 ~~for the entire study area. The uncertainty range results from the minimum, and maximum ground ice content estimates per soil~~
~~layer. These results~~ Our results point to substantial ground ice occurrences outside of rock glaciers. In direct comparison to the
rock glacier area of Site A, the total ~~water equivalents ice contents~~ calculated for Site D contain only ~~7-17% and 12-30% (for~~
~~10 and 30- 20 % (for the 10 m investigation depth scenario) of the water equivalent ground ice content~~ stored in rock glaciers
at Site A ~~(containing a maximum of 9.0 m w.e)~~. However, on the catchment scale, when ~~the total water equivalent of the two~~
445 ~~catchments is compared relative to the total area per km² comparing water equivalents between Site A and Site D,~~ the values
for Site D are several times larger (Figure 8). Based on ~~ground ice volume estimates contained in 1 km² of water equivalents~~
~~contained~~ the defined catchment areas, Site A ~~(for scenario SA1)~~ contains only about ~~107 - 12 %~~ of what is stored at Site D for
the ~~30 m depth scenario and about 18% for the 10 m depth scenario~~. This is considered reasonable, as most of the catchment
area of Site A (outside of rock glaciers) is expected to be unfrozen (permafrost-free).

450 5 Discussion

With the upscaling strategy established in the framework of this study, ground ice content estimations are now also possible
outside of rock glacier dominated catchments, and the relative importance of different permafrost occurrences for total ground

ice volumes can be evaluated. It is acknowledged, that the absolute numbers of our results should be considered with care, as there are many ~~different~~ uncertainties involved in the various steps of our upscaling ~~methodology: strategy.~~ These uncertainties are discussed below:

455 (i) The estimation of the ice content of the different layers in the conceptual soil stratigraphy models for each site is challenging, as the prescription of a realistic porosity model is a known weakness of the geophysics-based 4-phase model (4PM) (~~Pellet and Hauck, 2017; ?~~)(Mewes et al., 2017; Pellet et al., 2016). This has been addressed with the development of the so-called petrophysical joint inversion (PJI) scheme that is able to invert for porosity in addition to ice and
460 water content (Wagner et al. 2019, Mollaret et al. 2020). However, the successful application of the PJI to a large number of geologically and geomorphologically different profiles is still difficult due to convergence problems in the absence of a priori knowledge (see part I, Hilbich et al., submitted). For the calculations of this study, the porosity models were adjusted for each profile based on a) the landform and the surface substrate, b) the interpreted geophysical results, and
465 c) ground truthing information, where available. We assume porosity values around 50 % for the uppermost sediment layers at the sediment sites, and around 40 % for ~~a surface covered with~~ highly fractured bedrock (with a decreasing gradient for more consolidated bedrock that is expected underneath). For the rock glaciers of Site A, porosity values > 60 % were prescribed in case of highly resistive layers to allow for supersaturated conditions (cf. Hilbich et al., submitted). As these values are merely (field-based) assumptions, porosity estimates could be a ~~large~~ source of uncertainty by leading to either over- or underestimation of the ground ice contents.

470 (ii) Although being capable of estimating ground ice content from the geophysical results (~~e.g. Pellet et al., 2016; Halla et al., 2018; Mollaret et al., 2019; de Pasquale et al., 2020~~)(e.g. Pellet et al., 2016; Halla et al., 2021; Mollaret et al., 2019; de Pasquale et al., 2020), the success of the 4PM application depends on the available data. The identification of representative ground ice content of the different layers is therefore often not possible from the geophysical results alone. In addition, the 4PM requires co-located ERT and RST measurements, which are not always available in a sufficiently large number to be used in an upscaling approach as
475 presented in our study.

(iii) The conceptual soil stratigraphy models used here assume a constant thickness and ice content for the entire upscaling class (under the assumption of relatively homogeneous conditions at Site D). However, lateral differences are to be expected, and the derived ground ice content estimates can only serve as rough approximations on the catchment scale.

~~The main uncertainties of the ice content estimation result from a) the assumptions for min/max ice contents (partly related to uncertainties of the 4PM), and b) from the delineation of the upscaling classes. Especially in catchments, where landforms with clear morphological outlines are missing,~~

480 (iv) There are also various uncertainties with regard to the ~~latter may cause substantial uncertainties~~ classification of the study area into upscaling classes, especially where clear landforms (e.g. rock glaciers, talus slopes) are absent, as is the case for Site D. The classification process is of rather qualitative nature and strongly relies on field observations and expert knowledge. The large potential uncertainties associated with the classification step are indicated by the uncertainty

485

bars shown in Figure 8. These large error bars origin from the uncertainties regarding the spatial extent of the upscaling classes and corresponding subsurface ice occurrences and their maximal/minimal values. To assess the sensitivity of the calculated ground ice contents to the delineation of the upscaling classes, different possible delineation-classification scenarios were compared for Site D. The scenarios were established by either using different tomograms as reference for the ice content of an upscaling class, or by assuming more/less sub-classes (i.e. combining similar classes to larger upscaling classes, or subdividing upscaling classes to smaller ones). The resulting ranges of the ground ice contents calculated for each scenario reach from 14 % to 28 % (shown by black bars for S1 and S2 SD1 and SD2 in Figure 8), with lower uncertainty ranges for the 10 m investigation depth. Wherever The uncertainty range shown in this Figure represents the upper bound. In a purely applied study, one would narrow down this uncertainty range by choosing a best guess scenario for spatial extension, depth of layers and ice content ranges.

To resume, the main uncertainties of the ice content estimation result from a) the assumptions for min/max ice contents (partly related to uncertainties of the 4PM), and b) from the classification of the upscaling classes. Especially in catchments where landforms with clear morphological outlines are missing, the latter may cause substantial uncertainties regarding the spatial extension of the subsurface ice occurrences and their maximal or minimal ice content values. Nevertheless, wherever geophysical data are available in combination with the observations made during the field campaigns, we can be rather confident about the results for Site D for an investigation depth of 10 m (cf. comparison with ground truthing data in part I (Hilbich et al., submitted)), whereas the 30 m depth scenario should be considered with more caution as it is close to the limit of the penetration depth of the available geophysical surveys.

~~Even if the uncertainties remain considerable, the general permafrost distribution and the intra- and inter-landform heterogeneity of the considered catchments is much better captured by geophysical profiles than from remote sensing based approaches alone. This is especially important for catchments that do not contain any clear geomorphic permafrost indicators, such as rock glaciers, that could be delineated using remote sensing data. The upscaling methodology presented in this paper is a first attempt at estimating ground ice contents on a larger scale than just landforms (e.g. rock glaciers or talus slopes). Based on our findings, further studies should focus on improving the knowledge of the ground ice content distribution in different landforms aside from rock glaciers. Furthermore, the upscaling methodology could be improved by e.g. coupling the geophysical results to a spatially distributed ground thermal model or using machine learning image classification schemes that allow to estimate ground ice contents of larger areas by classifying landforms into potentially ice-rich or ice-poor classes.~~

~~Most studies published~~ To our knowledge, permafrost ground ice outside of rock glaciers has not yet been quantified or documented in discussions related to permafrost in the Andes before. Most published studies on estimating the hydrological importance of rock glaciers in the Andes base their estimations of the thickness of the ice-rich layer in rock glaciers on an empirical rule proposed by Brenning (2005) (e.g. Esper Angillieri, 2009; Rangecroft et al., 2015; Jones et al., 2018a, 2019). This empirical relation is based on rock glacier geometry obtained through geomorphological mapping and relates the thickness of the ice-rich layer to the rock glacier area (Brenning, 2005), without explicitly taking into account the potentially complex spatial variability and dependence on rock glacier kinematics and morphology. As shown by the extensive field-based data presented in the companion paper by Hilbich et al. (submitted), the resulting estimates of the thickness of the ice-rich layer

of active rock glaciers is in most cases overestimated when compared to what has been estimated from geophysical surveys and boreholes. For some rock glaciers the empirical rule based estimates of rock glacier thickness is twice as large as what has been found through the geophysical measurements, where the thickness of a potentially ice-rich core can be delimited with high confidence due to the large resistivity contrasts between the different layers. In consequence, area-thickness based calculations of ground ice volumes are significantly higher than calculations ~~that are~~ based on geophysics. The additional assumption of a general volumetric ice content of 50 % commonly used certainly adds to the overestimation of the ice volume contained in certain rock glaciers. Hilbich et al. (submitted) found that such previous ground ice content assumptions for rock glaciers correspond to field-based results only when their ice-rich zone alone is considered.




Figure 9. Comparison of water equivalent (m w.e.) estimations for Site A, scenario SA1 (considering only rock glaciers as ice-bearing class), for (i) the geophysically-based upscaling technique (blue bars) presented in this paper and (ii) the empirical rule (red bars) introduced by Brenning (2005).

Figure 9 compares the results of the water equivalents for Site A, scenario ~~S3~~ SA1 (only considering rock glaciers), using our field-based upscaling ~~methodology with the results of the approach with the~~ total water equivalents calculated using the empirical approach by Brenning (2005), both, per sub-catchment and for the total study site area. The geophysically based results are ~~again given in a range calculated from given as~~ minimum, mean and maximum ice content ~~estimations~~ per sub-surface layer, whereas ~~for the approach of Brenning (2005),~~ we used minimum, mean and maximum ice content assumptions

of 20%, 50% and 60% ,~~respectively, while using the empirical rule for the~~ in addition to the empirical thickness estimation
535 of the ~~thickness of the~~ ice-rich layer to illustrate the approach of Brenning (2005). As can be seen from Figure 9, the mean
water equivalent calculated using the empirical rule is much larger in ~~each case all cases~~ compared to the ~~calculations using the~~
results using the geophysics-based upscaling technique of this study, ~~which is based on the geophysical results~~. Although ac-
knowledging the fact that simplified empirical rules ~~as the one shown above~~ can be useful tools to cover large and remote areas,
they clearly ~~over-generalizes over-generalize~~ the complex and heterogeneous subsurface conditions and ground ice contents
540 of rock glaciers, ~~and as they do not recognise natural variability and actual periglacial processes, they should, in general, be~~
~~improved or complemented with~~. We believe that using field-based evidence, evidence (such as geophysical investigations) in
combination with large-scale remote sensing data ultimately leads to more realistic ground ice content quantification compared
to purely remote sensing approaches. For Site A, we can be confident that ground ice quantification based on the large number
of geophysical profiles carried out on the various rock glaciers are more realistic than the pure remote sensing estimates.
545 Furthermore, with the geophysical data presented in Part I of this study (Hilbich et al., submitted) it becomes clear, that rock
glacier ice contents may vary significantly from the commonly used 40 - 60 %. This is why our upscaling approach strongly
relies on in-situ data although it means that it is not directly transferable to other catchments where no geophysical data exist.
Nevertheless, we believe that field-based data (geophysical or other) are in any case essential for validating remote sensing
products or model-based estimates on ground ice contents.

550 Comparison of water equivalent (m w.e per 1 km²) estimations for Site A, scenario S3 (considering only rock glaciers as
ice-bearing class), for (i) the geophysically-based upscaling technique (blue bars) presented in this paper and (ii) the empirical
rule (red bars) introduced by Brenning (2005):

Furthermore, our results suggest that also non-rock glacier catchments may contain substantial volumes of ground ice. It
is, however, important to note that this water, stored in the ground, does not necessarily influence the annual variability in
555 ~~the water balance, as the presence or absence of a rock glacier in a watershed may only marginally influences the annual~~
~~watershed hydrology. Other factors~~ Even if the uncertainties remain considerable, the general permafrost distribution and
the intra- and inter-landform heterogeneity of the considered catchments is much better captured by geophysical profiles
than from remote sensing based approaches alone. This is especially important for catchments that do not contain any clear
geomorphic permafrost indicators, such as ~~the overall permafrost characteristics, terrain types and precipitation pattern, play~~
560 ~~more significant roles~~ rock glaciers, that could be delineated using remote sensing data.

6 Conclusion

In this paper we presented an upscaling methodology strategy for geophysical measurements with the aim to characterize the
subsurface conditions in high mountain areas with regard to the occurrence of permafrost. Further, the potential ground ice
content for the entire area of two study sites was calculated. The study is based on a series of geophysical measurements (ERT
565 and RST) that were carried out at several study sites in the dry Andes of Chile and Argentina (Central Andes), presented in
Hilbich et al. (submitted), of which we apply our methodology approach to one rock glacier dominated and a rock glacier

free site. The upscaling ~~methodology~~method is based on (a) a simple statistical approach that follows previous work on permafrost distribution models, using the geophysical data as permafrost evidence, and (b) a linkage between geophysical results, substrate type and surface geomorphology to define and delimit areas with presumably similar subsurface conditions (active layer thickness, ground ice content). The so-defined areas were ~~defined~~used as upscaling classes and further used to calculate total ground ice volumes for a larger area. The general approach and strong linkage to the extensive geophysical data sets allowed the estimation of ground ice volumes on the catchment scale. An important advantage of our approach is that it is not restricted to rock glaciers, as in most studies present to date (Azócar and Brenning, 2010; Rangescroft et al., 2015; Schaffer et al., 2019; Jones et al., 2018b; Croce and Milana, 2002). We demonstrate, that the estimation of large-scale ground ice volumes can be improved by including (i) non-rock glacier permafrost occurrences, and (ii) field evidence through a large number of geophysical surveys and ground truthing information (where available). The upscaling strategy presented in this paper is a first attempt at estimating ground ice contents on a larger scale than just landforms (e.g. rock glaciers or talus slopes). Based on our findings, further studies should focus on improving the knowledge of the ground ice content distribution in different landforms aside from rock glaciers. Furthermore, the upscaling approach could be improved by e.g. coupling the geophysical results to a spatially distributed ground thermal model or using machine learning to exploit relationships between surface and subsurface parameters. Our study presents one of the first estimations of the ground ice volume and corresponding water equivalents contained in ice-rich permafrost areas in rock-glacier free catchments. It has been shown that such areas can contain a substantial volume of ground ice that should not be neglected in cryospheric, hydrologic, climate or environmental studies. ~~However,~~It is, however, important to note that the water, stored in the ground ice, does not necessarily influence the annual variability in the water balance, as the presence or absence of a rock glacier in a watershed may only marginally influences the annual watershed hydrology. Other factors, such as the overall permafrost characteristics, terrain types and precipitation patterns, play more significant roles. Therefore, without further investigations and especially the assessment of the temporal scales of runoff contribution of ground ice melt from such extensive permafrost occurrences, its importance for the hydrological cycle cannot be addressed.

590 7 Appendix

~~Inverted tomograms of all ERT profiles for Site A and Site D~~

Author contributions. TM designed the study, participated in the geophysical campaigns, wrote the major part of the text, and made all figures. CHi planned, coordinated and participated at the geophysical campaigns, and did the processing of the geophysical data. CHa supervised and contributed to the study design. PW and LA coordinated the environmental impact assessment studies, which included the geophysical campaigns, borehole drilling and the excavation of test pits. They planned and coordinated the field logistics of the geophysical campaigns together with CHi, and provided further background information. All authors contributed actively to the discussion and interpretation of all data sets, and the intermediate and final version of the manuscript.

Competing interests. The authors declare that they have no conflict of interest.

600 *Acknowledgements.* The acquisition of the geophysical data set presented would not have been possible without the valuable support and hard work of numerous field helpers from Chile, Argentina and Switzerland. Therefore we sincerely thank all field helpers for their efforts in the field. The authors also would like to acknowledge the support from various private companies that agreed for having their data published, provided additional information and logistically supported the various field campaigns. Last but not least, we thank the two anonymous reviewers and the editor for their constructive feedback and suggestions that significantly improved the paper.

Appendix A: Appendix

605 **References**

- Arenson, L. U. and Jakob, M.: The significance of rock glaciers in the dry Andes - A discussion of Azócar and Brenning (2010) and Brenning and Azócar (2010), *Permafrost and Periglacial Processes*, 21, 282–285, <https://doi.org/10.1002/ppp.693>, 2010.
- Azócar, G. F. and Brenning, A.: Hydrological and geomorphological significance of rock glaciers in the dry Andes, Chile (27°–33° S), *Permafrost and Periglacial Processes*, 21, 42–53, <https://doi.org/10.1002/ppp.669>, 2010.
- 610 Azócar, G. F., Brenning, A., and Bodin, X.: Permafrost distribution modelling in the semi-arid Chilean Andes, *The Cryosphere*, 11, 877–890, <https://doi.org/10.5194/tc-11-877-2017>, 2017.
- Baldis, C. T. and Liaudat, D. T.: Permafrost model in coarse-blocky deposits for the Dry Andes, Argentina (28°–33° s), *Geographical Research Letters*, 46, 33–58, <https://doi.org/10.18172/cig.3802>, 2020.
- Bodin, X., Rojas, F., and Brenning, A.: Status and evolution of the cryosphere in the Andes of Santiago (Chile, 33.5° S.), *Geomorphology*, 615 118, 453–464, <https://doi.org/10.1016/j.geomorph.2010.02.016>, 2010.
- Brenning, A.: Climatic and geomorphological controls of rock glaciers in the Andes of Central Chile: Combining statistical modelling and field mapping, Dissertation, <https://doi.org/10.1002/ppp.528>, 2005.
- Buchli, T., Kos, A., Limpach, P., Merz, K., Zhou, X., and Springman, S. M.: Kinematic investigations on the Furggwanhorn Rock Glacier, Switzerland, *Permafrost and Periglacial Processes*, 29, 3–20, <https://doi.org/10.1002/ppp.1968>, 2018.
- 620 Bucki, A. K., Echelmeyer, K. A., and MacInnes, S.: The thickness and internal structure of Fireweed rock glacier, Alaska, U.S.A., as determined by geophysical methods, *Journal of Glaciology*, 50, 67–75, <https://doi.org/10.3189/172756504781830196>, 2004.
- Croce, F. A. and Milana, J. P.: Internal structure and behaviour of a rock glacier in the Arid Andes of Argentina, *Permafrost and Periglacial Processes*, 13, 289–299, <https://doi.org/10.1002/ppp.431>, 2002.
- Dafflon, B., Oktem, R., Peterson, J., Ulrich, C., Tran, A. P., Romanovsky, V., and Hubbard, S. S.: Coincident aboveground and belowground 625 autonomous monitoring to quantify covariability in permafrost, soil, and vegetation properties in Arctic tundra, *Journal of Geophysical Research: Biogeosciences*, 122, 1321–1342, <https://doi.org/10.1002/2016JG003724>, 2017.
- de Pasquale, G., Valois, R., Schaffer, N., and MacDonell, S.: Active and inactive Andean rock glacier geophysical signatures by comparing 2D joint inversion routines of electrical resistivity and refraction seismic tomography, *The Cryosphere Discussions*, pp. 1–33, <https://doi.org/10.5194/tc-2020-306>, 2020.
- 630 Devine, F., Kalanchey, R., Winkelmann, N., Gray, J., Melnyk, J., Elfen, S., Borntraeger, B., and Stillwell, I.: Pre-feasibility Study for the Filo del Sol Project Authored By, Tech. rep., 2019.
- Duguay, M., Edmunds, A., Arenson, L. U., and Wainstein, P.: Quantifying the significance of the hydrological contribution of a rock glacier – A review, GEOQuébec 2015, 68th Canadian Geotechnical Conferene, 7th Canadian Permafrost Conference, https://www.researchgate.net/publication/282402787_Quantifying_the_significance_of_the_hydrological_contribution_of_a_rock_glacier_-_A_review, 2015.
- 635 Esper Angillieri, M. Y.: A preliminary inventory of rock glaciers at 30° S latitude, Cordillera Frontal of San Juan, Argentina, *Quaternary International*, 195, 151–157, <https://doi.org/10.1016/j.quaint.2008.06.001>, 2009.
- Esper Angillieri, M. Y.: Permafrost distribution map of San Juan Dry Andes (Argentina) based on rock glacier sites, *Journal of South American Earth Sciences*, 73, <https://doi.org/10.1016/j.jsames.2016.12.002>, 2017.
- García, A., Ulloa, C., Amigo, G., Milana, J. P., and Medina, C.: An inventory of cryospheric landforms in the arid diagonal of South America 640 (high Central Andes, Atacama region, Chile), *Quaternary International*, 438, 4–19, <https://doi.org/10.1016/j.quaint.2017.04.033>, 2017.

- Gruber, S.: Derivation and analysis of a high-resolution estimate of global permafrost zonation, *The Cryosphere*, 6, 221–233, <https://doi.org/10.5194/tc-6-221-2012>, 2012.
- Gruber, S. and Hoelzle, M.: Statistical modelling of mountain permafrost distribution: Local calibration and incorporation of remotely sensed data, *Permafrost and Periglacial Processes*, 12, 69–77, <https://doi.org/10.1002/ppp.374>, 2001.
- 645 Gubler, S., Endrizzi, S., Gruber, S., and Purves, R. S.: Sensitivities and uncertainties of modeled ground temperatures in mountain environments, *Geoscientific Model Development*, 6, 1319–1336, <https://doi.org/10.5194/gmd-6-1319-2013>, 2013.
- Gude, M., Dietrich, S., Hauck, C., Kneisel, C., Krause, S., Molenda, R., Ruzicka, V., Zacharda, M., Kneisel, C., Krause, S., Molenda, R., Ruzicka, V., and Zacharda, M.: Evaluation of permafrost conditions in non-alpine scree slopes in Central Europe by geophysical methods, *EAEJA*, p. 8966, <https://ui.adsabs.harvard.edu/abs/2003EAEJA.....8966G/abstract>, 2003a.
- 650 Gude, M., Dietrich, S., Mäusbacher, R., Hauck, C., Molenda, R., Ruzicka, V., and Zacharda, M.: Permafrost conditions in non-alpine scree slopes in central Europe, in: *Permafrost* (eds. Phillips, Springman, Arenson), pp. 331–336, Swets & Zeitlinger, 2003b.
- Halla, C., Blothe, J., Liaudat, D. T., and Schrott, L.: Permafrost in the andes of Argentina -A significant hydrological resource, *Geographische Rundschau*, 70, 36–40, 2018.
- Halla, C., Henrik Blöthe, J., Tapia Baldis, C., Trombotto Liaudat, D., Hilbich, C., Hauck, C., and Schrott, L.: Ice content and interannual
655 water storage changes of an active rock glacier in the dry Andes of Argentina, *Cryosphere*, 15, 1187–1213, <https://doi.org/10.5194/TC-15-1187-2021>, 2021.
- Hauck, C. and Kneisel, C.: *Applied geophysics in periglacial environments*, Cambridge University Press, <https://doi.org/10.1017/CBO9780511535628>, 2008a.
- Hauck, C. and Kneisel, C.: Quantifying the ice content in low-altitude scree slopes using geophysical methods, in: C. Hauck & C. Kneisel
660 (eds.), *Applied Geophysics in Periglacial Environments*, pp. 153–164, Cambridge University Press, 2008b.
- Hauck, C., Böttcher, M., and Maurer, H.: A new model for estimating subsurface ice content based on combined electrical and seismic data sets, *The Cryosphere*, 5, 453–468, <https://doi.org/10.5194/tc-5-453-2011>, 2011.
- Hauck, C., Hilbich, C., and Mollaret, C.: A Time-lapse geophysical model for detecting changes in ground ice content based on electrical and seismic mixing rules, in: *23rd European Meeting of Environmental and Engineering Geophysics*, Malmö, 3.-7.9.2017,
665 <https://doi.org/10.3997/2214-4609.201702024>, 2017.
- Hausmann, H., Krainer, K., Brückl, E., and Mostler, W.: Internal structure and ice content of Reichenkar rock glacier (Stubai Alps, Austria) assessed by geophysical investigations, *Permafrost and Periglacial Processes*, 18, 351–367, <https://doi.org/10.1002/ppp.601>, 2007.
- Hilbich, C. and Hauck, C.: *Geophysical Surveys in Filo Del Sol*, unpublished internal report, Fribourg, 2018.
- Hilbich, C., Marescot, L., Hauck, C., Loke, M. H., and Mausbacher, R.: Applicability of electrical resistivity tomography monitoring to
670 coarse blocky and ice-rich permafrost landforms, *Tech. Rep. 3*, <https://doi.org/10.1002/ppp.652>, 2009.
- Hilbich, C., Fuss, C., and Hauck, C.: Automated time-lapse ERT for improved process analysis and monitoring of frozen ground, *Permafrost and Periglacial Processes*, 22, 306–319, <https://doi.org/10.1002/ppp.732>, 2011.
- Hilbich, C., Mollaret, C., and Hauck, C.: *Geophysical Surveys Mineras Los Pelambres (2016) and Rio Blanco (2017)*, Chile, unpublished internal report, Fribourg, 2018.
- 675 Hoelzle, M.: *Permafrost und Gletscher im Oberengadin*, Grundlagen und Anwendungsbeispiele für automatisierte Schätzverfahren, VAW Mitteilungen, ETH Zürich, 1994.
- Hoelzle, M., Scherler, M., and Hauck, C.: Permafrost ice as an important water resource for the future?, <https://www.openaccessgovernment.org/permafrost-ice-an-important-water-resource-future/39774/>, 2017.

- Hoelzle, M., Barandun, M., Bolch, T., Fiddes, J., Gafurov, A., Muccione, V., Saks, T., and Shahgedanova, M.: The status and role of the
680 alpine cryosphere in Central Asia, in: *The Aral Sea Basin*, pp. 100–121, Routledge, <https://doi.org/10.4324/9780429436475-8>, 2019.
- Hubbard, S. S., Gangodagamage, C., Dafflon, B., Wainwright, H., Peterson, J., Gusmeroli, A., Ulrich, C., Wu, Y., Wilson, C., Rowland, J.,
Tweedie, C., and Wullschleger, S. D.: Quantifying and relating land-surface and subsurface variability in permafrost environments using
LiDAR and surface geophysical datasets, *Hydrogeology Journal*, 21, 149–169, <https://doi.org/10.1007/s10040-012-0939-y>, 2013.
- IPCC: Special Report on the Ocean and Cryosphere in a Changing Climate (SROCC), Tech. rep., <https://www.ipcc.ch/srocc/>, 2019.
- 685 Jones, D. B., Harrison, S., Anderson, K., and Betts, R. A.: Mountain rock glaciers contain globally significant water stores, *Scientific Reports*,
8, 1–10, <https://doi.org/10.1038/s41598-018-21244-w>, 2018a.
- Jones, D. B., Harrison, S., Anderson, K., Selley, H. L., Wood, J. L., and Betts, R. A.: The distribution and hydrological significance of rock
glaciers in the Nepalese Himalaya, *Global and Planetary Change*, 160, <https://doi.org/10.1016/j.gloplacha.2017.11.005>, 2018b.
- Jones, D. B., Harrison, S., Anderson, K., and Whalley, W. B.: Rock glaciers and mountain hydrology: A review, *Earth-Science Reviews*, 193,
690 66–90, <https://doi.org/10.1016/j.earscirev.2019.04.001>, 2019.
- Kenner, R., Noetzli, J., Hoelzle, M., Raetzo, H., and Phillips, M.: Distinguishing ice-rich and ice-poor permafrost to map ground temperatures
and ground ice occurrence in the Swiss Alps, *Cryosphere*, 13, 1925–1941, <https://doi.org/10.5194/tc-13-1925-2019>, 2019.
- Koenig, C., Hilbich, C., Hauck, C., and Arenson, L.: Ground Temperature within Mountain Permafrost Zones of the Central Andes, in: *Swiss
Geoscience Meeting (SGM)*, 2019.
- 695 Liaudat, D. T., Sileo, N., and Dapeña, C.: Periglacial water paths within a rock glacier-dominated catchment in the Stepanek area, Central
Andes, Mendoza, Argentina, *Permafrost and Periglacial Processes*, <https://doi.org/10.1002/ppp.2044>, 2020.
- Masiokas, M., Rabatel, A., Rivera, A., Ruiz, L., Pitte, P., Ceballos, J. L., Barcaza, G., Soruco, A., and Bown, F.: A review of the current state
and recent changes of the Andean cryosphere, *Frontiers in Earth Science*, 8, 99, <https://doi.org/10.3389/FEART.2020.00099>, 2020.
- Maurer, H. and Hauck, C.: Instruments and methods: Geophysical imaging of alpine rock glaciers, *Journal of Glaciology*, 53, 110–120,
700 <https://doi.org/10.3189/172756507781833893>, 2007.
- Mewes, B., Hilbich, C., Delaloye, R., and Hauck, C.: Resolution capacity of geophysical monitoring regarding permafrost degradation
induced by hydrological processes, *Cryosphere*, 11, 2957–2974, <https://doi.org/10.5194/tc-11-2957-2017>, 2017.
- Minsley, B. J., Abraham, J. D., Smith, B. D., Cannia, J. C., Voss, C. I., Jorgenson, M. T., Walvoord, M. A., Wylie, B. K., Anderson, L.,
Ball, L. B., Deszcz-Pan, M., Wellman, T. P., and Ager, T. A.: Airborne electromagnetic imaging of discontinuous permafrost, *Geophysical
705 Research Letters*, 39, 2503, <https://doi.org/10.1029/2011GL050079>, 2012.
- Mollaret, C., Hilbich, C., Pellet, C., Flores-Orozco, A., Delaloye, R., and Hauck, C.: Mountain permafrost degradation documented through
a network of permanent electrical resistivity tomography sites, *The Cryosphere*, 13, 2557–2578, <https://doi.org/10.5194/tc-13-2557-2019>,
2019.
- Mollaret, C., Wagner, F. M., Hilbich, C., Scapozza, C., and Hauck, C.: Petrophysical Joint Inversion Applied to Alpine Permafrost Field
710 Sites to Image Subsurface Ice, Water, Air, and Rock Contents, *Frontiers in Earth Science*, 8, 85, <https://doi.org/10.3389/feart.2020.00085>,
2020.
- Monnier, S. and Kinnard, C.: Internal structure and composition of a rock glacier in the Andes (upper Choapa valley, Chile) using borehole
information and ground-penetrating radar, *Annals of Glaciology*, 54, 61–72, <https://doi.org/10.3189/2013AoG64A107>, 2013.
- Paterson, W.: *The Physics of Glaciers*, Elsevier, <https://doi.org/10.1016/c2009-0-14802-x>, 1994.
- 715 Pellet, C. and Hauck, C.: Monitoring soil moisture from middle to high elevation in Switzerland: Set-up and first results from the SOMO-
MOUNT network, *Hydrology and Earth System Sciences*, 21, 3199–3220, <https://doi.org/10.5194/hess-21-3199-2017>, 2017.

- Pellet, C., Hilbich, C., Marmy, A., and Hauck, C.: Soil moisture data for the validation of permafrost models using direct and indirect measurement approaches at three alpine sites, *Frontiers in Earth Science*, 3, <https://doi.org/10.3389/feart.2015.00091>, 2016.
- 720 Rangecroft, S., Harrison, S., and Anderson, K.: Rock Glaciers as Water Stores in the Bolivian Andes: An Assessment of Their Hydrological Importance, *Arctic, Antarctic, and Alpine Research*, 47, 89–98, <https://doi.org/10.1657/aaar0014-029>, 2015.
- Salzmann, N., Huggel, C., Rohrer, M., Silverio, W., Mark, B. G., Burns, P., and Portocarrero, C.: Glacier changes and climate trends derived from multiple sources in the data scarce Cordillera Vilcanota region, southern Peruvian Andes, *The Cryosphere*, 7, 103–118, <https://doi.org/10.5194/tc-7-103-2013>, 2013.
- 725 Schaffer, N., MacDonell, S., Réveillet, M., Yáñez, E., and Valois, R.: Rock glaciers as a water resource in a changing climate in the semi-arid Chilean Andes, *Regional Environmental Change*, 19, 1263–1279, <https://doi.org/10.1007/s10113-018-01459-3>, 2019.
- Schneider, S., Hoelzle, M., and Hauck, C.: Influence of surface and subsurface heterogeneity on observed borehole temperatures at a mountain permafrost site in the Upper Engadine, Swiss Alps, *The Cryosphere*, 6, 517–531, <https://doi.org/10.5194/tc-6-517-2012>, 2012.
- Staub, B.: The evolution of mountain permafrost in the context of climate change - towards a comprehensive analysis of permafrost monitoring data from the Swiss Alps, Tech. rep., 2015.
- 730 Tapia, J., Townley, B., Córdova, L., Poblete, F., and Arriagada, C.: Hydrothermal alteration and its effects on the magnetic properties of Los Pelambres, a large multistage porphyry copper deposit, *Journal of Applied Geophysics*, 132, 125–136, <https://doi.org/10.1016/J.JAPPGEO.2016.07.005>, 2016.
- Urrutia, R. and Vuille, M.: Climate change projections for the tropical Andes using a regional climate model: Temperature and precipitation simulations for the end of the 21st century, *Journal of Geophysical Research*, 114, D02 108, <https://doi.org/10.1029/2008JD011021>, 2009.
- 735 Wicky, J. and Hauck, C.: Numerical modelling of convective heat transport by air flow in permafrost talus slopes, *The Cryosphere*, 11, 1311–1325, <https://doi.org/10.5194/tc-11-1311-2017>, 2017.




Figure A1. Inverted tomograms of all ERT profiles for Site A and Site D